

Review

Arms Race between Enveloped Viruses and the Host ERAD Machinery

Dylan A. Frabutt and Yong-Hui Zheng *

Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing, MI 48824, USA

* Author to whom correspondence should be addressed: zhengyo@msu.edu; Tel.: +1-517-884-5314; Fax: +517-353-8957

Abstract: Enveloped viruses represent a significant category of pathogens that cause serious diseases in animals. These viruses express envelope glycoproteins that are singularly important during infection of host cells by mediating fusion between the viral envelope and host cell membranes. Despite low homology at protein levels, three classes of viral fusion proteins have, as of yet, been identified based on structural similarities. Their incorporation into viral particles is dependent upon their proper sub-cellular localization after being expressed and folded properly in the endoplasmic reticulum (ER). However, viral protein expression can cause stress in the ER, and host cells respond to alleviate the ER stress in the form of the unfolded protein response (UPR); the effects of which have been observed potentiating or inhibiting viral infection. One important arm of UPR is to elevate the capacity of the ER-associated protein degradation (ERAD) pathway, which is comprised of host quality control machinery that ensures proper protein folding. In this review, we provide relevant details regarding viral envelope glycoproteins, UPR, ERAD, and their interactions in host cells.

Keywords: enveloped viruses; viral glycoproteins; endoplasmic reticulum-associated degradation; ERAD; unfolded protein response; UPR; ER stress

1. Enveloped Viruses

Despite their vast diversity, animal viruses can be simply divided into two categories: non-enveloped viruses and enveloped viruses [1]. While non-enveloped viruses are wrapped with naked shells made of viral capsid proteins, enveloped viruses are covered with a lipid-bilayer on their matrix proteins, which is called viral envelope. The viral envelope is obtained from progenitor host cells during the budding process, which can be a portion of plasma membrane or intracellular membrane. On the surface of the enveloped viruses, there are peplomers that project from the viral envelope, which play a critical role in viral infection. These peplomers are also described as spikes, which are made of viral envelope glycoproteins. Envelope spikes serve to identify and bind to viral receptors on the host cell surface, allowing viral entry into cells and initiation of infection by mediating virus-cell fusion. Thus, the infectivity of enveloped viruses is absolutely dependent on the integrity of the viral envelope, and the functionality of the viral glycoproteins found therein.

Enveloped viruses are more stable than non-enveloped viruses under physiological conditions, at the expense of their sensitivity to high-temperature, low-pH, desiccation, or detergent-treatment, which limits their ability to withstand severe environments [2]. The entry of enveloped viruses requires formation of a fusion pore between viral envelope and cell membrane, through which viral genome is released into the cell. This fusion process is triggered by interactions between viral glycoproteins on viral envelope and viral receptors on the cell surface, which can occur directly at the plasma membrane at neutral pH or in endocytic compartments at either low or neutral pH [3]. In addition, enveloped viruses can also enter cells through direct cell-to-cell contacts via virological synapses to provide a

means by which the virus can cross the biophysical and immunological obstacles to infection [4]. The membrane penetration mechanism differs fundamentally in non-enveloped viruses, but similar strategies are used for their entry [5]. In general, enveloped animal viruses possess greater adaptability than non-enveloped animal viruses, and consequently, cause a number of severe diseases, such as acquired immunodeficiency syndrome (AIDS), influenza, severe acute respiratory syndrome (SARS), hemorrhagic fever, hepatitis, encephalitis, microcephaly, etc...

2. Virus Envelope Glycoproteins

The fusion between viral envelope and cell membranes is absolutely critical for the entry of enveloped viruses, which is usually triggered by the insertion of a viral envelope glycoprotein's (Env) fusion peptide into the host membrane. The vast majority of viral fusion proteins belong to type I transmembrane proteins, which have a single transmembrane domain (TMD), with their N-terminus outside cells and C-terminus inside cells. Most have been modeled as existing in a less stable prefusion state or a stable postfusion state. Many also oligomerize into trimeric fusogenic complexes in their postfusion states, forming trimeric hairpin structures on the viral envelope.

Based on their structural and mechanistic properties, viral fusion proteins have been classified into three distinct classes [6]. Class I fusion proteins are found in influenza viruses, paramyxovirus, retroviruses, and filoviruses viruses. They are first expressed as a polypeptide precursor and then cleaved by cellular proteases, yielding a transmembrane protein with an amino-terminal fusion peptide and a surface protein, which are attached either non-covalently or by a disulfide-bond. The core of the class I protein fusogenic domain is predominantly composed of α -helices, which contains a N-terminally located fusion peptide. These proteins trimerize and form a central coiled-coil structure with a three- α -helix bundle in the prefusion state, which refold into a six- α -helix bundle in the postfusion state.

Class II fusion proteins are found in flaviviruses, hepaciviruses, alphaviruses, togaviruses, and Rift Valley fever viruses. They depend on a second membrane glycoprotein as a chaperone, which is produced from the same polypeptide precursor where they are arrayed in tandem. When the chaperone is cleaved off, the fusion protein gains the fusogenic activity. These fusion proteins are mostly made of β -sheets and exist as homo- or heterodimers with the fusion peptides buried in internal loops in the prefusion state. In the postfusion state, these proteins undergo self-rearrangement into stable trimeric hairpins, exposing the fusion peptide and resulting in viral and host membrane fusion.

Class III fusion proteins are found in rhabdoviruses, herpesviruses, and baculoviruses. They are directly translated as a single protein from viral mRNA without involvement of protease cleavage, and trimerize in both pre- and postfusion states. Notably, they combine structural signatures found in both classes I and II, which include a central trimeric coiled coil, three domains predominantly made of β -sheets, and internal fusion peptides in the prefusion state. However, unlike in class I and II fusion proteins, the prefusion and postfusion states are reversible in class III fusion proteins.

In addition to these structure features, viral fusion proteins are subject to N-glycosylation at varying degrees. Although some viral envelope proteins such as the Dengue virus (DENV) E protein are glycosylated at relatively low levels [7], most of the other important human viruses are subject to heavy glycosylation. For example, HIV-1 Env precursor gp160 has ~35 potential N-linked glycosylation sites (PNGSs): ~30 in gp120 and ~4 in gp41 [8]; influenza HA molecules have 5 to 11 PNGSs depending on subtypes, with the majority of sites residing in the globular head of the molecule [9]; hepatitis C virus (HCV) E1 has 4 PNGSs and E2 has 9 PNGSs [10]; and Ebola virus GP has 17 PNGSs (15 in GP1 and 2 in GP2) [11]. HIV-1 Env and HCV E1 and E2 are so heavily glycosylated that ~50% of their respective molecular masses are derived from N-linked glycans. Most of the critical sites in these viral

glycoproteins are conserved during viral evolution, suggesting the important function of glycosylation in viral infections.

Glycosylation is one of the most common posttranslational modifications in eukaryotic cells, which is required for maintaining protein structure. Viruses have taken advantage of this benefit at nearly every step of the viral life cycle [12]. The immediate beneficiaries of glycosylation are the viral fusion proteins. *N*-glycosylation significantly promotes their folding and solubility, enhances subsequent trafficking of these viral proteins to their destinations, and ensures that they are properly processed and incorporated into virions. Nevertheless, glycosylation can have distinct effects that are both advantageous and detrimental to viral fitness. For example, if glycosylation occurs closely to the glycoprotein processing sites, it may block the precursor cleavage by proteases and inhibit viral infection [13]; if it occurs adjacent to the receptor-binding site, it may enhance the binding affinity and promote viral infection [14,15]. In addition, the high density of glycans on virions may form a shield to impede antibody attack and promote immune evasion. However, these glycans can also become epitopes for stimulating neutralizing antibodies and the innate immune response, making viruses more vulnerable to immune clearance [16]. Thus, there are multiple selective pressures on viral envelope glycosylation that will influence the pattern of glycosylation in order to achieve the optimal fitness in their hosts [17].

3. Glycosylation and Unfolded Protein Response (UPR)

Viruses are obligate intracellular parasites, and their glycoprotein biosynthesis and modification rely entirely on host cell machinery in the secretory pathway. Therefore, viral and host proteins are glycosylated in a similar manner by the same mechanism. Although glycans can be attached to polypeptide structures via several different mechanisms, Asparagine (*N*)-linked glycosylation represents a fundamental and the best characterized post-translational modification in eukaryotic organisms [18].

N-linked glycosylation starts from the membrane of the endoplasmic reticulum (ER), where the precursor tetradecasaccharide is assembled. This precursor consists of 2 *N*-acetylglucosamine (GlcNAc), 9 mannose (Man, 4 are α 1,2-linked), and 3 terminal glucose (Glc) residues distributed on three extended Man branches a, b, and c (Glc₃Man₉GlcNAc₂) (Figure 1A) [19,20]. When nascent polypeptides enter the ER lumen, the precursor is *en bloc* attached to Asn residues of a nascent polypeptide in a consensus Asn-X-(Ser/Thr) motif. After the attachment, these precursors are processed by a series of enzymes in both the ER and the Golgi apparatus to remold the core oligosaccharide into diverse *N*-linked glycan structures (Figure 1B). The 1st step in this process is the sequential removal of the two outermost Glc residues on branch A. The 1st Glc residue is removed by glucosidases I, resulting in di-glycosylated oligosaccharide Glc₂Man₉GlcNAc₂, which is recognized by an ER transmembrane lectin malectin [21]. The 2nd Glc residue is then removed by glucosidase II, resulting in mono-glycosylated oligosaccharide Glc₁Man₉GlcNAc₂, which is recognized by two other ER lectins, the membrane-bound calnexin (CNX) and/or soluble calreticulin (CRT). Interaction with these two chaperones segregates the newly formed glycoprotein and provides access to thiol-disulfide oxidoreductases such as ERp57, which promotes disulfide bond formation, resulting in protein folding into a native conformation. Once a protein is properly folded, glucosidase II cleaves the last Glc residue on branch A, which releases the protein from the CNX/CRT cycle. The ER class I α -mannosidase (ERManI) then cleaves the outermost Man residue on branch B on native proteins, resulting in oligosaccharide Man₈GlcNAc₂. These high-Man glycans are then recognized by lectins including ER-GIC53, vesicular integral membrane protein of 36Kda (VIP36), and VIP36-like (VIPL), which promote trafficking from the ER to the Golgi [22]. The remaining Man residues are cleaved by the Golgi mannosidases, and the glycan remodeling process is continued through the remainder of the *N*-glycosylation pathway, which generates functional glycoproteins that are secreted to cell surface (Figure 1B).

In addition to these chaperones and enzymes that promote protein folding, the ER is also equipped with a unique quality control mechanism that extracts and degrades proteins that are not correctly folded or assembled into their native conformation, which is called ER-associated protein degradation (ERAD) [23]. In fact, the folding efficiency of glycoproteins in the ER is very low, which requires cycles of association and dissociation from CNX/CRT to ensure the proper glycoprotein maturation. If glycoproteins with the Man₉GlcNAc₂ oligosaccharide display non-native conformations, they are reglucosylated by the UDP-Glc:unfolded glycoprotein glucosyltransferase (UGT1 or UGGT), and subject to additional rounds of re-engagement with the CNX/CRT machinery until folding is achieved. However, if a certain time frame for the folding is exceeded, proteins may never fold properly. Misfolded proteins are sequestered into in COP-II-dependent, highly mobile ER-derived quality control vesicles (QCVs), where ERManI is enriched (Figure 1B) [24]. Because ERManI is able to excise all α 1,2-Man residues when it is expressed at much higher levels *in vitro* [25], it may catalyze extensive demannosylation, resulting in production of low-Man oligosaccharide Man₅GlcNAc₂-containing glycoprotein precursors. The removal of the a branch terminal Man residue, which is the acceptor for Glc transferred by UGGT, disables these proteins from reengagement with CNX/CRT and re-entering into the folding cycle. Importantly, the low-Man N-glycans represent a tag for defective glycoproteins, targeting them to ERAD for degradation [26].

With only one-tenth of the total cell volume, the ER is responsible for the synthesis of the entirety of the secreted or membrane proteins, which account for one-third of total cellular proteins. Therefore, the ER has extremely high protein concentrations (100 mg/ml), which renders this organelle very susceptible to protein aggregation [27]. In addition, the protein folding is error prone, and this process can be further compromised by physiological and pathological perturbations. Moreover, genetic mutations may prohibit proteins from being folded properly. All these factors may cause accumulation of unfolded or misfolded proteins. When the level of these aberrant proteins exceeds the folding and clearance capacity of the ER, known as the ER homeostasis, it leads to a cellular stress response termed “ER stress”, which in turn activates the unfolded protein response (UPR) to restore the ER homeostasis [28]. ER stress is sensed by three ER transmembrane receptors: double-stranded RNA-activated protein kinase (PKR)-like ER kinase (PERK), inositol-requiring enzyme 1 (IRE1), and activating transcription factor 6 (ATF6), which are in association with another ER chaperone, the binding immunoglobulin protein (BiP, or GRP78). BiP dissociates from these sensors upon the ER stress, resulting in their activation and UPR to mitigate this stress [29,30]. IRE1 activates the transcription factor X-Box Binding Protein 1 (XBP-1), which in turn up-regulates expression of ERAD components to boost protein degradation; PERK phosphorylates the eukaryotic initiation factor (eIF)-2 α and halts protein translation; and ATF6 up-regulates expression of ER chaperones such as BiP, CNX, CRT, and protein disulfide isomerase (PDI) to increase folding efficiency. However, if these objectives are not achieved within a certain time span or the disruption is prolonged, UPR also activates pathways leading to cell death. Although PERK activation causes global inhibition of protein translation by blocking the eIF-2 α activity, it paradoxically enhances translation of the transcription factor ATF4. ATF4 then trans-activates the CCAAT/enhancer-binding protein-homologous protein (CHOP), which is a pro-apoptotic transcription factor, resulting in cell death by apoptosis [31].

Frabutt_Fig.1




Figure 1. (A) Schematic presentation of the N-linked core oligosaccharide structure. The core is composed of two N-acetylglucosamine (GlcNAc, blue), nine mannose (Man, red), and three glucose (Glc, yellow) residues. a, b, and c are three oligosaccharide branches. **(B) Schematic description of N-glycosylation, ERAD, and ER stress pathways.** Nascent polypeptide is translocated through sec61 into the rough ER, where the core oligosaccharide is transferred from a dolichol phosphate onto asparagine residues in NxS/T motifs (1). The two terminal glucose residues on the core oligosaccharide are trimmed by glucosidase I (GI) (2) and GII (3), respectively, allowing for the association with the chaperones, CNX and CRT, which promote folding to a native conformation. Eventually, the last terminal glucose residue will be trimmed by GII and the glycoprotein will maintain a native conformation (4), or misfold (7). Glycoproteins that reach a native conformation will have the terminal α1,2-Man residue on the b branch removed by ERMAnI, as a signal to allow it to traverse the canonical secretory pathway for surface presentation (6). Polypeptides unable to reach a native conformation (7) will engage in multiple rounds of the CNX/CRT cycle, facilitated by reglucosylation of the terminal glucose by UGGT (8), and trafficking between QCV (9) and the ERQC (10) under the ER stress condition. Terminally misfolded glycoproteins will be demannosylated to remove all α1,2-Man residues (11), followed by association with lectins OS9 and XTP3-B, for ERAD (12). Upon ER stress QCV will localize to the ERQC with their glycoprotein substrates (12). ERMAnI containing QCVs are rapidly recycled through autophagy (13). Without client glycoproteins to interact with, EDEMosome components are degraded through an autophagy-like mechanism (14). Viruses can hijack EDEMosomes to form double membrane vesicles (DMVs) that serve as platforms for their replication (15).

4. ER-Associated Protein Degradation (ERAD)

ERAD is a protein quality control mechanism conserved in all eukaryotic cells, which is an important arm of UPR, necessary to alleviate the ER stress [32]. ERAD results in selective dislocation of unfolded and misfolded proteins from the ER to the cytosol via specific membrane penetration machinery, which are subsequently degraded by the cytosolic ubiquitin proteasome system (UPS) [33]. It not only plays

an important role in the quality control of functional proteins produced from the ER, but also is critical for maintenance of the ER homeostasis by eliminating unfolded and misfolded proteins. Thus, ERAD is a central element of both the secretory pathway and UPR, and so far, more than 60 human diseases have been attributed to this pathway [34].

Although the vast majority of secreted proteins are glycosylated, the ER is responsible for the folding and assembly of both glycosylated and non-glycosylated proteins into functional complexes, which are subjected to ERAD quality control if they are misfolded. The process of ERAD can be divided into three steps: substrate recognition, retrotranslocation, and ubiquitylation/proteasomal degradation. In fact, extensive excision of α 1,2-Man residues from *N*-glycans sends an important signal to trigger misfolded glycoprotein degradation, which is catalyzed by class I mannosidases [35].

Class I mannosidases belong to the glycoside hydrolase family 47 (GH47), which are *exo-acting* α 1,2-mannosidases that are divided into three subfamilies [36]. The 1st subfamily consists of ERMAnI, which is supposed to cleave the outmost α 1,2-Man residue on the B branch from *N*-linked glycans in the ER. The 2nd subfamily consists of three Golgi α -mannosidase I, including GolgiManIA, GolgiManIB, and GolgiManIC, which cleave the remaining three α 1,2-Man residues in the Golgi complex for *N*-glycan maturation. The 3rd subfamily consists of the ER degradation-enhancing α -mannosidase-like proteins (EDEM) 1, 2, and 3. Although some EDEM orthologs in lower eukaryotes have detectable α 1,2-mannosidase activity, such activity has not been reported for any mammalian EDEM proteins *in vitro*. Nevertheless, there is evidence suggesting that these EDEM proteins should have the enzymatic activity *in vivo* [37,38]. Indeed, the extent of Man excision determines the fate of a glycoprotein, which could be either targeted to ERAD for degradation or sent to Golgi for normal secretion. ERMAnI exhibits a slow rate of enzymatic activity, which allows nascent proteins to perform multiple rounds of reglucosylation and achieve proper folding [39]. Properly folded glycoproteins should have one Man residue trimmed off from *N*-glycans by ERMAnI, which then interacts with the high-Man binding lectin, the ER-Golgi intermediate compartment-53 (ERGIC-53) [40], for trafficking from the ER to the Golgi (Figure 1B). However, if glycoproteins are misfolded terminally, the remaining three α 1,2-Man residues are excised from these molecules, which targets misfolded proteins for degradation [41].

It is still not completely understood how misfolded proteins are subjected to such extensive demannosylation in the ER and then targeted to ERAD. Although ERMAnI alone may be able to complete this task, there is evidence suggesting that additional GH47 enzymes are involved. It has been shown that elevation of the Golgi mannosidases could accelerate ERAD, so these enzymes may possibly be responsible for such extensive excision likely by trafficking back to the ER via an unknown mechanism [42]. In fact, the localization of the GolgiManIA has recently been observed in QCVs with other canonical ERAD machinery such as ERMAnI, and its overexpression and knockdown can, respectively, increase or retard trimming of misfolded glycoproteins from Man₉GlcNAc₂ to Man₅GlcNAc₂ *in vitro* [43]. In addition, upon ER stress, QCVs are converged to form the ER-derived quality compartments (ERQCs), where EDEM proteins are also sequestered (Figure 1B) [41]. EDEM1 and EDEM3 have been found to boost mannose trimming when overexpressed [38,44,45]. In addition, using a genomic knockout approach, it has been recently proposed that EDEM2 should play a central role in trimming of the outmost Man residue on b branch, whereas EDEM1 and EDEM3 should be responsible for trimming of the remaining α 1,2-Man residues. Accordingly, a “double check” model for misfolded glycoproteins has been proposed, which suggests that EDEM2 catalyzes the first step of Man trimming, and EDEM1 and EDEM3 contribute the second step [37]. Under these joint actions, all four α 1,2-Man residues are removed from the oligosaccharide, which is then recognized by the lectins OS-9 and XTP3-B via the mannose 6-phosphate receptor homology (MRH) domain (Figure 1B) [46,47]. Misfolded proteins are targeted to specific translocation channels (retrotranslocons) for retrotranslocation via an energy-dependent manner, which is facilitated by p97, a member of ATPases associated with diverse cellular activities (AAA) family, by catalyzing ATP hydrolysis [48].

It is still mysterious how these retrotranslocons are formed and how integral membrane and luminal ERAD substrates are exported across the ER membrane through these retrotranslocons. The 1st candidate channel is composed of Sec61 complex, which is comprised of α , β , and γ subunits. The α subunit crosses membrane 10 times, which forms a channel with the other subunits. The Sec61 heterotrimeric channel is the main translocon involved in co-translational protein transport into the ER [49]. However, there is evidence suggesting that the Sec61 translocon is also involved in retrotranslocation of ERAD substrates, implying the non-specificity and bi-directional property of this channel [50]. The 2nd candidate is Derlin-1, -2, or 3. Derlin-1 has 6 transmembrane domains, which may form channels via oligomerization [51]. The 3rd candidate is from the ERAD-specific E3 ligases. They have a large number of transmembrane domains, which are not only responsible for the polyubiquitylation, but could act as potential exit channels for ERAD substrates [52].

In *Saccharomyces cerevisiae*, there are two major types of RING-finger E3 ligase complexes Hrd1 and Doa10, which mediate ERAD by targeting discrete substrates [53]. Hrd1 was the 1st E3 enzyme identified in the ERAD pathway during study of HMG-CoA reductase degradation (Hrd) [54]. Hrd1 has 6 transmembrane helices in its N-terminal transmembrane domain and a catalytic RING domain in the soluble C-terminal region extended to the cytosol. Hrd1 has two mammalian orthologs named HRD1 and gp78, and its functioning E2 enzyme is known as Ubc7, which also has two mammalian orthologs Ube2g2 and Ube2g1 [55]. Hrd1 is unstable, so it is necessary for it to be stabilized by its co-factor Hrd3 in an equimolar ratio [56]. The mammalian ortholog of Hrd3 is SEL1L, which is required for ERAD substrate retrotranslocation [57]. Hrd1 interacts with Der1, the putative yeast channel protein that has 4 transmembrane helices. The Der1-Hrd1 interaction is bridged by another integral membrane protein Usa1, which is also required for Hrd1 oligomerization [58]. The Usa1 mammalian ortholog is called Herp, which also interacts with HRD1 and Derlin-1 and plays an important role in ERAD [59].

Doa10 was identified during studying the degradation of MAT- α 2-10 (doa), which is a yeast transcription factor. Doa10 is a ~150 kDa protein that has 14 transmembrane domains, which requires both Ubc6 and Ubc7 as E2 enzymes. The Doa10 mammalian ortholog is TEB4 (MARCH6), which plays a similar function in the ERAD pathway with similar subcellular distribution and topology [60]. The Ubc6 E2 enzyme has two mammalian orthologs, Ube2j1 and Ube2j2, both are involved in ERAD [61,62].

In *Saccharomyces cerevisiae*, ERAD is executed synergistically by Hrd1 and Doa10 with minimal redundancy because they exhibit different substrate specificity. Doa10 mainly triggers ubiquitylation of specific soluble proteins and membrane proteins with degrons exposed to the cytosol; a process referred to as ERAD-C [63,64]. Hrd1 is responsible for another two types of substrates, whose degradation is termed ERAD-L and ERAD-M. ERAD-L includes soluble luminal proteins in the ER and transmembrane proteins with degrons exposed to the ER lumen; ERAD-M includes transmembrane proteins with degrons embedded into the ER membrane [53]. Simultaneous inactivation of both genes has been shown to increase the sensitivity to heavy metal-induced cellular stress and exhibit an elevated UPR.

The regulation of protein folding and the functional relation between ERAD and UPR are much more complex in mammalian cells. In *Saccharomyces cerevisiae*, the CNX cycle does not exist due to lack of UGGT. In addition, protein synthesis is tightly controlled at the translational level by determination of the stoichiometry to avoid surplus production, resulting in minimal dependence on posttranslational regulation of protein expression. Moreover, although the yeast ERManI ortholog Mns1p and EDEM ortholog Htm1p are indispensable for ERAD, only one EDEM ortholog is present in yeast [65,66]. Because overly active ERAD may interfere with the regular protein folding process in the ER,

mammalian cells have evolved mechanisms to tightly regulate this quality control device by combination of compartmentalization and tuning.

EDEM1 is segregated into ER-derived, LC3-I-associated vesicles, which are called EDEMosomes, where EDEM1, OS-9, and SEL1L are concentrated when they lack client glycoproteins to dislocate (Figure 1B) [67]. Notably, unlike chaperones and the other enzymes, many ERAD regulators including ERManI, EDEM1, OS-9, HERP, and SEL1L are short-lived proteins, and ERManI, EDEM1, SEL1L, and OS-9 are targeted to the lysosomal pathway for degradation [67,68]. Thus, EDEMosomes are called ERAD tuning vesicles, which deliver their content to lysosomes for disposal via an autophagy-like pathway to reduce the ERAD capacity under natural conditions [69]. Additionally, lysosomal inhibitors are able to cause the accumulation of aggregating mutant of dysferlin in the ER when compared to the wild-type, which was used as evidence to propose that large protein aggregates are disposed of via an autophagy/lysosomal pathway, dubbed ERAD II [70]. However, under ER stress, most of these factors are highly induced, including the EDEM proteins, but not ERManI [38,71,72]. Under stress, QCVs are recruited to ERQC, resulting in accumulation of ERManI and its glycoprotein substrates [73]. Moreover, many other ERAD components are also concentrated in ERQC, which include EDEM1, HRD1, Derlin-1, Sec61 β , and Herp. Importantly, it has been found that EDEM1 stabilizes ERManI and increases its protein expression at steady-state levels [74]. Such enrichment of these critical components accelerates efficient assembly of the ERAD machinery, potentiating the degradation of misfolded glycoproteins and alleviating ER stress.

5. Viruses and UPR

During infection, viruses are able to hijack the host translational machinery and saturate the ER with viral proteins. Not only do viruses use the ER to generate their glycoproteins, but some even utilize the ER as their site to assemble progeny particles [75]. Such accumulation of viral proteins in the ER places a heavy demand on the protein folding machinery, which may cause ER stress, and in turn, activate UPR, resulting in restoration of the ER homeostasis or apoptosis. So far, at least 36 viruses have been found to be able to induce ER stress, and activate the three UPR stress signaling pathways [76].

Enveloped viruses may bud through the plasma membrane or an intracellular compartment. In addition, their envelope glycoproteins are targeted to the ER for post-translational modifications and folding. Not surprisingly, many viral envelope glycoproteins are significant inducers of the UPR, which includes HCV [77], hepatitis B virus (HBV) [78], coronaviruses [79], alphaviruses [80], and retroviruses [81]. As introduced earlier, the UPR utilizes three different mechanisms to alleviate ER stress: reducing global protein translation, increasing the ER folding capacity, and enhancing ERAD by activating the PERK, ATF6, or IRE1-XBP1 pathways, respectively.

Viral infections may activate these pathways, resulting in inhibition or enhancement of viral replication. For example, the PERK-mediated global translation shutdown is a very effective antiviral mechanism, and a similar shutdown by the double-stranded RNA (dsRNA)-activated protein kinase (PKR) has been used in the interferon pathway to defend against viral infection [82]. Conceivably, viruses have evolved a number of strategies to circumvent the detrimental effect of UPR to establish productive infection. Some viruses such as HCV and Semliki Forest virus (SFV) are still able to produce viral proteins even when the cellular translational machinery is shut down, because these viruses have their own internal ribosome entry site (IRES) to recruit and assemble the ribosomal initiation complex for protein expression [83]; Epstein-Barr virus (EBV), herpes simplex virus (HSV), and African swine fever virus (ASFV) can counteract the PERK-mediated eIF2 α phosphorylation by activating an eIF2 α phosphatase PP1 [84-86]; HCV E2 protein directly interacts with PERK to prevent ER stress sensing by acting as a pseudo-substrate to block the PERK activity [87]. In addition to combating the UPR, viruses also take advantages of UPR pathways to benefit their replication. For examples, influenza A virus (IAV)

replication is promoted by activation of the IRE1-XBP1 pathway [88]; ATF6 activation promotes ASFV, lymphocytic choriomeningitis virus (LCMV), DENV, human cytomegalovirus (HCMV), and Japan encephalitis virus (JEV) replication [89,90]; and ATF4 activation enhances human immunodeficiency virus type 1 (HIV-1) replication [91]. Thus, despite the detrimental effects, viruses have evolved to manipulate host UPR signaling pathways to promote viral infections.

Below, we will focus on the roles of ERAD played in virus replication, which is the main target of this review.

6. Roles of ERAD in Promotion of Virus Replication

As introduced earlier, ERAD transports unfolded/misfolded proteins from the ER into the cytosol for proteasomal degradation. Conceivably, viruses can manipulate and exploit this cellular machinery to degrade several important host factors to promote their propagation.

Herpesviruses have evolved multiple mechanisms to suppress the host immune response via ERAD. Major histocompatibility complex (MHC) molecules play an indispensable role in triggering immediate immune response to inhibit virus infections. Herpesviruses may inhibit MHC class I (MHC-I) expression by targeting these molecules to ERAD for degradation. For example, HCMV produces two transmembrane proteins US2 and US11, each is sufficient to bind to MHC-I heavy chains, causing their dislocation from the ER to the cytosol for degradation [92]. Notably, US2 and US11 use different mechanisms to degrade MHC-I. Although both US2 and US11 interact with the p97 ATPase, the MHC-I dislocation by US2 is dependent on signal peptide peptidase (SPP), whereas that by US11 is dependent on Derlin-1 and SEL1L [93]. In addition, the MHC-I degradation induced by US2 requires the TRC8/MARCH6 and that by US11 requires the MARCH7/axotrophin E3 ligase [94,95]. Mouse gammaherpesvirus 68 (MHV68) uses another mechanism to inhibit MHC-I. MHV68 produces a protein termed MK3, which is a Ring-finger E3 ligase anchored on the ER membrane. MK3 interacts with MHC-I heavy chain molecules, and it also associates with the transporter-associated with antigen processing (TAP), p97, and Derlin-1, resulting in rapid degradation of the MHC-I by proteasomes [96]. Thus, herpesviruses have evolved numerous strategies to block the MHC antigen presentation and evade host immune response for establishing persistent infection.

Primate lentiviruses also harness the ERAD pathway to promote their replication via downregulation of their receptor CD4. CD4 downregulation prevents superinfection and promotes viral release by interrupting viral receptor-envelope interactions on plasma membrane, leading to a controlled and productive viral infection and immunodeficiency [97]. These viruses produce two accessory proteins Nef and Vpu to trigger CD4 degradation via two distinctive mechanisms [98]. Nef uses the endocytic pathway to redirect CD4 from cell surface, or interfere with the transport of newly synthesized CD4 from the trans-Golgi network (TGN) to the cell surface, resulting in CD4 dislocation to endosomes and degradation by lysosomes [99]. However, Vpu interacts with CD4 in the ER and induces CD4 proteasomal degradation via ERAD [100]. Vpu is a small transmembrane protein encoded by HIV-1 and some simian immunodeficiency virus (SIV) isolates. Vpu forms ion conductive membrane pores; it also interacts with β -transducin repeat-containing proteins (β TrCP), which are F-box/WD repeat-containing proteins that are part of the Skp1-Cul1-F-box (SCF) E3 ubiquitin ligase complex [101]. The Vpu-induced CD4 degradation is strictly dependent on the SCF- β TrCP complex [102]. Notably, this E3 ligase complex is not associated with the ER membrane, and therefore does not normally function in ERAD. However, the degradation also requires the cytosolic ATPase p97 and its cofactors UFD1L and NPL4, which are key components of the ERAD machinery, suggesting that CD4 is degraded via ERAD [102]. Nevertheless, the degradation is not dependent on HRD1, SEL1L, and UBC7, suggesting that an atypical ERAD pathway is involved in the CD4 degradation.

In addition to degradation, viruses may harness ERAD components to benefit their replication. First, ERAD can promote viral protein expression. Mouse mammary tumor virus (MMTV) is a betaretrovirus, which produces a protein termed Rem to export viral unspliced RNAs from the nucleus to the cytosol for protein expression. Although Rem is synthesized in the ER, it escapes the secretory pathway via retrotranslocation to the cytosol through ERAD, and this trafficking process is dependent on the p97 ATPase and Derlin-1 [103]. Similarly, Hepatitis E virus (HEV) ORF2 is an N-linked glycoprotein, but functions as the major capsid protein. Although it is expressed in the ER, it depends on ERAD components to exit from the ER to the cytoplasm without being polyubiquitylated [104].

Second, ERAD can promote virus entry. Polyomaviruses (PyV) enter cells through the ER and then replicate in nuclei [105]. To get from the ER to the nucleus, these viruses can cross the ER membrane into the cytosol via the ERAD translocons [106]. An example of this is mouse PyV, which uses Derlin-2, whereas simian virus 40 (SV40) uses Derlin-1 and SEL1L complex for dislocation [106,107]. In addition, the proteasome machinery is also required for the human BK PyV exit from the ER [107].

Third, ERAD can promote virus replication. The replication of positive-strand RNA viruses normally involves formation of double-membrane vesicles (DMVs) and convoluted membranes (CMs) by rearrangement of cellular membranes, which segregates and protects viral proteins and genomes from the host's innate immune response. As introduced earlier, the ERAD activity can be adjusted by ERAD tuning vesicles termed EDEMosomes (Figure 1B), which display nonlipidated LC3 and segregate the ERAD factors EDEM1, OS-9, and SEL1L from the ER lumen [69]. By comparing the similarity between DMVs and EDEMosomes, it has been discovered that mouse hepatitis virus (MHV), equine arteritis virus (EAV), and JEV indeed replicate in these ERAD tuning vesicles [108]. Thus, these viruses can subvert EDEMosomes as their replication vesicles to promote infection [109].

7. Roles of ERAD in Inhibition of Virus Replication

Although ERAD has been frequently manipulated by a number of viruses to promote infection or attenuate immune response, it may also function directly as an antiviral device to protect host cells from infection. Because viral envelope glycoprotein production and folding take place in the ER, these viral proteins may become the primary targets for ERAD, resulting in inhibition of viral infection.

Primate lentiviruses including HIV and SIV have very lower number of envelope glycoproteins on their surface, and the average copy number is ~14 Env trimers per virion [110,111]. In contrast, IFA, Sendai virus, HSV, and Moloney murine leukemia virus (MoMuLV) have much more envelope glycoprotein on their surface [112-115]. It has been speculated that the exceptionally low number of Env spikes may protect HIV-1 from host immune response [116]. Consistently, it was observed decades ago that the efficiency of HIV-1 Env expression in infected cells is extremely low: almost 85% Env proteins are retained in the ER and get degraded [117-119]. This degradation mechanism was not clear until we recently reported that HIV-1 Env glycoproteins are targeted to ERAD for degradation.

From completely unrelated studies, we isolated HIV-1 non-permissive (NP) and permissive (P) T cell clones N2-NP and N5-P from the original CEM.NKR human T cell line [120]. Our initial analysis uncovered that HIV-1 replication is restricted from the 2nd round of viral life cycle in N2-NP cells, resulting in ~1,000-fold inhibition when compared in N5-P. Further transcriptome analysis by microarrays uncovered that N2-NP cells overexpress the mitochondrial translocator protein (TSPO), which strongly inhibits the HIV-1 Env expression [121]. TSPO interacts with the mitochondrial permeability transition pore (mPTP) complex, which includes the outer membrane protein voltage-dependent anion channel (VDAC) protein, the inner membrane protein adenine nucleotide translocase (ANT), and the mitochondrial matrix protein cyclophilin D (CypD) [122]. TSPO binds to VDAC and contributes to the regulation of the mitochondrial membrane permeability by the mPTP complex [123]. Our results suggested that TSPO overexpression could reduce the oxidative redox status in the ER,

which interferes with the Env oxidative folding process, resulting in Env degradation. Consistently, the rapid Env degradation in N2-NP cells was rescued by kifunessine, an effective ERAD inhibitor by inhibiting the GH47 enzyme activity [124], suggesting that HIV-1 is degraded via ERAD in N2-NP cells.

To further explore the Env degradation mechanism, we investigated which of those four ER-associated GH47 enzymes is responsible for the Env degradation. Notably, when ERManI, EDEM1, EDEM2, and EDEM3 were ectopically expressed in 293T cells, only ERManI strongly inhibited the Env expression in a dose-dependent manner. In addition, when the endogenous ERManI was knocked out by CRSPR/Cas9, TSPO was no longer able to suppress the Env expression [125]. These results demonstrated that ERManI should be responsible for the initiation of HIV-1 Env degradation via ERAD. Human ERManI is a 699-amino-acid, 79.5-kDa, type II membrane protein, which is divided into N-terminal cytoplasmic domain (CD), transmembrane (TM) helix, luminal 'stem' region, and catalytic domain [126,127]. Using immunoprecipitation assay, we found that HIV-1 Env interacts with the catalytic domain of ERManI [125]. The structure of this catalytic domain shows an (α)₇-barrel composed of 14 consecutive helices [128]. In the catalytic domain, there are seven residues that are critical for ERManI function: C527 and C556 form a highly conserved disulphide bond and were reportedly critical for the protein folding [129]; E330, D463, and E599 were proposed as catalytic residues [128]; R334C and E397K mutations are found in nonsyndromic autosomal-recessive intellectual disability (NS-ARID) disease [130]; and the R334C mutation is also found in the congenital disorders of glycosylation [131]. All these residues have been found to be required for the Env degradation, suggesting that the mannosidase activity is important for the ERManI activity. ERManI also targets the terminally misfolded human alpha1-antitrypsin variant null (Hong Kong) (NHK) for degradation via ERAD, but neither its catalytic activity nor its catalytic domain is required for this degradation, suggesting that different mechanisms are involved in HIV-1 Env and NHK degradations [132]. We have also found that the viral protein R (Vpr) of HIV-1 enhances viral replication in monocyte-derived macrophages (MDMs) and dendritic cells (MDDCs) by rescuing Env from ERAD degradation through the ERAD (II) autophagy pathway. We concluded this since compounds known to facilitate glycoprotein folding (PK11195 and As₂O₃), inhibit ER α -mannosidases crucial for ERAD (Kifunessine), and block lysosomal proteases (Bafilomycin) were able to rescue envelope expression and infectivity in Δ Vpr background to that of wild-type virus [133].

As aforementioned, unlike ERManI, whose expression is irresponsive to UPR, the expression of the EDEMs is induced upon UPR via the IRE1/XBP activation pathway, which boosts ERAD and alleviates the ER stress. Although ectopic expression of EDEMs did not inhibit HIV-1 Env expression [125], these proteins inhibit the expression of some other envelope glycoproteins. HBV expresses three surface glycoproteins, the large (L), middle (M), and small (S), which are translated from different initiation codons within the same open reading frame (ORF) and share the tetra-spanning transmembrane domains in the S protein. The N-terminus of the M and L protein contain additional preS2 and preS1-preS2 domains, respectively. The common S domain has an N-glycosylation site, and the M preS2 domain has another site. Overexpression of the surface proteins is sufficient to activate the IRE1/XBP1 pathway and elevate EDEM1, EDEM2, and EDEM3 expression. Importantly, EDEM1 overexpression destabilizes S, M, and L, and EDEM1 silencing stabilizes their expression [134]. In addition, it is the autophagy/lysosomal pathway, but not the proteasomal pathway that is involved in the degradation of HBV surface glycoproteins, further complicating our understanding of viral protein degradation process via ERAD [134].

HCV has two N-glycosylated envelope proteins E1 and E2 on the surface of virions, which are type I transmembrane proteins expressed from a common viral polyprotein precursor. HCV infection strongly induces the activation of the IRE1 stress sensor, resulting in elevation of EDEM1, EDEM2, and EDEM3, but not the ERManI expression. Both EDEM1 and EDEM3 but not EDEM2 interact with E2, and overexpression of these two proteins induces E2 polyubiquitylation and degradation. Conversely,

knockdown of EDEM1 expression or treatment with kifunesine increases E2 expression, and also reduces the interaction of EDEM1 and EDEM3 with SEL1L [135]. Taken together, these results strongly suggest that EDEM proteins are able to extract viral polypeptides from the ER quality control cycle, and degrade them via ERAD. However, since none of these proteins can target JEV E protein to ERAD for degradation, not every viral glycoprotein is recognizable by these proteins [135]. *In vivo* experiments on patients with chronic liver injury were unable to identify up-regulation of UPR and ERAD elements in diseased versus control patients, which adds further complexity to the potential role of UPR during HCV infection [136].

ERAD has also been implicated in the degradation of HCMV glycoproteins, gH and gL, via the 26S proteasome. HCMV produces at least 65 unique glycoproteins, with 4 homologues to the HSV glycoproteins, gH, gB, gL, and gM [137]. The glycoproteins, gH and gL, are constituents of the gCII type complexes found on the surface of HCMV virions. The gCII trimeric complex between gH, gL, and gO can initiate pH independent fusion [138]. In addition, a pentameric complex between gH, gL, and the gene products U128, U130, and U131 is able to mediate entry into different cell types via pH-dependent receptor-mediated endocytosis; a process that requires the trimeric gH/gL/gO complex [139]. While it has been known that the glycoprotein gL stabilizes the expression and potentiates surface localization of gH for many years [140], only recently has it been shown that gH is degraded via ERAD in the absence of gL [141]. Replacement of the cytoplasmic tail of gH with that of the human CD4 protein subverted gH degradation via ERAD, potentiating surface expression.

The current studies available describe two paradigms for ERAD to target viral glycoproteins for degradation: ERManI-mediated, which targets HIV-1 Env, and EDEM-mediated, which can target HCV and HBV surface glycoproteins. GH47 family members share a common catalytic mannosidase homology domain of ~440-residues [45], and the three catalytic residues E330, D463, and E599 found in ERManI are all conserved in these proteins [36]. Nevertheless, there is little protein sequence homology beyond this domain among these proteins. Unlike ERManI, all three EDEMs are ER-luminal proteins, although the signal sequence of EDEM1 is resistant to cleavage [142]. EDEM3 has two novel features including additional protease-associated domain of unknown function and a KDEL signal for ER retention [38]. Whether or how coordination between the EDEMs and ERManI facilitates ERAD is still a convoluted issue. Due to lysosomal degradation mediated by the N-terminal cytoplasmic tail, ERManI is expressed at very low basal levels in cells, and its expression is not induced by UPR [74]. Such proteolytically driven checkpoint control of ERManI expression may contribute to establish glycoprotein quality control at a baseline level, which maintains the ER homeostasis without activation of IRE1/XBP1. However, if this basic mechanism fails to restore the ER homeostasis, IRE1/XBP1 is induced to elevate expression of the EDEMs, which will increase ERAD. Consistently, it was found that unlike HCV and HBV, although HIV-1 induces UPR, it barely activates the IRE1/XBP1 pathway, which may explain why HIV-1 Env is not directly targeted by EDEM proteins [81]. Nevertheless, these two different arms of ERAD do not exclude the role of the EDEMs in ERManI-mediated degradation. EDEMs may accelerate release of terminally misfolded glycoproteins from the CNX/CRT cycle, and thereby help ERManI to conduct more extensive demannosylation [143]; and the association of EDEM with SEL1L may further accelerate the cytosolic delivery of misfolded proteins [144]. Moreover, EDEM1 may form a complex with ERManI, which stabilizes ERManI by suppression of its proteolytic degradation [74]. Discrepancies concerning the localization of ERManI with various labs determining colocalization with the ER, Golgi, or ER-Golgi intermediate compartments and quality control vesicles, lends credence to both current theories that ERManI is either a Golgi checkpoint in quality control that will return misfolded proteins back to the ER for further processing or that it resides in quality control vesicles with glycoprotein substrates as part of the CNX/CRT cycle [24,145].

8. Conclusion

It is well established that viruses have evolved to manipulate host UPR and ERAD to optimize their replication, whether they are 'tuning' host quality control to ensure the proper folding of their envelope glycoproteins, circumventing ERAD in order to prevent degradation of their viral envelope glycoproteins, or hijacking ERAD to dispose of host proteins. There are still many questions left to be answered, including the identities of the dislocons that each envelope glycoprotein is targeted to, the motifs or patterns that allow α 1,2-mannosidases to differentiate between native and misfolded glycoproteins, why some viral proteins are disproportionately targeted (HCMV gH), and the roles that the UPR and ERAD play *in vivo* during viral infections. In general, these are very exciting areas that need more extensive efforts to study.

Acknowledgments

YHZ is supported by National Institutes of Health grants AI106477 and AI120189. DAF is supported by Russell B. Duvall Award for graduate studies.

Author Contributions:

DAF and YHZ wrote and edited the manuscript.

Conflicts of Interest:

The authors declare no conflict of interest.

References

1. Knipe, D.M.; Howley, P.M. *Fields virology*. 6th ed.; Wolters Kluwer/Lippincott Williams & Wilkins Health,: Philadelphia, PA, 2013; p 2 volumes.
2. Dimitrov, D.S. Virus entry: Molecular mechanisms and biomedical applications. *Nature reviews. Microbiology* **2004**, *2*, 109-122.
3. Grove, J.; Marsh, M. The cell biology of receptor-mediated virus entry. *The Journal of cell biology* **2011**, *195*, 1071-1082.
4. McDonald, D.; Wu, L.; Bohks, S.M.; KewalRamani, V.N.; Unutmaz, D.; Hope, T.J. Recruitment of hiv and its receptors to dendritic cell-t cell junctions. *Science* **2003**, *300*, 1295-1297.
5. Tsai, B. Penetration of nonenveloped viruses into the cytoplasm. *Annual review of cell and developmental biology* **2007**, *23*, 23-43.
6. Podbilewicz, B. Virus and cell fusion mechanisms. *Annual review of cell and developmental biology* **2014**, *30*, 111-139.
7. Lee, E.; Leang, S.K.; Davidson, A.; Lobigs, M. Both e protein glycans adversely affect dengue virus infectivity but are beneficial for virion release. *Journal of virology* **2010**, *84*, 5171-5180.
8. Wang, W.; Nie, J.; Prochnow, C.; Truong, C.; Jia, Z.; Wang, S.; Chen, X.S.; Wang, Y. A systematic study of the n-glycosylation sites of hiv-1 envelope protein on infectivity and antibody-mediated neutralization. *Retrovirology* **2013**, *10*, 14.
9. Tate, M.D.; Job, E.R.; Deng, Y.M.; Gunalan, V.; Maurer-Stroh, S.; Reading, P.C. Playing hide and seek: How glycosylation of the influenza virus hemagglutinin can modulate the immune response to infection. *Viruses* **2014**, *6*, 1294-1316.
10. Helle, F.; Duverlie, G.; Dubuisson, J. The hepatitis c virus glycan shield and evasion of the humoral immune response. *Viruses* **2011**, *3*, 1909-1932.
11. Lennemann, N.J.; Rhein, B.A.; Ndungo, E.; Chandran, K.; Qiu, X.; Maury, W. Comprehensive functional analysis of n-linked glycans on ebola virus gp1. *mBio* **2014**, *5*, e00862-00813.
12. Zhang, M.; Gaschen, B.; Blay, W.; Foley, B.; Haigwood, N.; Kuiken, C.; Korber, B. Tracking global patterns of n-linked glycosylation site variation in highly variable viral glycoproteins: Hiv, siv, and hcv envelopes and influenza hemagglutinin. *Glycobiology* **2004**, *14*, 1229-1246.

13. Deshpande, K.L.; Fried, V.A.; Ando, M.; Webster, R.G. Glycosylation affects cleavage of an h5n2 influenza virus hemagglutinin and regulates virulence. *Proceedings of the National Academy of Sciences of the United States of America* **1987**, *84*, 36-40.
14. Klenk, H.D.; Wagner, R.; Heuer, D.; Wolff, T. Importance of hemagglutinin glycosylation for the biological functions of influenza virus. *Virus research* **2002**, *82*, 73-75.
15. Wagner, R.; Wolff, T.; Herwig, A.; Pleschka, S.; Klenk, H.D. Interdependence of hemagglutinin glycosylation and neuraminidase as regulators of influenza virus growth: A study by reverse genetics. *Journal of virology* **2000**, *74*, 6316-6323.
16. Doores, K.J. The hiv glycan shield as a target for broadly neutralizing antibodies. *The FEBS journal* **2015**, *282*, 4679-4691.
17. Vigerust, D.J.; Shepherd, V.L. Virus glycosylation: Role in virulence and immune interactions. *Trends in microbiology* **2007**, *15*, 211-218.
18. Moremen, K.W.; Tiemeyer, M.; Nairn, A.V. Vertebrate protein glycosylation: Diversity, synthesis and function. *Nat Rev Mol Cell Biol* **2012**, *13*, 448-462.
19. Helenius, A.; Aebi, M. Roles of n-linked glycans in the endoplasmic reticulum. *Annual review of biochemistry* **2004**, *73*, 1019-1049.
20. Lamriben, L.; Graham, J.B.; Adams, B.M.; Hebert, D.N. N-glycan-based er molecular chaperone and protein quality control system: The calnexin binding cycle. *Traffic* **2016**, *17*, 308-326.
21. Schallus, T.; Jaech, C.; Feher, K.; Palma, A.S.; Liu, Y.; Simpson, J.C.; Mackeen, M.; Stier, G.; Gibson, T.J.; Feizi, T., et al. Malectin: A novel carbohydrate-binding protein of the endoplasmic reticulum and a candidate player in the early steps of protein n-glycosylation. *Molecular biology of the cell* **2008**, *19*, 3404-3414.
22. Slominska-Wojewodzka, M.; Sandvig, K. The role of lectin-carbohydrate interactions in the regulation of er-associated protein degradation. *Molecules* **2015**, *20*, 9816-9846.
23. Meusser, B.; Hirsch, C.; Jarosch, E.; Sommer, T. Erad: The long road to destruction. *Nature cell biology* **2005**, *7*, 766-772.
24. Benyair, R.; Ogen-Shtern, N.; Mazkereth, N.; Shai, B.; Ehrlich, M.; Lederkremer, G.Z. Mammalian er mannosidase i resides in quality control vesicles, where it encounters its glycoprotein substrates. *Molecular biology of the cell* **2015**, *26*, 172-184.
25. Herscovics, A.; Romero, P.A.; Tremblay, L.O. The specificity of the yeast and human class i er alpha 1,2-mannosidases involved in er quality control is not as strict previously reported. *Glycobiology* **2002**, *12*, 14G-15G.
26. Frenkel, Z.; Gregory, W.; Kornfeld, S.; Lederkremer, G.Z. Endoplasmic reticulum-associated degradation of mammalian glycoproteins involves sugar chain trimming to man6-5glcnac2. *The Journal of biological chemistry* **2003**, *278*, 34119-34124.
27. Hetz, C.; Martinon, F.; Rodriguez, D.; Glimcher, L.H. The unfolded protein response: Integrating stress signals through the stress sensor ire1alpha. *Physiol Rev* **2011**, *91*, 1219-1243.
28. Gardner, B.M.; Pincus, D.; Gotthardt, K.; Gallagher, C.M.; Walter, P. Endoplasmic reticulum stress sensing in the unfolded protein response. *Cold Spring Harbor perspectives in biology* **2013**, *5*, a013169.
29. Bertolotti, A.; Zhang, Y.; Hendershot, L.M.; Harding, H.P.; Ron, D. Dynamic interaction of bip and er stress transducers in the unfolded-protein response. *Nature cell biology* **2000**, *2*, 326-332.
30. Walter, P.; Ron, D. The unfolded protein response: From stress pathway to homeostatic regulation. *Science* **2011**, *334*, 1081-1086.
31. Han, J.; Back, S.H.; Hur, J.; Lin, Y.H.; Gildersleeve, R.; Shan, J.; Yuan, C.L.; Krokowski, D.; Wang, S.; Hatzoglou, M., et al. Er-stress-induced transcriptional regulation increases protein synthesis leading to cell death. *Nature cell biology* **2013**, *15*, 481-490.
32. Romisch, K. Endoplasmic reticulum-associated degradation. *Annual review of cell and developmental biology* **2005**, *21*, 435-456.
33. Lemus, L.; Goder, V. Regulation of endoplasmic reticulum-associated protein degradation (erad) by ubiquitin. *Cells* **2014**, *3*, 824-847.

34. Guerriero, C.J.; Brodsky, J.L. The delicate balance between secreted protein folding and endoplasmic reticulum-associated degradation in human physiology. *Physiol Rev* **2012**, *92*, 537-576.
35. Hebert, D.N.; Molinari, M. Flagging and docking: Dual roles for n-glycans in protein quality control and cellular proteostasis. *Trends Biochem Sci* **2012**, *37*, 404-410.
36. Mast, S.W.; Moremen, K.W. Family 47 alpha-mannosidases in n-glycan processing. *Methods in enzymology* **2006**, *415*, 31-46.
37. Ninagawa, S.; Okada, T.; Sumitomo, Y.; Kamiya, Y.; Kato, K.; Horimoto, S.; Ishikawa, T.; Takeda, S.; Sakuma, T.; Yamamoto, T., et al. Edem2 initiates mammalian glycoprotein erad by catalyzing the first mannose trimming step. *The Journal of cell biology* **2014**, *206*, 347-356.
38. Hirao, K.; Natsuka, Y.; Tamura, T.; Wada, I.; Morito, D.; Natsuka, S.; Romero, P.; Sleno, B.; Tremblay, L.O.; Herscovics, A., et al. Edem3, a soluble edem homolog, enhances glycoprotein endoplasmic reticulum-associated degradation and mannose trimming. *The Journal of biological chemistry* **2006**, *281*, 9650-9658.
39. Aikawa, J.; Matsuo, I.; Ito, Y. In vitro mannose trimming property of human er alpha-1,2 mannosidase i. *Glycoconjugate journal* **2012**, *29*, 35-45.
40. Hauri, H.P.; Kappeler, F.; Andersson, H.; Appenzeller, C. Ergic-53 and traffic in the secretory pathway. *Journal of cell science* **2000**, *113* (Pt 4), 587-596.
41. Avezov, E.; Frenkel, Z.; Ehrlich, M.; Herscovics, A.; Lederkremer, G.Z. Endoplasmic reticulum (er) mannosidase i is compartmentalized and required for n-glycan trimming to man5-6glcnac2 in glycoprotein er-associated degradation. *Molecular biology of the cell* **2008**, *19*, 216-225.
42. Hosokawa, N.; You, Z.; Tremblay, L.O.; Nagata, K.; Herscovics, A. Stimulation of erad of misfolded null hong kong alpha1-antitrypsin by golgi alpha1,2-mannosidases. *Biochemical and biophysical research communications* **2007**, *362*, 626-632.
43. Ogen-Shtern, N.; Avezov, E.; Shenkman, M.; Benyair, R.; Lederkremer, G.Z. Mannosidase ia is in quality control vesicles and participates in glycoprotein targeting to erad. *Journal of molecular biology* **2016**.
44. Olivari, S.; Cali, T.; Salo, K.E.; Paganetti, P.; Ruddock, L.W.; Molinari, M. Edem1 regulates er-associated degradation by accelerating de-mannosylation of folding-defective polypeptides and by inhibiting their covalent aggregation. *Biochemical and biophysical research communications* **2006**, *349*, 1278-1284.
45. Olivari, S.; Molinari, M. Glycoprotein folding and the role of edem1, edem2 and edem3 in degradation of folding-defective glycoproteins. *FEBS letters* **2007**, *581*, 3658-3664.
46. Christianson, J.C.; Shaler, T.A.; Tyler, R.E.; Kopito, R.R. Os-9 and grp94 deliver mutant alpha1-antitrypsin to the hrd1-sel1 ubiquitin ligase complex for erad. *Nature cell biology* **2008**, *10*, 272-282.
47. Munro, S. The mrh domain suggests a shared ancestry for the mannose 6-phosphate receptors and other n-glycan-recognising proteins. *Current biology : CB* **2001**, *11*, R499-501.
48. Garza, R.M.; Sato, B.K.; Hampton, R.Y. In vitro analysis of hrd1p-mediated retrotranslocation of its multispansing membrane substrate 3-hydroxy-3-methylglutaryl (hmg)-coa reductase. *The Journal of biological chemistry* **2009**, *284*, 14710-14722.
49. Pfeffer, S.; Burbaum, L.; Unverdorben, P.; Pech, M.; Chen, Y.; Zimmermann, R.; Beckmann, R.; Forster, F. Structure of the native sec61 protein-conducting channel. *Nature communications* **2015**, *6*, 8403.
50. Kaiser, M.L.; Romisch, K. Proteasome 19s rp binding to the sec61 channel plays a key role in erad. *PloS one* **2015**, *10*, e0117260.
51. Hoelen, H.; Zaldumbide, A.; van Leeuwen, W.F.; Torfs, E.C.; Engelse, M.A.; Hassan, C.; Lebbink, R.J.; de Koning, E.J.; Resssing, M.E.; de Ru, A.H., et al. Proteasomal degradation of proinsulin requires derlin-2, hrd1 and p97. *PloS one* **2015**, *10*, e0128206.
52. Claessen, J.H.; Kundrat, L.; Ploegh, H.L. Protein quality control in the er: Balancing the ubiquitin checkbook. *Trends Cell Biol* **2012**, *22*, 22-32.
53. Hirsch, C.; Gauss, R.; Horn, S.C.; Neuber, O.; Sommer, T. The ubiquitylation machinery of the endoplasmic reticulum. *Nature* **2009**, *458*, 453-460.

54. Hampton, R.Y.; Gardner, R.G.; Rine, J. Role of 26s proteasome and hrd genes in the degradation of 3-hydroxy-3-methylglutaryl-coa reductase, an integral endoplasmic reticulum membrane protein. *Molecular biology of the cell* **1996**, *7*, 2029-2044.
55. Shibata, E.; Abbas, T.; Huang, X.; Wohlschlegel, J.A.; Dutta, A. Selective ubiquitylation of p21 and cdt1 by ubch8 and ube2g ubiquitin-conjugating enzymes via the crl4cdt2 ubiquitin ligase complex. *Molecular and cellular biology* **2011**, *31*, 3136-3145.
56. Gardner, R.G.; Swarbrick, G.M.; Bays, N.W.; Cronin, S.R.; Wilhovsky, S.; Seelig, L.; Kim, C.; Hampton, R.Y. Endoplasmic reticulum degradation requires lumen to cytosol signaling. Transmembrane control of hrd1p by hrd3p. *The Journal of cell biology* **2000**, *151*, 69-82.
57. Hosokawa, N.; Wada, I. Association of the sel1l protein transmembrane domain with hrd1 ubiquitin ligase regulates erad-l. *The FEBS journal* **2016**, *283*, 157-172.
58. Horn, S.C.; Hanna, J.; Hirsch, C.; Volkwein, C.; Schutz, A.; Heinemann, U.; Sommer, T.; Jarosch, E. Usa1 functions as a scaffold of the hrd-ubiquitin ligase. *Molecular cell* **2009**, *36*, 782-793.
59. Schulze, A.; Standera, S.; Buerger, E.; Kikkert, M.; van Voorden, S.; Wiertz, E.; Koning, F.; Kloetzel, P.M.; Seeger, M. The ubiquitin-domain protein herp forms a complex with components of the endoplasmic reticulum associated degradation pathway. *Journal of molecular biology* **2005**, *354*, 1021-1027.
60. Kreft, S.G.; Wang, L.; Hochstrasser, M. Membrane topology of the yeast endoplasmic reticulum-localized ubiquitin ligase doa10 and comparison with its human ortholog teb4 (march-vi). *The Journal of biological chemistry* **2006**, *281*, 4646-4653.
61. Wang, X.; Herr, R.A.; Rabelink, M.; Hoebe, R.C.; Wiertz, E.J.; Hansen, T.H. Ube2j2 ubiquitinates hydroxylated amino acids on er-associated degradation substrates. *The Journal of cell biology* **2009**, *187*, 655-668.
62. Burr, M.L.; Cano, F.; Svobodova, S.; Boyle, L.H.; Boname, J.M.; Lehner, P.J. Hrd1 and ube2j1 target misfolded mhc class i heavy chains for endoplasmic reticulum-associated degradation. *Proceedings of the National Academy of Sciences of the United States of America* **2011**, *108*, 2034-2039.
63. Carvalho, P.; Goder, V.; Rapoport, T.A. Distinct ubiquitin-ligase complexes define convergent pathways for the degradation of er proteins. *Cell* **2006**, *126*, 361-373.
64. Habeck, G.; Ebner, F.A.; Shimada-Kreft, H.; Kreft, S.G. The yeast erad-c ubiquitin ligase doa10 recognizes an intramembrane degron. *The Journal of cell biology* **2015**, *209*, 621.
65. Jakob, C.A.; Bodmer, D.; Spirig, U.; Battig, P.; Marcil, A.; Dignard, D.; Bergeron, J.J.; Thomas, D.Y.; Aebi, M. Htm1p, a mannosidase-like protein, is involved in glycoprotein degradation in yeast. *EMBO reports* **2001**, *2*, 423-430.
66. Burke, J.; Lipari, F.; Igldoura, S.; Herscovics, A. The *saccharomyces cerevisiae* processing alpha 1,2-mannosidase is localized in the endoplasmic reticulum, independently of known retrieval motifs. *European journal of cell biology* **1996**, *70*, 298-305.
67. Cali, T.; Galli, C.; Olivari, S.; Molinari, M. Segregation and rapid turnover of edem1 by an autophagy-like mechanism modulates standard erad and folding activities. *Biochemical and biophysical research communications* **2008**, *371*, 405-410.
68. Wu, Y.; Termine, D.J.; Swulius, M.T.; Moremen, K.W.; Sifers, R.N. Human endoplasmic reticulum mannosidase i is subject to regulated proteolysis. *The Journal of biological chemistry* **2007**, *282*, 4841-4849.
69. Bernasconi, R.; Molinari, M. Erad and erad tuning: Disposal of cargo and of erad regulators from the mammalian er. *Curr Opin Cell Biol* **2011**, *23*, 176-183.
70. Fujita, E.; Kouroku, Y.; Isoai, A.; Kumagai, H.; Misutani, A.; Matsuda, C.; Hayashi, Y.K.; Momoi, T. Two endoplasmic reticulum-associated degradation (erad) systems for the novel variant of the mutant dysferlin: Ubiquitin/proteasome erad(i) and autophagy/lysosome erad(ii). *Human molecular genetics* **2007**, *16*, 618-629.
71. Hosokawa, N.; Wada, I.; Hasegawa, K.; Yorihuzi, T.; Tremblay, L.O.; Herscovics, A.; Nagata, K. A novel er alpha-mannosidase-like protein accelerates er-associated degradation. *EMBO reports* **2001**, *2*, 415-422.

72. Olivari, S.; Galli, C.; Alanen, H.; Ruddock, L.; Molinari, M. A novel stress-induced edem variant regulating endoplasmic reticulum-associated glycoprotein degradation. *The Journal of biological chemistry* **2005**, *280*, 2424-2428.
73. Benyair, R.; Ogen-Shtern, N.; Lederkremer, G.Z. Glycan regulation of er-associated degradation through compartmentalization. *Seminars in cell & developmental biology* **2015**, *41*, 99-109.
74. Termine, D.J.; Moremen, K.W.; Sifers, R.N. The mammalian upr boosts glycoprotein erad by suppressing the proteolytic downregulation of er mannosidase i. *Journal of cell science* **2009**, *122*, 976-984.
75. Inoue, T.; Tsai, B. How viruses use the endoplasmic reticulum for entry, replication, and assembly. *Cold Spring Harbor perspectives in biology* **2013**, *5*, a013250.
76. Li, S.; Kong, L.; Yu, X. The expanding roles of endoplasmic reticulum stress in virus replication and pathogenesis. *Crit Rev Microbiol* **2015**, *41*, 150-164.
77. Chan, S.W. Unfolded protein response in hepatitis c virus infection. *Frontiers in microbiology* **2014**, *5*, 233.
78. Lazar, C.; Uta, M.; Branza-Nichita, N. Modulation of the unfolded protein response by the human hepatitis b virus. *Frontiers in microbiology* **2014**, *5*, 433.
79. Fung, T.S.; Liu, D.X. Coronavirus infection, er stress, apoptosis and innate immunity. *Frontiers in microbiology* **2014**, *5*, 296.
80. Rathore, A.P.; Ng, M.L.; Vasudevan, S.G. Differential unfolded protein response during chikungunya and sindbis virus infection: Chikv nsp4 suppresses eif2alpha phosphorylation. *Virology journal* **2013**, *10*, 36.
81. Borsa, M.; Ferreira, P.L.; Petry, A.; Ferreira, L.G.; Camargo, M.M.; Bou-Habib, D.C.; Pinto, A.R. Hiv infection and antiretroviral therapy lead to unfolded protein response activation. *Virology journal* **2015**, *12*, 77.
82. Williams, B.R. Signal integration via pkr. *Science's STKE : signal transduction knowledge environment* **2001**, *2001*, re2.
83. Komar, A.A.; Hatzoglou, M. Cellular ires-mediated translation: The war of itafs in pathophysiological states. *Cell Cycle* **2011**, *10*, 229-240.
84. Lee, D.Y.; Sugden, B. The lmp1 oncogene of ebv activates perk and the unfolded protein response to drive its own synthesis. *Blood* **2008**, *111*, 2280-2289.
85. Cheng, G.; Feng, Z.; He, B. Herpes simplex virus 1 infection activates the endoplasmic reticulum resident kinase perk and mediates eif-2alpha dephosphorylation by the gamma(1)34.5 protein. *Journal of virology* **2005**, *79*, 1379-1388.
86. Zhang, F.; Moon, A.; Childs, K.; Goodbourn, S.; Dixon, L.K. The african swine fever virus dp711 protein recruits the protein phosphatase 1 catalytic subunit to dephosphorylate eif2alpha and inhibits chop induction but is dispensable for these activities during virus infection. *Journal of virology* **2010**, *84*, 10681-10689.
87. Pavio, N.; Romano, P.R.; Graczyk, T.M.; Feinstone, S.M.; Taylor, D.R. Protein synthesis and endoplasmic reticulum stress can be modulated by the hepatitis c virus envelope protein e2 through the eukaryotic initiation factor 2alpha kinase perk. *Journal of virology* **2003**, *77*, 3578-3585.
88. Hassan, I.H.; Zhang, M.S.; Powers, L.S.; Shao, J.Q.; Baltrusaitis, J.; Rutkowski, D.T.; Legge, K.; Monick, M.M. Influenza a viral replication is blocked by inhibition of the inositol-requiring enzyme 1 (ire1) stress pathway. *The Journal of biological chemistry* **2012**, *287*, 4679-4689.
89. Galindo, I.; Hernaez, B.; Munoz-Moreno, R.; Cuesta-Geijo, M.A.; Dalmau-Mena, I.; Alonso, C. The atf6 branch of unfolded protein response and apoptosis are activated to promote african swine fever virus infection. *Cell death & disease* **2012**, *3*, e341.
90. Jheng, J.R.; Ho, J.Y.; Horng, J.T. Er stress, autophagy, and rna viruses. *Frontiers in microbiology* **2014**, *5*, 388.
91. Caselli, E.; Benedetti, S.; Gentili, V.; Grigolato, J.; Di Luca, D. Short communication: Activating transcription factor 4 (atf4) promotes hiv type 1 activation. *AIDS research and human retroviruses* **2012**, *28*, 907-912.

92. van der Wal, F.J.; Kikkert, M.; Wiertz, E. The hcmv gene products us2 and us11 target mhc class i molecules for degradation in the cytosol. *Current topics in microbiology and immunology* **2002**, *269*, 37-55.
93. Oresic, K.; Ng, C.L.; Tortorella, D. Tram1 participates in human cytomegalovirus us2- and us11-mediated dislocation of an endoplasmic reticulum membrane glycoprotein. *The Journal of biological chemistry* **2009**, *284*, 5905-5914.
94. van den Boomen, D.J.; Lehner, P.J. Identifying the erad ubiquitin e3 ligases for viral and cellular targeting of mhc class i. *Molecular immunology* **2015**, *68*, 106-111.
95. Rehm, A.; Engelsberg, A.; Tortorella, D.; Korner, I.J.; Lehmann, I.; Ploegh, H.L.; Hopken, U.E. Human cytomegalovirus gene products us2 and us11 differ in their ability to attack major histocompatibility class i heavy chains in dendritic cells. *Journal of virology* **2002**, *76*, 5043-5050.
96. Boname, J.M.; Stevenson, P.G. Mhc class i ubiquitination by a viral phd/lap finger protein. *Immunity* **2001**, *15*, 627-636.
97. Levesque, K.; Finzi, A.; Binette, J.; Cohen, E.A. Role of cd4 receptor down-regulation during hiv-1 infection. *Curr HIV Res* **2004**, *2*, 51-59.
98. Lindwasser, O.W.; Chaudhuri, R.; Bonifacino, J.S. Mechanisms of cd4 downregulation by the nef and vpu proteins of primate immunodeficiency viruses. *Curr Mol Med* **2007**, *7*, 171-184.
99. Foster, J.L.; Garcia, J.V. Hiv-1 nef: At the crossroads. *Retrovirology* **2008**, *5*, 84.
100. Margottin, F.; Bour, S.P.; Durand, H.; Selig, L.; Benichou, S.; Richard, V.; Thomas, D.; Strebel, K.; Benarous, R. A novel human wd protein, h-beta trcp, that interacts with hiv-1 vpu connects cd4 to the er degradation pathway through an f-box motif. *Molecular cell* **1998**, *1*, 565-574.
101. Strebel, K. Hiv-1 vpu - an ion channel in search of a job. *Biochimica et biophysica acta* **2014**, *1838*, 1074-1081.
102. Magadan, J.G.; Perez-Victoria, F.J.; Sougrat, R.; Ye, Y.; Strebel, K.; Bonifacino, J.S. Multilayered mechanism of cd4 downregulation by hiv-1 vpu involving distinct er retention and erad targeting steps. *PLoS pathogens* **2010**, *6*, e1000869.
103. Byun, H.; Halani, N.; Mertz, J.A.; Ali, A.F.; Lozano, M.M.; Dudley, J.P. Retroviral rem protein requires processing by signal peptidase and retrotranslocation for nuclear function. *Proceedings of the National Academy of Sciences of the United States of America* **2010**, *107*, 12287-12292.
104. Surjit, M.; Jameel, S.; Lal, S.K. Cytoplasmic localization of the orf2 protein of hepatitis e virus is dependent on its ability to undergo retrotranslocation from the endoplasmic reticulum. *Journal of virology* **2007**, *81*, 3339-3345.
105. Neu, U.; Stehle, T.; Atwood, W.J. The polyomaviridae: Contributions of virus structure to our understanding of virus receptors and infectious entry. *Virology* **2009**, *384*, 389-399.
106. Lilley, B.N.; Gilbert, J.M.; Ploegh, H.L.; Benjamin, T.L. Murine polyomavirus requires the endoplasmic reticulum protein derlin-2 to initiate infection. *Journal of virology* **2006**, *80*, 8739-8744.
107. Schelhaas, M.; Malmstrom, J.; Pelkmans, L.; Haugstetter, J.; Ellgaard, L.; Grunewald, K.; Helenius, A. Simian virus 40 depends on er protein folding and quality control factors for entry into host cells. *Cell* **2007**, *131*, 516-529.
108. Reggiori, F.; Monastyrska, I.; Verheije, M.H.; Cali, T.; Ulasli, M.; Bianchi, S.; Bernasconi, R.; de Haan, C.A.; Molinari, M. Coronaviruses hijack the lc3-i-positive edosomes, er-derived vesicles exporting short-lived erad regulators, for replication. *Cell host & microbe* **2010**, *7*, 500-508.
109. Noack, J.; Bernasconi, R.; Molinari, M. How viruses hijack the erad tuning machinery. *Journal of virology* **2014**, *88*, 10272-10275.
110. Zhu, P.; Chertova, E.; Bess, J., Jr.; Lifson, J.D.; Arthur, L.O.; Liu, J.; Taylor, K.A.; Roux, K.H. Electron tomography analysis of envelope glycoprotein trimers on hiv and simian immunodeficiency virus virions. *Proceedings of the National Academy of Sciences of the United States of America* **2003**, *100*, 15812-15817.
111. Zhu, P.; Liu, J.; Bess, J., Jr.; Chertova, E.; Lifson, J.D.; Grise, H.; Ofek, G.A.; Taylor, K.A.; Roux, K.H. Distribution and three-dimensional structure of aids virus envelope spikes. *Nature* **2006**, *441*, 847-852.

112. Forster, F.; Medalia, O.; Zauberman, N.; Baumeister, W.; Fass, D. Retrovirus envelope protein complex structure in situ studied by cryo-electron tomography. *Proceedings of the National Academy of Sciences of the United States of America* **2005**, *102*, 4729-4734.
113. Grunewald, K.; Desai, P.; Winkler, D.C.; Heymann, J.B.; Belnap, D.M.; Baumeister, W.; Steven, A.C. Three-dimensional structure of herpes simplex virus from cryo-electron tomography. *Science* **2003**, *302*, 1396-1398.
114. Loney, C.; Mottet-Osman, G.; Roux, L.; Bhella, D. Paramyxovirus ultrastructure and genome packaging: Cryo-electron tomography of sendai virus. *Journal of virology* **2009**, *83*, 8191-8197.
115. Yamaguchi, M.; Danev, R.; Nishiyama, K.; Sugawara, K.; Nagayama, K. Zernike phase contrast electron microscopy of ice-embedded influenza a virus. *J Struct Biol* **2008**, *162*, 271-276.
116. Schiller, J.; Chackerian, B. Why hiv virions have low numbers of envelope spikes: Implications for vaccine development. *PLoS pathogens* **2014**, *10*, e1004254.
117. Fennie, C.; Lasky, L.A. Model for intracellular folding of the human immunodeficiency virus type 1 gp120. *Journal of virology* **1989**, *63*, 639-646.
118. Hallenberger, S.; Tucker, S.P.; Owens, R.J.; Bernstein, H.B.; Compans, R.W. Secretion of a truncated form of the human immunodeficiency virus type 1 envelope glycoprotein. *Virology* **1993**, *193*, 510-514.
119. Willey, R.L.; Bonifacino, J.S.; Potts, B.J.; Martin, M.A.; Klausner, R.D. Biosynthesis, cleavage, and degradation of the human immunodeficiency virus 1 envelope glycoprotein gp160. *Proceedings of the National Academy of Sciences of the United States of America* **1988**, *85*, 9580-9584.
120. Zhou, T.; Han, Y.; Dang, Y.; Wang, X.; Zheng, Y.H. A novel hiv-1 restriction factor that is biologically distinct from apobec3 cytidine deaminases in a human t cell line cem.Nkr. *Retrovirology* **2009**, *6*, 31.
121. Zhou, T.; Dang, Y.; Zheng, Y.H. The mitochondrial translocator protein, tspo, inhibits hiv-1 envelope glycoprotein biosynthesis via the endoplasmic reticulum-associated protein degradation pathway. *Journal of virology* **2014**, *88*, 3474-3484.
122. Zamzami, N.; Kroemer, G. The mitochondrion in apoptosis: How pandora's box opens. *Nat Rev Mol Cell Biol* **2001**, *2*, 67-71.
123. Gatliff, J.; East, D.; Crosby, J.; Abeti, R.; Harvey, R.; Craigen, W.; Parker, P.; Campanella, M. Tspo interacts with vdac1 and triggers a ros-mediated inhibition of mitochondrial quality control. *Autophagy* **2014**, *10*, 2279-2296.
124. Wu, Y.; Swulius, M.T.; Moremen, K.W.; Sifers, R.N. Elucidation of the molecular logic by which misfolded alpha 1-antitrypsin is preferentially selected for degradation. *Proceedings of the National Academy of Sciences of the United States of America* **2003**, *100*, 8229-8234.
125. Zhou, T.; Frabutt, D.A.; Moremen, K.W.; Zheng, Y.H. Ermani (endoplasmic reticulum class i alpha-mannosidase) is required for hiv-1 envelope glycoprotein degradation via endoplasmic reticulum-associated protein degradation pathway. *The Journal of biological chemistry* **2015**, *290*, 22184-22192.
126. Gonzalez, D.S.; Karaveg, K.; Vandersall-Nairn, A.S.; Lal, A.; Moremen, K.W. Identification, expression, and characterization of a cDNA encoding human endoplasmic reticulum mannosidase i, the enzyme that catalyzes the first mannose trimming step in mammalian asn-linked oligosaccharide biosynthesis. *Journal of Biological Chemistry* **1999**, *274*, 21375-21386.
127. Tremblay, L.O.; Herscovics, A. Cloning and expression of a specific human alpha 1,2-mannosidase that trims man(9)glcnac(2) to man(8)glcnac(2) isomer b during n-glycan biosynthesis. *Glycobiology* **1999**, *9*, 1073-1078.
128. Vallee, F.; Karaveg, K.; Herscovics, A.; Moremen, K.W.; Howell, P.L. Structural basis for catalysis and inhibition of n-glycan processing class i alpha 1,2-mannosidases. *Journal of Biological Chemistry* **2000**, *275*, 41287-41298.
129. Lipari, F.; Herscovics, A. Role of the cysteine residues in the alpha1,2-mannosidase involved in n-glycan biosynthesis in *saccharomyces cerevisiae*. The conserved cys340 and cys385 residues form an essential disulfide bond. *The Journal of biological chemistry* **1996**, *271*, 27615-27622.

130. Rafiq, M.A.; Kuss, A.W.; Puettmann, L.; Noor, A.; Ramiah, A.; Ali, G.; Hu, H.; Kerio, N.A.; Xiang, Y.; Garshasbi, M., *et al.* Mutations in the alpha 1,2-mannosidase gene, *man1b1*, cause autosomal-recessive intellectual disability. *American journal of human genetics* **2011**, *89*, 176-182.
131. Gupta, S.; Fahiminiya, S.; Wang, T.; Dempsey Nunez, L.; Rosenblatt, D.S.; Gibson, W.T.; Gilfix, B.; Bergeron, J.J.; Jerome-Majewska, L.A. Somatic overgrowth associated with homozygous mutations in both *man1b1* and *sec23a*. *Cold Spring Harbor molecular case studies* **2016**, *2*, a000737.
132. Iannotti, M.J.; Figard, L.; Sokac, A.M.; Sifers, R.N. A golgi-localized mannosidase (*man1b1*) plays a non-enzymatic gatekeeper role in protein biosynthetic quality control. *The Journal of biological chemistry* **2014**, *289*, 11844-11858.
133. Zhang, X.; Zhou, T.; Frabutt, D.A.; Zheng, Y.H. Hiv-1 vpr increases env expression by preventing env from endoplasmic reticulum-associated protein degradation (erad). *Virology* **2016**, *496*, 194-202.
134. Lazar, C.; Macovei, A.; Petrescu, S.; Branza-Nichita, N. Activation of erad pathway by human hepatitis b virus modulates viral and subviral particle production. *PloS one* **2012**, *7*, e34169.
135. Saeed, M.; Suzuki, R.; Watanabe, N.; Masaki, T.; Tomonaga, M.; Muhammad, A.; Kato, T.; Matsuura, Y.; Watanabe, H.; Wakita, T., *et al.* Role of the endoplasmic reticulum-associated degradation (erad) pathway in degradation of hepatitis c virus envelope proteins and production of virus particles. *The Journal of biological chemistry* **2011**, *286*, 37264-37273.
136. McPherson, S.; Powell, E.E.; Barrie, H.D.; Clouston, A.D.; McGuckin, M.; Jonsson, J.R. No evidence of the unfolded protein response in patients with chronic hepatitis c virus infection. *Journal of gastroenterology and hepatology* **2011**, *26*, 319-327.
137. Britt, W.J.; Mach, M. Human cytomegalovirus glycoproteins. *Intervirology* **1996**, *39*, 401-412.
138. Compton, T.; Nepomuceno, R.R.; Nowlin, D.M. Human cytomegalovirus penetrates host cells by ph-independent fusion at the cell surface. *Virology* **1992**, *191*, 387-395.
139. Zhou, M.; Lanchy, J.M.; Ryckman, B.J. Human cytomegalovirus gh/gl/go promotes the fusion step of entry into all cell types, whereas gh/gl/ul128-131 broadens virus tropism through a distinct mechanism. *Journal of virology* **2015**, *89*, 8999-9009.
140. Hutchinson, L.; Browne, H.; Wargent, V.; Davis-Poynter, N.; Primorac, S.; Goldsmith, K.; Minson, A.C.; Johnson, D.C. A novel herpes simplex virus glycoprotein, gl, forms a complex with glycoprotein h (gh) and affects normal folding and surface expression of gh. *Journal of virology* **1992**, *66*, 2240-2250.
141. Gardner, T.J.; Hernandez, R.E.; Noriega, V.M.; Tortorella, D. Human cytomegalovirus gh stability and trafficking are regulated by er-associated degradation and transmembrane architecture. *Scientific reports* **2016**, *6*, 23692.
142. Tamura, T.; Cormier, J.H.; Hebert, D.N. Characterization of early edem1 protein maturation events and their functional implications. *The Journal of biological chemistry* **2011**, *286*, 24906-24915.
143. Molinari, M.; Calanca, V.; Galli, C.; Lucca, P.; Paganetti, P. Role of edem in the release of misfolded glycoproteins from the calnexin cycle. *Science* **2003**, *299*, 1397-1400.
144. Cormier, J.H.; Tamura, T.; Sunryd, J.C.; Hebert, D.N. Edem1 recognition and delivery of misfolded proteins to the sel1l-containing erad complex. *Molecular cell* **2009**, *34*, 627-633.
145. Pan, S.; Wang, S.; Utama, B.; Huang, L.; Blok, N.; Estes, M.K.; Moremen, K.W.; Sifers, R.N. Golgi localization of ermani defines spatial separation of the mammalian glycoprotein quality control system. *Molecular biology of the cell* **2011**, *22*, 2810-2822.

