

Supplementary Information

- Table S1.** ^{13}C NMR chemical shift of smenopyrone (**1**) in CDCl_3 compared with those of the eight diastereomers **4-11** of maurenone.
- Figure S1.** Difference in ^{13}C NMR chemical shift ($\Delta\delta$) between corresponding atoms of smenopyrone (**1**) and the eight stereoisomers **4-11** of the model compound maurenone.
- Figure S2.** UV and ECD spectra of smenopyrone (**1**).
- Figure S3.** Positive ion mode high-resolution ESI mass spectrum of smenopyrone (**1**).
- Figure S4.** Positive ion mode high-resolution ESI MS/MS spectrum (parent ion at m/z 419.28) and fragmentation of smenopyrone (**1**).
- Figure S5.** ^1H -NMR spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).
- Figure S6.** COSY spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).
- Figure S7.** NOESY spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).
- Figure S8.** HSQC spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).
- Figure S9.** HMBC spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).
- Figure S10.** Expansion of methyl region of the HMBC spectrum of smenopyrone (**1**) (700 MHz, CD_3OD).




Table S1. ^{13}C NMR chemical shifts of smenopyrone (**1**) in CDCl_3 compared with those of the eight diastereomers of maurenone, **4-11**.

Pos.		δ_{C} (CDCl_3)								
		1	4	5	6	7	8	9	10	11
9	CH	75.1	75.5	75.4	75.1	74.8	75.3	75.3	74.8	75.2
10	CH	38.6	41.1	41.4	41.6	41.5	41.2	41.2	41.5	41.8
11	C	172.4	172.4	172.3	173.0	173.1	172.6	172.3	173.2	173.2
12	C	108.5	109.9	109.7	108.6	108.5	109.8	109.8	108.6	108.7
13	C	195.4	195.6	195.6	195.7	195.7	195.5	195.5	195.7	195.7
14	CH	40.6	40.9	40.6	40.6	40.7	40.5	40.6	40.7	40.5
15	CH	87.4	87.8	87.0	87.0	87.4	84.5	84.5	84.7	84.2
16	CH	35.1	35.4	35.1	35.1	35.2	35.6	35.6	35.5	35.6
17	CH_2	22.1	22.0	22.2	21.9	21.9	26.8	26.8	26.5	26.5
18	CH_3	12.2	10.1	11.7	11.7	11.9	9.9	11.9	10.2	11.8
22	CH_3	13.9	14.7	14.7	13.3	12.9	14.6	14.6	12.9	13.4
23	CH_3	9.3	9.4	9.4	9.3	9.2	9.4	9.4	9.2	9.3
24	CH_3	10.6	10.0	11.0	10.6	10.1	9.9	9.9	9.5	9.7
25	CH_3	16.5	16.2	16.3	16.1	16.1	12.3	12.3	12.4	12.3
$\Sigma \Delta\delta $			6.6	4.0	3.1	3.7	17.8	15.6	17.1	15.5




Figure S1. Difference in ^{13}C NMR chemical shift ($\Delta\delta$) between corresponding atoms of smenopyrone (**1**) and the eight stereoisomers **4-11** of the model compound maurenone. The sum of absolute values of $\Delta\delta$ ($\Sigma|\Delta\delta|$) was used to evaluate the overall fit between **1** and **4-11**.





Figure S2. UV (top) and ECD (bottom) spectra (MeOH) of smenopyrone (**1**).




Figure S3. Positive ion mode high-resolution ESI mass spectrum of smenopyrone (**1**).




Figure S4. Positive ion mode high-resolution ESI MS/MS spectrum (parent ion at m/z 419.28) and fragmentation of smenopyrone (**1**).




Figure S5. ¹H-NMR spectrum of smenopyrone (**1**) (700 MHz, CD₃OD)




Figure S6. COSY spectrum of smenopyrone (1) (700 MHz, CD₃OD)




Figure S7. NOESY spectrum of smenopyrone (1) (700 MHz, CD₃OD)




Figure S8. HSQC spectrum of smenopyrone (**1**) (700 MHz, CD₃OD)




Figure S9. HMBC spectrum of smenopyrone (1) (700 MHz, CD₃OD)




Figure S10. Expansion of methyl region of the HMBC spectrum of smenopyrone (1) (700 MHz, CD₃OD)