

Preferred title: Using fish models to investigate the links between microbiome and social behaviour: the next step for translational microbiome research?

Running title: Links between microbiomes and sociality in fish

Marta Candeias Soares¹, Jo Cable², Monica Gomes Lima-Maximino³, Caio Maximino⁴, Raquel Xavier^{1*}

¹ CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos da Universidade do Porto, R. Padre Armando Quintas, 4485-661 Vairão, Portugal

² School of Biosciences, Cardiff University, Cardiff, CF10 3AX, U.K.

³ DMCF, Departamento de Morfologia e Ciências Fisiológicas, Universidade do Estado do Pará, Marabá, Brazil

⁴ IESB, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III, Marabá, Brazil

Abstract

Recent research has revealed surprisingly important connections between animals' microbiome and social behaviour. Social interactions can affect the composition and function of the microbiome; conversely, the microbiome affects social communication by influencing the hosts' central nervous system and peripheral chemical communication. These discoveries set the stage for novel research focusing on the evolution and physiology of animal social behaviour in relation to microbial transmission strategies. Here, we discuss the emerging roles of teleost fish models and their potential for advancing research fields linked to sociality and microbial regulation. We argue that fish models, such as the zebrafish, sticklebacks, guppies and cleaner-client dyads, will provide valuable insights into the roles of microbiome in shaping social behaviour and vice versa, while also being of direct relevance to the food and ornamental fish trades. The diversity of fish behaviour warrants more interdisciplinary research, including microbiome studies, which should have a strong ecological (field-derived) approach, together with laboratory-based cognitive and neurobiological experimentation. The implications of such integrated approaches may be of translational relevance, opening new avenues for future investigation using fish models.

Keywords: behaviour/sociality; cleaner fish; gut-brain axis; poecilids; sticklebacks; zebrafish

Table of contents:

1. The gut-brain axis and its importance for health, stress physiology and behaviour
2. From human to fish: widening microbiome-based behavioural neuroscience
 - 2.1. Fish microbiome: what we know
 - 2.2. A fishy absence: the gut-brain axis in fish
3. The usefulness of fish models to study microbiome-behaviour interactions
 - 3.1. Zebrafish
 - 3.2. Poeciliids
 - 3.3. Sticklebacks
 - 3.4. Cleaner fish
4. Concluding remarks

1. The gut-brain axis and its importance for health, stress physiology and behaviour

The microbiota-gut-brain axis (MGB axis) has a pivotal role in host health, stress physiology and behaviour (Foster, Rinaman, & Cryan, 2017). The interaction between the gut microbiome and the brain is now well established, at least in mammals, and involves multiple pathways, including immune signalling or neurotransmission through bacteria-derived metabolites (such as serotonin, histamine and catecholamine regulators; Sandhu et al., 2017). In humans, neonatal microbial colonization of the gut is influenced by hospital environment, antibiotic use, birth delivery mode, breastfeeding vs bottle feeding, and early life stress (e.g. see Yang et al., 2016). Concomitantly with early establishment of the gut microbiome are key neurodevelopmental landmarks

critical for establishing normal motor, cognitive and emotional function (Cong et al., 2016; Borre et al., 2014). The interaction between the microbiome, neurodevelopment and the central nervous system, i.e. the gut-brain axis, can be influenced by disease and stress, which cause an imbalance in the gut microbiome (DeGruttola, Low, Mizoguchi, & Mizoguchi, 2016; García-Ródenas et al., 2006). In healthy mammals, bacteria from the phyla Bacteroidetes, Firmicutes, Actinobacteria and Proteobacteria dominate the gut microbiome (e.g. Lozupone, Stombaugh, Gordon, Jansson, & Knight, 2012; Groussin et al., 2017; Yang et al., 2016). Disease-induced imbalances often result in increased intestinal permeability, due to the decrease in Firmicutes that are able to synthesize essential short-chain fatty acids (e.g. *Faecalibacterium* and *Eubacterium*), namely the butyrate, known to contribute to the maintenance of the gutbarrier (Rivière, Selak, Lantin, Leroy & De Vuyst, 2016; DeGruttola et al., 2016). Likewise, imbalances induced by stress are often associated with increased corticotropin-releasing hormone, which also causes increased intestinal permeability (e.g. Vanuytsel et al., 2014). This means that in diseased and/or stressed hosts there is an increased ability of gut bacteria and their metabolites to translocate through the intestinal mucosa into the bloodstream, accessing immune cells and neurons in the enteric nervous system (Gareau, Silva & Perdue, 2008; Teitelbaum, Gareau, Jury, Yang & Perdue, 2008; Tognini, 2017). Indeed, several psychiatric disorders related to stress have been linked with increased intestinal permeability (e.g. chronic depression and anxiety, Kelly et al., 2015; Kelly et al., 2016; van de Wouw et al., 2018) and in some cases diet supplementation with short-chain fatty acids was shown to ameliorate symptoms (e.g. van de Wouw et al., 2018). Depressive episodes, related to dysregulations of the hypothalamus-pituitary-adrenal (HPA) axis (Barden, 2004), also appear to have a close relationship with gut microbiome colonization. For instance, germ-free (GF) rats, that lack commensal gut bacteria – and, consequently, have a poorly developed immune system – exhibit elevated levels of plasma adrenocorticotrophic hormone and corticosterone (e.g. stronger HPA response) when subjected to restraint stress or to novel environments (Sudo et al., 2004; Clarke et al., 2013). In fact, neonatal stress leads to long-term changes in the diversity and

composition of the gut microbiome, resulting in a reduction of beneficial bacteria (e.g. *Lactobacillus*, see Bailey and Cole, 1999), which could contribute to long-term changes in stress reactivity and in the stress-related behaviour observed in these rats. Treatment with probiotics (mainly *Lactobacillus* spp.) through the initial stress stage normalizes basal corticosterone levels, which are elevated after maternal separation (Gareau et al., 2008). Likewise, pre-treatment of rats with *Lactobacillus farciminis* reduces the intestinal permeability that results from restraint stress, as well as the associated hyperactivity of the HPA axis (Ait-Belgnaoui et al., 2012) and potential stimulation of the MGB axis. In addition to these findings relating microbiome and HPA axis function, there is direct evidence that the microbiome can influence central nervous system activity through the activation of stress-related neurons; for example, orally administered bacteria induce c-Fos expression in afferent vagal neurons (Lyte, Li, Opitz, Gaykema, & Goehler, 2006) and in paraventricular hypothalamic neurons in GF rats (Sudo et al., 2004) in the absence of systemic immune responses, suggesting that commensal and pathogenic bacteria in the gastrointestinal tract release signals to the central stress circuits. Moreover, electrophysiological experiments demonstrated that vagal nerve endings in the gut are less excitable in GF rodents (McVey Neufeld, Mao, Bienenstock, Foster, & Kunze, 2013), or more excitable after probiotic administration (Kunze et al., 2009), suggesting changes in enteric neuronal responses due to variations in commensal microbiome.

The reciprocal relationship between the microbiome and social life of hosts arose as a surprising topic, emphasizing the impact of microbial communities on animal social communication and signaling. Microbial transfer between socially interacting partners is common and was recently reviewed by Archie and Tung (2015) and Vuong et al. (2017). Human families, for example, not only share a similar skin microbiome (e.g. Song et al., 2013; Lax et al., 2014), but pet dogs can facilitate homogenization of the microbiome between family members (Song et al., 2013). Besides the skin microbiome, oral and gut microbiomes were also more similar within a human

family, and especially between sexually interacting partners (Song et al., 2013). Moreover, microbial transfer between interacting individuals can occur even when the time frame for direct contact is reduced, for example the skin microbiome of players from opposing teams after a roller derby tournament revealed a significant homogenization (Meadow, Bateman, Herkert, O'Connor, & Green, 2013). Analyses of the gut microbiome in non-human primates, the Verreaux's sifaka lemurs, chimpanzees and baboons, showed that the best predictors for gut microbial composition were social group and physical interactions, in comparison to diet, kinship and habitat overlap (Moeller et al., 2016; Perofsky, Lewis, Abondano, Di Fiore, & Meyers, 2017; Tung et al., 2015). In birds, body contact also seems to be a good predictor for microbial diversity with higher diversities being positively associated with frequency of interactions in barn swallow females (Levin et al., 2016). In males this pattern was reversed, since frequent contact with other competing males caused significant stress, which, in turn, may have prompted a reduction in gut microbial diversity (Levin et al., 2016). Bacterial transmission can also be mediated through mating in kittiwakes, with White et al. (2010) demonstrating that the cloacal microbiome between mating partners is similar but can diverge over time if cloacas are covered thus preventing direct contact. (2017;). Disentangling the links between microbiome and fish health is a hot topic, particularly given the growth of the aquaculture industry. In this respect, the effects of pathogens, antibiotics and probiotics on microbiome composition has been subject of several studies showing that significant changes take place (e.g. Carlson, Leonard, Hyde, Petrosino, & Primm, 2017; Gonçalves & Gallardo-Escarate, 2017; Hennersdorf et al., 2016; Llewellyn et al., 2017; Reid et al., 2017; Li et al., 2016).

The few studies that have focused on the effects of stress on fish microbiomes found a significant impact on skin microbial diversities. Aquaculture conditions, such as reduced oxygen, overcrowding and frequent fish handling (but see Box 2), can significantly reduce skin bacteria in farmed Atlantic salmon and Brook charr, namely the genera *Methylobacterium*, *Sphingomonas* and *Propionibacterium* that have antibiotic properties against fish pathogens (Boutin,

Bernatchez, Audet, & Derôme, 2013; Minniti et al., 2017,). Horizontal microbial transfer between fish in close physical contact has also been reported in the gut microbiome of fish, since differences associated with zebrafish genotypes were diluted when fish were co-housed (Burns et al., 2017).

2.2. A fishy absence: the gut-brain axis in fish

As our understanding of fish host-microbiota interactions increases, so it becomes obvious that more research is necessary, particularly on the interdependence between the microbiota and fish behavioural responses. This contrasts with the rise in popularity of fish behavioural neuroscience, where several fish species are widely used as models to study social cognition and behaviour (Oliveira, 2013; Soares, Cardoso, Carvalho, & Maximino, 2018). While there is no doubt a significant bias towards using birds and mammals to study animal social behaviour and cognition, fish possess much more than just a predictable mixture of unlearned predispositions (Brown, 2015). Fish display sophisticated behaviours and complex social networks, such as Machiavellian strategies of deception and reconciliation (Bshary, Wickler, & Fricke, 2002), the ability to monitor the behaviour of others and their own reputation, to cooperate and to adjust their investment levels in accordance to context and even to use social tools to achieve direct goals (Brown & Laland, 2003; Soares, 2017; Soares et al., 2018). This complexity mirrors some homologous components of fish' neural representations with that of higher vertebrates (Oliveira, 2013). However, much remains to be discovered about the similarities and dissimilarities of behaviour and brain processes between fish and mammals.

3. The usefulness of fish models to study microbiome-behaviour interactions

Given that fish display a complex array of social behaviours, from solitary individuals to complex group behaviour based on reciprocity, we should expect a variety of effects on (and of) the microbiome. An emergent example is the use of social network analysis to describe the structure of

animal societies at different organizational levels (i.e. individual, dyad, group, community, population, metapopulation and species). These methodologies, however, have their limitations (reviewed in Wilson, Croft & Krause, 2014). For example, direct observations of social interactions can be challenging in aquatic environments, and these studies often rely on tagged individuals, and having to control for tagging effects (Wilson et al., 2014). For this reason, the application of social network analysis has so far been restricted to: a) small and easy monitored freshwater teleost species (e.g. Croft et al., 2011; Croft, Krause & James, 2004;); b) to larger species that can be marked or recognized through individual-specific colour patterns and with behaviour that can be observed from a distance (e.g. elasmobranchs, Guttridge et al. 2011; Mourier, Vercelloni & Plane, 2012); or c) to those that may bear sophisticated acoustic or radio tags (e.g. Guttridge, Gruber, Krause & Sims, 2010). Currently, the challenge is to extend this methodology to other social fish systems, in the lab and in wild conditions, but also to include a stronger integrative approach, with, for instance, the addition of physiological and microbial measures, enabling the evaluation of social transmission combined with individuals' neurochemical balance and welfare (e.g. disease spread). Another key absence is the lack of knowledge regarding the gut-brain functioning in fish species and the potential impact on social behaviour. We argue that fish models, such as the zebrafish, sticklebacks, guppies, cleaner-client dyads (Box 1), can help provide valuable insights into the roles of microbiome in shaping social behaviour and vice versa.

3.1 Zebrafish

The zebrafish is a key model organism in developmental biology and genetics and is increasingly used in behavioural neuroscience (Kalueff, Stewart, & Gerlai, 2014; Norton & Bally-Cuif, 2010; Stewart, Braubach, Spitsbergen, Gerlai & Kalueff, 2014). A wide array of tools, from behavioural assays to neurogenetic and neurogenomic tools (Gerlai, 2014; Rinkwitz, Mourrain, & Becker, 2011;) have been developed for zebrafish, including specific assays for anxiety- and fear-like behaviour, as well as for stress responses (Clark, Boczek, & Ekker, 2011; Jesuthasan & Mathuru,

2008; Maximino et al., 2010; Stewart et al., 2014). So far, the single study focusing on the role of sociality in horizontal bacterial transfer in piscine hosts was conducted in zebrafish, with results supporting a role of co-housing in microbial composition of the gut (Burns et al., 2017). Furthermore, there is mounting evidence for a species-specific “core” gut microbiota in zebrafish, as animals reared for generations in the laboratory show similar components in relation to animals captured in the wild (Roeslers et al., 2011).

As described for other vertebrates, zebrafish exhibit ontogenetic shifts in microbiome composition (Falcinelli et al., 2015; Roeslers et al., 2011; Hall, Tolonen, & Xavier, 2017; Shin, Whon, & Bae, 2015). Recently, the effects of probiotics on zebrafish, namely on their behaviour, have been assessed. Adult zebrafish treated with the probiotic *Lactobacillus rhamnosus* IMC 501 for 28 days showed increases in the abundance of taxa belonging to Phylum Firmicutes in the gut (Borrelli et al., 2016). Treatment of zebrafish larvae with *L. rhamnosus* IMC 501 from 0 to 8 days post-fertilization decreased the relative abundance of Proteobacteria and increased Firmicutes. Borrelli et al. (2016) found similar increases in Firmicutes after 28 days of treatment in adult animals. Importantly, adult zebrafish treated with *L. rhamnosus* IMC 501 displayed less shoaling and increased brain-derived neurotrophic factor (bDNF), mRNA in the brain, with upregulation of genes involved in the serotonergic system in the brain (tph1a, tph1b, tph2, htr1aa, slc6a4a, and mao) and gut (tph1a) (Borrelli et al., 2016). The reductions in shoaling could represent an anxiolytic-like effect; however, Schneider et al. (2016) did not find alterations in anxiety-like behaviour after 15 days of treatment with *L. rhamnosus* GG – although this probiotic prevented the anxiolytic-like effect of chronic ethanol treatment. These discrepancies could also be due to differences in treatment duration (28 vs. 15 days). Consistent with this hypothesis, Davis et al. (2016) found that treatment with *Lactobacillus plantarum* for 30 days decreased anxiety-like behaviour in the novel tank test, an effect that was accompanied by increased slc6a4a (serotonin transporter) expression in the brain (but no alterations in the GABAergic, neuropeptide Y, or oxytocin-like

pathways). *L. plantarum* did not alter gut microbiota diversity, but Mycoplasmataceae, Stenotrophomonas, Catenibacterium, unclassified Lactobacillaceae, and Achromobacter were enriched in Lactobacillus-treated animals. Interestingly, when *L. plantarum*-treated animals were exposed to a chronic unpredictable stress paradigm, serum cortisol and leukogram alterations were not altered, but the probiotic protected against the gut dysbiosis induced by stress. Interpreting these findings is hard, as the authors did not report any behavioural alterations in stressed animals. For now results based on probiotic treatment should be treated with caution, but suggest that, at least in zebrafish, the microbiome is associated with defensive (anxiety-like) behaviour, and therefore could indirectly mediate sociality by altering the anti-predatory component of social behaviour or by decreasing the neophobia associated with social novelty.

3.2. Poeciliids

The guppy (*Poecilia reticulata*) and indeed a range of other poeciliid fish are widely studied tropical fish species (Evans, Pilastro, & Schlupp, 2011), renowned models of ecology and evolution. The Trinidadian guppy system in particular has been used to study the complex role of gene flow on adaptation and fitness in a natural setting. Naturally isolated populations provide natural (replicated) study systems that have been used intensively to assess the impact of different environmental factors, and predation and parasitism pressure on host morphology, genotype and behaviour (e.g. Endler, 1995; Magurran, 2005; Reznick, Shaw, Rodd, & Shaw, 1977; Reznick & Ricklefs, 2009). While these wild guppies have been used to study micro-evolutionary speciation in field and laboratory studies, the ease with which they can be maintained and bred in captivity has meant they are also one of the most common fish in the world-wide ornamental trade. Their popularity relates to the availability of many different domestic colour and ornament varieties and hybrids (locally known as rainbowfish or millionsfish to reflect their colour and abundance). As the trade continues to expand, new strains are being selectively bred as these small viviparous

fish have a fast generation time and are relatively tolerant of poor water quality. This has provided the basis for numerous studies on the genetics, immunology, particularly the MHC-genes (e.g. Phillips et al., 2018) as well as the ecology and behaviour (Croft et al., 2011) of this model species.

Using the 'natural Trinidad laboratory' of replicated distinct up- and down-stream populations across four rivers, Sullam et al. (2015) were the first to assess the guppy microbiome, and they did so in relation to host adaptation. They found clear microbiome differences in the guts of wild-caught and laboratory bred fish, and from field specimens revealed that gut microbiome communities varied temporally, across streams and between ecotypes in a stream-specific manner. But this study also highlighted the myriad of other biotic and abiotic factors that could be driving microbiome divergence (Sullam et al., 2015).

Studies assessing the social behaviour of guppies have revealed how different life-history trajectories may shape sociality (e.g., Dugatkin, 1988; Farr, 1975) and the importance of cooperation in the establishment of these social groups (Croft et al., 2004; Dugatkin & Alfieri, 1991; Wilson et al, 2014). Social network analyses have shown that guppies occupy consistent social positions within a network (Krause et al., 2016) but that parasites can influence the guppy social networks (Croft et al., 2011). If the gut microbiome also influence's guppy behaviour and there is a higher probability of microbial transfer between conspecifics, we predict, based on similar studies of zebrafish, that social interactions and prosocial behaviour will be associated with microbiome composition.

3.3 Sticklebacks

The three-spined stickleback (*Gasterosteus aculeatus*) is the temperate equivalent of the guppy, being a small freshwater fish that is robust and easy to maintain and breed in large numbers in both the laboratory and field. But this fish is not just a leading model in ecology and evolutionary

research; following publication of its genome (Jones et al., 2012) it arguably has reached super-model status. Following glacial retreat in the north-ern hemisphere 12,000 years ago, repeated evolution from ancestral oceanic to derived freshwater forms resulted in many beneficial genetic variants surviving in thousands of distinct natural populations (Bell & Foster, 1994). Unlike laboratory genetic screening, which identifies mutants causing abnormal development, physiology or behaviour, these natural variants highlight biological innovations. Consequently, a large community of researchers utilises stickleback variants to understand adaptive speciation, behaviour, development, immunology (e.g. Cresko, McGuigan, Phillips, & Postlethwait, 2007; Huntingford, Ostlund-Nilsson, & Mayer, 2006; Scharsack, Kalbe, Harrod, & Rauch, 2007), and a range of projects are in progress assessing the microbiome of sticklebacks across a range of different environments. To date, such research has been largely confined to correlative approaches: QTL mapping, divergence mapping, and association studies.

Stickleback microbiome research has only just got off the starting block, but we already know that diet has a sex-specific effect on the gut microbiota (Bolnick et al., 2014b), and perhaps surprisingly individuals with more generalised diets have less di-verse microbiota than dietary specialists (Bolnick et al., 2014c). Of key importance is host genotype compared to transient environmental factors (Smith, Snowberg, Caporaso, Knight, & Bolnick, 2015), but systems-based approaches in the future are likely to reveal the true complexity of these interactions. Oceanic (ancestral) sticklebacks mount a strong immune response to residential microbiota compared to freshwater variants as revealed by the use of the stickleback as a newly developed gnotobiotic vertebrate model (Milligan-Myhre et al., 2016). Alongside zebrafish, sticklebacks could provide a powerful tool for mapping the natural genetic basis of the variation in immune response to microbes (Small, Milligan-Myhre, Bassham, Guillemin, & Cresko, 2017), and thus provide important clues regarding the role of intrinsic factors in microbiome composition. In the next few years, improved genome annotation will facilitate identification of candidate genes and experimental

gene manipulation will provide the tools to validate suspected candidate genes. Importantly, recent research has already shown that specific Major Histocompatibility Complex alleles and diversity are related to the composition and diversity of gut bacteria in sticklebacks (Bolnick et al., 2014a), as has previously been discerned for macroparasites (Matthews, Harmon, M'Gonigle, Marchinko, & Schaschl, 2010; Eizaguirre, Lenz, Kalbe, & Milinski, 2012).

3.4. Cleaner fish

Cleaning mutualisms are interspecific associations where cleaners remove ectoparasites and infected tissue from the body surface, mouth and gill chambers of their client fish (Côte, 2000). These interactions have been extensively studied, particularly in tropical regions, where obligate cleaner fish (cleaners that depend exclusively on client-gleaned material throughout their life cycle) occupy territories (known as cleaning stations) that are visited by fish species (often referred to as clients). Cleaner-client fish dyads are perfect candidate models to study the role of microbiome in mutualistic behaviour and physiology as both partners come in close physical contact during interactions, allowing direct microbe transmission (skin-to-skin) while potentially also modulating the gut microbiome of cleaners, which feed actively of clients' mucus, parasites and hyperparasites.

In recent years, a priority has been identification of neurohormonal candidates that may modulate behavioural changes in marine cleaning mutualisms, allowing us to relate social cognition to the underlying mechanistic and neural mechanisms (Soares, 2017). These studies have highlighted the importance of neuromodulators such as nonapeptide arginine-vasotocin (AVT), which has substantial motivational and learning effects on cleaners (Soares, Bshary, Mendonça, Grutter, & Oliveira, 2012a; Cardoso et al., 2015a, Cardoso, Paitio, Oliveira, Bshary, & Soares, 2012b). Monoamines such as serotonin and dopamine (DA) also have a crucial impact, with serotonin increases being responsible for changes in motivation to engage mutualistically (Paula, Messias,

Grutter, Bshary, & Soares, 2015), while disruption in dopamine transmission leads the cleaner to anticipate a lower probability of feeding or, alternatively, a higher likelihood of being punished by being chased or the client leaving (Messias, Paula, Grutter, Bshary, & Soares, 2016a). The effects of pharmacological stimulation of the DAergic system were most evident during the learning process, in which individual learning abilities of cleaners were tested in laboratory conditions (Messias, Santos, Pinto, & Soares, 2016b). The involvement of the dopaminergic system on cleaners' decision-making changed depending on context and the predictive condition of reward (for instance when time delays occur, Soares, Cardoso, Malato, & Messias, 2017a), or with familiarity/novelty (Soares, Santos, & Messias, 2017b). Moreover, the role of social stress on cleaning performance has been studied. Indeed, one of the most notable behaviours by these animals is the cleaning and entering of predators' mouths, a behaviour that has been interpreted as altruistic on behalf of the clients that may simply eat the cleaner (Trivers, 1971). This seemingly fearless behaviour became a paradoxical feature of cleaners' proactivity and 'gutsy' cooperative behaviour, but also of putative anxiety and stress control, particularly when cleaners were found to show a preference for predatory over herbivorous clients, despite no obvious increase in ectoparasitic crustaceans in these hosts (Soares, Cardoso, & Côté, 2007). The main hypothesis raised to explain this behaviour is that cleaners need to be more proactive and spend more time inspecting predators (therefore glean more parasites and other material), in response to an increase in cortisol levels of energetic expenditure and putative decrease of immune reactivity (Soares, Bshary, Cardoso, Côté, Oliveira, 2012b). The role of stress mechanisms mediating the cleaners' cooperative levels were further tested, as these mechanisms were discovered to be involved in cleaners' decisions, as the exogenous effects of cortisol rises were found to propitiate cleaners' behavioural switch from cooperation to cheating (e.g. more stress led to more cheating; Soares, Cardoso, Grutter, Oliveira, & Bshary, 2014).

Our current understanding of cleaner fishes' social behaviour and proximate mechanisms prompts us to hypothesise that the cleaners and clients' microbial composition may have a direct

role in determining the costs and benefits of these complex behavioural trade-offs. But at this point it is impossible to know which behaviours are influenced by cleaners' microbiota, especially given the diversity of social challenges cleaner must face.

4. Concluding remarks

The use of fish in microbiome-related research is expanding (Llewellyn et al., 2014), although currently the research is mostly ecological or diet focused on farmed or commercially valuable fish species (Box 2). Similar to mammals, the gut microbiome of fish seems to play an important role in the innate immune response (e.g. Gómez and Balcázar, 2008; Tarnecki et al., 2017; deBruijn et al., 2018), fish have also been shown to exhibit ontogenetic shifts in microbiome composition (Falcinelli et al., 2015; Roeselers et al., 2011; Hall et al., 2017; Shin et al., 2015), and there is also a diluting effect in microbiome differences caused by social proximity (Burns et al., 2017).

The use of fish as a complementary species in translational studies has been proposed before (Gerlai, 2014; Soares et al., 2018; Stewart et al., 2015) and one rationale for extending the breadth of model organisms in translational studies is that it increases the probability of finding shared mechanisms that represent "core" components of a given disease or function (van de Mooij-Malsen, Vinkers, Peterse, Olivier, & Kas, 2011; Gerlai, 2014). Recognized advantages of fish use in basic and applied neurobehavioural research include rapid generation time and fertility (typically external fertilization), small size, amenability to genetic manipulation, and finally intermediate physiological and behavioural complexity in relation to other model organisms (Friedrich, Genoud, Wanner, Ahrens, & Hughes 2013; Friedrich, Jacobson, & Zhu, 2010, Kalueff et al., 2012, 2014; Maximino et al., 2015; Norton & Bally-Cuif, 2010; Rinkwitz et al., 2011; Stewart et al., 2015; Soares et al., 2018). Moreover, complex neuroscientific tools and sophisticated behavioural paradigms have already been developed for the zebrafish model and behavioural studies of fish also cover a multitude of species. The combination of the laboratory tools available for zebrafish and the ecological and ethological perspectives gleaned from other fish species is there-

fore ideal (Hall, Serrano, Rodd, & Tropepe, 2014). Importantly, protein-coding gene orthology between the zebrafish and human genome is close to 70% (Howe et al., 2013) and several genomic, transcriptomic, and transgenics tools have been developed for some fish (Rinkwitz et al., 2011).

A key difference between fish and other terrestrial models for microbiome studies is the fact that living in aquatic conditions may increase the level of microbiota contamination from surrounding water. However, many studies have compared the surrounding water microbiome with that of the fish, and while for larvae colonization from water microbiota seems to occur, for adult fish the observed differences are enough to discard this hypothesis. At this point, the major drawback of our suggested systems is an overall lack of core information. So far microbiome research in fish disregards behavioural and physiological responses. Nevertheless, the potential of using fish models is endless considering their diversity, and the fact we may build from extensive knowledge regarding behavioural ecology (guppies, sticklebacks), physiology and genetics (cleaners and zebrafish) and animal production of certain key species. This opens the door for a wealth of research: for example, do most fish species share common behavioural and physiological endpoints arising from modulation by microbiota?

Summarizing, the diverse behavioural systems exhibit by fish sets the stage for future interdisciplinary research that should start with a strong ecological (field-derived) approach, being followed by laboratory-based cognitive and pharmacological experimentation. The implications of such integrated approaches may be of translational relevance to human behavioural deficits and psychiatric disorders, especially regarding the influence of social conditions to microbiome functionality (see Box 2).

Box list

Box 1. A move to therapy? The use of model fish species to unravel the effects of microbiome into human social behavioural disorders

Animal-focused research is pivotal to the advancement of scientific knowledge, but fish use in translational medical research has mostly focused on zebrafish. In neurosciences and behaviour, both larval and adult zebrafish have contributed to our understanding of the genetic control of brain processes, and the effects of pharmacological manipulations. In fact, the similarities between mammalian and zebrafish paradigms are a clear demonstration of the evolutionarily conserved nature of behaviour and physiological modulation across species (Kalueff et al., 2014). We expect fish to share an increasing role when it comes to microbiome-based research and, most specially, in contributing empirically to the influence of skin and gut microbiome on the onset and development of several human behavioural disorders.

The study of the microbiome of model fish organisms could generate interesting insights into conserved and derived features of host-microbiome interactions and their impact on behaviour. Importantly, while the study of fish models is likely to be useful in the study of the evolutionary biology of host-microbiome interactions, the availability of behavioural assays for anxiety in both adult and larval zebrafish offers potential biomedical applications. The development of models for stress-induced dysbiosis, novel developments regarding the zebrafish inflammasome (Angosto & Mulero, 2014), and the interactions between these mechanisms pave the way for innovative models for psychiatric disorders. In addition, fish are emerging models of social behaviour and associated pathological alterations. Sociality is a fundamental dimension of human interactions and altered social behaviour is a symptom of many psychiatric disorders. In fish, due to the widespread influence of ecological and ecophysiological research, the focus has been largely aggressive behaviour – a symptom of acute psychotic states, drug abuse and mood disorders. A recent review highlighted that the broader array of social behaviour and cognition – from social motivation, social anxiety and social avoidance through social cognition, reproductive behaviour, and altruism and cooperation – represent important endpoints which could be used as models

of specific disordered domains of psychiatric conditions related to mood disorders and social anxiety (Soares et al., 2018). Understanding how the microbiome interacts with these factors could open avenues for investigation on probiotics and other live biotherapeutics on psychiatric disorders (Marchesi et al., 2015; O'Toole, Marchesi, & Hill, 2017).

Box 2. The social life of farmed fish: the search for new boundaries of social learning and stress response

Fish farming is crucial for maintaining global protein sources for rising human consumption and for providing animals that can restock exploited or/and degraded habitats or for create new fishing areas (Fernö, Huse, Jakobsen, & Kristiansen, 2006). When these hatchery-reared fish, however, are released into the wild they tend to have low survival partially due to behavioural deficits, although other causes (and consequences) are still under debate (Young, 2013; NASCO, 2017). Overall, available information indicates that this is a product of rearing environment, which limits fish exposure to natural stimuli and has profound consequences in fish learning and decision-making skills. In summarizing the differences (and implications) between natural and farmed fish, we identify the following:

1. Behavioural shifts. i) Foraging: fish reared in restricted conditions are less able to adjust to novel situations, which may occur when reared fish are released into the wild, needing to identify and catch live prey; ii) Aggression and reproduction: implications of rapid growth and high density.

1. Spatial dynamics. Farmed fish live in high density groups, but because these constantly changing, the establishment of clear individual and collective behaviour may thus be limited. While exposure to a greater number of conspecifics could be a source of extra information and learning, in practice the ability to recognize and associate with conspecifics may also decrease with group

size for some species (Griffiths & Magurran, 1997; Ward, Holbrook, Krause, & Hart, 2005). Additionally, artificial grouping of similar sized animals may also modulate behaviour.

1. Coping and anti-predator responses. The monocultural and relatively sterile conditions of aquaculture tend to produce fish unable to appropriately cope with novel and complex (natural) environments. The lack of previous exposure to predators, limits individual recognition which may become fatal (Brown & Warburton, 1999).

2. Parasite resistance. Stressful conditions can lead to immunocompromised fish that are more prone to infection and directly transmitted pathogens can spread rapidly amongst high density hosts.

3. Poorer development and physiological state (ontogeny), due to differences in nutrition, exercise and putative chronic stress levels of tank reared fish. Indeed, a combination of factors during the development of these fish, such as high levels of stress exposure and absence of appropriate socio-environmental stimulation may lead to limited neural and sensory development (Fernö et al., 2006).

In addition, many farmed fish are genetically selected, typically targeting rapid growth traits, which may also result in behavioural changes. In this respect, condition of the skin microbiota could also play a role in their behavioural neurobiology and welfare. For instance, farmed fish are more susceptible to pathogens compared to wild fish (e.g. Pulkinen et al), which may be mediated via changes to their microbiomes (but see Minniti et al., 2017). Thus, the implications of fish domestication in terms of behavioural, microbial and physiological variation urgently warrants further research.

Acknowledgements

MCS and RX are currently supported by FCT under the Programa Operacional Potencial Humano – Quadro de Referência Estratégico Nacional funds from the European Social Fund and Portuguese Ministério da Educação e Ciência (MCS: SFRH/BPD/109433/2015; RX: IF/00359/2015). Research on cooperative fish was supported by the European Regional Development Fund (ERDF) through COMPETE program and by National Funds through FCT - Foundation for Science and Technology (projects PTDC/MAR-BIO/0902/2014- POCI-01-0145-FEDER-016550; PTDC/BIA-MIC/27995/2017- POCI-01-0145-FEDER-027995; and PTDC/MAR/105276/2008)). JC and RX were partially funded by the Welsh Government and Higher Education Funding Council for Wales (HEFCW) AquaWales Project through the Sêr Cymru National Research Network for Low Carbon Energy and Environment (NRN-LCEE). CM and MGLM are currently supported by CNPq/Brazil under Edital Universal 2016 (CM: 400726/2016-5; MGLM: 423735/2016-0). The authors have no conflicting interests.

References

- Adleberth, I., Cerquetti, M., Poilane, I., Wold, A., & Collignon, A. (2000). Mechanisms of colonisation and colonisation resistance of the digestive tract part 1: Bacteria/host interactions. *Microbial Ecology in Health and Disease*, 12, 223-239. doi: 10.1080/089106000750060486
- Ait-Belgnaoui, A., Durand, H., Cartier, C., Chaumaz, G., Eutamene, H., Ferrier, L., Houdeau, E., Fioramonti, J., Bueno, L., & Theodorou, V. (2012). Prevention of gut leakiness by a probiotic treatment leads to attenuated HPA response to an acute psychological stress in rats. *Psychoneuroendocrinology*, 37, 1885-1895. doi: 10.1016/j.psyneuen.2012.03.024.
- Angosto, D., & Mulero, V. (2014). The zebrafish as a model to study the inflammasome. *Inflammasome*, 1, 27-29. doi:10.2478/infl-2014-0002.

Archie, E. A., & Tung, J. (2015). Social behavior and the microbiome. *Current Opinion in Behavioral Sciences*, 6, 28-34. doi: 10.1016/j.cobeha.2015.07.008.

Barden, N. (2004). Implication of the hypothalamic–pituitary–adrenal axis in the physiopathology of depression. *Journal of Psychiatry and Neurosciences*, 29, 185-193.

Bailey, M. T., & Coe, C. L. (1999). Maternal separation disrupts the integrity of the intestinal microflora in infant rhesus monkeys. *Development Psychobiology*, 35, 146-155. doi:10.1002/(SICI)1098-2302(199909)35:2<146::AID-DEV7>3.0.CO;2-G

Bell, M. A., & Foster, S. A. (1994). *The Evolutionary Biology of the Threespine Stickleback*. Oxford; New York: Oxford University Press.

Binning, S. A., Rey, O., Wismer, S., Triki, Z., Glauser, G., Soares, M. C., & Bshary, R., (2017). Reputation management promotes strategic adjustment of service quality in cleaner wrasse. *Scientific Reports*, 7, 8425. doi:10.1038/s41598-017-07128-5.

Bolnick, D. I., Snowberg, L. K., Caporaso, J. G., Lauber, C., Knight, R., & Stutz, W. E. (2014a). Major Histocompatibility Complex class IIb polymorphism influences gut microbiota composition and diversity. *Molecular Ecology*, 23, 4831-4845. doi: 10.1111/mec.12846

Bolnick, D. I., Snowberg, L. K., Hirsch, P. E., Lauber, C. L., Org, E., Parks, B., Lusi, A. J., Knight, R., Caporaso, J. G., & Svanbäck, R. (2014b). Individual diet has sex-dependent effects on vertebrate gut microbiota. *Nature Communications*, 5, 4500. doi: 10.1038/ncomms5500.

Bolnick, D. I., Snowberg, L. K., Hirsch, P. E., Lauber, C. L., Knight, R., Caporaso, J. G., & Svanbäck, R. (2014c). Individuals' diet diversity influences gut microbial diversity in two freshwater fish (threespine stickleback and Eurasian perch). *Ecology Letters*, 17, 979-987. doi: 10.1111/ele.12301.

Bonder, M. J., Kurilshikov, A., Tigchelaar, E. F., Mujagic, Z., Imhann, F., Vila, A. V., Deelen, P., Vatanen, T., Schirmer, M., Smeekens, S. P., Zhernakova, D. V., Jankipersadsing, S. A., Jaeger,

M., Oosting, M., Cenit, M. C., Masclee, A. A. M., Swertz, M. A., Li, Y., Kumar, V., Joosten, L., Harm-
sen, H., Weersma, R. K., Franke, L., Hofker, M. H., Xavier, R. J., Jonkers, D., Netea, M. G., Wij-
menga, C., Fu, J., & Zhernakova, A. (2016). The effect of host genetics on the gut microbi-
ome. *Nature Genetics*, 48, 1407-1412. doi: 10.1038/ng.3663.

Borre, Y. E., O’Keeffe, G. W., Clarke, G., Stanton, C., Dinan, T. G., Cryan, J. F. (2014). Microbiota
and neurodevelopmental windows: Implications for brain disorders. *Trends in Molecular Medi-
cine*, 20, 509-518. doi: 10.1016/j.molmed.2014.05.002.

Borrelli, L., Aceto, S., Agnisola, C., De Paolo, S., Dipineto, L., Stilling, R. M., Dinan, T. G., Cryan, J.
F., Menna, L. F., & Fioretti, A. (2016). Probiotic modulation of the mi-crobiota-gut-brain axis and
behaviour in zebrafish. *Scientific Reports*, 6, 30046. doi:10.1038/srep30046.

Boutin, S., Bernatchez, L., Audet, C., & Derôme, N. (2013). Network analysis highlights complex
interactions between pathogen, host and commensal microbiota. *PloS one*, 8,
e84772. doi:10.1371/journal.pone.0084772

Brown, C. (2015). Fish intelligence, sentience and ethics. *Animal Cognition*, 18, 1-17. doi:
10.1007/s10071-014-0761-0.

Brown, C., & Laland, K. (2001). Social learning and life skills training for hatchery reared fish. *Jour-
nal of Fish Biology*, 59, 471–493. doi:10.1006/jfbi.2001.1689.

Brown, C., & Warburton, K. (1999). Differences in timidity and escape responses between pred-
ator-naive and predator-sympatric rainbowfish populations. *Ethology*, 105, 491-502. doi:
10.1046/j.1439-0310.1999.00442.x.

Brown, C., Laland, K., & Krause, J. (2006) *Fish Cognition and Behavior*. Black-well Publishing Ltd,
Oxford, UK. doi: 10.1002/9780470996058.fmatter.

Bshary, R., Wickler, W., & Fricke, H. (2002) Fish cognition: A primate's eye view. *Animal Cognition*,
5, 1-13. doi:10.1007/s10071-001-0116-5.

Bshary, R., & Côté, I. M. (2008). New perspectives on marine cleaning mutualism. In Magnhagen, C., Braithwaite, V. A., Forsgren, E., & Kapoor, B. G. (Eds.), *Fish Behaviour* (pp. 563–592). Science Publishers, New Hampshire.

Burns, A. R., Miller, E., Agarwal, M., Rolig, A. S., Milligan-Myhre, K., Seredick, S., Guillemin, K. & Bohannan, B. J. M. (2017). Interhost dispersal alters microbiome assembly and can overwhelm host innate immunity in an experimental zebrafish model. *Proceedings of the National Academy of Sciences*, 114, 11181. doi: 10.1073/pnas.1702511114

Cardoso, S. C., Bshary, R., Mazzei, R., Paitio, J. R., Oliveira, R. F., & Soares, M. C. (2015a). Arginine vasotocin modulates associative learning in a mutualistic cleaner fish. *Behavior Ecology and Sociobiology*, 69, 1173-1181. doi:10.1007/s00265-015-1931-z.

Cardoso, S. C., Paitio, J. R., Oliveira, R. F., Bshary, R., & Soares, M. C. (2015b). Arginine vasotocin reduces levels of cooperative behaviour in a cleaner fish. *Physiology and Behaviour*, 139, 314-320. doi:10.1016/j.physbeh.2014.11.052.

Carlson, J. M., Hyde, E. R., Petrosino, J. F., Manage, A. B., & Primm, T. P. (2015). The host effects of *Gambusia affinis* with an antibiotic-disrupted microbiome. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 178, 163-168. doi:10.1016/j.cbpc.2015.10.004.

Carlson, J. M., Leonard, A. B., Hyde, E. R., Petrosino, J. F., & Primm, T. P. (2017). Microbiome disruption and recovery in the fish *Gambusia affinis* following exposure to broad-spectrum antibiotic. *Infection and Drug Resistance*, 10, 143-154. doi:10.2147/IDR.S129055.

Clark, K. J., Boczek, N. J., & Ekker, S. C. (2011). Stressing zebrafish for behavioral genetics. *Reviews in the Neurosciences*, 22, 49-62. doi: 10.1515/rns.2011.007.

Clarke, G., Grenham, S., Scully, P., Fitzgerald, P., Moloney, R. D., Shanahan, F., Dinan, T. G., & Cryan, J. F. (2013). The microbiome-gut-brain axis during early life regulates the hippocampal

serotonergic system in a sex-dependent manner. *Molecular Psychiatry*, 18, 666-673. doi: 10.1038/mp.2012.77.

Cong, X., Xu, W., Janton, S., Henderson, W. A., Matson, A., McGrath, J. M., Maas, K., & Graf, J. (2016). Gut microbiome developmental patterns in early life of preterm in-fants: Impacts of feeding and gender. *PLoS ONE*, 11:e0152751. doi: 10.1371/journal.pone.0152751.

Côté, I. M. (2000). Evolution and ecology of cleaning symbioses in the sea. *Oceanography and Marine Biology*, 38, 311–355.

Côté, I. M., & Soares, M. C. (2011). Gobies as cleaners. In R. A. Patzner, J. L. Van Tas-sell, M. Kovačičc & B. G. Kapoor (Eds) *The Biology of Gobies* (pp. 563-592). Enfield, NH: Science Publishers Inc.

Cresko, W. A., McGuigan, K. L., Phillips, P. C., & Postlethwait, J. H. (2007). Studies of threespine stickleback developmental evolution: progress and promise. *Genetica*, 129: 105-126. doi:10.1007/s10709-006-0036-z.

Croft, D. P., Edenbrow, M., Darden, S.K., Ramnarine, I. W., van Oosterhout, C., & Cable, J. (2011). Effect of gyrodactylid ectoparasites on host behaviour and social network structure in guppies *Poecilia reticulata*. *Behavioral Ecology and Sociobiology*, 65, 2219-2227. doi: 10.1007/s00265-011-1230-2.

Croft D. P., Krause, J., & James, R. (2004). Social networks in the guppy (*Poecilia reticulata*). *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271, S516-S519. doi: 10.1098/rsbl.2004.0206

Davis, D. J., Doerr, H. M., Grzelak, A. K., Busi, S. B., Jasarevic, E., Ericsson, A. C., & Bryda, E. C. (2016). *Lactobacillus plantarum* attenuates anxiety-related behavior and protects against stress-induced dysbiosis in adult zebrafish. *Scientific Reports*, 6, 33726 doi: 10.1038/srep33726.

de Bruijn, I., Liu, Y., Wiegertjes, G. F., Raaijmakers, J. M. (2018) Exploring fish microbial communities to mitigate emerging diseases in aquaculture. *FEMS Microbiology Ecology*, 94, fix161. <https://doi.org/10.1093/femsec/fix161>

de Mooij-van Malsen, A. J. G., Vinkers, C. H., Peterse, D. P., Olivier, B., & Kas, M. J.

H. (2011). Cross-species behavioural genetics: A starting point for unravelling the neurobiology of human psychiatric disorders. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35, 1383-1390. doi: <https://doi.org/10.1016/j.pnpbp.2010.10.003>

DeGruttola, A. K., Low, D., Mizoguchi, A., & Mizoguchi, E. (2016). Current Understanding of Dysbiosis in Disease in Human and Animal Models. *Inflammatory bowel diseases*, 22, 1137-1150. doi: 10.1097/MIB.0000000000000750

Desbonnet, L., Clarke, G., Shahanan, F., Dinan, T. G., & Cryan, J. F. (2014). Microbiota is essential for social development in the mouse. *Molecular Psychiatry*, 19, 146-148. doi: 10.1038/mp.2013.65.

Dugatkin, L. A. (1988). Do guppies play TIT FOR TAT during predator inspection visits? *Behavioral Ecology and Sociobiology*, 23, 395. doi: 10.1007/BF00303714.

Dugatkin, L. A., & Alfieri, M. (1991). Guppies and the TIT FOR TAT strategy: preference based on past interaction. *Behavioral Ecology and Sociobiology*, 28, 243. doi:10.1007/BF00175096

Egerton, S., Culloty, S., Whooley, J., Stanton, C., & Ross, R. P. (2018). The Gut Microbiota of Marine Fish. *Frontiers in Microbiology*, 9, 873. doi:10.3389/fmicb.2018.00873

Eizaguirre, C., Lenz, T. L., Kalbe, M., & Milinski, M. (2012). Rapid and adaptive evolution of MHC genes under parasite selection in experimental vertebrate populations. *Nature Communications*, 3, 621. doi:10.1038/ncomms1632.

Elinav, E., Strowig, T., Henao-Mejia, J., & Flavell, R. A. (2011). Regulation of the anti-microbial response by NLR proteins. *Immunity*, 34, 665-679. doi: 10.1016/j.immuni.2011.05.007

Endler, J.A. (1995). Multiple-trait coevolution and environmental gradients in guppies. *Trends in Ecology and Evolution*, 10, 22-29.

Endt, K., Stecher, B., Chaffron, S., Slack, E., Tchitchek, N., Benecke, A., Van Maele, L., Sirard, J.-C., Mueller, A. J., Heikenwalder, M., Macpherson, A. J., Strugnell, R., von Mering, C., & Hardt, W.-D. (2010). The microbiota mediates pathogen clearance from the gut lumen after non-typhoidal *Salmonella* diarrhea. *PLoS Pathogens*, 6, e1001097. doi: 10.1371/journal.ppat.1001097.

Evans, J. P., Pilastro, P., & Schlupp, I. (2011). *Ecology and Evolution of Poeciliid Fishes*. The University of Chicago Press.

Ezenwa, V. O., & Williams, A. E. (2014). Microbes and animal olfactory communication: Where do we go from here? *BioEssays*, 36, 847-854. doi: 10.1002/bies.201400016.

Falcinelli, S., Picchietti, S., Rodiles, A., Cossignani, L., Merrifield, D. L., Taddei, A. R., Maradonna, F., Olivotto, I., Gioacchini, G., & Carnevali, O. (2015). *Lactobacillus rhamnosus* lowers zebrafish lipid content by changing gut microbiota and host transcription of genes involved in lipid metabolism. *Scientific Reports*, 5, 9336. doi: 10.1038/srep09336.

Fernö, A., Huse, G., Jakobsen, P. J. & Kristiansen, T. S. (2006). The role of fish learning skills in fisheries and aquaculture. In C. Brown, K. Laland & J. Krause (Eds) *Fish Cognition and Behavior*, Blackwell Publishing Ltd, Oxford, UK. doi: 10.1002/9780470996058.ch14

Farr, J. A. (1975). The role of predation in the evolution of social behaviour of natural populations of the guppy, *Poecilia reticulata* (Pisces: Poeciliidae). *Evolution*, 29, 151-158. doi: 10.1111/j.1558-5646.1975.tb00822.x.

Fons, M., Gomez, A., & Karjalainen T. (2000). Mechanisms of colonisation and colonisation resistance of the digestive tract part 2: Bacteria/bacteria interactions. *Microbial Ecology in Health and Disease*, 12, 240-246. doi: 10.1080/089106000750060495.

- Foster, J. A., Rinaman, L., & Cryan, J. F. (2017). Stress & the gut-brain axis: Regulation by the microbiome. *Neurobiology of Stress*, 7, 124-136. doi: 10.1016/j.ynstr.2017.03.001.
- Friedrich, R. W., Genoud, C., Wanner, A. A., Ahrens, M. B., & Hughes, H. (2013). Analyzing the structure and function of neuronal circuits in zebrafish. *Frontiers Neural Circuits*, 7, 71. doi:10.3389/fncir.2013.00071.
- Friedrich, R. W., Jacobson, G. A., & Zhu, P. (2010). Circuit neuroscience in zebrafish. *Current Biology*, 20, R371–R381. doi:10.1016/j.cub.2010.02.039.
- García-Ródenas, C. L., Bergonzelli, G. E., Nutten, S., Schumann, A., Cherbut, C., Turini, M., Ornstein, K., Rochat, F., & Corthésy-Theulaz, I. (2006). Nutritional approach to restore impaired intestinal barrier function and growth after neonatal stress in rats. *Journal of Pediatric Gastroenterology and Nutrition*, 43, 16-24. doi: 10.1097/01.mpg.0000226376.95623.9f.
- Gareau, M. G., Silva, M. A., & Perdue, M. H. (2008). Pathophysiological mechanisms of stress-induced intestinal damage. *Current Molecular Medicine*, 8: 274-281.
- Garrett, W. S., Gordon, J. I., & Glimcher, L. H. (2010). Homeostasis and inflammation in the intestine. *Cell*, 140, 859-870. doi: 10.1016/j.cell.2010.01.023.
- Gerlai, R. (2014). Fish in behavior research: Unique tools with a great promise! *Journal of Neuroscience Methods*, 234, 54-58. doi: 10.1016/j.jneumeth.2014.04.015.
- Gilbert, S. F. (2014). A holobiont birth narrative: the epigenetic transmission of the human microbiome. *Frontiers in Genetics*, 5, 282. doi: 10.3389/fgene.2014.00282.
- Gómez, G. D., & Balcázar, J. L. (2008). A review on the interactions between gut microbiota and innate immunity of fish. *FEMS Immunology & Medical Microbiology*, 52, 145-154. doi:10.1111/j.1574-695X.2007.00343.x

Gonçalves, A. T., & Gallardo-Escarate, C. (2017). Microbiome dynamic modulation through functional diets based on pre- and probiotics (mannan-oligosaccharides and *Saccharomyces cerevisiae*) in juvenile rainbow trout (*Oncorhynchus mykiss*). *Journal of Applied Microbiology*, 122, 1333–1347. doi: 10.1111/jam.13437.

Griffiths, S. W., & Magurran, A. E. (1997). Schooling preferences for familiar fish vary with group size in a wild guppy population. *Proceedings of the Royal Society B: Biological Sciences*, 264, 547-551. doi: 10.1098/rspb.1997.0078.

Groussin, M., Mazel, F., Sanders, J. G., Smillie, C. S., Lavergne, S., Thuiller, W., & Alm E. J. (2017). Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nature Communications*, 8, 14319. doi: 10.1038/ncomms14319

Grutter, A. S., Murphy, J. M., & Choat, H. (2003). Cleaner fish drives local fish diversity on coral reefs. *Current Biology*, 13, 64-67. doi: 10.1016/S0960-9822(02)01393-3.

Guarner, F. G., & Malagelada, J.-R. (2003). Gut flora in health and disease. *Lancet*, 361, 512-519. doi: 10.1016/S0140-6736(03)12489-0

Guttridge, T. L., Gruber, S. H., DiBattista, J. D., Feldheim, K. A., Croft, D. P., Krause, S., & Krause, J. (2011). Assortative interactions and leadership in a free-ranging population of juvenile lemon shark *Negaprion brevirostris*. *Marine Ecology Progress Series*, 423, 235-245. <https://doi.org/10.3354/meps08929>

Guttridge, T. L., Gruber, S. H., Krause, J., & Sims, D. W. (2010). Novel Acoustic Technology for Studying free-ranging shark social behaviour by recording individuals' interactions. *PLoS ONE*, 5, e9324. doi:10.1371/journal.pone.0009324

Hall, Z. J., De Serrano, A. R., Rodd, F. H., & Tropepe, V. (2014). Casting a wider fish net on animal models in neuropsychiatric research. *Progress in Neuropsychopharmacology and Biological Psychiatry*, 55, 7-15. doi: 10.1016/j.pnpbp.2014.04.003.

Hall, A. B., Tolonen, A. C., & Xavier, R. J. (2017). Human genetic variation and the gut microbiome in disease. *Nature Reviews Genetics*, 18, 690-699. doi: 10.1038/nrg.2017.63.

Hennersdorf, P., Kleinertz, S., Theisen, S., Abdul-Aziz, M. A., Mrotzek, G., Palm, H. W., & Saluz, H. P. (2016). Microbial diversity and parasitic load in tropical fish of different environmental conditions. *PLoS ONE*, 11, e0151594. doi:10.1371/journal.pone.0151594.

Howe, K., Clark, M. D., Torroja, C. F., Torrance, J., Berthelot, C., Muffato, M., Collins, J. E., Humphray, S., McLaren, K., Matthews, L., McLaren, S., Sealy, I., Caccamo, M., Churcher, C., Scott, C., Barrett, J. C., Koch, R., Rauch, G. J., White, S., Chow, W., Kilian, B., Quintais, L. T., Guerra-Assunção, J. A., Zhou, Y., Gu, Y., Yen, J., Vogel, J. H., Eyre, T., Redmond, S., Banerjee, R., Chi, J., Fu, B., Langley, E., Maguire, S. F., Laird, G. K., Lloyd, D., Kenyon, E., Donaldson, S., Sehra, H., Almeida-King, J., Loveland, J., Trevanion, S., Jones, M., Quail, M., Willey, D., Hunt, A., Burton, J., Sims, S., McLay, K., Plumb, B., Davis, J., Clee, C., Oliver, K., Clark, R., Riddle, C., Elliot, D., Threadgold, G., Harden, G., Ware, D., Begum, S., Mortimore, B., Kerry, G., Heath, P., Phillimore, B., Tracey, A., Corby, N., Dunn, M., Johnson, C., Wood, J., Clark, S., Pelan, S., Griffiths, G., Smith, M., Glithero, R., Howden, P., Barker, N., Lloyd, C., Stevens, C., Harley, J., Holt, K., Panagiotidis, G., Lovell, J., Beasley, H., Henderson, C., Gordon, D., Auger, K., Wright, D., Collins, J., Raisen, C., Dyer, L., Leung, K., Robertson, L., Ambridge, K., Leongamornlert, D., McGuire, S., Gilderthorp, R., Griffiths, C., Manthravadi, D., Nichol, S., Barker, G., Whitehead, S., Kay, M., Brown, J., Murnane, C., Gray, E., Humphries, M., Sycamore, N., Barker, D., Saunders, D., Wallis, J., Babbage, A., Hammond, S., Mashreghi-Mohammadi, M., Barr, L., Martin, S., Wray, P., Ellington, A., Matthews, N., Ellwod, M., Woodmansey, R., Clark, G., Cooper, J., Tromans, A., Grafham, D., Skuce, C., Pandian, R., Andrews, R., Harrison, E., Kimberley, A., Garnett, J., Fosker, N., Hall, R., Garner, P., Kelly, D., Bird, C., Palmer, S., Gehring, I., Berger, A., Dooley, C. M., Ersan-Ürün, Z., Eser, C., Geiger, H., Geisler, M., Karotki, L., Kirn, A., Konantz, J., Konantz, M., Oberländer, M., Rudolph-Geiger, S., Teucke, M., Lanz, C., Raddatz, G., Osoegawa, K., Zhu, B., Rapp, A., Widaa, S., Langford, C., Yang, F., Schuster, S. C., Carter, N. P., Harrow, J., Ning, Z., Herrero, J., Searle, S. M., Enright, A., Geisler, R., Plasterk, R. H.,

Lee, C., Westerfield, M., de Jong, P. J., Zon, L. I., Postlethwait, J. H., Nüsslein-Volhard, C., Hubbard, T. J., Roest Crolius, H., Rogers, J., & Stemple, D. L. (2013). The zebrafish reference genome sequence and its relationship to the human genome. *Nature*, 496, 498-503. doi: 10.1038/nature12111

Huntingford, F. A., Ostlund-Nilsson, S., & Mayer, I. (2006) *Biology of the Three Spined Stickleback*. Taylor & Francis Group, CRC Press.

Jesuthasan, S. J., & Mathuru, A. S. (2008). The alarm response in zebrafish: Innate fear in a vertebrate genetic model. *Journal of Neurogenetics*, 22, 221-228. doi:10.1080/01677060802298475

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., Swofford, R., Pirun, M., Zody, M. C., White, S., Birney, E., Searle, S., Schmutz, J., Grimwood, J., Dickson, M. C., Myers, R. M., Miller, C. T., Summers, B. R., Knecht, A. K., Brady, S. D., Zhang, H., Pollen, A. A., Howes, T., Amemiya, C., B. I. G. S., W. G. A. Team, Lander, E. S., Di Palma, F., Lindblad-Toh, K., & Kingsley, D. M. (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, 484, 55-61. doi: 10.1038/nature10944.

Kalueff, A. V., Stewart, A.M., & Gerlai, R. (2014). Zebrafish as an emerging model for studying complex brain disorders. *Trends in Pharmacological Sciences*, 35, 63-75. doi: 10.1016/j.tips.2013.12.002.

Kalueff, A. V., Stewart, A. M., Kyzar, E. J., Cachat, J., Gebhardt, M., Landsman, S., Robinson, K., Maximino, C., Herculano, A. M., Jesuthasan, S., Wisenden, B., Bally-Cuif, L., Lange, M., Vernier, P., Norton, W., Tierney, K., Tropepe, V., Neuhauss, S. C. F., & Zebrafish Neuroscience Research Consortium (2012). Time to recognize zerafish "affective" behavior. *Behaviour*, 149, 1019-1036. doi:10.1163/1568539X-00003030.

Kelly, C. J., Zheng, L., Campbell, E. L., Saeedi, B., Scholz, C. C., Bayless, A. J., Wilson, K. E., Glover, L. E., Kominsky, D. J., Magnuson, A., Weir, T. L., Ehrentraut, S. F., Pickel, C., Kuhn, K. A., Lanis, J. M., Nguyen, V., Taylor, C. T., & Colgan, S. P. (2015). Crosstalk between Microbiota-Derived Short-

Chain Fatty Acids and Intestinal Epithelial HIF Augments Tissue Barrier Function. *Cell host & Microbe*, 17, 662-671. doi: 10.1016/j.chom.2015.03.005

Kelly, J.R., Borre, Y., O'Brien, C., Patterson, E., El Aidy, S., Deane, J., Kennedy, P. J., Beers, S., Scott, K., Moloney, G., Hoban, A. E., Scott, L., Fitzgerald, P., Ross, P., Stanton, C., Clarke, G., Cryan, J. F., & Dinan, T. G. (2016). Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat. *Journal of Psychiatric Research*, 82, 109-118. doi: 10.1016/j.jpsychires.2016.07.019

Koch, H., & Schmid-Hempel, P. (2011). Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proceeding of Natural Academy of Sciences USA*, 108, 19288-19292. doi: 10.1073/pnas.1110474108.

Krause, S., Wilson, A. D. M, Ramnarine, I. W., Herbert-Read, J. E., Clement, R. J. G., & Krause, J. (2016). Guppies occupy consistent positions in social networks: mechanisms and consequences. *Behavioral Ecology*, 28, 429-438. <https://doi.org/10.1093/beheco/arw177>

Kunze, W. A., Mao, Y. K., Wang, B., Huizinga, J. D., Ma, X., Forsythe, P., & Bienenstock, J. (2009). *Lactobacillus reuteri* enhances excitability of colonic AH neurons by inhibiting calcium-dependent potassium channel opening. *Journal of Cellular and Molecular Medicine*, 13, 2261-2270. doi: 10.1111/j.1582-4934.2009.00686.x

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M., Gibbons, S. M., Larsen, P., Shogan, B. D., Weiss, S., Metcalf, J. L., Ursell, L. K., Vázquez-Baeza, Y., Van Treuren, W., Hasan, N. A., Gibson, M. K., Colwell, R., Dantas, G., Knight, R., & Gilbert, J. A. (2014). Longitudinal analysis of microbial interaction between humans and the indoor environment. *Science*, 345, 1048-1052. doi: 10.1126/science.1254529

Levin, I. I., Zonana, D. M., Fosdick, B. K., Song, S. J., Knight, R., & Safran, R. J. (2016). Stress response, gut microbial diversity and sexual signals correlate with social interactions. *Biology letters*, 12, 20160352. doi: 10.1098/rsbl.2016.0352

Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012). Diversity, stability and resilience of the human gut microbiota. *Nature*, 489, 220. doi:10.1038/nature11550

Li, T., Long, M., Ji, C., Shen, Z., Gatesoupe, F. -J., Zhang, X., Zhang, Q., Zhang, L., Zhao, Y., Liu, X., & Li, A. (2016). Alterations of the gut microbiome of largemouth bronze gudgeon (*Coreius guichenoti*) suffering from furunculosis. *Scientific Reports*, 6, 30606. doi: 10.1038/srep30606.

Llewellyn, M. S., Boutin, S., Hoseinifar, S. H., & Derome, N. (2014). Teleost microbiomes: The state of the art in their characterization, manipulation and importance in aquaculture and fisheries. *Frontiers in Microbiology*, 5, 207. doi: 10.3389/fmicb.2014.00207.

Llewellyn, M.S., Leadbeater, S., Garcia, C., Sylvain, F.E., Custodio, M., Ang, K.P., Powell, F., Carvalho, G.R., Creer, S., Elliot, J., & Derome, N. (2017). Parasitism perturbs the mucosal microbiome of Atlantic Salmon. *Scientific Reports*, 7, 43465. doi: 10.1038/srep43465.

Lyte, M., Li, W., Opitz, N., Gaykema, R. P., & Goehler, L. E. (2006). Induction of anxiety-like behavior in mice during the initial stages of infection with the agent of murine colonic hyperplasia *Citrobacter rodentium*. *Physiology and Behavior*, 89, 350-357. doi: 10.1016/j.physbeh.2006.06.019.

Magurran AE. (2005). *Evolutionary Ecology: The Trinidadian Guppy*. Oxford University Press: New York.

Marchesi, J. R., Adams, D. H., Fava, F., Hermes, G. D. A., Hirschfield, G. M., Hold, G., Quraishi, M. N., Kinross, J., Smidt, H., Tuohy, K. M., Thomas, L. V., Zoetendal, E. G., & Hart, A. (2015). The gut microbiota and host health: a new clinical frontier. *Gut*, 65, 330-339. doi: 10.1136/gutjnl-2015-309990.

Matthews B, Harmon LJ, M'Gonigle L, Marchinko KB, & Schaschl H (2010) Sympatric and allopatric divergence of MHC genes in threespine stickleback. *PLoS One*, 5, e10948. Doi:10.1371/journal.pone.0010948.

Maximino, C., de Brito, T. M., Batista, A. W. S., Herculano, A. M., Morato, S., & Gouveia Jr., A. (2010). Measuring anxiety in zebrafish: A critical review. *Behavioural Brain Research*, 214, 151-171. doi: 10.1016/j.bbr.2010.05.031.

Maximino, C., Silva, R.X. do C., da Silva, S. de N.S., Rodrigues, L. do S.D.S., Barbosa, H., de Carvalho, T.S., Leão, L.K.D.R., Lima, M.G., Oliveira, K.R.M., & Herculano, A.M. (2015). Non-mammalian models in behavioral neuroscience: Consequences for biological psychiatry. *Frontiers in Behavioral Neurosciences*, 9, 233. doi:10.3389/fnbeh.2015.00233

Mayer I, Meager J, Skjæraasen JE, Rodewald P, Sverdrup G, & Fernö A. (2010). Domestication causes rapid changes in heart and brain morphology in Atlantic cod (*Gadus morhua*). *Environmental Biology of Fishes*, 92, 181-186. doi: 10.1007/s10641-011-9831-1.

McVey Neufeld, K. A., Mao, Y. K., Bienenstock, J., Foster, J. A., & Kunze, W. A. (2013). The microbiome is essential for normal gut intrinsic primary afferent neuron excitability in the mouse. *Neurogastroenterology and Motility*, 25, 183-188. doi: 10.1111/nmo.12049.

Meadow, J. F., Bateman, A. C., Herkert, K. M., O'Connor, T. K., & Green, J. L. (2013). Significant changes in the skin microbiome mediated by the sport of roller derby. *PeerJ*, 1, e53. doi: 10.7717/peerj.53

Messias, J.P.M., Paula, J.R., Grutter, A.S., Bshary, R., & Soares, M.C. (2016a). Dopamine disruption increases negotiation for cooperative interactions in a fish. *Scientific Reports*, 6, 20817. doi:10.1038/srep20817.

Messias, J. P. M., Santos, T. P., Pinto, M., & Soares, M. C. (2016b). Stimulation of dopamine D1 receptor improves learning capacity in cooperating cleaner fish. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20152272. doi:10.1098/rspb.2015.2272.

Milligan-Myhre, K., Small, C. M., Mittge, E. K., Agarwal, M., Currey, M., Cresko, W. A., & Guillemin, K. (2016). Innate immune responses to gut microbiota differ between oceanic and freshwater

threespine stickleback populations. *Disease Models and Mechanisms*, 9, 187-198. doi: 10.1242/dmm.021881.

Minniti, G., Hagen, L. H., Porcellato, D., Jørgensen, S. M., Pope, P. B., & Vaaje-Kolstad, G. (2017). The Skin-Mucus Microbial Community of Farmed Atlantic Salmon (*Salmo salar*). *Frontiers in microbiology*, 8, 2043. doi:10.3389/fmicb.2017.02043

Moeller, A. H., Foerster, S., Wilson, M. L., Pusey, A. E., Hahn, B. H., & Ochman, H. (2016). Social behavior shapes the chimpanzee pan-microbiome. *Science advances*, 2, e1500997. doi:10.1126/sciadv.1500997

Mourier, J., Vercelloni, J., & Planes, S. (2012). Evidence of social communities in a spatially structured network of a free-ranging shark species. *Animal Behaviour*, 83, 389-401. doi: <https://doi.org/10.1016/j.anbehav.2011.11.008>

Narrowe, A. B., Albuti-Lantz, M., Smith, E. P., Bower, K. J., Roane, T. M., Vajda, A. M., & Miller, C. S. (2015). Perturbation and restoration of the fathead minnow gut microbiome after low-level triclosan exposure. *Microbiome*, 3, 6. doi: 10.1186/s40168-015-0069-6.

NASCO (2017). Understanding the risks and benefits of hatchery and stocking activities to wild Atlantic salmon populations. Chaput, G., Knight, P., Russell, I., Sivertsen, A., Hutchinson, P. & Forero Segovia, S. L. (Eds) *Report of a Theme-based Special Session of the Council of NASCO* (61-116pp). NASCO Council document CNL(17)

Norton, W., & Bally-Cuif, L. (2010). Adult zebrafish as a model organism for behavioural genetics. *BMC Neurosciences*, 11, 90. doi:10.1186/1471-2202-11-90.

Oliveira, R.F., 2013. Mind the fish: Zebrafish as a model in cognitive social neuroscience. *Frontiers in Neural Circuits*, 7, 131. doi:10.3389/fncir.2013.00131

O'Mahony, S. M., Marchesi, J. R., Scully, P., Codling, C., Ceolho, A. M., Quigley, E. M., Cryan, J. F., & Dinan, T. G. (2009). Early life stress alters behavior, immunity, and microbiota in rats: Implications for irritable bowel syndrome and psychiatric illnesses. *Biological Psychiatry*, 65, 263-267. doi: 10.1016/j.biopsych.2008.06.026.

O'Toole, P. W., Marchesi, J., & Hill, C. (2017). Next-generation probiotics: The spectrum from probiotics to live biotherapeutics. *Nature Microbiology*, 2, 17057. doi: 10.1038/nmicrobiol.2017.57.

Pascoe, E. L., Hauffe, H. C., Marchesi, J. R., & Perkins, S. E. (2017). Network analysis of gut microbiota literature: An overview of the research landscape in non-human animal studies. *ISME Journal*, 11, 2644-2651. doi: 10.1038/ismej.2017.133.

Paula, J. R., Messias, J. P., Grutter, A. S., Bshary, R., Soares, M. C. (2015). The role of serotonin in the modulation of cooperative behavior. *Behavioral Ecology*, 26, 1005-1012. doi: 10.1093/beheco/arv039

Perofsky, A. C., Lewis, R. J., Abondano, L. A., Di Fiore, A., & Meyers, L. A. (2017). Hierarchical social networks shape gut microbial composition in wild Verreaux's sifaka. *Proceedings. Biological sciences*, 284, 20172274. doi: 10.1098/rspb.2017.2274

Phillips KP, Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., van Oosterhout, C., & Radwan, J. (2018) Immunogenetic novelty confers a selective advantage in host-pathogen coevolution. *Proceedings of the Natural Academy of Science U.S.A*, 115, 1552-1557. doi:10.1073/pnas.1708597115.

Poulin, R. (2013). Parasite manipulation of host personality and behavioural syndromes. *Journal of Experimental Biological*, 216, 18-26. doi: 10.1242/jeb.073353.

Pulkkinen, K., Suomalainen, L. R., Read, A. F., Ebert, D., Rintamäki, P., & Valtonen, E. T. (2009). Intensive fish farming and the evolution of pathogen virulence: the case of columnaris disease in Finland. *Proceedings. Biological Sciences*, 277, 593-600. doi: 10.1098/rspb.2009.1659

Ray, C., Bujan, N., Tarnecki, A., Davis D. A., Browdy, C., & Arias, C. R. (2017). Analysis of the gut microbiome of Nile tilapia *Oreochromis niloticus* L. fed diets supplemented with Previda® and saponin. *Journal of Fisheries Sciences.com*, 11, 36.

Reid, K.M., Patel, S., Robinson, A.J., Bu, L., Jarungsriapisit, J., Moore, L.J., & Salinas, I. (2017). Salmonid alphavirus infection causes skin dysbiosis in Atlantic salmon (*Salmo salar* L.) post-smolts. *PLoS ONE*, 12, e0172856. doi:10.1371/journal.pone.0172856.

Reznick, D. N., Shaw, F. H., Rodd, F. H., & Shaw, R. G. (1997). Evaluation of the rate of evolution in natural populations of guppies (*Poecilia reticulata*). *Science*, 275, 1934-1937.

Reznick, D. N., & Ricklefs, R. E. (2009). Darwin's bridge between microevolution and macroevolution. *Nature*, 457, 837-842. doi: 10.1038/nature07894.

Rinkwitz, S., Mourrain, P., & Becker, T.S. (2011). Zebrafish: An integrative system for neurogenetics and neurosciences. *Progress in Neurobiology*, 93, 231-243. doi:10.1016/j.pneurobio.2010.11.003.

Rivière, A., Selak, M., Lantin, D., Leroy, F., & De Vuyst, L. (2016). Bifidobacteria and Butyrate-Producing Colon Bacteria: Importance and Strategies for Their Stimulation in the Human Gut. *Frontiers in microbiology*, 7, 979. doi:10.3389/fmicb.2016.00979

Roeslers, G., Mittge, E. K., Stephens, W. Z., Parichy, D. M., Cavanaugh, C. M., Guillemin, K., & Rawls, J. F. (2011). Evidence for a core gut microbiota in the zebrafish. *ISME Journal*, 5, 1595-1608. doi: 10.1038/ismej.2011.38.

- Rosado, D., Pérez-Losada, M., Severino, R., Cable, J., & Xavier, R. (2019). Characterization of the skin and gill microbiomes of the farmed seabass (*Dicentrarchus labrax*) and seabream (*Sparus aurata*). *Aquaculture*, 500, 57-64. <https://doi.org/10.1016/j.aquaculture.2018.09.063>
- Sampson, T. R., & Mazmanian, S. K. (2015). Control of brain development, function, and behavior by the microbiome. *Molecular Psychiatry*, 21, 786-796. doi: 10.1016/j.chom.2015.04.011.
- Sandhu, K. V., Sherwin, E., Schellekens, H., Stanton, C., Dinan, T. G., & Cryan, J. F. (2017). Feeding the microbiota-gut-brain axis: diet, microbiome, and neuropsychiatry. *Translational Research*, 179, 223-244. doi: 10.1016/j.trsl.2016.10.002.
- Scharsack, J. P., Kalbe, M., Harrod, C., & Rauch, G. (2007). Habitat-specific adaptation of immune responses of stickleback (*Gasterosteus aculeatus*) lake and river eco-types. *Proceeding of the Royal Society B Biological Sciences*, 274, 1523-1532. doi: 10.1098/rspb.2007.0210
- Schmidt, V. T., Smith, K. F., Melvin, D. W., Amaral-Zettler, L. A. (2015). Community assembly of a euryhaline fish microbiome during salinity acclimation. *Molecular Ecology*, 24, 2537-2550. doi: 10.1111/mec.13177.
- Schmidt, V., Gomez-Chiarri, M., Roy, C., Smith, K., & Amaral-Zettler, L. (2017). Subtle microbiome manipulation using probiotics reduces antibiotic-associated mortality in fish. *Msystems*, 2, e00133-17. doi: 10.1128/mSystems.00133-17.
- Schneider, A. C. R., Rico, E. P., de Oliveira, D. L., Rosemberg, D. B., Guizzo, R., Meurer, F., & da Silveira, T. R. (2016). *Lactobacillus rhamnosus* GG effect on behavior of zebrafish during chronic ethanol exposure. *BioResearch Open Access*, 5, 1-5. doi:10.1089/biores.2015.0026.
- Sharon, G., Segal, D., Ringo, J. M., Hefetz, A., Ziber-Rosenberg I., & Rosenberg, E. (2010). Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of the Natural Academy of Sciences USA*, 107, 20051-20056. doi:10.1073/pnas.1009906107.

Shin, N.-R., Whon, T. W., & Bae, J. -W. (2015). Proteobacteria: Microbial signature of dysbiosis in gut microbiota. *Trends in Biotechnology*, 33, 496-503. doi: 10.1016/j.tibtech.2015.06.011.

Sikkel, P. C., Cheney, K. L., & Côté, I. M. (2004). In situ evidence for ectoparasites as a proximate cause of cleaning interactions in marine reef fish. *Animal Behavior*, 68, 241-247.

Small, C. M., Milligan-Myhre, K., Bassham, S., Guillemin, K., & Cresko, W. A. (2017). Host genotype and microbiota contribute asymmetrically to transcriptional variation in the threespine stickleback gut. *Genome Biology and Evolution*, 9, 504-520. doi: 10.1093/gbe/evx014.

Smith, C. C., Snowberg, L. K., Gregory Caporaso, J., Knight, R., & Bolnick, D. I. (2015) Dietary input of microbes and host genetic variation shape among-population differences in stickleback gut microbiota. *ISME Journal*, 9, 2515-2526. doi: 10.1038/ismej.2015.64.

Soares, M. C., Cardoso, S. C., Carvalho, T. S., & Maximino, C. (2018). The use of model fish as tools for research the biological mechanisms of cooperative behaviour: a future for translational research concerning social anxiety disorders? *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 82, 205-215. doi: 10.1016/j.pnpbp.2017.11.014.

Soares, M. C. (2017). The neurobiology of cooperation: The cleanerfish swims into the spotlight. *Frontiers in Behavioral Neurosciences*, 11, 191. doi: 10.3389/fnbeh.2017.00191.

Soares, M.C., Bshary, R., Mendonça, R., Grutter, A.S., & Oliveira, R.F. (2012a). Arginine vasotocin regulation of interspecific cooperative behaviour in a cleaner fish. *PLoS One*, 7, e39583. doi:10.1371/journal.pone.0039583.

Soares, M.C., Bshary, R., Cardoso, S.C., Côté, I.M., Oliveira, R.F. (2012b). Face your fears: Cleaning gobies inspect predators despite being stressed by them. *PLoS One*, 7, e39781. doi:10.1371/journal.pone.0039781.

Soares, M.C., Cardoso, S.C., & Côté, I.M. (2007). Client preferences by caribbean cleaning gobies: Food, safety or something else? *Behavioral Ecology and Sociobiology*, 61, 1015-1022. doi: 10.1007/s00265-006-0334-6.

Soares, M.C., Cardoso, S.C., Grutter, A.S., Oliveira, R.F., & Bshary, R. (2014). Cortisol mediates cleaner wrasse switch from cooperation to cheating and tactical deception. *Hormones and Behavior*, 66, 346-350. doi:10.1016/j.yhbeh.2014.06.010.

Soares, M.C., Cardoso, S.C., Malato, J.T., & Messias, J.P.M. (2017a). Can cleanerfish overcome temptation? A selective role for dopamine influence on cooperative-based decision making. *Physiology and Behavior*, 169, 124-129. doi:10.1016/j.physbeh.2016.11.028.

Soares, M. C., Santos, T. P., & Messias, J. P. M. (2017b). Dopamine disruption increases cleanerfish cooperative investment in novel client partners. *Royal Society Open Science*, 4, 160609. doi:10.1098/rsos.160609.

Song, S. J., Lauber, C., Costello, E. K., Lozupone, C. A., Humphrey, G., Berg-Lyons, D., Caporaso, J. G., Knights, D., Clemente, J. C., Nakielny, S., Gordon, J. I., Fierer, N., & Knight, R. (2013). Cohabiting family members share microbiota with one another and with their dogs. *eLife*, 2, e00458. doi:10.7554/eLife.00458

Stecher, B., Chaffron, S., Käppeli, R., Hapfelmeier, S., Friedrich, S., Weber, T. C., Kirundi, J., Suar, M., McCoy, K. D., von Mering, C., Macpherson, A. J., & Hardt, W.-D. (2010). Like will to like: Abundances of closely related species can predict susceptibility to intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathogens*, 6, e1000711. doi: 10.1371/journal.ppat.1000711.

Stewart, A. M., Braubach, O., Spitsbergen, J., Gerlai, R., & Kalueff, A. V. (2014). Zebrafish models for translational neuroscience research: from tank to bedside. *Trends in Neurosciences*, 37, 264-278. <http://doi.org/10.1016/j.tins.2014.02.011>

Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H., Gerlai, R., & Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular Psychiatry*, 20, 2-17. doi:10.1038/mp.2014.128

Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.-N., Kubo, C., & Koga, Y. (2004). Postnatal microbial colonization programs the hypothalamic–pituitary–adrenal system for stress response in mice. *Journal of Physiology*, 558, 263-275. doi: 10.1113/jphysiol.2004.063388.

Sullam, K. E., Rubin, B. E., Dalton, C. M., Kilham, S. S., Flecker, A. S., & Russell, J. A. (2015). Divergence across diet, time and populations rules out parallel evolution in the gut microbiomes of Trinidadian guppies. *ISME Journal*, 9, 1-15. doi: 10.1038/ismej.2014.231.

Sylvain, F.-E., Cheaib, B., Llewellyn, M., Gabriel Correia, T., Barros Fagundes, D., Luis Val, A., & Derome, N. (2016). pH drop impacts differentially skin and gut micro-biota of the Amazonian fish tambaqui (*Colossoma macropomum*), *Scientific Reports*, 6, 32032. doi:10.1038/srep32032.

Tarnecki, A. M., Burgos, F. A., Ray, C. L. & Arias, C. R. (2017). Fish intestinal microbiome: Diversity and symbiosis unravelled by metagenomics. *Journal of Applied Microbiology*, 123, 2-17. doi: 10.1111/jam.13415.

Teitelbaum, A. A., Gareau, M. G., Jury, J., Yang, P. C., & Perdue, M. H. (2008). Chronic peripheral administration of corticotropin-releasing factor causes colonic barrier dysfunction similar to psychological stress. *American Journal of Physiology*, 295, G452-G459. doi: 10.1152/ajpgi.90210.2008.

Theis, K. R., Venkataraman, A., Dycus, J. A., Koonter, K. D., Schmitt-Matzen, E. N., Wagner, A. P., Holekamp, K. E., & Schmidt, T. M. (2013). Symbiotic bacteria appear to mediate hyena social odors. *Proceedings of the Natural Academy of Sciences USA*, 110, 19832-19837. doi: 10.1073/pnas.1306477110.

Tognini, P. (2017). Gut microbiota: A potential regulator of neurodevelopment. *Frontiers in Cellular Neurosciences*, 11, 25. doi:10.3389/fncel.2017.00025.

Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly Review in Biology*, 46, 35-57.

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J.-C., Lynch, J., Grieneisen, L. E., Alt-mann, J., Alberts, S. C., Blekhman, R., & Archie, E. A. (2015). Social networks predict gut microbiome composition in wild baboons. *eLife*, 4, e05224. doi:10.7554/eLife.05224.

van de Wouw, M., Boehme, M., Lyte, J. M., Wiley, N., Strain, C., O'Sullivan, O., Clarke, G., Stanton, C., Dinan, T. G., & Cryan, J. F. (2018). Short-chain fatty acids: microbial metabolites that alleviate stress-induced brain-gut axis alterations. *Journal of Physiology*, 596, 4923-4944. doi:[10.1113/JP276431](https://doi.org/10.1113/JP276431)

Vanuytsel, T., van Wanrooy, S., Vanheel, H., Vanormelingen, C., Verschueren, S., Houben, E., Salim Rasoel, S., Tóth, J., Holvoet, L., Farré, R., Van Oudenhove, L., Boeckstaens, G., Verbeke, K., & Tack, J. (2014). Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. *Gut*, 63, 1293. doi: 10.1136/gut-jnl-2013-305690

Vuong, H. E., Yano, J. M., Fung, T. C., & Hsiao, E. Y. (2017). The microbiome and host behavior. *Annual Review in Neurosciences*, 40, 21-49. doi: 10.1146/annurev-neuro-072116-031347.

Walker, W. A. (2013). Initial intestinal colonization in the human infant and immune homeostasis. *Annals of Nutrition and Metabolism*, 63, 8-15. doi: 10.1159/000354907.

Wang, A. R., Ran, C., Ringø, E. and Zhou, Z. G. (2018). Progress in fish gastrointestinal microbiota research. *Reviews in Aquaculture*, 10, 626-640. doi:[10.1111/raq.12191](https://doi.org/10.1111/raq.12191)

Wang, J., Tao, Q., Wang, Z., Mai, K., Xu, W., Zhang, Y., & Ai, Q. (2017). Effects of fish meal replacement by soybean meal with supplementation of functional compound additives on intestinal

morphology and microbiome of Japanese seabass (*Lateolabrax japonicus*). *Aquaculture Research*, 48, 2186-2197. doi:10.1111/are.13055.

Ward, A. J. W., Holbrook, R. I., Krause, J., & Hart, P. J. B. (2005). Social recognition in sticklebacks: the role of direct experience and habitat cues. *Behavioral Ecology and Sociobiology*, 57, 575. doi: 10.1007/s00265-004-0901-7.

White, J., Mirleau, P., Danchin, E., Mulard, H., Hatch, S. A., Heeb, P., & Wagner, R. H. (2010). Sexually transmitted bacteria affect female cloacal assemblages in a wild bird. *Ecology letters*, 13, 1515-1524. doi: 10.1111/j.1461-0248.2010.01542.x

Wilkins, L. G. E., Rogivue, A., Schütz, F., Fumagalli, L., & Wedekind, C. (2015). Increased diversity of egg-associated bacteria on brown trout (*Salmo trutta*) at elevated temperatures. *Scientific Reports*, 5, 17084. doi: 10.1038/srep17084.

Wilson, A. D., Croft, D. P. & Krause, J. (2014). Social networks in elasmobranchs and teleost fishes. *Fish and Fisheries*, 15, 676-689. doi:10.1111/faf.12046

Wilson, A. D. M., Krause, S., James, R., Croft, D. P., Ramnarine, I. W., Borner, K. K., Clement, R. J. G., & Krause, J. (2014). Dynamic social networks in guppies (*Poecilia reticulata*). *Behavioral Ecology and Sociobiology*, 68,915-925. doi:10.1007/s00265-014-1704-0

Wismer, S., Pinto, A. I., Vail, A. L., Grutter, A. S., & Bshary, R. (2014). Variation in cleaner wrasse cooperation and cognition: potentially caused by developmental environment? *Ethology*, 120, 519–531. doi: 10.1111/eth.12223

Yang, I., Corwin, E. J., Brennan, P. A., Jordan, S., Murphy, J. R., & Dunlop, A. (2016). The infant microbiome: Implications for infant health and neurocognitive development. *Nursing Research*, 65, 76-88. doi: 10.1097/NNR.0000000000000133.

Young, K.A. (2013) The balancing act of captive breeding programmes: Salmon stocking and angler catch statistics. *Fisheries Management and Ecology*, 20, 434-444. doi: 10.1111/fme.12032

Zarkazi, K., Abell, G. C., Taylor, R. S., Neuman, C., Hatje, E., Tamplin, M. L., Katouli, M., Bowman. (2014). Pyrosequencing-based characterization of gastrointestinal bacteria of Atlantic salmon (*Salmo salar* L.) within a commercial mariculture system. *Journal of Applied Microbiology*, 117: 18-27. doi: 10.1111/jam.12514.

Zheng, P., Zeng, B., Zhou, C., Liu, M., Fang, Z., Xu, X., Zeng, L., Chen, J., Fan, S., Du, X., Zhang, X., Yang, D., Yang, Y., Meng, H., Li, W., Melgiri, N. D., Licionio, J., Wei, H., & Xie, P. (2016). Gut microbiome remodeling induces depressive-like behaviors through a pathway mediated by the host's metabolism. *Molecular Psychiatry*, 21, 786-796. doi: 10.1038/mp.2016.44.