

Review

Understanding Root Rot Disease in Agricultural Crops

Bruce A. Williamson-Benavides¹ and Amit Dhingra^{1,2,*}

¹ Molecular Plant Sciences, Washington State University, Pullman, WA - 99164; b.williamsonbenavid@wsu.com

² Department of Horticulture, Washington State University, Pullman, WA - 99164; adhingra@wsu.com

* Correspondence: adhingra@wsu.com

Abstract: Root rot diseases remain a major global threat to the productivity of agricultural crops. They are usually caused by more than one type of pathogen and are thus often referred to as a root rot complex. Fungal and oomycete species are the predominant participants in the complex, while bacteria and viruses are also known to cause root rot. Incorporating genetic resistance in cultivated crops is considered as the most efficient and sustainable solution to counter root rot; however, resistance is often quantitative in nature. Several genetics studies in various crops have identified quantitative trait loci associated with resistance. With access to whole genome sequences, the identity of the genes within the reported loci is becoming available. Several of the identified genes have been implicated in pathogen response. However, it is becoming apparent that at the molecular level, each pathogen engages a unique set of proteins to either infest the host successfully or be defeated or contained in doing so. In this review, a comprehensive summary of the genes and potential mechanisms underlying resistance or susceptibility against the most investigated root rots of important agricultural crops is presented.

Keywords: Fungi, oomycetes, resistance, susceptibility, molecular mechanisms, quantitative trait loci, mapping

1. Introduction

Root rots have a significant impact on global crop production [1]. Depending on the causal agent, host susceptibility, and the environmental conditions, crop losses can range from slightly above the economic threshold to losing complete fields. Interestingly, legumes seem to be the most common host for these pathogens. However, monocots and dicots, cereals and legumes, fruit trees, and tubers also fall prey to root rots.

Fungi and oomycetes most commonly cause root rot disease. However, bacteria and even viruses can be the causal agents [2–8]. Due to more than one pathogen's involvement, the disease is commonly referred to as root rot complex. Some classic examples include the black root rot of strawberry attributed to *Pythium*, *Fusarium*, and *Rhizoctonia* pathogens [9–11], and the pea root rot complex caused by *A. euteiches*, *F. oxysporum*, *F. solani*, *F. avenaceum*, *Mycosphaerella pinodes*, *Pythium* spp., *R. solani*, and *Phytophthora* spp. [12–15].

Unless the root rot complex affects seed germination, the root-specific symptoms go unnoticed or are not visible. If symptoms appear aboveground, the plants usually fail to recover. Some of the symptoms associated with root rots include browning and softening of root tips, root lesions that vary in size and color (reddish, brown, and black), yellowing and wilting of leaves, stunted plant growth, reduced yield, and loss of crop [1,3,16–19]. Selected root rot pathogens can also cause post-harvest rots in beets, potato, and sweet potato. The proliferation of root rot pathogens is favored by moderate to high soil moisture, poor drainage conditions, soil compaction, the optimal temperature for pathogen growth, mono-cropping, and other factors that contribute to plant stress [1,20–22]. The unpredictable climatic conditions portend an increase in mean temperatures and other natural calamities such as droughts, floods, and storms. These conditions are expected to inflict constant stress on crops, which is expected to favor the increased activity of root rot pathogens [23–25].

Cultural, physical, biological, and chemical control methods have been used as management strategies to control root rot disease. However, to date, these strategies have only been partially successful. Most of the root rot pathogens are distributed globally, and some species can survive up to 10 years in the soil. Several root rot pathogens are host-specific; however, some have a wide range of hosts. Therefore, crop rotation is not a practical strategy [16,26]. Chemical control is often inefficient due to these pathogens' soil-borne nature and is not the most sustainable option as it also impacts beneficial microbes.

There is a critical need to understand the genetic basis of root rots and incorporate the information in breeding strategies to develop root rot-resistant crops. The current understanding of plant molecular defense responses is derived primarily from studies using foliar pathosystems [27]. Specific and unique genetic and molecular aspects of the host-pathogen interactions in the roots have been unraveled in the past few decades. This review summarizes the different groups of root rot species that affect crops, instances of host resistance and susceptibility, and the genes and proposed molecular mechanisms underlying host-pathogen interactions.

2. Common causal agents of root rots

2.1 Fungi

Fungi represent one of the most predominant root rot causing agents. The most studied and problematic fungal root rots are the *Rhizoctonia* root rot, *Fusarium* root rot, *Phoma* root rot, and Black root rot. They account for incalculable yield losses across agricultural and horticultural crops. These fungal pathogens also impact the wood industry.

Rhizoctonia root rot is caused by the soilborne fungus *Rhizoctonia solani* (Table 1). *R. solani* is a species complex because of the many related but genetically distant isolates. Isolates are classified into 12 anastomosis groups (AGs) based on their hyphal incompatibility and their host specificity (Table 1) [28]. For instance, AG2-1 and AG4 are associated with stem and root rot diseases in dicots such as Brassicaceae species, while AG8 causes root rot in monocots [19,29]. In general, the first four AGs (AG-1, -2, -3, and -4) cause important diseases in plants worldwide, whereas the remaining AGs (AG-5, -6, -7, -8, -9, -10, -11, -12) are less destructive and generally have a restricted geographic distribution (Table 1) [18].

Symptoms of *R. solani* vary among species, but it primarily affects underground tissues (seeds, hypocotyls, and roots); however, it can also infect above-ground plant parts such as pods, fruits, leaves, and stems. Pre- and/or post-emergence damping off is the most common symptom of *R. solani*. Surviving seedlings can often develop reddish-brown lesions (cankers) on stems and roots. This pathogen can occasionally infect fruit and leaf tissue near or on the soil surface [18,19,30]. *R. solani* is responsible for high yield losses in many crops, and some of the noteworthy examples are highlighted in Table 1.

Table 1. Crop species affected by *Rhizoctonia solani* root rot. A summary of reported resistant genotypes and *R. solani* anastomosis groups (AGs) per host species is presented.

Host	Disease name	AG	Resistance	References
Barley (<i>Hordeum vulgare</i>)	Root rot	5	No reports	[31–34]
		8	Moderate and high levels of resistance. High resistant transgenic lines	
Bean (<i>Phaseolus vulgaris</i>)	Root rot	2	Moderate and high levels of resistance	[35–38]
		4	Moderate and high levels of resistance	
		5	Moderate levels of resistance	

Cabbage (<i>Brassica oleracea</i> var. <i>capitata</i> L.)	Wirestem	2	Moderate levels of resistance	[39–41]
		4	Moderate levels of resistance	
Carrot (<i>Daucus carota</i>)	Crown and brace root rot	1	Moderate levels of resistance	[42–44]
		2	Moderate levels of resistance	
		4	Moderate levels of resistance. Moderately resistant transgenic line	
Faba bean (<i>Vicia faba</i>)	Root rot and stem canker	2	Moderate levels of resistance	[45–47]
		4	Moderate and high levels of resistance	
		5	Moderate levels of resistance	
Lettuce (<i>Lactuca sativa</i>)	Bottom rot	1	No reports	[48]
Maize (<i>Zea mays</i>)	Banded leaf and sheath blight disease	1	Moderate and high levels of resistance	[47,49–52]
		2	Moderate levels of resistance	
Oat (<i>Avena sativa</i>)	Root rot, Bare patch	8	Moderate levels of resistance	[33]
Oilseed rape (<i>Brassica napus</i>)	Root rot and damping-off	2	Moderate levels of resistance	[53–55]
		4	No reports	
Onion (<i>Allium cepa</i>)	Stunting	4	No reports	[56,57]
		8	Moderate levels of resistance	
Pea (<i>Pisum sativum</i>)	Root rot	4	Moderate levels of resistance	[58,59]
Potato (<i>Solanum tuberosum</i>)	Black scurf and Stem canker	2	Moderate levels of resistance	[60–63]
		3	Moderate and high levels of resistance	
Rice (<i>Oryza sativa</i>)	Sheath blight	1	Moderate levels of resistance. Moderate levels of resistant transgenic lines.	[64–67]
Rye (<i>Secale cereale</i>)	Root rot, Bare patch	8	Moderate levels of resistance	[33,68]
Soybean (<i>Glycine max</i>)	Root rot	1	Moderate levels of resistance	[69–72]
		2	Moderate and high levels of resistance	
		4	Moderate and high levels of resistance	
Sugarbeet (<i>Beta vulgaris</i>)	Root rot	2	Moderate levels of resistance	[73–75]
Tomato (<i>Solanum lycopersicum</i>)	Foot and Root rot	3	Moderate levels of resistance	[76–78]
	Foot and Root rot	4	Moderate levels of resistance	
Triticale (<i>Triticale hexaploide</i>)	Root rot, Bare patch	8	Moderate levels of resistance	[33]
Wheat (<i>Triticum aestivum</i>)	Root rot, Bare patch	8	Moderate levels of resistance	[33,34,79,80 1]

The genus *Fusarium* constitutes a sizeable monophyletic group of several hundred species that includes agriculturally important plant pathogens, endophytes, saprophytes, and emerging pathogens of clinical importance [81]. The most important *Fusarium* species causing root rot is *F. solani* (Table 2). Other *Fusarium* spp that can cause root rot are *F. avenaceum*, *F. graminearum*, *F. culmorum*, *F. verticillioides*, and *F. pseudograminearum* (Table 2). However, the latter five are mainly associated with head blight or ear mold in different small-grain cereals [82].

F. solani (sexual morph *Nectria haematococca*) is a filamentous fungus of significant agricultural importance. This species is classified as *F. solani* species complex (FSSC) because it contains 60 phylogenetically distinct species [81,83]. Most of the studies on FSSC have been carried out while investigating host-pathogen interactions. Therefore, the group has been subdivided into *formae speciales* (f. sp.) based on host specificity [81].

Phytopathogenic FSSC species include some of the most economically significant plant pathogens associated with root rots and vascular wilts in over 100 crops [84]. Some of the most important *F. solani* that cause problems in agriculture are presented in Table 2.

FSSC causes foot and/or root rot of the infected host plant, and the degree of necrosis correlates with the severity of the disease [81]. Symptoms on above-ground portions may manifest as wilting, stunting, and chlorosis or lesions on the stem and/or leaves. However, symptoms vary depending on the specific FSSC pathogen and plant host involved [81].

Other *Fusarium* species that cause root rots of minor economic importance are *F. chlamydosporum*, which infects coleus and other ornamentals [85,86]. *F. oxysporum* can cause root rot in the Cactaceae family members [87], as well as stem and root rot in melons [88].

Table 2. Main crop species affected by *Fusarium* root rot. Table shows reported resistant genotypes per host species and host range and distribution per pathogen species.

Fungi spp	Plant attacked	Disease Name	Resistance reported in the literature	Host Range	Distribution	References
<i>Fusarium avenaceum</i>	Wheat (<i>Triticum aestivum</i>)	Head blight	Moderate levels of resistance	Wide range with over 200 species, including pulses, cereals, canola (<i>Brassica napus</i>), flax (<i>Linum</i> spp), and alfalfa (<i>Medicago truncatula</i>)	Worldwide	[89–91]
	Barley (<i>Hordeum vulgare</i>)	Head blight	Moderate levels of resistance			[92,93]
	Pea (<i>Pisum sativum</i>)	Root rot	Moderate levels of resistance			[94–96]
	Alfalfa (<i>Medicago sativa</i>)	Root and crown rot	Moderate levels of resistance			[97]
	Clover (<i>Trifolium subterraneum</i>)	Root rot	Moderate levels of resistance			[98]
<i>F. culmorum</i>	Wheat (<i>Triticum aestivum</i>)	Root rot and head blight	Moderate levels of resistance	Wide range of host plants, including rye (<i>Secale cereale</i>), corn (<i>Zea mays</i>), sorghum (<i>Sorghum</i> spp.), various grasses, sugar beet (<i>Beta vulgaris</i>), bean (<i>Phaseolus</i> spp.), pea (<i>Pisum sativum</i>), asparagus (<i>Asparagus</i> spp), hop (<i>Humulus lupulus</i>), strawberry (<i>Fragaria × ananassa</i>), and potato (<i>Solanum tuberosum</i>)	Worldwide	[89,90,99–101]
	Barley (<i>Hordeum vulgare</i>)	Head blight	Moderate levels of resistance			[102,103]
	Oat (<i>Avena sativa</i>)	Head blight	Moderate levels of resistance			[104]
<i>F. graminearum</i>	Barley (<i>Hordeum vulgare</i>)	Head blight	Moderate levels of resistance	Wide range, especially many species of cereals and grasses such as oat (<i>Avena sativa</i>), rice (<i>Oryza</i>), cucumber (<i>Cucumis sativus</i>), soy (<i>Glycine max</i>), tomato (<i>Lycopersicon</i> spp.), alfalfa	Worldwide	[102,105–107]
	Wheat (<i>Triticum aestivum</i>)	Head blight	Moderate levels of resistance			[102,108–112]
	Soybean (<i>Glycine max</i>)	Pod blight and root rot	High levels of resistance			[113–115]

	Maize (<i>Zea mays</i>)	Ear mold and root rot	Moderate levels of resistance	(<i>Medicago truncatula</i>), sorghum (<i>Sorghum</i> spp.)		[116–120]
<i>F. pseudograminearum</i>	Wheat (<i>Triticum aestivum</i>)	Crown rot	Moderate levels of resistance	All major winter cereals barley (<i>Hordeum vulgare</i>), oats (<i>Avena sativa</i>) and grass genera, such as <i>Phalaris</i> , <i>Agropyron</i> and <i>Bromus</i>	All areas cultivated with wheat and barley	[121,122]
	Barley (<i>Hordeum vulgare</i>)	Crown rot	Moderate levels of resistance			[123,124]
<i>F. solani</i> f. sp. <i>batatas</i>	Sweetpotato (<i>Ipomoea batatas</i>)	Storage root	Moderate levels of resistance	Not well known	China	[125]
<i>F. solani</i> f. sp. <i>glycines</i>	Soybean (<i>Glycine max</i>)	Sudden death syndrome	Moderate levels of resistance	Broad range including bean (<i>Phaseolus</i> spp.), Soybean (<i>Glycine max</i>), alfalfa (<i>Medicago truncatula</i>), clover (<i>Trifolium</i> spp), pea (<i>Pisum sativum</i>), corn (<i>Zea mays</i>), wheat (<i>Triticum</i> spp), ryegrass (<i>Lolium</i> spp), pigweed (<i>Amaranthus</i> spp), and lambsquarters (<i>Chenopodium album</i>)	America, Asia, and Africa	[126–128]
<i>F. solani</i> f. sp. <i>phaseoli</i>	Bean (<i>Phaseolus vulgaris</i>)	Root rot	Moderate ¹ and high ² levels of resistance	Pea (<i>Pisum sativum</i>)	All continents except Australia	[129–131]
<i>F. solani</i> f. sp. <i>pisi</i>	Pea (<i>Pisum sativum</i>)	Root rot	Moderate levels of resistance	Chickpea (<i>Cicer arietinum</i>), clover, soybean, as well as several other non-legume hosts, such as ryegrass (<i>Lolium</i> spp), potato (<i>Solanum tuberosum</i>) ginseng (<i>Panax ginseng</i>) and mulberry tree (<i>Morus alba</i>)	Worldwide	[132,133]
<i>F. verticillioides</i>	Maize (<i>Zea mays</i>)	Root and ear rot	Moderate levels of resistance	Wide range of hosts such as rice (<i>Oryza sativa</i>), sorghum (<i>Sorghum</i> spp), soybean (<i>Glycine max</i>), alfalfa (<i>Medicago truncatula</i>), bean (<i>Phaseolus</i> spp), wheat (<i>Triticum</i> spp), ryegrass (<i>Lolium</i> spp)	Worldwide	[134]

Some species belonging to the genus *Phoma* are known to cause root rots [135] (Table 3). Although their economic impact is not as significant as the root rots caused by the fungi species mentioned above, considerable yield losses have been reported in alfalfa, sugar beet, corn, and onion (Table 3). *Thielaviopsis basicola* is another global soil-borne fungus that causes black root rot disease. This disease is characterized by necrotic lesions on various parts of the host roots [136–138]. Most reports highlighted the effect of this root rot

in cotton (Table 3). Crops such as legumes, tobacco, carrot, citrus, groundnut, and chicory have also been reported to be impacted.

Some other fungal root rot pathogens affect crops with less frequency. These pathogens include *Aspergillus* spp., *Alternaria* spp., *Curvularia* spp., *Rhizopus* spp. and *Penicillium* spp., in fruit trees [139]; *Rigidoporus lignosus* and *Phellinus noxius* in rubber trees [140]; and *Macrophomina phaseolina* in chickpeas [141]. Armillaria root rot is a threat in apples, walnut, kiwi, and grapes [142,143]. The ascomycete *Rosellinia necatrix* is known to cause white root rot in trees such as apple in the Kashmir valley [144] and avocado in the Mediterranean [145].

Table 3. Main crop species affected by fungus *Phoma* and *Thielaviopsis basicola* root rot. A summary of reported resistant genotypes per host species and host range and distribution per pathogen species, is presented.

Fungi spp	Plant attacked	Disease Name	Resistance reported in the literature	Host range	Distribution	References
<i>Phoma betae</i>	Sugar beet (<i>Beta vulgaris</i>)	Crown and root rot	Moderate levels of resistance	Different varieties of <i>Beta vulgaris</i> such as table beet, sugar beet, Swiss chard	World-wide distribution, found in all beet-growing areas.	[146]
<i>P. terrestris</i> (<i>Setophoma terrestris</i>)	Corn (<i>Zea mays</i>)	Red root rot	Moderate levels of resistance	45 genera including cereals, vegetables and grasses such as soybean (<i>Glycine max</i>), pea (<i>Pisum sativum</i>), sugarcane (<i>Saccharum</i> spp), oats (<i>Avena sativa</i>), barley (<i>Hordeum vulgare</i>), wheat (<i>Triticum</i> spp), cucumber (<i>Cucumis sativus</i>), tomato (<i>Lycopersicon</i> spp.), pepper (<i>Capsicum annuum</i>)	World-wide distribution	[147]
	Onion (<i>Allium cepa</i>)	Pink root rot	Moderate levels of resistance			[148,149]
<i>P. sclerotioides</i>	Alfalfa (<i>Medicago sativa</i>)	Brown root rot	Moderate levels of resistance	Wheat (<i>Triticum</i> spp), barley (<i>Hordeum vulgare</i>), and oat (<i>Avena sativa</i>) .	North America and Australia	[150,151]
<i>Thielaviopsis basicola</i>	Cotton (<i>Gossypium herbaceum</i>)	Black root rot	Moderate levels of resistance. Moderate levels of resistance in transgenic lines	Wide range of hosts, plants from at least 15 families including horseradish (<i>Armoracia rusticana</i>), carrot (<i>Daucus carota</i>), strawberry (<i>Fragaria x ananassa</i>), tomato (<i>Lycopersicon</i> spp.), bean (<i>Phaseolus</i> spp), pea (<i>Pisum sativum</i>)	World-wide distribution	[152–155]
	Tobacco (<i>Nicotiana</i> spp)	Black root rot	Moderate levels of resistance			[156,157]

2.2 Oomycetes

Oomycetes resemble fungi in their growth habits and nutritional strategies. However, they are evolutionarily distant from fungi and belong to the kingdom Stramenopiles

[158]. Oomycetes are a large group of terrestrial and aquatic eukaryotic organisms. They are dispersed via zoospores, generate thick-walled sexual oospores, possess cellulose in their cell walls, are vegetatively diploid, have heterokont flagellae (one tinsel and one whiplash), and have tubular mitochondrial cristae [159].

The terrestrial oomycetes are mainly parasites of the vascular plants and include several important pathogens such as *Aphanomyces* spp., *Pythium* spp., and *Phytophthora* spp that cause root rot (Table 4). These oomycetes appear to have extraordinary genetic flexibility, enabling them to adapt rapidly and overcome chemical control measures and genetic resistance in host plant [160–162].

Among *Aphanomyces* spp that cause root rots, *A. cochlioides* and *A. euteiches* cause significant agricultural concerns (Table 4). *A. cochlioides* causes damping off and chronic root rot in sugar beet, spinach, cockscomb, among other species of Chenopodiaceae and Amaranthaceae [163,164] (Table 4). Due to the extended prevalence of the disease in soil and severity in the field, outbreaks of *A. cochlioides* root rot have become a severe problem in many sugar beet growing areas [165]. *A. cochlioides* root rot, when severe, can lead to death and drastically reduced recoverable sugar per ton [166]. Little is known about the genetic basis of resistance to *A. cochlioides* root rot. Still, several sugar beet genotypes have been released and also used in the development of molecular markers that have been found associated with disease resistance genes [165,167,168] (Table 4).

Table 4. Main oomycete species that cause root rot disease in crop species. A summary of reported symptoms and resistant genotypes per host species and pathogen-host range and distribution is presented.

Oomycetes spp	Plant attacked	Reported symptoms	Resistance reported in the literature	Host range	Distribution	References
<i>Aphanomyces cochlioides</i>	Sugar beet (<i>Beta vulgaris</i>)	Root rot and damping off. Infected hypocotyl and root rapidly turn black and shrink to a dark, slender thread at junctures of lateral roots and/or at the root tip. The disease can occur from the seedling stage until harvest. Undersized plants with yellowed lower leaves. Severely infected plants die. Postharvest reduction of sugar yield	High levels of resistance	Spinach (<i>Spinacia oleracea</i>), and several wild species of Beta, including <i>B. maritima</i> and <i>B. patellaris</i>	Across all sugar beet plantations	[165–169]
<i>Aphanomyces euteiches</i>	Pea (<i>Pisum sativum</i>)	Damping off and root rot. Root tissue becomes honey-brown or blackish-brown. Reduced nodulation. Chlorosis of cotyledons and necrosis of epicotyls or hypocotyls. Chlorosis, necrosis, and wilting of the foliage. Severely infected plants die.	High levels of partial resistance	Faba bean (<i>Vicia faba</i>), red clover (<i>Trifolium pratense</i>), white clover (<i>Trifolium repens</i>), <i>Medicago truncatula</i> , lentil (<i>Lens culinaris</i>)	Asia, Europe, Oceania, North America	[130,170–174]
	Alfalfa (<i>Medicago sativa</i>)		High levels of resistance			
	Bean (<i>Phaseolus vulgaris</i>)		Levels of partial and complete resistance			

<i>Phytophthora citrophthora</i>	Citrus spp.	Serious gummosis of citrus trees, root rot, stem necrosis, canker, fruit rot, twig blight, and seedling blight.	Tolerant transgenic C. <i>sinensis</i> . Partial levels of resistance in citrus rootstocks. High levels of resistant citrus rootstocks	88 genera including: kiwifruit (<i>Actinidia deliciosa</i>), watermelon (<i>Citrullus lanatus</i>) strawberry (<i>Fragaria ananassa</i>), walnut (<i>Juglans regia</i>), apricot (<i>Prunus armeniaca</i>), sweet cherry (<i>P. avium</i>), almond (<i>P. dulcis</i>), potato (<i>Solanum tuberosum</i>), cocoa (<i>Theobroma cacao</i>), blueberries (<i>Vaccinium</i>)	Cosmopolitan	[175–180]
<i>Phytophthora nicotianae</i>	Citrus spp.	Symptoms vary per host. Damping-off, crown rot, leaf blight, fruit rot. Occasionally, it attacks aerial parts of the plant and can cause brown rot of fruit.	Tolerant transgenic C. <i>limonia</i> . Partial levels of resistance in citrus rootstocks	255 genera in 90 families. including tobacco, citrus, cotton, and orchids	Cosmopolitan	[176,177,181–185]
	Tomato (<i>Solanum lycopersicum</i>)		Partial levels of resistance			
<i>Phytophthora cactorum</i>	Apple (<i>Malus domestica</i>)	Damping off of seedlings, fruit rot, leaf, stem and root rot, collar and crown rot, stem canker.	Partial levels of resistance	154 genera of vascular plants in 54 families	Cosmopolitan	[177,186–191]
	Strawberry (<i>Fragaria ananassa</i>) ×	Infected apple trees present reduced tree vigor and growth, chlorosis of leaves, and eventual collapse or death of the tree. Infected trees may decline slowly over one or more years, or they may collapse and die rapidly after resuming growth in the spring.	Partial levels of resistance			

<i>Phytophthora cinnamomi</i>	Avocado (<i>Persea americana</i>)	Root rot, heart rot, wilt. Primary infection at the feeder roots, resulting in a brownish black and brittle appearance. There is almost no progression into the larger roots. Root rot leads to death of the feeder roots which results in insufficient water and nutrient uptake, eventually leading to tree mortality	Partial levels of resistance	266 genera in 90 families, commonly hardwood trees.	Cosmopolitan	[177,192]
<i>Phytophthora fragariae</i>	Strawberry (<i>Fragaria</i> × <i>ananassa</i>)	Red stele or red core root rot. Symptoms also include wilting of leaves, reduced flowering, stunting, and bitter fruit	Partial levels of resistance	-	Asia, Australia, New Zealand, Europe, North America	[177,193–196]
	Raspberry (<i>Rubus</i> spp)		Partial and high levels of resistance			
<i>Phytophthora sojae</i>	Soybean (<i>Glycine max</i>)	Root and stem rot; pre- and post-emergence damping-off, seedling wilt, seedling blight. Symptoms may also include necrosis of hypocotyls, chlorosis of leaves, and plant may turn reddish-orange to orange-brown in color.	Partial levels of resistance	Lupine (<i>Lupinus</i> spp.); also reported in six other genera in five families	Australia, North America (Canada, USA), South America (Chile), Asia (Korea, China) and New Zealand	[177,197,198]
<i>Phytophthora capsici</i>	Pepper (<i>Capsicum annuum</i>)	Fruit, stem, and root rot. Symptoms may include also seedling damping-off and leaf wilt. Leaf tissue becomes wilted, light green or gray-green, and later tan to white. Fruit rots are irregular in shape and olive green or light green in color. Infected seeds are brown and shriveled.	Partial levels of resistance	51 genera in 28 families, including tomatoes (<i>Lycopersicon esculentum</i>), other Solanaceae spp, Macadamia spp., cacao (<i>Theobroma cacao</i>)	Cosmopolitan	[177,199–201]
<i>Phytophthora medicaginis</i>	Alfalfa (<i>Medicago sativa</i>)	Root rot, damping-off of seedlings. Older plants are also infected. Reddish-brown or black root lesions. Mature plants exhibit chlorosis, desiccation of	High levels of resistance	Sainfoin (<i>Onobrychis viciifolia</i>), chickpea (<i>Cicer arietinum</i>),	Cosmopolitan, throughout the range of the host	[177,202–206]
	Soybean (<i>Glycine max</i>)		Partial levels of resistance			

	Chickpea (<i>Cicer arietinum</i>)	foliage, and reduced growth but may also collapse.	Partial levels of resistance	cherry (<i>Prunus mahaleb</i>)		
<i>Pythium ultimum</i>	Cucumber (<i>Cucumis sativus</i>)	Damping off and root rot. Disease can manifest as seed rot, preemergence and postemergence damping-off, root rot, dark brown or reddish roots, and sunken lesions on lower hypocotyls. Plants are stunted or chlorotic. Severe infections often lead to plants collapse and dead. Root tips of diseased plants appear as brown.	No records	Cabbage (<i>Brassica oleracea</i> var. <i>capitata</i>), carrot (<i>Daucus carota</i>), melon (<i>Cucumis melon</i>), wheat (<i>Triticum aestivum</i>)	Cosmopolitan	[207–211]
	Sugar beet (<i>Beta vulgaris</i>)		Partial levels of resistance			
	Sorghum (<i>Sorghum bicolor</i>)		No records			
	Tomato (<i>Solanum lycopersicum</i>)		No records			
	Soybean (<i>Glycine max</i>)		Partial levels of resistance			
	Bean (<i>Phaseolus vulgaris</i>)		High levels of resistance			
<i>Pythium irregulare</i>	Soybean (<i>Glycine max</i>)	Pre- and post-emergence damping off of seedlings (greenhouse) and root rot (field) of older plants. Contaminated seeds and seedlings will quickly turn brown and soft before decomposing. The roots become brown, may appear water soaked, and the tips necrotic. Water soaked areas of the hypocotyl observed above the soil may develop which eventually dry to a brown lesion, foliage may turn chlorotic or a greenish-grey and/or wilt. Plants may become stunted or collapse.	Moderate to high levels of resistance	Over 200 host species including pineapple (<i>Ananas comosus</i>), cereals, grasses, celery (<i>Apium graveolens</i>), pepper (<i>Capsicum annuum</i>), pecan trees (<i>Carya illinoensis</i>), Citrus spp, strawberries (<i>Fragaria × ananassa</i>), lentils (<i>Lens culinaris</i>), corn (<i>Zea mays</i>), soybean (<i>Glycine max</i>), cucumber (<i>Cucumis sativus</i>), onion (<i>Allium cepa</i>), carrot (<i>Daucus carota</i>) and a number of floricultural crops	Cosmopolitan in greenhouses and field systems	[98,212–214]
	Clover (<i>Trifolium subterraneum</i>)		Moderate resistance			

<i>Pythium aphanidermatum</i>	Lettuce (<i>Lactuca sativa</i>)	Causes root and stem rots, as well as pre- and post- emergence damping off. It causes blights of grasses and fruit. Symptoms begin at the root tips. Root growth is inhibited, and roots are blackened, mushy and rotten. The rotten part of the roots may slough off exposing the inner root core. It causes wilting, loss of vigor, stunting, chlorosis and leaf drop. If the lesion expands to encircle the stem, the plant collapses. Mature tissue can be infected and destroyed. Beets and other fleshy plant organs are susceptible to rot in the field and during storage.	No reports	Broad host range, including cotton (<i>Gossypium</i> spp), grasses, papaya (<i>Carica papaya</i>), , cereals, Brassica species and beans (<i>Phaseolus vulgaris</i>)	Cosmopolitan in greenhouses and field systems	[215–219]
	Cucumber (<i>Cucumis sativus</i>)		No reports			
	Pepper (<i>Capsicum annuum</i>)		Moderate levels of tolerance			
	Tomato (<i>Solanum lycopersicum</i>)		No reports			
	Soybean (<i>Glycine max</i>)		High resistance			
	Sugar beet (<i>Beta vulgaris</i>)		Partial levels of resistance			

A. euteiches causes seedling damping-off and root rot disease in a variety of field crops worldwide. It first causes softened and water-soaked roots that result in stunted seedlings and yellow leaves. Then the pathogen spreads rapidly, and the cortical tissue and the delicate branches of feeding rootlets are destroyed. In severe cases, the plants collapse and die (Table 4) [16,17].

Main yield losses caused by *A. euteiches* are observed in legumes. *A. euteiches* is the most devastating pathogen of pea in several countries, with yearly losses that average 10% to 80% each year [16,172]. Significant yield losses have also been reported in alfalfa [220], clover [221], fava beans [222], and lentils [223]. Like *A. cochlioides*, *A. euteiches* is a strictly soil-borne pathogen that may survive up to 10 years in soil [26], and no efficient chemical control is currently available. The only way to control the disease is to avoid cultivating legumes in infested fields for ten years [16,26]. To date, no fully resistant pea cultivars have been developed. In 2012, eight pea germplasm lines carrying partial resistance to *A. euteiches* and acceptable agronomic characteristics were released for fresh, frozen, and dry pea production (Table 4) [173]. Resistant lines of alfalfa are also available to growers [170,171].

Pythium genus possesses over two hundred described species, and at least ten *Pythium* spp. cause *Pythium* damping-off and root rot in various legumes and monocots (Table 4). *Pythium* root rot infection symptoms are similar to other root rots; however, only the root tips show necrosis during early infection [224]. It is also typical of this pathogen that the entire primary root's rapid black rot moves up to the stem [29]. *P. ultimum* and *P. irregulare* have been reported as the most ubiquitous pathogens in this group, regularly found in the field, sand, pond and stream water, and decomposing vegetation [29,224]. In the greenhouse industry, the three most commonly encountered root-rot *Pythium* species are *P. ultimum*, *P. irregulare*, and *P. aphanidermatum* [224].

P. ultimum is a principal causal agent of seed decay and pre-emergence/postemergence damping-off in bean [211,225]. A study found that only cream-seeded beans exhibited high resistance levels, while all the white-seeded accessions were susceptible [211]. *P. irregulare* is often the most common pathogenic species of *Pythium* in soybean farms

[226,227]. These studies found that *P. irregulare*, compared to other *Pythium* spp, had the highest pathogenicity levels in soybean. A total of sixty-five soybean genotypes were evaluated for resistance to *P. irregulare*, and about a third showed moderate to high levels of resistance (Table 4) [212].

Pythium aphanidermatum is the predominant pathogen in greenhouse-grown cucumber. It can rapidly spread through zoospores in a recirculating nutrient film culture system [228–230]. *P. aphanidermatum* is also one of the most critical sugar beet diseases in temperate areas with high soil moisture levels. In addition to direct damage to plants in the fields, this pathogen also causes root rot in storage [231]. Several sugar beet genotypes have been found to be partially resistant to *P. aphanidermatum* root rot (Table 4) [216,231]. Other economically important plant species affected by *Pythium* spp are parsnip and parsley [232], wheat [233], and sugarcane [234]. *P. aphanidermatum* and *P. ultimum* mediated root rot has been reported in ornamental plants [235].

Phytophthora spp. represent more than 100 species, and most of them have been classified as aggressive plant pathogens that cause extensive losses in agricultural and horticultural crops [236]. *Phytophthora* means “plant destroyer,” a name coined in the 19th century when the potato disease caused by *Phytophthora infestans* (causal agent of potato late blight) set the stage for the Great Irish Famine [237]. General symptoms of *Phytophthora* infection include wilting, yellow or sparse foliage, and branch dieback [3].

P. citrophthora is the most wide-spread oomycete pathogen in citrus growing areas accounting for millions of dollars in crop losses annually [238,239]. In Citrus, *P. citrophthora* causes gummosis, root rot, and during winter, it causes brown rot of the fruit. *P. nicotianae* also causes foot rot and root rot in Citrus. This pathogen is more commonly found in subtropical areas of the world [239,240]. Nursery- and large-trees can be rapidly girdled and killed by both pathogens [239].

P. cactorum is another pathogen capable of producing high yield losses in fruit trees. In Canada and certain US regions, it has been identified as the most important cause of crown rot of apple [186]. The use of rootstocks resistant to *P. cactorum* and other *Phytophthora* spp has been considered a good management practice. Sources with different resistance levels have been identified since 1959; however, no highly resistant rootstocks have been found (Table 4) [186,188,190,191]. In strawberry, *P. cactorum* crown rot is also considered a disease of commercial importance worldwide. Several strawberry lines with partial resistance to the disease have been identified recently (Table 4) [187,189].

P. cinnamomic is another *Phytophthora* pathogen that affects fruit trees. This disease causes problems mainly in avocado. On average, this disease leads to an annual loss of 10% of the world's avocado crop [241]. This disease has eliminated commercial avocado production in many Latin American regions and is the major limiting factor of production in Australia, South Africa, and California [192]. Therefore, the development of resistant *P. cinnamomic* rootstocks is currently one of the most important goals for the avocado industry. *P. cinnamomi* is also a problem for pineapple production in Australia since it reduces plant growth and yield. *P. cinnamomi* root rot may result in total loss of this crop, especially for the new pineapple hybrids, which are susceptible to *P. cinnamomic* [242]. Other *Phytophthora* species that significantly affect agriculturally important crops are *P. fragariae*, *P. sojiae*, *P. capsici*, and *P. medicaginis* (Table 4).

2.3 Bacteria and Viruses

Bacteria are not a significant root rot causing agent. However, these root rots can cause substantial economic damage. Main yield losses occur in potato and sweet potato. Considerable losses have also been reported for green peppers and Chinese cabbages (*Brassica campestris* subsp. *pekinensis*) [2–6,243].

Bacteria commonly gain entry into the host through wounds in the roots [3,244]. They may also be able to gain access through the leaves, where bacteria develop under aerobic conditions in the aerial parts or migrate to the bulb, rhizome, or directly infect the storage organ [3,243]. These bacteria are characterized by the production of large quantities of extracellular enzymes that include pectinases, cellulases, proteases, and xylanases, which

digest the host cell walls and cause disease [245,246]. From this set of enzymes, the pectinases are believed to be the most important in pathogenesis, causing tissue maceration and cell death [243]. The ability to produce a broader range of enzymes more rapidly and larger quantities than pectolytic saprophytic microorganisms enables bacterial root rots to invade living plants more readily [243,247].

The number of identified bacterial root rot pathogens belong to two genera, *Pectobacterium* and *Dickeya*. Four pathogens, *Pectobacterium carotovorum* subsp. *Carotovorum* (formerly *Erwinia carotovora* subsp. *Carotovora*), *P. atrosepticum* (formerly *E. carotovora* subsp. *Atroseptica*), *Dickeya dianthicola*, and *D. solani* (both previously known as *E. chrysanthemi*), cause wilt and rot diseases in monocot and dicot plants worldwide. Of these pathogens, *P. carotovorum* subsp. *Carotovorum* has the broadest host range worldwide. *P. atrosepticum* is restricted to potato. *D. dianthicola* and *D. solani* are pathogenic to many plants in the tropical and subtropical regions and affect maize and dahlia in the temperate regions.

Symptoms of bacterial root rot, mainly characterized in potato and sweet potato, include chlorosis of leaf tissue and a black, water-soaked decay at the bottom of the stems that gradually extends to the top. In severe cases, the entire plant collapses [248]. Fibrous roots have localized lesions with a characteristic black appearance. In storage roots, sunken brown lesions with black margins can be observed at the surface [249].

The information regarding viruses as the causal agents of root rot is limited. Some studies have reported the effect of Cassava brown streak virus [7,8] and Ugandan cassava brown streak virus [250] in root rot development. However, sparse or no evidence is available in yield losses and molecular mechanisms of resistance within the host. Information regarding these non-traditional root rot agents is expected to increase as detection methods evolve [3].

3. Molecular mechanisms of resistance against root rot pathogens

When a pathogen attacks a plant, several molecular mechanisms are activated. Plants first respond to pathogen infection via pathogen-associated molecular pattern (PAMP) triggered immunity (PTI) [251]. Pathogens respond with a variety of effector proteins to counter the PTI [252]. Plants can sometimes detect these effectors and respond with the commonly known effector-triggered immunity (ETI).

Most plant-pathogen interactions characterized so far fall under the gene-for-gene interaction model. According to this model, a dominant or semi-dominant resistant (R) gene from the host and a corresponding avirulence (AVR) gene from the pathogen interact and activate further downstream reactions (PTI and/or ETI). The R-AVR interaction concludes with an incompatible response in which no disease symptoms are produced [253]. The most abundant R genes code for the plant nucleotide binding site leucine-rich repeat (NBS-LRR) proteins that are responsible for detecting potential pathogens and triggering a defense response [254].

Lack of corresponding R-AVR interactions leads to a compatible response leading to pathogenesis. Both compatible and incompatible interactions result in the recruitment of different sets of proteins that determine total and partial resistance or susceptibility.

The molecular mechanisms of root rot pathogens-mediated interactions are very diverse. The variations in plant-pathogen interaction are dependent on the species and race of the pathogen and the specific host genotype involved. The following sections describe validated and proposed molecular defense mechanisms against fungal – *Rhizoctonia*, *Fusarium*, *Phoma*, and *Thielaviopsis basicola*; and oomycete – *Aphanomyces*, *Pythium*, and *Phytophthora*, root rot pathogens.

4. Common causal agents of root rots

4.1. Fungal root rot

4.1.1. *Rhizoctonia solani*

Rhizoctonia solani is a pathogen with a broad host range. Resistance to *R. solani* has been studied mainly in sugar beet and rice with additional reports in potato and model organisms such as *Arabidopsis* and tobacco.

In sugar beet, the genetic basis for *Rhizoctonia* resistance is considerably narrow. The GWS 359-52R genotype is the universal parental line for essentially all resistant cultivars [255–257]. An early study suggested that the resistance to *R. solani* is associated with at least two loci with two or three alleles [258]. This hypothesis was supported by a recent QTL analysis, which localized the resistance loci on chromosomes 4, 5, and 7. These QTLs collectively explain 71% of the total phenotypic variation [257]. Genes involved in pathogen recognition and responses downstream of R-genes co-segregated with the resistance QTL located on chromosomes 4 and 7, respectively [257]. Genes that show similarity with the Xa21 and Pto were found to co-segregate with the QTL on chromosome 5. Xa21 is a well-characterized cell membrane receptor that, through phosphorylation and cleavage of its intracellular kinase domain, perceives the presence of pathogens [259]. Xa21 relays the signal to the nucleus through multi-step signal cascades, involving MAPK and WRKY signaling [260,261]. Xa21 is known to trigger hormone signaling, especially cytokinins [259]. The role of cytokinins in defense response remains elusive; however, studies have reported that cytokinins prompt salicylic acid (SA) accumulation [262,263]. The Pto gene encodes a cytoplasmic serine-threonine kinase that interacts with avirulence proteins and confers HR-mediated resistance [264]. Pto has been considered an important candidate gene for broad-spectrum resistance in molecular breeding approaches [265,266].

Metabolomic profiling aided in characterizing the changes in sugarbeet 0 and 7 days after inoculation (dai) with *R. solani* [267]. N1-caffeoyl-N10-feruloylspermidine and codonocarpine, both alkaloids, showed higher levels in resistant germplasm roots than susceptible germplasm at 0 dai. The role of alkaloids has been suggested to be a conserved defense response to *R. solani* and other necrotrophic fungal pathogens [267,268]. N1-caffeoyl-N10-feruloylspermidine and codonocarpine alkaloids have multiple and complex roles; therefore, it is difficult to predict their specific function against *R. solani*. Furthermore, two oleanic acid-like compounds (saponins) were found in the resistant germplasm, and their abundance continued to increase after infection with *R. solani*. Saponins are known to have antifungal activity [269,270]. Thus, three metabolites, N1-caffeoyl-N10-feruloylspermidine, codonocarpine, and oleanic acid-like compounds, are important candidates for follow-up studies on the interaction between sugar beet and *R. solani*.

No rice cultivar shows complete resistance, but partial resistance to *R. solani* has been reported. These studies have proposed that different defense mechanisms are activated in the partially resistant rice genotypes. A summary of changes detectable following *R. solani* inoculations in the partially resistant rice genotypes is presented in Figure 1.

Twenty-five genes were found to be differentially expressed in rice after infection with *R. solani* [271]. These same genes were also differentially expressed when rice was challenged with *Magnaporthe grisea* and *Xanthomonas oryzae* suggesting a conserved defense response to different pathogens. This analysis showed that Pathogenesis-Related (PR) 1b and PBZ1 genes were detected at 12 hours post-infection (hpi) when the *R. solani* mycelium started to grow on the surface of the plant [271]. The expression of PR1b increased gradually from 12 to 72 hpi. A few lesions began to develop at 36 hpi, and typical lesions developed at 48 hpi. Meanwhile, the expression of PBZ1 increased to its maximum level at 48 hpi. PR1b gene is induced by pathogens commonly associated with SA-related systemic acquired resistance (Figure 1) [272,273]. Further downstream function or signaling effects of the PR1b protein remain unknown. The PBZ1 gene, a PR10 family protein, has been shown to induce cell death in rice, *Nicotiana tabacum*, and *Arabidopsis* lines [274]. Cell death is caused by PBZ1-RNase activity inside the plant cell (Figure 1) [274,275]. On the other hand, the gene glutathione peroxidase 1 (GP1), which protects cells against both oxidative stresses and inhibits oxidative stress-induced cell death, was found to be induced at four hpi, reaching a maximum at 24 hpi upon *R. solani* infection (Figure 1) [276].

Feedback signaling potentially provides an equilibrium between the antagonistic action of PR1b and PBZ1 versus GP1 during the defense response against *R. solani*.




Figure 1. Model of resistance response following inoculation with *Rhizoctonia solani* in rice. Events include the recognition of PAMPs by the host (A); cell signaling induced by both jasmonate (JA) and salicylic acid (SA) hormones (B); cell death (C); ROS scavenging (D); synthesis and action of enzymes that attack the pathogen as well prepare the host for the attack (E); and systemic acquired resistance (F). The specific cell receptors participating in recognition of PAMPs are still unknown. SA participates directly in cell death and SAR activation, which are triggered by the PBZ1-RNase and PR1b enzymes, respectively. ROS scavenging is performed by the glutathione peroxidase 1 (GP1), which protects cells against both oxidative stresses and inhibits oxidative stress-induced cell death. JA triggers the activation of the phenylpropanoid pathway for the lignification of the host cell walls. Chitinases are synthesized and released to combat the *R. solani*. Defense against *R. solani* is best achieved by early action against the young hyphae. Model adapted and summarized from S. Chen et al., (2004); Shrestha et al., (2008); Taheri & Tarighi, (2010); C.-J. Zhao et al., (2008).

Another study demonstrated that chitinase levels correlated with resistance to *R. solani* in rice cultivars (Figure 1) [277]. Chitinase activity was detected 24 h after inoculation of seven moderately resistant cultivars. However, in a susceptible genotype, chitinase activity was delayed and was seen only after 36 h post-inoculation. Moderately resistant rice cultivars had higher levels of chitinase activity and lower disease severity and number of infection cushions formed than the susceptible genotype [277]. Studies indicate that protection in potatoes against *R. solani* is enhanced by co-expression of chitinases, 1,3- β -glucanases, and osmotin proteins [279,280]. It has been hypothesized previously that co-expression of these enzymes is needed to speed up the destruction of *R. solani* hyphae [280,281]. Newly synthesized chitin in cell walls of young hyphae is more sensitive to enzymatic degradation [282]. Therefore, the defense against *R. solani* is best achieved by early action against the young hyphae.

Resistance to *R. solani* in rice has also been associated with the jasmonate (JA) mediated priming of the phenylpropanoid pathway and the resultant enhanced lignification (Figure 1) [278]. A gene that rapidly accumulated to high, sustained levels in rice after *R. solani* challenge was the disease resistance response protein 206 [271]. This gene

participates in the production of active lignans, thus playing a central role in plant secondary metabolism. It was proposed that it be worth evaluating this protein's role in defense against *R. solani* in rice [271].

Resistance response in potato, bean, and cowpea seems to be dependent on SA [280,283,284]. On the other hand, a screening of 36 *Arabidopsis thaliana* ecotypes with differences in auxin, camalexin, SA, abscisic acid (ABA), and JA/ethylene (Et) acid pathways did not reveal any variation in response to *R. solani*. It demonstrated that resistance to *R. solani* was independent of these metabolic pathways [285]. In *A. thaliana*, it has been shown that NADPH oxidases mediate the resistance to *Rhizoctonia solani* [285]. The NADPH oxidase double mutant resulted in an almost complete loss of resistance. This last observation highlights a unique target to be evaluated or incorporated in crop plants such as sugar beet and rice.

4.1.2. Transgenic approach to combat *Rhizoctonia solani*

A polygalacturonase-inhibiting protein (PGIP) from sugar beet introduced into *Nicotiana benthamiana* resulted in enhanced resistance to *R. solani* [286]. Crude PGIP protein extracts from transgenic *N. benthamiana* plants significantly inhibited *R. solani* polygalacturonase. The crude extracts also inhibited polygalacturonase from *Fusarium solani* and *Botrytis cinerea*. Transgenic plants were also significantly more resistant to these three fungi [286]. Similarly, the expression of a common bean-PGIP also conferred strong resistance against *R. solani* in tobacco [287]. Transgenic tobacco expressing the bean PGI also expressed enhanced resistance against *Phytophthora parasitica* and *Peronospora hyoscyami* [287]. Transgenic sugar beet expressing the bean PGIP gene showed only minor quantitative effects in enhancing resistance against *R. solani* [288].

Transgenic rice expressing ACS2 (1-aminocyclopropane-1-carboxylic acid synthase, a key enzyme of Et biosynthesis) gene exhibited increased resistance to a field isolate of *R. solani*, as well as different races of *M. oryzae* [289]. This study showed an increased expression of PR1b (10 to 60-fold) and PR5 (2.0 to 7.9-fold) genes in the transgenic lines, as well as no negative impact on crop productivity [289]. Rice transgenic lines expressing broad-spectrum resistance 2 (BSR2) [290], thaumatin-like proteins [64], and a chitinase [291] have also exhibited enhanced resistance against *R. solani* as well.

4.1.3. *Fusarium solani* root rot: The case of pea and similitudes with soybean

One of the predominant causal agent of root rots in *P. sativum* is *Fusarium solani* f. sp. *pisi* (*Fsp*). The molecular responses to *Fsp* infection have been reported in pea since the late 1970's. A model of partially resistant and susceptible reactions against *Fsp* in pea is presented in Figure 2. Some of these studies have reported the association between pea and its non-host pathogen *F. solani* f. sp. *phaseoli* (*Fsph*). Generally, non-host resistance is more durable due to the involvement of multiple mechanisms making it an important model to study [292].

Experiments examining the interaction between pea-*Fsph* and pea-*Fsp* showed that *Fsph* and *Fsp* elicitors such as chitosan and DNase are released and directly affect the chromatin structure of the plant host (Figure 2) [293–296]. In turn, chromatin structure changes result in the alteration of gene expression patterns (Figure 2). Changes in chromatin structure, such as the decrease of the expression of High-Mobility Group (HMG) A transcription factor and modification of histones H2A/H2B, have been temporarily associated with the onset of PR gene activation (Isaac *et al.* 2009) (Figure 2). Pretreatment of pea tissue with chitosan and *Fsph* DNase has been shown to enhance protection against *Fsp* [297,298].




Figure 2. Model of reported changes detected following *Fusarium solani* f. sp. *pisi* (*Fsp*) inoculation in pea. Events include the action of *Fsp* DNase, chitosan, and PAMPs and/or their detection by the host. DNase and chitosan are associated with nuclear fragmentation in the plant nucleus affecting chromatin structure. These changes, along with the host's detection of PAMPs, trigger the defense responses such as the accumulation of pathogenesis-related (PR) genes. Specific responses in partially resistant and susceptible pea genotypes are depicted. Upwards and downwards pointing arrows represent overexpression and underexpression of genes, respectively. Model adapted and summarized from Hadwiger, (2008); B. A. Williamson-Benavides et al., (2020).

The accumulation of PR RNA seems crucial to acquiring resistance against *Fsph* [295]. PR proteins, such as the defensins DRR230 and DRR39, and the RNase PR-10 have a direct antifungal effect [296,300]. Other PR proteins, such as PR-1, a homolog of PR1b in *Arabidopsis*, act as positive regulators of plant immunity [296]. Chitinase and β -glucanase are constitutively expressed, but their basal expression increases 10 hours post-inoculation with *Fsph* [301]. These PR proteins' expression occurs within the crucial period for developing a resistance response against *Fsph* [294]. Similar mechanisms of resistance are triggered in pea to halt *F. oxysporum* pv. *pisi* infection [295,302].

There are significant similarities in the biochemical responses induced by the non-host pathogen *Fsph* and the host-pathogen *Fsp* in pea. In both cases, there is a nearly complete suppression of the phosphorylation of chromatin proteins, which leads to the elimination of HMG A from the cell nuclei and alteration of the histone biochemical structure [293–296]. Also, the same PR genes, such as DRR230, DRR39, RNase PR-10, and PR-1, seem to be upregulated when challenged with the two pathogens. The major difference in the biochemical responses induced by *Fsph* and *Fsp* is the speed at which the plants react. The type of response exhibited by pea varies with the rate of induction of PR genes and other associated biochemical pathways. In case of either the *Fsph* or *Fsp* infection, the fungus releases DNAses extracellularly, which localize to the host nuclei and degrades the nuclear DNA (Figure 2) [295,296,303]. Fungal DNAses can also impact the nuclei in the fungal mycelia and trigger their deterioration [295]. In case of a compatible interaction between *Fsp* and a pea genotype, the host's slower reaction rate allows *Fsp* to protect a small number of its nuclei from fungal DNAses, allowing the growth of *Fsp* to resume after 12 hours post-inoculation [296,298]. In contrast, the relatively rapid response generated in the host against *Fsph* terminates the fungi's development at 6 hrs post-inoculation [295,296].

In pea, phenylalanine ammonia-lyase and chalcone synthase enzymes are upregulated two hours post-inoculation with *Fsph* and *Fsp*. These two enzymes participate in the phenylpropanoid pathway and play a significant role in producing flavonoids and isoflavonoids (Figure 2). The phenylpropanoid pathway potentially plays a role in partial

resistance to *Fsp* in pea (Figure 2) [299] and partial resistance to *Fusarium solani* f. sp. *glycines* (*Fsg*) in soybean [304,305].

Pea contains an isoflavone synthase enzyme, which redirects phenylpropanoid pathway intermediate naringenin (4',5,7-trihydroxyflavanone) to synthesize pisatin (Sreedhya *et al.* 2006). Pisatin is an extensively studied phytoalexin from pea, and its production increases in the presence of *Fsp*, *Fsph*, and chitosan [293]. Interestingly, *Fsp* isolates incapable of demethylating pisatin are low in virulence and susceptible to pisatin [295,306]. The phytoalexin glyceollin levels increased in the *Fsp*-inoculated roots of two partially resistant soybean cultivars compared to a susceptible one [304]. The role of these two phytoalexins in the interaction between plant host and *Fusarium* remains to be elucidated.

A time-course RNAseq compared the expression of *Fsp*-responsive genes in four partially resistant and four susceptible pea genotypes after 0, 6, and 12 hours post pathogen challenge [299]. *Fsp* challenge produced a more intense and diverse overexpression of genes in the susceptible genotypes. In contrast, the partially resistant genotypes showed fewer changes in the expression of defense-related genes and a faster reset to a basal metabolic state (Figure 2). In the partially resistant genotypes, gene expression and GO enrichment analyses revealed that genes involved in exocytosis and secretion by cell, the anthocyanin synthesis pathway, as well as the DRR230 PR gene were overexpressed (Figure 2) [299]. Genes coding for receptor-mediated endocytosis, sugar transporters, SA synthesis, and signaling, and cell death were overexpressed in the susceptible genotypes (Figure 2).

A total of five RIL populations have been analyzed to identify QTLs in response to *Fsp* challenge [307–310]. A major QTL, named *Fsp-Ps2.1*, has been found on chromosome 6 that explains 39.0–53.4% of the phenotypic variance [307–309]. The *A* (pigmented flower/anthocyanin pigmentation) gene was mapped within the interval of *Fsp-Ps2.1*. However, *Fsp-Ps2.1* was mapped in a white flower (*aa* × *aa*) cross [308]. Therefore, it has been hypothesized that the resistance gene(s) responsible for *Fsp-Ps2.1* effect may not necessarily be *A* since *Fsp-Ps2.1* was initially identified in a white (*a*) flowered cross. *Fsp-Ps2.1* co-located with the *Aphanomyces* root rot partial resistance QTL *Ae-Ps2.1* [307,308]. A second QTL, *Fsp-Ps6.1*, explained 17.3% of the phenotypic variance. Three defensin family genes, pI39 and DRR230-A and DRR230-B, mapped near *Fsp-Ps3.1* [308]. A different subset of parental white-flowered genotypes was crossed to developed two populations that segregate for *Fsp* resistance [310]. QTL analysis of these two populations identified five QTLs that explain 5.26–14.76% of the resistance to *Fsp*. Three of these are considered newly reported QTLs. The recently identified QTLs and the absence of a major QTL on chromosome 6, reported in previous studies, reflects the wide degree of genetic resources of resistance available to combat *Fsp* in pea [310].

4.1.4. Transgenic approach to counter *Fusarium solani* root rot

An antibacterial peptide-encoding gene from alfalfa seeds, alfAFP, was fused to the C-terminal of the rice chitinase-encoding gene and introduced into tobacco [311]. The recombinant protein enhanced resistance against *F. solani* in transgenic tobacco plants. Transgenic lines did not exhibit wilting symptoms, even 30 days post-inoculation with *F. solani* [311]. In a different approach, the Et-responsive factor *StERF94* was expressed in potatoes. The transgenic lines exhibited enhanced resistance to *F. solani* [312]. Transgenic potato plants showed a limited production of H₂O₂ and increased expression of antioxidant enzymes and PR proteins [312].

4.1.5. *Fusarium graminearum* root rot

As is the case with other root rots, the mechanism of interaction of soybean and *Fusarium graminearum* has not been studied in depth. However, several QTL analyses have identified the potential loci responsible for resistance, and in some cases, several candidate genes have been mapped to these QTL regions.

Five putative QTLs were identified in a RIL population derived from a cross between Conrad (resistant) × Sloan (susceptible) parents [114]. These QTLs explained a small percentage of the phenotypic variance (3.6-9.2%) and were located on chromosomes 8, 13, 15, 16, and 19. These QTLs were not the same as those that confer resistance to *Phytophthora sojae*, suggesting that different loci are involved in resistance against these root rot pathogens [114]. Similar results were obtained from a genome-wide association study using cultivated and landraces of soybean [313]. This study identified twelve SNPs associated with *F. graminearum* resistance, which explained only a small percentage (5.53–14.71%) of the observed phenotypic variation.

A major QTL on chromosome 8 that explained 38.5% of the phenotypic variance was found in a RIL population derived from a cross between 'Wyandot' (partially resistant) × PI 567301B (highly resistant) [113]. This QTL harbored 39 genes, including the *Rhg4* locus for soybean cyst nematode (SCN) resistance [113]. Nine genes coding for hydroxymethylglutaryl-CoA, a key enzyme in flavonoid biosynthesis pathway, were found in this QTL. In addition, there were three rapid alkanization factor (RALF) genes that can initiate a signal transduction pathway and two genes coding for subtilisin-like proteases. Subtilisin proteases are believed to be secreted into the extracellular matrix and function to reorganize cell wall components during defense response [113,314]. A subsequent study identified four differentially expressed genes that mapped to this QTL located on chromosome 8. These genes included an actin-related protein 2/3 complex subunit, an unknown protein, a hypothetical protein, and a chalcone synthase 3 [315]. This study demonstrated that removal of the seed coat of highly resistant soybean lines makes them susceptible to *F. graminearum*, indicating that proteins or secondary compounds in the seed coat may be involved in resistance [315].

4.1.6. *Fusarium root rot in cereals*

The *Fusarium* species *F. avenaceum*, *F. graminearum*, *F. culmorum*, *F. verticillioides*, *F. pseudograminearum* are ubiquitous soil-borne fungus able to cause foot and root rot and *Fusarium* head blight or ear mold on different small-grain cereals such as wheat, barley, maize, and oat [316–318]. The emphasis of this review is limited to horticultural crops. For a comprehensive review of *Fusarium* disease in cereals, the reader is directed to previous studies and reviews [102,319–326].

4.1.7. *Phoma root rot*

Studies related to the understanding of the molecular mechanism underlying *Phoma* resistance are scarce. A few studies were identified that focused on JA and thiabendazole's use to combat postharvest rots caused by *Phoma betae* and *P. sclerotioides* [327,328] or on the assessment of alfalfa cultivars for resistance against *P. sclerotioides* [151,329].

A recent study in onion utilized different sources of genetic resistance to *P. terrestris* to develop segregating families. One segregating family was scored for resistance and susceptibility with the resultant ratio of segregants fitting a single dominant locus. However, in another segregating family, the resulting segregation ratio did not fit a single dominant or a recessive locus [330]. The severity of root rot was mapped to a locus on chromosome 4, and it explained 28%-35% of the phenotypic variation. Estimates of additive and dominance effects revealed that this source of resistance is co-dominantly inherited [330].

P. terrestris resistance was also assessed from a different source of resistance, and the resulting, co-dominantly inherited, QTL mapped to the same region on chromosome 4 and explained 54% of the phenotypic variation [330]. This study demonstrated that resistance from different genetic sources mapped to the same chromosome region and showed similar modes of inheritance [330].

4.1.8. *Thielaviopsis basicola root rot*

A QTL mapping study [331] and a proteomic analysis [138] are the only reports that provide some insight into the sources of resistance and defense response mechanisms against *Thielaviopsis basicola* in cotton. Phenotypic variation between resistant and susceptible cotton lines was associated with three QTLs that explained 19.1, 10.3, and 8.5% of the total phenotypic variation [331]. This study provided a list of 624 candidate genes that were located within the identified QTL regions. The list included 22 pathogen defense and 36 stress-responsive genes. Fine mapping is required to narrow down this list of candidate genes for each QTL.

A time-course analysis of cotton root proteomes was performed during a compatible interaction with *T. basicola* [138]. Analysis of root extracts was conducted at 1, 3, 5, and 7 days post-inoculation. The study found that more plant proteins were down-regulated, especially in the early stages of infection, than upregulated. Fifty-eight protein clusters were found to be upregulated across the time-course analysis. According to their putative biological role, these fifty-eight protein clusters were further identified and classified into five major categories: defense, stress, primary and secondary metabolism, and diverse function. A number of the upregulated proteins corresponded to PR, with the majority of them belonging to the PR-10 family. The function of PR-10 genes is still unknown; however, the authors suggested that these proteins may be involved in hormone-mediated disease resistance in cotton [138]. A putative thaumatin protein, another PR protein, was also upregulated during *T. basicola* infection. Two additional pathogen defense proteins corresponded to a Meloidogyne-induced protein MIC-3, which were originally correlated with the disruption of nematode development in cotton [332]. The molecular function of the MIC-3 and the MIC family remains unknown, mainly due to the absence of known functional motifs and domains [333].

4.1.9. A transgenic approach to counter *Thielaviopsis basicola* root rot

Transgenic cotton lines expressing AtNPR1 (nonexpresser of PR1) gene were found to be significantly tolerant to *T. basicola* [152]. The roots of the transgenic lines tended to recover faster after *T. basicola* infection. Transgenic plants also showed higher shoot and root mass, longer shoot length, and a greater number of boll-set than wild-type plants after *T. basicola* infection. NPR1 is a regulatory protein that participates as a critical positive regulator of the SA-dependent signaling pathway and systemic acquired resistance. Transcriptional analysis of transgenic roots exhibited stronger and faster induction of PR proteins such as PR1, thaumatin, glucanase, LOX1, and chitinase [152].

4.2. Oomycete root rot

4.2.1. *Aphanomyces* root rot:

A. euteiches cause high yield losses in legumes such as pea and alfalfa. It has been challenging to investigate the genetic basis of resistance in these two plant species due to their complex and partial genome information, the polygenic inheritance of resistance, and difficulties in field-based phenotyping. *Medicago truncatula*, with a much simpler genome, has been used as a surrogate model to understand the molecular interactions and resistance mechanism against *A. euteiches* [334]. The key molecular responses associated with *A. euteiches* resistance in *Medicago truncatula* are presented in Figure 3.




Figure 3. Model of resistance response following inoculations with *Aphanomyces euteiches* in *Medicago truncatula*. Events include the synthesis and signaling of abscisic acid (ABA) and reactive oxygen species (ROS). ROS are regulated by a small GTPase, named MtROP9. ABA and ROS signaling result in the overexpression of PR5 gene and the phenylpropanoid pathway. The lignification and synthesis of 2'-O-methylisiquiritigenin are positively correlated with disease resistance against *A. euteiches*. 2'-O-methylisiquiritigenin was shown to significantly impede *A. euteiches* development and zoospore germination. Model adapted and summarized from Badis et al., (2015); Colditz et al., (2004); Djéballi et al., (2009); Leonard M Kiirika et al., (2014); Leonard Muriithi Kiirika et al., (2012); Nyamsuren et al., (2003); Trapphoff et al., (2009).

A monogenic control of resistance against *A. euteiches* has been reported in several studies. A QTL named AER1 was mapped to the distal part of chromosome 3 [337,342]. The genomic region corresponding to the QTL contains a supercluster of nucleotide-binding site leucine-rich repeat (NBS-LRR) genes [337,342]. This region also included proteasome-related genes, a cluster of nine F-box protein-encoding genes, and one gene coding for a ubiquitin-associated enzyme [343]. qRT-PCR data showed that the ubiquitin-associated enzyme and three F-box-encoding genes were induced in a resistant line (A17) following pathogen inoculation but not in the susceptible line. F-box proteins are known to be involved in hormone regulation and in plant immunity [344,345], regulation of pericycle cell divisions [346], and lateral root production [347].

The highly redundant presence of ABA-responsive proteins indicates that ABA-mediated signaling is involved in the interaction between *M. truncatula* and *A. euteiches* (Figure 3) [336,340]. ABA production and signaling are known to contribute to JA accumulation as well as for the activation of resistance against *Phytophthora irregularis* in *Arabidopsis* [348,349]. Furthermore, PR10 family proteins, interspersed within the ABA-responsive genes, increased in *M. truncatula* after *A. euteiches* infection. However, a later study reported that the accumulation of PR10 protein was mainly correlated with *A. euteiches* proliferation and not plant resistance [343]. RNAi-mediated suppression of several PR10 genes led to a reduced *A. euteiches* colonization, which was linked with a parallel induction of a set of PR5 proteins (thaumatin-like proteins) (Figure 3) [350]. Therefore, PR10 and PR5 proteins act antagonistically.

A significant link has been found between enhanced synthesis and accumulation of flavonoid compounds and resistance against *A. euteiches* (Figure 3) [335]. Transcriptome and proteome analyses revealed a strong induction of chalcone synthase and isoflavone reductase genes after pathogen challenge [341,343,351,352]. Furthermore, the gene coding for isiquiritigenin 2'-O-methyltransferase showed the highest induction after *A.*

euteiches infection in the most resistant lines [336]. The metabolite 2'-O-methylisoliquiritigenin was shown to significantly impede *A. euteiches* development and zoospore germination (Figure 3) [335].

The accumulation of lignin has also been linked with partial resistance against *A. euteiches* [335,337]. Furthermore, the higher accumulation of lignin in resistant plants is associated with more efficient hydrogen peroxide (H₂O₂) scavenging mechanisms that are activated due to infection (Figure 3) [353]. ROS are activated following *A. euteiches* inoculation and are regulated by a small GTPase, named MtROP9 [338,339,341]. The knock-down of MtROP9 in *M. truncatula* resulted in three primary outcomes: 1-) prevented the detection of respiratory burst oxidase homologs, 2-) led to reduced activity of enzymes involved in the primary antioxidative processes; and 3-) promoted *A. euteiches* hyphal root colonization [338,339].

4.2.2. Role of Nodulation and mycorrhizas in *Aphanomyces* root rot:

The Nod Factor perception (NFP) gene involved in nodulation was reported to confer resistance against *A. euteiches* in *M. truncatula* [351]. NFP knockout mutants were significantly more susceptible to *A. euteiches* than wild-type, while NFP overexpressing lines showed increased resistance. Transcriptome analyses showed that knockout of the NFP gene led to changes in the expression of more than 500 genes involved in dynamic cell processes associated with disease response [351].

The NF-YA1 gene, a central transcriptional regulator of symbiotic nodule development, determined susceptibility to *A. euteiches* [354]. The *Mtnf-ya1-1* mutant plants showed a better survival rate and reduced symptoms as compared to their wild-type background. Comparative analysis of the transcriptome of wild-type and *Mtnf-ya1-1* mutant lines resulted in identifying 1509 differentially expressed genes. Among these differentially expressed genes, 36 defense-related genes were constitutively expressed in *Mtnf-ya1-1*, while 20 genes linked to hormonal, notably auxins, Et and ABA, pathways were repressed [354].

Mycorrhiza seems to impart a bioprotective effect, observed earlier in pea roots against *A. euteiches* [355,356]. The increased resistance is postulated to be due to the following reasons: (I) enhanced physical resistance and damage compensation capability of the plant due to improved nutritional status, (II) changes in the microbial populations of the mycorrhizosphere, (III) competition between invading microorganisms (IV), increased production of secondary metabolites that have antimicrobial properties (V) activation of plant defense mechanism via accumulation of defense-related proteins [343,355]. Histochemical analysis of both microorganisms in the roots revealed a competition for physical space, which likely reduces *A. euteiches* hyphae or oospores, resulting in diminished disease symptoms [343].

4.2.3. *Pythium* root rot

The molecular mechanism of resistance to *Pythium* root rot has been primarily investigated in soybean and the common bean; however, recently, *Pythium*-responsive genes have been reported in apples as well.

In soybean, five QTLs associated with resistance to *P. sojae* were mapped to chromosomes 1, 6, 8, 11, and 13 [212]. Each QTL explained 7.9 to 17.8% of the phenotypic variation. QTLs associated with resistance to other root rot pathogens colocalize with the QTLs associated with *P. sojae* resistance. Chromosome 1 QTL colocalized with a QTL associated with resistance for *Phytophthora sojae* [357]. The chromosome 6 QTL was closely located to a QTL reported for *Phytophthora sojae* [358] and *Fusarium virguliforme* [359]. The QTL on chromosome 8 was found in a region associated with resistance for *F. virguliforme* [360]. The QTL on chromosome 13 was located in an area associated with resistance to several other soybean pathogens, including *Phytophthora sojae* [361], *F. virguliforme* [362], and *F. graminearum* [114]. The QTLs on chromosomes 6 and 8 also colocalized with two QTLs found associated with resistance to *Pythium ultimum*. These QTLs, associated with *P.*

ultimum resistance, on chromosomes 6 and 8 explain 7.5–13.5% and 6.3–16.8% of the phenotypic variance, respectively [363].

In common bean, the response of a set of 40 common genotypes to *P. ultimum* was investigated [364]. The emergence rate showed a significant association between seed coat color and response to this pathogen. Eleven bean genotypes with colored seeds exhibited a high percentage of emergence. A major gene (Py-1) controlling the emergence rate was mapped to the region of the gene P, an essential color gene involved in the control of seed coat color, located on LG 7. Using a RIL population of colored seeds, other two QTLs associated with the emergence rate and another with seedling vigor were identified on LG 3 and 6, respectively. QTL on LG6 was mapped to the gene V region, which is another gene involved in the genetic control of seed color.

The transcriptomic changes in apple root tissue when infected with *P. ultimum* were analyzed using tolerant and susceptible rootstock lines [365,366]. The mechanism of defense response involving the recognition of PAMPs, hormone signaling, and synthesis of PR genes was identified. Genes coding for proteins with predicted function of pathogen detection such as the chitin elicitor receptor kinase (CERK) and wall-associated receptor kinase (WAK) were among the differentially expressed genes identified in the resistant line. Genes associated with the biosynthesis and signaling of several phytohormones including Et, JA, and cytokinins were specifically induced in response to *P. ultimum* inoculation. The strong induction of cytokinin hydroxylase encoding genes suggests that cytokinin signaling may play a unique role in the defense response in apple roots. Furthermore, genes coding for secondary metabolism enzymes, cell wall fortification, PR proteins, laccase, mandelonitrile lyase, and cyanogenic beta-glucosidase were consistently up-regulated in the later stages of infection [365]. Like apple, in *Zingiber zerumbet* (shampoo ginger or wild ginger), high differential modulation of genes involved in cell wall fortification, lignin biosynthesis, and SA/JA hormone indicates that these genes play a central role in restricting *P. myriotylum* proliferation [367].

On a global scale, delayed and/or interrupted activation of multiple defense pathways seems to underlie susceptibility. This has been observed in various transcriptome analysis studies against root rots [295,296,299,353]. Similar observations were discernible from transcriptomic and microscopic data in the susceptible B.9 roots [27,366,368]. Microscopy data on the pathogen growth progress revealed a swift development of root necrosis in the most susceptible genotypes, with the entire root system becoming necrotic within a period of 24 h after initial infection [368]. The necrosis progression could be delayed for several days without the whole root tissues being engulfed for the most resistant genotypes.

4.2.4. *Phytophthora root rot:*

Soybean is the species of choice to understand the molecular interactions between *Phytophthora* and the plant host. *Phytophthora root rot* (PRR) of soybean is the second leading cause of yield loss in soybean in North America, surpassed only by soybean cyst nematode [369].

More than 20 dominant genes, known as RPS genes, associated with PRR resistance have been identified in the soybean genome, with most of them mapping to Chromosome 3 [197,358,359,370–374].

Once incorporated into soybean cultivars, *Phytophthora* race-specific resistance genes have a useful life of only 8 to 15 years before new virulent races of the pathogen evolve [197,371,375]. Intensive use of race-specific genetic resistance for control of *P. sojae* has resulted in the emergence of new races that are virulent to the current resistance genes [376]. Over 50 races of *P. sojae* have been reported in the literature [377,378]. Currently, none of the single host-resistance genes can counter all *P. sojae* races. Several reports have highlighted the importance of marker-assisted selection (MAS) to pyramid several QTLs in soybean cultivars to reduce losses by PRR [358,359,370,371,379]. This approach would help in reducing the selection pressure for new virulent races of *P. sojae*. Partial resistance or tolerance, also called quantitative disease resistance, generates a lower selection

pressure on the pathogen population; therefore, it is expected that partial resistance will be more durable than general race-specific resistance.

The *Rps1k* gene in soybean has garnered significant interest because it confers stable resistance to broad-spectrum *P. sojae* strains in the USA. The *Rps1k* gene locus, cloned as part of a BAC, carries two classes of coiled coil-nucleotide binding-leucine rich repeat ((CC)-NBS-LRR) genes [380,381]. *Rps2* and *Rps4* gene loci were cloned, and they were also characterized as NBS-LRR genes [373,382]. In *RpsS* gene locus, 14 predicted genes exist, with three being NBS-LRR type genes [383]. The *RpsYD29* gene was mapped to a region with two NBS-LRR type genes [384]. These two genes showed high similarity to the NBS-LRR present in the *Rps1k* gene locus. *Rps10* gene has been mapped to chromosome 17, where eight putative candidate genes were found. Two candidate genes encoding serine/threonine (Ser/Thr) protein kinases were identified [385]. The identity and function of the remaining RPS genes remain unknown.

In soybean roots, a strong correlation between the extent of preformed suberin in soybean roots and the resistance to *P. sojae* was observed [386]. As a cell wall component, suberin is known to constitute a barrier to the pathogen and also acts as a toxin to microbes due to its high concentration of phenolic compounds [386,387]. To colonize the root, *P. sojae* hyphae grow through the suberized middle lamellae between epidermal cells. This process took 2 to 3 h longer in Conrad (resistant genotype) than in OX760-6 (partially resistant genotype) [375]. Subsequent growth of hyphae through the endodermis was also delayed in Conrad. The delay in the progression of *P. sojae* in the resistant cultivar provides this genotype with more time to activate and establish its chemical defenses. Additionally, Conrad had more preformed aliphatic suberin and was induced to form more aliphatic suberin after initial infection than OX760-6. The authors concluded that suberin's synthesis provides a target for the selection and development of new soybean cultivars with higher levels of partial resistance to *P. sojae*.

4.2.5. *Phytophthora* root rot in other crops

P. rubiis is one of the most serious and destructive diseases of raspberry [388,389]. Two major QTLs, located on LG 3 and 6, associated with *Phytophthora* resistance, have been identified [389]. Root vigor and disease resistance mapped to the same major QTL on LG 3. An auxin receptor or germin-like protein mapped to this LG 3 QTL. This QTL is possibly involved in the initiation of new axes of growth as a defense response. The effect of the LG 6 QTL has only been identified at the infected site. Therefore, LG 6 QTL may be better interpreted as a resistance locus rather than a vigor-related gene [389].

In avocado, the transcript levels of defense-related genes were characterized and compared among five rootstocks with varying resistance to root rot [390]. The results indicated the involvement of PR-5 and endochitinase in the defense response. However, neither of the genes could be directly linked to the observed resistance. The difference in transcript abundance of phenylalanine ammonia-lyase and lipoxygenase genes was also observed when comparing resistant and less resistant rootstocks, indicating their potential involvement in the resistance.

In strawberry, five genes for resistance to thirty races of *P. fragariae* have been identified, including Rpf1 that was characterized as a dominant monogenic gene that confers resistance to at least 18 races of *P. fragariae* [193,391]. However, none of the five genes have been characterized or associated with any known defense mechanism.

5. Conclusions

Moderate to high levels of resistance have been identified in breeding lines and/or cultivars of most crops affected by root rots. Furthermore, significant progress has been made in identifying genes that respond to fungi and oomycete root rot pathogens. It is evident that no universal response controls the resistance against this heterogeneous group of pathogens. The reactions are highly dependent on the host genetics and the

pathogen involved. Resistant responses are governed, mostly, by multiple mechanisms and genes and, on rare occasions, by a single, independent, dominant gene. Hormones that drive responses against root rots vary for each individual host-pathogen interaction. The role of SA, JA, Et, ABA, cytokinins, or any other hormone, should be studied on an individual basis.

Hundreds of studies have opted for the screening of resistance QTLs. In some of these studies, a major resistance QTL was identified. However, for most of these QTLs, the genes underlying resistance remain to be elucidated. The identification of these genes might help confer broad resistance against root rots.

The host's rapid response seems to be a shared feature that determines resistance against root rot pathogens. Resistant and susceptible genotypes commonly respond with the same set of defense genes; however, the slower speed of response in the susceptible genotypes results in pathogenesis. This standard feature was documented in the rice-*R. solani*, potato-*R. solani*, pea-*F. solani*, *M. truncatula*-*A. euteiches*, and apple-*Phytophthora ultimum* interactions.

Interestingly, legumes are predominant hosts of root rot pathogens. A shared factor among legumes might explain this susceptibility. A vital factor to consider is that legumes are the only known plant taxon that forms a symbiotic relationship with *Rhizobium* spp. Symbiosis-related mechanisms might be a gateway hijacked by root rot pathogens.

Author Contributions: BWB and AD wrote, have read, and agreed to the published version of the manuscript.

Funding: BWB acknowledges graduate research assistantship support from Washington State University Graduate School. Work in the Dhingra lab in the area of crop improvement is supported in part by Washington State University Agriculture Research Center Hatch grant WNP00011, and pea improvement by USA Dry Pea & Lentil Council.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kumari, N.; Katoch, S. Wilt and Root Rot Complex of Important Pulse Crops: Their Detection and Integrated Management. In *Management of Fungal Pathogens in Pulses*; Springer, 2020; pp. 93–119.
2. Bhat, K.A.; Masood, S.D.; Bhat, N.A.; Bhat, M.A.; Razvi, S.M.; Mir, M.R.; Sabina, A.; Wani, N.; Habib, M. Current status of post harvest soft rot in vegetables: a review. *Asian J. Plant Sci.* **2010**, *9*, 200–208.
3. Bodah, E.T. Root rot diseases in plants: a review of common causal agents and management strategies. *Agri Res Tech Open Access J* **2017**, *5*, 555661.
4. KIKUMOTO, T. Ecology and biocontrol of soft rot of Chinese cabbage. *Japanese J. Phytopathol.* **2000**, *66*, 60–62.
5. Liao, C.-H. Control of foodborne pathogens and soft-rot bacteria on bell pepper by three strains of bacterial antagonists. *J. Food Prot.* **2009**, *72*, 85–92.
6. Perombelon, M.C.M.; Kelman, A. Ecology of the soft rot erwinias. *Annu. Rev. Phytopathol.* **1980**, *18*, 361–387.
7. Bock, K.R. Studies on cassava brown streak virus disease in Kenya. *Trop. Sci.* **1994**, *34*, 134–145.
8. Hillocks, R.J.; Raya, M.D.; Mtunda, K.; Kiozia, H. Effects of brown streak virus disease on yield and quality of cassava in Tanzania. *J. Phytopathol.* **2001**, *149*, 389–394.
9. Louws, F.; Sun, J.; Whittington, H.; Driver, J.; Peeden, K.; Liu, B. Evaluation of fungicides and mustard meal to manage black root rot of strawberry and analysis of *Pythium*, *Fusarium*, and *Rhizoctonia* on strawberry roots. In Proceedings of the PHYTOPATHOLOGY; AMER PHYTOPATHOLOGICAL SOC 3340 PILOT KNOB ROAD, ST PAUL, MN 55121 USA, 2012; Vol. 102, p. 72.
10. Manici, L.M.; Caputo, F.; Baruzzi, G. Additional experiences to elucidate the microbial component of soil suppressiveness

- towards strawberry black root rot complex. *Ann. Appl. Biol.* **2005**, *146*, 421–431.
11. Particka, C.A.; Hancock, J.F. Field evaluation of strawberry genotypes for tolerance to black root rot on fumigated and nonfumigated soil. *J. Am. Soc. Hortic. Sci.* **2005**, *130*, 688–693.
 12. Xue, A.G. Biological control of pathogens causing root rot complex in field pea using *Clonostachys rosea* strain ACM941. *Phytopathology* **2003**, *93*, 329–335.
 13. Hosseini, S.; Elfstrand, M.; Heyman, F.; Jensen, D.F.; Karlsson, M. Deciphering common and specific transcriptional immune responses in pea towards the oomycete pathogens *Aphanomyces euteiches* and *Phytophthora pisi*. *BMC Genomics* **2015**, *16*, 627.
 14. Tu, J.C. Effects of soil compaction, temperature, and moisture on the development of the *Fusarium* root rot complex of pea in southwestern Ontario. *Phytoprotection* **1994**, *75*, 125–131.
 15. Zitnick-Anderson, K.; Simons, K.; Pasche, J.S. Detection and qPCR quantification of seven *Fusarium* species associated with the root rot complex in field pea. *Can. J. Plant Pathol.* **2018**, doi:10.1080/07060661.2018.1429494.
 16. Gaulin, E.; Jacquet, C.; Bottin, A.; Dumas, B. Root rot disease of legumes caused by *Aphanomyces euteiches*. *Mol. Plant Pathol.* **2007**.
 17. Hamon, C.; Baranger, A.; Coyne, C.J.; McGee, R.J.; Le Goff, I.; L'Anthoëne, V.; Esnault, R.; Riviere, J.-P.; Klein, A.; Mangin, P. New consistent QTL in pea associated with partial resistance to *Aphanomyces euteiches* in multiple French and American environments. *Theor. Appl. Genet.* **2011**, *123*, 261–281.
 18. Carling, D.E.; Baird, R.E.; Gitaitis, R.D.; Brainard, K.A.; Kuninaga, S. Characterization of AG-13, a newly reported anastomosis group of *Rhizoctonia solani*. *Phytopathology* **2002**, *92*, 893–899.
 19. Tewoldemedhin, Y.T.; Lamprecht, S.C.; McLeod, A.; Mazzola, M. Characterization of *Rhizoctonia* spp. recovered from crop plants used in rotational cropping systems in the Western Cape province of South Africa. *Plant Dis.* **2006**, *90*, 1399–1406.
 20. Allmaras, R.R.; Fritz, V.A.; Pflieger, F.L.; Copeland, S.M. Impaired internal drainage and *Aphanomyces euteiches* root rot of pea caused by soil compaction in a fine-textured soil. *Soil Tillage Res.* **2003**, *70*, 41–52.
 21. Falcon, M.F.; Fox, R.L.; Trujillo, E.E. Interactions of soil pH, nutrients and moisture on *Phytophthora* root rot of avocado. *Plant Soil* **1984**, *81*, 165–176.
 22. Rhoades, C.C.; Brosi, S.L.; Dattilo, A.J.; Vincelli, P. Effect of soil compaction and moisture on incidence of *phytophthora* root rot on American chestnut (*Castanea dentata*) seedlings. *For. Ecol. Manage.* **2003**, *184*, 47–54.
 23. La Porta, N.; Capretti, P.; Thomsen, I.M.; Kasanen, R.; Hietala, A.M.; Von Weissenberg, K. Forest pathogens with higher damage potential due to climate change in Europe. *Can. J. Plant Pathol.* **2008**, *30*, 177–195.
 24. Kubiak, K.; Żółciak, A.; Damszel, M.; Lech, P.; Sierota, Z. *Armillaria* pathogenesis under climate changes. *Forests* **2017**, *8*, 100.
 25. Klopfenstein, N.B. *Approaches to predicting potential impacts of climate change on forest disease: an example with Armillaria root disease*; US Department of Agriculture, Forest Service, Rocky Mountain Research Station, 2009;
 26. Papavizas, G.C.; Ayers, W.A. *Aphanomyces species and their root diseases in pea and sugarbeet*; 1974;
 27. Zhu, Y.; Saltzgeber, M. A systematic analysis of apple root resistance traits to *Pythium ultimum* infection and the underpinned molecular regulations of defense activation. *Hortic. Res.* **2020**, *7*, 1–11.
 28. Anderson, N.A. The genetics and pathology of *Rhizoctonia solani*. *Annu. Rev. Phytopathol.* **1982**, *20*, 329–347.
 29. Paulitz, T.C.; Adams, K. Composition and distribution of *Pythium* communities in wheat fields in eastern Washington State. *Phytopathology* **2003**, *93*, 867–873.
 30. Sneh, B.; Jabaji-Hare, S.; Neate, S.M.; Dijst, G. *Rhizoctonia species: taxonomy, molecular biology, ecology, pathology and disease control*; Springer Science & Business Media, 2013; ISBN 9401729018.
 31. Burton, R.J.; Coley-Smith, J.R.; Wareing, P.W.; Gladders, P. *Rhizoctonia oryzae* and *R. solani* associated with barley stunt disease in the United Kingdom. *Trans. Br. Mycol. Soc.* **1988**, *91*, 409–417.
 32. McKinley, A. Evaluation of progeny of genetically engineered barley plants for resistance to *rhizoctonia oryzae* and

- rhizoctonia solani AG-8. **2003**.
33. Neate, S.M. A comparison of controlled environment and field trials for detection of resistance in cereal cultivars to root rot caused by *Rhizoctonia solani*. *Plant Pathol.* **1989**, *38*, 494–501.
 34. Rush, C.M.; Carling, D.E.; Harveson, R.M.; Mathieson, J.T. Prevalence and pathogenicity of anastomosis groups of *Rhizoctonia solani* from wheat and sugar beet in Texas. *Plant Dis.* **1994**, *78*, 349–352.
 35. Karaca, G.H.; Ozkoc, I.; Erper, I. Determination of the anastomosis grouping and virulence of *Rhizoctonia solani* Kühn isolates associated with bean plants grown in Samsun/Turkey. *Pakistan J. Biol. Sci.* **2002**, *5*, 434–437.
 36. Muyolo, N.G.; Lipps, P.E.; Schmitthenner, A.F. Reactions of dry bean, lima bean, and soybean cultivars to *Rhizoctonia* root and hypocotyl rot and web blight. *Plant Dis.* **1993**, *77*, 234–238.
 37. Oladzad, A.; Zitnick-Anderson, K.; Jain, S.; Simons, K.; Osorno, J.M.; McClean, P.E.; Pasche, J. Genotypes and genomic regions associated with *Rhizoctonia solani* resistance in common bean. *Front. Plant Sci.* **2019**, *10*, 956.
 38. Peña, P.A.; Steadman, J.R.; Eskridge, K.M.; Urrea, C.A. Identification of sources of resistance to damping-off and early root/hypocotyl damage from *Rhizoctonia solani* in common bean (*Phaseolus vulgaris* L.). *Crop Prot.* **2013**, *54*, 92–99.
 39. Keinath, A.P.; Farnham, M.W. Differential cultivars and criteria for evaluating resistance to *Rhizoctonia solani* in seedling *Brassica oleracea*. *Plant Dis.* **1997**, *81*, 946–952.
 40. Rollins, P.A.; Keinath, A.P.; Farnham, M.W. Effect of inoculum type and anastomosis group of *Rhizoctonia solani* causing wirestem of cabbage seedlings in a controlled environment. *Can. J. plant Pathol.* **1999**, *21*, 119–124.
 41. Yanga, G.H.; Chenb, J.Y.; Pua, W.Q. NEW DISEASE REPORT. *Plant Pathol.* **2007**, *56*, 351.
 42. Grisham, M.P.; Anderson, N.A. Pathogenicity and host specificity of *Rhizoctonia solani* isolated from carrots. *Phytopathology* **1983**, *73*, 1564–1569.
 43. Naito, S.; Kanematsu, S. Characterization and pathogenicity of a new anastomosis subgroup AG-2-3 of *Rhizoctonia solani* Kühn isolated from leaves of soybean. *Japanese J. Phytopathol.* **1994**, *60*, 681–690.
 44. Punja, Z.K. Transgenic carrots expressing a thaumatin-like protein display enhanced resistance to several fungal pathogens. *Can. J. Plant Pathol.* **2005**, *27*, 291–296.
 45. Assunção, I.P.; Nascimento, L.D.; Ferreira, M.F.; Oliveira, F.J.; Michereff, S.J.; Lima, G.S.A. Reaction of faba bean genotypes to *Rhizoctonia solani* and resistance stability. *Hortic. Bras.* **2011**, *29*, 492–497.
 46. Rashid, K.Y.; Bernier, C.C. Genetic diversity among isolates of *Rhizoctonia solani* and sources of resistance in *Vicia faba*. *Can. J. Plant Pathol.* **1993**, *15*, 23–28.
 47. VALKONEN, J.P.T.; von HEIROTH, W.; SAVELA, M. Fungi and Gram-negative bacteria as soilborne minor pathogens of goat's rue (*Galega orientalis* Lam.). *Ann. Appl. Biol.* **1993**, *123*, 257–269.
 48. Dijst, G.; Schneider, J.H.M. Flowerbulbs diseases incited by *Rhizoctonia* species. In *Rhizoctonia species: taxonomy, molecular biology, ecology, pathology and disease control*; Springer, 1996; pp. 279–288.
 49. Fähler, B.; Petersen, P. Rapid greenhouse screening of maize for resistance to *Rhizoctonia solani* AG2-2IIIB/Ein schnelles Screening von Mais auf Resistenz gegenüber *Rhizoctonia solani* AG2-2IIIB im Gewächshaus. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz/Journal Plant Dis. Prot.* **2004**, 292–301.
 50. Garg, A.; Prasanna, B.M.; Sharma, R.C.; Rathore, R.S.; Saxena, S.C.; Chauhan, S.V.S. Identification of resistance sources to banded leaf and sheath blight (*Rhizoctonia solani* f. sp. sasakil) in maize. *Indian Phytopathol.* **2011**.
 51. Sharma, R.C.; Rai, S.N.; Batsa, B.K. Identifying resistance to banded leaf and sheath blight of maize. *Indian Phytopathol.* **2012**.
 52. Zhao, M.; Zhang, Z.; Zhang, S.; Li, W.; Jeffers, D.P.; Rong, T.; Pan, G. Quantitative trait loci for resistance to banded leaf and sheath blight in maize. *Crop Sci.* **2006**, *46*, 1039–1045.
 53. Kataria, H.; Verma, P.R.; Gisi, U. Variability in the sensitivity of *Rhizoctonia solani* anastomosis groups to fungicides. *J. Phytopathol.* **1991**, *133*, 121–133.
 54. Yang, J.; Verma, P.R. Screening genotypes for resistance to pre-emergence damping-off and postemergence seedling root rot

- of oilseed rape and canola caused by *Rhizoctonia solani* AG-2-1. *Crop Prot.* **1992**, *11*, 443–448.
55. Verma, P.R. Biology and control of *Rhizoctonia solani* on rapeseed: a review. *Phytoprotection* **1996**, *77*, 99–111.
56. Erper, I.; Karaca, G.H.; Turkkan, M.; Ozkoc, I. Characterization and pathogenicity of *Rhizoctonia* spp. from onion in Amasya, Turkey. *J. Phytopathol.* **2006**, *154*, 75–79.
57. Sharma-Poudyal, D.; Paulitz, T.C.; du Toit, L.J. Evaluation of onion genotypes for resistance to stunting caused by *Rhizoctonia solani* AG 8. *HortScience* **2015**, *50*, 551–554.
58. Hwang, S.F.; Gossen, B.D.; Conner, R.L.; Chang, K.F.; Turnbull, G.D.; Lopetinsky, K.; Howard, R.J. Management strategies to reduce losses caused by *Rhizoctonia* seedling blight of field pea. *Can. J. plant Sci.* **2007**, *87*, 145–155.
59. McCoy, R.J.; Kraft, J.M. Comparison of techniques and inoculum in evaluating peas (*Pisum sativum*) for resistance to stem rot caused by *Rhizobium solani*. *Plant Dis.* **1984**, *68*, 53–55.
60. Leach, S.S.; Webb, R.E. Evaluation of potato cultivars, clones and a true seed population for resistance to *Rhizoctonia solani*. *Am. Potato J.* **1993**, *70*, 317–328.
61. Naz, F.; Rauf, C.A.; Abbasi, N.A.; Haque, I.; Ahmad, I. Influence of inoculum levels of *Rhizoctonia solani* and susceptibility on new potato germplasm. *Pak. J. Bot* **2008**, *40*, 2199–2209.
62. Zhang, X.-Y.; Yu, X.-X.; Yu, Z.; Xue, Y.-F.; Qi, L.-P. A simple method based on laboratory inoculum and field inoculum for evaluating potato resistance to black scurf caused by *Rhizoctonia solani*. *Breed. Sci.* **2014**, *64*, 156–163.
63. Yanar, Y.; Yilmaz, G.; Cesmeli, I.; Coskun, S. Characterization of *Rhizoctonia solani* isolates from potatoes in turkey and screening potato cultivars for resistance to AG-3 isolates. *Phytoparasitica* **2005**, *33*, 370–376.
64. Datta, K.; Velazhahan, R.; Oliva, N.; Ona, I.; Mew, T.; Khush, G.S.; Muthukrishnan, S.; Datta, S.K. Over-expression of the cloned rice thaumatin-like protein (PR-5) gene in transgenic rice plants enhances environmental friendly resistance to *Rhizoctonia solani* causing sheath blight disease. *Theor. Appl. Genet.* **1999**, *98*, 1138–1145.
65. Eizenga, G.C.; Lee, F.N.; Rutger, J.N. Screening *Oryza* species plants for rice sheath blight resistance. *Plant Dis.* **2002**, *86*, 808–812.
66. Jha, S.; Tank, H.G.; Prasad, B.D.; Chattoo, B.B. Expression of Dm-AMP1 in rice confers resistance to *Magnaporthe oryzae* and *Rhizoctonia solani*. *Transgenic Res.* **2009**, *18*, 59–69.
67. Li, Z.; Pinson, S.R.M.; Marchetti, M.A.; Stansel, J.W.; Park, W.D. Characterization of quantitative trait loci (QTLs) in cultivated rice contributing to field resistance to sheath blight (*Rhizoctonia solani*). *Theor. Appl. Genet.* **1995**, *91*, 382–388.
68. Mazzola, M.; Wong, O.T.; Cook, R.J. Virulence of *Rhizoctonia oryzae* and *R. solani* AG-8 on wheat and detection of *R. oryzae* in plant tissue by PCR. **1996**.
69. Bradley, C.A.; Hartman, G.L.; Nelson, R.L.; Mueller, D.S.; Pederson, W.L. Response of ancestral soybean lines and commercial cultivars to *Rhizoctonia* root and hypocotyl rot. *Plant Dis.* **2001**, *85*, 1091–1095.
70. Muyolo, N.G.; Lipps, P.E.; Schmitthenner, A.F. Anastomosis grouping and variation in virulence among isolates of *Rhizoctonia solani* associated with dry bean and soybean in Ohio and Zaire. *Phytopathology* **1993**, *83*, 438–444.
71. Rahman, M.T.; Bhuiyan, M.K.A.; Karim, M.A.; Rubayet, M.T. SCREENING OF SOYBEAN RESISTANCE GENOTYPES AGAINST *Fusarium oxysporum*, *Macrophomina phaseolina*, *Rhizoctonia solani* AND *Sclerotium rolfsii*.
72. Yang, X.B.; Berggren, G.T.; Snow, J.P. Types of *Rhizoctonia* foliar blight on soybean in Louisiana. *Plant Dis.* **1990**, *74*, 501–504.
73. Nagendran, S.; Hammerschmidt, R.; McGrath, J.M. Identification of sugar beet germplasm EL51 as a source of resistance to post-emergence *Rhizoctonia* damping-off. *Eur. J. plant Pathol.* **2009**, *123*, 461–471.
74. Panella, L.; Frese, L.; Srivastava, H.M.; Lange, W. Screening and utilizing Beta genetic resources with resistance to *Rhizoctonia* root rot and *Cercospora* leaf spot in a sugar beet breeding program. In Proceedings of the International Beta Genetic Resources Network: A report on the 4th International Beta Genetic Resources Workshop and World Beta Network Conference Held at the Aegean Agricultural Research Institute, Izmir, Turkey. 28 Feb.–3 Mar. 1996.; International Plant Genetic Resources Institute Rome, 1998; pp. 62–72.

75. Scholten, O.E.; Panella, L.W.; De Bock, T.S.M.; Lange, W. A greenhouse test for screening sugar beet (*Beta vulgaris*) for resistance to *Rhizoctonia solani*. *Eur. J. Plant Pathol.* **2001**, *107*, 161–166.
76. Date, H.; Yagi, S.; Okamoto, Y.; Oniki, M. On the leaf blight of tomatoes by *Thanatephorus cucumeris* (Frank) Donk (*Rhizoctonia solani*). *Ann. Phytopathol. Soc. Jpn* **1984**, *50*, 1375–1381.
77. Nikraftar, F.; Taheri, P.; Rastegar, M.F.; Tarighi, S. Tomato partial resistance to *Rhizoctonia solani* involves antioxidative defense mechanisms. *Physiol. Mol. Plant Pathol.* **2013**, *81*, 74–83.
78. Taheri, P.; Tarighi, S. The role of pathogenesis-related proteins in the tomato-*Rhizoctonia solani* interaction. *J. Bot.* **2012**, *2012*.
79. Mahoney, A.K.; Babiker, E.M.; Paulitz, T.C.; See, D.; Okubara, P.A.; Hulbert, S.H. Characterizing and mapping resistance in synthetic-derived wheat to *Rhizoctonia* root rot in a green bridge environment. *Phytopathology* **2016**, *106*, 1170–1176.
80. Okubara, P.A.; Leston, N.; Micknass, U.; Kogel, K.-H.; Imani, J. Rapid Quantitative Assessment of *Rhizoctonia* Resistance in Roots of Selected Wheat and Barley Genotypes. *Plant Dis.* **2016**, *100*, 640–644.
81. Coleman, J.J. The *Fusarium solani* species complex: ubiquitous pathogens of agricultural importance. *Mol. Plant Pathol.* **2016**, *17*, 146–158.
82. Foroud, N.A.; Chatterton, S.; Reid, L.M.; Turkington, T.K.; Tittlemier, S.A.; Gräfenhan, T. *Fusarium* diseases of Canadian grain crops: impact and disease management strategies. In *Future Challenges in Crop Protection Against Fungal Pathogens*; Springer, 2014; pp. 267–316.
83. Zhang, N.; O'Donnell, K.; Sutton, D.A.; Nalim, F.A.; Summerbell, R.C.; Padhye, A.A.; Geiser, D.M. Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. *J. Clin. Microbiol.* **2006**, *44*, 2186–2190.
84. Šišić, A.; Bačanović-Šišić, J.; Al-Hatmi, A.M.S.; Karlovsky, P.; Ahmed, S.A.; Maier, W.; De Hoog, G.S.; Finckh, M.R. The 'forma specialis' issue in *Fusarium*: a case study in *Fusarium solani* f. sp. *pisii*. *Sci. Rep.* **2018**, *8*, 1–17.
85. Singh, R.; Soni, S.K.; Awasthi, A.; Kalra, A. Evaluation of vermicompost doses for management of root-rot disease complex in *Coleus forskohlii* under organic field conditions. *Australas. Plant Pathol.* **2012**, *41*, 397–403.
86. Singh, R.; Kalra, A.; Ravish, B.S.; Divya, S.; Parameswaran, T.N.; Srinivas, K.; Bagyaraj, D.J. Effect of potential bioinoculants and organic manures on root-rot and wilt, growth, yield and quality of organically grown *Coleus forskohlii* in a semiarid tropical region of Bangalore (India). *Plant Pathol.* **2012**, *61*, 700–708.
87. Lops, F.; Cibelli, F.; Raimondo, M.L.; Carlucci, A. First report of stem wilt and root rot of *Schlumbergera truncata* caused by *Fusarium oxysporum* f. sp. *Opuntiarum* in Southern Italy. *Plant Dis.* **2013**, *97*, 846.
88. Cohen, R.; Orgil, G.; Burger, Y.; Saar, U.; Elkabetz, M.; Tadmor, Y.; Edelstein, M.; Belausov, E.; Maymon, M.; Freeman, S. Differences in the responses of melon accessions to *Fusarium* root and stem rot and their colonization by *Fusarium oxysporum* f. sp. *radicis-cucumerinum*. *Plant Pathol.* **2015**, *64*, 655–663.
89. Golinski, P.; Kaczmarek, Z.; Kiecana, I.; Wisniewska, H.; Kaptur, P.; Kostecki, M.; Chelkowski, J. *Fusarium* head blight of common Polish winter wheat cultivars—comparison of effects of *Fusarium avenaceum* and *Fusarium culmorum* on yield components. *J. Phytopathol.* **2002**, *150*, 135–141.
90. Wojciechowski, S.; Chelkowski, J.; Ponitka, A.; Šlusarkiewicz-Jarżina, A. Evaluation of spring and winter wheat reaction to *Fusarium culmorum* and *Fusarium avenaceum*. *J. Phytopathol.* **1997**, *145*, 99–103.
91. Mesterhazy, A.; Bartók, T.; Kászonyi, G.; Varga, M.; Tóth, B.; Varga, J. Common resistance to different *Fusarium* spp. causing *Fusarium* head blight in wheat. *Eur. J. Plant Pathol.* **2005**, *112*, 267–281.
92. Linkmeyer, A.; Götz, M.; Hu, L.; Asam, S.; Rychlik, M.; Hausladen, H.; Hess, M.; Hüchelhoven, R. Assessment and introduction of quantitative resistance to *Fusarium* head blight in elite spring barley. *Phytopathology* **2013**, *103*, 1252–1259.
93. Xue, A.; Ho, K.; Butler, G.; Vigier, B.; Babcock, C. Pathogenicity of *Fusarium* species causing head blight in barley. *Phytoprotection* **2006**, *87*, 55–61.
94. Chittam, K.; Porter, L.; McPhee, K.; Khan, M.; Goswami, R.S. *Fusarium avenaceum* as causal agent of root rot in field peas

- and its control. In Proceedings of the Phytopathology; AMER PHYTOPATHOLOGICAL SOC 3340 PILOT KNOB ROAD, ST PAUL, MN 55121 USA, 2010; Vol. 100, pp. S25–S25.
95. Eranthodi, A.; Schneiderman, D.; Harris, L.J.; Witte, T.E.; Sproule, A.; Hermans, A.; Overy, D.P.; Chatterton, S.; Liu, J.; Li, T. Enniatin Production Influences *Fusarium avenaceum* Virulence on Potato Tubers, but not on Durum Wheat or Peas. *Pathogens* **2020**, *9*, 75.
96. Li, W.J.; Feng, J.; Chang, K.F.; Conner, R.L.; Hwang, S.F.; Strelkov, S.E.; Gossen, B.D.; McLaren, D.L. Microsatellite DNA markers indicate quantitative trait loci controlling resistance to pea root rot caused by *Fusarium avenaceum* (Corda ex Fries) Sacc. *Plant Pathol. J.* **2012**, *11*, 114–119.
97. Miller-Garvin, J.E.; Viands, D.R. Selection for resistance to *Fusarium* root rot, and associations among resistances to six diseases in alfalfa. *Crop Sci.* **1994**, *34*, 1461–1465.
98. You, M.P.; Barbett, M.J.; Nichols, P.G.H. New sources of resistance identified in *Trifolium subterraneum* breeding lines and cultivars to root rot caused by *Fusarium avenaceum* and *Pythium irregulare* and their relationship to seedling survival. *Australas. Plant Pathol.* **2005**, *34*, 237–244.
99. Browne, R.A.; Cooke, B.M. Resistance of wheat to *Fusarium* spp. in an in vitro seed germination assay and preliminary investigations into the relationship with *Fusarium* head blight resistance. *Euphytica* **2005**, *141*, 23–32.
100. Erginbas-Orakci, G.; Morgounov, A.; Dababat, A.A. Determination of resistance in winter wheat genotypes to the dryland root rots caused by *Fusarium culmorum* in Turkey. *Int. J. Agric. Wildl. Sci.* **2018**.
101. Scholten, O.E.; Steenhuis-Broers, G.; Osman, A.; Bremer, E. Screening for resistance to *Fusarium* head blight in spring wheat cultivars. **2006**.
102. Bai, G.; Shaner, G. Management and resistance in wheat and barley to *Fusarium* head blight. *Annu. Rev. Phytopathol.* **2004**, *42*, 135–161.
103. ŪSELE, G.; BEINAROVIČA, I.; MEŽAKA, I.; LEGZDIŅA, L. Comparison of spring barley (*Hordeum vulgare* L.) screening methods for *Fusarium* head blight resistance breeding. *Zemdirbyste-Agriculture* **2013**, *100*, 317.
104. Gavrilova, O.P.; Gagkaeva, T.Y.; Loskutov, I.G. Screening of parent material for breeding oat varieties resistant to *Fusarium* disease and accumulation of mycotoxins in grain. *Russ. Agric. Sci.* **2012**, *38*, 33–35.
105. Hori, K.; Kobayashi, T.; Sato, K.; Takeda, K. QTL analysis of *Fusarium* head blight resistance using a high-density linkage map in barley. *Theor. Appl. Genet.* **2005**, *111*, 1661–1672.
106. Mamo, B.E.; Steffenson, B.J. Genome-wide association mapping of *Fusarium* head blight resistance and agromorphological traits in barley landraces from Ethiopia and Eritrea. *Crop Sci.* **2015**, *55*, 1494–1512.
107. Massman, J.; Cooper, B.; Horsley, R.; Neate, S.; Dill-Macky, R.; Chao, S.; Dong, Y.; Schwarz, P.; Muehlbauer, G.J.; Smith, K.P. Genome-wide association mapping of *Fusarium* head blight resistance in contemporary barley breeding germplasm. *Mol. Breed.* **2011**, *27*, 439–454.
108. Anderson, J.A.; Stack, R.W.; Liu, S.; Waldron, B.L.; Fjeld, A.D.; Coyne, C.; Moreno-Sevilla, B.; Fetch, J.M.; Song, Q.J.; Cregan, P.B. DNA markers for *Fusarium* head blight resistance QTLs in two wheat populations. *Theor. Appl. Genet.* **2001**, *102*, 1164–1168.
109. Campbell, K.A.G.; Lipps, P.E. Allocation of resources: sources of variation in *Fusarium* head blight screening nurseries. *Phytopathology* **1998**, *88*, 1078–1086.
110. Del Blanco, I.; Froberg, R.; Stack, R.; Berzonsky, W.; Kianian, S. Detection of QTL linked to *Fusarium* head blight resistance in Sumai 3-derived North Dakota bread wheat lines. *Theor. Appl. Genet.* **2003**, *106*, 1027–1031.
111. Fuentes, R.G.; Mickelson, H.R.; Busch, R.H.; Dill-Macky, R.; Evans, C.K.; Thompson, W.G.; Wiersma, J. V.; Xie, W.; Dong, Y.; Anderson, J.A. Resource allocation and cultivar stability in breeding for *Fusarium* head blight resistance in spring wheat. *Crop Sci.* **2005**, *45*, 1965–1972.
112. He, X.; Singh, P.K.; Duveiller, E.; Schlang, N.; Dreisigacker, S.; Singh, R.P. Identification and characterization of international

- Fusarium head blight screening nurseries of wheat at CIMMYT, Mexico. *Eur. J. plant Pathol.* **2013**, *136*, 123–134.
113. Acharya, B.; Lee, S.; Mian, M.A.R.; Jun, T.-H.; McHale, L.K.; Michel, A.P.; Dorrance, A.E. Identification and mapping of quantitative trait loci (QTL) conferring resistance to Fusarium graminearum from soybean PI 567301B. *Theor. Appl. Genet.* **2015**, *128*, 827–838.
114. Ellis, M.L.; Wang, H.; Paul, P.A.; St. Martin, S.K.; McHale, L.K.; Dorrance, A.E. Identification of soybean genotypes resistant to Fusarium graminearum and genetic mapping of resistance quantitative trait loci in the cultivar Conrad. *Crop Sci.* **2012**, *52*, 2224–2233.
115. Zhang, J.X.; Xue, A.G.; Zhang, H.J.; Nagasawa, A.E.; Tambong, J.T. Response of soybean cultivars to root rot caused by Fusarium species. *Can. J. Plant Sci.* **2010**, *90*, 767–776.
116. Ali, M.L.; Taylor, J.H.; Jie, L.; Sun, G.; William, M.; Kasha, K.J.; Reid, L.M.; Pauls, K.P. Molecular mapping of QTLs for resistance to Gibberella ear rot, in corn, caused by Fusarium graminearum. *Genome* **2005**, *48*, 521–533.
117. Asran, M.R.; Buchenauer, H. Pathogenicity of Fusarium graminearum isolates on maize (*Zea mays* L.) cultivars and relation with deoxynivalenol and ergosterol contents/Pathogenität von Fusarium graminearum Isolaten an Mais-(*Zea mays* L.) Sorten und Beziehung zu Deoxynivalenol-und Ergost. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz/Journal Plant Dis. Prot.* **2003**, 209–219.
118. du Toit, L.J.; Kirby, H.W.; Pedersen, W.L. Evaluation of an aeroponics system to screen maize genotypes for resistance to Fusarium graminearum seedling blight. *Plant Dis.* **1997**, *81*, 175–179.
119. Löffler, M.; Kessel, B.; Ouzunova, M.; Miedaner, T. Population parameters for resistance to Fusarium graminearum and Fusarium verticillioides ear rot among large sets of early, mid-late and late maturing European maize (*Zea mays* L.) inbred lines. *Theor. Appl. Genet.* **2010**, *120*, 1053–1062.
120. Silva, E.; Mora, E.A.; Medina, A.; Vásquez, J.; Valdez, D.; Danial, D.L.; Parlevliet, J.E. Fusarium ear rot and how to screen for resistance in open pollinated maize in the Andean regions. *Euphytica* **2007**, *153*, 329–337.
121. Poole, G.J.; Smiley, R.W.; Paulitz, T.C.; Walker, C.A.; Carter, A.H.; See, D.R.; Garland-Campbell, K. Identification of quantitative trait loci (QTL) for resistance to Fusarium crown rot (*Fusarium pseudograminearum*) in multiple assay environments in the Pacific Northwestern US. *Theor. Appl. Genet.* **2012**, *125*, 91–107.
122. Wildermuth, G.B.; Morgan, J.M. Genotypic differences in partial resistance to crown rot caused by *Fusarium pseudograminearum* in relation to an osmoregulation gene in wheat. *Australas. Plant Pathol.* **2004**, *33*, 121–123.
123. Liu, Y.; Zheng, Y.; Wei, Y.; Zhou, M.; Liu, C. Genotypic differences to crown rot caused by *Fusarium pseudograminearum* in barley (*Hordeum vulgare* L.). *Plant Breed.* **2012**, *131*, 728–732.
124. Liu, Y.; Ma, J.; Yan, W.; Yan, G.; Zhou, M.; Wei, Y.; Zheng, Y.; Liu, C. Different tolerance in bread wheat, durum wheat and barley to Fusarium crown rot disease caused by *Fusarium pseudograminearum*. *J. Phytopathol.* **2012**, *160*, 412–417.
125. da Silva, W.L.; Clark, C.A. Infection of sweetpotato by *Fusarium solani* and *Macrophomina phaseolina* prior to harvest. *Plant Dis.* **2013**, *97*, 1636–1644.
126. Klingelfuss, L.H.; Yorinori, J.T.; Arias, C.A.A.; Destro, D. Reaction of soybean cultivars to sudden death syndrome and disease scoring methods for screening resistance. *Embrapa Soja-Artigo em periódico indexado* **2002**.
127. Mueller, D.S.; Hartman, G.L.; Nelson, R.L.; Pedersen, W.L. Evaluation of Glycine max germ plasm for resistance to *Fusarium solani* f. sp. glycines. *Plant Dis.* **2002**, *86*, 741–746.
128. Mueller, D.S.; Nelson, R.L.; Hartman, G.L.; Pedersen, W.L. Response of commercially developed soybean cultivars and the ancestral soybean lines to *Fusarium solani* f. sp. glycines. *Plant Dis.* **2003**, *87*, 827–831.
129. Bilgi, V.N.; Bradley, C.A.; Khot, S.D.; Grafton, K.F.; Rasmussen, J.B. Response of dry bean genotypes to *Fusarium* root rot, caused by *Fusarium solani* f. sp. phaseoli, under field and controlled conditions. *Plant Dis.* **2008**, *92*, 1197–1200.
130. Hagerty, C.H.; Cuesta-Marcos, A.; Cregan, P.B.; Song, Q.; McClean, P.; Noffsinger, S.; Myers, J.R. Mapping *Fusarium solani* and *Aphanomyces euteiches* root rot resistance and root architecture quantitative trait loci in common bean. *Crop Sci.* **2015**,

- 55, 1969–1977.
131. Schneider, K.A.; Grafton, K.F.; Kelly, J.D. QTL analysis of resistance to Fusarium root rot in bean. *Crop Sci.* **2001**, *41*, 535–542.
132. Hagedorn, D.J. Testing commercial Pea varieties for reaction to Fusarium root rot, *Fusarium solani* f. pisi. *Phytopathology* **1960**, *50*.
133. Porter, L.D.; Kraft, J.M.; Grünwald, N.J. Release of pea germplasm with Fusarium resistance combined with desirable yield and anti-lodging traits. *J. Plant Regist.* **2014**, *8*, 191–194.
134. Miedaner, T.; Bolduan, C.; Melchinger, A.E. Aggressiveness and mycotoxin production of eight isolates each of *Fusarium graminearum* and *Fusarium verticillioides* for ear rot on susceptible and resistant early maize inbred lines. *Eur. J. plant Pathol.* **2010**, *127*, 113–123.
135. De Gruyter, J.; Aveskamp, M.M.; Woudenberg, J.H.C.; Verkley, G.J.M.; Groenewald, J.Z.; Crous, P.W. Molecular phylogeny of *Phoma* and allied anamorph genera: towards a reclassification of the *Phoma* complex. *Mycol. Res.* **2009**, *113*, 508–519.
136. Kirkpatrick, T.L.; Rockroth, C.S. *Compendium of cotton diseases.*; American Phytopathological Society (APS Press), 2001; ISBN 0890542791.
137. Geldenhuis, M.M.; Roux, J.; Cilliers, A.J.; Wingfield, B.D.; Wingfield, M.J. Clonality in South African isolates and evidence for a European origin of the root pathogen *Thielaviopsis basicola*. *Mycol. Res.* **2006**, *110*, 306–311.
138. Coumans, J.V.F.; Poljak, A.; Raftery, M.J.; Backhouse, D.; Pereg-Gerk, L. Analysis of cotton (*Gossypium hirsutum*) root proteomes during a compatible interaction with the black root rot fungus *Thielaviopsis basicola*. *Proteomics* **2009**, *9*, 335–349.
139. Zaman, N.; Ahmed, S. Survey of root rot of groundnut in rainfed areas of Punjab, Pakistan. *African J. Biotechnol.* **2012**, *11*, 4791–4794.
140. Nandris, D.; Chadoeuf, J.; Pierrat, J.C.; Joannes, H.; Geiger, J.-P.; Nicole, M. Modelling rubber-tree root diseases, simulations of various inoculum rates and methods of control. *Eur. J. For. Pathol.* **1996**, *26*, 25–44.
141. Manjunatha, S. V; Naik, M.K.; Khan, M.F.R.; Goswami, R.S. Evaluation of bio-control agents for management of dry root rot of chickpea caused by *Macrophomina phaseolina*. *Crop Prot.* **2013**, *45*, 147–150.
142. Thomidis, T.; Exadaktylou, E. Effectiveness of cyproconazole to control *Armillaria* root rot of apple, walnut and kiwifruit. *Crop Prot.* **2012**, *36*, 49–51.
143. Aguín-Casal, O.; Sáinz-Osés, M.J.; Mansilla-Vázquez, J.P. *Armillaria* species infesting vineyards in northwestern Spain. *Eur. J. plant Pathol.* **2004**, *110*, 683–687.
144. Bhat, Z.A.; Sheikh, F.A.; Mubarak, T.; Bhat, J.A.; Zargar, M.A.; Wani, A.A.; Rather, G.H.; Itoo, H.U. On Farm Testing and Popularization of Integrated Management Module of Apple Root Rot Under High Altitude Temperate Conditions. *J. Krishi Vigyan* **2012**, *1*, 54–57.
145. Sánchez, M.A.G.; Cazorla, F.M.; Cayo, R.; de Vicente, A.; Jiménez, R.M.P. Studies of soil and rhizosphere bacteria to improve biocontrol of avocado white root rot caused by *Rosellinia necatrix*. *S. Michele all'Adige, Italy* **2004**, *27*, 169–172.
146. Bugbee, W.M.; Campbell, L.G. Combined resistance in sugar beet to *Rhizoctonia solani*, *Phoma betae*, and *Botrytis cinerea*. *Plant Dis.* **1990**, 353.
147. Mao, W.; Carroll, R.B.; Whittington, D.P. Association of *Phoma terrestris*, *Pythium irregulare*, and *Fusarium acuminatum* in causing red root rot of corn. *Plant Dis.* **1998**, *82*, 337–342.
148. Coleman, P.M.; Ellerbrock, L.A.; Lorbeer, J.W. Reaction of selected onion cultivars to pink root under field conditions in New York. *Plant Dis.* **1997**, *81*, 138–142.
149. Wiriyaitsomboon, P. *Characterization of Setophoma terrestris causing pink root in onion, disease management, and age-related resistance*; Michigan State University, 2015; ISBN 1339303280.
150. Hollingsworth, C.R.; Gray, F.A.; Groose, R.W. Evidence for the heritability of resistance to brown root rot of alfalfa, caused by *Phoma sclerotoides*. *Can. J. plant Pathol.* **2005**, *27*, 64–70.
151. Berkenkamp, B.; McCartney, D.; Bittman, S. Resistance of alfalfa cultivars to brown root rot. *Can. J. Plant Sci.* **1991**, *71*, 211–

- 213.
152. Kumar, V.; Joshi, S.G.; Bell, A.A.; Rathore, K.S. Enhanced resistance against *Thielaviopsis basicola* in transgenic cotton plants expressing Arabidopsis NPR1 gene. *Transgenic Res.* **2013**, *22*, 359–368.
153. Rajasekaran, K.; Cary, J.W.; Jaynes, J.M.; Cleveland, T.E. Disease resistance conferred by the expression of a gene encoding a synthetic peptide in transgenic cotton (*Gossypium hirsutum* L.) plants. *Plant Biotechnol. J.* **2005**, *3*, 545–554.
154. Wheeler, T.A.; Gannaway, J.R. Identification of germplasm resistant to *Thielaviopsis basicola* in the USDA cotton germplasm collection. In Proceedings of the World cotton conference-4; 2007; pp. 10–14.
155. Wheeler, T.A.; Gannaway, J.R.; Keating, K. Identification of resistance to *Thielaviopsis basicola* in diploid cotton. *Plant Dis.* **1999**, *83*, 831–833.
156. Bai, D.; Reeleder, R.; Brandie, J.E. Identification of two RAPD markers tightly linked with the *Nicotiana debneyi* gene for resistance to black root rot of tobacco. *Theor. Appl. Genet.* **1995**, *91*, 1184–1189.
157. Trojak-Goluch, A.; Berbec, A. Potential of *Nicotiana glauca* (Grah.) as a source of resistance to black root rot *Thielaviopsis basicola* (Berk. and Broome) Ferr. in tobacco improvement. *Plant Breed.* **2005**, *124*, 507–510.
158. Harper, J.T.; Waanders, E.; Keeling, P.J. On the monophyly of chromalveolates using a six-protein phylogeny of eukaryotes. *Int. J. Syst. Evol. Microbiol.* **2005**, *55*, 487–496.
159. van West, P.; Appiah, A.A.; Gow, N.A.R. Advances in research on oomycete root pathogens. *Physiol. Mol. Plant Pathol.* **2003**, *62*, 99–113.
160. Chamnanpant, J.; Shan, W.; Tyler, B.M. High frequency mitotic gene conversion in genetic hybrids of the oomycete *Phytophthora sojae*. *Proc. Natl. Acad. Sci.* **2001**, *98*, 14530–14535.
161. Fry, W.E.; Goodwin, S.B. Re-emergence of potato and tomato late blight in the United States. *Plant Dis.* **1997**, *81*, 1349–1357.
162. Schettini, T.M.; Legg, E.J.; Michelmore, R.W. Insensitivity to metalaxyl in California populations of *Bremia lactucae* and resistance of California lettuce cultivars to downy mildew. *Phytopathology* **1991**, *81*, 64–70.
163. Harveson, R.M.; Rush, C.M. The influence of irrigation frequency and cultivar blends on the severity of multiple root diseases in sugar beets. *Plant Dis.* **2002**, *86*, 901–908.
164. Akamatsu, H.O.; Grünwald, N.J.; Chilvers, M.I.; Porter, L.D.; Peever, T.L. Development of codominant simple sequence repeat, single nucleotide polymorphism and sequence characterized amplified region markers for the pea root rot pathogen, *Aphanomyces euteiches*. *J. Microbiol. Methods* **2007**, *71*, 82–86.
165. Taguchi, K.; Okazaki, K.; Takahashi, H.; Kubo, T.; Mikami, T. Molecular mapping of a gene conferring resistance to *Aphanomyces* root rot (black root) in sugar beet (*Beta vulgaris* L.). *Euphytica* **2010**, *173*, 409–418.
166. Campbell, L.G.; Klotz, K.L. Postharvest storage losses associated with *Aphanomyces* root rot in sugarbeet. **2006**.
167. Taguchi, K.; Ogata, N.; Kubo, T.; Kawasaki, S.; Mikami, T. Quantitative trait locus responsible for resistance to *Aphanomyces* root rot (black root) caused by *Aphanomyces cochlioides* Drechs. in sugar beet. *Theor. Appl. Genet.* **2009**, *118*, 227–234.
168. Saunders, J.W.; Mcgrath, J.M.; Theurer, J.C.; Halloin, J.M. Registration of SR87 Sugarbeet Germplasm with Low Tare and Resistances to *Cereospora* and *Aphanomyces*. *Crop Sci.* **2000**, *40*, 1833.
169. Windels, C.E. *Aphanomyces* root rot on sugar beet. *Plant Heal. Prog.* **2000**, *1*, 8.
170. Fitzpatrick, S.; Brummer, J.; Hudelson, B.; Malvick, D.; Grau, C. *Aphanomyces* root rot resistance (Races 1 and 2). In Proceedings of the Page D-2 in: Standard Tests Bull. N. Am. Alfalfa Improvement Conference; 1998.
171. Grau, C.R.; Muehlchen, A.M.; Tofte, J.E.; Smith, R.R. Variability in virulence of *Aphanomyces euteiches*. *Plant Dis.* **1991**, *75*, 1153–1156.
172. Hughes, T.J.; Grau, C.R. *Aphanomyces* root rot (common root rot) of legumes. *Aphanomyces root rot (common root rot) Legum.* **2007**.
173. McGee, R.J.; Coyne, C.J.; Pilet-Nayel, M.-L.; Moussart, A.; Tivoli, B.; Baranger, A.; Hamon, C.; Vandemark, G.; McPhee, K. Registration of pea germplasm lines partially resistant to *Aphanomyces* root rot for breeding fresh or freezer pea and dry pea

- types. *J. Plant Regist.* **2012**, *6*, 203–207.
174. Pfender, W.F.; Hagedorn, D.J. *Aphanomyces euteiches* f. sp. *phaseoli*, a causal agent of bean root and hypocotyl rot. *Phytopathology* **1982**, *72*, 306–310.
175. Fagoaga, C.; Rodrigo, I.; Conejero, V.; Hinarejos, C.; Tuset, J.J.; Arnau, J.; Pina, J.A.; Navarro, L.; Peña, L. Increased tolerance to *Phytophthora citrophthora* in transgenic orange plants constitutively expressing a tomato pathogenesis related protein PR-5. *Mol. Breed.* **2001**, *7*, 175–185.
176. Matheron, M.E.; Wright, G.C.; Porchas, M. Resistance to *Phytophthora citrophthora* and *P. parasitica* and nursery characteristics of several citrus rootstocks. *Plant Dis.* **1998**, *82*, 1217–1225.
177. Park, J.; Park, B.; Veeraraghavan, N.; Jung, K.; Lee, Y.-H.; Blair, J.E.; Geiser, D.M.; Isard, S.; Mansfield, M.A.; Nikolaeva, E. *Phytophthora* database: a forensic database supporting the identification and monitoring of *Phytophthora*. *Plant Dis.* **2008**, *92*, 966–972.
178. Thomidis, T.; Exadaktylou, E.; Sotiropoulos, T. Susceptibility of three citrus rootstocks towards *Phytophthora cactorum*, *P. citrophthora*, *P. parasitica* and *P. citricola*/Anfälligkeit dreier Citrus-Veredelungsunterlagen gegenüber *Phytophthora cactorum*, *P. citrophthora*, *P. parasitica* und *P. citricola*. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz/Journal Plant Dis. Prot.* **2005**, 204–207.
179. Tuzcu, Ö.; Cinar, A.; Göksedef, M.O.; Özsan, M.; Biçici, M. Resistance of citrus rootstocks to *Phytophthora citrophthora* during winter dormancy. *Plant Dis.* **1984**, *68*, 502–505.
180. Yildirim, B.; Yeşiloğlu, T.; Incesu, M. Fruit yield and quality of Santa Teresa lemon on seven rootstocks in Adana (Turkey). *African J. Agric. Res.* **2010**, *5*, 1077–1081.
181. Azevedo, F. Al; Mourão Filho, F.A.A.; Mendes, B.M.J.; Almeida, W.A.B.; Schinor, E.H.; Pio, R.; Barbosa, J.M.; Guidetti-Gonzalez, S.; Carrer, H.; Lam, E. Genetic transformation of Rangpur lime (*Citrus limonia osbeck*) with the *tbO* (bacterio-opsin) genes and its initial evaluation for *Phytophthora nicotianae* resistance. *Plant Mol. Biol. Report.* **2006**, *24*, 185–196.
182. Benfradj, N.; Metoui, N.; Boughalleb-M'Hamdi, N. Screening for tolerance of different citrus rootstocks against zoospores of *Phytophthora nicotianae* in infested soil. *J. Phytopathol. Pest Manag.* **2016**, *3*, 63–75.
183. Graham, J.H. Root regeneration and tolerance of citrus rootstocks to root rot caused by *Phytophthora nicotianae*. *Phytopathology* **1995**, *85*, 111–117.
184. Sakupwanya, M.N.; Labuschagne, N.; Loots, T.; Apostolides, Z. Towards developing a metabolic-marker based predictive model for *Phytophthora nicotianae* tolerance in citrus rootstocks. *J. Plant Pathol.* **2018**, *100*, 269–277.
185. Washington, W.S.; McGee, P.; Flett, S.P.; Jerie, P.H.; Ashcroft, W.J. Cultivars and fungicides affect *Phytophthora* root rot in processing tomatoes. *Australas. Plant Pathol.* **2001**, *30*, 309–315.
186. Carisse, O.; Khanizadeh, S. Relative resistance of newly released apple rootstocks to *Phytophthora cactorum*. *Can. J. plant Sci.* **2006**, *86*, 199–204.
187. Eikemo, H.; Brurberg, M.B.; Davik, J. Resistance to *Phytophthora cactorum* in diploid *Fragaria* species. *HortScience* **2010**, *45*, 193–197.
188. McIntosh, E.D.L. of apple trees. *Can. Plant Dis. Surv.* **1975**, *55*, 109–116.
189. Mangandi, J.; Verma, S.; Osorio, L.; Peres, N.A.; van de Weg, E.; Whitaker, V.M. Pedigree-based analysis in a multiparental population of octoploid strawberry reveals QTL alleles conferring resistance to *Phytophthora cactorum*. *G3 Genes, Genomes, Genet.* **2017**, *7*, 1707–1719.
190. Sewell, G.W.F.; Wilson, J.F. Resistance Trials of Some Apple Rootstock Varieties to *Phytophthora Cactorum* (L. & C.) Schroet. *J. Hortic. Sci.* **1959**, *34*, 51–58.
191. Utkhede, R.S.; Quamme, H.A. Use of the excised shoot assay to evaluate resistance to *Phytophthora cactorum* of apple rootstock cultivars. *Can. J. Plant Sci.* **1988**, *68*, 851–857.
192. Douhan, G.W.; Fuller, E.; McKee, B.; Pond, E. Genetic diversity analysis of avocado (*Persea americana* Miller) rootstocks

- selected under greenhouse conditions for tolerance to phytophthora root rot caused by *Phytophthora cinnamomi*. *Euphytica* **2011**, *182*, 209.
193. Haymes, K.M.; Van de Weg, W.E.; Arens, P.; Maas, J.L.; Vosman, B.; Den Nijs, A.P.M. Development of SCAR markers linked to a *Phytophthora fragariae* resistance gene and their assessment in European and North American strawberry genotypes. *J. Am. Soc. Hortic. Sci.* **2000**, *125*, 330–339.
194. Rugienius, R.; Siksnianas, T.; Stanys, V.; Gelvonauskiene, D.; Bendokas, V. Use of RAPD and SCAR markers for identification of strawberry genotypes carrying red stele (*Phytophthora fragariae*) resistance gene Rpf1. *Agron Res* **2006**, *4*, 335–339.
195. Pattison, J.A.; Wilcox, W.F.; Weber, C.A. Assessing the resistance of red raspberry (*Rubus idaeus* L.) genotypes to *Phytophthora fragariae* var. *rubi* in hydroponic culture. *HortScience* **2004**, *39*, 1553–1556.
196. Weber, C.A.; Pattison, J.; Samuelian, S. Marker assisted selection for resistance to root rot in red raspberry caused by *Phytophthora fragariae* var. *rubi*. In Proceedings of the IX International Rubus and Ribes Symposium 777; 2005; pp. 311–316.
197. Burnham, K.D.; Dorrance, A.E.; VanToai, T.T.; St. Martin, S.K. Quantitative trait loci for partial resistance to *Phytophthora sojae* in soybean. *Crop Sci.* **2003**, *43*, 1610–1617.
198. Dorrance, A.E.; Schmitthenner, A.F. New sources of resistance to *Phytophthora sojae* in the soybean plant introductions. *Plant Dis.* **2000**, *84*, 1303–1308.
199. Bosland, P.W.; Lindsey, D.L. *A seedling screen for Phytophthora root rot of pepper, Capsicum annuum*; 1991;
200. Kim, H.-J.; Nahm, S.-H.; Lee, H.-R.; Yoon, G.-B.; Kim, K.-T.; Kang, B.-C.; Choi, D.; Kweon, O.Y.; Cho, M.-C.; Kwon, J.-K. BAC-derived markers converted from RFLP linked to *Phytophthora capsici* resistance in pepper (*Capsicum annuum* L.). *Theor. Appl. Genet.* **2008**, *118*, 15.
201. Oguniwin, E.A.; Berke, T.F.; Massoudi, M.; Black, L.L.; Huestis, G.; Choi, D.; Lee, S.; Prince, J.P. Construction of 2 intraspecific linkage maps and identification of resistance QTLs for *Phytophthora capsici* root-rot and foliar-blight diseases of pepper (*Capsicum annuum* L.). *Genome* **2005**, *48*, 698–711.
202. Cerra, S.M. *Phytophthora* root and stem rot of soybean in Iowa: Minimizing losses through an improved understanding of population structure and implementation of novel management strategies. **2007**.
203. Dale, M.L.; Irwin, J.A.G. Glasshouse and field screening of chickpea cultivars for resistance to *Phytophthora megasperma* f. sp. *medicaginis*. *Aust. J. Exp. Agric.* **1991**, *31*, 663–667.
204. Knights, E.J.; Southwell, R.J.; Schwinghamer, M.W.; Harden, S. Resistance to *Phytophthora medicaginis* Hansen and Maxwell in wild *Cicer* species and its use in breeding root rot resistant chickpea (*Cicer arietinum* L.). *Aust. J. Agric. Res.* **2008**, *59*, 383–387.
205. Vandemark, G.J.; Barker, B.M. Quantifying *Phytophthora medicaginis* in susceptible and resistant alfalfa with a real-time fluorescent PCR assay. *J. Phytopathol.* **2003**, *151*, 577–583.
206. Wiersma, D.W.; Grau, C.R.; Undersander, D.J. Alfalfa cultivar performance with differing levels of resistance to *Phytophthora* and *Aphanomyces* root rots. *J. Prod. Agric.* **1995**, *8*, 259–264.
207. Abrinbana, M.; Babai-Ahary, A.; Heravan, I.M. Assessment of resistance in sugarbeet lines to damping-off caused by *Pythium ultimum* Trow var. *ultimum* under greenhouse conditions. *Plant Pathol. J.* **2007**.
208. Bates, G.D.; Rothrock, C.S.; Rupe, J.C. Resistance of the Soybean Cultivar Archer to *Pythium* Damping-Off and Root Rot Caused by Several *Pythium* spp. *Plant Dis.* **2008**, *92*, 763–766.
209. Balk, C.S. Assessment of resistance in soybean to *Pythium ultimum* and sensitivity of Ohio's diverse *Pythium* species towards metalaxyl 2014.
210. Cheng, L. *Pythium ultimum*. *NC State Univ. Dep. Plant Pathol.* **2007**.
211. Lucas, B.; Griffiths, P.D. Evaluation of common bean accessions for resistance to *Pythium ultimum*. *HortScience* **2004**, *39*, 1193–1195.
212. Ellis, M.L.; McHale, L.K.; Paul, P.A.; St Martin, S.K.; Dorrance, A.E. Soybean germplasm resistant to *Pythium irregulare* and

- molecular mapping of resistance quantitative trait loci derived from the soybean accession PI 424354. *Crop Sci.* **2013**, *53*, 1008–1021.
213. Farr, D.F.; Rossman, A.Y.; Palm, M.E.; McCray, E.B. Fungal databases, systematic botany & mycology laboratory, ARS, USDA. 2007 2008.
214. Katawczik, M. *Pythium irregulare*. *NCState Dep. Plant Pathol.* **2008**.
215. Aliyu, T.H.; Balogun, O.S.; Adesina, O.M. EFFECT OF PYTHIUM APHANIDERMATUM ON TWO CULTIVARS OF PEPPER (*Capsicum* spp). **2012**.
216. FATTAHI, S.H.; Zafari, D.; Mahmoudi, B. Evaluation of superior sugar beet genotypes for resistance to important root rot pathogens in the greenhouse. **2011**.
217. Mahmoudi, S.B.; Koulaei, H.E.; Hasani, M.; Alaghebandzade, N.; Soltani, J.; Kakueinezhad, M. Development of sugar beet S1 pollinator lines resistant to *Pythium* root rot. In Proceedings of the Proceedings of the 1st International and 13th Iranian Crop Science Congress 3rd Iranian Seed Science and Technology Conference. Seed and Plant improvement Institute Karaj, Iran; 2014.
218. Parker, K.C. *Pythium aphanidermatum*. *Soilborne plant Pathog. NC State Univ. Consult.* **2009**, *20*.
219. Rosso, M.L.; Rupe, J.C.; Chen, P.; Mozzoni, L.A. Inheritance and genetic mapping of resistance to *Pythium* damping-off caused by *Pythium aphanidermatum* in 'Archer' soybean. *Crop Sci.* **2008**, *48*, 2215–2222.
220. Richard, C.; Beghdadi, A.; Martin, J.G. *Aphanomyces euteiches*, a novel root pathogen to alfalfa in Québec. *Plant Dis.* **1991**, *75*.
221. Tofte, J.E.; Smith, R.R.; Grau, C.R. Reaction of red clover to *Aphanomyces euteiches*. *Plant Dis.* **1992**, *76*, 39–42.
222. Van Leur, J.A.G.; Southwell, R.J.; Mackie, J.M. *Aphanomyces* root rot on faba bean in northern NSW. *Australas. Plant Dis. Notes* **2008**, *3*, 8–9.
223. Chen, W.; Sharma, H.C.; Muehlbauer, F.J. *Compendium of chickpea and lentil diseases and pests.*; American Phytopathological Society (APS Press), 2011; ISBN 0890543836.
224. Beckerman, J. Disease Management Strategies for Horticultural Crops: *Pythium* Root rot of Herbaceous Plants. *Purdue Univ.* **2011**.
225. Nzungize, J.; Gepts, P.; Buruchara, R.; Male, A.; Ragama, P.; Busogoro, J.P.; Baudoin, J.-P. Introgression of *Pythium* root rot resistance gene into Rwandan susceptible common bean cultivars. *African J. Plant Sci.* **2011**, *5*, 193–200.
226. Broders, K.D.; Wallhead, M.W.; Austin, G.D.; Lipps, P.E.; Paul, P.A.; Mullen, R.W.; Dorrance, A.E. Association of soil chemical and physical properties with *Pythium* species diversity, community composition, and disease incidence. *Phytopathology* **2009**, *99*, 957–967.
227. Broders, K.D.; Lipps, P.E.; Paul, P.A.; Dorrance, A.E. Characterization of *Pythium* spp. associated with corn and soybean seed and seedling disease in Ohio. *Plant Dis.* **2007**, *91*, 727–735.
228. Menzies, J.G.; Ehret, D.L.; Stan, S. Effect of inoculum density of *Pythium aphanidermatum* on the growth and yield of cucumber plants grown in recirculating nutrient film culture. *Can. J. Plant Pathol.* **1996**, *18*, 50–54.
229. Utkhede, R.S.; Koch, C.A. Rhizobacterial growth and yield promotion of cucumber plants inoculated with *Pythium aphanidermatum*. *Can. J. plant Pathol.* **1999**, *21*, 265–271.
230. Utkhede, R.S.; Levesque, C.A.; Dinh, D. *Pythium aphanidermatum* root rot in hydroponically grown lettuce and the effect of chemical and biological agents on its control. *Can. J. Plant Pathol.* **2000**, *22*, 138–144.
231. Kakueinezhad, M.; Taheri, P.; Mahmoudi, S.B.; Tarighi, S. Resistance assessment and biochemical responses of sugar beet lines against *Pythium aphanidermatum*, causing root rot. *Eur. J. Plant Pathol.* **2018**, *151*, 307–319.
232. Petkowski, J.E.; de Boer, R.F.; Norng, S.; Thomson, F.; Minchinton, E.J. *Pythium* species associated with root rot complex in winter-grown parsnip and parsley crops in south eastern Australia. *Australas. Plant Pathol.* **2013**, *42*, 403–411.
233. Mavrodi, O. V.; Walter, N.; Elateek, S.; Taylor, C.G.; Okubara, P.A. Suppression of *Rhizoctonia* and *Pythium* root rot of wheat

- by new strains of *Pseudomonas*. *Biol. Control* **2012**, *62*, 93–102.
234. Dissanayake, N.; Hoy, J.W.; Griffin, J.L. Herbicide effects on sugarcane growth, Pythium root rot, and Pythium arrhenomanes. *Phytopathology* **1998**, *88*, 530–535.
235. Elmer, W.H.; Gent, M.P.N.; McAvoy, R.J. Partial saturation under ebb and flow irrigation suppresses Pythium root rot of ornamentals. *Crop Prot.* **2012**, *33*, 29–33.
236. Hardham, A.R.; Blackman, L.M. Molecular cytology of Phytophthora-plant interactions. *Australas. Plant Pathol.* **2010**, *39*, 29–35.
237. Kroon, L.P.N.M.; Brouwer, H.; de Cock, A.W.A.M.; Govers, F. The genus Phytophthora anno 2012. *Phytopathology* **2012**, *102*, 348–364.
238. Dirac, M.F.; Menge, J.A. High temperatures are not responsible for lack of infection of citrus roots by Phytophthora citrophthora during the summer, but suppressive soil microorganisms may inhibit infection by P. citrophthora. *Plant Soil* **2002**, *241*, 243–249.
239. Alvarez, L.A.; Vicent, A.; De la Roca, E.; Bascón, J.; Abad-Campos, P.; Armengol, J.; García-Jiménez, J. Branch cankers on citrus trees in Spain caused by Phytophthora citrophthora. *Plant Pathol.* **2008**, *57*, 84–91.
240. Graham, J.H.; Menge, J.A. Root diseases. *Citrus Heal. Manag.* **1999**, 126–135.
241. Bekker, T.F.; Kaiser, C.; Labuschagne, N. Efficacy of water soluble silicon against Phytophthora cinnamomi root rot of avocado: A progress report. *South African Avocado Grow. Assoc. Yearb.* **2006**, *29*, 58–62.
242. Anderson, J.M.; Pegg, K.G.; Scott, C.; Drenth, A. Phosphonate applied as a pre-plant dip controls Phytophthora cinnamomi root and heart rot in susceptible pineapple hybrids. *Australas. Plant Pathol.* **2012**, *41*, 59–68.
243. Pérombelon, M.C.M. Potato diseases caused by soft rot erwinias: an overview of pathogenesis. *Plant Pathol.* **2002**, *51*, 1–12.
244. Zhao, Y.; Li, P.; Huang, K.; Wang, Y.; Hu, H.; Sun, Y. Control of postharvest soft rot caused by Erwinia carotovora of vegetables by a strain of Bacillus amyloliquefaciens and its potential modes of action. *World J. Microbiol. Biotechnol.* **2013**, *29*, 411–420.
245. Collmer, A.; Keen, N.T. The role of pectic enzymes in plant pathogenesis. *Annu. Rev. Phytopathol.* **1986**, *24*, 383–409.
246. Charkowski, A.O. The soft rot Erwinia. In *Plant-associated bacteria*; Springer, 2007; pp. 423–505.
247. Liao, C.-H. Analysis of pectate lyases produced by soft rot bacteria associated with spoilage of vegetables. *Appl. Environ. Microbiol.* **1989**, *55*, 1677–1683.
248. Huang, L.F.; Fang, B.P.; Luo, Z.X.; Chen, J.Y.; Zhang, X.J.; Wang, Z.Y. First report of bacterial stem and root rot of sweetpotato caused by a Dickeya sp. (Erwinia chrysanthemi) in China. *Plant Dis.* **2010**, *94*, 1503.
249. Muimba-Kankolongo, A. *Food Crop Production by Smallholder Farmers in Southern Africa: Challenges and Opportunities for Improvement*; Academic Press, 2018; ISBN 0128143843.
250. Bigirimana, S.; Barumbanze, P.; Ndayinzamaso, P.; Shirima, R.; Legg, J.P. First report of cassava brown streak disease and associated Ugandan cassava brown streak virus in Burundi. *New Dis. Reports* **2011**, *24*, 588–2044.
251. Grant, M.; Lamb, C. Systemic immunity. *Curr. Opin. Plant Biol.* **2006**, *9*, 414–420.
252. Takemoto, D.; Jones, D.A. Particle bombardment-mediated transient expression to identify localization signals in plant disease resistance proteins and target sites for the proteolytic activity of pathogen effectors. In *Plant-Pathogen Interactions*; Springer, 2014; pp. 91–101.
253. Hammond-Kosack, K.E.; Jones, J.D.G. Plant disease resistance genes. *Annu. Rev. Plant Biol.* **1997**, *48*, 575–607.
254. Harris, C.J.; Sloatweg, E.J.; Goverse, A.; Baulcombe, D.C. Stepwise artificial evolution of a plant disease resistance gene. *Proc. Natl. Acad. Sci.* **2013**, *110*, 21189–21194.
255. Gaskill, J.O. Breeding for Rhizoctonia resistance in sugarbeet. *J Am Sugar Beet Technol* **1968**, *15*, 107–119.
256. Panella, L. Registration of FC709-2 and FC727 sugarbeet germplasms resistant to Rhizoctonia root rot and Cercospora leaf spot. *Crop Sci.* **1999**, *39*, 298–299.

257. Lein, J.C.; Sagstetter, C.M.; Schulte, D.; Thurau, T.; Varrelmann, M.; Saal, B.; Koch, G.; Borchardt, D.C.; Jung, C. Mapping of rhizoctonia root rot resistance genes in sugar beet using pathogen response-related sequences as molecular markers. *Plant Breed.* **2008**, *127*, 602–611.
258. Hecker, R.J.; Ruppel, E.G. Inheritance of Resistance to Rhizoctonia Root Rot in Sugarbeet 1. *Crop Sci.* **1975**, *15*, 487–490.
259. Peng, H.; Chen, Z.; Fang, Z.; Zhou, J.; Xia, Z.; Gao, L.; Chen, L.; Li, L.; Li, T.; Zhai, W. Rice Xa21 primed genes and pathways that are critical for combating bacterial blight infection. *Sci. Rep.* **2015**, *5*, 12165.
260. Seo, Y.-S.; Chern, M.; Bartley, L.E.; Han, M.; Jung, K.-H.; Lee, I.; Walia, H.; Richter, T.; Xu, X.; Cao, P. Towards establishment of a rice stress response interactome. *PLoS Genet* **2011**, *7*, e1002020.
261. Peng, Y.; Bartley, L.E.; Canlas, P.; Ronald, P.C. OsWRKY IIa transcription factors modulate rice innate immunity. *Rice* **2010**, *3*, 36–42.
262. Choi, J.; Choi, D.; Lee, S.; Ryu, C.-M.; Hwang, I. Cytokinins and plant immunity: old foes or new friends? *Trends Plant Sci.* **2011**, *16*, 388–394.
263. Argueso, C.T.; Ferreira, F.J.; Epple, P.; To, J.P.C.; Hutchison, C.E.; Schaller, G.E.; Dangl, J.L.; Kieber, J.J. Two-component elements mediate interactions between cytokinin and salicylic acid in plant immunity. *PLoS Genet* **2012**, *8*, e1002448.
264. Kim, Y.J.; Lin, N.-C.; Martin, G.B. Two distinct Pseudomonas effector proteins interact with the Pto kinase and activate plant immunity. *Cell* **2002**, *109*, 589–598.
265. Tang, X.; Xie, M.; Kim, Y.J.; Zhou, J.; Klessig, D.F.; Martin, G.B. Overexpression of Pto activates defense responses and confers broad resistance. *Plant Cell* **1999**, *11*, 15–29.
266. Zamora, M.G.M.; Castagnaro, A.P.; Ricci, J.C.D. Genetic diversity of Pto-like serine/threonine kinase disease resistance genes in cultivated and wild strawberries. *J. Mol. Evol.* **2008**, *67*, 211–221.
267. Webb, K.M.; Freeman, C.; Broeckling, C.D. Metabolome profiling to understand the defense response of sugar beet (*Beta vulgaris*) to *Rhizoctonia solani* AG 2-2 IIIB. *Physiol. Mol. Plant Pathol.* **2016**, *94*, 108–117.
268. Aliferis, K.A.; Jabaji, S. FT-ICR/MS and GC-EI/MS metabolomics networking unravels global potato sprout's responses to *Rhizoctonia solani* infection. *PLoS One* **2012**, *7*.
269. Bednarek, P.; Osbourn, A. Plant-microbe interactions: chemical diversity in plant defense. *Science (80-.)*. **2009**, *324*, 746–748.
270. Sparg, S.; Light, M.E.; Van Staden, J. Biological activities and distribution of plant saponins. *J. Ethnopharmacol.* **2004**, *94*, 219–243.
271. Zhao, C.-J.; Wang, A.-R.; Shi, Y.-J.; Wang, L.-Q.; Liu, W.-D.; Wang, Z.-H.; Lu, G.-D. Identification of defense-related genes in rice responding to challenge by *Rhizoctonia solani*. *Theor. Appl. Genet.* **2008**, *116*, 501–516.
272. Bertini, L.; Palazzi, L.; Proietti, S.; Pollastri, S.; Arrigoni, G.; Polverino de Laureto, P.; Caruso, C. Proteomic analysis of MeJa-induced defense responses in rice against wounding. *Int. J. Mol. Sci.* **2019**, *20*, 2525.
273. Van Loon, L.C.; Bakker, P.; Pieterse, C.M.J. Systemic resistance induced by rhizosphere bacteria. *Annu. Rev. Phytopathol.* **1998**, *36*, 453–483.
274. Kim, S.G.; Kim, S.T.; Wang, Y.; Yu, S.; Choi, I.S.; Kim, Y.C.; Kim, W.T.; Agrawal, G.K.; Rakwal, R.; Kang, K.Y. The RNase activity of rice probenazole-induced protein1 (PBZ1) plays a key role in cell death in plants. *Mol. Cells* **2011**, *31*, 25–31.
275. Park, C.; Kim, K.; Shin, R.; Park, J.M.; Shin, Y.; Paek, K. Pathogenesis-related protein 10 isolated from hot pepper functions as a ribonuclease in an antiviral pathway. *Plant J.* **2004**, *37*, 186–198.
276. Chen, S.; Vaghchhipawala, Z.; Li, W.; Asard, H.; Dickman, M.B. Tomato phospholipid hydroperoxide glutathione peroxidase inhibits cell death induced by Bax and oxidative stresses in yeast and plants. *Plant Physiol.* **2004**, *135*, 1630–1641.
277. Shrestha, C.L.; Ona, I.; Muthukrishnan, S.; Mew, T.W. Chitinase levels in rice cultivars correlate with resistance to the sheath blight pathogen *Rhizoctonia solani*. *Eur. J. plant Pathol.* **2008**, *120*, 69–77.
278. Taheri, P.; Tarighi, S. Riboflavin induces resistance in rice against *Rhizoctonia solani* via jasmonate-mediated priming of phenylpropanoid pathway. *J. Plant Physiol.* **2010**, *167*, 201–208.

279. Punja, Z.K. Genetic engineering of plants to enhance resistance to fungal pathogens—a review of progress and future prospects. *Can. J. Plant Pathol.* **2001**, *23*, 216–235.
280. Lehtonen, M.J.; Somervuo, P.; Valkonen, J.P.T. Infection with *Rhizoctonia solani* induces defense genes and systemic resistance in potato sprouts grown without light. *Phytopathology* **2008**, *98*, 1190–1198.
281. Takemoto, D.; Furuse, K.; Doke, N.; Kazuhito, K. Identification of chitinase and osmotin-like protein as actin-binding proteins in suspension-cultured potato cells. *Plant cell Physiol.* **1997**, *38*, 441–448.
282. Lorito, M.; Woo, S.L.; Fernandez, I.G.; Colucci, G.; Harman, G.E.; Pintor-Toro, J.A.; Filippone, E.; Muccifora, S.; Lawrence, C.B.; Zoina, A. Genes from mycoparasitic fungi as a source for improving plant resistance to fungal pathogens. *Proc. Natl. Acad. Sci.* **1998**, *95*, 7860–7865.
283. Chandra, A.; Saxena, R.; Dubey, A.; Saxena, P. Change in phenylalanine ammonia lyase activity and isozyme patterns of polyphenol oxidase and peroxidase by salicylic acid leading to enhance resistance in cowpea against *Rhizoctonia solani*. *Acta Physiol. Plant.* **2007**, *29*, 361–367.
284. Wen, K.; Seguin, P.; St.-Arnaud, M.; Jabaji-Hare, S. Real-time quantitative RT-PCR of defense-associated gene transcripts of *Rhizoctonia solani*-infected bean seedlings in response to inoculation with a nonpathogenic binucleate *Rhizoctonia* isolate. *Phytopathology* **2005**, *95*, 345–353.
285. Foley, R.C.; Gleason, C.A.; Anderson, J.P.; Hamann, T.; Singh, K.B. Genetic and genomic analysis of *Rhizoctonia solani* interactions with *Arabidopsis*; evidence of resistance mediated through NADPH oxidases. *PLoS One* **2013**, *8*.
286. Li, H.; Smigocki, A.C. Sugar beet polygalacturonase-inhibiting proteins with 11 LRRs confer *Rhizoctonia*, *Fusarium* and *Botrytis* resistance in *Nicotiana* plants. *Physiol. Mol. Plant Pathol.* **2018**, *102*, 200–208.
287. Borrás-Hidalgo, O.; Caprari, C.; Hernández-Estevéz, I.; De Lorenzo, G.; Cervone, F. A gene for plant protection: expression of a bean polygalacturonase inhibitor in tobacco confers a strong resistance against *Rhizoctonia solani* and two oomycetes. *Front. Plant Sci.* **2012**, *3*, 268.
288. Kalantari, M.; Motallebi, M.; Zamani, M.R. Bean Polygalacturonase-Inhibiting Protein Expressed in Transgenic Sugar Beet Inhibits Polygalacturonase from *Rhizoctonia Solani*. *Biosci. Biotechnol. Res. Asia* **2016**, *8*, 19–28.
289. Helliwell, E.E.; Wang, Q.; Yang, Y. Transgenic rice with inducible ethylene production exhibits broad-spectrum disease resistance to the fungal pathogens *Magnaporthe oryzae* and *Rhizoctonia solani*. *Plant Biotechnol. J.* **2013**, *11*, 33–42.
290. Maeda, S.; Dubouzet, J.G.; Kondou, Y.; Jikumaru, Y.; Seo, S.; Oda, K.; Matsui, M.; Hirochika, H.; Mori, M. The rice CYP78A gene BSR2 confers resistance to *Rhizoctonia solani* and affects seed size and growth in *Arabidopsis* and rice. *Sci. Rep.* **2019**, *9*, 1–14.
291. Kalpana, K.; Maruthasalam, S.; Rajesh, T.; Poovannan, K.; Kumar, K.K.; Kokiladevi, E.; Raja, J.A.J.; Sudhakar, D.; Velazhahan, R.; Samiyappan, R. Engineering sheath blight resistance in elite indica rice cultivars using genes encoding defense proteins. *Plant Sci.* **2006**, *170*, 203–215.
292. Gill, U.S.; Lee, S.; Mysore, K.S. Host versus nonhost resistance: distinct wars with similar arsenals. *Phytopathology* **2015**, *105*, 580–587.
293. Hadwiger, L.A.; Beckman, J.M. Chitosan as a component of pea-*Fusarium solani* interactions. *Plant Physiol.* **1980**, *66*, 205–211.
294. Isaac, J.; Hartney, S.L.; Druffel, K.; Hadwiger, L.A. The non-host disease resistance response in peas; alterations in phosphorylation and ubiquitination of HMG A and histones H2A/H2B. *Plant Sci.* **2009**, *177*, 439–449.
295. Hadwiger, L.A. Pea-*Fusarium solani* interactions contributions of a system toward understanding disease resistance. *Phytopathology* **2008**, *98*, 372–379.
296. Hadwiger, L.A. Anatomy of a nonhost disease resistance response of pea to *Fusarium solani*: PR gene elicitation via DNase, chitosan and chromatin alterations. *Front. Plant Sci.* **2015**, *6*, 373.
297. Kendra, D.F.; Hadwiger, L.A. Characterization of the smallest chitosan oligomer that is maximally antifungal to *Fusarium solani* and elicits pisatin formation in *Pisum sativum*. *Exp. Mycol.* **1984**, doi:10.1016/0147-5975(84)90013-6.

298. Klosterman, S.J.; Chen, J.; Choi, J.J.; Chinn, E.E.; Hadwiger, L.A. Characterization of a 20 kDa DNase elicitor from *Fusarium solani* f. sp. *phaseoli* and its expression at the onset of induced resistance in *Pisum sativum*. *Mol. Plant Pathol.* **2001**, *2*, 147–158.
299. Williamson-Benavides, B.A.; Sharpe, R.M.; Nelson, G.M.; Bodah, E.T.; Porter, L.D.; Dhingra, A. Identification of *Fusarium solani* f. sp. *pisii* (Fsp) responsive genes in *Pisum sativum*. *Front. Genet.* **2020**, *11*, 950, doi:10.3389/FGENE.2020.00950.
300. Almeida, M.S.; Cabral, K.M.S.; Zingali, R.B.; Kurtenbach, E. Characterization of two novel defense peptides from pea (*Pisum sativum*) seeds. *Arch. Biochem. Biophys.* **2000**, *378*, 278–286.
301. Mauch, F.; Hadwiger, L.A.; Boller, T. Ethylene: symptom, not signal for the induction of chitinase and β -1, 3-glucanase in pea pods by pathogens and elicitors. *Plant Physiol.* **1984**, *76*, 607–611.
302. Chang, M.-M.; Hadwiger, L.A.; Horovitz, D. Molecular characterization of a pea β -1, 3-glucanase induced by *Fusarium solani* and chitosan challenge. *Plant Mol. Biol.* **1992**, *20*, 609–618.
303. Hadwiger, L.A.; Adams, M.J. Nuclear changes associated with the host-parasite interaction between *Fusarium solani* and peas. *Physiol. Plant Pathol.* **1978**, *12*, 63–72.
304. Lozovaya, V. V.; Lygin, A. V.; Zernova, O. V.; Li, S.; Hartman, G.L.; Widholm, J.M. Isoflavonoid accumulation in soybean hairy roots upon treatment with *Fusarium solani*. *Plant Physiol. Biochem.* **2004**, *42*, 671–679.
305. Iqbal, M.J.; Yaegashi, S.; Ahsan, R.; Shopinski, K.L.; Lightfoot, D.A. Root response to *Fusarium solani* f. sp. *glycines*: temporal accumulation of transcripts in partially resistant and susceptible soybean. *Theor. Appl. Genet.* **2005**, *110*, 1429–1438.
306. Mackintosh, S.F.; Matthews, D.E.; VanEtten, H.D. Two additional genes for pisatin demethylation and their relationship to the pathogenicity of *Nectria haematococca* on pea. *Mol. Plant-Microbe Interact* **1989**, *2*, 354–362.
307. Coyne, C.J.; Porter, L.D.; Boutet, G.; Ma, Y.; McGee, R.J.; Lesné, A.; Baranger, A.; Pilet-Nayel, M.-L. Confirmation of *Fusarium* root rot resistance QTL Fsp-Ps 2.1 of pea under controlled conditions. *BMC Plant Biol.* **2019**, *19*, 98.
308. Coyne, C.J.; Pilet-Nayel, M.; McGee, R.J.; Porter, L.D.; Smýkal, P.; Grünwald, N.J. Identification of QTL controlling high levels of partial resistance to *Fusarium solani* f. sp. *pisii* in pea. *Plant Breed.* **2015**, *134*, 446–453.
309. Feng, J.; Hwang, R.; Chang, K.F.; Conner, R.L.; Hwang, S.F.; Strelkov, S.E.; Gossen, B.D.; McLaren, D.L.; Xue, A.G. Identification of microsatellite markers linked to quantitative trait loci controlling resistance to *Fusarium* root rot in field pea. **2011**, *91*, 199–204, doi:10.4141/CJPS09176.
310. Williamson-Benavides, B.; Sharpe, R.; Nelson, G.; Bodah, E.T.; Porter, L.D.; Amit Dhingra1, 2§Bruce A. Williamson-Benavides1, Richard Sharpe2, Grant Nelson1, Eliane T. Bodah3, Lyndon D. Porter4, and Amit Dhingra1, 2§Bruce A. Williamson-Benavides1, Richard Sharpe2, Grant Nelson1, Eliane T. Bodah3, Lyndon D. Porter4, and Amit Dhingra, 2§ Identification of root rot tolerance QTLs in pea using *Fusarium solani* f. sp. *pisii*-responsive differentially expressed genes. *bioRxiv* **2021**, In Review, doi:10.1101/2020.11.13.382077.
311. Badrhaddad, A.; Nazarian-Firouzabadi, F.; Ismaili, A. Fusion of a chitin-binding domain to an antibacterial peptide to enhance resistance to *Fusarium solani* in tobacco (*Nicotiana tabacum*). *3 Biotech* **2018**, *8*, 391.
312. Charfeddine, M.; Samet, M.; Charfeddine, S.; Bouaziz, D.; Bouzid, R.G. Ectopic Expression of StERF94 Transcription Factor in Potato Plants Improved Resistance to *Fusarium solani* Infection. *Plant Mol. Biol. Report.* **2019**, *37*, 450–463.
313. Zhang, C.; Zhao, X.; Qu, Y.; Teng, W.; Qiu, L.; Zheng, H.; Wang, Z.; Han, Y.; Li, W. Loci and candidate genes in soybean that confer resistance to *Fusarium graminearum*. *Theor. Appl. Genet.* **2019**, *132*, 431–441.
314. Nelsen, N.S.; Li, Z.; Warner, A.L.; Matthews, B.F.; Knap, H.T. Genomic Polymorphism Identifies a Subtilisin-Like Protease near the *Rhg4* Locus in Soybean. *Crop Sci.* **2004**, *44*, 265–273.
315. Million, C.R.; Wijeratne, S.; Cassone, B.J.; Lee, S.; Rouf Mian, M.A.; McHale, L.K.; Dorrance, A.E. Hybrid genome assembly of a major quantitative disease resistance locus in soybean toward *Fusarium graminearum*. *Plant Genome* **2019**, *12*, 1–17.
316. Scherm, B.; Balmas, V.; Spanu, F.; Pani, G.; Delogu, G.; Pasquali, M.; Migheli, Q. *Fusarium culmorum*: causal agent of foot and root rot and head blight on wheat. *Mol. Plant Pathol.* **2013**, *14*, 323–341.

317. Osborne, L.E.; Stein, J.M. Epidemiology of Fusarium head blight on small-grain cereals. *Int. J. Food Microbiol.* **2007**, *119*, 103–108.
318. Aoki, T.; Ward, T.J.; Kistler, H.C.; O'donnell, K. Systematics, phylogeny and trichothecene mycotoxin potential of Fusarium head blight cereal pathogens. *JSM Mycotoxins* **2012**, *62*, 91–102.
319. Mesterhazy, A.; Lemmens, M.; Reid, L.M. Breeding for resistance to ear rots caused by Fusarium spp. in maize—a review. *Plant Breed.* **2012**, *131*, 1–19.
320. Saharan, M.S. Current status of resistant source to Fusarium head blight disease of wheat: a review. *Indian Phytopathol.* **2020**, *73*, 3–9.
321. Gilbert, J.; Haber, S. Overview of some recent research developments in Fusarium head blight of wheat. *Can. J. Plant Pathol.* **2013**, *35*, 149–174.
322. Lanubile, A.; Muppirala, U.K.; Severin, A.J.; Marocco, A.; Munkvold, G.P. Transcriptome profiling of soybean (*Glycine max*) roots challenged with pathogenic and non-pathogenic isolates of *Fusarium oxysporum*. *BMC Genomics* **2015**, *16*, 1089.
323. Kolb, F.L.; Bai, G.H.; Muehlbauer, G.J.; Anderson, J.A.; Smith, K.P.; Fedak, G. Symposium on genetic solutions to Fusarium head blight in wheat and barley: Challenges, opportunities, and imperatives. *Crop Sci.* **2001**, *41*, 611.
324. Rudd, J.C.; Horsley, R.D.; McKendry, A.L.; Elias, E.M. Host plant resistance genes for Fusarium head blight: sources, mechanisms, and utility in conventional breeding systems. *Crop Sci.* **2001**, *41*, 620–627.
325. Buerstmayr, H.; Ban, T.; Anderson, J.A. QTL mapping and marker-assisted selection for Fusarium head blight resistance in wheat: a review. *Plant Breed.* **2009**, *128*, 1–26.
326. Ma, Z.; Xie, Q.; Li, G.; Jia, H.; Zhou, J.; Kong, Z.; Li, N.; Yuan, Y. Germplasm, genetics and genomics for better control of disastrous wheat Fusarium head blight. *Theor. Appl. Genet.* **2020**, 1–28.
327. Fugate, K.K.; Ferrareze, J.P.; Bolton, M.D.; Deckard, E.L.; Campbell, L.G. Postharvest jasmonic acid treatment of sugarbeet roots reduces rot due to *Botrytis cinerea*, *Penicillium claviforme*, and *Phoma betae*. *Postharvest Biol. Technol.* **2012**, *65*, 1–4.
328. Bugbee, W.M.; Cole, D.F. Comparison of thiabendazole and genetic resistance for control of sugar beet storage rot. *Phytopathology* **1979**, *69*, 1230–1232.
329. Wang, H.; Hwang, S.F.; Chang, K.F.; Gossen, B.D.; Turnbull, G.D.; Howard, R.J. Assessing resistance to spring black stem and leaf spot of alfalfa caused by *Phoma* spp. *Can. J. Plant Sci.* **2004**, *84*, 311–317.
330. Marzu, J.C.; Straley, E.; Havey, M.J. Genetic Analyses and Mapping of Pink-Root Resistance in Onion. *J. Am. Soc. Hortic. Sci.* **2018**, *143*, 503–507.
331. Niu, C.; Lister, H.E.; Nguyen, B.; Wheeler, T.A.; Wright, R.J. Resistance to *Thielaviopsis basicola* in the cultivated A genome cotton. *Theor. Appl. Genet.* **2008**, *117*, 1313.
332. Callahan, F.E.; Jenkins, J.N.; Creech, R.G.; Lawrence, G.W. Changes in cotton root proteins correlated with resistance to root knot nematode development. **1997**.
333. Wubben, M.J.; Callahan, F.E.; Velten, J.; Burke, J.J.; Jenkins, J.N. Overexpression of MIC-3 indicates a direct role for the MIC gene family in mediating Upland cotton (*Gossypium hirsutum*) resistance to root-knot nematode (*Meloidogyne incognita*). *Theor. Appl. Genet.* **2015**, *128*, 199–209.
334. Jacquet, C.; Bonhomme, M. Deciphering resistance mechanisms to the root rot disease of legumes caused by *Aphanomyces euteiches* with *Medicago truncatula* genetic and genomic resources. *Model Legum. Medicago truncatula* **2019**, 307–316.
335. Badis, Y.; Bonhomme, M.; Lafitte, C.; Huguet, S.; Balzergue, S.; Dumas, B.; Jacquet, C. Transcriptome analysis highlights preformed defences and signalling pathways controlled by the *pr A e1* quantitative trait locus (QTL), conferring partial resistance to *Aphanomyces euteiches* in *Medicago truncatula*. *Mol. Plant Pathol.* **2015**, *16*, 973–986.
336. Colditz, F.; Nyamsuren, O.; Niehaus, K.; Eubel, H.; Braun, H.-P.; Krajinski, F. Proteomic approach: identification of *Medicago truncatula* proteins induced in roots after infection with the pathogenic oomycete *Aphanomyces euteiches*. *Plant Mol. Biol.* **2004**, *55*, 109–120.

337. Djébali, N.; Jauneau, A.; Ameline-Torregrosa, C.; Chardon, F.; Jaulneau, V.; Mathé, C.; Bottin, A.; Cazaux, M.; Pilet-Nayel, M.-L.; Baranger, A. Partial resistance of *Medicago truncatula* to *Aphanomyces euteiches* is associated with protection of the root stele and is controlled by a major QTL rich in proteasome-related genes. *Mol. Plant-Microbe Interact.* **2009**, *22*, 1043–1055.
338. Kiirika, L.M.; Schmitz, U.; Colditz, F. The alternative *Medicago truncatula* defense proteome of ROS—defective transgenic roots during early microbial infection. *Front. Plant Sci.* **2014**, *5*, 341.
339. Kiirika, L.M.; Bergmann, H.F.; Schikowsky, C.; Wimmer, D.; Korte, J.; Schmitz, U.; Niehaus, K.; Colditz, F. Silencing of the Rac1 GTPase MtROP9 in *Medicago truncatula* stimulates early mycorrhizal and oomycete root colonizations but negatively affects rhizobial infection. *Plant Physiol.* **2012**, *159*, 501–516.
340. Nyamsuren, O.; Colditz, F.; Rosendahl, S.; Tamasloukht, M.; Bekel, T.; Meyer, F.; Kuester, H.; Franken, P.; Krajinski, F. Transcriptional profiling of *Medicago truncatula* roots after infection with *Aphanomyces euteiches* (oomycota) identifies novel genes upregulated during this pathogenic interaction. *Physiol. Mol. Plant Pathol.* **2003**, doi:10.1016/j.pmpp.2003.09.001.
341. Trapphoff, T.; Beutner, C.; Niehaus, K.; Colditz, F. Induction of distinct defense-associated protein patterns in *Aphanomyces euteiches* (Oomycota)-elicited and-inoculated *Medicago truncatula* cell-suspension cultures: A proteome and phosphoproteome approach. *Mol. plant-microbe Interact.* **2009**, *22*, 421–436.
342. Pilet-Nayel, M.-L.; Prospéri, J.-M.; Hamon, C.; Lesne, A.; Lecoïnte, R.; Le Goff, I.; Hervé, M.; Deniot, G.; Delalande, M.; Huguet, T. AER1, a major gene conferring resistance to *Aphanomyces euteiches* in *Medicago truncatula*. *Phytopathology* **2009**, *99*, 203–208.
343. Colditz, F.; Braun, H.-P.; Jacquet, C.; Niehaus, K.; Krajinski, F. Proteomic profiling unravels insights into the molecular background underlying increased *Aphanomyces euteiches*-tolerance of *Medicago truncatula*. *Plant Mol. Biol.* **2005**, *59*, 387–406.
344. Lechner, E.; Achard, P.; Vansiri, A.; Potuschak, T.; Genschik, P. F-box proteins everywhere. *Curr. Opin. Plant Biol.* **2006**, *9*, 631–638.
345. Guo, H.; Ecker, J.R. Plant responses to ethylene gas are mediated by SCFEBF1/EBF2-dependent proteolysis of EIN3 transcription factor. *Cell* **2003**, *115*, 667–677.
346. del Pozo, J.C.; Diaz-Trivino, S.; Cisneros, N.; Gutierrez, C. The balance between cell division and endoreplication depends on E2FC-DPB, transcription factors regulated by the ubiquitin-SCFSKP2A pathway in *Arabidopsis*. *Plant Cell* **2006**, *18*, 2224–2235.
347. Ivanchenko, M.G.; Muday, G.K.; Dubrovsky, J.G. Ethylene–auxin interactions regulate lateral root initiation and emergence in *Arabidopsis thaliana*. *Plant J.* **2008**, *55*, 335–347.
348. Cao, F.Y.; Yoshioka, K.; Desveaux, D. The roles of ABA in plant–pathogen interactions. *J. Plant Res.* **2011**, *124*, 489–499.
349. Adie, B.A.T.; Pérez-Pérez, J.; Pérez-Pérez, M.M.; Godoy, M.; Sánchez-Serrano, J.-J.; Schmelz, E.A.; Solano, R. ABA is an essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defenses in *Arabidopsis*. *Plant Cell* **2007**, *19*, 1665–1681.
350. Colditz, F.; Niehaus, K.; Krajinski, F. Silencing of PR-10-like proteins in *Medicago truncatula* results in an antagonistic induction of other PR proteins and in an increased tolerance upon infection with the oomycete *Aphanomyces euteiches*. *Planta* **2007**, *226*, 57–71.
351. Rey, T.; Nars, A.; Bonhomme, M.; Bottin, A.; Huguet, S.; Balzergue, S.; Jardinaud, M.; Bono, J.; Cullimore, J.; Dumas, B. NFP, a Lys M protein controlling nod factor perception, also intervenes in *Medicago truncatula* resistance to pathogens. *New Phytol.* **2013**, *198*, 875–886.
352. Schenkluhn, L.; Hohnjec, N.; Niehaus, K.; Schmitz, U.; Colditz, F. Differential gel electrophoresis (DIGE) to quantitatively monitor early symbiosis-and pathogenesis-induced changes of the *Medicago truncatula* root proteome. *J. Proteomics* **2010**, *73*, 753–768.
353. Djébali, N.; Mhadhbi, H.; Lafitte, C.; Dumas, B.; Esquerré-Tugayé, M.-T.; Aouani, M.E.; Jacquet, C. Hydrogen peroxide

- scavenging mechanisms are components of *Medicago truncatula* partial resistance to *Aphanomyces euteiches*. *Eur. J. Plant Pathol.* **2011**, *131*, 559.
354. Rey, T.; Laporte, P.; Bonhomme, M.; Jardinaud, M.-F.; Huguet, S.; Balzergue, S.; Dumas, B.; Niebel, A.; Jacquet, C. MtNF-YA1, a central transcriptional regulator of symbiotic nodule development, is also a determinant of *Medicago truncatula* susceptibility toward a root pathogen. *Front. Plant Sci.* **2016**, *7*, 1837.
355. Kjølner, R.; Rosendahl, S. The presence of the arbuscular mycorrhizal fungus *Glomus intraradices* influences enzymatic activities of the root pathogen *Aphanomyces euteiches* in pea roots. *Mycorrhiza* **1997**, *6*, 487–491.
356. Bødker, L.; Kjølner, R.; Rosendahl, S. Effect of phosphate and the arbuscular mycorrhizal fungus *Glomus intraradices* on disease severity of root rot of peas (*Pisum sativum*) caused by *Aphanomyces euteiches*. *Mycorrhiza* **1998**, *8*, 169–174.
357. Wang, H. Identification and dissection of soybean QTL conferring resistance to *Phytophthora sojae* 2011.
358. Li, X.; Han, Y.; Teng, W.; Zhang, S.; Yu, K.; Poysa, V.; Anderson, T.; Ding, J.; Li, W. Pyramided QTL underlying tolerance to *Phytophthora* root rot in mega-environments from soybean cultivars 'Conrad' and 'Hefeng 25.' *Theor. Appl. Genet.* **2010**, *121*, 651–658.
359. Njiti, V.N.; Meksem, K.; Iqbal, M.J.; Johnson, J.E.; Kassem, M.A.; Zobrist, K.F.; Kilo, V.Y.; Lightfoot, D.A. Common loci underlie field resistance to soybean sudden death syndrome in Forrest, Pyramid, Essex, and Douglas. *Theor. Appl. Genet.* **2002**, *104*, 294–300.
360. Hashmi, R.Y. *Inheritance of resistance to soybean sudden death syndrome (SDS) in Ripley x Spencer F (5) derived lines*; Southern Illinois University at Carbondale, 2004; ISBN 0496969455.
361. Wang, H.; Waller, L.; Tripathy, S.; St. Martin, S.K.; Zhou, L.; Krampis, K.; Tucker, D.M.; Mao, Y.; Hoeschele, I.; Saghai Maroof, M.A. Analysis of genes underlying soybean quantitative trait loci conferring partial resistance to *Phytophthora sojae*. *Plant Genome* **2010**, *3*, 23–40.
362. Kazi, S.; Shultz, J.; Afzal, J.; Johnson, J.; Njiti, V.N.; Lightfoot, D.A. Separate loci underlie resistance to root infection and leaf scorch during soybean sudden death syndrome. *Theor. Appl. Genet.* **2008**, *116*, 967–977.
363. Klepadlo, M.; Balk, C.S.; Vuong, T.D.; Dorrance, A.E.; Nguyen, H.T. Molecular characterization of genomic regions for resistance to *Pythium ultimum* var. *ultimum* in the soybean cultivar Magellan. *Theor. Appl. Genet.* **2019**, *132*, 405–417.
364. Campa, A.; Pérez-Vega, E.; Pascual, A.; Ferreira, J.J. Genetic analysis and molecular mapping of quantitative trait loci in common bean against *Pythium ultimum*. *Phytopathology* **2010**, *100*, 1315–1320.
365. Shin, S.; Zheng, P.; Fazio, G.; Mazzola, M.; Main, D.; Zhu, Y. Transcriptome changes specifically associated with apple (*Malus domestica*) root defense response during *Pythium ultimum* infection. *Physiol. Mol. Plant Pathol.* **2016**, *94*, 16–26.
366. Zhu, Y.; Shao, J.; Zhou, Z.; Davis, R.E. Genotype-specific suppression of multiple defense pathways in apple root during infection by *Pythium ultimum*. *Hortic. Res.* **2019**, *6*, 1–17.
367. Nath, V.S.; Koyyappurath, S.; Alex, T.E.; Geetha, K.A.; Augustine, L.; Nasser, A.; Thomas, G. Transcriptome-based mining and expression profiling of *Pythium* responsive transcription factors in *Zingiber* sp. *Funct. Integr. Genomics* **2019**, *19*, 249–264.
368. Zhu, Y.; Zhao, J.; Zhou, Z. Identifying an elite panel of apple rootstock germplasm with contrasting root resistance to *Pythium ultimum*. *J. Plant Pathol.* **2018**, *9*, 1000461.
369. Wrather, J.A.; Anderson, T.R.; Arsyad, D.M.; Tan, Y.; Ploper, L.D.; Porta-Puglia, A.; Ram, H.H.; Yorinori, J.T. Soybean disease loss estimates for the top ten soybean-producing countries in 1998. *Can. J. Plant Pathol.* **2001**, *23*, 115–121.
370. Han, Y.; Teng, W.; Yu, K.; Poysa, V.; Anderson, T.; Qiu, L.; Lightfoot, D.A.; Li, W. Mapping QTL tolerance to *Phytophthora* root rot in soybean using microsatellite and RAPD/SCAR derived markers. *Euphytica* **2008**, *162*, 231–239.
371. Dorrance, A.E.; McClure, S.A.; St. Martin, S.K. Effect of partial resistance on *Phytophthora* stem rot incidence and yield of soybean in Ohio. *Plant Dis.* **2003**, *87*, 308–312.
372. Weng, C.; Yu, K.; Anderson, T.R.; Poysa, V. Mapping genes conferring resistance to *Phytophthora* root rot of soybean, *Rps1a* and *Rps7*. *J. Hered.* **2001**, *92*, 442–446.

373. Sandhu, D.; Schallock, K.G.; Rivera-Velez, N.; Lundeen, P.; Cianzio, S.; Bhattacharyya, M.K. Soybean Phytophthora resistance gene Rps8 maps closely to the Rps3 region. *J. Hered.* **2005**, *96*, 536–541.
374. Dorrance, A.E. Management of Phytophthora sojae of soybean: a review and future perspectives. *Can. J. Plant Pathol.* **2018**, *40*, 210–219.
375. Ranathunge, K.; Thomas, R.H.; Fang, X.; Peterson, C.A.; Gijzen, M.; Bernards, M.A. Soybean root suberin and partial resistance to root rot caused by Phytophthora sojae. *Phytopathology* **2008**, *98*, 1179–1189.
376. Drenth, A.; Whisson, S.C.; Maclean, D.J.; Irwin, J.A.G.; Obst, N.R.; Ryley, M.J. The evolution of races of Phytophthora sojae in Australia. *Phytopathology* **1996**, *86*, 163–169.
377. Leitz, R.A.; Hartman, G.L.; Pedersen, W.L.; Nickell, C.D. Races of Phytophthora sojae on soybean in Illinois. *Plant Dis.* **2000**, *84*, 487.
378. Malvick, D.K.; Grunden, E. Characteristics of Phytophthora sojae populations in Illinois and implications for management of Phytophthora rot of soybean. In Proceedings of the PHYTOPATHOLOGY; AMER PHYTOPATHOLOGICAL SOC 3340 PILOT KNOB ROAD, ST PAUL, MN 55121 USA, 2004; Vol. 94, pp. S65–S65.
379. Sugimoto, T.; Kato, M.; Yoshida, S.; Matsumoto, I.; Kobayashi, T.; Kaga, A.; Hajika, M.; Yamamoto, R.; Watanabe, K.; Aino, M. Pathogenic diversity of Phytophthora sojae and breeding strategies to develop Phytophthora-resistant soybeans. *Breed. Sci.* **2012**, *61*, 511–522.
380. Gao, H.; Narayanan, N.N.; Ellison, L.; Bhattacharyya, M.K. Two classes of highly similar coiled coil-nucleotide binding-leucine rich repeat genes isolated from the Rps1-k locus encode Phytophthora resistance in soybean. *Mol. plant-microbe Interact.* **2005**, *18*, 1035–1045.
381. Gao, H.; Bhattacharyya, M.K. The soybean-Phytophthora resistance locus Rps1-k encompasses coiled coil-nucleotide binding-leucine rich repeat-like genes and repetitive sequences. *BMC Plant Biol.* **2008**, *8*, 29.
382. Graham, M.A.; Marek, L.F.; Shoemaker, R.C. Organization, expression and evolution of a disease resistance gene cluster in soybean. *Genetics* **2002**, *162*, 1961–1977.
383. Sun, J.; Li, L.; Zhao, J.; Huang, J.; Yan, Q.; Xing, H.; Guo, N. Genetic analysis and fine mapping of RpsJS, a novel resistance gene to Phytophthora sojae in soybean [Glycine max (L.) Merr.]. *Theor. Appl. Genet.* **2014**, *127*, 913–919.
384. Zhang, J.; Xia, C.; Wang, X.; Duan, C.; Sun, S.; Wu, X.; Zhu, Z. Genetic characterization and fine mapping of the novel Phytophthora resistance gene in a Chinese soybean cultivar. *Theor. Appl. Genet.* **2013**, *126*, 1555–1561.
385. Zhang, J.; Xia, C.; Duan, C.; Sun, S.; Wang, X.; Wu, X.; Zhu, Z. Identification and candidate gene analysis of a novel Phytophthora resistance gene Rps10 in a Chinese soybean cultivar. *PLoS One* **2013**, *8*, e69799.
386. Thomas, R.; Fang, X.; Ranathunge, K.; Anderson, T.R.; Peterson, C.A.; Bernards, M.A. Soybean root suberin: anatomical distribution, chemical composition, and relationship to partial resistance to Phytophthora sojae. *Plant Physiol.* **2007**, *144*, 299–311.
387. Fang, X. Chemical composition of soybean root epidermal cell walls 2006.
388. Valenzuela-Estrada, L.R.; Bryla, D.R.; Hoashi-Erhardt, W.K.; Moore, P.P.; Forge, T.A. Root traits associated with Phytophthora root rot resistance in red raspberry. In Proceedings of the X International Rubus and Ribes Symposium 946; 2011; pp. 283–287.
389. Graham, J.; Hackett, C.A.; Smith, K.; Woodhead, M.; MacKenzie, K.; Tierney, I.; Cooke, D.; Bayer, M.; Jennings, N. Towards an understanding of the nature of resistance to Phytophthora root rot in red raspberry. *Theor. Appl. Genet.* **2011**, *123*, 585–601.
390. Engelbrecht, J.; Van den Berg, N. Expression of defence-related genes against Phytophthora cinnamomi in five avocado rootstocks. *S. Afr. J. Sci.* **2013**, *109*, 1–8.
391. Van de Weg, W.E.; Henken, B.; Giezen, S. Assessment of the resistance to Phytophthora fragariae var. fragariae of the USA and Canadian differential series of strawberry genotypes. *J. Phytopathol.* **1997**, *145*, 1–6.

1.