

Article

Multidimensional statistical technique for interpreting the spontaneous breakthrough cancer pain phenomenon. A secondary analysis from the IOPS-MS study

Marco Cascella ¹, Anna Crispo ^{2,*}, Gennaro Esposito ¹, Cira Antonietta Forte ¹, Sergio Coluccia ², Giuseppe Porciello ², Alfonso Amore ³, Sabrina Bimonte ¹, Sebastiano Mercadante ⁴, Augusto Caraceni ⁵, Massimo Mammucari ⁶, Paolo Marchetti ⁷, Rocco Domenico Mediati ⁸, Silvia Natoli ⁹, Giuseppe Tonini ¹⁰ and Arturo Cuomo ¹

¹ Division of Anesthesia and Pain Medicine, Istituto Nazionale Tumori - IRCCS - "Fondazione G.Pascale", Naples, Italy

² Epidemiology and Biostatistics Unit, Istituto Nazionale Tumori, IRCCS Fondazione G. Pascale, Naples, Italy

³ Melanoma and Sarcoma Surgery Unit, Istituto Nazionale Tumori, IRCCS Fondazione G. Pascale, Naples, Italy

⁴ Anesthesia and Intensive Care & Pain Relief and Supportive Care, 00185 La Maddalena, Italy

⁵ Palliative Care, Pain Therapy and Rehabilitation, National Cancer Institute, IRCCS Foundation, 20133 Milan, Italy; augusto.caraceni@istitutotumori.mi.it

⁶ Primary Care Unit, ASL RM1, 00193 Rome, Italy; massimo.mammucari@libero.it

⁷ Department of Clinical and Molecular Medicine, La Sapienza University of Rome, 00185 Rome, Italy; paolo.marchetti@uniroma1.it

⁸ Palliative Care and Pain Therapy Unit, Careggi Hospital, 50139 Florence, Italy; mediatir@aou-careggi.toscana.it

⁹ Department of Clinical Science and Translational Medicine—University of Rome Tor Vergata, 00133 Rome, Italy; silvia.natoli@uniroma2.it

¹⁰ Medical Oncology Department, Campus Bio-Medico University of Rome, 00128 Rome, Italy; g.tonini@unicampus.it

* Correspondence: a.crispo@istitutotumori.na.it; Tel.: +39-081-5903597

Simple Summary: Pain is one of the most common and debilitating symptoms in cancer patients. A clinical peculiarity of cancer pain is the breakthrough cancer pain (BTcP) which is defined as a temporary exacerbation of pain that "breaks through" a phase of adequate pain control by an opioid-based therapy. The NP-BTcP occurs in the absence of any specific activity. In this paper, we addressed the topic through a mathematical approach to provide many indications for identifying the diagnostic and therapeutic gaps on NP-BTcP management.

Abstract: Breakthrough cancer pain (BTcP) is a temporary exacerbation of pain that "breaks through" a phase of adequate pain control by an opioid-based therapy. The non-predictable BTcP (NP-BTcP) is a subtype of BTcP that occurs in the absence of any specific activity. Since NP-BTcP has an important clinical impact, this analysis is aimed at characterizing the NP-BTcP phenomenon through a multidimensional statistical technique. This is a secondary analysis based on the Italian Oncologic Pain multiSetting - Multicentric Survey (IOPS-MS). A correlation analysis was performed to characterize NP-BTcP profile about its intensity, number of episodes per day, and type. The Multidimensional Correspondence Analysis (MCA) determined the identification of 4 groups (Phenotypes). A univariate analysis was performed to assess differences between the 4 Phenotypes and selected covariates. The four phenotypes represent the hierarchical classification according to the status of NP-BTcP: from the best (Phenotype 1) to the worst (Phenotype 4). The univariate analysis found a significant association between the onset time >10 min in the Phenotype 1 (37.3%) vs. the onset > 10 min in Phenotype 4 (25.8%) ($p < 0.001$). The Phenotype 1 was characterized by gastrointestinal type of cancer (26.4%) respect to Phenotype 4 where the most frequent cancer affected the lung (28.8%) ($p < 0.001$). Phenotype 4 was mainly managed with rapid onset opioids, while in Phenotype 1 many patients were treated with oral, subcutaneous, or intravenous morphine (56.4% and 44.4%, respectively; $p = 0.008$). The ability to characterize NP-BTcP can offer enormous benefits for the management of this serious aspect of cancer pain. This strategy can provide many indications for identifying the diagnostic and therapeutic gaps on NP-BTcP management.

Keywords: cancer pain; breakthrough cancer pain; cluster analysis

1. Introduction

Despite the many therapeutic possibilities, pain remains one of the most common and debilitating symptoms in cancer patients. Its prevalence is approximately 55% during anticancer treatment, 40% after therapy, and about 66% in the advanced and progressive phase of disease stages [1]. Although the numerical data are useful for obtaining a quantitative estimate of the problem, the precise characterization of this complex symptom is mandatory. This involves defining the various clinical presentations.

A clinical peculiarity of cancer pain is the breakthrough cancer pain (BTcP). It is a temporary exacerbation of pain that "breaks through" a phase of adequate pain control by an opioid-based therapy [2]. Notably, BTcP can affect up to 70% of cancer patients [3]. Clinically, an episode of BTcP is of short duration (15-30 min), moderate-to-severe intensity, and short-latency between onset and peak of intensity (a few minutes) [4]. Its clinical presentations, however, may vary depending on multiple factors, such as the clinical setting, and type and degree of the background pain [5].

BTcP is divided into two categories including the predictable and the non-predictable BTcP (NP-BTcP) type. The predictable, or incident type, is further categorized into three subclasses including the volitional BTcP (caused by a voluntary act), the non-volitional subtype (caused by an involuntary act), and the procedural pain. The NP-BTcP occurs in the absence of any specific activity. It is also indicated as spontaneous or idiopathic BTcP [2,4].

The clinical and therapeutic implications, and probably the pathophysiology, are very different. For example, since the predictable BTcP is usually triggered by voluntary (e.g., dressing changes), not involuntary movements (e.g., bowel spasm), or therapeutic procedures, adequate knowledge of the phenomenon can facilitate the prevention and implementation of effective therapeutic strategies. On the other hand, since it is not possible to establish its triggering events, the clinical and therapeutic implications of NP-BTcP are of fundamental importance. The idiopathic form has many aspects yet to be defined. However, since it accounts for about 70% of the total cases of BTcP [5], and is in nature unpredictably, with little or no warning, it should be carefully addressed.

The study of the phenomenon can be performed through clinical investigations. Another strategy is the analysis of high sample size datasets, using ad hoc statistical methods. On these premises, this work aims to characterize the NP-BTcP phenomenon through a multidimensional method applied to the most copious database collected so far.

2. Materials and Methods

2.1. The IOPS-MS dataset

This is a secondary analysis based on the Italian Oncologic Pain multiSetting - Multicentric Survey (IOPS-MS) that was carried out for dissecting the BTcP phenomenon [3,5]. Details concerning enrollment of patients and all recorded variables are described elsewhere [3,5]. The local Ethical Committee approved the protocol, and each patient signed an informed consent.

Inclusion criteria were

- age greater than 18 years;
- diagnosis of cancer at any stage; well-controlled and stable background pain with an intensity ≤ 4 (on a 0-10 numerical rating scale, NRS);
- presence of BTcP episode of moderate-severe intensity, clearly distinguished from background pain.

Exclusion criteria were:

- no cancer diagnosis;
- unstable and/or uncontrolled background pain (>4/10);
- no relevant peaks in pain intensity (<5/10);
- poor collaboration or refusal to participate

Among all recorded variables from the original study, some variables were selected for this secondary analysis: age, gender, setting, type of primary tumor, onset, type of BTcP pain, Karnofsky status, and type of physician were recorded. Patients were asked about the average time of meaningful pain relief after their BTcP medication (≤ 10 min, >10 min). The study was observational, and pharmaceutical therapies were used according to local policy, without following strict protocols.

In the original IOPS-MS dataset, $n=4016$ cases were entered. Of these, 30.5% ($n=1225$) were incident BTcP and 69.5% ($n=2791$) NP-BTcP [5]. The statistical analysis of this study concerned the latter subgroup.

2.2. Statistical analysis

Descriptive statistics was provided for BTcP characteristics and main variables. At first, to characterize BTcP profile a Pearson correlation was performed among the following continuous variables:

- a) BTcP intensity score (NRS 5-10)
- b) number of BTcP episodes per day (1-4)
- c) duration of BTcP.

A multidimensional analysis was performed using the continuous variables with the highest correlation (a and b variables; $p < 0.001$), and the categorical variable "BTcP type" (nociceptive, neuropathic, mixed: presence or absence). Consequently, the Multidimensional Correspondence Analysis (MCA) determined the identification of groups (termed as 'Phenotypes'). The CA-plot showed the theoretical clusters and a further confirmative hierarchical analysis (HCPC) [6] was implemented to compare the theoretical findings through an automatic classification criterion [7].

An automatic procedure that maximize the inertia between centroids-barycenter (Ward method) was implemented to classify the groups.

Finally, an univariate analysis was performed to observe differences between the 4 Phenotypes; a Chi-Square test was used to measure categorical variables, and a p -value less than 0.05 was considered statistically significant (Figure 1). The statistical analysis was carried out using R software by using the FactoMineR package [8] and SPSS version 26.




Figure 1. Flow chart. Abbreviations. IOPS=Italian Oncological Pain Survey. BTcP = breakthrough cancer pain. NRS=numerical rating scale.

3. Results

This section may be divided by subheadings. It should provide a concise and precise description of the experimental results, their interpretation, as well as the experimental conclusions that can be drawn. Among patients with NP-BTcP (n=2791) from the original study, one hundred and twenty presented more than 4 episodes per day. Those patients were excluded from this secondary analysis, thus 95.7% of the total set was analyzed (n=2671).

Patients' characteristics are described in Table 1. Fifty five percent of patients were male, over 52% were above 65 years of age; the most frequent setting was the Day-Hospital setting (47%), and the most frequent primary tumor was the lung carcinoma (25%). The number, intensity and type of BTcP were described; about 68% of the onset was less than 10 minutes, and regarding the therapy 49% of patients received rapid onset opioids (ROOs) for BTcP management.

Table 1. Descriptive Statistics.

	N(total = 2671)	%
Gender		
Male	1465	54.8
Female	1206	45.2
Age		
<55	589	22.1
55-64	696	26.1
65-74	810	30.3

≥75	576	21.6
Setting		
Inpatients	1015	36.0
Outpatients / DH	1250	46.8
Hospice / Home palliative care	406	15.2
Primary Tumor		
Lung	665	24.9
Gastrointestinal	470	17.6
Breast	278	10.4
Pancreas	237	8.9
Uro-Gynecological	401	15.0
Other	620	23.2
N° of BTcP		
1	1082	40.5
2	621	23.2
3	177	6.6
4		
Intensity of BTcP		
5	118	4.4
6	394	14.8
7	840	31.4
8	849	31.8
9	277	10.4
10	193	7.2
Type of BtcP Pain		
Nociceptive	672	25.2
Neuropatic	214	8.0
Mixed pain	1785	66.8
Onset		
≤10 min	1806	67.6
>10 min	865	32.4
BTcP therapy		
None	366	13.8
ROOs^	1302	48.7

Morphine ^o	545	20.4
Other	458	17.1

Abbreviations: RT = Radiotherapy; DH=Day Hospital; ROOs = (rapid onset opioids. Legend: ^ROOs include: OTFC = Oral transmucosal fentanyl citrate; FBT=Fentanyl buccal tablet; FBST=Sublingual fentanyl; FPNS = Fentanyl Pectin Nasal Spray. ^oMorphine given by the oral, subcutaneous, intramuscular or intravenous route.

MCA was shown in Figure 2. The 4-clusters (Phenotypes) represent a hierarchical classification according to the BTcP status which from P1 (Phenotype 1) to P4 (Phenotypes 4) gradually get worse.




Figure 2. Multidimensional Correspondence Analysis (MCA). Identification of 4 groups (Phenotypes, P) of breakthrough cancer pain (BTcP) from P1 to P4 by combining the continuous variables BTcP intensity score (NRS 5-10), and number of BTcP episodes per day (1-4) with the categorical variable BTcP type (nociceptive, neuropathic, mixed: presence or absence).

An automatic procedure recognizes the 6-theoretical clusters groups (within groups sum of squares 6, $WSS_6 = 509.13$). The K-Means method, represented by the Elbow plot (Figure 3) suggests that a good representation of clusters is given from 4 to 10 groups. So a parsimonious point of view has been adopted and 4-clusters final groups (Phenotypes) have been selected ($WSS_4=838.93$). According to cluster estimation the agreement between induced classification and automatic classification was 31.6%.

Elbow Plot: Optimal number of clusters




Figure 3. Elbow Plot. An automatic procedure recognizes the 6-theoretical clusters groups (within groups sum of squares, $WSS_6 = 509.13$). The K-Means method, suggests that a good representation

of clusters is given from 4 to 10 groups. By using a parsimonious point of view, 4-clusters (Phenotypes) have been selected (WSS4 = 838.93). According to cluster estimation the agreement between induced classification and automatic classification was 31.6%.

The features of the 4 clusters/phenotypes are synthesized in Figure 4.

PHENOTYPES	EPISODES/ DAY				BTcP Intensity (0-10 NRS)								
	1	2	3	4	5-6		7		8-9		10		
					Type of PAIN		Type of PAIN		Type of PAIN		Type of PAIN		
					Noc	NoB	Noc	NoB	Noc	NoB	Noc	NoB	
P1	X				X	X	X						
		X			X								
			X		X								
				X	X								
P2	X						X	X					
		X					X	X	X				
			X		X	X							
				X	X								
P3	X								X	X			
		X							X	X			
			X				X	X	X				
				X			X	X					
P4	X											X	
		X										X	
			X									X	X
				X					X	X	X		

Figure 4. Phenotypes of non-predictable breakthrough cancer pain. Four phenotypes (from P1 to P4) are described according to the pain intensity, number of episodes a day, and type of pain. Abbreviations: NRS = numerical rating scale; Noc = Nociceptive; NoB = Neuropathic or Both.

The univariate analysis shows the associations between the 4 phenotypes and selected variables (Table 2); a borderline significance was observed for age: young patients (<55 years) were more favorable to be in P4 (24.2%) compared with the elderly (≥ 75 years) who were allocated in P1 (25.9%) ($p=0.055$). A significant association was assessed between the onset time >10 min in the Phenotype 1 (37.3%) vs. the onset > 10 min in Phenotype 4 (25.8%) ($p<0.001$); inpatients were mainly allocated into the cluster P4, respect to those in palliative care settings who expressed a phenotype 1 (43.6% vs. 26.6%, respectively $p<0.001$). About cancer type, P1 was characterized by gastrointestinal (26.4%) respect to P4 where the most frequent cancer affected was the lung (28.8%) ($p<0.001$).

Phenotype 4 was mainly managed with rapid onset opioids (ROOs), while in phenotype 1 many patients were treated with oral, subcutaneous, or intravenous morphine (56.4% and 44.4%, respectively; $p=0.008$). Moreover number of patients who did not receive therapy (ROOs, morphine and other) decreased from P1 to P4 (16.5% and 8.1%, respectively $p=0.003$) (Figure 5).

Table 2. Univariate analysis.

	PHENOTYPES				<i>p-value*</i>
	1 (N=394)	2 (N=962)	3 (N=1079)	4 (N=236)	
Age					0.055
<55	68 (17.3%)	220 (22.9%)	244 (22.6%)	57 (24.2%)	
55-64	96 (24.4%)	229 (23.8%)	305 (28.3%)	66 (26.1%)	
65-74	128 (32.5%)	301 (31.3%)	313 (29.0%)	68 (28.8%)	
≥75	102 (25.9%)	212 (22.0%)	217 (20.1%)	45 (19.1%)	
Gender					0.5
Female	177 (44.9%)	452 (47%)	476 (44.1%)	101 (42.8%)	
Male	217 (55.1%)	510 (53%)	603 (55.9%)	135 (57.2%)	
Onset					<0.001
≤10 min	247 (62.7%)	552 (57.4%)	832 (77.1%)	175 (74.2%)	
>10 min	147 (37.3%)	410 (42.6%)	247 (22.9%)	61 (25.8%)	
Setting					<0.001
Inpatients	167 (42.2%)	308 (32.0%)	437 (40.5%)	103 (43.6%)	
Outpatients / DH	122 (31.0%)	487 (50.6%)	539 (50.0%)	102 (43.2%)	
Hospice / Home palliative care	105 (26.6%)	167 (17.4%)	103 (9.5%)	31 (13.1%)	
Cancer type					<0.001
Lung	67 (17.0%)	232 (24.1%)	298 (27.6%)	68 (28.8%)	
Gastric	104 (26.4%)	173 (18.0%)	161 (14.9%)	31 (13.1)	
Breast	41 (10.4%)	110 (11.4%)	104 (9.6%)	23 (9.7%)	
Pancreatic	38 (9.6%)	80 (8.3%)	101 (9.4%)	18 (7.6%)	
Uro-Gynecological	42 (10.7%)	146 (15.2%)	172 (15.9%)	41 (17.4%)	
All other cancers	102 (25.9%)	407 (42.5%)	459 (42.5%)	101 (42.8%)	
Therapy					0.008
None**	65 (16.5%)	150 (15.6%)	132 (12.2%)	19 (8.1%)	
ROOs	175 (44.4%)	441 (45.8%)	553 (51.3%)	133 (56.4%)	
Morphine ^o	91 (23.1%)	182 (18.9%)	211 (19.6%)	43 (18.2%)	
Other therapies	63 (16%)	200 (20.8%)	183 (17%)	41 (17.4%)	

Abbreviations: RT=Radiotherapy; DH=Day Hospital; ROOs=Rapid Onset Opioids Legend: Orally Morphine Sulfate, Morphine Chloridrate subcutaneous or intravenous. ** Chi-square test between "None" vs. "ROOs, Morphine and other therapies" statistically significant.




Figure 5. Distribution of therapies according to Phenotypes. The NP-BTcP management differs between Phenotypes. Orange-Trend indicates the significant increase of ROOs from P1 to P4 (‡, $p=0.008$); Blu-Trend shows the significant decrease of none therapy from P1 to P4 (**, $p=0.003$). Abbreviations: NP-BTcP= non-predictable breakthrough cancer pain; ROOs=Rapid Onset Opioids.

4. Discussion

To our knowledge, this is the first study aimed at characterizing the NP-BTcP phenomenon. Several pieces of evidence showed that BTcP cannot be intended as a single nosological entity, but includes different and changing pathogenetic mechanisms justifying its sub-typing [2,3]. Consequently, methods useful for dissecting the NP-BTcP forms can offer great help for clinicians, and important benefits for patients.

The main challenge for BTcP management is the implementation of new procedures that can be useful for both diagnostic and therapeutic approaches. Recently, Pantano et al. [9] performed a machine learning approach to identify possible subgroups of both types of BTcP. Their algorithm was based on BTcP therapy satisfaction, clinical features, basal pain, and rapid-onset opioids. This approach allowed the identification of 12 distinct BTcP clusters. In our analysis, through the automatic classification method, 4 clusters of NP-BTcP were identified. The clusters were obtained by combining two continuous variables (BTcP intensity and number of BTcP episodes per day) with the categorical variable BTcP type (nociceptive, neuropathic, mixed) (Figure 2).

Subsequently, the characteristics assumed by the individuals in each group were projected. In turn, a rule of classification was developed (Figure 4). This approach could facilitate rapid identification of the patient who is potentially at higher risk of developing more severe forms of NP-BTcP, in terms of pain intensity, type, and number of episodes per day. Notably, although the method initially suggested a classification of 6 groups, the choice was given to the 4 groups, given the best goodness of estimate ($WSS_4 = 838.93$). For all these reasons, the goodness of fit was 32% which represents the explained variation of the entire phenomenon. Furthermore, it must be considered that the model was based on the 3 variables and a large size of the sample. According to the hierarchical order assumed by the Phenotypes, it was easier to recognize the best group (P1) versus the worst one (P4).

The data emerging from the categorization indicate that P1 expresses nociceptive pain. Although even 4 episodes per day can occur, the NRS (0-10) is of intensity between 5-6. On the other hand, P4 is characterized by episodes of neuropathic or mixed pain of

severe intensity. The intermediate clusters (P3 and P4), may have a nociceptive type of NP-BTcP, but of high intensity, or they express neuropathic/mixed pain of submaximal intensity.

Subsequently, the univariate analysis showed that phenotype 1 is characterized by older age, slow onset, gastrointestinal as primary tumor, and greater propensity to be treated in the context of the palliative care setting. On the contrary, the main features of phenotype 4 are younger age, rapid onset and furthermore it most frequently concerns inpatients affected by lung cancer. Moreover, regarding the NP-BTcP therapy, the worst phenotype (P4) was mainly managed with rapid onset opioids (ROOs); on the contrary, in phenotype 1 many patients were treated with oral, subcutaneous, or intravenous morphine. Interestingly, guidelines on cancer pain treatment do not refer to the severity of BTcP and provide different guidance regarding its treatment. For instance, the European Society for Medical Oncology (ESMO) guidelines recommended the use of fentanyl formulations for the treatment of BTcP (level of evidence I, degree of recommendation A). In the same guidelines, morphine received a level of evidence of II and a degree of recommendation of B [10]. On the other side, despite the WHO guidelines stated that BTcP should always be relieved with rescue medicine based on clinical experience and patient need, immediate-release or slow-release morphine was recommended [11]. Nevertheless, generic guidelines continue to suggest the use of oral opioids whereas specific BTcP guidelines recommend the use of ROOs as rescue medication. According to Davis et al. [12], the different attitudes towards BTcP management do not reflect research evidence but personal opinions.

A great concern is the lack of treatment of NP-BTcP in each phenotype. Notably, about 17% of patients in P1 did not receive any treatment and, even if this percentage decreases up to P4, there is a statistically significant difference compared with those who received ROOs, morphine, and all other therapies. Despite the guidelines on the topic, proper cancer pain treatment remains a paramount unmet medical need for cancer patients [13].

The ability to characterize NP-BTcP can offer enormous benefits for the management of this serious aspect of cancer pain. This strategy can provide many indications for identifying the diagnostic and therapeutic gaps in NP-BTcP management.

Particular care perspectives are very interesting. The possibility of developing a mobile application to quickly allocate the patient in one of the 4 clusters is one of these perspectives. This method could implement systems already in use or being tested. The Pain ASsessment in CAncer Patients by Machine LEarning (PASCALe) study (NCT04726228) is ongoing; it is aimed at evaluating the possibility that telemedicine can improve the quality of life of cancer patients through self-management and remote monitoring solutions. The study involves the development of an application that also records information on the BTcP features. After the first phase of implementation of the database and machine learning for identifying objective characters of pain, the authors plan to include the 4 clusters to study multiple features of NP-BTcP (clinical data, therapy, impact on quality of life, and others).

In addition to experimental approaches, telemedicine tools could be used to assist those patients who are immediately allocated to the worst clusters (P3 and P4). For these patients, the management of NP-BTcP must be very aggressive to counteract its impact on the quality of life.

If adequate resources are available, even the so-called best phenotypes (P1 and P2) can benefit from telemedicine. The experience accumulated in the management of patients with cancer pain during the pandemic suggests that many episodes of NP-BTcP are not reported during periodic consultations, but are recorded by the patient in the monitoring system. In other words, the impact on quality of life is always present, but both patients and clinicians tend to underestimate the effect.

This study has multiple limitations. The most important limitation concerns the assumption that it is a secondary analysis based on a survey. However, our starting point

remains the most abundant dataset on the subject. Moreover, the original IOPS study was not drawn to perform this analysis. This gap could be overcome by the development of prospective studies to verify the reliability (validation) of the proposed classification system (Table 2). Another limitation regards the mathematical approach as the method works only with complete data. Nevertheless, we adopted a dataset with no or few missing values.

5. Conclusion

The idiopathic form of BTcP has many aspects yet to be defined and the characterization of the phenomenon means being able to improve its management. In-depth knowledge is also mandatory because NP-BTcP accounts for over two-thirds of all BTcP cases. While waiting to obtain data from prospective studies, dataset analyses carried out according to ad hoc statistical approaches, can allow us to define clusters (phenotypes) that differ in type of pain, intensity, and number of daily episodes. It can be deduced, for instance, that those suffering from lung cancer may be classified in the worst phenotype of NP-BTcP (P4) and this prerequisite presupposes greater diagnostic attention. Finally, a more precise definition of the clusters could lead to important therapeutic perspectives. For example, a mobile application could be useful for the clinician to frame individual patients. To be more precise, patients who fall into the worst phenotypes could be followed up under telemedicine programs. The aim is to provide tailored management of the NP-BTcP.

Author Contributions: Conceptualization, A.C., M.C., Arturo C.; methodology, M.C., A.C., software, S.C., formal analysis, S.C., A.C., M.C., investigation M.C., G.E., C.A.F., A.A., S.B., S.M., Augusto C., M.M., P.M., R.D.M., S.N., G.T., Arturo C.; data curation, A.C. C.A.F., S.C., G.P.; writing—original draft preparation, M.C., A.C., S.M., Augusto C., M.M., P.M., R.D.M., S.N., G.T., Arturo C.; writing—review and editing, M.C., A.C.; supervision, M.C., S.B., Arturo C.; project administration, M.C., A.C. S.M., Arturo C. All authors have read and agreed to the published version of the manuscript.

Funding: The original study was sponsored by Molteni Farmaceutici, Italy. Data were independently analyzed and managed by authors. The funders had no role in the design of this secondary analyses or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (PROTOCOLLO DI STUDIO IOPS-MS) Policlinico Tor Vergata (REGISTRO SPERIMENTAZIONI n.21/13) Roma, 20 febbraio 2013.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study. Written informed consent has been obtained from the patient(s) to publish this paper”.

Data Availability Statement: The datasets generated during/and or analyzed during the current study are available from the corresponding author on request.

Acknowledgments: The authors thank Mrs Maria Cristina Romano for data curation and the IOPS-MS group for their work in the original study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. van den Beuken-van Everdingen, M.H.; Hochstenbach, L.M.; Joosten, E.A.; Tjan-Heijnen, V.C.; Janssen, D.J. Update on Prevalence of Pain in Patients With Cancer: Systematic Review and Meta-Analysis. *J Pain Symptom Manage* **2016**, *51*, 1070-1090 e1079, doi:10.1016/j.jpainsymman.2015.12.340.
2. Mercadante, S.; Portenoy, R.K. Breakthrough cancer pain: twenty-five years of study. *Pain* **2016**, *157*, 2657-2663, doi:10.1097/j.pain.0000000000000721.

3. Mercadante, S.; Lazzari, M.; Reale, C.; Cuomo, A.; Fusco, F.; Marchetti, P.; Mediati, R.D.; Chiurazzi, B.; Ciuffedra, L.; Caraceni, A.; et al. Italian Oncological Pain Survey (IOPS): a multicentre Italian study of breakthrough pain performed in different settings. *Clin J Pain* **2015**, *31*, 214-221, doi:10.1097/AJP.0000000000000161.
4. Davies, A.N.; Dickman, A.; Reid, C.; Stevens, A.M.; Zeppetella, G.; Science Committee of the Association for Palliative Medicine of Great, B.; Ireland. The management of cancer-related breakthrough pain: recommendations of a task group of the Science Committee of the Association for Palliative Medicine of Great Britain and Ireland. *Eur J Pain* **2009**, *13*, 331-338, doi:10.1016/j.ejpain.2008.06.014.
5. Mercadante, S.; Marchetti, P.; Cuomo, A.; Caraceni, A.; Mediati, R.D.; Vellucci, R.; Mammucari, M.; Natoli, S.; Lazzari, M.; Dauri, M.; et al. Factors Influencing the Clinical Presentation of Breakthrough Pain in Cancer Patients. *Cancers (Basel)* **2018**, *10*, doi:10.3390/cancers10060175.
6. Husson, F., Josse, J. and Pagès, J. Principal Component Methods Hierarchical Clustering Partitional Clustering: Why Would We Need to Choose for Visualizing Data? **2010**.
7. Le, S.; Josse, J.; Husson, F. FactoMineR: An R package for multivariate analysis. *J Stat Softw* **2008**, *25*, 1-18, doi:DOI 10.18637/jss.v025.i01.
8. R Development Core Team R: *A language and Environment for Statistical Computing.*, R Foundation for Statistical Computing: Vienna, Austria, 2019.
9. Pantano, F.; Manca, P.; Armento, G.; Zeppola, T.; Onorato, A.; Iuliani, M.; Simonetti, S.; Vincenzi, B.; Santini, D.; Mercadante, S.; et al. Breakthrough Cancer Pain Clinical Features and Differential Opioids Response: A Machine Learning Approach in Patients With Cancer From the IOPS-MS Study. *JCO Precis Oncol* **2020**, *4*, doi:10.1200/PO.20.00158.
10. Fallon, M.; Giusti, R.; Aielli, F.; Hoskin, P.; Rolke, R.; Sharma, M.; Ripamonti, C.I.; clinicalguidelines@esmo.org, E.G.C.E.a. Management of cancer pain in adult patients: ESMO Clinical Practice Guidelines. *Ann Oncol* **2018**, *29 Suppl 4*, iv166-iv191, doi:10.1093/annonc/mdy152.
11. WHO guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents. Available online: <https://apps.who.int/iris/bitstream/handle/10665/279700/9789241550390-eng.pdf?ua=1> (accessed on 6/6/2021).
12. Davies, A.N.; Elsner, F.; Filbet, M.J.; Porta-Sales, J.; Ripamonti, C.; Santini, D.; Webber, K. Breakthrough cancer pain (BTcP) management: a review of international and national guidelines. *BMJ Support Palliat Care* **2018**, *8*, 241-249, doi:10.1136/bmjspcare-2017-001467.
13. Jiang, C.; Wang, H.; Wang, Q.; Luo, Y.; Sidlow, R.; Han, X. Prevalence of Chronic Pain and High-Impact Chronic Pain in Cancer Survivors in the United States. *JAMA Oncol* **2019**, *5*, 1224-1226, doi:10.1001/jamaoncol.2019.1439.