

Table S2. List of antibodies used in this study.

Antibody	Protein I.D.	origin	serum	dilution	Reference / source
AtpA	Mitochondrial ATP-synthase subunit α	<i>Zea mays</i>	Mouse (monoclonal)	1/500	Thomas Elthon collection, PM014
AtpB	Mitochondrial ATP-synthase subunit β	<i>Zea mays</i>	Mouse (monoclonal)	1/5,000	(Michael <i>et al.</i> 1993)
CA2	γ -carbonic anhydrase-like subunit 2	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1/1,000	(Perales <i>et al.</i> 2005, Sunderhaus <i>et al.</i> 2006)
Cox2	Cytochrome oxidase subunit-2	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1/5,000	Agrisera antibodies, AS04 053A
Nad1	NADH-dehydrogenase complex subunit-1	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1/1000	Gift of Dr. Etienne Meyer, Halle U.
Nad9	NADH-dehydrogenase complex subunit-9	<i>Triticum spp.</i>	Rabbit (polyclonal)	1/50,000	(Lamattina <i>et al.</i> 1993)
RISP	Rieske iron-sulfur protein	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1/5,000	Gift of Prof. Ian Small, UWA
VDAC (PORIN)	Mitochondrial membrane-associated β -barrel proteins	<i>Zea mays</i>	Mouse (monoclonal)	1/1000	Thomas Elthon collection, PM035
AOX1/2	Alternative oxidase	<i>Sauromatum guttatum</i>	Rabbit (polyclonal)	1/500	Agrisera antibodies, AS04 054
Nad6	NADH-ubiquinone oxidoreductase chain 6	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1:1000	Agrisera (AS15 2926)
Cob	Apocytochrome B	<i>Beta vulgaris ssp. maritima</i>	Rabbit (polyclonal)	1:4000	Meyer et al. 2018
CYTC1	Cytochrome C1	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1:1000	Giégé et al. 2005
CYTC	Cytochrome C	<i>Arabidopsis thaliana</i>	Rabbit (polyclonal)	1:1000	Agrisera (AS08 343A)

References

- Giege. P, Sweetlove LJ, Cognat V, Leaver CJ. 2005.** Coordination of nuclear and mitochondrial genome expression during mitochondrial biogenesis in Arabidopsis. *Plant cell*, **17**, 1497-1512.
- Lamattina, L., Gonzalez, D., Gualberto, J. and Grienenberger, J.-M.** (1993) Higher plant mitochondria encode an homologue of the nuclear-encoded 30-kDa subunit of bovine mitochondrial complex I. *Eur J Biochem*, **217**, 831-838.
- Meyer EH, Lehmann C, Boivin S, Brings L, De Cauwer I, Bock R, Kuhn K, Touzet P.** (2018) CMS-G from Beta vulgaris ssp. maritima is maintained in natural populations despite containing an atypical cytochrome c oxidase. *Biochem J*, **475**, 759–773.
- Michael, H.L., Arnost, H. and Elthon, T.E.** (1993) Monoclonal Antibodies to the a- and b- Subunits of the Plant Mitochondrial F₁-ATPase. *Plant Physiology*, **101**, 931-937.
- Perales, M., Eubel, H., Heinemeyer, J., Colaneri, A., Zabaleta, E. and Braun, H.-P.** (2005) Disruption of a nuclear gene encoding a mitochondrial gamma carbonic anhydrase reduces complex I and supercomplex I+III2 levels and alters mitochondrial physiology in Arabidopsis. *J Mol Biol*, **350**, 263-277.
- Sunderhaus, S., Dudkina, N.V., Jansch, L., Klodmann, J., Heinemeyer, J., Perales, M., Zabaleta, E., Boekema, E.J. and Braun, H.P.** (2006) Carbonic anhydrase subunits form a matrix-exposed domain attached to the membrane arm of mitochondrial complex I in plants. *J Biol Chem*, **281**, 6482-6488.