

Article

High Resistance to Antibiotics Recommended in Standard Treatment Guidelines in Ghana: A Cross-Sectional Study of Antimicrobial Resistance Patterns in Patients with Urinary Tract Infections Between 2017 -2021

Benjamin Asamoah^{1*}, Appiah-Korang Labi², Himanshu A. Gupte³, Hayk Davtyan⁴, Georgette Marfo Peprah¹, Forster Adu-Gyan¹, Divya Nair⁵, Karlos Muradyan⁴, Nasreen S Jessani^{6,7} and Paul Sekyere Nyantakyi¹

1. MDS-Lancet laboratories, Accra, Ghana: paul.s-nyantakyi@cerbalancetafrica.com (P.S.N.); forster.adu-ghan@cerbalancetafrica.com (F.A.G.); georgette.peprah@cerbalancetafrica.com (G.M.P.)
 2. World Health Organization, Country office, Accra, Ghana: labia@who.int (A.K.L.)
 3. Narotam Sekhsaria Foundation, Mumbai, India himanshu.gupte@gmail.com (H.G.)
 4. Tuberculosis Research and Prevention Center, Yerevan, Armenia: haykdav@gmail.com (H.D.); carlosmuradyan2@gmail.com (K.M.)
 5. International Union Against TB and Lung Disease (The Union), 75006 Paris, France: Divya.Nair@theunion.org (D.N.)
 6. Centre for Evidence based Health Care, Stellenbosch University South Africa: njessani@sun.ac.za (N.S.J.)
 7. Department of International Health, John Hopkins Bloomberg School of Public Health, USA:
- * Correspondence: benjamin.asamoah@cerbalancetafrica.com (B.A.); +233 27 321 4641

Abstract: Management of urinary tract infection is challenged by increasing antimicrobial resistance (AMR) worldwide. In this study we describe the trends in antimicrobial resistance of uropathogens isolated from the largest private sector laboratory in Ghana over a five-year period. We reviewed positive urine cultures at the MDS Lancet Laboratories from 2017 to 2021. Proportions of uropathogens with antimicrobial resistance to oral and parenteral antimicrobials recommended by the Ghana standard treatment guidelines were determined. The proportion of multi-drug resistant isolates, ESBL and carbapenemase-producing phenotypes were determined. Of 94,134 urine specimens submitted for culture, 20,010 (22.1%) were culture positive. *Enterobacterales* were the commonest group of organisms, *E. coli* (70.6) being the commonest isolate and *Enterococcus* spp. the commonest gram positive (1.3%) organisms. Among oral antimicrobials the highest resistance was observed to ciprofloxacin (62.3%) and cefuroxime (60.2) % and the least resistance to Fosfomycin (1.9%). The least resistance among parenteral antimicrobials was to meropenem (0.3%). Highest multi-drug resistance levels were observed among *Klebsiella* spp. (68.6%) and *E. coli* (64.0%). Extended spectrum beta-lactamase (ESBL) positivity was highest in *Klebsiella* spp. (58.6%) and *E. coli* (50.0%). There may be a need to review the Ghana standard treatment guidelines to reflect increased resistance among uropathogens to recommended antimicrobials.

Keywords: Urinary tract infection; Ghana; Antimicrobial resistance (AMR); Multi-drug resistance (MDR); MDS Lancet Laboratories; Access, Watch and Reserved (AWaRE) classification; Uropathogens;

1. Introduction

Antimicrobial resistance (AMR) threatens human health by limiting the options for treating infections [1]. AMR has been declared as one of the top 10 public health threats to humanity by the World Health Organization, and a direct obstacle to achieving the Sustainable Development Goals [1]. Illnesses resulting from organisms with AMR lead to long hospital stays and require broad spectrum and expensive antimicrobials for treatment, resulting in large financial burdens for patients [1].

Urinary tract infections (UTIs) are one of the leading causes of morbidity and a growing health care expenditure globally affecting 150 million people each year [2][3]. Morbidity and mortality from UTIs is higher in lower middle-income countries (LMICs) than in higher income countries[4]. UTIs are most commonly caused by uropathogenic *Escherichia coli* (UPEC) *Klebsiella pneumoniae* and *Enterococcus faecalis* [5].

Widespread and indiscriminate use of antibiotics has led to the development of an alarming level of AMR among uropathogens, including the emergence of extended spectrum beta-lactamase (ESBL) and carbapenemase producing organisms [6]. ESBL and carbapenemase producing organisms are typically multi-drug resistant[6]. This mandates the use of the Watch and Reserve group of antibiotics from the AWaRe (Access, Watch, Reserve) classification of the WHO [7]: these antibiotics are typically parenteral and may require hospitalization for the purposes of administration [8]. Common risk factors associated with resistant UTI are urinary catheterization, previous hospitalization, previous antibiotic use, and residence in a nursing home facility[9].

Although the burden of UTIs in Africa is not clearly documented, prevalence rates ranging between 10.1% and 76.6% have been reported in different sub-populations from across the continent [9–11]. In the same studies AMR rates as high as 95% have been reported to antibiotics used for treating UTIs [9–11]. Whilst there is limited information about the general national prevalence of UTIs in Ghana, two studies have reported 42.75% prevalence in pregnant women [13] and 86% among hospitalized adults [14]. The proportion of multi-drug resistant (MDR) UTIs have been reported to be as high as 93.6% in Ghana [10], [12].

AMR is a recognized threat in Ghana and there is a national policy and action plan based on the global action plan on AMR[15]. Ghana is currently amongst the countries that is reporting to the global AMR surveillance system (GLASS) starting from 2021[16].

A nation-wide study exploring AMR patterns in various uropathogens across several years and in different regions of Ghana can strengthen the knowledge and evidence base to guide policy and practice in line with five objectives (Awareness, Surveillance, Infection prevention and control, Antimicrobial usage and Research and innovation) of the global action plan of AMR by World Health Organization [16].

MDS Lancet Laboratories Gh. LTD is a member of the Cerba Lancet Africa group, a leading network of clinical pathology and medical diagnostics in Africa. MDS Lancet Laboratories Gh. LTD is currently the largest private pathology laboratory in Ghana, providing service to healthcare facilities in the private and public sectors

In this study, we aimed to identify the AMR patterns and trends of MDR among uropathogens isolated from urine samples of patients with suspected UTI processed at MDS Lancet Laboratories, Ghana, over a five-year period, from 2017 to 2021. Specifically, we sought to determine trends in i) the yield of culture testing and proportions of all identified species, ii) the resistance patterns of antibiotics grouped by route of administration and WHO AwaRe classification, iii) factors associated with AMR and iv) trends in *E.coli* antibiotic resistance over the study period.

2. Materials and Methods

2.1 Study Design

This was a cross-sectional study involving secondary analysis of laboratory data of urine cultures and antimicrobial susceptibility testing conducted at the MDS Lancet Laboratories, Ghana, between 2017 and 2021.

2.2. Study Setting

2.2.1. General Setting

Ghana is classified as a low-middle income Country (LMIC) in West Africa with a total population around 30.8 million. Accra is the capital city located at the coastal belt and has a total population of about 5 million [17]. Health care in Ghana is mainly provided by public sector facilities but with significant contribution from private sector facilities including laboratories. The public healthcare system consists of tertiary care hospitals, district (secondary) hospitals, and primary health care facilities [18]. According to the law, patients need prescriptions from a medical doctor to get antibiotics [15]. However, it is possible to get antibiotics without prescriptions from private pharmacies. The Ghana Standard Treatment Guidelines (2017) suggest the following antibiotics for the treatment of UTI: ciprofloxacin, cefuroxime and amoxicillin/clavulanic acid for uncomplicated infections and gentamicin and ceftriaxone for complicated infections [19].

2.2.2. Specific setting

MDS Lancet Laboratory Gh LTD is a private medical laboratory with 29 locations dotted 10 out of 16 administrative regions of Ghana. The Laboratory processes about 148,500 samples per month and receives more than 50,000 urine (routine examination and cultures) samples per year. All testing and information management processes are conducted under strict quality control mechanisms as defined in the laboratory's standard operating procedures. The laboratory holds ISO 15189 2012 certification. The headquarters/main laboratory is located in Accra The microbiology department where this research was carried out has standard quality assurance mechanisms in place at pre analytical, analytical and post analytical stages.

2.2.3. Sample Collection and Processing

Urine samples for this study originated from patients diagnosed with suspected UTI in health care facilities (hospitals and clinics) in both the public and private sectors. Urine samples are usually transported by patients or their relatives in sterile urine containers. All microbiology samples from the different branches that are meant for culture are transported to the microbiology laboratory in Accra in less than 24 hours for processing using temperature controlled sample transportation bags

All urine samples received in the laboratory are registered and plated on cysteine lactose electrolyte deficient (CLED) agar, McConkey, blood agar and antimicrobial plates [20]. The first three media are used to determine the presence of microorganisms in the sample, while the latter is used to determine the presence of antibiotics in the urine and provides information on the use of antibiotics prior to sample collection. After 24 hours of incubation, cultures are assessed and positive samples are further processed for identification of the specific microorganisms by performing indole and oxidase and other biochemical tests, as well as the Analytical Profile Index (API) or Microscan [21]. Antimicrobial susceptibility testing is performed using the Kirby Bauer disc diffusion method on Mueller Hinton agar [22]. Zones of inhibition are interpreted using the clinical laboratory standard institute (CLSI) guidelines [23].

2.3. Study Population

The study population included all urine samples submitted for culture and susceptibility testing at the MDS Lancet Laboratories in Ghana from January 2017 to December 2021.

2.4. Data variables

For this study the following variables were extracted from the laboratory database and used in the analysis: Patient ID and demographic characteristics, sample collection date and results entry date, the isolated organisms, ESBL status (for gram negatives), and antibiotic susceptibility patterns.

2.5. Sources of data, data collection and validation

The MDS Lancet Laboratory uses an electronic laboratory information system (LIS) where all the data related to the patient and samples are entered. Right from the collection of the sample, information is entered into the system and verified at each stage from transportation to processing of the samples. Results of the tests are reported onto the same LIS and the information becomes readily available for collection at any MDS Lancet branch. The LIS of the MDS Lancet Laboratories was used to extract data for the study period. It was extracted into MS Excel format and analyzed using STATA® (version 16.0 Copyright 1985-2019 StataCorp LLC).

2.6. Statistical Analysis

Frequency and proportions were used to summarize demographic and microbiological characteristics. MDR was defined as resistance to at least one antimicrobial agent from three or more classes of antibiotics [24]. Presence of resistance to any antibiotic and MDR among all culture positive samples were summarized as proportions with 95% confidence interval (95% CI). Resistance to commonly prescribed antibiotics (individually and grouped according to AWaRe classification) for different isolates was reported as a percentage. To identify factors independently associated with UTI due to MDR organisms in a patient, modified Poisson regression with robust variance estimator was carried out. A patient was considered to have UTI due to an MDR organism if any of the isolates from their urine sample turned out to be MDR. As this was a cross-sectional study, prevalence ratios (PR) and adjusted prevalence ratios (aPR) were reported as measures of association. Levels of significance for all statistical tests was set at a p-Value of 0.05 [25].

3. Results

3.1. Patient characteristics and proportions of all identified bacterial isolates in urine samples

Over the five years, a total of 328,073 urine samples were submitted for processing. Of these, 94,134 samples were submitted for culture and susceptibility testing at the MDS Lancet laboratories and 20,010 (22.1%) yielded at least one organism on culture. Among the culture positive samples, 793 (4%) yielded more than one organism. Table 1 shows the characteristics of patients with culture positive urine samples. Most culture positive samples (38.8%) were from patients aged 15-44 years. Positive urine samples were more common in females (72.5%) compared to males, and the majority of positive urine samples were obtained from the Greater Accra region (63.5%).

Antibiotic residue was present in 1,986 (9.9%) culture positive samples. Isolates resistant to at least one antibiotic were found in 65.3% (95% CI: 64.6-66.0%) of cultures and multi-drug resistant isolates in 63% (95% CI: 62.3-63.6%) of cultures.

Table 1. Socio-demographic and microbiological characteristics of patients with urinary tract infections with culture positive samples¹ analyzed at MDS Lancet Laboratories, Ghana, from 2017 to 2021.

Characteristics	N	(%)
Total number of patients	20,010	(100)
Age in years		
<15	1,474	(7.4)
15-44	7,771	(38.8)
45-64	4,539	22.7

	≥65	6,040	(30.2)
	Unknown	186	0.9
Sex	Male	5,484	(27.4)
	Female	14,505	(72.5)
	Unknown	21	(0.1)
Geographic location of urine specimens	Accra	12,702	(63.5)
	Ashanti	4,266	(21.3)
	Others	3,042	(15.2)
Antimicrobial substance in urine ²	Present	1,986	(9.9)
Number of bacterial isolates identified	One	19,217	(96.0)
	Two	786	(3.9)
	Three	7	(<0.1)
Antimicrobial resistance	Resistance to at least one anti-microbial	13,079	(65.3)
	MDR ³ in at least one isolate	12,609	(63.0)

¹Culture positive samples mean the urine sample yielded at least one bacterial isolate; ²Presence of antimicrobial substance in urine sample implies that patients were on antibiotics before their urine sample was taken; ³MDR- Multi-drug resistance defined as resistance to three or more antibiotic drug classes.

3.2. Uropathogen Characteristics

Table 2 shows the distribution of uropathogens from urine cultures. *Enterobacterales* were the commonest group of organisms with *Escherichia coli* (70.6%), *Klebsiella* spp. (15.0%) and *Proteus* spp. (5.4%), being the most common, while *Enterococcus* spp (1.3%) was the commonest Gram-positive organism isolated. This pattern of organism distribution was similar across all age groups, region of sample origin and year of specimen collection. The full list of isolated pathogen and patient characteristics are provided in supplementary Table S1.

Table 2. Distribution of common uropathogens isolated from urine samples of patients with urinary tract infections analyzed at MDS Lancet Laboratories Ghana from 2017 to 2021 based on age and region.

	Gram negative bacteria (GNB)											Gram positive bacteria (GPC)		
	<i>E. coli</i>		<i>Klebsiella</i> spp.		<i>Proteus</i> spp.		<i>Pseudomonas</i> spp.		<i>Acinetobacter</i> spp.		Other GNB			
	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ¹	n (%) ²	n (%) ²
Age groups in years														
<15	997	(6.8)	272	(8.7)	104	(9.3)	5	(5.0)	9	(18.8)	143	(9.9)	26	(8.8)
15-44	5,887	(40.1)	1,141	(36.7)	374	(33.4)	15	(15.0)	17	(35.4)	501	(34.8)	137	(46.4)
45-64	3,329	(22.7)	735	(23.6)	237	(21.2)	21	(21.0)	4	(8.3)	309	(21.5)	64	(21.7)
≥65	4,341	(29.5)	942	(30.3)	395	(35.3)	59	(59.0)	18	(37.5)	472	(32.8)	65	(22.0)
Regions														
Accra	9,437	(64.2)	1,921	(61.7)	723	(64.6)	65	(65.0)	27	(56.3)	824	(57.2)	186	(63.1)
Ashanti	3,080	(21.0)	659	(21.2)	280	(25.0)	18	(18.0)	6	(12.5)	348	(24.2)	77	(26.1)
Others	2,178	(14.8)	532	(17.1)	117	(10.4)	17	(17.0)	15	(31.3)	268	(18.6)	32	(10.8)
Total	14,695	(100)	3,112	(100)	1,120	(100)	100	(100)	48	(100)	1,440	(100)	295	(100)

¹Percentages reported out of all culture positive samples with gram negative bacilli (GNB) isolates reported for each category; ²Percentages reported out of all culture positive samples with gram positive cocci (GPC) isolates reported for each category.

3.3. Trends in urinary bacterial isolates over five years, 2017-2021

Annual trends in urinary bacterial isolates for the top five GNBs and gram positive organisms over five years (2017-2021) are shown in Figure 1.




Figure 1. Annual trends in number of isolates for the top five GNBs and gram positives from urine samples of patients with urinary tract infections at MDS Lancet laboratories Ghana between 2017 to 2021.

Total number of isolates over the years per year are as follows. 2017 (1,568), 2018 (2,456), 2019 (4,193), 2020 (5,050), and 2021 (7,543)

Figure 1 shows a general increase in the number of isolates for the top five GNBs over the study period.

¹Number for each year is the total of each isolate for the particular year

²Other GNB: all gram negative bacilli isolated in the laboratory apart from *E.coli*, *Klebsiella* spp., *Proteus* spp., *Pseudomonas* spp. and *Acinetobacter* spp.

GPC: Gram positive cocci

3.4. Resistance patterns of antibiotics grouped by route of administration and WHO AWaRe classification

Tables 3 and 4 show the proportion of the most common uropathogens resistant to oral antibiotics as recommended by the national standard treatment guidelines including fosfomycin, and common parenteral antibiotics which were likely to be used as second line agents for treatment. Among *Enterobacteriales*, the least resistance was observed to fosfomycin - *E. coli* (2.9%), *Klebsiella* spp. (9.9%) and *Proteus* spp. (14.0%). The highest resistance was observed to ciprofloxacin (62.3%) and cefuroxime (60.2%). Among *Enterococcus* spp., the least resistance was observed to nitrofurantoin (1.5%).

The least resistance to parenteral antibiotics among *Enterobacteriales* was to meropenem (0.3%) and tigecycline (1.7%). Among isolated *Enterococcus* spp., no resistance was observed to vancomycin. Among ESBL positive organisms the least resistance observed among oral antibiotics was to fosfomycin (5.4%) and nitrofurantoin (41.6%). Among the parenteral antibiotics, the least resistance was observed to amikacin (5.3%) and meropenem (1.2%). Details of the antimicrobial resistance patterns among ESBL positive organisms are provided in supplementary Table 2.

Table 3. Prevalence of antimicrobial resistance to common oral antibiotics used in treating UTI in Ghana among the top six GNBs and *Enterococcus spp* isolated from urine samples of patients with UTI at MDS Lancet laboratories Ghana between 2017 to 2021.

	Ampicillin	Amoxicillin clavulanate	Nitrofurantoin	Ciprofloxacin	Cefuroxime	Fosfomycin
	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹
<i>E. coli</i>	N/A	9,411 (64.0)	3,575 (24.3)	9,161 (62.3)	8,851 (60.2)	435 (2.9)
<i>Klebsiella spp</i>	N/A	2,083 (66.9)	1,639 (52.7)	1,659 (53.3)	1,986 (63.8)	308 (9.9)
<i>Proteus spp</i>	N/A	105 (9.4)	N/A	151 (13.5)	91 (8.13)	6 (14.0)
<i>Acinetobacter spp</i>	N/A	N/A	N/A	9 (18.8)	N/A	N/A
<i>Pseudomonas spp</i>	N/A	N/A	N/A	41 (41)	N/A	N/A
<i>Enterococcus faecalis</i>	9 (3.3)	N/A	4 (1.5)	N/A	N/A	5 (1.9)

¹Percentages calculated with total number of urine samples which yielded the particular species as denominator. Antibiotics in **Green** belong to Access group and those in **Yellow** belong to Watch group as per WHO AWaRe classification 2021; UTI- urinary tract infection, GNB- gram negative bacilli, GPC-gram positive cocci, N/A-Not applicable (not tested for the isolate).

Table 4. Prevalence of antimicrobial resistance to common parenteral antibiotics used in treating UTI in Ghana among top uropathogens isolated from urine samples of patients with UTI at MDS Lancet Laboratories, Ghana, between 2017 to 2021.

Organism	Ak	Caz	Cro	Mem	Piptaz	Tige
	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹	N (%) ¹
<i>E. coli</i>	461 (3.1)	7,147 (48.62)	7,362 (50.1)	42 (0.3)	7,148 (48.6)	252 (1.7)
<i>Klebsiella spp</i>	116 (3.7)	1,802 (57.9)	1,834 (58.9)	56 (1.8)	1,803 (57.9)	173 (5.6)
<i>Proteus spp</i>	3 (0.3)	46 (4.1)	46 (4.1)	3 (0.3)	17 (39.5)	N/A
<i>Acinetobacter spp</i>	2 (4.2)	17 (35.4)	N/A	1 (2.1)	14 (29.2)	N/A
<i>Pseudomonas spp</i>	21 (21)	30 (30)	N/A	19 (19)	27 (27.0)	N/A

¹Percentages calculated with total number of urine samples which yielded the particular species as denominator. Antibiotics in **Green** belong to Access group, those in **Yellow** belong to Watch group and those in **Red** belong to Reserve Group as per WHO AWaRe classification 2021. Cro: Ceftriaxone, Cp: Cefepime, Piptaz: Piperacillin tazobactam, Caz: Ceftazidime, Mem: Meropenem, Ak: Amikacin, Vanc: Vancomycin, Tige: Tigecycline, UTI- urinary tract infection, GNB- gram negative bacilli, N/A- not applicable.

Resistance of *E. coli* to oral antibiotics (amoxicillin/clavulanic acid, cefuroxime, ciprofloxacin, fosfomycin and nitrofurantoin) was constant over the study period as shown in Figure 2.




Figure 2. Annual trends in proportion of resistance to selected antimicrobials among *E. coli* isolated from urine samples of patients with urinary tract infections at MDS Lancet laboratories Ghana between 2017 to 2021. ¹Proportion for each year calculated with number of samples in which *E. coli* was isolated in the year as denominator (2017= 1256, 2018=1877, 2019=3078, 2020=3469, 2021=5015); AUG: Amoxicillin/clavulanic acid, CXM: Cefuroxime, CIP: Ciprofloxacin, FOS: Fosfomycin, NIT: Nitrofurantoin, ESBL: Extended spectrum beta lactamase.

3.5. Multi-drug resistance and ESBL positivity

The proportion of isolates, which showed MDR was highest in *Klebsiella* spp. (68.6%) and *E. coli* (64.0%). Among *Enterobacterales*, ESBL positivity was highest among *Klebsiella* spp. (58.6%) followed by *E. coli* (50.0%) and least in *Proteus* spp. (5.3%). In addition, the proportions of ESBL were stable across the study years. Details of trends in MDR and ESBL positivity among commonly isolated uropathogens are provided in Table 5. Gram positive bacteria are not presented due to low number of resistant isolates.

Table 5. Prevalence of multi drug resistance (MDR) and extended spectrum beta lactamase (ESBL) positivity among commonly isolated bacteria causing UTI in patients at MDS Lancet Laboratories in Ghana from 2017 to 2021.

Organism	MDR						ESBL					
	2017 N (%) ¹	2018 N (%) ¹	2019 N (%) ¹	2020 N (%) ¹	2021 N (%) ¹	Overall N (%) ¹	2017 N (%) ²	2018 N (%) ²	2019 N (%) ²	2020 N (%) ²	2021 N (%) ²	Overall N (%) ²
Gram negatives												
<i>E. coli</i>	871 (69.4)	1,309 (69.7)	1,958 (63.6)	2,182 (62.9)	3,087 (61.6)	9,407 (64.0)	600 (47.8)	975 (51.9)	1,585 (51.5)	1,692 (48.8)	2,492 (49.7)	7,344 (50.0)
<i>Klebsiella</i> spp.	137 (74.5)	216 (77.4)	376 (65.2)	522 (68.3)	885 (67.7)	2,136 (68.6)	106 (57.6)	181 (64.9)	329 (57.0)	441 (57.7)	765 (58.5)	1,822 (58.6)
<i>Proteus</i> spp.	4(6.8)	14 (11.6)	33 (15.8)	32 (11.4)	59 (13.1)	142 (12.68)	4 (6.8)	6 (5.0)	16 (7.7)	11 (3.9)	22 (4.9)	59 (5.3)
<i>Pseudomonas</i> spp.	0 (0.0)	1 (16.7)	9 (40.9)	14 (51.8)	16 (35.6)	40 (40)	N/A	N/A	N/A	N/A	N/A	N/A
<i>Acinetobacter</i> spp.	0 (0.0)	0 (0)	0 (0.0)	6 (31.6)	4 (18.2)	10 (20.8)	N/A	N/A	N/A	N/A	N/A	N/A
Other Gram-negative rods	65 (94.3)	140 (89.1)	220 (85.3)	348 (86.1)	425 (77.0)	1,198 (83.2)	44 (63.8)	90 (57.3)	131 (50.8)	126 (31.2)	185 (33.5)	576 (40)

¹Percentage of MDR isolates for each bacterial species is calculated with number of samples which yielded the particular species as denominator; ²Percentages of ESBL producing isolates for each bacterial species is calculated with number of samples which yielded the particular species as denominator; ³ N/A =Not applicable.

3.6. Factors associated with antimicrobial resistance

Age, gender, specimen location and presence of antibiotic residue in urine were found to be independently associated with a patient having an MDR uropathogen (Table 6). Patients aged 65 years and above (aPR 1.16, 95%CI: 1.11-1.22) and males (aPR: 1.13, 95%CI: 1.11-1.16) had significantly higher risk of UTI due to a multi-drug resistant uropathogen. Patients with antibiotic residues in their urine (aPR 1.40, 95%CI: 1.37-1.43) had significantly higher risk of UTI due to a multi-drug resistant uropathogen.

Table 6. Factors associated with multi-drug resistance in patients with urinary tract infections with culture positive samples tested at MDS Lancet Laboratories in Ghana between 2017-2021.

Factors	Total N ¹	MDR n(%) ²	PR (95% CI)	aPR (95% CI)	p-value
Age in years					
<15	1,474	854 (57.9)	Ref	Ref	
15-44	7,771	4,595 (59.1)	1.01 (0.97-1.06)	1.05 (0.99-1.09)	0.06
45-64	4,539	3,020 (66.5)	1.14 (1.09-1.20)	1.14 (1.08-1.19)	<0.001
>=65	6,040	4,207 (69.6)	1.20 (1.14-1.25)	1.16 (1.11-1.22)	<0.001
Sex					
Male	5,484	3,989 (72.7)	1.20 (1.17-1.22)	1.13 (1.11-1.16)	<0.001
Female	14,505	8,783 (60.5)	Ref	Ref	
Specimen location³					
Accra	12,702	7,931 (62.4)	Ref	Ref	
Ashanti	4,266	2,722 (63.8)	1.02 (0.99-1.05)	1.04 (1.01-1.07)	0.002
Others	3,042	2,131 (63.8)	1.12 (1.09-1.15)	1.09 (1.07-1.13)	<0.001
Antimicrobial substance in urine					
Present	1,986	1,742 (87.1)	1.43 (1.40-1.46)	1.40 (1.37-1.43)	<0.001
Absent	18,007	11,030 (61.2)	Ref	Ref	

¹Missing values in age (186), sex (21) and antimicrobial substance in urine (17) dropped from analysis; ²Row percentages; ³Specimen locations are the regions from where the samples were collected. Samples from regions outside greater Accra were well packaged and transported in temperature regulated specimen bags; MDR-Multidrug resistance; PR- prevalence ratio; CI - confidence interval; aPR- adjusted prevalence ratio.

4. Discussion

In this study, the culture yield over the study period was 22.1%. This is similar to results obtained in other studies in which the positivity ranged from 17% to 37% [26],[27],[27]. *Enterobacteriales* were the commonest group of pathogens isolated with *E. coli* and *Klebsiella* spp. being the commonest isolates and *Enterococcus* spp. being the commonest Gram-positive organisms. These organisms predominate in the human gut flora, which is known to be the common source of autoinfection for UTIs. These findings are similar to the results in other studies where *E. coli* at 68.3% and *Klebsiella pneumoniae* at 31.7% were the most predominant isolates from UTI [29] In another study from India, *E.coli* was the predominant isolate at 77.9% followed by *Klebsiella* spp. at 22.1% [30]

High levels of resistance to Standard Treatment Guidelines (STG) recommended antibiotics for treating urinary tract infection was observed in this study. This situation was more acute among *Enterobacteriales*. For example, among *E. coli* isolates resistance to cefuroxime and ciprofloxacin was 60.2% and 62.3% respectively. This situation can result in poor treatment outcomes including prolonged treatment for empirically treated UTIs. Oral antibiotics with least resistance in this study were nitrofurantoin and fosfomycin. Fosfomycin is a new agent on the Ghanaian market and is not widely available. Thus, its use is restricted, and it is expected that resistance to this antibiotic would be low. Although nitrofurantoin is an old drug, side effects arising from its use has may have resulted in its limited use and consequently the relatively low resistance observed compared to other agents [31],[32],[32].

In a recent review of AMR among uropathogens in the Asia-Pacific region a similar pattern was observed; resistance to commonly used drugs such as ciprofloxacin, sulfamethoxazole-trimethoprim and ceftriaxone ranged from 33% to 90%, with low resistance seen to nitrofurantoin between 2.7% to 34% and to fosfomycin between 1.7%-1.8% [34]

Resistance to parenteral antibiotics such as ceftriaxone and gentamicin recommended for the treatment of complicated UTI's was also high. This may be due to the relatively high prevalence of ESBL observed among *Enterobacterales*. ESBL genes are typically borne on plasmids which are known to also carry multiple resistance genes making affected organisms multi-drug resistant [35]. The proportion of ESBL among *Enterobacterales* in this study was high, similar to findings from other studies from Ghana [12].

The proportion of ESBL among organisms seemed to be constant over the 5 years under consideration. Our study suggests that agents useful against ESBL producing organisms are fosfomycin, meropenem and amikacin. These agents may be considered for empirical treatment of UTI when risk factors for MDR infection such as previous hospitalizations or antibiotic use exist. MDR was a common characteristic among the majority of uropathogens isolated. This suggests AMR may be a major problem in Ghana.

Among the different age groups the risk of having an MDR infection was associated with older patients. Increasing age is associated with increased use and misuse of antibiotics, a situation likely to increase MDR risk. In other studies, increasing age has been found to be associated with increased risk of having a MDR UTI and hence increased use of antibiotics [35],[37]. A study in the United Kingdom (UK) showed an increased risk of MDR with age with odds ratios of 1.8, 2.69, 3.22 and 3.62 for the age ranges 50-59, 60-69, 70-79 and >80 years respectively [37]

Whilst the prevalence of UTI was common in women in this study, the risk of MDR infection was significantly higher in males compared to women. UTI in males is usually associated with a urological abnormality such as prostatic hyperplasia or posterior urethral valves which are usually associated with instrumentation and risk of infection [38][39][40]. Lastly, the highest risk of MDR was found in patients with the presence of antibiotics in the urine. This presupposes that most of these patients have difficulty to treat infections and have probably been started on ineffective antibiotics or have misused antibiotics, all of which may have contributed to the development of resistance.

A major strength of this study is the large number of isolates included from different regions of Ghana, and the study represents one of the largest datasets on AMR in uropathogens on the African continent as well as in Ghana. This data may give a better understanding of AMR in uropathogens in Ghana, although the preponderance of data from the Greater Accra and Ashanti regions may introduce some bias into data representativeness. Another strength of the study is that we followed STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) guidelines to report findings of our study [41].

A major limitation of this study, however, was the inability to segregate the data into outpatient and inpatient because of data unavailability. Such data would have allowed for the segregation of community acquired and hospital acquired infections and their association with AMR. This study being a laboratory only based surveillance has the potential to overestimate the level of resistance among uropathogens as has been shown in other studies [42].

This study has significant implications for policy and practice. Based on the current data, there may be a need to revise the STG guidelines on antibiotics to be used for treating UTIs, especially complicated infections to reflect the large proportions of AMR observed among uropathogens. As these guideline changes are considered, there will be a need to implement antibiotic stewardship measures to prolong the usefulness of agents such as fosfomycin, nitrofurantoin, amikacin and meropenem which may be recommended as alternate antibiotics. Also, there will be a need to investigate the utility of other oral antibiotics for treating UTIs to offer clinicians and patients more therapeutic options in the face of growing MDR infections. In the long term, there is a need to establish laboratory-based

surveillance of uropathogens nationwide, augmented by large population-based studies covering large geographical regions to inform policy and practice. This is important to correct for the selection bias of laboratory surveillance for MDR organisms. Also, clinicians should be encouraged to utilize culture and susceptibility testing routinely during the management of UTIs to avoid poor treatment outcomes because of resistant strains.

5. Conclusions

Over the five-year period, Gram negative pathogens especially *E. coli* were the commonest cause of UTIs. High levels of resistance to STG recommended oral and parenteral antibiotics for the management of UTIs were observed. There may be a need to review the current STG guidelines to reflect the high numbers of AMR in the most common isolates.

Author Contributions: Conceptualization, B.A., H.G., H.D. and A.K.L.; methodology, B.A., H.G., H.D. and A.K.L.; software, B.A., D.N., K.M. and A.K.L.; data curation and validation, B.A., F.A.G., D.N. and G.P.M.; formal analysis; D.N., B.A. and K.M.; resources, P.S.N., H.G. and H.D.; data curation; writing—original draft preparation, B.A., H.G., H.D., D.N., K.M. and A.K.L.; writing—review and editing, all authors; visualization, N.S.J.; supervision, A.K.L., P.S.N., H.G. and H.D.; project administration, B.A., A.K.L. and P.S.N. All authors have read and agreed to the published version of the manuscript.

Funding: This SORT IT AMR Programme was funded by the National Institute of Health Research, Department of Health & Social Care of the United Kingdom and supported by implementing partners

Institutional Review Board Statement: Permission to use the data was obtained from the CEO of MDS-Lancet laboratories Gh LTD. The study received approval from the Noguchi Memorial Institute for Medical Research (NMIMR) NMIMR-IRB CPN 041/21-22) dated 2nd March 2022). The Ethics Advisory Group of the International Union against Tuberculosis and Lung Disease, Paris, France (approval No 41/21 dated 13 December 2021), also approved the study

Informed Consent Statement: No patient consent was required as secondary data was extracted from the electronic database of MDS-lancet Laboratories.

Data Availability Statement: The data presented in this study are available on request from the corresponding author

Acknowledgments: This research was conducted through the global partnership of Structured Operational Research and Training Initiative (SORT IT), coordinated by TDR, the Special Programme for Research and Training in Tropical Diseases at the World Health Organization (TDR). The specific SORT IT program that led to these publications included a partnership of TDR with WHO Ghana country office, The International Union Against Tuberculosis and Lung Diseases, Paris; Institute of Tropical Medicine. MDS-Lancet laboratories, Accra, Ghana; Narotam Sekhsaria Foundation, Mumbai, India; Tuberculosis Research and Prevention Center, Yerevan, Armenia; The INCLIN Trust International, New Delhi, India; Nasreen for the Review of this manuscript.

The authors wish to acknowledge Ntethelelo Mthimunya and the entire team of Lancet laboratories ELAB, South Africa for their support with extracting data from MEDITECH

Conflicts of Interest: The authors declare no conflict of interest. The authors of the paper designed, this study, collected the data, analyzed, interpreted and did all the writing without the funders playing any role

References

- [1] World Health Organization (WHO), 'Antimicrobial resistance', *Factsheet*, 2020. .
- [2] B. Zalewska-pi, 'Phage Therapy as a Novel Strategy in the Treatment of Urinary Tract Infections Caused by E. Coli', 2020.
- [3] M. Joshua and M. Raymond, 'Virulence factors and antibiotic resistance patterns of uropathogenic *Escherichia coli*', *African J. Microbiol. Res.*, vol. 8, no. 43, pp. 3678–3686, 2014, doi: 10.5897/ajmr2014.7034.
- [4] S. WE and N. SR, 'Urinary tract infections: disease panorama and challenges', *J. Infect. Dis.*, vol. 183 Suppl, pp. S1–S4, Mar.

- 2001, doi: 10.1086/318850.
- [5] A. L. Flores-Mireles, J. N. Walker, M. Caparon, and S. J. Hultgren, 'Urinary tract infections: Epidemiology, mechanisms of infection and treatment options', *Nat. Rev. Microbiol.*, vol. 13, no. 5, pp. 269–284, May 2015, doi: 10.1038/NRMICRO3432.
- [6] R. Bartoletti *et al.*, 'EAU GUIDELINES ON UROLOGICAL INFECTIONS', no. March, pp. 247–269, 2017.
- [7] G. A. Plan, A. Resistance, I. H. Regulations, T. Atb, and N. Atb, 'A Technical Guide to Implementing the World Health Organization 's AWARe Antibiotic Classification in MTaPS Program Countries', no. May 2015, pp. 1–7, 2019.
- [8] K. S. Kaye, H. H. Nguyen, and M. J. Rybak, 'Clinical Review GUIDELINES FOR MANAGEMENT AND CARE TRANSITIONS IN THE EMERGENCY', *J. Emerg. Med.*, vol. 48, no. 4, pp. 508–519, 2015, doi: 10.1016/j.jemermed.2014.12.001.
- [9] J. Tenney, N. Hudson, H. Alnifaidy, J. T. C. Li, and K. H. Fung, 'Risk factors for acquiring multidrug-resistant organisms in urinary tract infections: A systematic literature review', *Saudi Pharm. J. SPJ*, vol. 26, no. 5, p. 678, Jul. 2018, doi: 10.1016/J.JSPS.2018.02.023.
- [10] O. Patients and E. S. Donkor, 'Urinary Tract Infections among Bladder Outlet Obstruction Patients in Accra, Ghana: Aetiology, Antibiotic Resistance, and Risk Factors', 2018, doi: 10.3390/diseases6030065.
- [11] R. Z. Sangeda, F. Paul, and D. M. Mtweve, 'Prevalence of urinary tract infections and antibiogram of uropathogens isolated from children under five attending Bagamoyo District Hospital in Tanzania : A cross-sectional study [version 1 ; peer review : awaiting peer review]', pp. 1–10, 2021.
- [12] P. Z. Horlortu, N. T. K. D. Dayie, N. Obeng-nkrumah, and A. Labi, 'Community acquired urinary tract infections among adults in Accra , Ghana', 2019.
- [13] A. O. Forson, W. B. Tsidi, D. N. Adjei, M. N. Quarchie, and N. O. Nkrumah, '*Escherichia coli* bacteriuria in pregnant women in Ghana : antibiotic resistance patterns and virulence factors', *BMC Res. Notes*, no. 2018, pp. 1–7, 2019, doi: 10.1186/s13104-018-3989-y.
- [14] E. Gambrah, A. Owusu-ofori, E. Biney, C. Opong, and S. E. Cof, 'International Journal of Infectious Diseases Diagnosis and treatment of urinary tract infections in hospitalized adults in Ghana : The role of the clinical microbiology laboratory in improving antimicrobial stewardship', vol. 102, pp. 497–500, 2021, doi: 10.1016/j.ijid.2020.10.068.
- [15] S. Kwame Yevutsey *et al.*, 'Situational analysis of antibiotic use and resistance in Ghana: policy and regulation', doi: 10.1186/s12889-017-4910-7.
- [16] 'Global action plan on antimicrobial resistance'. [Online]. Available: https://apps.who.int/iris/bitstream/handle/10665/193736/9789241509763_eng.pdf?sequence=1.
- [17] E. B. Agyekum, F. Amjad, M. Mohsin, and M. N. S. Ansah, 'A bird's eye view of Ghana's renewable energy sector environment: A Multi-Criteria Decision-Making approach', *Util. Policy*, vol. 70, no. June 2020, p. 101219, 2021, doi: 10.1016/j.jup.2021.101219.
- [18] 'Ministry of Health | Ghana official website [internet]. Role and function'. 2015.
- [19] Ministry of Health - Ghana National Drugs Programme (GNDP), *Standard Treatment Guidelines*, no. May. 2017.
- [20] B. V. S. and Prashanth, 'Urine culture contamination: a one-year retrospective study at the tertiary care hospital', *Dec. 29, 2014*.
<https://go.gale.com/ps/i.do?id=GALE%7CA469639434&sid=googleScholar&v=2.1&it=r&linkaccess=abs&issn=22784748&p=HRCA&sw=w&userGroupName=anon-70fd146c> (accessed Nov. 05, 2021).
- [21] S. T. Altheide, 'Biochemical and Culture-Based Approaches to Identification in the Diagnostic Microbiology Laboratory'. <http://clsjournal.ascls.org/content/early/2020/01/20/ascls.119.001875> (accessed Nov. 05, 2021).
- [22] 'Antimicrobial Susceptibility Testing by the Kirby-Bauer Disc Diffusion Method'. <http://www.annclinlabsci.org/content/3/2/135.short> (accessed Nov. 05, 2021).
- [23] CLSI, A. L. Dolinsky, R. K. Ohiro, W. Fan, C. Xiao, and F. Wu, 'National Committee for Clinical Laboratory Standards. 2000.

- Performance standard for antimicrobial susceptibility testing. Document M100–S10.’, *J. Int. Med. Res.*, vol. 46, no. September, p. 18, 2019, [Online]. Available: http://www.epa.gov/nerlcwww/1604sp02.pdf%0Ahttp://www.emeraldinsight.com/doi/10.1108/08876049410065598%0Ahttp://clsi.org/media/1469/m100s27_sample.pdf%0Ahttp://shop.clsi.org/site/Sample_pdf/M100S27_sample.pdf.
- [24] A. P. Magiorakos *et al.*, ‘Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance’, *Clin. Microbiol. Infect.*, vol. 18, no. 3, pp. 268–281, 2012, doi: 10.1111/j.1469-0691.2011.03570.x.
- [25] L. A. McNutt, C. Wu, X. Xue, and J. P. Hafner, ‘Estimating the relative risk in cohort studies and clinical trials of common outcomes’, *Am. J. Epidemiol.*, vol. 157, no. 10, pp. 940–943, 2003, doi: 10.1093/aje/kwg074.
- [26] R. De Rosa, S. Grosso, G. Lorenzi, G. Bruschetta, and A. Camporese, ‘Clinica Chimica Acta Evaluation of the new Sysmex UF-5000 fluorescence flow cytometry analyser for ruling out bacterial urinary tract infection and for prediction of Gram negative bacteria in urine cultures’, *Clin. Chim. Acta*, vol. 484, no. May, pp. 171–178, 2018, doi: 10.1016/j.cca.2018.05.047.
- [27] K. J. Wojno *et al.*, ‘Multiplex PCR Based Urinary Tract Infection (UTI) Analysis Compared to Traditional Urine Culture in Identifying Significant Pathogens in Symptomatic Patients’, *Urology*, vol. 136, pp. 119–126, 2019, doi: 10.1016/j.urology.2019.10.018.
- [28] M. L. Yarbrough, W. Lainhart, A. R. McMullen, N. W. Anderson, and C. D. Burnham, ‘Impact of total laboratory automation on workflow and specimen processing time for culture of urine specimens’, 2018.
- [29] M. B. Jalil, M. Younus, and N. Al, ‘The prevalence of multiple drug resistance *Escherichia coli* and *Klebsiella pneumoniae* isolated from patients with urinary tract infections’, no. June, pp. 1–7, 2022, doi: 10.1002/jcla.24619.
- [30] M. Rizwan, M. Akhtar, A. K. Najmi, K. Singh, D. Res, and N. Delhi, ‘*Escherichia coli* and *Klebsiella pneumoniae* Sensitivity / Resistance Pattern Towards Antimicrobial Agents in Primary and Simple Urinary Tract Infection Patients Visiting University Hospital of Jamia Hamdard New Delhi’, 2018.
- [31] N. N. Maredia, M. J. Fanning, A. L. Christie, B. C. Prokesch, and P. E. Zimmern, ‘Adverse effects of chronic nitrofurantoin therapy in women with recurrent urinary tract infections in an outpatient setting’, *World J. Urol.*, no. 0123456789, 2020, doi: 10.1007/s00345-020-03464-w.
- [32] Y. Madani and B. Mann, ‘Nitrofurantoin-induced lung disease and prophylaxis of urinary tract infections’, *Nat. Publ. Gr.*, vol. 21, no. 3, pp. 337–341, 2012, doi: 10.4104/pcrj.2012.00059.
- [33] M. J. Munoz-davila, ‘Role of Old Antibiotics in the Era of Antibiotic Resistance. Highlighted Nitrofurantoin for the Treatment of Lower Urinary Tract Infections’, no. Figure 1, pp. 39–48, 2014, doi: 10.3390/antibiotics3010039.
- [34] A. K. Sugianli, F. Ginting, I. Parwati, M. D. De Jong, and F. Van Leth, ‘JAC- Antimicrobial Resistance Antimicrobial resistance among uropathogens in the Asia-Pacific region : a systematic review’, pp. 1–12, 2021, doi: 10.1093/jacamr/dlab003.
- [35] M. Gekenidis, A. Kläui, K. Smalla, and D. Drissner, ‘Transferable Extended-Spectrum β -Lactamase (ESBL) Plasmids in Enterobacteriaceae from Irrigation Water’, 2020.
- [36] J. E. Simmering, F. Tang, J. E. Cavanaugh, L. A. Polgreen, and P. M. Polgreen, ‘The Increase in Hospitalizations for Urinary Tract Infections and the Associated Costs in the United States, 1998–2011’, *Open Forum Infect. Dis.*, vol. 4, no. 1, pp. 1–7, 2017, doi: 10.1093/ofid/ofw281.
- [37] M. Pujades-rodriguez, R. M. West, M. H. Wilcox, and J. Sandoe, ‘EClinicalMedicine Lower Urinary Tract Infections : Management , Outcomes and Risk Factors for Antibiotic Re-prescription in Primary Care’, *EClinicalMedicine*, vol. 14, pp. 23–31, 2019, doi: 10.1016/j.eclinm.2019.07.012.
- [38] H. F. Health, K. D. Suson, and H. F. Health, ‘Henry Ford Health Scholarly Commons Re : Luke Harper , T . Blanc , M . Peycelon , et al . Circumcision and Risk of Febrile Urinary Tract Infection in Boys with Posterior Urethral Valves : Result of the CIRCUP Randomized Trial . Eur Urol . Letter to the Ed’, pp. 2021–2022, 2021.

-
- [39] R. Klaus and B. Lange-sperandio, 'Chronic Kidney Disease in Boys with Posterior Urethral Valves – Pathogenesis , Prognosis and Management', 2022.
- [40] L. E. Nicolle, 'Catheter associated urinary tract infections', pp. 1–8, 2014.
- [41] E. Von Elm, D. G. Altman, M. Egger, S. J. Pocock, P. C. Gøtzsche, and J. P. Vandenbroucke, 'The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement : guidelines for reporting observational studies', pp. 1453–1457, 2007.
- [42] A. K. Sugianli *et al.*, 'Laboratory-based versus population-based surveillance of antimicrobial resistance to inform empirical treatment for suspected urinary tract infection in Indonesia', *PLoS One*, vol. 15, no. 3, pp. 1–10, 2020, doi: 10.1371/journal.pone.0230489.