

Article

Not peer-reviewed version

Insecticidal Traits of Variants in a Genotypically Diverse Natural Isolate of *Anticarsia gemmatalis* Multiple Nucleopolyhedrovirus (AgMNPV)

Ana Parras-Jurado , Delia Muñoz , [Inés Beperet](#) , [Trevor Williams](#) , [Primitivo Caballero](#) *

Posted Date: 19 June 2023

doi: 10.20944/preprints202306.1363.v1

Keywords: Baculoviridae; bioinsecticide; genotypic variant; pathogenicity; virulence; occlusion body production



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Insecticidal traits of variants in a genotypically diverse natural isolate of *Anticarsia gemmatalis* multiple nucleopolyhedrovirus (AgMNPV)

Ana Parras-Jurado^{1,2}, Delia Muñoz¹, Inés Beperet², Trevor Williams³ and Primitivo Caballero^{1,2,*}

¹ Institute for Multidisciplinary Research in Applied Biology, Universidad Pública de Navarra, 31006 Pamplona, Navarra, Spain

² Departamento de Investigación y Desarrollo, Bioinsectis SL, Pol. Ind. Mocholi Plaza Cein 5, Nave A14, 31110 Noain, Navarra, Spain

³ Instituto de Ecología AC (INECOL), Xalapa, Veracruz 91073, Mexico

* Correspondence: pcm92@unavarra.es; Tel.: +34-948-168-004

Abstract: Outbreaks of *Anticarsia gemmatalis* (Hübner, 1818) (Lepidoptera: Erebidae), a major pest of soybean, can be controlled below economic thresholds with methods that do not involve the application of synthetic insecticides. Formulations based on natural isolates of the *Anticarsia gemmatalis* multiple nucleopolyhedrovirus (AgMNPV) (*Baculoviridae*: *Alphabaculovirus*), played a significant role in integrated pest management programs in the early 2000s, but a new generation of chemical insecticides and transgenic soybean, have displaced AgMNPV-based products over the past decade. However, the marked genotypic variability present among and within alphabaculovirus isolates suggests that highly insecticidal genotypic variants can be isolated and used to reduce virus production costs or overcome isolate-dependent host resistance. This study aimed to select novel variants of AgMNPV with suitable insecticidal traits that could complement the existing AgMNPV active ingredients. Three distinct AgMNPV isolates were compared by restriction endonuclease profile and in terms of their occlusion body (OB) pathogenicity. One isolate was selected (AgABB51) from which eighteen genotypic variants were plaque purified and characterized in terms of their insecticidal properties. The five most pathogenic variants varied in OB pathogenicity although none of them was faster-killing or had higher OB production characteristics than the wild-type isolate. We conclude that the AgABB51 wild-type isolates appear to be genotypically structured for fast speed-of-kill and high OB production, both of which would favor transmission. Interactions among the component variants are likely to influence this insecticidal phenotype.

Keywords: *Baculoviridae*; bioinsecticide; genotypic variant; pathogenicity; virulence; occlusion body production

1. Introduction

The velvetbean caterpillar, *Anticarsia gemmatalis* (Hübner, 1818) (Lepidoptera: Erebidae), is one of the main species of the soybean pest complex [1,2]. It causes major economic damage to this crop across the Americas [1–4]. The main strategies employed to control *A. gemmatalis* outbreaks rely on the use of synthetic insecticides, a range of selective or biorational products and the cultivation of transgenic plants expressing Bt insecticidal proteins [5–8].

The *Anticarsia gemmatalis* multiple nucleopolyhedrovirus (AgMNPV) (*Baculoviridae*: *Alphabaculovirus*), a natural pathogen of this pest, has played a significant role in integrated pest control programs, and was applied over ~2 million hectares annually in the early 2000s. The high pathogenicity of the virus results in the control of this pest below economic thresholds following a single application of virus occlusion bodies (OBs) to soybean crops [9]. However, the use of AgMNPV-based

insecticides has been drastically reduced over the past decade due to the widespread adoption of transgenic soybean [10–12].

In nature, nucleopolyhedroviruses exist as heterogeneous populations composed of different genotypes [13,14]. Genotypic variants arise from random molecular events involving recombination, insertion, duplication, the deletion of genomic sequences or horizontal gene transfer [15,16]. Some of these variations confer distinct insecticidal properties, usually measured in terms of OB pathogenicity metrics, speed of kill and OB production, which can directly influence virus survival and transmission between susceptible hosts [17–21].

Intraspecific genotypic diversity and phenotypic variation has been described among geographical isolates of a wide array of alphabaculovirus species [19,21–23]. Characterization of the genotypic variants present in a wild-type isolate has proved to be important for the selection and development of some of these variants as the active ingredients of virus-based insecticides [24–26].

Indeed, the prototype AgMNPV-2D, is a majority variant present AgMNPV formulations against *A. gemmatalis* [27], was originally identified by plaque purification of a wild type isolate from Brazil [28,29]. However, from field experiences in other host-virus systems [30–32], access to a range of highly insecticidal genotypic variants is important in case of isolate-dependent host resistance [33–36]. Variants can also differ in their OB production traits, which lend some variants to be more amenable than others to mass-production processes necessary for the commercialization of virus insecticides [37].

In this study, we compared several natural isolates of AgMNPV and characterized the genotypic variants present in the most pathogenic isolate, with the aim of comparing the insecticidal traits of the individual variants with those of commercial products that were previously available against *A. gemmatalis*. Specifically, comparative molecular and biological analyses of three AgMNPV isolates resulted in the selection of one of them, from which eighteen genotypic variants were obtained and characterized in terms of OB pathogenicity, speed-of-kill, and OB productivity.

2. Materials and Methods

2.1. Insects, cells and virus isolates

The *Anticarsia gemmatalis* population was established from larvae collected in soybean fields of Tamaulipas, Mexico and maintained at 25 ± 1 °C, 75% relative humidity and 16 h light: 8 h dark photoperiod on a wheatgerm-based semi-synthetic diet [38].

Sf9 cells (Gibco, Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA) were maintained in Sf-900 II medium (Gibco) at 28 °C.

The *Anticarsia gemmatalis* multiple nucleopolyhedrovirus (AgMNPV) isolates used in this study were AgMNPV-30WT (hereinafter known as Ag30WT) isolated in Mexico [21], and AgMNPV-ABB15 (AgABB15) and AgMNPV-ABB51 (AgABB51), which both originated from the virus collection of the Institut National de la Recherche Agronomique (INRA), France and were kindly provided by Miguel López-Ferber. These INRA isolates were likely to have been deposited during the studies on AgMNPV by Crozier and Ribeiro [16]. OBs were amplified in fourth instars of *A. gemmatalis* and purified by homogenization of each virus-killed larva in 1 mL of 0.1% sodium dodecyl sulfate (SDS) followed by filtration through muslin and centrifugation at 2400 *g* for 5 min to eliminate insect debris. The resulting pellets were washed and resuspended in 2 vol. milli-Q water. Purified OB suspensions were titrated under phase-contrast microscopy at $\times 400$ using an improved hemocytometer (Paul Marienfeld GmbH, Lauda-Königshofen, Germany) and aliquots of each isolate were stored at -20 °C until required.

Unless otherwise stated, all bioassays and related procedures described in the following sections were performed at 25 °C, 75% relative humidity, 16 h light: 8 h dark photoperiod.

2.2. DNA extraction from OBs

Occlusion derived virions (ODVs) were released by incubating 40 μL of OB suspension (10^{10} OBs/mL) with 100 μL of 0.5 M Na_2CO_3 and 360 μL distilled water at 60 °C for 30 min. A supernatant containing the released virions was obtained by centrifugation at 5900 g for 5 min and immediately transferred to a new microcentrifuge tube and incubated with 25 μL 10% SDS, 25 μL 0.5 M EDTA and 15 μL proteinase K (20 mg/mL) at 65 °C for 1 h to degrade the virion membrane and the nucleocapsid and, hence, release the virus genome. Viral DNA was then separated from proteins by adding 150 μL MPC Protein Precipitation Reagent (Epicentre, Illumina Inc., San Diego, California, USA), vortexing vigorously for 10 s, and pelleting the debris by centrifugation at 10000 g and 4 °C for 10 min. The supernatant was transferred to a new microcentrifuge tube and DNA was pelleted by adding 1 mL ice-cold absolute ethanol and centrifuged at 16200 g at 4 °C for 10 min. The pelleted DNA was washed with 500 μL 70% ethanol, and the DNA resuspended in 50 μL bidistilled water and incubated at 60 °C for 15 min.

2.3. Cloning of genotypic variants

Individual genotypes were obtained from AgABB51 by plaque purification [39]. A concentration of 10^8 OB/mL was used to inoculate fifth instar *A. gemmatalis* larvae by the droplet feeding method [40]. At 48 h post infection (hpi), infected larvae were bled and hemolymph containing budded virions was collected and stored at -20 °C. Hemolymph was then passed through a 0.45 μm filter and used to prepare six serial dilutions in Sf-900 II (Gibco) medium with antibiotics. A 200 μL volume from each dilution was used to inoculate 5×10^5 Sf9 cells (Gibco) and, at 10 d post infection (dpi), clearly separated individual plaques containing individual clones were collected with a sterile Pasteur pipette and diluted in 300 μL Sf-900 II medium. Clones were then injected into the hemocoel of *A. gemmatalis* fifth instar larvae that were individually placed in the wells of a cell culture plate with semi-synthetic diet and incubated at 25 °C until death or pupation.

2.4. Viral DNA Restriction Endonuclease Analysis

For restriction endonuclease analysis, 2 μg viral DNA were digested with two units of HindIII (Fast Digest, Thermofisher) for 2 h at 37 °C. DNA fragments were separated by electrophoresis in a 1% agarose gel immersed in TAE buffer (40 mM Tris, 20 mM acetic acid, 1 mM EDTA pH 8.0) running at 18V for 15 h. DNA fragments were stained with GelRed (Biotium, Fremont, CA, USA) and photographed on a transilluminator (Gel Doc EZ Imager, Bio-Rad Inc., Hercules, CA, USA).

2.5. Biological activity

The three isolates employed, Ag30WT, AgABB15 and AgABB51, were used to inoculate groups of 28 newly molted second instar larvae of *A. gemmatalis* at three different concentrations (10^3 , 10^5 and 10^7 OBs/mL) by the droplet feeding method [40]. Larvae that drank the OB suspensions within 10 min were individualized in wells of a cell culture plate with semi-synthetic diet. Control larvae drank a solution that contained no OBs. These assays were performed in triplicate using different insect batches. Mortality was recorded every 24 hours until all larvae were dead or had pupated.

The pathogenicity of AgABB51 OBs, expressed as the median lethal concentration (LC_{50}), was determined by the droplet feeding method [40]. Groups of 28 newly molted second instar larvae were allowed to drink OB suspensions of one of the following concentrations, expected to cause between 10% and 90% mortality: 1.2×10^3 , 3.7×10^3 , 1.1×10^4 , 3.3×10^4 and 1.0×10^5 OBs/mL. Control larvae drank a solution that contained no OBs. Larvae that drank the OB suspensions within 10 min were individualized in wells of a cell culture plate with semi-synthetic diet and mortality was recorded at 24-h intervals until all larvae had died or pupated. This assay was performed in triplicate with different insect batches. Concentration-mortality data were analyzed by Probit regression to estimate LC_{50} values using the software POLO Plus (Leora) [41].

Median time to death (MTD) was estimated in AgABB51-infected second instars that consumed a suspension of 6.8×10^4 OBs/mL (estimated to result in ~90% mortality). Larvae that drank the inoculum within 10 min were individualized with semi-synthetic diet and mortality was recorded every

8 h until all larvae had died or pupated. Control larvae consumed a solution without OBs. The study was performed on three batches of insects. In order to estimate MTD values, a survival analysis was performed using the 'Survival' package [42] in R (v4.2.2) [43]. The Akaike Information Criterion (AIC) was calculated in order to identify the best fitting model.

OB production was determined in fifth instars inoculated with an LC₉₉ (10⁸ OBs/mL) concentration of AgABB51 OBs by the droplet feeding method [40]. In each of the three biological replicates, dead larvae were collected individually in a 1.5 mL microcentrifuge tube and OBs were purified in a total volume of 1 mL with milli-Q water. OB suspensions from 26 larvae from each replicate were titered in a Neubauer hemocytometer. Data were subjected to a Shapiro-Wilk normality test and analysis of variance (ANOVA) followed by post-hoc Tukey's HSD test in R (v4.2.2) [43].

An inoculum concentration equal to the LC₅₀ of AgABB51 OBs (1.1 × 10⁴ OBs/mL) was used to evaluate the mortality response of *A. gemmatalis* second instar larvae to each of the genotypic variants isolated in section 2.3. Larvae were inoculated by the droplet feeding method [40] and incubated individually on semi-synthetic diet in the wells of a cell culture plate. Mortality was registered daily until all individuals were dead or pupated. The bioassay was performed on three batches of insects with appropriate controls. Percentage of mortality values were analyzed by Kruskal-Wallis test in R (v4.2.2) [43].

2.6. Genotypic variant selection and biological characterization

The genotypic variants that induced higher mortality responses than the wild-type isolate (AgABB51) were selected for further characterization in terms of LC₅₀, MTD and OB production as described in section 2.5., whereas MTD values were subjected to a log-rank test using the package 'Survival' [42] and a post hoc Bonferroni-corrected pairwise *t*-test. OB production data were subjected to a Shapiro-Wilk normality test and one-way ANOVA followed by a post-hoc Tukey HSD test. All analyses were performed using R (v4.2.2) [43].

3. Results

3.1. Identification of AgMNPV isolates by restriction endonuclease analysis

Analysis of the genomic DNA of three different AgMNPV isolates (Ag30WT, AgABB15 and AgABB51) revealed three slightly different REN profiles, as judged by the distinct restriction fragment length polymorphisms (Figure 1). All the isolates appeared to generate restriction fragments in submolar concentrations, indicating the presence of different genotypic variants within each isolate (Figure 1).




Figure 1. Restriction endonuclease profiles of the genomic DNA of three different AgMNPV isolates following treatment with HindIII. m denotes the molecular marker. Fragment size in kilobases (Kb) is shown on the left. Red arrowheads indicate restriction fragment length polymorphisms (RFLPs) and asterisks on the left of each lane indicate the presence of submolar bands.

3.2. Mortality response to AgMNPV isolates

Similar mortality values were observed at the lowest and highest viral concentrations used for the three candidate isolates, but isolate AgABB51 resulted in higher larval mortality at 10^5 OBs/mL (Table 1). Thus, we selected AgABB51 for further characterization.

Table 1. Percentage of mortality (\pm SE) observed for each AgMNPV isolate in *A. gemmatalis* second instars. Values in parentheses indicate the numbers of larvae tested at each inoculum concentration.

Virus	10^3 OBs/mL	10^5 OBs/mL	10^7 OBs/mL
Ag30WT	8 ± 4 (78)	58 ± 8 (73)	98 ± 2 (83)
AgABB15	11 ± 5 (78)	72 ± 8 (81)	100 ± 0 (79)
AgABB51	10 ± 6 (82)	87 ± 1 (77)	100 ± 0 (74)

None of the control insects ($n = 72$ -78 larvae per isolate) died of polyhedrosis disease

3.3. Biological and genotypic characterization of AgABB51

The pathogenicity of AgABB51 OBs, in terms of the LC_{50} , was estimated at 1.1×10^4 OBs/mL. A median time to death value of 138.6 hours post inoculum (hpi) was estimated through a survival analysis using the log-normal model, which was identified as the best fitting model by comparison of distribution-dependent AIC values. The mean OB yield was 1.23×10^9 OBs in *A. gemmatalis* fifth instars (Table 2).

Table 2. Estimated median lethal concentration (LC₅₀) and median time to death (MTD) values for AgABB51 in *A. gemmatalis* second instars and OB production values for AgABB51 in *A. gemmatalis* fifth instars.

Virus	LC ₅₀ * (OBs/mL)	95% Confidence Limits		MTD (h)	95% Confidence Limits		OB Production (OBs/larva)	95% Confid
		Low	High		Low	High		
AgABB51	1.1 × 10 ⁴	8.5 × 10 ³	1.5 × 10 ⁴	138.6	134.3	143.0	1.23 × 10 ⁹	1.15 × 10 ⁹

*LC₅₀ estimated by Probit regression with a slope (± SE) 1.634 ± 0.148 and intercept (± SE) -6.612 ± 0.612 (goodness-of-fit test $\chi^2 = 0.2198$, d.f. = 3, p = 0.974, heterogeneity = 0.0733).

A total of 128 different clones were obtained from the plaque assay and were each amplified by injection into *A. gemmatalis* fifth instar larvae. OBs from virus-killed larvae were collected and the viral DNA digested with HindIII to determine differences in their genomic profile. Eighteen different genotypic variants were obtained and named A to R (Figure 2).




Figure 2. Restriction endonuclease profiles of the genomic DNA of the 18 different AgABB51 genotypic variants obtained with HindIII. m denotes the molecular marker. Fragment size in kilobases (Kb) is shown on the left. The presence (red arrowheads) or absence (green arrowheads) of characteristic restriction fragments are indicated compared to genotype A, as this was the most similar profile to that of the wild-type isolate (AgABB51).

The frequency of each variant was determined as the number of times each characteristic restriction profile was observed in the 128 clones (Figure 3).




Figure 3. Frequency of AgABB51 genotypic variants A through R in terms of number of clones comprising a particular variant profile from the total of 128 clones analyzed.

3.4. Biological characterization of AgABB51 genotypic variants

The mean percentage of mortality in larvae inoculated with each of the different genotypic variants and the wild-type AgABB51 varied significantly ($\chi^2 = 46.99$, $df = 18$, $p < 0.005$). However, only larvae inoculated with five genotypic variant OBs (D, E, I, L, M) showed higher mortalities than larvae inoculated with AgABB51 OBs in the initial bioassays, at the concentration equal to the LC_{50} of AgABB51 (1.1×10^4 OBs/mL). All five of these variants were selected for further characterization (Table 3). Of these, variants E and I caused the highest mean (\pm SE) larval mortalities of $61.0 \pm 7.4\%$ and $61.8 \pm 1.1\%$, respectively.




Figure 4. Mean percentage of mortality caused by AgABB51 wild-type isolate and each of the genotypic variants A-R on *A. gemmatalis* second instars inoculated with 1.1×10^4 OBs/mL. Error bars indicate the standard error.

The pathogenicity of AgABB51 and the five selected genotypic variants ranged from 1×10^4 OBs/mL (AgABB51) to 7.6×10^3 OBs/mL (variant I), there were no significant differences detected among these values (Table 3).

Table 3. Lethal concentration estimates and relative potency values for AgABB51 and selected individual genotypic variants in *A. gemmatalis* second instars.

Variant	LC ₅₀ (OBs/mL)	Relative potency	95% Confidence Limits		LC ₉₀ (OBs/mL)	Relative potency	95% Confidence Limits		Slop
			Low	High			Low	High	
AgABB51	1.0×10^4	1	-	-	6.9×10^4	1	-	-	1.54
D	8.8×10^3	1.15	0.84	1.56	5.3×10^4	1.30	0.77	2.11	1.64
E	8.0×10^3	1.26	0.92	1.72	4.4×10^4	1.55	0.91	2.65	1.72
I	7.6×10^3	1.33	0.97	1.81	4.1×10^4	1.68	1.00	2.9	1.75
L	9.2×10^3	1.10	0.80	1.50	5.6×10^4	1.21	0.70	2.13	1.62
M	9.2×10^3	1.10	0.74	1.36	4.9×10^4	1.40	0.83	2.35	1.65

Data did not differ significantly as hypotheses for parallelism ($\chi^2 = 2.67$, $df = 5$, $p > 0.05$) and equality ($\chi^2 = 7.83$, $df = 10$, $p > 0.05$) were not rejected. The relative potencies were calculated as the ratio of effective concentrations relative to the wild-type AgABB51.

Survival analysis using the log-normal model revealed MTD values that varied between 155.0 h (variant E) and 138.9 h (variant M), compared to 139.0 h for the reference AgABB51 isolate (log-rank $\chi^2 = 16.3$, $d.f. = 5$, $p < 0.05$). However, a post hoc Bonferroni analysis revealed that only the MTD values of AgABB51 and variant M differed significantly compared to variant E, whereas the other variants showed intermediate MTD values (Figure 4a).

In terms of OB production, none of the selected genotypic variants was more productive than the AgABB51 isolate and larvae infected by the variants D, E, I and L produced fewer OBs than those infected by the AgABB51 isolate ($F_{5,121} = 8.533$; $p < 0.001$) (Figure 4b).




Figure 5. (a) Median time to death (MTD) values for the AgABB51 isolate and selected genotypic variants in *A. gemmatalis* second instars. Error bars indicate standard error and different lowercase letters indicate significant differences between variants (Bonferroni-adjusted t-test; $p < 0.05$). (b) OB production values obtained for AgABB51 and the selected genotypic variants in *A. gemmatalis* fifth instars. Error bars indicate standard error and different lowercase letters indicate significant differences between variants (Tukey HSD test; $p < 0.05$).

4. Discussion

Three field-collected AgMNPV isolates were compared in terms of restriction endonuclease profile characteristics and their respective insecticidal properties. AgABB51 was recognized as the most genotypically diverse, as judged by the greater number of RFLPs and submolar fragments within its restriction profile, suggesting the presence of more than one genotypic variant in the wild-type isolate. The *A. gemmatalis* mortality response induced by these isolates was tested at three different concentrations. Albeit similar mortality values were observed at the lowest and highest viral concentrations, AgABB51 resulted in the highest mortality response (87%) at an intermediate inoculum concentration (Table 1). AgABB51 was therefore selected for further biological and genotypic characterization. The OB pathogenicity (LC_{50}) and speed-of-kill (MTD) values of this isolate were 1.1×10^4 OBs/mL and 138.6 hours in second instars, respectively, whereas OB production averaged 1.23×10^9 OBs in each virus-killed fifth instar (Table 2). Although the AgMNPV-2D variant was not included in our bioassays and comparisons of the results of different experimental events tend to be difficult, the AgABB51 LC_{50} and MTD values seemed lower than those of AgMNPV-2D, suggesting that AgABB51 might possess traits that would favor its use as a biological insecticide [21,27,44,45].

A total of 128 clones were obtained by plaque purification of the AgABB51 isolate. From these, 18 different genotypic variants (AgABB51-A to R) were identified based on their HindIII restriction profiles (Figure 2). A similar heterogeneity has been reported in the population structure of other alphabaculoviruses [20,46,47], including other AgMNPV isolates [16,21,29,48].

Minor variation in the genome sequence can have a direct impact on the insecticidal phenotype of genotypic variants [37,49,50]. In line with this idea, 13 of the AgABB51 variants caused a mortality similar to or lower than that of AgABB51-wt, whereas five variants (D, E, I, L and M) caused mortality slightly higher (56.8 - 61.8%) than the wild-type isolate.

There was no clear correlation between the frequency of the variant in the isolate (Figure 3) and virus-induced mortality (Figure 4), as previously observed by other authors [20,21,50,51]. One reason may be that variants that are amenable to cell culture conditions are not necessarily the most pathogenic or transmissible variants in nature [22,51–53]. The alternative *in vivo* cloning method was originally developed to purify variants from alphabaculovirus isolates [54,55], but is a highly labor-intensive method that has largely been abandoned in favor of plaque purification [20].

The five selected variants from AgABB51 were not significantly more pathogenic than the AgABB51-wt (Table 3). None of these variants displayed a faster speed-of-kill or higher OB production than AgABB51-wt (Figure 4). Indeed, for most of these variants slower speed of kill was associated with lower OB production per larva, which deviates from the finding that speed of kill is often correlated with higher OB production, presumably because the virus has more time to replicate and the insect can continue to grow during the infection period [56–58]. It appears therefore, that the AgABB51 isolate is genotypically structured so that the speed-of-kill favors rapid insect-to-insect transmission in combination with high OB production which also increases the probability of transmission. These emergent traits likely arise from interactions among the component genotypes, a phenomenon also observed in experiments involving the production of variant mixtures in other alphabaculoviruses [59].

On a more general note, the study of the highly diverse AgABB51 population also highlights the importance of collective infectious units in virus transmission [13,60,61]. The polyploid nature of viruses that disperse in groups, such as alphabaculoviruses, has important consequences for viral evolution, as it increases the probability of coinfection and recombination and complementation among coinfecting variants. Cells infected with multiple virus genomes will favor interactions between viruses, that may result in changes in viral pathogenesis, the diversity of the virus progeny and the evolution of host resistance [61].

Author Contributions: Conceptualization, P.C.; formal analysis, A.P.; funding acquisition, P.C.; investigation, A.P.; resources, T.W.; methodology, A.P., I.B. and P.C.; writing - original draft preparation, A.P.; writing - review and editing, A.P., D.M., T.W. and P.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Ministerio de Ciencia e Innovación, grant number AGL2017-83498-C2-1-R. A.P. received a predoctoral grant PRE2018-086829 from the Agencia Estatal de Investigación (Ministerio de Ciencia e Innovación) of the Spanish Government.

Acknowledgments: We thank Miguel López-Ferber for providing two of the AgMNPV isolates and Noelia Gorriá and Itxaso Ibáñez for insect rearing. J. Sebastián Gómez Díaz (INECOL) provided logistical support to T.W.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Bueno, R.C.O.; Bueno, A. de F.; Moscardi, F.; Postali Parra, J.R.; Hoffmann-Campo, C.B. Lepidopteran Larva Consumption of Soybean Foliage: Basis for Developing Multiple-Species Economic Thresholds for Pest Management Decisions. *Pest Management Science* **2011**, *67*, 170–174, doi:10.1002/ps.2047.
2. Moscardi, F.; Carvalho, R.C.Z. de Consumo e utilização de folhas da soja por *Anticarsia gemmatalis* Hübn. (Lepidoptera: Noctuidae) infectada, em diferentes estádios larvais, por seu vírus de poliedrose nuclear. *Anais da Sociedade Entomológica do Brasil* **1993**, *22*, 267–280, doi:10.37486/0301-8059.v22i2.847.
3. Bayer Crop Science Bayer - Crop Pests Compendium - *Anticarsia Gemmatalis* by Bayer Crop Science AG Available online: <https://www.agriculture-xprt.com/products/bayer-anticarsia-gemmatalis-617566> (accessed on 24 May 2023).
4. Musser, F.R.; Catchot, A.L.; Conley, S.P.; Davis, J.A.; Difonzo, C.; Greene, J.K.; Lorenz, G.M.; Owens, D.; Reisig, D.D.; Roberts, P.; et al. 2019 Soybean Insect Losses in the United States. *Midsouth Entomologist* **2020**, *13*, 1–23.

5. Fernandes, F.O.; Abreu, J.Á. de; Christ, L.M.; Rosa, A.P.S.A. da Insecticides Management Used in Soybean for the Control of *Anticarsia gemmatalis* (Hübner, 1818) (Lepidoptera: Eribyidae). *Journal of Agricultural Science* **2018**, *10*, 223, doi:10.5539/jas.v10n11p223.
6. Degrande, P.E.; Vivian, L.M. Pragas Da Soja. *Tecnologia e Produção: Soja e Milho 2011/2012* **2012**, 155–206.
7. Murúa, M.G.; Vera, M.A.; Herrero, M.I.; Fogliata, S.V.; Michel, A. Defoliation of Soybean Expressing Cry1Ac by Lepidopteran Pests. *Insects* **2018**, *9*, 4–8, doi:10.3390/insects9030093.
8. Bueno, R.C.O.F.; Raetano, C.G.; Junior, J.D.; Carvalho, F.K. Integrated Management of Soybean Pests: The Example of Brazil. *Outlooks on Pest Management* **2017**, *28*, 149–153, doi:10.1564/v28_aug_02.
9. Moscardi, F. Assessment of the Application of Baculoviruses for Control of Lepidoptera. *Annual Review of Entomology* **1999**, *44*, 257–289, doi:10.1146/annurev.ento.44.1.257.
10. Sosa-Gómez, D.R. Microbial Control of Soybean Pest Insects and Mites. In *Microbial Control of Insect and Mite Pests: From Theory to Practice*; Elsevier Inc.: Londrina, Paraná, Brazil, **2017**; pp. 199–208 ISBN 978-0-12-803566-5.
11. Moscardi, F.; de Souza, M.L.; de Castro, M.E.B.; Lara Moscardi, M.; Szewczyk, B. Baculovirus Pesticides: Present State and Future Perspectives. In *Microbes and Microbial Technology*; Ahmad, I., Ahmad, F., Pichtel, J., Eds.; Springer: New York, **2011**; pp. 415–445 ISBN 978-1-4419-7930-8.
12. Valicente, F.H. Entomopathogenic Viruses. In *Natural Enemies of Insect Pests in Neotropical Agroecosystems*; Souza, B., Vázquez, L.L., Marucci, R.C., Eds.; Springer Nature Switzerland AG: Switzerland, **2019**; pp. 137–150 ISBN 978-3-030-24733-1.
13. Clavijo, G.; Williams, T.; Muñoz, D.; Caballero, P.; López-Ferber, M. Mixed Genotype Transmission Bodies and Virions Contribute to the Maintenance of Diversity in an Insect Virus. *Proc Biol Sci* **2010**, *277*, 943–951, doi:10.1098/rspb.2009.1838.
14. Erlandson, M.A. Genetic Variation in Field Populations of Baculoviruses: Mechanisms for Generating Variation and Its Potential Role in Baculovirus Epizootiology. *Virol. Sin.* **2009**, *24*, 458–469, doi:10.1007/s12250-009-3052-1.
15. de Brito, A.F.; Braconi, C.T.; Weidmann, M.; Dilcher, M.; Pereira Alves, J.M.; Gruber, A.; de Andrade Zannotto, P.M. The Pangenome of the *Anticarsia gemmatalis* multiple nucleopolyhedrovirus (AgMNPV). *Genome Biology and Evolution* **2016**, *8*, 94–108, doi:10.1093/gbe/evv231.
16. Croizier, G.; Ribeiro, H.C.T. Recombination as a Possible Major Cause of Genetic Heterogeneity in *Anticarsia gemmatalis* nuclear polyhedrosis virus Wild Populations. *Virus Research* **1992**, *26*, 183–196, doi:http://dx.doi.org/10.1016/0168-1702(92)90012-X.
17. Cabodevilla, O.; Ibañez, I.; Simón, O.; Murillo, R.; Caballero, P.; Williams, T. Occlusion Body Pathogenicity, Virulence and Productivity Traits Vary with Transmission Strategy in a Nucleopolyhedrovirus. *Biological Control* **2011**, *56*, 184–192, doi:10.1016/j.biocontrol.2010.10.007.
18. Harrison, R.L.; Puttler, B.; Popham, H.J.R. Genomic Sequence Analysis of a Fast-Killing Isolate of *Spodoptera frugiperda* multiple nucleopolyhedrovirus. *Journal of General Virology* **2008**, *89*, 775–790, doi:10.1099/vir.0.83566-0.
19. Rowley, D.L.; Popham, H.J.R.; Harrison, R.L. Genetic Variation and Virulence of Nucleopolyhedroviruses Isolated Worldwide from the Heliothine Pests *Helicoverpa armigera*, *Helicoverpa zea*, and *Heliothis virescens*. *Journal of Invertebrate Pathology* **2011**, *107*, 112–126, doi:10.1016/j.jip.2011.03.007.
20. Aguirre, E.; Beperet, I.; Williams, T.; Caballero, P. Genetic Variability of *Chrysodeixis includens* nucleopolyhedrovirus (ChinNPV) and the Insecticidal Characteristics of Selected Genotypic Variants. *Viruses* **2019**, *11*, 581, doi:10.3390/v11070581.
21. Del-Angel, C.; Lasa, R.; Rodríguez-del-Bosque, L.A.; Mercado, G.; Beperet, I.; Caballero, P.; Williams, T. *Anticarsia gemmatalis* nucleopolyhedrovirus from Soybean Crops in Tamaulipas, Mexico: Diversity and Insecticidal Characteristics of Individual Variants and Their Co-Occluded Mixtures. *Florida Entomologist* **2018**, *101*, 404–410, doi:10.1653/024.101.0319.
22. Cory, J.S.; Green, B.M.; Paul, R.K.; Hunter-Fujita, F. Genotypic and Phenotypic Diversity of a Baculovirus Population within an Individual Insect Host. *Journal of Invertebrate Pathology* **2005**, *89*, 101–111, doi:10.1016/j.jip.2005.03.008.
23. Lee, H.H.; Miller, L.K. Isolation of Genotypic Variants of *Autographa californica* nuclear polyhedrosis virus. *J Virol* **1978**, *27*, 754–767, doi:10.1128/JVI.27.3.754-767.1978.
24. Arrizubieta, M.; Simón, O.; Caballero, P.; Williams, T. Nuevos genotipos del nucleopoliedrovirus simple de *Helicoverpa armigera* (HearSNPV), procedimiento para su producción y uso como agente de control biológico. Patent ES2555165, 29 December 2015.
25. Caballero, P.; Bernal, A.; Simón, O.; Carnero, A.; Hernández-Suárez, E.; Williams, T.G. Nuevos genotipos del nucleopoliedrovirus simple de *Chrysodeixis chalcites* (ChchSNPV), procedimiento para su producción y uso como agente de control biológico. Patent ES2504866, 8 October 2014.
26. Caballero, P.; Murillo, R.; Muñoz, D.; Williams, T. El Nucleopoliedrovirus de *Spodoptera exigua* (Lepidoptera: Noctuidae) Como Bioplaguicida: Análisis de Avances Recientes En España. *Revista Colombiana de Entomología* **2009**, *35*, 105–115.

27. Oliveira, J.V.; Wolff, J.L.C.; Garcia-Maruniak, A.; Ribeiro, B.M.; de Castro, M.E.B.; de Souza, M.L.; Moscardi, F.; Maruniak, J.E.; de Andrade Zanotto, P.M. Genome of the Most Widely Used Viral Biopesticide: *Anticarsia gemmatalis* multiple nucleopolyhedrovirus. *Journal of General Virology* **2006**, *87*, 3233–3250, doi:10.1099/vir.0.82161-0.
28. Allen, G.E.; Knell, J.D. A Nuclear polyhedrosis virus of *Anticarsia gemmatalis*: I. Ultrastructure, Replication, and Pathogenicity. *The Florida Entomologist* **1977**, *60*, 233, doi:10.2307/3493914.
29. Maruniak, J.E.; Garcia-Maruniak, A.; Souza, M.L.; Zanotto, P.M.A.; Moscardi, F. Physical Maps and Virulence of *Anticarsia gemmatalis* nucleopolyhedrovirus Genomic Variants. *Archives of Virology* **1999**, *144*, 1991–2006, doi:10.1007/s007050050720.
30. Fan, J.; Jehle, J.A.; Rucker, A.; Nielsen, A.L. First Evidence of CpGV Resistance of Codling Moth in the USA. *Insects* **2022**, *13*, 533, doi:10.3390/insects13060533.
31. Jehle, J.A.; Schulze-Bopp, S.; Undorf-Spahn, K.; Fritsch, E. Evidence for a Second Type of Resistance against *Cydia pomonella* granulovirus in Field Populations of Codling Moths. *Applied and Environmental Microbiology* **2016**, *83*, e02330-16, doi:10.1128/AEM.02330-16.
32. Hoover, K.; Washburn, J.O.; Volkman, L.E. Midgut-Based Resistance of *Heliothis virescens* to Baculovirus Infection Mediated by Phytochemicals in Cotton. *Journal of Insect Physiology* **2000**, *46*, 999–1007, doi:10.1016/S0022-1910(99)00211-5.
33. Eberle, K.E.; Asser-Kaiser, S.; Sayed, S.M.; Nguyen, H.T.; Jehle, J.A. Overcoming the Resistance of Codling Moth against Conventional *Cydia pomonella* granulovirus (CpGV-M) by a New Isolate CpGV-I12. *Journal of Invertebrate Pathology* **2008**, *98*, 293–298, doi:10.1016/j.jip.2008.03.003.
34. Gebhardt, M.M.; Eberle, K.E.; Radtke, P.; Jehle, J.A. Baculovirus Resistance in Codling Moth Is Virus Isolate-Dependent and the Consequence of a Mutation in Viral Gene Pe38. *Proc Natl Acad Sci U S A* **2014**, *111*, 15711–15716, doi:10.1073/pnas.1411089111.
35. Abot, A.R.; Moscardi, F.; Fuxa, J.R.; Sosa-Gómez, D.R.; Richter, A.R. Development of Resistance By *Anticarsia gemmatalis* from Brazil and the United States to a Nuclear Polyhedrosis Virus under Laboratory Selection Pressure. *Biological Control* **1996**, *7*, 126–130, doi:10.1006/bcon.1996.0075.
36. Levy, S.M.; Falleiros, A.M.F.; Moscardi, F.; Gregório, E.A. Susceptibility/Resistance of *Anticarsia gemmatalis* Larvae to Its nucleopolyhedrovirus (AgMNPV): Structural Study of the Peritrophic Membrane. *J Invertebr Pathol* **2007**, *96*, 183–186, doi:10.1016/j.jip.2007.04.007.
37. Barrera, G.; Williams, T.; Villamizar, L.; Caballero, P.; Simón, O. Deletion Genotypes Reduce Occlusion Body Potency but Increase Occlusion Body Production in a Colombian Spodoptera frugiperda nucleopolyhedrovirus Population. *PLOS ONE* **2013**, *8*, e77271, doi:10.1371/journal.pone.0077271.
38. Greene, G.L.; Leppla, N.C.; Dickerson, W.A. Velvetbean Caterpillar: A Rearing Procedure and Artificial Medium123. *Journal of Economic Entomology* **1976**, *69*, 487–488, doi:10.1093/jee/69.4.487.
39. King, L.A.; Possee, R.D. *The Baculovirus Expression System. A Laboratory Guide*; Chapman & Hall: London, UK, **1992**; ISBN 0-412-37150-2.
40. Hughes, P.R.; Wood, H.A. In Vivo and in Vitro Bioassay Methods for Baculoviruses. In *The Biology of Baculoviruses*; Granados, R.R., Federici, B.A., Eds.; CRC Press: Boca Raton, FL, USA, **1986**; Vol. 2, pp. 1–30.
41. *Polo Plus*, version 1.0; LeOra Software: Parma, MO, USA, 2003.
42. Therneau, T.M. *A Package for Survival Analysis in R*; 2023; <https://CRAN.Rproject.org/package=survival>
43. R Core Team R: *A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2022.
44. Ferreira, B.C.; Melo, F.L.; Silva, A.M.R.; Sanches, M.M.; Moscardi, F.; Ribeiro, B.M.; Souza, M.L. Biological and Molecular Characterization of Two *Anticarsia gemmatalis* multiple nucleopolyhedrovirus Clones Exhibiting Contrasting Virulence. *Journal of Invertebrate Pathology* **2019**, *164*, 23–31, doi:10.1016/j.jip.2019.03.008.
45. da Silveira, E.B.; Cordeiro, B.A.; Ribeiro, B.M.; de Castro, M.E.B.; Soares, E.F.; Bão, S.N. An *Anticarsia gemmatalis* multiple nucleopolyhedrovirus Mutant, VApAg, Induces Hemocytes Apoptosis in Vivo and Displays Reduced Infectivity in Larvae of *Anticarsia gemmatalis* (Hübner) (Lepidoptera: Noctuidae). *Virus Research* **2007**, *130*, 182–192, doi:10.1016/j.virusres.2007.06.010.
46. Corsaro, B.G.; Fraser, M.J. Characterization of Genotypic and Phenotypic Variation in Plaque-Purified Strains of HzSNPV Elkar Isolate. *Intervirology* **1987**, *28*, 185–198, doi:10.1159/000150016.
47. Hodgson, D.J.; Vanbergen, A.J.; Watt, A.D.; Hails, R.S.; Cory, J.S. Phenotypic Variation between Naturally Co-Existing Genotypes of a Lepidopteran Baculovirus. *Evolutionary Ecology Research* **2001**, *3*, 687–701.
48. Ribeiro, H.C.T.; Pavan, O.H.O.; Muotri, A.R. Comparative Susceptibility of Two Different Hosts to Genotypic Variants of the *Anticarsia gemmatalis* nuclear polyhedrosis virus. *Entomologia Experimentalis et Applicata* **1997**, *83*, 233–237, doi:10.1046/j.1570-7458.1997.00177.x.
49. Ogembo, J.G.; Chaeychomsri, S.; Kamiya, K.; Ishikawa, H.; Katou, Y.; Ikeda, M.; Kobayashi, M. Cloning and Comparative Characterization of Nucleopolyhedroviruses Isolated from African Bollworm, *Helicoverpa armigera*, (Lepidoptera: Noctuidae) in Different Geographic Regions. *Journal of Insect Biotechnology and Sericology* **2007**, *76*, 1_39-1_49, doi:10.11416/jibs.76.1_39.

-
50. Bernal, A.; Simón, O.; Williams, T.; Muñoz, D.; Caballero, P. A Chrysodeixis chalcites single-nucleocapsid nucleopolyhedrovirus Population from the Canary Islands Is Genotypically Structured To Maximize Survival. *Applied and Environmental Microbiology* **2013**, *79*, 7709–7718, doi:10.1128/AEM.02409-13.
 51. Simon, O.; Williams, T.; Lopez-Ferber, M.; Caballero, P. Genetic Structure of a Spodoptera frugiperda nucleopolyhedrovirus Population: High Prevalence of Deletion Genotypes. *Applied and Environmental Microbiology* **2004**, *70*, 5579–5588, doi:10.1128/AEM.70.9.5579–5588.2004.
 52. Pijlman, G.P.; van den Born, E.; Martens, D.E.; Vlak, J.M. Autographa californica baculoviruses with Large Genomic Deletions Are Rapidly Generated in Infected Insect Cells. *Virology* **2001**, *283*, 132–138, doi:10.1006/viro.2001.0854.
 53. Kool, M.; Voncken, J.W.; van Lier, F.L.; Tramper, J.; Vlak, J.M. Detection and Analysis of Autographa californica nuclear polyhedrosis virus Mutants with Defective Interfering Properties. *Virology* **1991**, *183*, 739–746, doi:10.1016/0042-6822(91)91003-y.
 54. Smith, I.R.L.; Crook, N.E. In Vivo Isolation of Baculovirus Genotypes. *Virology* **1988**, *166*, 240–244, doi:10.1016/0042-6822(88)90165-1.
 55. Muñoz, D.; Castillejo, J.I.; Caballero, P. Naturally Occurring Deletion Mutants Are Parasitic Genotypes in a Wild-Type Nucleopolyhedrovirus Population of *Spodoptera exigua*. *Applied and Environmental Microbiology* **1998**, *64*, 4372–4377, doi:10.1128/AEM.64.11.4372-4377.1998.
 56. Kumar, C.; Sathiah, N.; Rabindra, R. Optimizing the Time of Harvest of Nucleopolyhedrovirus Infected *Spodoptera litura* (Fabricius) Larvae under in Vivo Production Systems. *Current Science* **2005**, *88*.
 57. Cory, J.S.; Myers, J.H. Adaptation in an Insect Host–Plant Pathogen Interaction. *Ecology Letters* **2004**, *7*, 632–639, doi:10.1111/j.1461-0248.2004.00617.x.
 58. Simón, O.; Williams, T.; López-Ferber, M.; Taulemesse, J.-M.; Caballero, P. Population Genetic Structure Determines Speed of Kill and Occlusion Body Production in *Spodoptera frugiperda* multiple nucleopolyhedrovirus. *Biological Control* **2008**, *44*, 321–330, doi:10.1016/j.biocontrol.2007.12.005.
 59. Williams, T.; López-Ferber, M.; Caballero, P. Nucleopolyhedrovirus Cocclusion Technology: A New Concept in the Development of Biological Insecticides. *Frontiers in Microbiology* **2022**, *12*, doi:10.3389/fmicb.2021.810026.
 60. Sanjuán, R. Collective Infectious Units in Viruses. *Trends Microbiol* **2017**, *25*, 402–412, doi:10.1016/j.tim.2017.02.003.
 61. Leeks, A.; Sanjuán, R.; West, S.A. The Evolution of Collective Infectious Units in Viruses. *Virus Res* **2019**, *265*, 94–101, doi:10.1016/j.virusres.2019.03.013.