

A new *Pseudomonas aeruginosa* isolate enhances its unusual 1,3-propanediol generation from glycerol in bioelectrochemical system

Julia Pereira Narcizo^a, Lucca Bonjy Kikuti Mancilio^{b,c}, Matheus Pedrino^b, María-Eugenia Guazzaroni^{b*}, Adalgisa Rodrigues de Andrade^a, Valeria Reginatto^a

^aDepartment of Chemistry, Faculty of Philosophy, Sciences and Letters of Ribeirão Preto (FFCLRP) - University of São Paulo (USP) - Ribeirão Preto, SP, Brazil

^bDepartment of Biology, Faculty of Philosophy, Sciences and Letters of Ribeirão Preto (FFCLRP) - University of São Paulo (USP) - Ribeirão Preto, SP, Brazil.

^cDepartment of Microbiology, University of Massachusetts – Amherst, Massachusetts, USA.

*Correspondence to: María-Eugenia Guazzaroni, meguazzaroni@ffclrp.usp.br

Supplementary Materials




Figure S1. Taxonomic affiliation of strain EL14 inferred with TYGS (FastME 2.1.6.1) (Lefor and Gascuel, 2015) from GBDP (Genome Blast Distance Phylogeny) distances calculated from strain EL14 genome. The branch lengths are scaled in terms of GBDP distance formula d5. The numbers above branches are GBDP pseudo-bootstrap support values > 60 % from 100 replications, with an average branch support of 95.5 %. The tree was rooted at the midpoint.