

Communication

Not peer-reviewed version

A New Fluorescent Probe for Hydrogen Sulfide Detection in Solution and Living Cells

[Wei Feng](#)^{*}, [Qicai Xiao](#), Lu Wang, [Yuanyong Yang](#)^{*}

Posted Date: 4 July 2023

doi: 10.20944/preprints202307.0213.v1

Keywords: Hydrogen Sulfide; Fluorescent Probe; H₂S Selective Detection; H₂S Quantification; Az-ide-Reduction



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Communication

A New Fluorescent Probe for Hydrogen Sulfide Detection in Solution and Living Cells

Wei Feng ^{2,t,*}, Qicai Xiao ^{3,†} Lu Wang ⁴ and Yuanyong Yang ^{1,*}

¹ School of Pharmacy, Guizhou Medical University; Yangyuanyong@gmc.edu.cn

² BGI - Shenzhen, China; fengweinku@sina.com

³ School of Pharmaceutical Sciences (Shenzhen), Sun Yat-sen University; xiaoqicai@mail.sysu.edu.cn

⁴ Department of Physiology, College of Basic Medicine, Guizhou Medical University; 2899374290@qq.com

* Correspondence: fengweinku@sina.com (W.F.); Yangyuanyong@gmc.edu.cn (Y.Y.)

Abstract: Since Hydrogen Sulfide (H₂S) was recognized as a gas transmitter, its detection and quantification have become a hot research topic among chemists and biologists. In this area, fluorescent probes have shown great advantages: fast and strong responds, low detection limit and easy manipulation. Here we developed a new fluorescent probe that detected H₂S selectively among various bioactive and inorganic salts. This probe was based on the core structure of fluorescein and reacted with H₂S through azide-reduction. Great linearity was achieved correlating fluorescence intensity and H₂S concentrations in solution. The detection of H₂S in cancer cells was also achieved.

Keywords: hydrogen sulfide; fluorescent probe; H₂S selective detection; H₂S quantification; azide-reduction

1. Introduction


The introduction Since recognized an important cell-signaling gasotransmitter [1,2], hydrogen sulfide (H₂S) has attracted great interest from research area of both biology [3,4] and chemistry [5,6]. H₂S has been reported playing important roles in many physiological and pathological conditions, such as inflammation regulation [7,8], cardioprotection [9], neuromodulation[1], hypertension [10], pain perception [11,12] and even cancer [13]. It has been found that amounts of H₂S that is endogenously generated are different in different tissues. H₂S amount in healthy cells is also different from that in abnormal cells. As a result, detection, and quantification of H₂S are of great importance in understanding its biological effects. Among various technologies for H₂S detection [14], fluorescent probes show great potential in detecting H₂S both in vitro and in vivo due to fast response, excellent selectivity, high sensitivity and real time imaging [15]. Several strategies have been developed for the design of fluorescent probes of H₂S [16], including reduction of aryl azide [17] and aryl nitro compounds [18], H₂S-specific reactions based on its nucleophilicity [19], H₂S induced metal displacement approach [20], and disulfide exchange reaction [21]. Among these strategies, azide-reduction has attracted extensive attention due to its fast reaction rate, high selectivity and harmless byproduct N₂. In 2011, Chang group reported a series of rhodamine-based fluorescent probes (SF1 and SF2), which can successfully and selectively detect H₂S via azide reduction in both water and live cells [22]. In the same year, Wang group developed a fluorescent probe DNS-Az which reached the maximum fluorescent intensity within 30 seconds upon mixing with sulfide anion [23]. The extremely fast reaction was believed to favor the real time detection of H₂S in vivo because of its rapid metabolism. Almost in the same year or around, various fluorophores were modified with azide group for H₂S detection, such as Cyanines-N₃ (Cy-N₃) [24], Phenanthroimidazole-N₃ (PI-N₃) [25], 7-nitrobenz-2-oxa-[1,3]diazole-N₃ (NBD-N₃) [26], Pyrene-1,3,6-trisulfonate-N₃ [27], Coumarin-N₃ [28,29], Naphthalimide-N₃ [30] and so on. In all these probes, an azide group was attached to the aromatic ring or the conjugate system. The azide functionality was either formed by oxidation of primary amine or installed by azide nucleophilic replacement of halide.

Azidomethyl has been used to protect amine or alcohol in the synthesis of carbohydrate[31,32], nucleosides[33], triazoles [34] and peptides [32]. Nowadays, the most important application of azidomethyl is as a reversible terminator in DNA sequencing by synthesis (SBS) [35]. The azidomethyl capped 3'-OH can be easily and quickly deprotected with reducing agents, and thus the SBS process can be continued. Because DNA sequencing is such a big project that it takes a long time, chemical reaction of each step is required to be as fast as possible. Azidomethyl reduction meets all requirements perfectly. Inspired by these works, we developed a turn-on fluorescent probe (FL-N₃) for H₂S detection by protecting fluorescein with azidomethyl and tested the probe in aqueous solution and cancer cells.

2. Results

2.1. Synthesis

The probe FL-N₃ was based on fluorescein with one of the hydroxyl groups protected by a polyethylene glycol (PEG) side chain which is expected to increase the probe's hydrophilicity [36]. The other hydroxyl group was caged with azidomethyl which reacts with sulfide to release free hydroxyl and turn fluorescence on (Scheme 1, FL-N₃ to FL-ONa). We synthesized this probe starting from commercially available fluorescein sodium in 5 steps including two processes of two-step in one-pot. Fluorescein sodium and tosylated hexaethylene glycol monomethyl ether were heated together in DMF. The free carboxylic acid and hydroxyl were both protected with hexaethylene glycol monomethyl ether by forming an ester and an ether respectively. Without purification of the intermediate, Hydrolysis of the newly formed ester with sodium hydroxide and subsequent acidification with HCl led to the formation of compound B. The re-produced phenolic hydroxyl was protected with methylthio-methyl (as in compound C), which was then converted to chloromethyl by reacting with NCS and TMSCl. The probe (FL-N₃) was achieved after nucleophilic substitution of the chloride with azide. Although 5 steps were required, two intermediates were not necessarily to be separated, which made the synthesis quite straight forward.



Scheme 1. Synthetic scheme for the probe FL-N₃ and its reaction with sulfide.

2.2. Fluorescence Response

We first tested the UV absorbance of the probe in solution with or without Na₂S. The probe alone showed no absorbance in the range from 300 to 600 nm. After incubated with sodium sulfide for 30 minutes, a strong absorbance peak appeared at 455nm (Supporting information, Figure S1) which indicated the formation of fluorescent product. The fluorescence properties of this probe were then tested at excitation wavelength of 455 nm. Although a PEG side chain has been attached, the solubility of the probe is still not good enough in pure water. As a result, tests were carried out in water and

methanol mixture. When the probe FL-N₃ was mixed with Na₂S at concentration of 1 mM, the fluorescence intensity increased by 32-fold at λ_{em} of 515 nm after 20 minutes (Figure 1a). The fluorescence response over time was also examined (Figure 1b). FL-N₃ and Na₂S were mixed at concentrations of 5 mM and 10 mM respectively. The fluorescence intensity increased quickly in the first twenty minutes. It reached the max after a slow increase during the period from 20 to 60 minute. The maximum fluorescence intensity kept almost consistent for one hour from 60 to 120 minute. The long-lasting fluorescence of the probe could be important in a lot of applications [37]. HPLC analysis demonstrated the formation of a major product (supporting information, Figure S2) and MS analysis indicated that the fluorescence increase was due to the formation of FL-ONa (Scheme 1, mass spectra data in supporting information, Figure S3).




Figure 1. Probe response to Na₂S: (a) Fluorescence response of 1 mM FL-N₃ to 20 mM Na₂S. Data were acquired after 20 minutes of reaction at 25 °C in water/MeOH (1/1, v/v) with excitation at λ_{ex} =455 nm, emission was collected between 490 and 600 nm; (b) Fluorescence response over time of 5 mM FL-N₃ to 10 mM Na₂S. Data were acquired at 25 °C in water/MeOH (1/1, v/v) with excitation at λ_{ex} =455 nm, emission was collected at λ_{em} =515 nm.

2.3. Selectivity

To test the probe's selectivity, we selected a series of biothios and bio-relevant sulfur-containing anions. Only HS⁻ induced a significant increase of fluorescence (Figure 2). Bio-thiols including cysteine, glutathione and dithiothreitol led to negligible fluorescence intensity change. Bio-relevant reducing salts such as sodium L-ascorbate, sodium thiocyanate and sodium thiosulfate also failed to turn the probe on. Other inorganic salts that were commonly used in buffer were also tested (SI, Figure S4). Ammonium acetate, sodium citrate, lithium chloride and dipotassium phosphate did not increase the fluorescence intensity. However, basic salts like potassium carbonate and lithium hydroxide led to about three- and five-fold fluorescence increase respectively comparing with the blank. This is probably due to the acidic property of the C-H bond on the carbon of azidomethyl group, which was deprotonated by hydroxide. One molecule of nitrogen and formamide were released to turn the fluorescence on. A possible mechanism was predicted as in SI (SI, Figure S5).




Figure 2. Fluorescence intensity of the probe (0.1 mM probe, 30 minutes after mixing) in the presence of sulfide or other species in water and methanol (10% MeOH, v/v). 1. Glutathione, 2. L-Cysteine, 3. Dithiothreitol, 4. Sodium L-Ascorbate, 5. Sodium Thiocyanate, 6. Sodium Thiosulfate, 7. Sodium sulfide, 8. Blank, (1-7: 1mM).

2.4. Linearity

It has been found that the bioactivity of H₂S is concentration dependent [38], which means quantification of H₂S in solution or cells is of great importance in H₂S study. When FL-N₃ was exposed to Na₂S of different concentrations, a linear correlation between fluorescence responses of the probe and Na₂S concentrations was observed, enabling a calibration equation to be established and applicable to the analysis of H₂S concentrations (Figure 3).




Figure 3. Linear correlation between fluorescence responses of FL-N₃ and Na₂S concentrations in water and methanol (10% MeOH, v/v). The probe concentration was 1 mM and data were acquired 30 minutes after mixing with Na₂S.

2.5. Detection in Cells

Since the probe FL-N₃ has well-behaved fluorescent property and selectivity for H₂S, it is probably suitable for the detection of H₂S in cultured cells. Confocal imaging to visualize H₂S inside cells was carried out. MCF-7 cells were incubated together with FL-N₃, after washing free molecules of probe away, Na₂S solution was added. Confocal images were taken after another 60-minute incubation and results were shown in Figure 4. The probe successfully entered cells and reacted with sulfide to show red fluorescence (Figure 4a). In contrast, the sample of control (Figure 4b) showed no fluorescence at all. We also tested compound D which contains a methyl group instead of PEG side chain. Surprisingly, compound D failed to enter cells and little fluorescence was observed after treatment with Na₂S (Figure 4c). This result proved the importance of the PEG side chain for probe's hydrophilicity and cell permeability, which are key factors in bio-applications.




Figure 4. Confocal images showing signals in MCF-7 cells exposed to Na₂S. MCF-7 cells were incubated with 20 μM probe FL-N₃ or comparing compound D for 2 h at 37 °C, washed, and then incubated with or without Na₂S for 1 h before imaging: (a) FL-N₃ with 100 μM Na₂S (b) FL-N₃ without Na₂S (c) Compound D with 100 μM Na₂S (d) Compound D without Na₂S.

3. Discussion

We designed and synthesized a turning-on probe for the detection of H₂S in solution and in cells. The probe works through the reduction of azidomethyl group by H₂S. It also inspired us that azidomethyl group could be an ideal choice in the modification of turning-on dyes. When fluorescence is turned off by capping different dyes with azidomethyl, it can be turned on easily by adding certain reductant, which is potential in biology research.

4. Materials and Methods

Procedure for probe response to sodium sulfide test: The probe FL-N₃ was dissolved in methanol at a concentration of 2 mM. Sodium sulfide was dissolved in water at a concentration of 40 mM. Two wells of a corning 96-well plate were selected and to each well was added 100 μL probe solution. 100 μL Na₂S solution was added to one of the wells and to the other was added 100 μL water. The plate was incubated at RT for 20 minutes before reading fluorescence on a plate reader. The fluorescence was read at λ_{ex}=455 nm with λ_{em} set from 500 nm to 600 nm.

Procedure for fluorescence response over time test: The probe FL-N₃ was dissolved in methanol at a concentration of 10 mM. Sodium sulfide was dissolved in water at a concentration of 20 mM. To a corning 96-well plate was added 100 μL probe solution and 100 μL Na₂S solution. The fluorescence was read on a plate reader every minute in 2 hours with λ_{ex}=455 nm and λ_{em}= 515 nm.

Procedure for probe selectivity test: The probe FL-N₃ was dissolved in methanol at a concentration of 1 mM. Other tested compounds or salts were dissolved in water at a concentration of 10 mM. To a corning 96-well plate, was added 160 μL water and 20 μL probe solution. Then 20 μL aqueous solution of one tested compound or salt was added to one well. To the well of blank, 20 μL water was added. After 30-minute incubation at RT, fluorescence was read on a plate reader (brand) with λ_{ex}=455 nm and λ_{em}= 515 nm.

Procedure for linear correlation test: The probe FL-N₃ was dissolved in methanol at a concentration of 10 mM. To 6 wells of a corning 96-well plate was added 20 μL of the probe solution. Following this, to the 6 wells was added 180 μL, 160 μL, 140 μL, 100 μL, 60 μL, 20 μL water in order. Then 0 μL, 20 μL, 40 μL, 80 μL, 120 μL, 160 μL of sodium sulfide solution (1 mM) was added to the 6 wells in order. After incubation at RT for 20 minutes, the fluorescence was read on a plate reader with λ_{ex}=455 nm and λ_{em}= 515 nm.

Procedure for confocal imaging: MCF7 cells were seeded at 20000 cells/well in an 8-chamber plate (0.8 cm²). Probe FL-N₃ (20 μM in DMSO) was added and cells were incubated for 2 h at 37 °C. Cells were then washed with PBS (3×), and Na₂S solution (0.1 mM) was added and incubated for 5 h. Fluorescent images of the live cells were obtained using a confocal fluorescence microscope FV 1000 and processed using Olympus Fluoview Ver.3.1. Viewer (Olympus, Tokyo, Japan).

Procedure for the synthesis of compound B: Fluorescein sodium salt (135 mg, 0.36mmol) was added to a two necked rbf, after evacuation, the rbf was recharged with N₂, and anhydrous DMF (3 mL) was added followed by addition of 2,5,8,11,14,17-hexaoxonadecan-19-yl 4-methylbenzenesulfonate (450mg, 1mmol). The mixture was heated to 90 degree and stirred for 36 hours. The mixture was then cooled to room temperature, and DMF was removed by evaporation and the residual was re-dissolved in 10 mL 10% NaHCO₃ solution. Extracted with EA for three times and the organic layer was combined, washed with brine, dried over Na₂SO₄ and concentrated. The residual was re-dissolved in 5 mL methanol, 2 M NaOH (2 mL) was added, and stirred at room temperature for 2 hours. 1M HCl was added slowly to adjust PH to 2, and the mixture was stirred for 30 minutes. Methanol was removed via rotary evaporation and the residual was extracted with EA for three times. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The product was isolated via silica gel chromatography (DCM/MeOH = 100/1) with a yield of 65%.

Procedure for the synthesis of compound C: Compound B (122mg, 0.2mmol) was dissolved in 10 mL anhydrous MeCN under Ar. Silver oxide (92 mg, 0.4 mmol) was added followed by addition of chloromethyl methyl thioether (167 μ L, 2 mmol) and one drop of pyridine. The mixture was heated to 50 degree and stirred for 48 hours. Cooled to room temperature and solvent was removed via rotary evaporation. The product was isolated via silica gel chromatography (DCM/MeOH = 100/1) with a yield of 81%.

Procedure for the synthesis of probe FL-N₃: Compound C (67mg, 0.1 mmol) was dissolved in 5 mL anhydrous DCM under Ar. N-chlorosuccinimide (16 mg, 0.12 mmol) was added and the mixture was stirred at room temperature for 3 hours. Then trimethyl silane chloride (13 mg, 0.12 mmol) was added and the mixture was stirred at room temperature further for 6 hours. After removal of solvent via rotary evaporation, the residual was dried under vacuum and re-dissolved in dry THF under Ar, following this, TMSN₃ (0.2 mmol) and TBAF (0.2 mmol, 1M in THF) were added and the mixture was stirred at room temperature overnight. Remove solvent via rotary evaporation and the product was isolated via silica gel chromatography (DCM/MeOH = 100/1) as a yellow oil. The product was further purified via prep HPLC in the case of impure

5. Conclusions

In summary, we have developed a new strategy to construct fluorescein-based probe for the detection of H₂S in solution and in cells. The probe showed fast response and excellent selectivity to sulfide over a series of bio-relevant sulfur containing compounds and inorganic salts. Linear correlation of Na₂S concentration and probe fluorescent response made the quantification of H₂S in solution possible. Its successful application in cell confocal imaging may provide a new tool in the study of biological effects of hydrogen sulfide.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

Author Contributions: Wei Feng and Qicai Xiao made equal contribution to this work. Conceptualization, Wei Feng and Yuanyong Yang.; methodology, Wei Feng and Qicai Xiao; investigation, Qicai Xiao and Lu Wang; writing—original draft preparation, Wei Feng; writing—review and editing, Yuanyong Yang; supervision, Yuanyong Yang; funding acquisition, Wei Feng and Yuanyong Yang. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by National Natural Science Foundation of China, grant number 21807120, 22061012 and Department of Science & Technology of Guizhou Province ([2020]4Y208).

Data Availability Statement: The data presented in this study are available in supplementary information.

Acknowledgments: We acknowledge the support from Sun Yat-sen University and BGI-Shenzhen.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are available from the authors.

References

1. Abe, K.; Kimura, H. The possible role of hydrogen sulfide as an endogenous neuromodulator. *J. Neurosci.* **1996**, *16*, 1066-1071. DOI: 10.1523/JNEUROSCI.16-03-01066.1996.
2. Wang, R. Two's company, three's a crowd: can H₂S be the third endogenous gaseous transmitter? *FASEB J.* **2002**, *16*, 1792-1798. DOI: 10.1096/fj.02-0211hyp.
3. Xiao, Q.; Ying, J.; Xiang, L.; Zhang, C. The biologic effect of hydrogen sulfide and its function in various diseases. *Medicine* **2018**, *97*, E13065. DOI: 10.1097/MD.00000000000013065.
4. Cao, X.; Ding, L.; Xie, Z.-z.; Yang, Y.; Whiteman, M.; Moore, P. K.; Bian, J.-S. A Review of Hydrogen Sulfide Synthesis, Metabolism, and Measurement: Is Modulation of Hydrogen Sulfide a Novel Therapeutic for Cancer? *Antioxid. Redox Sign.* **2019**, *31*, 1-38. DOI: 10.1089/ars.2017.7058.
5. Song Z. J.; Ng, M. Y.; Lee, Z.-W.; Dai, W.; Hagen, T.; Moore, P. K.; Huang, D.; Deng, L.-W.; Tan, C.-H. Hydrogen sulfide donors in research and drug development. *MedChemComm* **2014**, *5*, 557-570. DOI: 10.1039/c3md00362k.

6. Jose, D. A.; Sakla, R.; Sharma, N.; Gadiyaram, S.; Kaushik, R.; Ghosh, A. Sensing and Bioimaging of the Gaseous Signaling Molecule Hydrogen Sulfide by Near-Infrared Fluorescent Probes. *ACS Sensors* **2020**, *5*, 3365-3391. DOI: 10.1021/acssensors.0c02005.
7. Li L., Bhatia, M.; Zhu, Y. Z.; Zhu, Y. C.; Ramnath, R. D.; Wang, Z. J.; Anuar, F. B. M.; Whiteman, M.; Salto-Tellez, M.; Moore, P. K. Hydrogen sulfide is a novel mediator of lipopolysaccharide-induced inflammation in the mouse. *FASEB J.* **2005**, *19*, 1196-1198. DOI: 10.1096/fj.04-3583fje.
8. Fiorucci S. Hydrogen Sulfide: From Physiology to Pharmacology. *Inflamm. Allergy Drug Targets* **2011**, *10*, 77-84. DOI: 10.2174/187152811794776277.
9. Pan, T.-T.; Feng, Z.-N.; Lee, S. W.; Moore, P. K.; Bian, J.-S. Endogenous hydrogen sulfide contributes to the cardioprotection by metabolic inhibition preconditioning in the rat ventricular myocytes. *J. Mol. Cell. Cardiol.* **2006**, *40*, 119-130. DOI: 10.1016/j.yjmcc.2005.10.003.
10. Zhong, G.; Chen, F.; Cheng, Y.; Tang, C.; Du, J. The role of hydrogen sulfide generation in the pathogenesis of hypertension in rats induced by inhibition of nitric oxide synthase. *J. hypertens.* **2003**, *21*, 1879-1885. DOI: 10.1097/01.hjh.0000084760.37215.75.
11. Kawabata, A.; Ishiki, T.; Nagasawa, K.; Yoshida, S.; Maeda, Y.; Takahashi, T.; Sekiguchi, F.; Wada, T.; Ichida, S.; Nishikawa, H. Hydrogen sulfide as a novel nociceptive messenger. *Pain* **2007**, *132*, 74-81. DOI: 10.1016/j.pain.2007.01.026.
12. Distrutti, E.; Sediari, L.; Mencarelli, A.; Renga, B.; Orlandi, S.; Antonelli, E.; Roviezzo, F.; Morelli, A.; Cirino, G.; Wallace, J. L. Evidence that hydrogen sulfide exerts antinociceptive effects in the gastrointestinal tract by activating KATP channels. *J. Pharmacol. Exp. Ther.* **2006**, *316*, 325-335. DOI: 10.1124/jpet.105.091595.
13. Feng, W.; Teo, X.-Y.; Novera, W.; Ramanujulu, P. M.; Liang, D.; Huang, D.; Moore, P. K.; Deng, L.-W.; Dymock, B. W. Discovery of New H₂S Releasing Phosphordithioates and 2,3-Dihydro-2-phenyl-2-sulfanylenebenzo[d][1,3,2]oxazaphospholes with Improved Antiproliferative Activity. *J. Med. Chem.* **2015**, *58*, 6456-6480. DOI: 10.1021/acs.jmedchem.5b00848.
14. Pandey, S. K.; Kim, K.-H.; Tang, K.-T. A review of sensor-based methods for monitoring hydrogen sulfide. *TrAC-Trend. Anal. Chem.* **2012**, *32*, 87-99. DOI: 10.1016/j.trac.2011.08.008.
15. Li, J.; Yin, C.; Huo, F. Chromogenic and fluorogenic chemosensors for hydrogen sulfide: review of detection mechanisms since the year 2009. *Rsc Adv.* **2015**, *5*, 2191-2206. DOI: 10.1002/chin.201511350.
16. Feng, W.; Dymock, B. W. Fluorescent Probes for H₂S Detection and Quantification. In *Chemistry, Biochemistry Pharmacology of Hydrogen Sulfide* **2015**, 291-323. DOI: 10.1007/978-3-319-18144-8_15.
17. Lin, V. S.; Lippert, A. R.; Chang, C. J. Chapter Four. Azide-Based Fluorescent Probes. In *Methods in Enzymology, Vol. 554*, Elsevier, **2015**, pp. 63-80. DOI: 10.1016/bs.mie.2014.11.011.
18. Bae, J.; Choi, M. G.; Choi, J.; Chang, S.-K. Colorimetric signaling of hydrogen sulfide by reduction of a phenylseleno-nitrobenzoxadiazole derivative. *Dyes Pigments* **2013**, *99*, 748-752. DOI: 10.1016/j.dyepig.2013.06.018.
19. Liu, C.; Chen, W.; Shi, W.; Peng, B.; Zhao, Y.; Ma, H.; Xian, M. Rational Design and Bioimaging Applications of Highly Selective Fluorescence Probes for Hydrogen Polysulfides. *J. Am. Chem. Soc.* **2014**, *136*, 7257-7260. DOI: 10.1021/ja502968x.
20. Qu, X.; Li, C.; Chen, H.; Mack, J.; Guo, Z.; Shen, Z. A red fluorescent turn-on probe for hydrogen sulfide and its application in living cells. *Chem. Commun.* **2013**, *49*, 7510-7512. DOI: 10.1039/c3cc44128h.
21. Liu, C.; Pan, J.; Li, S.; Zhao, Y.; Wu, L. Y.; Berkman, C. E.; Whorton, A. R.; Xian, M. Capture and Visualization of Hydrogen Sulfide by a Fluorescent Probe. *Angew. Chem.* **2011**, *123*, 10511-10513. DOI: 10.1002/anie.201104305.
22. Lippert, A. R.; New, E. J.; Chang, C. J. Reaction-based fluorescent probes for selective imaging of hydrogen sulfide in living cells. *J. Am. Chem. Soc.* **2011**, *133*, 10078-10080. DOI: 10.1021/ja203661j.
23. Peng, H.; Cheng, Y.; Dai, C.; King, A. L.; Predmore, B. L.; Lefer, D. J.; Wang, B. A Fluorescent Probe for Fast and Quantitative Detection of Hydrogen Sulfide in Blood. *Angew. Chem.* **2011**, *123*, 9846-9849. DOI: 10.1002/anie.201104236
24. Yu, F.; Li, P.; Song, P.; Wang, B.; Zhao, J.; Han, K. Facilitative functionalization of cyanine dye by an on-off-on fluorescent switch for imaging of H₂O₂ oxidative stress and thiols reducing repair in cells and tissues. *Chem. Commun.* **2012**, *48*, 2852-2854. DOI: 10.1039/C2CC30985H.
25. Zheng, K.; Lin, W.; Tan, L. A phenanthroimidazole-based fluorescent chemosensor for imaging hydrogen sulfide in living cells. *Org. Biomol. Chem.* **2012**, *10*, 9683-9688. DOI: 10.1039/c2ob26956b.
26. Zhou, G.; Wang, H.; Yang M.; Chen, X. An NBD fluorophore-based colorimetric and fluorescent chemosensor for hydrogen sulfide and its application for bioimaging. *Tetrahedron* **2013**, *69*, 867-870. DOI: 10.1016/j.tet.2012.10.106.
27. Hartman, M. C.; Dcona, M. M. Chemodosimeter for direct measurement of hydrogen sulfide in biological fluids. *Analyst* **2012**, *137*, 4910-4912. DOI:10.1039/c2an35870k.
28. Li, W.; Sun, W.; Yu, X.; Du, L.; Li, M. Coumarin-based Fluorescent Probes for H₂S Detection. *J. fluoresc.* **2013**, *23*, 181-186. DOI: 10.1007/s10895-012-1131-4.

29. Chen, B.; Li, W.; Lv, C.; Zhao, M.; Jin, H.; Jin, H.; Du, J.; Zhang, L.; Tang, X. Fluorescent probe for highly selective and sensitive detection of hydrogen sulfide in living cells and cardiac tissues. *Analyst* **2013**, *138*, 946-951. DOI: 10.1039/c2an36113b.
30. Montoya, L. A.; Pluth, M. D. Selective turn-on fluorescent probes for imaging hydrogen sulfide in living cells. *Chem. Commun.* **2012**, *48*, 4767-4769. DOI: 10.1039/c2cc30730h.
31. Xu, J.; Guo, Z. (2-Azidomethyl)phenylacetyl as a new, reductively cleavable protecting group for hydroxyl groups in carbohydrate synthesis. *Carbohydrate research* **2002**, *337*, 87-91. DOI: 10.1016/S0008-6215(01)00292-0.
32. Pothukanuri, S.; Winssinger, N. A Highly Efficient Azide-Based Protecting Group for Amines and Alcohols. *Org. Lett.* **2007**, *9*, 2223-2225. DOI: 10.1002/chin.200739049.
33. Wada, T.; Ohkubo, A.; Mochizuki, A.; Sekine, M. 2-(Azidomethyl)benzoyl as a new protecting group in nucleosides. *Tetrahedron Lett.* **2001**, *42*, 1069-1072. DOI: 10.1016/S0040-4039(00)02183-3.
34. Loren, J. C.; Krasinski, A.; Fokin, V. V.; Sharpless, K. B. NH-1,2,3-Triazoles from Azidomethyl Pivalate and Carbamates: Base-Labile N-Protecting Groups. *Synlett* **2005**, *2005*, 2847-2850. DOI: 10.1002/chin.200614133.
35. Guo, J.; Xu, N.; Li, Z.; Zhang, S.; Wu, J.; Kim, D. H.; Marma, M. S.; Meng, Q.; Cao, H.; Li, X.; Shi S.; Yu L.; Kalachikov S.; Russo J. J.; Turro N. J. Four-color DNA sequencing with 3'-O-modified nucleotide reversible terminators and chemically cleavable fluorescent dideoxynucleotides. *P. Natl. Acad. Sci. USA* **2008**, *105*, 9145-9150. DOI: 10.1073/pnas.0804023105.
36. Kiani, S.; Mousavi, S. M.; Shahtahmassebi, N.; Saljoughi, E. Hydrophilicity improvement in polyphenylsulfone nanofibrous filtration membranes through addition of polyethylene glycol. *Appl. Surf. Sci.* **2015**, *359*, 252-258. DOI: 10.1016/j.apsusc.2015.10.107.
37. Fu, Y.; Zhang, J.; Lakowicz, J. R. Reduced blinking and long-lasting fluorescence of single fluorophores coupling to silver nanoparticles. *Langmuir* **2008**, *24*, 3429-3433. DOI: 10.1021/la702673p.
38. Lee, Z. W.; Teo, X. Y.; Tay, Y. W.; Tan, C. H.; Hagen, T.; Moore, P.; Deng, L.-W. *Brit. J. Pharmacol.* **2014**, *171*, 4322-4336. DOI: 10.1111/bph.12773.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.