

Article

Not peer-reviewed version

---

# Quantum Purity as an Information Measure and Nernst law

---

[Angelo Plastino](#)<sup>\*</sup> and [F. Pennini](#)

Posted Date: 7 July 2023

doi: 10.20944/preprints202307.0515.v1

Keywords: Nernst law, statistical mechanics, purity



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# Quantum Purity as an Information Measure and Nernst Law

F. Pennini <sup>1,2,\*</sup> and A. Plastino <sup>3</sup>

<sup>1</sup> Departamento de Física, Universidad Católica del Norte, Av. Angamos 0610, Antofagasta, Chile

<sup>2</sup> Departamento de Física, Facultad de Ingeniería, Universidad Nacional de Mar del Plata (UNMDP), CONICET, Mar del Plata, Argentina

<sup>3</sup> Instituto de Física La Plata–CCT-CONICET, Universidad Nacional de La Plata, C.C. 727, 1900, La Plata, Argentina

**Abstract:** We attempt to re-express Nernst law in terms of a suitable information measure (IM). This is done on the basis of adapting the idea of purity to a thermal Gibbs environment, which yields what we might call a “purity” (and symbolize it by the symbol  $D$ ). We further apply the properly generalization of this  $D$ -IM to a classical scenario. This generalization turns out to have interesting consequences, when used in conjunction with the classical Shannon entropy  $S$ . These consequences are related to Nernst law.

**Keywords:** Nernst law; statistical mechanics; purity

## 1. Introduction

In quantum mechanics, the purity measures how similar is a given quantum state to a pure one. In other words, it provides insights into the degree to which a quantum state is pure or mixed, and hence the amount of information that can be extracted from it. Mathematically, the purity  $D$  of a quantum state is given by the trace of the square of its density operator  $\rho$ , i.e.,  $Tr(\rho^2)$ . For a pure state, the purity is 1, indicating that the state is entirely non-mixed. On the other hand, for a maximally mixed state (also known as a completely mixed state), the purity is  $1/M$ , where  $M$  is the dimensionality of the Hilbert space of the system. In a mixed state, the purity lies between these two extremes.

Both the purity and degree of mixture  $1 - D$  provide measures of the information content present in a quantum state. Pure states, with a purity of 1, typically exhibit a coherent superposition and can be entangled. Mixed states, on the other hand, have reduced purity and contain a mixture of quantum states with varying probabilities. The degree of mixture quantifies the extent of the mixture and reflects the amount of information and entanglement present in the state. These information measures are widely used in quantum information theory, quantum state estimation, and the characterization of quantum systems, as they provide valuable insights into the purity and degree of mixture of quantum states [1–4].

### 1.1. Our goals and organization

The Third Law of thermodynamics (Nernst law) implies that it is impossible to reach absolute zero temperature (AZT) by any finite number of processes, and that at AZT the entropy of a perfectly ordered crystalline substance. In other words, a perfectly ordered crystal at absolute zero temperature would have no disorder or randomness, and would be in a state of minimum energy [5].

This law has important implications for the behavior of matter at low temperatures and is essential for understanding the behavior of condensed matter physics, including the properties of solids, liquids, and super-fluids. The third Law is also used in the design and optimization of refrigeration and cooling systems, as well as in the study of chemical reactions and the behavior of materials at extreme conditions [5].

Here, instead, we discuss an alternative viewpoint. We will try to embark into an information theory discussion in which the protagonist in the notion of purity  $D$  regarded as an information measure (IM).

We will focus our attention on the harmonic oscillator (HO) and on the ideal gas, to provide an understandable and clear picture of our state of affairs. The simplicity of the systems enables one to easily visually what the alternative  $D$  interpretation here to be advanced means in a physical scenario.

Thus, we will try to link thermodynamics third law to the notion of purity within the framework of 1) the harmonic oscillator and 2) the ideal gas. In more detail:

- In Section II we consider the purity notion first to a quantum thermal scenario and then to a classical environment.
- In Section III we obtain a useful expression for the purity quantifier  $D$ , that relates it to the Helmholtz free energy. This relation serves to considerably simplify later on our  $D$  manipulations.
- In Section IV we consider the quantum Harmonic Oscillator (HO) and concoct a spacial way of expressing its Shannon entropy  $S$  solely in  $D$  terms, which illuminates some aspects of Nernst law.
- In Section V, the nucleus of this effort, we consider the classical HO and comment on some interesting traits.
- Section VI is devoted to the ideal gas. Some new insight is gained.
- Section VII is devoted to conclusions.

## 2. A generalization of the purity notion to a finite temperature scenario

The density operator that describes the statistical ensemble of quantum states in a system at finite temperature incorporates the probabilities associated with different energy eigenstates of the system. In the case of a discrete set of energy eigenstates, the density operator can be expressed as:

$$\rho = \sum_i P_i |i\rangle\langle i|, \quad (1)$$

where  $P_i$  represents the probability associated with the  $i$ -th energy eigenstate. It is determined by the thermal equilibrium, canonical distribution of Gibbs' that uses the so called exponential Boltzmann factors [5]

$$P_i = (1/Z) \exp(-\beta E_i), \quad (2)$$

so that our principal system-input is the set of level energies  $E_i$ . We denote  $\beta = 1/kT$  ( $k$  is the Boltzmann-constant). this entails a set of canonical probabilities  $P_i = \exp(-\beta E_i)/Z$ ,  $Z$  being the partition function. Several quantifiers like the entropy  $S$  or the free energy  $F$  are built up with the  $P_i$  [5]. By summing over all possible energy eigenstates and incorporating the corresponding probabilities, the appropriate density operator captures the ensemble average behavior of the system at finite temperature. The diagonal elements of the density operator provide the populations of the energy eigenstates, while the off-diagonal elements capture the coherence and quantum correlations between different eigenstates. The density operator at finite temperature provides a powerful tool for analyzing and characterizing the behavior of quantum systems under thermal equilibrium conditions [5], accounting for the probabilities associated with different energy eigenstates and their corresponding energies. The purity  $D$  is obviously

$$D = \text{Tr}\rho^2 = \sum_i P_i^2. \quad (3)$$

Notice that small  $D$  might be said to entail a kind of disorder, as we have then only very low probability elements  $P_i$  that can not differ too much from one another- That is, we are very close to an uniform distribution representing maximum randomness. On the other hand, large  $D$  might be said to imply order as one must encounter some (relatively) very large  $P_i$ s, indicating preferred micro-states and

thus some kind of structure. Thus,  $D$  might be regarded as an order-disorder indicator. The classical counterpart of  $D$ , to be discussed below, is known to be an indicator of such kind.

### 2.1. Generalization of the purity notion to a classical scenario

The pertinent discussion can be made in terms of an statistical concept advanced around 25 years ago, called the disequilibrium, an order-disorder quantifier [6]. Low disequilibrium entails disorder and large disequilibrium entails order. The central idea is that, statistically, maximum disorder is represented by a uniform distribution (UF) [6]. As a consequence, the more different our current probability distribution (CPD) is from the UF, the more order this CPD represents [6]. The associated Euclidean distance in probability space between the CPD and the UF is the disequilibrium [6]. We will baptize it also as  $D$ . The product of  $D$  with the Shannon entropy  $S$

$$C = D S, \quad (4)$$

is called the statistical complexity. This  $C$ -expression is in many instances the standard complexity form employed in several current publications. For a small sample, see, for instance, Refs. [7–14]. Why repeating the symbol  $D$  for naming two different notions? Because, formally, both equal the sum of the squares of probability elements  $P_i$ . At finite temperature and within Gibbs' canonical ensemble, two two pertinent types of probability coincide.

### 3. Relation between $D$ and the Helmholtz free energy $F$

Given the partition function  $Z$  we have the sequence [5]

$$Z(T) = \sum_i \exp(-\beta E_i), \quad (5)$$

$$F(T) = -kT \ln(Z(T)), \quad (6)$$

$$F(T/2) = -k(T/2) \ln(Z(T/2)), \quad (7)$$

$$2\beta[F(T) - F(T/2)] = 2\beta kT [(1/2) \ln(Z(T/2)) - \ln(Z(T))], \quad (8)$$

$$2\beta[F(T) - F(T/2)] = \ln \left[ \frac{Z(T/2)}{Z(T)^2} \right], \quad (9)$$

$$\exp(2\beta[F(T) - F(T/2)]) = \frac{Z(T/2)}{Z(T)^2}, \quad (10)$$

$$\frac{Z(T/2)}{Z(T)^2} = \left[ \sum_i \exp(-\beta E_i) / Z \right]^2 = \sum_i P_i^2 = D, \quad (11)$$

so that we finally arrive at our desired relation that connects  $D$  with  $F$ :

$$D = \exp[2\beta(F(T) - F(T/2))]. \quad (12)$$

We will use this important expression in the following sections.

### 4. Purity and the quantum harmonic oscillator

Consider a three-dimensional harmonic oscillators of frequency  $\omega$  in equilibrium at temperature  $T$ , whose energies are quantified according to  $\epsilon_n = \hbar\omega(n + 3/2)$ , with  $n = 0, 1, 2, \dots$ . The canonical partition function for a single oscillator is [5]

$$Z_1 = \frac{e^{-3\beta\hbar\omega/2}}{(1 - e^{-\beta\hbar\omega})^3}, \quad (13)$$

while the Shannon entropy reads [5]

$$S = 3k \left[ \frac{\beta\hbar\omega e^{-\beta\hbar\omega}}{1 - e^{-\beta\hbar\omega}} - \ln(1 - e^{-\beta\hbar\omega}) \right], \quad (14)$$

which is definite positive for all temperature  $T$ . Also, the Helmholtz free energy is

$$F = \frac{3}{2}\hbar\omega + 3kT \ln(1 - e^{-\beta\hbar\omega}). \quad (15)$$

Now, we follow the indications given in Ref. [7] in order to obtain the purity  $D$  that, as we just saw above, is a simple function of the canonical Helmholtz free energy  $F$ . We had in (12) for the purity

$$D = \exp[2\beta(F(T) - F(T/2))]. \quad (16)$$

Therefore, considering that  $F$  is given by (15), then one finds that the purity is given by

$$D = \left( \frac{1 - e^{-\beta\hbar\omega}}{1 + e^{-\beta\hbar\omega}} \right)^3 = \tanh^3 \left( \frac{\beta\hbar\omega}{2} \right). \quad (17)$$

Note that, when  $T = 0$ , then  $D = 1$ , and for  $T$  tending towards infinity we have  $D = 0$ . Therefore, the purity is of course bounded, i.e.,

$$0 \leq D \leq 1. \quad (18)$$

The above relation (17) shows that for small  $\beta$  the hyperbolic tangent above approaches  $\beta\hbar\omega/2$ . From Eq. (17), we can isolate the variable  $\beta\hbar\omega$  in the fashion

$$\beta\hbar\omega = -\ln \left( \frac{1 - D^{1/3}}{1 + D^{1/3}} \right) = 2 \operatorname{arctanh} \left( D^{1/3} \right). \quad (19)$$

Now, introducing Eq. (19) into Eq. (14), we find the entropy  $S$  can be cast as a function of just  $D$  [15]. Thus, we reach the wished for possibility of expressing the entropy solely in terms of the purity

$$S = -3k \left[ \left( \frac{1 - D^{1/3}}{2D^{1/3}} \right) \ln \left( \frac{1 - D^{1/3}}{1 + D^{1/3}} \right) + \ln \left( \frac{2D^{1/3}}{1 + D^{1/3}} \right) \right]. \quad (20)$$

We clearly see that maximum  $D$  yields zero entropy, the third law of thermodynamics cast in terms of purity. Let us insist on this point: *When we express  $S$  in terms of  $D$ , we realize that, at maximum purity, the entropy vanishes. This may seem rather obvious, but it opens the door for us to ask: what happens classically when we express the classical  $S$  in terms of the classical  $D$ ? Could it be that some similar connection  $S$ - $D$  might emerge?. We will see below that the answer is a positive one.*

## 5. Classical harmonic oscillator and classically generalized $D$

We saw above that the classically generalized  $D$  is called disequilibrium, a notion advanced some 25 years ago, independently of the purity idea [6]. The purity-disequilibrium connection is being made in this article. Let us then consider the three-dimensional harmonic oscillators of frequency  $\omega$  in a Gibbs' canonical ensemble at the temperature  $T$ , whose respective Hamiltonian is  $\mathcal{H} = \sum_{i=1}^{3N} p_i^2 / (2m) + m\omega^2 x_i^2 / 2$  [5]. *We wish to ascertain what happens when we attempt to express  $S$  in terms of  $D$ .*

The partition function of each oscillator is

$$Z = \left( \frac{kT}{\hbar\omega} \right)^3, \quad (21)$$

and contains all the statistical HO information necessary for our ends. The entropy is [5]

$$S = 3k \left[ 1 - \ln \left( \frac{\hbar\omega}{kT} \right) \right], \quad (22)$$

which is positive definite whenever

$$T \geq \hbar\omega / (ek). \quad (23)$$

This yields the condition to avoid negative entropies.

Remember that we saw above that  $D$  is a simple function of  $F$ , of a form given by Eq. (16) [7]. Then, considering that  $F = -kT \ln Z$ , one finds

$$D = \left( \frac{\hbar\omega}{2kT} \right)^3. \quad (24)$$

and also

$$Z = \frac{1}{2D}, \quad (25)$$

which assures us that the whole classical HO statistical mechanics is governed by  $D$ .

Remark that, as in the quantum instance, both the pertinent partition function and  $S$  are simple functions of  $D$  and, importantly enough (see Eq. (23) and Eq. (24)),

$$0 \leq D \leq D_{max}, \quad (26)$$

since, let us repeat, the temperature  $\hbar\omega / (ek) \leq T < \infty$ . In addition, the classical statistical complexity is of the form

$$C = D(S/k) = 3 \left( \frac{\hbar\omega}{2kT} \right)^3 \left[ 1 - \ln \left( \frac{\hbar\omega}{kT} \right) \right]. \quad (27)$$

### 5.1. Present results for the HO

Let us first insist on (a) that the generalized  $D$  is a measure of order, and (b) repeating Eq. (24)

$$\left( \frac{\hbar\omega}{2kT} \right)^3 = D, \quad (28)$$

to emphasize that **the generalized purity, here called disequilibrium, is the ratio between the HO's vibrational and kinetic energies**. Replacing this into Eq. (27) we are led, for the single HO, to the desideratum of expression the Shannon entropy entirely in terms of  $D$ , i.e.,

$$S = 3k[1 - \ln(2D^{1/3})]. \quad (29)$$

It is clear that the entropy will vanish for a special  $D$  value. We need to ascertain the meaning of such special value. Now, for the statistical complexity we find

$$C = 3D \left[ 1 - \ln(2D^{1/3}) \right]. \quad (30)$$

It is important now to note that, given that the  $C$ -measure must be a positive quantity, as stipulated in Ref. [16], then the maximum classical value for  $D$  can not be infinite. It must at most adopt that value which makes  $\ln(2D^{1/3}) = 1$ , entailing, as we saw above and repeat here because of its relevance,

$$0 \leq D \leq D_{max}, \quad (31)$$

with

$$D_{max} = (e/2)^3 \sim 2.50. \quad (32)$$

This provides us with a maximal classical degree of order. Also, it results in several interesting and hopefully new classical features. The C-positivity argument also applies to the entropy, of course, with identical results.

*Additionally, for such  $D_{max}$ -disequilibrium-value the entropy vanishes according to Eq. (29). Thus, maximum order implies zero entropy as in the quantum instance above, which can be rephrased as an order-based third law.* The special quantum connection S-D of the preceding Section has indeed a classical counterpart. We can thus loosely speak of a kind of classical Nernst law of vanishing entropy at the situation of maximum order (as measured by  $D$ ).

Let us here insist again on a previous assertion by reiterating that from Eq. (24) we saw that

$$\frac{\hbar\omega}{2kD^{1/3}} = T, \quad (33)$$

which, using our above result for  $D_{max}$  connects temperature with our kind of disorder measured by the generalized  $D$ . Temperature grows as the disorder degree augments, as it is intuitively obvious. But using now Eq. (32) we see that this last assertion implies that as order, that grows with  $D$ , has a maximum, classical temperature has a frequency-depending (non zero!) minimum (see Eq. (35) below). For

$$D > (e/2)^3 \quad \text{one finds } S < 0. \quad (34)$$

**Negative entropy entails using an incorrect value for  $D$**  or, which is the same, a too low energy, according to the prescription  $u < u_{minimum}$  with

$$u_{minimum} = \frac{\hbar\omega}{e}. \quad (35)$$

Moreover, we realize that

$$\beta = \frac{2D^{1/3}}{\hbar\omega}, \quad (36)$$

so that the Gibbs canonical probability  $P$  for an HO Hamiltonian  $\mathcal{H}$  and a partition function  $Z$  reads

$$P = \frac{\exp(-\frac{2D^{1/3}}{\hbar\omega}\mathcal{H})}{Z}, \quad (37)$$

which tells us that the HO physics is fully determined by the ratio  $D^{1/3}/\omega$ . We duly note that  $D$  has a maximum possible value. Given Eqs. (33) and (34), we discover that, for fixed frequency (neither too high nor too low) the permissible minimum temperature is of the order of  $10^{-10}$  Kelvin. As an additional comment we revisit the mean energy of the harmonic oscillator [5]

$$u = 3kT. \quad (38)$$

Thus, in view of Eq. (28), for the single HO, we have, for  $N$  HOs, a mean energy  $U$

$$u = \frac{3\hbar\omega/2}{D^{1/3}}, \quad (39)$$

The above classical (and of statistical origin) minimum energy has a value slightly smaller than the well known quantum one.

We see that the HO energy can be thought of as being purely originated by disorder. If the order degree indicator  $D$  is large enough, the system will be encountered at a very low energy, but not arbitrarily low. This is indeed a new result. The classical HO seems to have a minimum energy, a minimum temperature, and a maximal order degree at which  $S = 0$ , if we respect the condition that the statistical complexity can not be negative.

## 6. The ideal gas

The system consists of  $N$  mono-atomic identical particles contained in a volume  $V$ . They are in thermal equilibrium at temperature  $T$ . The pertinent Hamiltonian reads  $\mathcal{H} = \sum_{i=1}^N p_i^2/2m$ , where  $m$  is the mass of the particles and  $p_i$  the concomitant momenta, with  $i = 1, \dots, N$  [5,17]. The associated canonical partition function adopts the appearance [5,17]

$$Z_N = \frac{1}{N!} \left( \frac{V}{\lambda^3} \right)^N. \quad (40)$$

$\lambda = h/\sqrt{2\pi mkT}$  stands for the particles' mean thermal wavelength [5]. The Helmholtz free energy  $F$  can be cast as [5]

$$F = NkT \left[ \ln \left( n \lambda^3 \right) - 1 \right], \quad (41)$$

with  $n = N/V$  and  $v = V/N$  the molar density and volume per particle [5]. The classical entropy is provided by the well known Sackur-Tetrode equation that reads [5]

$$S = Nk \ln \left[ e^{5/2} \left( n \lambda^3 \right)^{-1} \right], \quad (42)$$

which is positive definite if we fulfill the requirement  $n \lambda^3 \ll e^{5/2}$ .

### 6.1. The connection S-D

According to Ref. [17] one has for  $D$  and the complexity  $C$  per particle

$$D = n \lambda^3 e^{-1} 2^{-3/2}, \quad (43)$$

and then

$$C = n \lambda^3 e^{-1} 2^{-3/2} \ln \left[ e^{5/2} \left( n \lambda^3 \right)^{-1} \right]. \quad (44)$$

Remark that  $\lambda$  is the thermal de Broglie wavelength, that is, roughly, the mean de Broglie wavelength of the gas molecules at the temperature  $T$ . The mean inter-particle spacing  $d_T$  is, approximately,  $v^{1/3}$ . Whenever  $\lambda$  is much smaller than this  $d_T$ , the gas can be regarded as classical. Contrarily, when  $\lambda$  is of the order of or larger than  $d_T$ , quantum effects should dominate and the gas ought to be treated as quantal.

Notice that, on the basis of the above relations, both  $S$  and  $C$  become negative for  $n \lambda^3 > e^{5/2}$  which it involves  $D > (e/2)^{3/2} \sim 1.59$ . If we agree in wishing to avoid this negativity, then there is a maximum degree of order  $D = 1.59$ , an important result. Clearly, for  $D = 1.59$  the complexity  $C = 0$ . A simple manipulation yields the mean energy  $u = U/N$  for  $D = 1.59$  as

$$u_{\text{minimum}} = \frac{6\pi\hbar^2}{mv^{2/3}e^{5/3}}, \quad (45)$$

a very small but non-zero quantity. Thus, in these circumstances we can speak of a maximum degree of order for which the entropy vanishes, an order related Nernst law.

## 7. Conclusions

What this paper offers is a choice regarding the classical entropy's  $S$  nature. We can either accept the possibility of having  $S < 0$ , as most people seem to do, or reject it. If we reject it, then the degree of order exhibits a maximum, at which  $S = 0$ , an order-based Nernst law.

Based on statistical mechanics considerations revolving on the notion of purity and its quantum counterpart (both symbolized by the same letter  $D$ ), we have developed some interesting, and hopefully new, classical traits of the harmonic oscillator and of the ideal gas. They derive from the fact that a complexity measure can not be a negative quantity [16]. This could perhaps might also be said of  $S$ . Our protagonist role is placed by the notions of purity-disequilibrium  $D$ .

On such a basis we discovered that, in both the classical HO and ideal gas statistics, the traits described below became apparent.

- Quantum purity is an indicator of how different is a quantum state from the totally mixed one. Its classical counterpart is called disequilibrium and is an indicator of how different is the classical probability distribution from the uniform one. We are talking of the same idea as expressed in two quite distinct scenarios. Notably, the same mathematical expression, that we call  $D$ , can be used in the two contexts.
- $D$  has a maximum possible value. In the quantum case, this maximum value is unity. Classically, we saw in two examples that there is a maximum value  $D_{max} = (e/2)^3$  for the HO and  $D_{max} = (e/2)^{3/2}$  for the ideal gas.
- We conjecture that, classically, the maximum value of  $D$ , that can be thought to represent maximum order, is equal to  $(e/2)^{f/2}$ ,  $f$  being the number of degrees of freedom of the system. For a 3-dimensional HO  $f = 6$  and for the ideal gas  $f = 3$ .
- For such maximum order values, the classical HO entropy and the ideal gas one vanish at a temperature  $T_0$  (with  $T_0 \neq 0$ ), a kind of order based third law that applies not at zero temperature but a finite one.
- There exist minimum possible classical mean energies (HO and ideal gas) values. This seems to be a new and surprising result.
- Either the classical HO or the ideal gas entropies are negative for a  $D > D_{maximum}$  situation. Such situation would clearly be a contradictory one because  $D$  can not be larger than  $D_{maximum}$ . Thus, negative entropies should be unacceptable.
- If we reject the possibility of negative entropies, this fact by itself impedes the classical temperature to reach the zero value, as we already know by appeal to conventional treatments.

**Author Contributions:** All authors participated in equal fashion in the research and planning of this paper. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding

**Data Availability Statement:** Everything that might be needed is here.

**Acknowledgments:** We thank Conicet (Argentine Agency).

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. F. Pennini, A. Plastino, Thermal effects in quantum phase-space distributions, *Phys. Lett. A* **23**, 374 (2010) 1927.
2. D. Giulini, et al., *Decoherence and the Appearance of a Classical World in Quantum Theory*, Springer, Berlin, 1996; M. Namiki, S. Pascazio, H. Nakazato, *Decoherence and Quantum Measurements*, World Scientific, Singapore, 1997.

3. P. Facchi, A. Mariano, S. Pascazio, *Decoherence versus entropy in neutron interferometry*, Phys. Rev. A **63** (2001) 052108;
4. S. Watanabe, Z. Phys. **113** (1939) 482; W.H. Furry, Boulder Lectures in Theoretical Physics, vol. 8A, University of Colorado Press, 1966.
5. R.K. Pathria, Statistical Mechanics, 2nd., Butterworth-Heinemann, Oxford, UK, 1996.
6. R. López-Ruiz, H. Mancini, X. Calbet, A statistical measure of complexity. Phys. Lett. A **209** (1995) 321-326.
7. R. López-Ruiz, Complexity in some physical systems, International Journal of Bifurcation and Chaos **11** (2001) 2669-2673.
8. M.T. Martin, A. Plastino, O.A. Rosso, Statistical complexity and disequilibrium, Phys. Lett. A **311** (2003) 126-132.
9. L. Rudnicki, I.V. Toranzo, P. Sánchez-Moreno, J.S. Dehesa, Monotone measures of statistical complexity, Phys. Lett. A **380** (2016) 377-380.
10. R. López-Ruiz, A Statistical Measure of Complexity in Concepts and recent advances in generalized information measures and statistics, A. Kowalski, R. Rossignoli, E.M.C. Curado (Eds.), Bentham Science Books, pp. 147-168, New York, 2013.
11. K.D. Sen (Editor), Statistical Complexity, Applications in electronic structure, Springer, Berlin 2011.
12. M.T. Martin, A. Plastino, O.A. Rosso, Generalized statistical complexity measures: Geometrical and analytical properties, Physica A **369** (2006) 439-462.
13. C. Anteneodo, A.R. Plastino, Some features of the López-Ruiz-Mancini-Calbet (LMC) statistical measure of complexity, Phys. Lett. A **223** (1996) 348-354.
14. R. Piasecki, A. Plastino, Entropic descriptor of a complex behaviour, Physica A **389** (2010) 397-407.
15. F. Pennini, A. Plastino, A.R. Plastino, and A. Hernando, Structural Statistical Quantifiers and Thermal Features of Quantum Systems, Entropy **23** (2021) 1-19.
16. R. López-Ruiz, H. Mancini, X. Calbet, A statistical measure of complexity. ArXiv 1009.1498 (2010).
17. R. Branada, F. Pennini, A. Plastino, Physica A **511** (2018) 18-26.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.