

Article

Not peer-reviewed version

The Utility of 4D-CT Imaging in Primary Hyperparathyroidism Management in a Low Volume Center

[Marko Murruste](#)^{*}, Martin Kivilo, Karri Kase, Ülle Kirsimägi, Annika Tähepõld, Kaia Tammiksaar

Posted Date: 17 July 2023

doi: 10.20944/preprints2023071038.v1

Keywords: Primary hyperparathyroidism; parathyroid imaging; parathyroidectomy; four-dimensional computed tomography



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The Utility of 4D-CT Imaging in Primary Hyperparathyroidism Management in a Low Volume Center

Marko Murruste ^{1,*}, Martin Kivilo ², Karri Kase ², Ülle Kirsimägi ¹, Annika Tähepõld ¹ and Kaia Tammiksaar ¹

¹ Tartu University Hospital

² University of Tartu

* Correspondence: marko.murruste@kliinikum.ee

Abstract: Background: Parathyroid imaging techniques are used in patients with biochemically confirmed primary hyperparathyroidism (pHPT) to detect the localization of enlarged parathyroid gland(s) (PTG) and thereby to allow more focused, unilateral or minimally invasive parathyroidectomy (PTX), instead of classical bilateral neck exploration (BNE). Although the preferred sequence of imaging continues to evolve, ultrasonography (US) is usually the first line modality, followed by ^{99m}Tc sestamibi parathyroid scan (SPS). However, both tests may have suboptimal localizing accuracy with considerable rate of non-localized cases. Therefore, more accurate four-dimensional computed tomography scan (4D-CT) has been employed for PTG imaging. Currently, there is a paucity of data evaluating the utility of 4D-CT in low case-load settings. Aim: To evaluate the impact of PTG imaging, using 4D-CT in conjunction with its intraoperatively displayed results, on the outcomes of surgical PTX. Our primary outcome was sensitivity of 4D-CT in comparison to US and SPS for preoperative PTG adenoma localization. The secondary outcomes were the need for redo surgery and morbidity. Additionally, the characteristics of surgical PTX, i.e. operative time, number of removed parathyroid glands, and rate of simultaneous thyroid surgery, were assessed. Methods: A single-center retrospective analysis of surgically treated patients with pHPT from 01/2010 to 01/2021 was conducted. Evaluation of impact of the preoperative imaging modalities on the results of surgical treatment was carried out. Results: During the study period 290 PTX were performed, 45 cases were excluded due to surgery for secondary, tertiary or recurrent HPT, or due to use of alternative imaging techniques; the remaining 245 patients were included in the study. US was carried out for PTG imaging in 236 (96.3%), SPS in 93 (38.0%), and 4D-CT in 52 patients (21.2%). The use of 4D-CT was associated with a significantly higher rate of successful localization of enlarged PTG (49 cases, 94.2%) compared to US and SPS (74 cases, 31.4%, and 54 cases, 58.1%, respectively). We distinguished between three groups of patients based on preoperative imaging: 1) PTG lateralization via US or SPS in 106 (43.3%) cases, 2) precise localization of PTG via 4D-CT in 49 (20.0%) patients, and 3) in 90 cases (36.7%) PTG imaging failed to localize enlarged gland. The group of 4D-CT localization had significantly shorter operative time, lower rate of simultaneous thyroid resections, as well as lower rate of removal of ≥ 2 PTG, compared to the other groups. The 4D-CT imaging was also associated with the lowest perioperative morbidity and with the lowest median PTH in one month follow-up; however, compared to the other groups, these differences were statistically not significant. Implementation of 4D-CT (since 01/2018) decreased the need for redo surgery (from 11.5% to 7.3%), and significantly increased the annual case load of PTX at our institution (from 15.3 to 41.0), compared to the period before 4D-CT diagnostics. Conclusions: 4D-CT imaging enabled to precisely locate almost 95% of enlarged PTG in patients with pHPT. Accurate localization and intraoperatively displayed imaging results are useful guides for surgeons in making PTX a faster and safer procedure in a low volume center.

1. Introduction

Primary hyperparathyroidism (pHPT) is an increasingly common endocrine disorder, reaching a prevalence of 0.3% in the general European population [1,2]. Furthermore, in the Scandinavian countries, the prevalence of pHPT has reached 2%-5% in peri- and postmenopausal women [3]. The wider availability of blood calcium screening tests and ultrasonography (US) has probably led to this increase in the recent decades [4]. As surgical treatment is the only definitive cure for pHPT, a parallel increase in the prevalence of parathyroidectomies (PTX) has been seen in the past decades [5,6].

When compared to conservative management, PTX is a relatively safe and cost-effective procedure [7]. Cure rates of surgical PTX exceed 95% at centers with high expertise [8]. Although this procedure is associated with a relatively low risk of complications, the ultimate goal of surgery – normal function of the remaining parathyroid glands (PTG) manifesting as persistent normocalcemia, is not always met. This is often due to difficulties in preoperative and intraoperative localization of pathologic PTG, leading to persistence of pHPT and the need for redo surgery [9]. Although the need for preoperative PTG imaging has been a subject of numerous debates, it has been demonstrated that preoperative localization of enlarged PTG facilitates intraoperative exploration of the culprit gland. What's more, PTG imaging also has the potential to prevent a large number of bilateral neck explorations (BNE) and to promote unilateral neck exploration or minimally invasive PTX instead [10]. Therefore, current guidelines recommend preoperative localization of hyperfunctioning PTG for the selection of suspect PTX candidates, as it allows for a more focused approach, reduces operation time, complications, and results in a better cosmetic outcome and greater patient satisfaction [11,12]. Although the preferred sequence of imaging continues to evolve, US is usually the first line modality, followed by ^{99m}Tc sestamibi parathyroid scan with subtraction imaging (SPS). However, both tests may have suboptimal localizing accuracy with considerable rate of failed localizations. Therefore, more accurate four-dimensional computed tomography scan (4D-CT) has been suggested for PTG imaging.

Another aspect affecting successful PTX and biochemical cure is surgeons' experience. Multiple studies have reported better outcomes of parathyroid surgery in high volume centers, however no definite volume threshold has been established [13]. Although some studies have proposed a threshold of 20 PTX per year, this number is purely based on professional opinion, a so called 'pragmatic and achievable target' [14,15].

The above described problems were also seen in our center. We had a relatively high percentage of patients with failed preoperative localization of PTG, and our annual case-load was relatively low (ca 10-15 cases of surgically treated patients with pHPT) [16]. Thus, in 2018 we attempted to improve PTX success rate by implementing 4D-CT imaging for localization of pathologically enlarged PTG from 2018. The aim of the present study was to evaluate the impact 4D-CT imaging on the outcomes of surgical PTX in a low volume PTX center.

2. Methods

Ethics. The present study is a single-center retrospective analysis of surgically treated patients with pHPT. The Research Ethics Committee of the University of Tartu approved this clinical research (protocol 300/T-4).

Patients. All adult patients aged 18 years or older, who were operated due to pHPT at the Department of Surgery of Tartu University Hospital between 01/2010 and 01/2021, and in whom US, SPS, or 4D-CT for PTG imaging was used, were included in the study. The exclusion criteria were surgery for secondary, tertiary or recurrent HPT. We also excluded a few patients whose PTG imaging investigations were magnetic resonance imaging (MRI) or conventional computed tomography (CT).

Aim of the present study was to evaluate the impact of using 4D-CT as PTG imaging modality on the outcomes of surgical PTX.

Our primary outcome was sensitivity of 4D-CT in comparison to US and SPS for preoperative PTG adenoma localization. The secondary outcomes were the need for redo surgery and morbidity.

Additionally, the characteristics of surgical PTX, i.e. operative time, number of removed parathyroid glands, and rate of simultaneous thyroid surgery, were assessed.

We hypothesized that the success rate of 4D-CT in detecting enlarged PTG is higher compared to US and SPS, which results in better outcomes of surgical treatment. Furthermore, we hoped that implementation of 4D-CT will make PTX outcomes comparable to large volume centers.

PTG imaging. In the present study the routine management of patients with suspected pHPT consisted of preoperative biochemical confirmation of pHPT, followed by localization investigations, including US (performed by endocrinologists) and SPS in cases of inconclusive US imaging. Since 2018, we have replaced SPS with 4D-CT imaging and SPS was only used in cases where 4D-CT was contraindicated (e.g. renal insufficiency, intolerance of iodine based contrast media). To facilitate better intraoperative localization of enlarged PTG, the results of 4D-CT imaging (precise localization of suspected pathological PTG) were always displayed intraoperatively using the axial, coronal and/or sagittal CT-planes.

Surgical treatment consisted of a small Kocher incision (usually up to 5 cm) for access, followed by the classical medial approach to explore the thyroid and parathyroid regions, mobilizing the strap muscles from their midline position by progressive lateral mobilization.

For patients with failed preoperative localization or with multiglandular PTG involvement, a classical BNE was carried out. In the case of successful lateralization of enlarged PTG using US or SPS, a unilateral neck exploration was employed. For patients with precise preoperative localization of pathologic PTG via 4D-CT imaging, a one-region dissection in the suspected area was undertaken. In the case of failed focused exploration of PTG, conversion to classical BNE was employed.

Database. The study database comprises data about surgical treatment (operative time, number of removed PTG, simultaneous thyroid surgery, complications, and length of stay), and biochemical characteristics before and after surgical treatment (serum calcium, ionized calcium, parathyroid hormone (PTH), 25-hydroxyvitamin D, phosphate, and creatinine levels). Biochemical follow-up data were obtained one month and one year after surgical treatment.

Persistence of pHPT was defined as presence of one of the following: reoperation for pHPT during 2 years of follow-up, or ionized calcium levels above 1.32 mmol/L with synchronous PTH levels over 6.9 pmol/L.

Data analysis. The impact of the preoperative imaging modalities on the results of surgical treatment were evaluated in three groups: 1) PTG localization via US or SPS, as these modalities enabled to detect the affected side ('lateralization'), but precise three-dimensional localization was not employed, 2) precise localization of PTG via 4D-CT, and 3) cases where preoperative PTG localization was not successful ('image negative patients').

Additional assessment of the results was carried out in two time frames: before implementation of 4D-CT (from 01/2010 to 12/2017) and after it (01/2018 to 01/2021).

Statistical analysis. All perioperative and follow-up data were entered in a computerized database (Microsoft Access 2016). The software package Statistica version 13.3 (TIBCO Software, California, USA) was used for statistical calculations.

The main characteristics are presented as medians (with 25% and 75% percentiles) in the case of non-normal distribution of variables; dichotomous variables are reported as counts and percentages. Comparison between the study groups was made using the following tests: Fisher's exact test in the case of percentages, non-parametric Mann-Whitney U-test in the case of medians for non-normally distributed variables. The level of statistical significance was set at $p < .05$.

3. Results

Patients. During the study period 290 parathyroid operations were carried out, 32 cases were excluded due to surgery for secondary, tertiary or recurrent HPT, and 13 cases due to PTG imaging via MRI or conventional CT. The remaining 245 patients, who were operated for pHPT and whose imaging investigations were US, SPS or 4D-CT, were included in the study (Fig. 1).

Preoperative data. The median age of the patients at the time of surgical treatment was 66 years; there were 218 female (89.0%) and 27 male patients (Table 1). The preoperative biochemical

characteristics were: median ionized calcium level 1.51 mmol/L; PTH 15.0 pmol/L; and phosphate 0.86 mmol/L.

Table 1. Preoperative characteristics of the patients.

Characteristic	Patients, n=245
Age (y)	66 (58-74)
Female gender, n (%)	218 (89.0%)
iCa*	1.51 (1.44-1.61)
PTH [#]	15.0 (10.4-25.3)
Phosphate ^{##}	0.86 (0.73-0.95)
Parathyroid imaging:	
- US, n (%)	236 (96.3)
- successful localization, n (%)	74 (31.4)
- SPS, n (%)	93 (38.0)
- successful localization, n (%)	54 (58.1)
- 4D-CT, n (%)	52 (21.2)
- successful localization, n (%)	49 (94.2)

Data are presented as median (with interquartile range) or percentages
 Y years; iCa ionized calcium; PTH parathyroid hormone; US ultrasonography; SPS sestamibi parathyroid scan; 4D-CT four-dimensional computed tomography
 *Reference value 1.16-1.32 mmol/L, #Reference value 1.6-6.9 pmol/L, ##Reference value 0.81-1.45 mmol/L

PTG imaging. For PTG imaging, US was carried out in 236 (96.3%, Fig. 2), SPS in 93 (38.0%), and 4D-CT in 52 patients (21.2%) since its implementation in 2018. There occurred no complications associated with the use of 4D-CT imaging (e.g. development or exacerbation of renal insufficiency; intolerance of intravenous iodine based contrast media etc.). However, for 6 patients 4D-CT was contraindicated due to underlying renal insufficiency or known allergy to iodine based contrast media.

The use of 4D-CT was associated with a significantly higher rate of successful localization of enlarged PTG (49 cases, 94.2%) compared to US and SPS (74 cases, 31.4%; and 54 cases, 58.1%, respectively).

Results of surgical treatment.

According to the preoperative best knowledge of PTG localization, three groups of patients were distinguished: 1) PTG lateralization (via UH or SPS), 2) precise PTG localization (via 4D-CT), and 3) patients with non-successful localization of PTG. There were 106 patients (43.3%) in the first group, 49 in the second (20.0%), and 90 (36.7%) in the third (Table 2).

Table 2. Comparison of operative and follow-up characteristics of the patients according to the results of preoperative imaging.

	Localized via US or SPS (106)	Localized via 4D-CT (49)	Non-localized PTG (90)	p-value
Operative characteristics:				
- Operative time (min)	53 (36-71)	35 (24-56)	51 (41-64)	0.004
- Removed \geq 2 PTG, n (%)	25 (23.6)	7 (14.3)	32 (35.6)	0.018
- True multiglandular disease ^{&} , n (%)	17 (16.0)	5 (10.2)	16 (17.8)	0.505
- PTG size (mm)	15 (10-20)	13 (10-19)	15 (8-20)	0.728
- Simultaneous thyroid surgery, n (%)	30 (28.3)	14 (28.6)	47 (52.2)	0.001
Complications, n (%)	14 (13.2)	1 (2.0)	9 (10.0)	0.076
- Wound hematoma, n	0	0	3	-

- RLN injury, n	3	1	1	-
- Subjective dysphonia, n	5	0	0	-
- Other	6	0	5	-
Length of stay (d)	2 (2-2)	2 (2-2)	2 (2-2)	0.272
Follow-up (30 day):				
- iCa*	1.29 (1.23-1.35)	1.25 (1.22-1.32)	1.29 (1.23-1.33)	0.318
- PTH [#]	7.1 (5.1-12.3)	5.6 (4.3-7.6)	6.2 (3.8-11.0)	0.082
Follow-up (1 year):				
- iCa*	1.27 (1.23-1.30)	1.25 (1.20-1.32)	1.26 (1.22-1.33)	0.798
- PTH [#]	5.4 (4.5-9.9)	5.7 (3.8-8.3)	5.4 (3.7-9.5)	0.735
Redo surgery, n (%)	13 (12.3)	3 (6.1)	7 (7.8)	0.467

Data are presented as median (with interquartile range) or percentages
 US ultrasonography; SPS sestamibi parathyroid scan; 4D-CT four-dimensional computed tomography; PTG parathyroid gland; RLN recurrent laryngeal nerve; iCa ionized calcium; PTH parathyroid hormone
 & Histologically proven multiglandular parathyroid gland involvement, *Reference value 1.16-1.32 mmol/L, [#]Reference value 1.6-6.9 pmol/L, [°]Reference value 0.81-1.45 mmol/L

More precise localization of PTG with the use of 4D-CT led to significantly shorter median operative time, being 35 min (Fig. 3), compared to 51 min if PTG localization was not successful, and 53 min after localization via US/SPS.

Assessment of the characteristics of PTX revealed a significantly higher proportion of patients in the group of non-localized PTG, undergoing removal of ≥ 2 PTG (35.6%), compared to 23.6% and 14.3% in the US/SPS and 4D-CT groups, respectively. While the rate of true histologically proven multiglandular disease was similar between the groups, it is obvious that in the remaining cases 'innocent' PTG were removed.

Unsuccessful preoperative localization of PTG was also associated with a significantly higher rate of simultaneous thyroid resections (52.2%) compared to US/SPS (28.3%) and 4D-CT (28.6%) groups.

4D-CT imaged patients showed the lowest complications rate, at 2.0%, compared to 13.2% and 10.0% for the US/SPS imaged patients and the patients with failed preoperative imaging, respectively. However, the difference between the groups remained statistically non-significant ($p=0.076$).

Short-term biochemical follow-up (one month after surgery) revealed the lowest PTH and ionized calcium levels in the group of 4D-CT imaged cases. Nonetheless, compared to the US-SPS imaged and non-localized cases, the difference between the groups was not significant ($p=0.082$ for PTH, and $p=0.318$ for ionized calcium).

Persistence of pHPT was diagnosed in three patients (6.1%) of the 4D-CT group, and in 13 (12.3%) and 7 patients (7.8%) of the US/SPS and the non-localized groups, respectively. A reoperation for persistence of hyperparathyroidism was undertaken in most cases during the first year after initial surgery. As a result of successful reoperation, one-year biochemical follow-up demonstrated similar ionized calcium and PTH levels in all three groups.

Next we evaluated the prevalence of complications and failure to cure as a combined metric of negative outcome suggested by Ullmann et al [9]. There was no significant difference between the groups. Yet the lowest rate of one or both negative events occurred in the group of 4D-CT localized PTG, being 8.2%, compared to 23.6% and 16.7% in the groups of UH/SPS localized and of non-localized patients, respectively.

Additionally, analysis of the results for two time frames was performed. During the first period (from 2010 to 2017) PTG was localized via US or SPS; the second period (from 2018 to 2020) started after the implementation of 4D-CT diagnostics, which was indicated for patients with inconclusive UH results. This analysis revealed two important effects of implementation of 4D-CT (Fig. 4). The need for redo surgery fell from 11.5% during the first period to 7.3% during the second period, while

the annual case-load of PTX almost tripled at our institution: from 15.3 cases per year for the first period to 41 cases per year for the second period.

4. Discussion

The present retrospective study reports an impact of implementation of 4D-CT imaging of PTG on the results of surgical treatment of pHPT in a low volume center.

The need for PTG imaging has been a subject of debate over decades. It has been suggested repeatedly that an experienced parathyroid surgeon is capable of localizing a vast majority of parathyroid glands without imaging assistance [17]. It has even been stated that the greatest challenge in preoperative localization of PTG adenoma is locating an experienced parathyroid surgeon [12]. However, as parathyroid surgery has to be performed also at low or medium case-load institutions, expert surgeons are not always available [18]. Things are further complicated, as this experience can only be gained by parathyroid surgery. It has been demonstrated that the surgical experience gained from non-parathyroid surgery in the same anatomical region (e.g. thyroid surgery) does not improve outcomes in parathyroid surgery. For example, centers performing large number of thyroid but low number of parathyroid surgical procedures have a significantly higher rate of reoperative parathyroid surgery [19]. So, other methods need to compliment the surgeon's experience.

Preoperative imaging is a good way to compensate for the experience cap in lower volume centers. Frank et al demonstrated that negative preoperative imaging correlates with reduced success rates of surgical exploration and higher need for redo surgery [20]. An even more convincing argument for the use of extended localization diagnostics is that focused dissection in the area of enlarged PTG is associated with reduced morbidity [21]. Therefore, as preoperative precise anatomical localization of enlarged PTG facilitates intraoperative exploration of the culprit gland, PTG imaging has the potential to prevent a large number of BNE, as well as to promote unilateral neck exploration or minimally invasive PTX instead [10].

Although the preferred sequence of imaging continues to evolve and there exists significant institutional variation, a common approach is that following neck US, SPS is performed to confirm the identification of enlarged PTG, or to exclude an ectopic localization before surgical treatment is intended [22]. A meta-analysis assessing the value of US imaging in pHPT reported an overall pooled sensitivity of 76.1% and a positive predictive value of 93.2%, however, significant operator and center dependency was emphasized [23]. According to the common sequence, US is followed by SPS imaging [24]. Nevertheless, despite the wide availability of SPS, several studies have reported great variation in its diagnostic accuracy, with reported sensitivity from 25.4% for US-negative patients to 78.9% for single-gland disease [10,25]. Hence, the quality of both imaging investigations and their interpretation vary dramatically between institutions and both common localizing tests (US and SPS) may have suboptimal localizing accuracy with considerable rate of non-localized cases [26].

In the present study, endocrinologists performed all US tests, however, the rate of successful localization of enlarged PTG was relatively low, while the results of SPS diagnostics were only slightly better. This was a cause of a high rate of cases with failed preoperative imaging of enlarged PTG. Therefore, we implemented 4D-CT scan for preoperative PTG imaging in 2018 at our institution. Owing to this, we achieved a 94.2% success rate of detecting enlarged PTG. Even more importantly, there occurred several favorable changes from the clinical point of view. After the implementation of 4D-CT imaging the need for redo surgery decreased from 11.5% to 7.3%. We also saw a move towards less extensive and more focused surgical dissection. This was illustrated by shorter operative times of PTX; lower number of patients undergoing removal of two or more PTG; lower rate of simultaneous thyroid surgery; and lower morbidity. However, the last difference remained statistically non-significant.

Shorter operative time for PTX in the 4D-CT imaged patients was expected, as precise preoperative localization of enlarged PTG enabled us to use unilateral focused exploration instead of BNE in most cases. We mostly used a small (4-5 cm) central Kocher's skin incision, followed by a unilateral dissection on the side of enlarged PTG. The potential benefit of using Kocher's incision is the easy conversion from unilateral dissection to classical BNE in the case of failed one-side

exploration. However, according to consensus, PTX is minimally invasive when performed through an incision of less than 3 cm. So, our approach has to be defined as non-minimally invasive [27,28]. Despite this, advances in preoperative imaging certainly promote the use of minimally invasive PTX with lower complication rate and better cosmetic effect compared to BNE [29].

An additional tool facilitating a more focused dissection and better guidance of surgery was the intraoperative display of the results of 4D-CT PTG imaging. However, this statement is based on professional experience.

We believe that significantly higher rate of extirpation of two or more PTG in the group of preoperatively non-localized PTG was caused by difficulties of intraoperative localization of culprit PTG, especially in cases of small adenomas. Exploration of a slightly enlarged PTG resulted in its extirpation. If the macroscopic changes were 'not convincing', or the frozen section did not confirm the diagnosis of PTG adenoma, further explorative dissection and extirpation of a second PTG, whether with minimal or more 'convincing' changes, was undertaken. As the rate of true histologically proven multiglandular disease for the groups was similar, we could conclude that the use of 4D-CT enables to reduce the rate of removal of 'innocent PTG'.

The uncertain location of PTG probably led to the almost twofold rate of simultaneous thyroid surgery (52.2%) in patients with failed preoperative imaging of the PTG pathology, compared to cases of successfully localized enlarged PTG (28.3%-28.6%). The higher rate of concurrent thyroidectomies might have been due to the removal of the protruding thyroid tissue, especially in the retrothyroideal position, mistaken for PTG adenomas. Ryan et al reported a rate of 25% for simultaneous thyroid surgery, and they performed thyroid resection to optimize operative access, or to ensure complete removal of involved PTG without the risk of seeding the parathyroid tissue, or more commonly, to remove synchronous thyroid pathology [30]. Thus, the optimal rate for a concurrent thyroid surgery seems to be one in every four cases of PTX. In our study we only saw similar rates for image-positive patients. However, as the rate of simultaneous thyroid surgery reported in other studies has been relatively high as well (up to 50%), we believe that this is a place for further research and discussion [25].

After the implementation of 4D-CT, the observed decrease in morbidity was most probably attributable to the less extensive surgical exploration, and discarding routine BNE. Although the difference in morbidity between the groups was obvious (2.0% for 4D-CT, 13.2% for US or SPS localized, and 10.0% for non-localized cases), it remained on the verge of statistical significance ($p=0.076$). Nevertheless, we share the opinion that surgeons should not perform blind exploration, as positive imaging investigation can almost always be obtained to make the surgery safer [24].

There are several other diagnostic options suggested to improve the results of surgical treatment of pHPT. Positron emission tomography (PET) with ^{11}C -choline has shown outstanding results with an accuracy of almost 99% [31]. However, despite its excellent accuracy, PET has considerable limitations mostly related to its cost and availability [32]. In the present study intraoperative diagnostic tests (e.g. PTH measurement and frozen section consultation) were used relatively rarely, as they are deemed unacceptable at our institution for being too time consuming. However, we are looking forward to implementing rapid intraoperative PTH assay, as encouraged by a systematic review by Medas et al reporting a significantly lower rate of persistent and recurrent pHPT with the use of this test [33]. This technique eliminates the need for intraoperative frozen section consultation, thereby saving time and expense [24]. In summary, we think that other novel diagnostic modalities should be used in concurrence with 4D-CT in low volume centers.

This study had some limitations. First, the retrospective nature of this study means that there were some missing data points. Second, from a personal experience and epidemiological data, we believe that PTH is underdiagnosed in Estonia. These cases are usually asymptomatic, which means that the adenoma at fault could be smaller and harder to localize. So, if more attention is drawn to PTH and its diagnosis improves, the benefits of 4D-CT can become more evident.

5. Conclusions

4D-CT imaging enabled to precisely locate almost 95% of enlarged PTG in patients with pHPT. Accurate localization and intraoperatively displayed imaging results are useful guides for surgeons in making PTX a faster and safer procedure in a low volume center.

Author Contributions: MM, MK, KK, ÜK, AT and KT conceptualized the study. MM prepared all the documents. MK led data collection process. MM drafted the manuscript, assisted by MK, KK and AT. ÜK performed statistical analysis. All the authors critically revised the manuscript and approved the final manuscript.

Funding: None.

Ethics Approval: Approval for the study was obtained from the Research Ethics Committee of the University of Tartu (number 300-T4).

Consent for Publication: Not applicable.

Availability of Data and Materials: Study data can be made available on a reasonable request for future collaborative analyses.

Competing interests: No competing interests.

List of Abbreviations

BNE	bilateral neck exploration
CT	computed tomography
HPT	hyperparathyroidism
iCa	ionized calcium
MRI	magnetic resonance imaging
pHPT	primary hyperparathyroidism
PET	positron emission tomography
PTG	parathyroid gland
PTH	parathyroid hormone
PTX	parathyroidectomy
RLN	recurrent laryngeal nerve
SPS	^{99m} Tc sestamibi parathyroid scan
US	ultrasonography
4D-CT	four-dimensional computed tomography

References

1. Minisola S, Arnold A, Belaya Z, et al. Epidemiology, Pathophysiology, and Genetics of Primary Hyperparathyroidism. *J Bone Miner Res.* 2022;37(11):2315-2329. doi:10.1002/jbmr.4665
2. Donatini G, Marciniak C, Lenne X, et al. Risk Factors of Redo Surgery After Unilateral Focused Parathyroidectomy: Conclusions From a Comprehensive Nationwide Database of 13,247 Interventions Over 6 Years. *Ann Surg.* 2020;272(5):801-806. doi:10.1097/SLA.0000000000004269
3. Bollerslev J, Schalin-Jääntti C, Rejnmark L, et al. Unmet therapeutic, educational and scientific needs in parathyroid disorders: Consensus statement from the first European Society of Endocrinology Workshop (PARAT). *Eur J Endocrinol.* 2019;181(3):P1-P19. doi:10.1530/EJE-19-0316
4. Clarke BL. Asymptomatic primary hyperparathyroidism. *Parathyroid Disord.* 2019;51:13-22.
5. Wilhelm SM, Wang TS, Ruan DT, et al. The American association of endocrine surgeons guidelines for definitive management of primary hyperparathyroidism. *JAMA Surg.* 2016;151(10):959-968. doi:10.1001/jamasurg.2016.2310
6. Khokar AM, Kuchta KM, Moo-Young TA, Winchester DJ, Prinz RA. Parathyroidectomy is Safe in Elderly Patients: A National Surgical Quality Improvement Program Study. *World J Surg.* 2020;44(2):526-536. doi:10.1007/s00268-019-05280-9

7. Zanocco K, Heller M, Sturgeon C. Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism. *Endocr Pract.* 2011;17:69-74.
8. Wilhelm SM, Wang TS, Ruan DT, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg.* 2016;151(10):959-968. doi:10.1001/jamasurg.2016.2310
9. Ullmann TM, Adam MA, Sosa JA. Surgeon Volume and Outcomes in Primary Hyperparathyroidism—What Is Old Is New Again. *JAMA Surg.* 2022;157(7):589.
10. Naik M, Khan SR, Owusu D, et al. Contemporary Multimodality Imaging of Primary Hyperparathyroidism. *Radiographics.* 2022;42(3):841-860. doi:10.1148/rg.210170
11. Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg.* 2011;253(3):585-591. doi:10.1097/SLA.0b013e318208fed9
12. Bilezikian JP, Potts JT, El-Hajj Fuleihan G, et al. Summary statement from a workshop on asymptomatic primary hyperparathyroidism: A perspective for the 21st century. *J Clin Endocrinol Metab.* 2002;87(12):5353-5361. doi:10.1210/jc.2002-021370
13. Erinjeri NJ, Udelsman R. Volume–outcome relationship in parathyroid surgery. *Best Pract Res Clin Endocrinol Metab.* 2019;33(5):101287. doi:10.1016/j.beem.2019.06.003
14. Gray WK, Navaratnam A V, Day J, Wass JAH, Briggs TWR, Lansdown M. Volume-outcome associations for parathyroid surgery in England: Analysis of an administrative data set for the Getting It Right First Time program. *JAMA Surg.* 2022;157(7):581-588.
15. Sosa JA, Powe NR, Levine MA, Udelsman R, Zeiger MA. Profile of a clinical practice: Thresholds for surgery and surgical outcomes for patients with primary hyperparathyroidism: A national survey of endocrine surgeons. *J Clin Endocrinol Metab.* 1998;83(8):2658-2665. doi:10.1210/jc.83.8.2658
16. Nikkolo C, Saar S, Sokirjanski M, Junkin LK, Lepner U. Retrospektiivne uuring primaarse hüperparatüreosi kirurgilise ravi tulemustest Tartu Ülikooli Kliinikumis. *Eesti Arst.* 2014;93(11):622-626.
17. Kettle AG, O'Doherty MJ. Parathyroid Imaging: How Good Is It and How Should It Be Done? *Semin Nucl Med.* 2006;36(3):206-211. doi:10.1053/j.semnuclmed.2006.03.003
18. Iacobone M, Scerrino G, Palazzo FF. Parathyroid surgery: an evidence-based volume—outcomes analysis: European Society of Endocrine Surgeons (ESES) positional statement. *Langenbeck's Arch Surg.* 2019;404(8):919-927. doi:10.1007/s00423-019-01823-9
19. Mitchell J, Milas M, Barbosa G, Sutton J, Berber E, Siperstein A. Avoidable reoperations for thyroid and parathyroid surgery: Effect of hospital volume. *Surgery.* 2008;144(6):899-907. doi:10.1016/j.surg.2008.07.022
20. Frank E, Watson WA, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope.* 2020;130(12):E963-E969. doi:10.1002/lary.28566
21. Minisola S, Cipriani C, Diacinti D, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-D8.
22. Morris MA, Saboury B, Ahlman M, et al. Parathyroid Imaging: Past, Present, and Future. *Front Endocrinol (Lausanne).* 2022;12(February):1-23. doi:10.3389/fendo.2021.760419
23. Cheung K, Wang TS, Farrokhyar F, Roman SA, Sosa JA. A meta-analysis of preoperative localization techniques for patients with primary hyperparathyroidism. *Ann Surg Oncol.* 2012;19(2):577-583. doi:10.1245/s10434-011-1870-5
24. Udelsman R. Approach to the patient with persistent or recurrent primary hyperparathyroidism. *J Clin Endocrinol Metab.* 2011;96(10):2950-2958. doi:10.1210/jc.2011-1010
25. Lenschow C, Wennmann A, Hendricks A, et al. Questionable value of [99mTc]-sestamibi scintigraphy in patients with pHPT and negative ultrasound. *Langenbeck's Arch Surg.* 2022;407(8):3661-3669. doi:10.1007/s00423-022-02648-9
26. Pappu S, Donovan P, Cheng D, Udelsman R. Sestamibi scans are not all created equally. *Arch Surg.* 2005;140(4):383-386. doi:10.1001/archsurg.140.4.383
27. Brunaud L, Zarnegar R, Wada N, Ituarte P, Clark OH, Duh QY. Incision length for standard thyroidectomy and parathyroidectomy: when is it minimally invasive? *Arch Surg.* 2003;138(10):1140-1143.
28. Bellantone R, Raffaelli M, Crea C De, Traini E, Lombardi CP. Minimally-invasive parathyroid surgery. Published online 2011:207-215.
29. Ishii H, Mihai R, Watkinson JC, Kim DS. Systematic review of cure and recurrence rates following minimally invasive parathyroidectomy. *BJS Open.* 2018;2(6):364-370. doi:10.1002/bjs5.77
30. Ryan S, Courtney D, Moriariu J, Timon C. Surgical management of primary hyperparathyroidism. *Eur Arch Oto-Rhino-Laryngology.* 2017;274(12):4225-4232. doi:10.1007/s00405-017-4776-4
31. Orevi M, Freedman N, Mishani E, Bocher M, Jacobson O, Krausz Y. Localization of parathyroid adenoma by 11C-choline PET/CT: preliminary results. *Clin Nucl Med.* 2014;39(12):1033-1038.
32. Tay D, Das JP, Yeh R. Preoperative localization for primary hyperparathyroidism: A clinical review. *Biomedicines.* 2021;9(4). doi:10.3390/biomedicines9040390

33. Medas F, Cappellacci F, Canu GL, Noordzij JP, Erdas E, Calò PG. The role of Rapid Intraoperative Parathyroid Hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis. *Int J Surg.* 2021;92(July). doi:10.1016/j.ijssu.2021.106042

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.