

Article

Not peer-reviewed version

Validation and Evaluation of the Psychometric Properties of an Obstetric Violence Scale

[Héctor González-de la Torre](#)*, Paula Nikola González-Artero, Daniel Muñoz de León-Ortega, María Reyes Lancha-de la Cruz, [José Verdú-Soriano](#)

Posted Date: 28 August 2023

doi: 10.20944/preprints202308.1854.v1

Keywords: obstetrics labor; obstetric violence; surveys and questionnaires; validation Studies as topic



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Validation and Evaluation of the Psychometric Properties of an Obstetric Violence Scale

Héctor González-de la Torre ^{1,2,*}, Paula Nikola González-Artero ³,
Daniel Muñoz de León-Ortega ³, María Reyes Lancha-de la Cruz ⁴ and José Verdú-Soriano ⁵

¹ Department of Nursing. University of Las Palmas de Gran Canaria. Edificio Ciencias de la Salud. C/ Blas Cabrera Felipe s/n. CP:35016, Las Palmas de Gran Canaria-Canary Islands, Spain.

hector.gonzalez@ulpgc.es

² Research Support Unit of Insular Maternal and Child University Hospital Complex of Gran Canaria, Canary Health Service. Avda Marítima del Sur S/N.CP:35016, Las Palmas de Gran Canaria-Canary Islands, Spain. hgontor@gobiernodecanarias.org

³ Department of Obstetrics and Gynaecology. Insular Maternal and Child University Hospital Complex of Gran Canaria-Canary Health Service. Avda Marítima del Sur S/N.CP:35016, Las Palmas de Gran Canaria-Canary Islands, Spain.

⁴ Delivery room service. General Hospital of Fuerteventura Virgen de la Peña-Canary Health Service. Carretera del aeropuerto, Km 1.CP:35600 - Puerto del Rosario, Canary Islands, Spain.

⁵ Department of Community Nursing, Preventive Medicine, Public Health and History of Science, Faculty of Health Sciences, University of Alicante (UA), Alicante, Spain.pepe.verdu@ua.es

* Correspondence: hector.gonzalez@ulpgc.es; hegontor@gmail.com; Tel.: +34-928448880

Abstract: Obstetric Violence refers to dehumanized or derogative treatment of women in their pregnancy, childbirth or postpartum periods and may be manifested through the attitudes of health professionals or the performance of unjustified or outdated practices without maternal consent. Currently, there is no tool validated in Spain to measure women's perception of obstetric violence. The objective of this study was to carry out the cultural adaptation and validation of an existing 14-item Obstetric Violence Scale to the Spanish context, and to evaluate its psychometric properties. The research was conducted in two phases: first, a methodological study designed to evaluate content validity, through assessments by 8 experts (calculating the Aiken V coefficient) and face validity in a sample of 20 women; second, a cross-sectional study to evaluate construct validity (through confirmatory factor analysis and Rasch analysis), divergent validity against a scale of birth satisfaction, known groups validity and reliability. In Phase 1, Aiken V values higher than 0.71 were obtained for all items. Phase 2 was conducted on a sample of 256 women and the fit values for the unidimensional model were RMSEA: 0.070 [95%CI: 0.059-0.105] and GFI: 0.982 [95%CI: 0.823-0.990]. The Rasch analysis indicated poor performance of item 2, which was removed. The Omega and Cronbach's Alpha coefficients were 0.863 and 0.860, respectively. A final 13-item version of the Obstetric Violence scale was produced, with total score ranging from 0 (no obstetric violence perception) to 52 (maximum obstetric violence perception). The Obstetric Violence Scale is a reliable and useful tool to measure women's perception of obstetric violence.

Keywords: obstetrics labor; obstetric violence; surveys and questionnaires; validation studies as topic

1. Introduction

The choice of an adequate term to designate derogative or inhumane practices and attitudes towards women in their pregnancy, childbirth or post-partum periods is a much debated and controversial issue [1,2]. In the English-speaking countries like the United States, such practices are often called "mistreatment at childbirth", "disrespect" or "abuse" [1,3–5]. In other countries, the term "obstetric violence" is preferred [1,6,7].

According to the World Health Organization (WHO), obstetric violence (OV) "is defined as a specific form of violence from health professionals (mainly doctors and nursing staff) towards pregnant women, during childbirth or puerperium. It is a violation of women's reproductive and

sexual rights” [8]. Several forms of obstetric violence are recognized, including five highlighted categories, which are used in legal definitions: routine and unnecessary interventions or medication (on the mother or the infant); verbal abuse, humiliation or physical abuse; lack of suitable equipment or facilities; conducting practices without maternal informed consent (i.e., consent after receiving complete, truthful and sufficient information), and discrimination on cultural, economic, religious or ethnic grounds[8].

OV also includes any practice or attitude affecting a woman’s psychological wellbeing, e.g. treating her as a child, adopting paternalistic or authoritarian attitudes, behaving in a derogatory way, humiliating or insulting her, etc [1,3,6,7,9].

Both terms, maltreatment-abuse and OV, share the idea of violence based on gender inequality and effects of a patriarchal biomedical system, which usually denies women their autonomy and control over their maternal process [5,6,10].

OV negatively affects the physical, psychological and emotional health of women who experience it, their relatives and the health professionals that witness it. Excessive, unnecessary or unjustified interventions may often harm women’s health [11]. Performing episiotomy in the absence of fetal risk or without maternal consent is a clear example of this. Episiotomy has been associated with chronic pain and dyspareunia, while it has not been proven to prevent severe perineal trauma [12–14]. Therefore, performing routine episiotomy is unjustified and can be considered OV [11,12]. Further examples include obstetric techniques or practices not supported by scientific evidence, like the Kristelle or the Hamilton maneuvers, or excessive vaginal examinations [11,15].

Besides physical harm, there are well documented psychological and emotional aftereffects of OV [15,16]. Victims may experience emotional alterations such as feelings of loneliness or isolation, stress or insecurity [15]. Occasionally, OV causes shame and directly harms women’s self-image and body perception [15]. Such effects may impair women’s sexual and affective dimensions, what makes OV a form of sexual violence [15,17].

Post-traumatic stress disorder is another OV consequence affecting the psycho-emotional dimension. Several studies describe high post-traumatic stress prevalence after childbirth, with rates from 1-6% to 35%[18,19], one of the main risk factors being traumatic experiences during labor due to health providers’ actions or attitudes[20]. This condition also affects women’s partners and children and has an impact on family relationships [21]. Women experiencing verbal or psycho-affective OV are also at higher risk of post-partum depression [22].

OV is a worldwide phenomenon [8,10,17]. Bohren et al [17] published a review of 65 studies from 34 countries in all continents. Despite considerable heterogeneity, they found that the childbirth experiences of women around the world were often stained with OV. However, the prevalence figures vary largely in different countries [1,23–25]. Reported prevalence of mistreat or abuse to women during labor range from 11% in Mexico [26] or 49.4% in Latin America [27] to more than 70% in some African countries [28–30]. High OV prevalence rates have also been reported in Europe [1,23,25].

Analyzing the differences in OV prevalence must take into account the complexity of the construct [2,4,31] which, as mentioned, includes several attitudes and practices deeply influenced by social or cultural aspects, as well as by medical factors[6,7]. Certain practices considered as OV, may be not perceived as such by women or health professionals in certain contexts, which consequently influences the reported rates [2,6,9,32].

The lack of tools specifically validated to measure OV is a further related problem. For example, OV has been evaluated with the Norvold Abuse Questionnaire (Nor-AQ) [33], which is not a specific tool for this type of violence. Recently, a more specific tool has been developed, the Students’ Perceptions of Respectful Maternity Care Scale [34,35], which was validated for some countries [36], although it fails to collect some OV-related aspects.

The PercOV-S Questionnaire (PercOV-S) was designed and validated in Spain for measuring OV as perceived by students of nursery, midwifery and medicine [37]. This tool, which was validated with a sample of 169 students, includes 33 items grouped into two domains: “invisible OV practices” and “visible OV practices”, scored on a Likert-type scale. The PercOV-S shows high reliability and

internal validity and has been used in several studies [38,39]. However, all these tools have been designed for healthcare staff and do not evaluate women's OV perception.

There is a specific scale to evaluate OV as perceived by women, called *Escala de Violencia Obstétrica* (Obstetric Violence Scale), developed by Cárdenas and Salinero [40] based on the *Test de violencia obstétrica* (Obstetric violence test) developed by the association *El Parto es Nuestro* [41]. This test was adapted as a scale to measure women's OV perception based on their memories of certain aspects and situations during labor, which are considered related to OV. It is a unidimensional 14-item tool, of which some psychometric properties have been evaluated (reliability and construct validity) [40]. Items are scored from 1 (It does not describe what happened to me at all) to 5 (This is definitely what happened to me). The scale was validated with a sample of 367 Chilean women in the Valparaíso region (Chile)[40].

Given the lack of tools to measure OV perception from the women's point of view and since the above-mentioned tool has not been adapted to the Spanish context, the objective of this study was to carry out a cultural adaptation and validation of the Obstetric Violence Scale developed by Cárdenas and Salinero [40] to the Spanish context and to evaluate its psychometric properties.

2. Materials and Methods

The research was conducted in 2 phases:

Phase 1: content validation through expert judgement, assessment of face validity through a pilot study on a target population and assessment of the understandability of the OV scale.

Phase 2: assessment of construct validity in a sample through factorial analysis and Rasch analysis, divergent validity against a tool that measures satisfaction with labor, assessment of the scale's reliability (internal validity) and known groups validity.

2.1. Phase 1

In the first phase, a methodological study was conducted to evaluate content validity and face validity of the OV scale proposed by Cárdenas and Salinero [40].

2.1.1. Content validity

Through assessment by a panel of 8 experts from different areas. Experts were selected with the aim of gathering a range of professionals with different views of the studied issue. The experts assessed the pertinence-relevance of every item (whether an item actually evaluated what it was intended to evaluate and the importance of that item in the studied construct). These criteria were expressed in scores from 1 (not pertinent-relevant item) to 4 (very pertinent-relevant item). Based on the experts' scores, a content validity index was calculated for every item (CVI-i), using the Aiken test with the corresponding 95% confidence intervals (95%CI) [42]. This coefficient ranges between 0 and 1 with values closer to 1 indicating better agreement between the judges, which is considered better content validity [42].

2.1.2. Face validity

A pilot test (pre-test) was carried out in a target population to assess the understandability and acceptability of the scale. This pilot test was carried out in a sample of 20 puerperal women, selected by non-probabilistic convenience sampling at the discretion of the research team, provided that they fulfilled the study's inclusion criteria (older than 18 years, in her first 6 weeks postpartum; last delivery in week 37-42 of pregnancy, live newborn). Participants were asked to quantitatively evaluate the acceptability of items by scoring them from 1 to 4 (1 = poorly acceptable/poorly related to OV; 4 = very acceptable/very related to OV), and the mean acceptability scores were calculated for every item. Additionally, the degree of agreement was calculated for each item as the percentage of participants who scored that item with 3 or 4 over the total of women. Mean scores over 2 plus agreement over 80% were considered adequate for an item to be acceptable. Participants were also

asked to assess the understandability of the scale and to suggest changes if they deemed it appropriate.

Additionally, the final version of the scale was evaluated with INFLESZ [43], a tool validated in the Spanish context, used to evaluate text readability and ease of reading by users of healthcare services. It is based on the Szigriszt Pazos' Perspicuity Formula and establishes the following classification system: 0 to 40, very difficult; 40 to 55, moderately difficult; 55 to 65, average difficulty; 65 to 80, fairly easy and 80 to 100, very easy [43].

2.2. Phase 2

2.2.1. Design

Cross-sectional study designed to obtain a sample for assessing construct validity (through factorial analysis and Rasch analysis), calculation of reliability, evaluation of divergent validity against a scale of satisfaction with labor and known groups validation.

2.2.2. Study population

The study population included puerperal women from the province of XXXXX, who visited the office of their primary care midwife in the first 6 weeks postpartum. The inclusion criteria were: older than 18 years, in her first 6 weeks postpartum; last delivery in week 37-42 of pregnancy, live newborn. The exclusion criteria were: inability to read/understand the scale, home birth and/or last delivery by planned caesarean section. Withdrawal criteria were established as: incorrect completion of the questionnaire and/or participant's desire to leave the study.

Eligible participants had given birth at one of the following 4 centers: Center 1 (a public tertiary level hospital and provincial reference center for high-risk births), Center 2 (a private hospital), Center 3 and Center 4 (regional public hospitals for low-risk pregnancies).

2.2.3. Sampling and sample size

Participants were selected through a non-probability convenience sampling procedure. The Cárdenas and Salinero Obstetric Violence Scale include 14 items [40]. According to the classical theory of factor analysis (FA), there must be at least 10-15 subjects per item in the analyzed tool [44,45]. Moreover, since using a polychoric correlation matrix requires a minimum of 200 subjects [44], a minimum sample size of 200 women was estimated for this phase of the study, to allow Confirmatory Factor Analysis (CFA).

2.2.4. Data collection tool and variables of the study

A self-administered physical questionnaire was prepared, with a first part collecting several sociodemographic and obstetric variables and a second part containing the Obstetric Violence Scale developed by Cárdenas and Salinero [40].

The variables collected in the first part were: age, education level (no education, primary school, secondary school, university), number of deliveries, type of last delivery (eutocic vaginal delivery, dystocic vaginal delivery with forceps, dystocic vaginal delivery with vacuum, not-planned cesarean section), center (Center 1, xxxxxx; Center 2 xxxxxx; Center 3 xxxxxxx; Center 4, xxxxxxx), episiotomy (Yes/No), consent to perform episiotomy (Yes/No), artificial rupture of membranes (Yes/No), consent to artificial rupture of membranes (Yes/No), labor induction (Yes/No), consent to labor induction (Yes/No), prohibition of receiving food or drink during labor (Yes/No), epidural analgesia (Yes/No), presentation of a childbirth plan (Yes/No) and observation of the childbirth plan (Yes/No).

The second part of the questionnaire included the Obstetric Violence Scale, a tool designed to assess the occurrence of certain OV-related aspects based on postpartum women's reported memories of their childbirth experience. The 14 items of the scale were scored in a Likert-type format from 1 = "It does not describe what happened to me at all" to 5 = "This definitely happened to me".

Additionally, to assess divergent validity, 20 women were provided with a version of the questionnaire that included a third part, consisting of a tool to measure maternal satisfaction with labor; namely the Spanish version of the Childbirth Experience Questionnaire (CEQ-E) [46]. The CEQ-E, composed of 22 items grouped into 4 domains, has been reported to reach an overall Cronbach's alpha coefficient of 0.88 and was validated and used in the Spanish context [46,47].

2.2.5. Data collection

Data were collected from December 1st, 2022, to March 31st, 2023. Participants were recruited through collaborating midwives in 21 primary care centers of the province of XXXXX. All women attending their first to third postpartum control visit, who met the inclusion criteria, were willing to participate and accepted to take part in the study were offered participation. Once they completed the questionnaires, they handed them to their midwife, who kept them until the end of the data collection process.

2.2.6. Data analysis and interpretation

Data analysis in phase 2 was carried out in the following steps:

-Descriptive analysis of the sample and the items: A descriptive analysis of the studied variables and the involved items was conducted. Qualitative variables were expressed in percentages and frequencies; quantitative variables were expressed in measures of central tendency (mean) and dispersion (standard deviation; minimum-maximum values). Symmetry and kurtosis were calculated for each item on the scale.

-Construct validity by factor analysis: A CFA was carried out based on the unidimensional model proposed by Cárdenas and Salinero [40]. The suitability of data for a FA was assessed with the Kaiser Meyer Olkin index (KMO) and the Bartlett's statistic. KMO values higher than 0.75 were considered adequate and p values ≤ 0.05 were considered statistically significant for the Bartlett's statistic [44].

The suitability of the factorial solution was evaluated through the following indices: Root Mean Square of Residuals (RMSR), Root Mean Square Error of Approximation (RMSEA), Non Normed Fit Index (NNFI), Comparative Fit Index (CFI), Goodness of Fit Index (GFI) and Adjusted Goodness of Fit Index (AGFI). For the RMSR, a value of 0.05 was considered an acceptable fit; for the RMSEA, values below 0.05 were considered a good fit and values between 0.05 and 0.08, a reasonable fit [44]. NNFI and CFI values of 0.95 or higher, and GFI and AGFI values higher than 0.90 were considered to indicate good fit of the model [44].

The items' symmetry and kurtosis values indicated the suitability of using a polychoric correlation matrix for the CFA. Unweighted least squares factor extraction (ULS) and PROMIN oblique rotation were used [48]. The number of factors to be retained was established through a parallel analysis, and the consistency of the retained factors was calculated; 95% confidence intervals were calculated for the item scores and the model measures.

The Generalized G-H index was calculated to evaluate the degree to which the items reflected a common factor. This index is used to measure the highest percentage of factor variation that the items can measure and two characteristics of the factor analysis: a) the quality of the items as indicators, and b) the expected replicability of the solution in different studies. It ranges between 0 and 1 and approaches 1 with increasing magnitudes of factorial load and/or increasing number of items. Values higher than 0.80 are considered to indicate a well-defined latent variable, likely to remain stable in different studies, while low H values suggest a poorly defined latent variable, likely to change in different studies [49]. H-Latent assesses how correctly a factor can be identified through the continuous latent response variables that underlie the observed item scores, while H-Observed assesses how correctly it can be identified from the observed item scores [49].

To evaluate unidimensionality, the Unidimensional Congruence (UniCo), Explained Common Variance (ECV) and Mean of Item RESidual Absolute Loadings (MIREAL) indices were used [50]. UniCo values higher than 0.95, ECV values higher than 0.85 and MIREAL values lower than 0.30 indicated that the data could be considered as essentially unidimensional [50].

-Construct-structural validity by Rasch analysis: After verifying that the model was unidimensional, a Rasch analysis was performed. To that end, the OV scores were adapted from a 1-5 range to a 0-4 range.

To estimate the parameters, the Joint Maximum Likelihood estimation (JMLE) method was used for an Andrich's Rating Scale Model. Items' fit and people's fit were estimated through outfit - Unweighted Mean Square fit statistic (UMS) and infit -Weighted Mean Square Fit statistic (WMS). Fit index values between 0.8 and 1.2 were considered good fit, while values between 0.5 and 1.5 were considered acceptable fit [51].

To establish the quality statistics, the reliability and separation indices were calculated, both for items and people. For people, reliability values higher than 0.8 and separation values higher than 2 are desirable. Item reliability refers to the degree to which items' difficulty can be ordered. Item separation provides information on the quality of item location in the latent variable. The assumption of local independence of items was tested with the Yen's Q3 test [52], which evaluates the correlation matrix of residuals. A graphical assessment was also carried out by making an item-map.

-Reliability: The Omega and Alpha coefficients were calculated, as well as the Bayes expected a posteriori estimation (EAP scores)[50] reliability of the factorial structure.

-Divergent validity: To explore a possible negative correlation between the OV scale and the CEQ-E questionnaire, the Spearman correlation coefficient was used \neg since the normality test (Shapiro-wilk test) showed non-symmetry for the OV scale scores distribution – considering negative correlation for coefficients (r) higher than -0.5. Standardized scores (z-values) were used to compare both tools.

-Final proposed scale and known groups validation: Once a final structure of the OV scale was reached, an inferential analysis was conducted to explore the association between the different variables and the score, by comparing groups of women likely to have experienced OV according to several aspects described in the literature. Since data were asymmetric, as evidenced by the Kolmogorov-Smirnov test, the non-parametric Mann-Whitney U test was used to compare means between two groups and the Kruskal Wallis contrast test, was used to compare means between more than two groups, followed by a post hoc contrast (Dwass-Steel-Critchlow-Fligne test) to identify the group-comparisons that showed the differences; α values ≤ 0.05 were considered statistically significant. The effect size was calculated for each evaluated association using the Hedges' formula (Hedges' g) and the Kelley's epsilon squared measure.

The descriptive and inferential analyses were carried out with the statistical package JAMOVI®v.2.3.24. The FA and model reliability were carried out with the free-access software FACTOR® Release Version 12.02.01x64 bits, and the Rasch analysis was conducted with the J Metrik® Software.

2.2.7. Ethical considerations

This research was approved by the Research Ethics Committee CEIm XXXXXX, registration number 2022-XXX-X. All women willing to participate were given an information sheet, in which the objectives of the study were explained, an informed consent form that they had to sign and a revocation form. Data collection forms were anonymous, without names or identification data. All databases were blinded.

3. Results

This section may be divided by subheadings. It should provide a concise and precise description of the experimental results, their interpretation, as well as the experimental conclusions that can be drawn.

3.1. Phase 1

3.1.1. Content Validity

The professional profile of the selected experts (7 women, 1 man) can be found in Supplementary Material 1. All items received CVI-i values higher than 0.70. Table 1 show the scores assigned to each item by each of the experts, as well as the CVI-i values and their corresponding 95%CI.

Table 1. Scores and content validity index for each item obtained by Aiken's test.

	Expert 1	Expert 2	Expert 3	Expert 4	Expert 5	Expert 6	Expert 7	Expert 8	CVI-i[CI95%]*
<i>Item 1</i>	4	4	4	4	3	4	4	3	0.92 [0.74-0.98]
<i>Item 2</i>	3	4	4	2	3	4	3	2	0.71 [0.51-0.85]
<i>Item 3</i>	3	4	4	1	3	4	4	4	0.79 [0.60-0.91]
<i>Item 4</i>	3	4	4	2	4	4	4	3	0.83 [0.64-0.93]
<i>Item 5</i>	4	4	4	2	4	4	4	3	0.88 [0.69-0.96]
<i>Item 6</i>	4	4	4	4	4	4	4	4	1.00 [0.86-1.00]
<i>Item 7</i>	3	4	4	3	3	4	3	3	0.79 [0.60-0.91]
<i>Item 8</i>	4	4	4	3	3	4	3	3	0.83 [0.74-0.93]
<i>Item 9</i>	4	4	4	3	3	4	4	3	0.88 [0.69-0.96]
<i>Item 10</i>	4	4	4	3	3	4	3	3	0.83 [0.64-0.93]
<i>Item 11</i>	4	4	4	3	4	4	4	3	0.92 [0.74-0.98]
<i>Item 12</i>	3	4	4	4	3	3	3	2	0.75 [0.55-0.88]
<i>Item 13</i>	4	4	4	3	2	3	3	3	0.75[0.55-0.88]
<i>Item 14</i>	4	4	4	3	3	4	4	3	0.88[0.69-0.96]

*Content Validity Index-item [Confidence intervals at 95%].

3.1.2. Face validity and pilot study in a target population

All items received mean scores higher than 3 and degrees of agreement higher than 80%. Item number 12 (Your childbirth care experience made you feel vulnerable, guilty or insecure in any sense) received the lowest mean score and degree of agreement (3.2 and 80%, respectively). The scores assigned by the 20 participants in the pilot study can be found in Supplementary Material Table S2, together with the mean score and degree of agreement of each item of the scale.

Participants did not report difficulties understanding any of the items; therefore, item modifications were not introduced. The perspicuity level for the scale was 64.74 points, which corresponds to a text with a normal level of readability, according to the INFLESZ scale.

3.2. Phase 2

3.2.1. Descriptive analysis of the sample and the items

A total of 271 questionnaires were collected; however, 15 of them were withdrawn due to poor data completion; thus, the sample was finally composed of 256 women (n=256). Their mean age was 31.66 years (SD=5.54) [range 18-46 years]; 159 (62.1%) of them were primiparous, and 97(37.9%) were multiparous (mean number of deliveries 1.32, SD=0.80). Regarding the level of education, 61 (23.8%) women had primary education, 103 (40.2%) secondary education and 92 (35.9%) university studies.

Regarding the type of delivery in last childbirth, 199 women (77.7%) had a normal vaginal delivery, 33 women had an unplanned cesarean section (12.9%), 17 (6.6%) had a dystocic vaginal delivery with forceps and 7 (2.7%), a dystocic delivery with vacuum. Regarding epidural analgesia, 186 women (72.7%) received analgesia while 70 women (27.3%) did not; 64 women (25.0%) were prohibited to take food or drink during childbirth.

Regarding other obstetric variables, 32 women (12.5%) underwent episiotomy, 83 (32.4%) underwent artificial rupture of membranes and 116 (45.3%) underwent labor induction. Regarding consent, 17 women with episiotomy (53.1%), 65 women with artificial rupture of membranes (78.3%) and 99 women with labor induction (84.6%) were asked for consent before the interventions.

Finally, participants were asked if they had presented a birth plan, and if so, if the plan had been observed. Most women, 182 (71.1%), had not presented a birth plan. From the 74 women (28.9%) who did present a plan, 62 (24.2%) reported that it was observed and 12 (4.7%) reported it was not.

Raw scores and response percentages of each item can be found in Supplementary Material S3. Table 2 shows a descriptive analysis of the items, as well as the symmetry and kurtosis values.

Table 2. Descriptive analysis of the items in the OV scale.

Item	M [CI95%]*	SD**	Symmetry***	Kurtosis***
1- Members of the healthcare staff made ironic or derogative comments or made jokes about your behavior.	1.22 [1.09-1.36]	0.85	3.90	13.91
2- You were addressed to with nicknames or diminutives (e.g., mummy, chubby, etc.) or treated as if you were unable to understand the processes you were going through.	1.10 [1.02-1.19]	0.55	5.90	35.06
3- You felt treated as a child or neglected by the staff, as if you were unable to make decisions about what was happening to you before, during or after delivery.	1.25 [1.13-1.39]	0.82	3.55	12.11
4- You were somehow criticized for expressing your emotions (cry, scream of pain, etc.) during labor or delivery.	1.24 [1.10-1.38]	0.86	3.70	12.54
5- It was impossible for you to ask queries or express your fears or concerns because nobody answered, or they answered in a bad way.	1.18 [1.07-1.31]	0.74	4.34	18.20
6- You were subjected to medical procedures without asking your consent or without explaining why such procedures were needed.	1.37 [1.21-1.54]	1.03	2.79	6.47
7- At the moment of delivery, you were compelled to keep lying on your back despite you expressed you discomfort with that position.	1.27 [1.13-1.42]	0.90	3.32	9.83
8- You were compelled to stay in bed and prevented from walking or seeking the position you needed.	1.21 [1.09-1.35]	0.79	3.88	14.26

9- You were not allowed to be accompanied by someone you trusted in.	1.15 [1.04-1.27]	0.71	4.75	21.46
10- You were prevented from having immediate contact with your newborn, before the doctor took him/her away (caressing, holding him/her in your arms, etc.).	1.21 [1.09-1.35]	0.83	3.83	13.36
11- After delivery, they make you feel you had not behaved up to what was expected of you (that you had not "helped").	1.08 [1.00-1.16]	0.48	6.55	44.21
12- Your childbirth care experience made you feel vulnerable, guilty or insecure in any sense.	1.23 [1.11-1.37]	0.80	3.59	12.22
13- After delivery, you were denied the opportunity to use a birth control device or procedure (IUD, tubal ligation, etc.).	1.09 [1.01-1.17]	0.50	6.17	39.37
14- During or after labor, you felt exposed to the gaze of other people unknown to you (exposure to strangers).	1.46 [1.28-1.65]	1.16	2.41	4.24

*Mean [95% confidence interval], **Standard deviation, ***Polychoric correlation is recommended for univariate item distributions that are asymmetric or with excessive kurtosis. In case both indices are lower than one, Pearson's correlation is recommended.

3.2.2. Construct validity by factor analysis

A CFA was performed based on the proposed 14-item unifactorial model. The KMO values and the Bartlett's statistic indicated an adequate fit of the sample (KMO=0.824, 95%CI: 0.824-0.888; Bartlett= $p \leq 0.001$). The one-factor solution showed 60.72% explained variance, with the parallel analysis indicating a single-factor solution. Fit values for this model were RMSEA=0.070 (95%CI: 0.059-0.105), NNFI=0.985 (95%CI: 0.898-0.990), IFC=0.987 (95%CI: 0.914-0.991), GFI=0.982 (95%CI: 0.823-0.990) and AGFI=0.979 (95%CI: 0.791-0.989). The Root Mean Square of Residuals (RMSR) yielded a value of 0.085 (95%CI: 0.041-0.121). According to the Kelley's criteria for an acceptable model, the expected RMSR value in this model was 0.062.

Table 3 shows the factorial loads after model rotation and the 95%CIs. All items showed loads over 0.450, although the confidence intervals were very wide, and 3 items (numbers 2, 7 and 13) showed loads lower than 0.350 in their confidence intervals.

Table 3. Unrotated loading matrix with factor loadings of the one-dimensional model for OV scale.

Items	Factor 1	IC95%*
1- Members of the healthcare staff made ironic or derogative comments or made jokes about your behavior.	0.721	[0.354-0.878]
2- You were addressed to with nicknames or diminutives (e.g., mummy, chubby, etc.) or treated as if you were unable to understand the processes you were going through.	0.645	[0.299-0.892]
3- You felt treated as a child or neglected by the staff, as if you were unable to make decisions about what was happening to you before, during or after delivery.	0.842	[0.594-0.963]
4- You were somehow criticized for expressing your emotions (cry, scream of pain, etc.) during labor or delivery.	0.748	[0.387-0.912]
5- It was impossible for you to ask queries or express your fears or concerns because nobody answered, or they answered in a bad way.	0.884	[0.470-1.000]
6- You were subjected to medical procedures without asking your consent or without explaining why such procedures were needed.	0.745	[0.364-0.868]
7- At the moment of delivery, you were compelled to keep lying on your back despite you expressed you discomfort with that position.	0.486	[0.188-0.739]

8- You were compelled to stay in bed and prevented from walking or seeking the position you needed.	0.790	[0.377-0.918]
9- You were not allowed to be accompanied by someone you trusted in.	0.695	[0.334-0.898]
10- You were prevented from having immediate contact with your newborn, before the doctor took him/her away (caressing, holding him/her in your arms, etc.).	0.771	[0.375-0.911]
11- After delivery, they make you feel you had not behaved up to what was expected of you (that you had not "helped").	0.891	[0.491-1.000]
12- Your childbirth care experience made you feel vulnerable, guilty or insecure in any sense.	0.957	[0.734-1.000]
13- After delivery, you were denied the opportunity to use a birth control device or procedure (IUD, tubal ligation, etc.).	0.473	[-0.489-0.764]
14- During or after labor, you felt exposed to the gaze of other people unknown to you (exposure to strangers).	0.755	[0.460-0.949]

*95% confidence interval.

The results of the analysis showed 0.969 H-latent (95%CI: 0.939-1.000) and 0.513 observed H (95%CI: 0.223-0.743) for a single factor. The unidimensionality analysis showed the following results: UniCo=0.984 (95%CI: 0.957-0.996), ECV=0.900 (95%CI: 0.850-0.943), and MIREAL=0.190 (95%CI: 0.140-0.232). These results supported the model unidimensionality for the scale.

3.2.3. Construct-structural validity by Rasch analysis

Given that the factor analysis supported a unidimensional solution, a Rasch analysis was additionally performed to complement the construct validity of the scale. This required adapting the scores to a 0-4 range (0="It does not describe what happened to me at all", 1 = "I'm not sure but I believe/feel that this didn't happen to me", 2 = "I'm not sure", 3 = "I'm not sure but I believe/feel that this happened to me", 4 = "This definitely happened to me.").

An initial analysis revealed poor performance of item number 2, which indicated the convenience or removing it from the matrix (Supplementary Material 4). A new analysis without item 2 was carried out, and the infit-(WMS) and outfit-UMS values were obtained (Table 4). Infit-(WMS) values indicated good fit for all items except numbers 5 and 12, which presented acceptable fit. Outfit-UMS values showed acceptable fit for items 5 and 8 and bad fit for items 11 and 12.


Table 4. Fit values of the items according to the Joint Maximum Likelihood estimation method in the Rasch analysis.

Item	Difficulty Index*	Infit-WMS**	Outfit-UMS**
1-Members of the healthcare staff made ironic or derogative comments or made jokes about your behavior.	-0.09	1.09	1.17
3-You felt treated as a child or neglected by the staff, as if you were unable to make decisions about what was happening to you before, during or after delivery.	-0.08	0.85	0.83
4-You were somehow criticized for expressing your emotions (cry, scream of pain, etc.) during labor or delivery.	-0.11	1.18	1.39
5-It was impossible for you to ask queries or express your fears or concerns because nobody answered, or they answered in a bad way.	0.02	0.77	0.58
6-You were subjected to medical procedures without asking your consent or without explaining why such procedures were needed.	-0.32	1.08	1.06

7-At the moment of delivery, you were compelled to keep lying on your back despite you expressed you discomfort with that position.	-0.11	1.06	1.08
8-You were compelled to stay in bed and prevented from walking or seeking the position you needed.	-0.00	0.93	0.76
9-You were not allowed to be accompanied by someone you trusted in.	0.10	1.16	1.16
10-You were prevented from having immediate contact with your newborn, before the doctor took him/her away (caressing, holding him/her in your arms, etc.).	-0.03	1.02	0.95
11-After delivery, they make you feel you had not behaved up to what was expected of you (that you had not "helped").	0.54	0.86	0.37
12-Your childbirth care experience made you feel vulnerable, guilty or insecure in any sense.	0.02	0.58	0.45
13-After delivery, you were denied the opportunity to use a birth control device or procedure (IUD, tubal ligation, etc.).	0.53	1.09	1.13
14-During or after labor, you felt exposed to the gaze of other people unknown to you (exposure to strangers).	-0.46	1.21	1.17

* Difficulty index: Indicates in this case the highest values of violence. **Outfit-Unweighted Mean Square fit statistic (UMS) and infit - Weighted Mean Square Fit Statistic (WMS). Fit index values between 0.8 and 1.2 meant a good fit and values between 0.5 and 1.5 meant an acceptable fit.

Graph 1 shows a person-item map of the scale. Regarding the quality statistics, the reliability values of items and people were 0.841 and 0.262 respectively, while the separation index values were 2.300 and 0.596, respectively, which indicated acceptable reliability for the items but not for the people. Regarding the assumption of local independence (Yen Q3 test), most values in the correlation matrix were below 0.2-0.3, so that local independence of the items was considered valid (Supplementary Material Table S5).




Graph 1. Person-item map of Violence Obstetric Scale.

3.2.4. Reliability

Based on the FA, the EAP was 0.969, with a Factor Determinacy Index of 0.984; the Omega and Cronbach's Alpha coefficients were 0.863 (95%CI: 0.839-0.888) and 0.860 (95%CI: 0.834-0.883), respectively.

3.2.5. Divergent validity

The mean total score of the CEQ-E questionnaire – completed by 20 women – was 2.91 (SD=0.47). The Spearman's correlation coefficient was $r = -0.794$ ($p < 0.001$), evidencing a negative correlation between the total score of the OV scale and the total score of the CEQ-E questionnaire. Such correlation is illustrated in the scatter plot in Graph 2 (the higher the OV scores, the lower the satisfaction scores).



Graph 2. Divergence graph between CEQ-E scores and the Violence Obstetric Scale scores.

3.2.6. Final proposed scale and known groups validation

A final version of OV scale was proposed, with 13 items (original item 2 was removed) to be scored between 0 and 4, where 0 = "It does not describe what happened to me at all", 1 = "I'm not sure but I believe/feel that this did not happen to me", 2 = "I'm not sure", 3 = "I'm not sure but I think/feel that this did happen to me", 4 = "This definitely happened to me". Since the total OV score is calculated by adding individual item-scores, it ranges from 0 (no OV perception) to 52 (maximum OV perception). The mean score recorded in the sample for the final version of the scale was 3.03 (SD=6.55; minimum score=0; maximum score=52). The final scale obtained in Spanish language can be consulted in Supplementary Material S6.

For known groups validation, the association between certain variables and the scale's score was analyzed and the effect size was measured for each inference. Statistically significant differences were found for all associations (with varying effect size magnitudes), except for the association "undergoing labor induction" and "presenting a birth plan", which showed no differences related to the OV scores. Table 5 shows all the inferences for the bivariate variables.

Table 5. Mean, standard deviation, p-value and effect size for every inference considered in the known-group validation.

Variables	M(SD)*	p-value**	Effect size***
Parity			
<i>Primiparous (n=159)</i>	3.59 (7.39)		
<i>Multiparous (n=97)</i>	2.11 (4.75)	0.040**	0.23
Episiotomy			
<i>No (n=224)</i>	2.63 (5.97)		
<i>Yes (n=32)</i>	5.84 (9.32)	0.012**	0.50
You were asked consent for episiotomy			
<i>No (n=15)</i>	8.93 (9.87)		
<i>Yes (n=17)</i>	3.12 (8.13)	0.010**	0.65

Artificial rupture of membranes (ARM)			
<i>No (n=173)</i>	2.38 (6.43)		
<i>Yes (n=83)</i>	4.40 (6.62)	≤0.001**	0.31
You were asked consent for ARM			
<i>No (n=18)</i>	9.78 (7.26)		
<i>Yes (n=65)</i>	2.91 (5.63)	≤0.001**	1.14
Inducing labor			
<i>No (n=140)</i>	2.81 (6.80)		
<i>Yes (n=116)</i>	3.29 (6.24)	0.195	0.07
You were asked consent for inducing labor			
<i>No (n=18)</i>	6.83 (7.23)		
<i>Yes (n=99)</i>	2.70 (5.85)	≤0.001**	0.68
Prohibition of receiving food			
<i>No (n=192)</i>	2.74 (6.52)		
<i>Yes (n=64)</i>	3.89 (6.61)	0.018**	0.18
Epidural analgesia			
<i>No (n=70)</i>	1.56 (3.84)		
<i>Yes (n=186)</i>	3.59 (7.24)	0.006**	0.31
Presentation of a childbirth plan			
<i>No (n=182)</i>	2.40 (5.75)		
<i>Yes (n=74)</i>	4.58 (8.03)	0.066	0.34
Your childbirth plan was observed			
<i>No (n=12)</i>	12.42 (12.93)		
<i>Yes (n=62)</i>	3.04 (5.70)	≤0.001**	1.28

*Mean (standard deviation). ** Statistically significant $p \leq 0.05$ (Mann-Whitney's U-test). *** Effect size according to Hedges (Hedges's g): it considers both groups' variances and sizes. Values < 0.2 indicate small effects, 0.5 indicates medium effect and 0.8, large effect.

Finally, in the comparison of means with more than two groups (using the Kruskal-Wallis contrast test followed by a post-hoc contrast), no statistically significant differences were found for the education level or the center. However, differences were significant for the type of delivery, with the differences occurring between normal vaginal delivery versus unplanned cesarean section, in the post hoc tests ($p = 0.001$) (Supplementary Material Table S7).

4. Discussion

Although OV is an increasingly debated topic [6,53,54], there is currently no adequately validated tool to specifically measure women's perception of OV. With the aim of solving this problem in our context, this study was focused on the validation of a perceived-OV measurement tool, evaluating as many as possible psychometric properties with an as robust as possible methodological approach. We preferred to use an already developed tool and to improve certain methodological aspects of the original study by Cardenas and Salinero [40].

In phase 1, face validity evaluation, most items were well accepted and no one was considered difficult to understand. Additionally, content validity was analyzed by a group of 8 experts from different related disciplines (obstetricians, midwives and nurses). Aiken's V coefficients above 0.75 were obtained for all items except number 2, with a value of 0.71 (although lower values were found for some items in their confidence intervals).

The interpretation of Aiken's V is not free of controversy [55]. While some authors consider that values over 0.5 are acceptable, most of them only accept values over 0.70 [56]. Given that confidence intervals are very sensitive to the sample size [55], recruiting more experts may be a solution. Even so, the recorded values provided useful information for making certain decisions in the tool validation process.

The fit values recorded for the model indicated an acceptable fit. Cardenas and Salinero reported RMSEA, Tucker Lewis Index (TLI; also known as Non-Normed Fit Index, NNFI) and CFI, but not GFI, in their study [40]. Today, using indices that evaluate different aspects and avoiding to report redundant information is recommended [44]. Ferrando et al. proposed that indices should be communicated based on three criteria: (a) fit of the solution per se (for example, the Goodness of Fit Index GFI), (b) comparative fit of the proposed solution against the null model of independence (Non-Normed Fit Index, NNFI or Comparative Fit Index, CFI), and (c) relative fit of the model according to its complexity (Root Mean Square Error of Approximation, RMSEA) [44].

In addition, it is currently recommended to report the RMSR value independently of the model and estimated solution [44], which was not reported by Cárdenas and Salinero [40]. This value allows to calculate the Kelley criterion, where the RMSR value is compared with the standard error for 0-correlation in the population, thus allowing to evaluate the suitability of the factorial solution (since if the RMSR is much higher than the expected value, the model should not be considered good)[44,57].

A further aspect to be discussed in the FA of Cárdenas and Salinero [40] is that the authors did not report whether they used linear or nonlinear approximation, which affects the decision of using Pearson or polychoric correlation matrix [44,45,48,58]. Given the marked asymmetry and kurtosis of the scores recorded for the item scores in our questionnaire (higher than those reported by Cardenas and Salinero [40]), it was clear that a polychoric matrix should be used, which entails a more complex model [44,45,48,58].

Furthermore, Cárdenas and Salinero [40] did not indicate whether they performed tests of sample adequacy to a FA, for example by using the KMO or the Bartlett's statistic [45].

Something that has been verified is the unidimensionality of the scale, from the values of the three used indices. This aspect could be analyzed with a RASCH approach. The RASCH approach is based on the classical item response theory (IRT) and allows to assess a tool from two perspectives: the inherent functioning of the scale and the people who complete it [51,59].

After this analysis, item number 2 (You were addressed to with nicknames or diminutives, e.g., "mommy", "chubby", etc., or treated as if you were unable to understand the processes you were going through) was removed from the scale. Such decision was not only based on its poor performance in the analysis, but also on the low scores it received in the content validation by experts. In addition, its low factorial load (less than 0.300) in the confidence interval and its marked asymmetry-kurtosis (also present in the study by Cardenas and Salinero, although to a lesser degree [40]) undoubtedly affected this result. A probable reason for this finding is that some expressions of the item are not used in the context, in which this study was carried out. This finding illustrates the importance of the cultural context and peculiarities in the adaptation and validation of tools in different countries, even within the same language [60].

Using a Rasch analysis involves certain assumptions, such as the unidimensionality of the model [51,61], or the existence of minimum scores of 0 for every item. Therefore, the OV scale scoring system was accordingly adapted to a 0-4 range. Thus, the overall score of the scale ranged from 0, which means no perceived OV, to a maximum of 52, which means maximum perceived OV. In our opinion, such adaptation improves scale interpretation and offers better guidance for the practical use of the tool. Cardenas and Salinero [40] did not establish a clear measurement system nor provided measurement cut-off points for the scale.

Finally, the Rasch approach requires checking the local independence of items, for example, by using the Yen's Q3 test [52]. Traditionally, reference values of 0.2-0.3 have been used; however, there is no homogeneous criterion, since this value depends on the size of the sample, the number of items and the number of responses involved [51,52].

From the usual point of view, the OV scale reliability (internal consistency) analyzed with the Omega and Cronbach's Alpha coefficients was adequate, with values above 0.80 for both coefficients [62], similar to those reported by Cárdenas and Salinero (0.83 and 0.88, respectively)[40].

In this study, the Rasch analysis allowed to evaluate the person-separation reliability, thus describing people separation according to their scoring pattern [61]. Better separation entails more

accurate measurement [61]. The higher the value, the better the separation; usually a minimum value of 2 or higher is used for the index, which indicates that the tool can separate people from at least two strata, for example, low and high capacity [51,61].

The separation indexes lead to the conclusion that reliability was acceptable for the items but not for the people, which suggests that this questionnaire may not have sensitivity enough to measure OV in our context. This finding might be due to different reasons: first, a larger sample may be needed; second, OV may be absent or little prevalent in our context; third, women may be unaware or have a little knowledge of OV, which would still be an invisible practice in our health system. These three possibilities should be addressed in future studies.

The psychometric assessment of the tool suggests that, in contexts with low OV levels, its usefulness for the fine measurement of OV is limited. However, the evaluation of other psychometric properties indicates that it may be actually useful to detect OV. Thus, the evaluation of divergent validity against a scale of satisfaction with labor confirmed that, the greater the satisfaction, the lower the OV perception, which is consistent with the theoretical model.

The results of the known groups validation were also clear. An inferential analysis showed that women undergoing certain interventions perceived more OV, especially if they had not given consent, with statistically significant differences and considerable effect sizes (consent to episiotomy, consent to artificial rupture of membranes and consent to induction of labor). These findings are in line with other studies conducted in our context [11,63] and with the results of Cardenas and Salinero [40].

In this regard, it is important to point out that informed consent (regulated in Spain by Law 41/2002 November 14th, 2002 on patient's autonomy and rights and duties regarding information and clinical documentation [64]) must be requested from all patients before any intervention and not doing so entails OV [4,6,8,9].

A further related aspect concerns the presentation of a birth plan. It was evident that this tool is still underused by women in our context (only 28.9% of the total sample presented a birth plan). It was found that women presenting a birth plan, which was not observed perceived more OV, with statistically significant differences and considerable effect size. This might be due to the fact that those women had more information on the delivery and postpartum processes and were thus able to identify OV-related situations or procedures, especially those considered invisible practices [1,6,31,37].

The use of birth plans is controversial. Some studies indicate that it is associated with greater women's dissatisfaction due to unfulfilled expectations during labor [65,66]. However, the problem may lay in the lack of effective communication between healthcare users and providers. Birth plans may serve as a vehicle for such communication and improve women's satisfaction and feeling of control over the process [67]. The results support their potential importance in the management of OV-related situations.

While in the study by Cárdenas and Salinero, the highest score was assigned to item 5 (It was difficult or impossible for you to ask questions or express your fears or concerns because nobody answered, or they answered in a bad way) [40], in this study it was assigned to item 14 (During or after labor you felt exposed to the gaze of other people unknown to you (exposure to strangers)). In the PercOV-S scale, failure to preserve women's privacy during childbirth is included into the invisibles practices domain of OV [37]; therefore, it could remain unnoticed by health professionals, though not by women, as found in this study and described in other ones [38]. This aspect should be taken into account and policies should be applied to warrant women's privacy during childbirth, especially in centers with educative activities and high turnover of trainees. In the context of OV, intimacy has not been given proper attention as compared to other practices [68,69]; however, it is one of the reasons for many women to seek certain settings to give birth e.g., their homes [70].

It has been described that the hospital setting is directly related to OV in childbirth. A study by Mena de Tudela et al. showed that private healthcare centers were more prone to invasive practices, something directly related to higher perceived OV [11]. However, no significant differences in OV perception were found in our study, neither between public or private centers, nor between centers

with different levels of care. These results should be interpreted with caution though, due to the reduced sample size.

No statistically significant perceived OV differences were found in connection with the education level. However, there was a tendency to higher perceived OV in women with higher education (primary studies $M=1.48$, secondary studies $M=3.06$, university studies $M=4.03$). This finding supports the hypothesis that information is a key factor in women's awareness and perception of OV, although no studies on such relationship have been found in the literature.

Results evidence that some women claim having suffered OV to different degrees, both through practices or techniques applied without prior consent and through disrespectful treatment from health professionals. Having a reliable tool to quantify such phenomenon can help detect and alleviate this problem. Further studies are required to propose cut-off points for this scale. However, any score other than zero should be worrying and would require intervention.

Regarding the limitations of the study, several points should be highlighted. Since the temporal reliability of the tool was not assessed, it cannot be ruled out that women's OV perception is a construct affected by time. Furthermore, analyzing the convergent validity against a different tool could also be of interest. Although, there are hardly tools to specifically assess OV perception, convergent validity could be evaluated against the PercOV-S to investigate similarities or differences between the perceptions of users and health professionals.

Finally, further studies with different populations are required, since the scale reliability was acceptable for the items but not for the women in the sample. It should be taken into account that the studied construct is rather complex [2–5] and involves both women's internal factors (perceptions according to education, knowledge, previous experiences and feelings) and external factors (provided healthcare, which may vary in different contexts, centers, regions, countries and/or involved health professionals). New studies with larger samples and women from other regions are needed to verify good performance of the scale in settings with different OV levels.

5. Conclusions

OV is a frequent problem with serious physical and psychological consequences for women. Despite being a topic of much debate in recent years, there are no specific validated tools to assess women's perception of OV. The Obstetric Violence Scale is a tool with adequate psychometric properties and is reliable and useful to measure women's OV perception, despite some limitations, especially in contexts of low OV. Tools with higher sensitivity are needed to evaluate and measure all OV-related aspects.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Table S1: Professional profiles of the experts participating in the content validation process; Table S2: Scores from the pilot test with 20 postpartum women, means scores and degree of agreement of each item of the scale; Table S3: Frequencies and response percentages of each item of Obstetric Violence Scale; Table S4: First Rasch analysis by Joint Maximum Likelihood estimation (JMLE) with all the items.; Table S5: Correlation matrix. Yen Q3 Test; Table S6: Obstetric Violence Scale (Spanish); Table S7: Means, standard deviations, p-value, contrast statistic and effect size for variables in more than two groups.

Author Contributions: Conceptualization: H.G.-d.l.T., P.N.G-A, D.M.-d.L.O and M.R.L.-d.l.C.; Methodology: H.G.-d.l.T. and J.V,-S; formal analysis: H.G.-d.l.T. and J.V,-S; investigation; P.N.G-A, D.M.-d.L.O and M.R.L.-d.l.C.; writing—original draft preparation: P.N.G-A, D.M.-d.L.O and M.R.L.-d.l.C.; writing—review and editing: H.G.-d.l.T.; Supervision: H.G.-d.l.T. and M.R.L.-d.l.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Research Ethics Committee (CEI/CEIm) Hospital Dr. Negrín, Las Palmas (Code N° 2022-457-1). Date: 21/10/2022.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data used in this research are confidential and are protected in a coded and anonymized database kept by the research group in accordance with Spanish regulations. However, the raw data from the Violence Obstetric Scale (response to each item) and without the rest of the sociodemographic-obstetric variables could be shared with those researchers who contact the corresponding author if requested with a reasoned and logical request.

Public Involvement Statement: No public involvement in any aspect of this research.

Guidelines and Standards Statement: This manuscript was drafted against the (STROBE) for a cross-sectional study, descriptive research (Phase 2) and COSMIN (CONsensus-based Standards for the selection of health Measurement Instruments) guidelines for reporting results of measurement instrument studies.

Acknowledgments: To all the women who participated in this study. To all the midwives who participated in the data collection. To all the experts for participating in the questionnaire validation process. To the Research Support Unit of the Insular Maternal and Child University Hospital Complex of Gran Canaria. To the Multiprofessional Teaching Unit of Obstetrics and Gynecology of the Insular Maternal and Child University Hospital Complex of Gran Canaria.

Conflicts of Interest: The authors declare no conflict of interests.

References

1. Mena-Tudela, D.; Iglesias-Casás, S.; González-Chordá, V.M.; Cervera-Gasch, Á.; Andreu-Pejó, L.; Valero-Chilleron, M.J. Obstetric violence in Spain (Part I): Women's perception and interterritorial differences. *Int J Environ Res Public Health*. 2020,17,1–14. doi: 10.3390/ijerph17217726.
2. Mena-Tudela, D.; Roman, P.; González-Chordá, V.M.; Rodriguez-Arrastia, M.; Gutiérrez-Cascajares, L.; Ropero-Padilla, C. Experiences with obstetric violence among healthcare professionals and students in Spain: A constructivist grounded theory study. *Women Birth*. 2023,36, e219-e226. doi: 10.1016/j.wombi.2022.07.169.
3. Darilek, U.; A Woman's Right to Dignified, Respectful Healthcare During Childbirth: A Review of the Literature on Obstetric Mistreatment. *Issues Ment Health Nurs*. 2018,39,538-541. doi: 10.1080/01612840.2017.1368752.
4. Savage, V.; Castro, A. Measuring mistreatment of women during childbirth: a review of terminology and methodological approaches. *Reprod Health*. 2017,14,138. doi: 10.1186/s12978-017-0403-5
5. Betron, M.L.; McClair, T.L.; Currie, S.; Banerje, J. Expanding the agenda for addressing mistreatment in maternity care: a mapping review and gender analysis. *Reprod Health*. 2018,15,143. doi: 10.1186/s12978-018-0584-6.
6. Rodríguez Mir, J.; Martínez Gandolfi, A.; La violencia obstétrica: una práctica invisibilizada en la atención médica en España [Obstetric violence. A hidden practice in medical care in Spain]. *Gac Sanit*. 2021,35,211-212. doi:10.1016/j.gaceta.2020.06.019.
7. Araujo-Cuauro, J.C. Obstetric violence: a hidden dehumanizing practice, exercised by medical care personnel: Is it a public health and human rights problem?. *Rev Mex Med Forense*. 2019,4,1-11. WHO. The prevention and elimination of disrespect and abuse during facility-based childbirth:
8. WHO statement. World Health Organ [Internet]. 2015,4. Available from: https://apps.who.int/iris/bitstream/handle/10665/134588/WHO_RHR_14.23_cze.pdf
9. Martínez-Galiano, J.M.; Rodríguez-Almagro, J.; Rubio-Álvarez, A.; Ortiz-Esquinas, I.; Ballesta-Castillejos, A.; Hernández-Martínez, A. Obstetric Violence from a Midwife Perspective. *Int J Environ Res Public Health*. 2023,20,4930. doi: 10.3390/ijerph20064930.
10. Faheem, A. The nature of obstetric violence and the organisational context of its manifestation in India: a systematic review. *Sex Reprod Health Matters*. 2021,29,2004634. doi: 10.1080/26410397.2021.2004634.
11. Mena-Tudela, D.; Iglesias-Casás, S.; González-Chordá, V.M.; Cervera-Gasch, Á.; Andreu-Pejó, L.; Valero-Chilleron, M.J. Obstetric Violence in Spain (Part II): Interventionism and Medicalization during Birth. *Int J Environ Res Public Health*. 2020,18,199. doi: 10.3390/ijerph18010199.
12. Jiang, H.; Qian, X.; Carroli, G.; Garner, P. Selective versus routine use of episiotomy for vaginal birth. *Cochrane Database Syst Rev*. 2017,2,CD000081. doi: 10.1002/14651858.CD000081.pub3.
13. Lurie, J.M.; Weidman, A.; Huynh, S.; Delgado, D.; Easthausen, I.; Kaur, G. Painful gynecologic and obstetric complications of female genital mutilation/cutting: A systematic review and meta-analysis. *PLoS Med*. 2020,17,e1003088. doi: 10.1371/journal.pmed.1003088.

14. Pereira, G.M.V.; Hosoume, R.S.; de Castro Monteiro, M.V.; Juliato, C.R.T.; Brito, L.G.O. Selective episiotomy versus no episiotomy for severe perineal trauma: a systematic review with meta-analysis. *Int Urogynecol J.* 2020,31,2291-2299. doi: 10.1007/s00192-020-04308-2.
15. Shabot, S.C. We birth with others: Towards a Beauvoirian understanding of obstetric violence. *Eur J Womens Stud.* 2021, 28, 213–228. <https://doi.org/10.1177/13505068209194>.
16. Sadler, M.; Santos, M.J.; Ruiz-Berdún, D.; Rojas, G.L.; Skoko, E.; Gillen, P.; Clausen, J.A. Moving beyond disrespect and abuse: addressing the structural dimensions of obstetric violence. *Reprod Health Matters.* 2016,24,47-55. doi: 10.1016/j.rhm.2016.04.002.
17. Bohren, M.A.; Vogel, J.P.; Hunter, E.C.; Lutsiv, O.; Makh, S.K.; Souza, J.P.; Aguiar, C.; Saraiva Coneglian, F.; Diniz, A.L.; Tunçalp, Ö.; Javadi, D.; Oladapo, O.T.; Khosla, R.; Hindin, M.J.; Gülmezoglu, A.M. The Mistreatment of Women during Childbirth in Health Facilities Globally: A Mixed-Methods Systematic Review. *PLoS Med.* 2015,12,e1001847. doi: 10.1371/journal.pmed.1001847.
18. Lai, X.; Chen, J.; Li, H.; Zhou, L.; Huang, Q.; Liao, Y.; Krewski, D.; Wen, S.W.; Zhang, L.; Xie, R.H. The incidence of post-traumatic stress disorder following traumatic childbirth: A systematic review and meta-analysis. *Int J Gynaecol Obstet.* 2023,162,211-221. doi: 10.1002/ijgo.14643.
19. Ertan, D.; Hingray, C.; Burlacu, E.; Sterlé, A.; El-Hage, W. Post-traumatic stress disorder following childbirth. *BMC Psychiatry.* 2021,21,155. doi: 10.1186/s12888-021-03158-6.
20. Beck, C.T.; Gable, R.K. A mixed methods study of secondary traumatic stress in labor and delivery nurses. *J Obstet Gynecol Neonatal Nurs.* 2012,41,747-60. doi: 10.1111/j.1552-6909.2012.01386.x.
21. Reshef, S.; Mouadeb, D.; Sela, Y.; Weiniger, F.C.; Freedman, S.A. Childbirth, trauma and family relationships. *Eur J Psychotraumatol.* 2023,14,2157481. doi: 10.1080/20008066.2022.2157481.
22. Martínez-Vázquez, S.; Hernández-Martínez, A.; Rodríguez-Almagro, J.; Delgado-Rodríguez, M.; Martínez-Galiano, J.M. Relationship between perceived obstetric violence and the risk of postpartum depression: An observational study. *Midwifery.* 2022,108,103297. doi: 10.1016/j.midw.2022.103297.
23. Reuther, M.L. Prevalence of Obstetric Violence in Europe: Exploring Associations with trust, and care-seeking intention. Bachelor's thesis, University of Twente, Enschede, Netherlands, August 3, 2021.
24. Miller, S.; Lalonde, A. The global epidemic of abuse and disrespect during childbirth: History, evidence, interventions, and FIGO's mother-baby friendly birthing facilities initiative. *Int J Gynaecol Obstet.* 2015, 131, S49-52. doi: 10.1016/j.ijgo.2015.02.005.
25. Khalil M.; Carasso, K.B.; Kabakian-Khasholian T. Exposing Obstetric Violence in the Eastern Mediterranean Region: A Review of Women's Narratives of Disrespect and Abuse in Childbirth. *Front Glob Womens Health.* 2022,3,850796. doi: 10.3389/fgwh.2022.850796.
26. Santiago, R.V.; Solórzano, E.H.; Iñiguez, M.M.; Monreal, L.M.A. Nueva evidencia a un viejo problema: El abuso de las mujeres en las salas de parto. *Rev. Conamed.* 2015,18, 14–20.
27. Terán, P.; Castellanos, C.; Gonzalez Blanco, M.; Ramos, D. Violencia obstétrica: Percepción de las usuarias. *Rev Obstet Ginecol Venez.* 2013, 73, 171–180.
28. Molla, W.; Wudneh, A.; Tilahun, R. Obstetric violence and associated factors among women during facility based childbirth at Gedeo Zone, South Ethiopia. *BMC Pregnancy Childbirth.* 2022, 22, 565 (2022). doi:10.1186/s12884-022-04895-6.
29. Sando, D.; Ratcliffe, H.; McDonald, K.; Spiegelman, D.; Lyatuu, G.; Mwanyika-Sando, M.; Emil, F.; Wegner, M.N.; Chalamilla, G.; Langer, A. The prevalence of disrespect and abuse during facility-based childbirth in urban Tanzania. *BMC Pregnancy Childbirth.* 2016, 16, 236. doi: 10.1186/s12884-016-1019-4.
30. Yalley, A.A. "We Beat Them to Help Them Push": Midwives' Perceptions on Obstetric Violence in the Ashante and Western Regions of Ghana. *Women.* 2023, 3, 22-40. doi:10.3390/women3010002.
31. Ferrão, A.C.; Sim-Sim, M.; Almeida, V.S.; Zangão, M.O. Analysis of the Concept of Obstetric Violence: Scoping Review Protocol. *J Pers Med.* 2022,12,1090. doi: 10.3390/jpm12071090.
32. Mena-Tudela, D.; Iglesias-Casás, S.; González-Chordá, V.M.; Valero-Chillerón, M.J.; Andreu-Pejó, L.; Cervera-Gasch, Á. Obstetric Violence in Spain (Part III): Healthcare Professionals, Times, and Areas. *Int J Environ Res Public Health.* 2021,18,3359. doi: 10.3390/ijerph18073359.
33. Swahnberg, I.M.; Wijma, B. The NorVold Abuse Questionnaire (NorAQ): validation of new measures of emotional, physical, and sexual abuse, and abuse in the health care system among women. *Eur J Public Health.* 2003,13, 361-6. doi: 10.1093/eurpub/13.4.361.
34. Dhakal, P.; Gamble, J.; Creedy, D.K.; Newnham, E. Development of a tool to assess students' perceptions of respectful maternity care. *Midwifery.* 2022, 105, 103228. doi: 10.1016/j.midw.2021.103228.

35. Dhakal, P.; Creedy, D.K.; Gamble, J.; Newnham, E.; McInnes, R. Effectiveness of an online education intervention to enhance student perceptions of Respectful Maternity Care: A quasi-experimental study. *Nurse Educ Today*. 2022, 114, 105405. doi: 10.1016/j.nedt.2022.105405.
36. Çamlıbel, M.; Uludağ, E. The Turkish version of the students' perceptions of respectful maternity care scale: An assessment of psychometric properties. *Nurse Educ Pract*. 2023,70,103684. doi: 10.1016/j.nepr.2023.103684.
37. Mena-Tudela, D.; Cervera-Gasch, A.; Alemany-Anchel, M.J.; Andreu-Pejó, L.; González-Chordá, V.M. Design and Validation of the PercOV-S Questionnaire for Measuring Perceived Obstetric Violence in Nursing, Midwifery and Medical Students. *Int J Environ Res Public Health*. 2020,17,8022. doi: 10.3390/ijerph17218022.
38. Mena-Tudela, D.; Cervera-Gasch, Á.; Andreu-Pejó, L.; Alemany-Anchel, M.J.; Valero-Chillerón, M.J.; Peris-Ferrando, E.; Mahiques-Llopis, J.; González-Chordá, V.M. Perception of obstetric violence in a sample of Spanish health sciences students: A cross-sectional study. *Nurse Educ Today*. 2022, 110, 105266. doi: 10.1016/j.nedt.2022.105266.
39. Biurrun-Garrido, A.; Brigidi, S.; Mena-Tudela, D. Perception of health sciences and feminist medical students about obstetric violence. *Enferm Clin (Engl Ed)*. 2023, 33, 234-243. doi: 10.1016/j.enfcl.2023.02.007.
40. Cárdenas, M.; Salinero, S. Validación de la escala de violencia obstétrica y pruebas de la invarianza factorial en una muestra de mujeres chilenas. *Interdisciplinaria*. 2021, 38, 209-223, doi:10.16888/interd.2021.38.2.14.
41. El Parto es nuestro. Test de violencia obstétrica. Available online: <https://www.elpartoesnuestro.es/blog/2014/08/18/test-de-violencia-obstetrica>. (Accessed on 25 September 2022).
42. Penfield, R.D.; Giacobbi, P.R. Jr. Applying a score confidence interval to Aiken's item content-relevance index. *Measurement in Physical Education and Exercise Science*.2004, 8, 213-225. doi:10.1207/s15327841mpee0804_3.
43. Barrio-Cantalejo, I.M.; Simón-Lorda, P.; Melguizo, M.; Escalona, I.; Marijuán, M.I.; Hernando, P. Validación de la Escala INFLEZ para evaluar la legibilidad de los textos dirigidos a pacientes. *Anales Sis San Navarra*. 2008,31, 135-152.
44. Ferrando, P.J.; Lorenzo-Seva, U. Hernández-Dorado, A.; Muñoz, J. [Decalogue for the Factor Analysis of Test Items]. *Psicothema*. 2022,34, 7-17. doi: 10.7334/psicothema2021.456.
45. Lloret-Segura, S.; Ferreres-Traver, A.; Hernández-Baeza, A.; Tomás-Marco, I. Exploratory Item Factor Analysis: a practical guide revised and updated. *Anal. Psicol*. 2014, 30, 1151-1169.doi: 10.6018/analesps.30.3.199361.
46. Soriano-Vidal, F.J.; Oliver-Roig, A.; Cabrero-García, J.; Congost-Maestre, N.; Dencker, A.; Richart-Martínez, M. The Spanish version of the Childbirth Experience Questionnaire (CEQ-E): reliability and validity assessment. *BMC Pregnancy Childbirth*. 2016,16, 372. doi: 10.1186/s12884-016-1100-z.
47. González-de la Torre, H.; Miñarro-Jiménez, S.; Palma-Arjona, I.; Jeppesen-Gutierrez, J.; Berenguer-Pérez, M.; Verdú-Soriano, J. Perceived satisfaction of women during labour at the Hospital Universitario Materno-Infantil of the Canary Islands through the Childbirth Experience Questionnaire (CEQ-E). *Enferm Clin (Engl Ed)*. 2021,31,21-30. doi: 10.1016/j.enfcli.2020.05.022.
48. Lloret-Segura, S.; Ferreres-Traver, A.; Hernández-Baeza, A.; Tomás-Marco, I. The exploratory factor analysis of items: guided analysis based on empirical data and software. *Anal. Psicol*. 2017, 33,417-432. doi: 10.6018/analesps.33.2.270211.
49. Hancock, G.R.; Mueller, R.O. Rethinking Construct Reliability within Latent Variable Systems. In: *Structural equation modeling: Present and future.*; Cudeck, R.,; du Toit, S.; Sörbom, D., Eds; Lincolnwood, IL: Scientific Software, United States of America, 2001; pp. 195 -216.
50. Ferrando, P. J.; Lorenzo-Seva, U. Assessing the quality and appropriateness of factor solutions and factor score estimates in exploratory item factor analysis. *Educ Psychol Meas*.2018,78,762-780. doi:10.1177/0013164417719308.
51. Bond, T.G.; Fox, C.M. Applying the Rasch model: Fundamental measurement in the human sciences,2nd ed.; Mahwah, NJ: Lawrence Erlbaum, United States of America, 2007. Doi:10.4324/9781410614575.
52. Christensen, K.B.; Makransky, G.; Horton, M. Critical Values for Yen's Q3: Identification of Local Dependence in the Rasch Model Using Residual Correlations. *Appl Psychol Meas*. 2017, 41,178-194. doi: 10.1177/0146621616677520.

53. Rodríguez Mir, J.; Martínez Gandolfi, A. Obstetric violence denied in Spain. *Enferm Clin (Engl Ed)*. 2022,32 Suppl 1,S82-S83. doi: 10.1016/j.enfcle.2021.09.004.
54. Khsim, I.E.F.; Rodríguez, M.M.; Riquelme Gallego, B.; Caparros-González, R.A.; Amezcua-Prieto, C. Risk Factors for Post-Traumatic Stress Disorder after Childbirth: A Systematic Review. *Diagnostics (Basel)*. 2022,12,2598. doi: 10.3390/diagnostics12112598.
55. Soto, C.M.;Segovia, J.L. [Confidence intervals for the content validity: A Visual Basic computer program for the Aiken's V]. *Anal. Psicol*. 2009,25, 169–171.
56. Charter, R. A. A breakdown of reliability coefficients by test type and reliability method and the clinical implications of low reliability. *J Gen Psychol*. 2003 Jul;130(3):290-304. doi: 10.1080/00221300309601160.
57. Fraser, C.; McDonald, R.P. NOHARM: Least squares item factor analysis. *Multivariate Behav Res*. 1988,23, 267-9. doi: 10.1207/s15327906mbr2302_9.
58. Ferrando P.J.; Lorenzo-Seva, U. El Análisis Factorial Exploratorio de los Ítems: algunas consideraciones adicionales. *Anal. Psicol*.2014,30,1170-1175. Doi:10.6018/analesps.30.3.199991.
59. Lord FM. Applications of Item Response Theory to Practical Testing Problems, 1st. ed.; Routledge Taylor and Francis Group; New York, USA, 2012.
60. Escobar Bravo, M.A. Adaptación transcultural de instrumentos de medida relacionados con la salud. *Enferm Clin*. 2004,14,102-6.
61. Stolt, M.; Kottorp, A.; Suhonen, R. The use and quality of reporting of Rasch analysis in nursing research: A methodological scoping review. *Int J Nurs Stud*. 2022,132,104244. doi: 10.1016/j.ijnurstu.2022.104244.
62. McNeish, D. Thanks coefficient alpha, we'll take it from here. *Psychol Methods*. 2018, 23, 412-433. doi: 10.1037/met0000144.
63. Martínez-Galiano, J.M.; Martínez-Vazquez, S.; Rodríguez-Almagro, J.; Hernández-Martinez, A. The magnitude of the problem of obstetric violence and its associated factors: A cross-sectional study. *Women Birth*. 2021, 34, e526-e536. doi: 10.1016/j.wombi.2020.10.002.
64. BOE-A-2002-22188. Ley 41/2002, de 14 de noviembre, básica reguladora de la autonomía del paciente y de derechos y obligaciones en materia de información y documentación clínica. Spain,2002.
65. Medeiros, R.M.K.; Figueiredo, G.; Correa, Á.C.P; Barbieri, M. Repercussions of using the birth plan in the parturition process. *Rev Gaucha Enferm*. 2019;40,e20180233. doi: 10.1590/1983-1447.2019.20180233.
66. Mirghafourvand, M.; Mohammad Alizadeh Charandabi, S.; Ghanbari-Homayi, S.; Jahangiry, L.; Nahae, J.; Hadian,T. Effect of birth plans on childbirth experience: A systematic review. *Int J Nurs Pract*. 2019,25,e12722. doi: 10.1111/ijn.12722.
67. Bell, C.H.; Muggleton, S.; Davis, D.L. Birth plans: A systematic, integrative review into their purpose, process, and impact. *Midwifery*. 2022,111,103388. doi: 10.1016/j.midw.2022.103388.
68. Baranowska, B.; Doroszewska, A.; Kubicka-Kraszyńska, U.; Pietrusiewicz, J.; Adamska-Sala, I.; Kajdy, A.; Sys, D.; Tataj-Puzyna, U.; Bączek, G.; Crowther, S. Is there respectful maternity care in Poland? Women's views about care during labor and birth. *BMC Pregnancy Childbirth*. 2019,19,520. doi: 10.1186/s12884-019-2675-y.
69. Brazy-Nancy, E.; Mattern, C.; Rakotonandrasana, B.I.; Ravololomihanta, V.; Norolalao, P.; Kapesa, L. A qualitative analysis of obstetric violence in rural Madagascar. *Heliyon*. 2023, 9, e13905. doi: 10.1016/j.heliyon.2023.e13905.
70. Galera-Barbero, T.M.; Aguilera-Manrique, G. Women's reasons and motivations around planning a home birth with a qualified midwife in Spain. *J Adv Nurs*. 2022, 78, 2608-2621. doi: 10.1111/jan.15225.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.