

Supplementary Materials

(3-(4-chlorophenyl)-4,5-dihydroisoxazol-5-yl)methyl benzenesulfonate

Loubna Mokhi ¹, Karim Chkirate ², Xiaodong Zhang ³, Mohsine Driowya ^{1,4} and Khalid Bougrin ^{1,5,*}

¹ *Equipe de Chimie des Plantes et de Synthèse Organique et Bioorganique, URAC23, Faculty of Science, B.P. 1014, Geophysics, Natural Patrimony and Green Chemistry (GEOPAC) Research Center, Mohammed V University in Rabat, Rabat 10010, Morocco*

² *Laboratory of Heterocyclic Organic Chemistry URAC 21, Pharmacochimie Competence Center, Av. Ibn Battouta, BP 1014, Faculty of Sciences, Mohammed V University in Rabat, Rabat 10010, Morocco*

³ *Department of Chemistry, Tulane University, New Orleans, LA 70118, USA*

⁴ *Higher School of Technology, Sultan Moulay Slimane University, B.P. 170, Khenifra 54006, Morocco.*

⁵ *Chemical & Biochemical Sciences Green-Process Engineering (CBS), Mohammed VI Polytechnic University, Lot 660, Hay Moulay Rachid, Ben Guerir 43150, Morocco*

* Corresponding author:

Prof. Khalid Bougrin

Email address: kbougrin@yahoo.fr or k.bougrin@um5.ac.ma

Table of contents

1. ^1H NMR and ^{13}C NMR spectra of compounds 5.....	3
2. ESI-MS spectra of compounds 5	4
3. FT-IR spectra of compounds 5	4

1. Copies of characteristic ^1H NMR and ^{13}C NMR spectra




Figure S1. ^1H NMR spectrum of (5)




Figure S2. ^{13}C NMR spectrum of (5)

2. Copy of characteristic ESI-MS spectra




Figure S3. ESI⁺-MS spectrum of (5)

3. Copy of characteristic FT-IR spectra




Figure S4. FT-IR spectrum of (5)