

Brief Report

Not peer-reviewed version

---

# Studying the Effect of Anthropogenic Activities on the Presence of Nitrogen Dioxide (NO<sub>2</sub>) through Remote Sensing and Ground Based Observations over Emirate of Fujairah, UAE

---

Reem Yaqoub Yousif Ahmed Abdalla , Aseela AL Moalla , Mohamed Ateeg , Marwa Hossny , Fouad Lamghari , [Shaher Bano Mirza](#) \*

Posted Date: 25 October 2023

doi: 10.20944/preprints202310.1543.v1

Keywords: atmospheric pollution; NO<sub>2</sub>; air quality; Sentinel-5p; TROPOMI; remote sensing data; covid-19; pandemic, outbreak



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Brief Report

# Studying the Effect of Anthropogenic Activities on the Presence of Nitrogen Dioxide (NO<sub>2</sub>) through Remote Sensing and Ground Based Observations over Emirate of Fujairah, UAE

Reem Yaqoub Yousif Ahmed Abdalla<sup>1</sup>, Aseela Al Moalla<sup>2</sup>, Mohamed Ateeg<sup>2</sup>, Marwa Hossny<sup>2</sup>, Shafer Bano Mirza<sup>1,\*</sup> and Fouad Lamghari Ridouane<sup>1</sup>

<sup>1</sup> Fujairah Research Centre, Sakamkam Road, Fujairah, United Arab Emirates.

<sup>2</sup> Fujairah Environment Authority, Sakamkam Road, Fujairah, United Arab Emirates.

\* Correspondence: shaher.bano@frc.ae; Phone: +971 508011873

**Abstract:** Air pollution is one of the biggest problems of this age, not only it contributes to climate change but also it has a negative effect on public and individual health due to rising morbidity and mortality. There are various contaminants that play a significant role in human disease. Numerous toxins are released into the environment as a result of anthropogenic activities including burning fossil fuels for transportation and electricity generation. In order to assist decision-makers in finding a long-term solution for improving environmental quality and population health status, NO<sub>2</sub> pollution monitoring and regulation is an essential task. moreover, efforts to reduce NO<sub>2</sub> emissions are important for improving air quality and public health. Based on data acquired from Sentinel-5P, the study presented a comparison of NO<sub>2</sub> measurements in Sentinel-5P and ground-based stations over the urban area of Fujairah during 2021 and comparison analysis of the tropospheric NO<sub>2</sub> column spatial configuration over urban area of Fujairah during comparable periods of pre, mid and post COVID-19 pandemic era as 2019, 2020 and 2021, respectively. These findings demonstrate remarkable tropospheric NO<sub>2</sub> column number density decreases even of 85% in the busier areas of emirate of Fujairah during the pandemic era. Sentinel-5P measurements draw attention to the significant reduction in NO<sub>2</sub> pollution over Fujairah during the COVID-19 lockdown where we had less traffic, which is supported by ground stations-based calculations.

**Keywords:** Atmospheric pollution; NO<sub>2</sub>; air quality; Sentinel-5p; TROPOMI; remote sensing data; COVID-19; Pandemic; outbreak

---

## Introduction

Urban air pollution is made of extremely complex mixture of gaseous and particulate substances [1]. Anthropogenic activities such as combustion of fossil fuels for energy production and transportation entails as the major cause of releasing various pollutants in the environment. These pollutants cause atmospheric composition change which negatively impact human health and its ecosystem [2]. Vehicle emissions are anticipated to worsen from what they are now by 2030. Numerous studies have reported the negative influence of vehicle emissions like hydrocarbons, nitrogen oxide, carbon monoxide, and particulates which are anticipated to worsen by 2030 [3]. Nitrogen dioxide (NO<sub>2</sub>) is the air pollutant species that is most critical to human health of its direct respiratory irritation, exacerbation of existing health conditions, long-term negative health impacts, and contribution to other harmful pollutants. Air quality guideline by World Health Organization (WHO) suggests NO<sub>2</sub> level shouldn't exceed 10 µg/m<sup>3</sup> annual mean and 25 µg/m<sup>3</sup> 24-hour mean. In 2019, 99% of the world population was living in places where these criteria were not meet [4]. According to WHO, 91 percent of the world population breathe air with high levels of pollutants which results in death of 7 million people/year [5]. According to epidemiological studies, bronchitis symptoms increased in asthmatic children due to long-term exposure of NO<sub>2</sub> [5]. NO<sub>2</sub> has capacity to create secondary pollutants like nitric acid and ozone (O<sub>3</sub>), which help to create smog, acid rain, and

the greenhouse effect [6]. Therefore, global warming increases when greenhouse gas concentrations rise in the atmosphere. The cumulative warming effect to the earth's atmosphere from greenhouse gases increased by 45 percent from 1990 to 2019 [7]. Lowering anthropogenic emissions of air pollutants and atmospheric particles from sources like vehicle traffic may help to enhance the world's air quality [8].

Monitoring air quality has become a crucial public health and environmental protection activity. In addition to ground-based monitoring stations, satellites are increasingly utilised for tracing air pollution sources and monitoring air quality. Satellite-based data are incredibly useful for monitoring air pollutants and the physical conditions of the Earth's surface due to the increasing availability of remotely sensed data derived from sophisticated sensors with high resolutions. In addition, metrological and satellite-based derivatives are incorporated into a variety of studies to elucidate the spatiotemporal patterns of air pollutants, biophysical parameters, and respiratory disease [9].

The COVID-19 outbreak, which was first reported in late December 2019 in Wuhan, China, has become one of the most grievous health crises of the 21st century, posing grave risks to all nations. Diverse drastic measures were taken on a global scale to control and mitigate the virus's rapid spread and to reduce its mortality rate. Particularly, the transport sector was severely impacted by curfews imposed as a result of the adoption of remote work systems by educational institutions, public and private sectors [10]. As a consequence, during the first few months of the pandemic, people's mobility changed drastically as the majority stayed at home during working hours. In addition, researchers began to report improvements in air quality based on information acquired from satellite images and monitoring stations on the ground. Using Sentinel-5 satellite data, numerous studies evaluated the changes in various air pollutants during the 2020 COVID-19 lockdown (March to June 2020) in 21 metropolitan cities spanning 14 Middle Eastern nations. Significant decreases in various air pollutants including NO<sub>2</sub> levels were observed primarily in minor cities and, to a lesser extent, in metropolitan cities. Other researchers analysed the effect of the COVID-19 lockdown on air quality in the Middle East, including Saudi Arabia, Kuwait, Iraq, Iran, and the United Arab Emirates [11–13].

However, the available regional studies primarily concentrated on the early phases of the pandemic and did not adequately represent the recovery phase or provide comprehensive and quantitative analyses of the changes in various air pollutants and mobility patterns. The Fujairah emirate is not investigated extensively in the United Arab Emirates, and very little effort has been made to monitor environmental pollution here. This study documents the impact of the COVID-19 lockdown and subsequent resuming on mobility and air quality in the Fujairah region of the United Arab Emirates, compares the impact of the lockdown on air quality as reflected by satellite data and ground-based observations, and correlates the mobility patterns of people with the observed changes in air quality during the lockdown and subsequent recovery.

The data mining process will be applied to collect data from satellite and four different ground-based station located in the Fujairah. The point-based ground measurements will be taken and applied in spatial modelling to accurately quantify the amount of air pollution. The distribution modelling of NO<sub>2</sub> is useful for epidemiological studies, management of the urban air which further helps in government planning and regulations [14]. These efforts will assist the government in finding a long-term solution for improving environmental quality and the health of the population.

## **Materials and Methods**

### *Study area*

Fujairah is one of the seventh Emirates in the United Arab Emirates. It is the only Emirate to be located on the eastern side of the UAE, near to Gulf of Oman. The total area of Fujairah is 1450 Km<sup>2</sup> which covers 1.8% of the whole area of UAE (About Fujairah, 2022). The population of the Emirate of Fujairah is reported as 292,358 in 2020, according to the Fujairah Statistics Centre (Fujairah's Population, 2021). Our analysis focuses on the urban area of Fujairah in 2021 which covers 450.05 Km<sup>2</sup> as shown in Figure 1.



**Figure 1.** The selected study area in Fujairah.

#### *Data Mining:*


The data has been collected from ground-based stations in urban Fujairah provided by Fujairah Environment Authority (FEA) and satellite Sentinel-5 Precursor (Table 1). These ground-based weather stations are equipped with Analyzer sensors capable of capturing a diverse array of atmospheric constituents, including  $\text{NO}_2$ . These sensors are strategically positioned at heights ranging between 5 to 10 meters above ground level, facilitating the collection of accurate and representative air quality measurements.

**Table 1.** Location and classification of spatial nature of the Areas where ground based stations are installed.

Station Name	Classification Of Spatial Nature
Point 1	Residential – Industrial
Point 2	Residential – Industrial
Point 3	Residential – Industrial
Point 4	Residential – Urban

#### *Nitrogen dioxide ( $\text{NO}_2$ ) data from ground-based stations*

These data include  $\text{NO}_2$  concentrations in microgram per cubic meter ( $\mu\text{g}/\text{m}^3$ ) measured hourly through day and night and average temperature in Celsius ( $^{\circ}\text{C}$ ) during each month at four different stations in the selected Fujairah urban area in 2021 (Table 1, Figure 2). An algorithm written in python has been applied to measure the mean of the concentration in each day. These data were used as reference to identify the differences in  $\text{NO}_2$  levels in various stations and compare it to the Sentinel-5P data.




**Figure 2.** The stations under study are marked on the map using GPS coordinates in the Google Earth Pro.


#### *Sentinel-5 Precursor satellite*

The data of Sentinel-5P has been collected via Sentinel Hub using EO Browser in near real time (NRTI). Sentinel-5P is an atmospheric monitoring satellite and has only one sensor called TROPospheric Monitoring Instrument (TROPOMI), which is the most developed multispectral imaging spectrometer. It monitors the sunlight that is reflected to space from the earth's surface and atmosphere and finds the special fingerprints of different gases in various regions of the electromagnetic spectrum. The gases accounted for special fingerprints include nitrogen dioxide, ozone, formaldehyde, methane, carbon monoxide, etc, and aerosols [17]. The Sentinel-5 Precursor satellite contributes valuable information for a wide range of applications related to environmental protection, public health, climate change mitigation, and scientific research. The data of NO<sub>2</sub> tropospheric column in the unit of mol per square meter (mol/m<sup>2</sup>) has been collected for 2021. The images of spatial distribution of NO<sub>2</sub> concentration over the study area has been downloaded for the following dates: 26/11/2019, 29/12/2019, 21/1/2020, 9/2/2020, 17/3/2020, 27/3/2020, 12/1/2021 and 27/3/2021 based on Mohammed bin Rashid space centre survey on NO<sub>2</sub> concentration before and during the lockdown [18] (Figure 3).







**Figure 4.** Comparative dynamics of NO<sub>2</sub> daily values for the year 2021 in Fujairah urban area, Point 3 ground-based station measurements are mentioned as orange line in µg/m<sup>3</sup>, Satellite data (Sentinel-5P) mentioned as blue line in mol/m against the dates on horizontal axis.




**Figure 5.** Comparative dynamics of NO<sub>2</sub> daily values for the year 2021 in Fujairah urban area, Point 4 ground-based station measurements are mentioned as orange line in µg/m<sup>3</sup>, Satellite data (Sentinel-5P) mentioned as blue line in mol/m against the dates on horizontal axis.



**Figure 6.** Comparative dynamics of NO<sub>2</sub> daily values for the year 2021 in Fujairah urban area, Point 2 ground-based station measurements are mentioned as orange line in  $\mu\text{g}/\text{m}^3$ , Satellite data (Sentinel-5P) mentioned as blue line in  $\text{mol}/\text{m}^2$  against the dates on horizontal axis.



**Figure 7.** Comparative dynamics of NO<sub>2</sub> daily values for the year 2021 in Fujairah urban area, Point 1 ground-based station measurements are mentioned as orange line in  $\mu\text{g}/\text{m}^3$ , Satellite data (Sentinel-5P) mentioned as blue line in  $\text{mol}/\text{m}^2$  against the dates on horizontal axis.




**Figure 8.** Average temperature in Celsius for each month of 2021 retrieved from Skamkam station data, Average temperatures in Celsius ( $^{\circ}\text{C}$ ) are shown in vertical axis and the months of 2021 on horizontal axis.

#### *Comparison of NO<sub>2</sub> pollution and covid-19 pandemic lockdown*

The COVID-19 pandemic led to significant and temporary reductions in NO<sub>2</sub> emissions and concentrations due to lockdowns, reduced economic activity, and changes in transportation patterns. The episode provided valuable insights into the relationship between human activities and air quality and prompted discussions on future strategies for achieving cleaner and healthier urban environments. The spatial distribution of NO<sub>2</sub> concentrations from Sentinel-5P is presented here for different dates to compare average tropospheric NO<sub>2</sub> column number density (Figure 9). Each colour



represents various levels of concentration. The dark red colour indicates the highest values of NO<sub>2</sub> concentrations which equals to 1.0E-4 mol/m<sup>2</sup> and the dark blue represents the least concentration of NO<sub>2</sub> which equals to 0 mol/m<sup>2</sup> (Figure 10). NO<sub>2</sub> concentration is highest in the east, then the north and west shows less concentration of NO<sub>2</sub>, and the south has the least concentration in the pre Covid-19 era presented here as the date of 26/11/2019. In contrast to that, on 29/12/2019 the north, northeast and northwest have higher NO<sub>2</sub> concentration and the south and southwest shows less NO<sub>2</sub> concentration. During 2020 pandemic, the level of NO<sub>2</sub> decreased dramatically although the north, northwest and northeast yet had higher NO<sub>2</sub> concentration compared to other locations on the map as seen on 27/3/2020 had the lowest level of NO<sub>2</sub>. The dark red colour for the southeast, the centre and east displayed for the day 27/3/2021 depicts the highest NO<sub>2</sub> concentration.



**Figure 9.** spatial distribution of NO<sub>2</sub> concentrations from Sentinel-5P. comparative maps on selected dates of average tropospheric NO<sub>2</sub> column number density over Fujairah urban area. On the right it shows the average tropospheric NO<sub>2</sub> column number density index.

## Discussion

The comparison of NO<sub>2</sub> concentration measurements acquired from Sentinel-5P as well as from ground-based stations reveals the significant impact weather has on NO<sub>2</sub> levels. The results exhibit pattern of NO<sub>2</sub> concentration which falls lower in the hotter months and higher in the colder months. Explanation may encompass on the increased traffic in colder months, as for UAE, the weather is much pleasant with average temperature in winters falling up to 19 °C as opposed to summer where average temperature may increase up to 41.3 °C. Point 3 station exhibit the highest average concentration level of NO<sub>2</sub> which equals to 33.24 µg/m<sup>3</sup>. The area is known to have dense zone with schools, offices, and industries. Consequently, pulling more traffic of motor vehicle including loader trucks towards this area during the daytime. Point 1 station had the least average concentration level of NO<sub>2</sub> which equals to 23.12 µg/m<sup>3</sup>. Point 1 area has less traffic compared to the other areas. The results highlighted that the traffic has huge impact on the NO<sub>2</sub> levels which bring above the speculation of motor vehicles as being the principal urban anthropogenic source of primary NO<sub>2</sub>. A greater percentage of primary NO<sub>2</sub> is present in the exhaust from diesel-fuelled vehicles compared to petrol-fuelled vehicles. Primary NO<sub>2</sub> is created through high combustion, from ground-level emissions connected to the burning of fossil fuels in automobiles [19]. While comparing results of NO<sub>2</sub> pollution with pre, mid and post pandemic era, we must point out that at the time of COVID-19 pandemic lockdown in 2020, significantly lower levels of NO<sub>2</sub> concentrations has been recorded where we have less traffic compared to 2019 and 2021. The result shows that the COVID-19 pandemic indeed had a positive effect on air governance which was beneficial for human health.

The distribution of the NO<sub>2</sub> pollution can be affected by the local climatic conditions. For example, pollutants can be spread and moved away from their source by strong winds and temperature is another climatic parameter to consider [20]. Which can tell that the emissions of NO<sub>2</sub> can be distributed to near areas of the original source.

To our knowledge, this is the first report of this kind of analysis from the region of Fujairah, UAE which instigate, the positive environmental impact of reduced level of NO<sub>2</sub> during the pandemic show that it is still possible to make great and rapid positive change. The conclusion follows from the fact that the highest levels of NO<sub>2</sub> to appear are in crowded areas where we have more traffic and near to industrial area which are the north and the northeast areas of the Fujairah urban area.

**Conflicts of interest:** The authors declare that they have no conflict of interests.

**Acknowledgments:** The authors thank the Fujairah Research Centre (FRC) for their financial and technical support and Fujairah Environment Authority (FEA) for technical support. Authors also extends their gratitude to Muhammed Sirajul Huda Kalathingal for technical support in experimental work. Authors appreciate the continuous support of Ms. Aseela Al Moalla throughout the project and initiating this project.

## References

1. J. Seinfeld, "Urban Air Pollution: State of the Science," 1989. [Online]. Available: [www.sciencemag.org](http://www.sciencemag.org)
2. M. Kampa and E. Castanas, "Human health effects of air pollution," *Environmental Pollution*, vol. 151, no. 2, pp. 362–367, Jan. 2008. doi: 10.1016/j.envpol.2007.06.012.
3. N. R. S. Olmo, P. H. do N. Saldiva, A. L. F. Braga, C. A. Lin, U. de P. Santos, and L. A. A. Pereirai, "A review of low-level air pollution and adverse effects on human health: Implications for epidemiological studies and public policy," *Clinics*, vol. 66, no. 4, pp. 681–690, 2011. doi: 10.1590/S1807-59322011000400025.
4. WHO, "WHO global air quality guidelines," 2021.
5. "Ambient (outdoor) air pollution," Sep. 22, 2021. [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health) (accessed Sep. 25, 2022).
6. WHO, "Health Aspects of Air Pollution with Particulate Matter, Ozone and Nitrogen Dioxide Report on a WHO Working Group OZONE-adverse effects NITROGEN DIOXIDE-adverse effects AIR POLLUTANTS, ENVIRONMENTAL-adverse effects META-ANALYSIS AIR-standards GUIDELINES," 2003.
7. EPA, "climate change," 2022. <https://www.epa.gov/climate-indicators/greenhouse-gases> (accessed Sep. 25, 2022).
8. H. C. Loh, I. Looi, A. S. H. Ch'ng, K. W. Goh, L. C. Ming, and K. H. Ang, "Positive global environmental impacts of the COVID-19 pandemic lockdown: a review," *GeoJournal*. Springer Science and Business Media Deutschland GmbH, 2021. doi: 10.1007/s10708-021-10475-6.
9. A. Shanableh *et al.*, "Effects of the COVID-19 lockdown and recovery on People's mobility and air quality in the United Arab Emirates using satellite and ground observations," *Remote Sens Appl*, vol. 26, p. 100757, 2022, doi: <https://doi.org/10.1016/j.rsase.2022.100757>.
10. H. Zheng *et al.*, "Significant changes in the chemical compositions and sources of PM<sub>2.5</sub> in Wuhan since the city lockdown as COVID-19," *Science of the Total Environment*, vol. 739, p. 140000, 2020.
11. H. Bherwani *et al.*, "Valuation of air pollution externalities: comparative assessment of economic damage and emission reduction under COVID-19 lockdown," *Air Qual Atmos Health*, vol. 13, pp. 683–694, 2020.
12. M. G. Del Buono, G. Iannaccone, M. Camilli, R. Del Buono, and N. Aspromonte, "The Italian outbreak of COVID-19: conditions, contributors, and concerns," in *Mayo Clinic Proceedings*, Elsevier, 2020, pp. 1116–1118.
13. O. Teixidó *et al.*, "The influence of COVID-19 preventive measures on the air quality in Abu Dhabi (United Arab Emirates)," *Air Qual Atmos Health*, vol. 14, pp. 1071–1079, 2021.
14. I. Suryati, H. Khair, and D. Gusrianti, "Distribution analysis of nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) in Medan city with Geographic Information System (GIS)", doi: 10.1051/mateconf/20192760.
15. "About Fujairah," Sep. 20, 2022. <http://fujegovt.ae/aboutfujairah.html> (accessed Sep. 20, 2022).
16. "Fujairah's population," May 02, 2021. <https://www.wam.ae/en/details/1395302931971> (accessed Sep. 20, 2022).
17. A. Castro Gómez, "ESA UNCLASSIFIED-For Official Use Exercise with EO Browser: Air Pollution (Sentinel-2, Sentinel-5P)," 2019.
18. Mohammed Bin Rashid Space Centre, "NO<sub>2</sub> Concentration in the Gulf Region & the UAE," Sep. 29, 2021. <https://storymaps.arcgis.com/stories/246700bd610b4cf682f31fd5118b659c> (accessed Sep. 24, 2022).

19. M. Hallquist and E. Mc Queen, "DEPARTMENT OF BIOLOGICAL AND ENVIRONMENTAL SCIENCES INVESTIGATION OF THE NOX-NO2-WIND SPEED-RELATIONSHIP AND PRIMARY NO2 AT TWO MONITORING STATIONS IN GOTHENBURG, 2010 AND 2019," 2020.
20. L. Contreras Ochando, C. I. Font Julián, and F. Contreras Ochando, "Airvlc: An application for real-time forecasting urban air pollution C`esar Ferri," 2015. [Online]. Available: <http://www.aemet.es/>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.