

Article

Not peer-reviewed version

Predictors of Vaccination Intentions and Behaviour during the COVID-19 pandemic in Italy

[Amanda Nerini](#)*, [Mirko Duradoni](#), [Camilla Matera](#), [Andrea Guazzini](#), Monica Paradisi, Adriele Sghembri

Posted Date: 13 November 2023

doi: 10.20944/preprints202311.0763.v1

Keywords: vaccination hesitancy; fear; Protection Motivation Theory; psychology; prediction; COVID-19 vaccination; health behavior



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Predictors of Vaccination Intentions and Behaviour during the COVID-19 Pandemic in Italy

Amanda Nerini ^{1*}, Mirko Duradoni ¹, Camilla Matera ¹, Andrea Guazzini ¹, Monica Paradisi ¹ and Adriele Sghembri ²

¹ Department of Education, Languages, Intercultures, Literatures and Psychology, University of Florence, Via San Salvi, 12 (pad.26), 50135, Florence, Italy, mirko.duradoni@unifi.it; camilla.matera@unifi.it; andrea.guazzini@unifi.it; monica.paradisi@unifi.it

² School of Psychology, University of Florence, Via della Torretta, 16, 50135, Florence, Italy
adriele.sghembri@stud.unifi.it

* Correspondence: amanda.nerini@unifi.it

Abstract: The present research aimed at understanding individuals' vaccination intentions and behaviours against COVID-19 through two different studies. In Study 1 (N = 213, 73% women; mean age = 24.03) the Protection Motivation Theory model was tested considering the fear of COVID-19 as a possible mediator between threat appraisal (in terms of both health and social life) and intentions to get vaccinated when vaccination was not already available. Study 2 (N = 1111, 68.9% women; mean age = 38.33) was conducted when the vaccine became available for the entire population. Through this study, by adopting the 5C model of vaccine hesitancy as a theoretical framework, we aimed to understand how people who got vaccinated and the ones who did not differed, considering fear of vaccination, personality and vaccination hesitancy. In Study 1, social limitations perceived severity and COVID-19 perceived vulnerability were significantly and positively related to fear of COVID-19. Contrary to what we expected, fear of Covid-19 was not a significant predictor of vaccination intention, which was predicted by both response efficacy and self-efficacy. In Study 2, in line with previous studies, vaccine hesitancy was negatively related to vaccination. More specifically, the social-oriented dimension of collective responsibility was the strongest predictor of effective behaviour. Our findings provide insights into the complexity of vaccine acceptance and emphasise the need for targeted interventions to promote vaccination and mitigate the spread of infectious diseases.

Keywords: vaccination hesitancy; fear; Protection Motivation Theory; COVID-19 vaccination; health behaviour; prediction; psychology

1. Introduction

On March 11 2020, afterward the rapid spread of COVID-19 around the world, the World Health Organization [1] proclaimed pandemic status. On December 27 2020, following European Medicines Agency approval, the COVID-19 vaccination campaign was launched in Europe to achieve herd immunity as quickly as possible [2], which allowed for a gradual and steady relaxation of containment measures. However, there was a significant portion of the population that did not want to receive vaccination against COVID-19, although available. The analysis of factors promoting the intention to get vaccinated is still ongoing to help health professionals in planning intervention that can foster behavioural change in order to vaccinate as many people as possible. Adopting the Protection Motivation Theory (PMT) [3, 4] and the 5C model [5] as theoretical frameworks, the present study aimed to identify factors that might motivate individuals to receive such a vaccination.

According to PMT [3, 4] the motivation to protect oneself, considered to be the proximate cause of health behaviour, is affected by threat and coping appraisal. Threat appraisal is determined by perceived threat severity, which refers to the beliefs about the negative consequences that may occur as a result of the maladaptive behaviour, and threat vulnerability, that consists of the probability of being susceptible to the negative consequences of the maladaptive behaviour. Coping appraisal is determined by response efficacy, which refers to the personal perceptions that adaptive behaviour

can eliminate or reduce the negative consequences of maladaptive behaviour, and self-efficacy, which refers to the individual's ability to implement the recommended behaviour. Fear, which is elicited in response to a situation judged as dangerous and towards which a protective action is taken [3], is considered to mediate the relationship between threat appraisal and the intention to implement a protective behaviour [6, 7]. In recent years, several studies applied PMT to the intention to be vaccinated against COVID-19, partially confirming its effectiveness in predicting behavioural intention [8-12]. Indeed, among the PMT variables, the response efficacy was found to be significantly related to received vaccination against COVID-19 [8-10; 12] and seemed to be the most decisive predictor; in contrast, the role of self-efficacy, perceived threat severity, and threat vulnerability are unclear, as these variables have not always emerged as significant predictors of the intention to be vaccinated against COVID-19 [8-12]. Xiao and colleagues [12] found a significant role of the variables related to the coping appraisal in the prediction of the intention to get vaccinated against COVID-19; specifically, response efficacy was the most powerful predictor of the criterion variable, while no significant association emerged between threat appraisal (threat severity and threat vulnerability) and the intention to be vaccinated against COVID-19. In the study by Wang and colleagues [11] on a sample of 3,145 university students, perceived severity of COVID-19 was positively associated with the motivation to get vaccinated; nevertheless, the coping appraisal dimensions of PMT and threat vulnerability were not significant predictors of motivation. Nerini and colleagues [13] showed that the fear of COVID-19 was relevant in motivating people to follow the measures set by the Italian government to reduce the spread of the pandemic and mediated the relationship between threat assessment and intentions. Feelings of fear arising from the perception that COVID-19 was a serious disease that could infect everyone, motivated individuals to adopt protective behaviours [13].

Scholars considered fear of vaccination as an important factor in vaccine hesitancy dynamics worldwide [14]. In the first months of 2021, due also to the widespread of fake news (e.g., the harmless nature of COVID-19 and poorly tested and experimental vaccines with hidden side effects) [15, 16], fear extended beyond the virus itself to include concerns about the vaccine [17]. The fear of experiencing adverse reactions or developing vaccine-related diseases, together with distrust of pharmaceutical companies and politicians, were a common reason for vaccine refusal [18-21]. All these factors contributed to the development of vaccine hesitancy, which refers to the hesitation or delay in accepting vaccination, even when vaccination services are readily available [22]. World Health Organization identifies vaccine hesitancy as one of the top 10 threats to global health in 2019 [23]. Apart from fear of vaccination, vaccine hesitancy is associated with socio-demographic (i.e., age and sex) and psychological (e.g., personality) variables [24, 25]. Women and younger-aged individuals seemed to be more hesitant and resistant [25]. Moreover, conscientiousness, narcissism, and psychopathy emerged as significant predictors of hesitation in accepting the COVID-19 vaccine [24].

Previous studies about COVID-19 considered perceived threat and associated feelings of fear as exclusively related to health (e.g., risk of death or long-term consequences). Nevertheless, both fear and vaccination hesitancy might be also influenced by the pandemic negative consequences that people perceive with respect not only to their health status, but also to their social life (e.g., limitations to individuals' social activities). The role of social factors in determining the intention to get vaccinated was considered in Betsch and colleagues' 5C model [5]. The authors identified five psychological factors as the main antecedents of vaccination behaviour: confidence, complacency, calculation, constraints, and collective responsibility. Confidence refers to individuals' trust in the effectiveness and safety of vaccines, as well as their faith in the recommendations made by health professionals and policymakers. Complacency reflects a low perceived risk and a diminished sense of threat regarding vaccine-preventable diseases. Calculation captures the extensive information-seeking behaviour, which can lead to vaccine hesitancy due to exposure to anti-vaccination messages. Constraints encompass both physical and psychological barriers to vaccine accessibility (e.g., geographical, financial, and comprehension issues). Collective responsibility refers to the willingness to protect others through vaccination, contributing to herd immunity. Differently from the other four factors, collective responsibility has a strong social valence. Individuals with high levels of confidence

and collective responsibility tend to exhibit positive attitudes towards vaccination; in contrast, those with high levels of complacency and constraints are more likely to hold negative attitudes. Empirical evidence regarding the calculation factor remains inconclusive [5].

2. Research overview

Even though previous studies confirmed the validity of PMT in predicting the intention to engage in preventive health behaviour, mixed results emerged about the role played by PMT dimensions in the prediction of the intention to get vaccinated against COVID-19. In addition, in these studies threat appraisal was considered exclusively with regard to the negative consequences of being infected by COVID-19 (such as risk of death or long-term consequences). To the best of our knowledge, no study assessed perception of threat with regard to the negative consequences that may occur in terms of limitations to one's social life. As COVID-19 pandemic has been an unprecedented event that has affected the entire globe, politicians, health experts but also pundits and public opinionists filled the media with different, and sometimes opposite, positions about containment measures and vaccination. In order to motivate people to protect themselves and others, it is crucial to understand if and how classic theoretical models are able to explain individuals' vaccination intentions and behaviours and if factors that are more related to the broader social context should be considered together with more strictly cognitive ones.

Starting from these considerations, we conducted two studies: a preliminary study (study 1) aimed at analyzing the intention to get vaccinated in a specific segment of the population (young adults) for whom the vaccine, at the time in which the study was conducted, was not yet available; a second study aimed at understanding individuals' vaccination behaviour when the vaccine became available for the entire population, so that we moved from analysing the intention to get vaccinated to actual vaccination.

Specifically, in the first study the PMT model was used to explain the intention to get vaccinated when the vaccination was not already available. We also analysed the mediational role of fear of COVID-19 in the relationship between threat appraisal and intentions, considering threat related not only to one's health, but also to one's social life.

In the second study we examined vaccination hesitancy in people who got and did not get vaccinated. By adopting the 5C model of vaccine hesitancy as a theoretical framework, we considered factors that, although still cognitive in nature, capture the influence of the social context on the decision to receive the COVID-19 vaccine (e.g., collective responsibility). Moving beyond Study 1's emphasis on fear of the disease, Study 2 focused on the fear associated with the recommended response, namely vaccination. Specifically, we examined the role of vaccination fear in relation to vaccine hesitancy dimensions, along with sociodemographic variables such as sex and age, as well as personality traits.

In Study 1 we tested the following hypotheses. Both COVID-19 threat appraisal (i.e., COVID-19 perceived severity and COVID-19 perceived vulnerability) and social limitations threat appraisal (i.e., social limitations perceived severity and social limitations perceived vulnerability) would predict the intention to get vaccinated via fear of COVID-19 (Hypothesis 1). Coping appraisal (i.e., response efficacy and self-efficacy) would be directly associated with the intention to get vaccinated (Hypothesis 2).

In Study 2 we tested the following hypotheses. Vaccine hesitancy would be negatively associated with actual vaccination (Hypotheses 3). Vaccine fear would be positively associated with vaccine hesitancy even after controlling for participants' age, sex, and personality traits based on the 5C model (Hypotheses 4).

Study 1

Methods

Participants and procedure

Participants were 213 Italian young adults (73% women; mean age = 24.03, $SD = 3.03$). As we planned to analyse the data through path analysis, following the indications by Weston and Gore [26], we aimed to recruit a sample greater than 200 participants. To be eligible for the study, the participants had to be aged between 18 and 30 years, as vaccination against COVID-19 was not available for this group in Italy at the time the questionnaire was administered (spring of 2021). Despite that, as some groups of professionals had been vaccinated independently from the age (e.g., teachers, health professionals), to be eligible for the study participants had to confirm that they had not received any doses of vaccination against COVID-19.

An online anonymous survey was shared through social networking sites (i.e., Facebook groups, Instagram, LinkedIn) between 21 May 2021 and 14 June 2021, inviting participants to take part in a study on Italian youth intentions of getting vaccinated against COVID-19. No incentive was provided and all participants gave their informed consent to the study procedures, which adhered to the Helsinki Declaration, Italian legal requirements on privacy and informed consent (Law Decree DL-101/2018), EU regulations (2016/699), and guidelines from the American Psychological Association (APA). The Ethical Committee of the University of Florence approved the study procedures (Ref. No. 0096243). The survey took about 15 min to complete. To ensure participants had complete control over whether to submit their responses or not, we did not record responses from participants who did not complete the survey. Measures

For the following measures all the items were rated on a five-point Likert scale (1 = strongly disagree; 5 = strongly agree). Higher scores indicated higher levels in the measured variables.

COVID-19 and social limitations threat appraisals. COVID-19 and social limitations threat appraisals were assessed in terms of both perceived severity and vulnerability through ten items adapted from previous studies conducted in different countries, including Italy [13, 27, 28]. COVID-19 perceived severity was measured using a three-item scale (e.g., “Those who do not get vaccinated in the event of contagion can put the health of their loved ones at risk”; $\alpha = .86$) and COVID-19 perceived vulnerability through two items (e.g., “The possibility that I could contract COVID-19 is high”; $r = .57$; $p < .001$). Social limitations perceived severity was measured using a three-item scale (e.g., “those who do not get vaccinated have a more limited social life”; $\alpha = .84$) and social limitations perceived vulnerability through two items (e.g., “I suffer as I feel my social life is limited”; $r = .45$; $p < .001$).

Coping Appraisal. Coping appraisal was measured through eight items adapted from previous studies [13, 27, 28] intended to capture both response efficacy and self-efficacy. Response efficacy was assessed using a four-item scale (e.g., “Vaccination will reduce the likelihood of getting sick”; $\alpha = .88$), as well as self-efficacy (e.g., “When vaccination will be available, I will be able to get vaccinated”; $\alpha = .86$).

Fear of COVID-19. Feelings of fear were assessed using the 7-item (e.g., “My heart races when I think I might get COVID-19”; $\alpha = .86$) Italian version [29] of the Fear of COVID-19 Scale (FCV-19S) [30].

Intentions to get vaccinated. The intention to get vaccinated against COVID-19 was measured through 3 items (e.g., “When it will be possible, I will make myself available to get vaccinated against COVID-19”; $\alpha = .93$) adapted from previous studies conducted in different countries, including Italy [13, 27, 28].

Sociodemographic details. All the participants reported their age, sex, and nationality.

Data Analyses

We tested a model in which both COVID-19 and social limitations threat appraisals were posited as predictors of the intention to get vaccinated via fear of COVID-19, while coping appraisal was directly associated with the intention to perform the behaviour. We used bootstrapping to test mediation by estimating the presence and size of the indirect (i.e., mediated) effects [31]. As specified in the paragraph devoted to participants and procedure, we considered that a sample size of at least 200 participants would be required [26]. We adopted the following goodness-of-fit indices: the χ^2/df ratio, a good score of which was 2 or below; the Comparative Fit Index (CFI); the Incremental Fit

Index (IFI), the value of which should be higher than .95; the Normed Fit Index (NFI), a good score of which is more than .90; the Root Mean Square Error of Approximation (RMSEA); a 90% confidence interval for RMSEA (RMSEA 90% CI); and the Standardised Root Mean square Residual (SRMR). RMSEA and SRMR are considered acceptable if they are .08 or lower [32].

Results

Descriptive statistics are displayed in Table 1. For almost all variables participants reported a medium-high mean score.

Table 1. Descriptive statistics: Observed scores (*Min e Max*), Means (*M*) and Standard Deviations (*SD*) (N = 213).

Variables	Min-Max	M(<i>SD</i>)
TA- Social limitations Severity	1-5	3.94 (1.10)
TA- Social limitations Vulnerability	1-5	4.38 (.69)
TA-COVID-19 Severity	1-5	4.59 (.72)
TA-COVID-19 Vulnerability	1-5	3.14 (1.03)
CA-Self-efficacy	1-5	4.33 (.81)
CA-Response Efficacy	1-5	4.29 (.87)
Fear of COVID-19	1-5	2.47 (.86)
Intention to get vaccinated	1-5	4.54 (.89)

Note: TA= Threat Appraisal; CA= Coping Appraisal.

Table 2 presents the intercorrelations between variables, which were all positive and significant. The strongest intercorrelations were found between the two coping appraisals variables (self-efficacy and response efficacy) and the intention to get vaccinated. Fear was strongly associated with COVID-19 perceived vulnerability.

Table 2. Intercorrelation between all variables (N=213).

Variables	1.	2.	3.	4.	5.	6.	7.
1.TA-Social limitations Vulnerability	-						
2. TA-Social limitations Severity	.23**	-					
3. TA-COVID-19 Severity	.23**	.50**	-				
4. TA-COVID-19 Vulnerability	.23**	.33**	.32**	-			
5. CA-Self-efficacy	.29**	.42**	.42**	.29**	-		
6. CA-Response Efficacy	.36**	.60**	.57**	.37**	.62**	-	
7. Fear of COVID-19	.22**	.27**	.20*	.59**	.19**	.35**	-
8. Intention to get vaccinated	.24**	.47**	.47**	.21**	.66**	.71**	.26**

Note: **p < .01; ***p < .001. TA= Threat Appraisal; CA= Coping Appraisal.

The proposed model (Figure 1) fit very well with the data [$\chi^2/df = 1.98$, $p = .06$; RMSEA = .07 (CI = .01; .12); SRMR = .02; CFI = .99; IFI = .99; NFI = .98] and accounted for an important percentage of the variance of intentions (59%).

Partially in line with Hypothesis 1, people who believed that COVID-19 could have negative consequences for their social life and perceived themselves susceptible to the negative consequences of contracting the virus were more afraid about COVID-19. Specifically, social limitations perceived severity and COVID-19 perceived vulnerability were significantly and positively related to fear of COVID-19, and COVID-19 perceived vulnerability emerged as the strongest predictor of fear. Contrary to Hypothesis 1, COVID-19 perceived severity and social limitations perceived

vulnerability were not significantly related to fear of COVID-19. Moreover, fear did not predict the intention to get vaccinated.

In line with Hypothesis 2, individuals who were confident regarding their ability to get vaccinated and perceived that vaccine could reduce the negative consequences of COVID-19 were more predisposed to implement this adaptive behavior. Indeed, both perceived self-efficacy and response efficacy were significantly and positively associated with the intention to get vaccinated.




Figure 1. Final model

Note: * $p < .05$; *** $p < .001$. TA= Threat Appraisal; CA= Coping Appraisal

Study 2

Methods

Participants and procedure

The data were collected between 20 December 2021 and 10 January 2022, after the vaccine became available but before it became mandatory. The final sample was composed of 1111 Italian-speaking participants who completed an online survey (68.9% women; mean age = 38.33, $SD = 13.94$). For some analyses (i.e., discriminant analysis and hierarchical regression) we had to exclude approximately 5% of the initial sample due to the necessity of maintaining a balanced representation across different sex groups in the analyses, particularly when considering individuals who identified as transgender or non-binary. Given the limitations in obtaining a sufficiently large sample for these specific sex identities, we opted to exclude these cases from the analyses while retaining a focus on age and binary sex. This was to prevent the introduction of substantial biases due to the inherent imbalance in the group sizes. Moreover, since we aimed to detect even relatively small effects while modeling for the impact of sociodemographic and personality variables, we conducted a preliminary power analysis. The analysis revealed that, given the chosen set of predictors, a sample size of 1000 participants would be necessary to achieve a statistical power of 0.90, enabling us to identify even relatively small effects ($f^2 = 0.02$) as statistically significant. Since we were able to secure more than 1000 participants, we deemed our sample size adequate. The procedure was the same as Study 1. The survey took approximately 20 min to complete. To ensure participants had complete control over

whether to submit their responses or not, we did not record responses from participants who did not complete the survey. The Ethical Committee of the University of Florence approved the study procedures (Ref. No. 0092811).

Measures

In the following measures all the items were rated on a five-point Likert scale (1 = strongly disagree; 5 = strongly agree). Higher scores indicated higher levels in the measured variables.

Personality. The Italian version [33] of the Ten-Item Personality Inventory (I-TIPI) [34] was used. The scale is composed of ten items assessing five dimensions: extraversion (e.g., extraverted, enthusiastic), agreeableness (e.g., sympathetic, warm), conscientiousness (e.g., dependable, self-disciplined), emotional stability (e.g., calm, emotionally stable), and openness to experience (e.g., open to new experiences, complex).

Fear of vaccination. The Italian version [17] of the Fear of Vaccination Scale (VFS-6) [35] was used. The scale consists of six items comprising two dimensions: cognitive ($\alpha = .70$) and somatic ($\alpha = .74$).

Vaccination hesitancy. The 5-C scale [5] was used to assess vaccine hesitancy. The scale consists of 15 items which comprise 5 dimensions: confidence (i.e., trust in the effectiveness and safety of vaccines); complacency (i.e., vaccination not deemed as a necessary preventive action); constraints (i.e., reporting issues that prevent one from getting vaccinated); calculation (i.e., extensive engagement in evaluating the risks of infections and vaccination to derive a good decision); and collective responsibility (i.e., the willingness to protect others through one's own vaccination). Cronbach's α for the 5-C scale ranges from 0.67 to 0.88.

Sociodemographic details. All the participants reported their age, sex, and nationality.

Results

Descriptive statistics disaggregated by sex are presented in Table 3.

Table 3. Descriptive statistics of the collected variables for the entire sample and disaggregated by sex.

Variables	Entire Sample	Females	Males
	M (SD)	M (SD)	M (SD)
Age	38.33 (13.94)	38.05 (13.82)	38.67 (14.44)
Extraversion	8.19 (2.76)	8.38 (2.85)	7.74 (2.50)
Agreeableness	10.38 (2.03)	10.49 (2.01)	10.05 (2.08)
Conscientiousness	10.42 (2.36)	10.57 (2.28)	10.12 (2.44)
Neuroticism	7.62 (2.77)	7.90 (2.73)	6.94 (2.74)
Openness	9.07 (2.12)	9.09 (2.09)	4.37 (2.31)
VF Cognitive	7.61 (4.07)	7.74 (4.19)	6.94 (3.66)
VF Somatic	5.00 (3.01)	5.18 (3.19)	4.37 (2.31)
VH Confidence	10.21 (1.35)	10.18 (3.30)	10.53 (3.46)
VH Complacency	5.44 (2.83)	5.17 (2.60)	5.94 (3.20)
VH Constraints	4.77 (2.29)	4.66 (2.20)	4.82 (2.35)
VH Calculation	11.03 (2.65)	11.00 (2.53)	11.00 (2.91)
VH Collective responsibility	11.95 (3.00)	12.15 (2.84)	11.66 (3.26)

Note: M = Mean; s.d. = standard deviation; VF= Vaccination Fear; VH = Vaccination Hesitancy.

To test how much vaccine hesitancy, in its 5 components, was related to actual vaccination, we relied on discriminant analysis [36]. This technique allowed us to investigate the efficacy of a linear combination of vaccine hesitancy dimensions in differentiating between vaccinated and unvaccinated people. The data were therefore analysed using stepwise discriminant analysis and thus excluding non-statistically significant factors. The five selected indicators were entered

concurrently in the discriminant function analysis. The linear equation performed quite well after excluding the vaccination hesitancy constraints dimension which resulted as a non-statistically significant factor. In fact, Wilk's Lambda ($\Lambda=0.58$), chi-square values ($\chi^2_{(4)}=577.8$), and the level of statistical significance ($p<.001$) provided robust support for a linear discriminating function together with a canonical correlation of 0.65 and a predictive accuracy of the model of 88.1%. The relative contribution to the equation of each variable is reported in Table 4.

Table 4. Standardised discriminant function coefficients.

Predictors	Coefficient
VH Confidence	0.31
VH Complacency	-0.23
VH Constraints	ns
VH Calculation	-0.12
VH Collective responsibility	0.57

Note. N = 1061; VH = Vaccination Hesitancy; ns = not statistically significant.

The derived function was mainly composed of collective responsibility ($\beta=0.57$) and confidence dimensions ($\beta=0.31$). Complacency dimension appeared to make a secondary contribution to the equation ($\beta=-0.23$) while calculation was of less importance ($\beta=-0.12$).

Subsequently, we performed multiple regression to assess the best set of predictors for each of the vaccine hesitancy dimension that emerged connected to actual vaccination in discriminant analysis (see Table 5).

Table 5. Multiple regression of identified predictors with Vaccine Hesitancy dimensions.

Predictors	VH Confidence	VH Complacency	VH Calculation	VH Collective responsibility
Age	-0.04	-0.01	0.07*	-0.06*
Sex	0.01	-0.16**	-0.04	0.12**
Extraversion	0.02	0.01	0.02	-0.01
Agreeableness	0.01	-0.06*	0.07*	0.02
Conscientiousness	0.02	-0.02	0.04	0.01
Neuroticism	0.06**	-0.16**	-0.02	0.12**
Openness	-0.02	0.05*	0.01	-0.02
VF Cognitive	-0.76***	0.57***	0.38**	-0.60***
VF Somatic	-0.02	0.11***	0.01	-0.11***
R ²	0.60	0.46	0.17	0.49

Note: N = 1061; * = $p < .05$; ** = $p < .01$; *** = $p < .001$; VH = Vaccination Hesitancy; VF = Vaccination Fear Scale.

As shown in Table 5, the role of both socio-demographic variables (i.e., age and sex) and personality traits was quite limited in terms of the number of statistically significant results and their effect size. Age was positively correlated with the extent of calculation, suggesting that older participants tended to engage in a more thorough evaluation of the advantages and disadvantages of vaccination. Conversely, age exhibited a negative correlation with collective responsibility, indicating that older individuals might be somewhat less inclined to protect others through their own vaccination efforts.

Regarding gender, cisgender women displayed lower levels of hesitancy in two out of four dimensions of vaccination hesitancy. In contrast, traits such as extraversion and conscientiousness did not demonstrate a significant association with vaccination hesitancy, particularly when accounting for vaccine fear. Similarly, openness showed no consistent relationship with vaccination hesitancy, except for a positive association with a higher level of complacency.

Conversely, agreeableness was linked to reduced complacency and increased calculation. Lastly, neuroticism consistently exhibited a negative association with vaccination hesitancy, implying that individuals with higher Neuroticism scores tend to have lower levels of vaccination hesitancy. Instead, vaccination fear, mostly in its cognitive component, appeared to substantially explain the variance of the vaccination hesitancy subscales (more than 50% for confidence, complacency, and collective responsibility; 17% for calculation).

Discussion

Our first study showed that vaccination intentions were significantly predicted by coping appraisal determinants, namely response efficacy and self-efficacy, in line with PMT [3]. These findings suggest that people who believed the vaccine was useful and felt more confident regarding their ability to get vaccinated were also more willing to perform the behaviour. Notably, our findings showed that fear was associated not only with the perception of negative consequences of COVID-19 for one's health, but also for one's social life. Individuals who were worried about the consequences of not getting vaccinated in terms of limitations to mobility and social experiences reported higher levels of fear. Nevertheless, contrary to our hypothesis, fear of Covid-19 did not predict vaccination intention. While in previous research fear of Covid-19 mediated the link between threat appraisal and intention to follow the measures to reduce the spread of the virus [13], this same role did not emerge in relation to vaccination intention. This unexpected result may be attributed to the age of the participants, for whom levels of fear might be not so high to trigger motivation. Moreover, during the period in which the data were collected, the situation might have been evaluated as less dangerous than one year before, when Nerini and colleagues conducted their study (2021).

In Study 2 we moved from analysing the intention to get vaccinated to actual vaccination. In line with previous studies, vaccine hesitancy was negatively related to vaccination [37, 38]. More specifically, the more social-oriented dimension of collective responsibility was the strongest predictor of vaccination, thus highlighting that vaccination should not be considered a pure individual cognitive-based phenomenon. Indeed, the individual and social sphere are intertwined and interdependent with one another in defining human behaviour [39]. Notably, the cognitive component of vaccine fear was strongly and negatively associated with the collective responsibility dimension of vaccine hesitancy, so that perceived social pressure to comply with vaccination (e.g., being a good and respected member of the community) might push people to overlook their fears [40, 41].

We should acknowledge some limitations of our studies. First, in Study 1 perceived threat might have changed during the data collection period. Future studies could explore the temporal dynamics of fear in relation to vaccination intention to better understand the role of fear as a mediator. Second, although we considered the role of some social factors, we focused mostly on individual-level psychological ones. Future studies could investigate the interplay between individual and sociocultural variables (e.g., social norms, peer influence, community support), together with structural ones, in shaping vaccination behaviour. Just as with other non-pharmaceutical interventions (NPI) implemented during the pandemic, such as contact tracing methods, the influence of the social context emerged as a pivotal factor in either encouraging or impeding the adoption of specific behaviors. This context likely also contributed to the normalization of certain fears and beliefs (42 Guazzini et al., 2021). Moreover, the correlational nature of the studies limits causal interpretations. Longitudinal or experimental designs could be employed to establish causal and temporal relationships among variables. Finally, the present studies focused on specific populations and contexts. Replication of the findings would contribute to their generalizability thus providing a more comprehensive understanding of vaccination behaviour.

In conclusion, our research contributed to identify predictors of vaccination behaviour during the COVID-19 pandemic. Coping appraisal determinants, but not fear, emerged as predictors of vaccination intention when the vaccine was not available. When vaccination was possible, we found a negative relationship between vaccine hesitancy and actual vaccination, with collective

responsibility being a key predictor. These findings provide insights into the complexity of vaccine acceptance and emphasise the need for targeted interventions which consider not only individual but also social factors to promote vaccination and mitigate the spread of infectious diseases. A focus on social and interpersonal factors can be useful not only for increasing vaccination levels, but also for improving people's wellbeing during periods in which social life is limited (43 Matera et al., 2021). More specifically, the results suggested that messages emphasizing the aspect of collective responsibility towards vaccination might be highly desirable. Furthermore, by incorporating references to vaccine safety (confidence) and crafting messages and interventions that reassure individuals about the exaggeration of the likelihood and severity of adverse outcomes resulting from vaccination, along with providing cognitive coping tools to address such fears, we should be able to achieve more effective and evidence-based health communication. Finally, effective messages should address beliefs related to one's ability in performing the recommended behavior (i.e., get vaccinated), describing it as an easy behavior to implement, thus making people confident that they can get vaccinated without too much efforts and difficulties.

Author Contributions: Amanda Nerini, Conceptualization; Formal analysis; Methodology; Supervision; Roles/Writing - original draft; and Writing - review & editing; Mirko Duradoni, Formal analysis; Methodology; Roles/Writing - original draft; and Writing - review & editing; Camilla Matera, Writing - review & editing; Andrea Guazzini, Writing - review & editing; Monica Paradisi, Methodology; Roles/Writing - original draft; Adriele Schembri, Investigation, Roles/Writing - original draft

Funding: This research received no external funding

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of the University of Florence (Study 1: Ref.No.0096243, 8th July 2020; Study 2: Ref.No.0092811, 12th June 2020)

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data availability statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization. *WHO Director-General's opening remarks at the media briefing on COVID-19*. Retrieved on March 11, 2020. <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>
2. Ministry of Health. *Fase 2, Governo vara nuovo Dpcm su riapertura*. Retrieved on November 19, 2021. <https://www.salute.gov.it/portale/nuovocoronavirus/dettaglioNotizieNuovoCoronavirus.jsp?lingua=italiano&menu=notizie&p=dalministero&id=4772>
3. Rogers, R.W. A protection motivation theory of fear appeals and attitude change, *Journal of Psychology*, **1975**, 91, 93–114.
4. Rogers, R.W. Cognitive and physiological processes in fear appeals and attitude change: a revised theory of protection motivation. In J.T. Cacioppo and R.E. Petty (eds) *Social Psychophysiology: A Source Book*. Guilford Press, New York, 1983, 153–176.
5. Betsch, C., Schmid, P., Heinemeier, D., Korn, L., Holtmann, C., & Böhm, R. Beyond confidence: Development of a measure assessing the 5C psychological antecedents of vaccination. *PLoS one*, **2018**, 13(12), e0208601.
6. Boss, S. R., Galletta, D. F., Lowry, P. B., Moody, G. D., & Polak, P. What do systems users have to fear? Using fear appeals to engender threats and fear that motivate protective security behaviors. *MIS Quarterly*, **2015**, 39(4), 837–864.
7. Floyd, D. L., Prentice-Dunn, S., & Rogers, R. W. A meta-analysis of research on protection motivation theory. *Journal of applied social psychology*, **2000**, 30(2), 407–429.
8. Ansari-Moghaddam, A., Seraji, M., Sharafi, Z., Mohammadi, M., & Okati-Aliabad, H. The protection motivation theory for predict intention of COVID-19 vaccination in Iran: a structural equation modeling approach. *BMC Public Health*, **2021**, 21(1), 1–9.
9. Li, L., Wang, J., Nicholas, S., Maitland, E., Leng, A., & Liu, R. The Intention to Receive the COVID-19 Vaccine in China: Insights from Protection Motivation Theory. *Vaccines*, **2021**, 9(5), 445.

10. Tong, K. K., He, M., Wu, A., Dang, L., & Chen, J. H. Cognitive Factors Influencing COVID-19 Vaccination Intentions: An Application of the Protection Motivation Theory Using a Probability Community Sample. *Vaccines*, **2021**, 9(10), 1170.
11. Wang, P. W., Ahorsu, D. K., Lin, C. Y., Chen, I. H., Yen, C. F., Kuo, Y. J., ... & Pakpour, A. H. Motivation to have covid-19 vaccination explained using an extended protection motivation theory among university students in china: The role of information sources. *Vaccines*, **2021**, 9(4), 380.
12. Xiao, Q., Liu, X., Wang, R., Mao, Y., Chen, H., Li, X., ... & Zheng, P. (2021). Predictors of Willingness to Receive the COVID-19 Vaccine after Emergency Use Authorization: The Role of Coping Appraisal. *Vaccines*, **2021**, 9(9), 967.
13. Nerini, A., Matera, C., Policardo, G. R., & Di Gesto, C. Containment measures against COVID-19 in Italy: The role of protection motivation and values. *Journal of Human Behavior in the Social Environment*, **2021**, 1-14.
14. Diaz, P., Zizzo, J., Balaji, N. C., Reddy, R., Khodamoradi, K., Ory, J., & Ramasamy, R. Fear about adverse effect on fertility is a major cause of COVID-19 vaccine hesitancy in the United States. *Andrologia*, **2022**, 54(4), e14361.
15. Montagni, I., Ouazzani-Touhami, K., Mebarki, A., Texier, N., Schüeck, S., Tzourio, C., & Confins Group. Acceptance of a Covid-19 vaccine is associated with ability to detect fake news and health literacy. *Journal of public health*, **2021**, 43(4), 695-702.
16. Muric, G., Wu, Y., & Ferrara, E. COVID-19 vaccine hesitancy on social media: building a public Twitter data set of antivaccine content, vaccine misinformation, and conspiracies. *JMIR public health and surveillance*, **2021**, 7(11), e30642.
17. Duradoni, M., Veloso, M. V., La Gamma, M., Monciatti, A. M., & Guazzini, A. (2022). Italian version of the Vaccination Fear Scale (VFS-6): internal and external validation. *Mediterranean Journal of Clinical Psychology*, **2022**, 10(2).
18. Hortal, M., & Di Fabio, J. L. Vaccine rejection and vaccination management: the grey areas Recusa vacinal e gestão da imunização: nuances e contrastes. *Revista panamericana de salud publica= Pan American journal of public health*, **2019**, 43, e54-e54.
19. Mellers, B. A., & McGraw, A. P. Anticipated emotions as guides to choice. *Current directions in psychological science*, **2001**, 10(6), 210-214.
20. Sato, R., & Fintan, B. Fear, knowledge, and vaccination behaviors among women in Northern Nigeria. *Human Vaccines & Immunotherapeutics*, **2020**, 16(10), 2438-2448.
21. Tahir, A. I., Ramadhan, D. S., Taha, A. A., Abdullah, R. Y., Karim, S. K., Ahmed, A. K., & Ahmed, S. F. Public fear of COVID-19 vaccines in Iraqi Kurdistan region: a cross-sectional study. *Middle East Current Psychiatry*, **2021**, 28, 1-8.
22. SAGE Working Group. Report of the SAGE working group on vaccine hesitancy. *World Health Organization*, **2014**.
23. World Health Organization. *Ten Threats to Global Health in 2019*. Retrieved on October 3, **2022**. <https://www.who.int/news-room/spotlight/ten-threats-to-global-health-in-2019>
24. Howard, M. C. The good, the bad, and the neutral: Vaccine hesitancy mediates the relations of Psychological Capital, the Dark Triad, and the Big Five with vaccination willingness and behaviors. *Personality and individual differences*, **2022**, 190, 111523.
25. Murphy, J., Vallières, F., Bentall, R. P., Shevlin, M., McBride, O., Hartman, T. K., ... & Hyland, P. Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom. *Nature communications*, **2021**, 12(1), 29.
26. Weston, R., & Gore Jr, P. A. A brief guide to structural equation modeling. *The counseling psychologist*, **2006**, 34(5), 719-751.
27. Bashirian, S., Jenabi, E., Khazaei, S., Barati, M., Karimi-Shahanjarini, A., Zareian, S., ... & Moeini, B. Factors associated with preventive behaviours of COVID-19 among hospital staff in Iran in 2020: an application of the Protection Motivation Theory. *Journal of Hospital Infection*, **2020**, 105(3), 430-433.
28. Roma, P., Monaro, M., Muzi, L., Colasanti, M., Ricci, E., Biondi, S., ... & Mazza, C. How to improve compliance with protective health measures during the COVID-19 outbreak: testing a moderated mediation model and machine learning algorithms. *International journal of environmental research and public health*, **2020**, 17(19), 7252.
29. Soraci, P., Ferrari, A., Abbiati, F. A., Del Fante, E., De Pace, R., Urso, A., & Griffiths, M. D. Validation and psychometric evaluation of the Italian version of the Fear of COVID-19 Scale. *International journal of mental health and addiction*, **2020**, 1-10.
30. Ahorsu, D. K., Lin, C. Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. The fear of COVID-19 scale: development and initial validation. *International journal of mental health and addiction*, **2020**, 1-9.
31. Rucker, D. D., Preacher, K. J., Tormala, Z. L., & Petty, R. E. Mediation analysis in social psychology: Current practices and new recommendations. *Social and personality psychology compass*, **2011**, 5(6), 359-371.

32. Hooper, D., Coughlan, J., & Mullen, M.R. Structural equation modelling: Guidelines for determining model fit. *Electronic Journal of Business Research Methods*, **2008**, *6*, 53–60.
33. Chiorri, C., Bracco, F., Piccinno, T., Modafferi, C., & Battini, V. Psychometric properties of a revised version of the Ten Item Personality Inventory. *European Journal of Psychological Assessment*, **2015**.
34. Gosling, S. D., Rentfrow, P. J., & Swann, W. B. A Very Brief Measure of the Big Five Personality Domains. *Journal of Research in Personality*, **2003**, *37*, 504–528.
35. Malas, O., & Tolsá, M. D. Vaccination Fear Scale (VFS-6): Development and Initial Validation. *Mediterranean Journal of Clinical Psychology*, **2021**, *9*(2).
36. Everitt, B. S., Landau, S., Leese, M., & Stahl, D. *Cluster analysis*. John Wiley & Sons, 2011.
37. Brumbaugh, J. T., Sokoto, K. C., Wright, C. D., Francis, S. E., Hubbard, J., Alexander, L., & McNeil, D. W. Vaccination intention and uptake within the Black community in Appalachia. *Health Psychology*, **2023**.
38. Simone, L., Vagni, M., Maiorano, T., Giostra, V., & Pajardi, D. How Implicit Attitudes toward Vaccination Affect Vaccine Hesitancy and Behaviour: Developing and Validating the V-IRAP. *International Journal of Environmental Research and Public Health*, **2022**, *19*(7), 4205.
39. Lewin, K. Field theory in social science: selected theoretical papers. Ed: Dorwin Cartwright, 1951.
40. Deutsch, M., & Gerard, H. B. A study of normative and informational social influences upon individual judgment. *The journal of abnormal and social psychology*, **1955**, *51*(3), 629.
41. Kelman, H. C. Compliance, identification, and internalization three processes of attitude change. *Journal of conflict resolution*, **1958**, *2*(1), 51–60.
42. Guazzini, A., Fiorenza, M., Panerai, G., & Duradoni, M. What went wrong? Predictors of contact tracing adoption in Italy during COVID-19 pandemic. *Future Internet*, **2021**, *13*(11), 286.
43. Matera, C., Paradisi, M., Boin, J., & Nerini, A. Perceived mattering in the face of COVID-19: links with emotion regulation and psychological well-being. *Psychology & Health*, **2022**, *37*(12), 1471–1491.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.