

Article

Not peer-reviewed version

Identification of Chromoblastomycosis and Phaeohyphomycosis Agents Through ITS-RFLP

[Gabriel S M. Sousa](#)*, Rodrigo De Oliveira, Alex B. De Souza, Ruan C. Monteiro, Elaine P T E. Santo, Luciano Franco Filho, Silvia H M. Da Silva

Posted Date: 26 December 2023

doi: 10.20944/preprints202312.1779.v1

Keywords: chromoblastomycosis; phaeohyphomycosis; PCR-RFLP; Herpotrichiellaceae; sequencing; phenotypic; molecular



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Identification of Chromoblastomycosis and Phaeohyphomycosis Agents through ITS-RFLP

Gabriel S M. Sousa ^{1,*}, Rodrigo De Oliveira ², Alex B. De Souza ², Ruan C. Monteiro ³, Elaine P T E. Santo ², Luciano Franco Filho ² and Silvia H M. Da Silva ²

¹ Programa de Pós-graduação em Biologia de Agentes Infecciosos e Parasitários, Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém 66075750, Brasil.

² Laboratório de Micoses Superficiais e Sistêmicas, Sessão de Bacteriologia e Micologia, Instituto Evandro Chagas, Ananindeua 67030000, Brasil; rodrigooliveira@iec.gov.br (R.O); alexbritosouza@yahoo.com.br (A.B.S); elainetavares@iec.gov.br (E.P.T.E.S); lucianofranco6@gmail.com (L.F.F); silviasilva@iec.gov.br (S.H.M.S).

³ Laboratory of Emerging Fungal Pathogens, Universidade Federal de São Paulo, São Paulo 1108723, Brasil; rcmonteiro@unifesp.br (R.C.M).

* Correspondence: gabriel.sousa@icb.ufpa.br

Abstract: Chromoblastomycosis (CBM) and phaeohyphomycosis (FEO) are infections caused by melanized filamentous fungal agents, primarily found in tropical and subtropical regions. Both infections pose significant challenges for the correct identification of the causative agent due to their morphological similarity, making conventional methods of morphological analysis highly subjective. Therefore, molecular techniques are necessary for the precise determination of these species. In this regard, this study aimed to contribute to a new methodology based on PCR-RFLP for the identification of agents causing CBM and FEO. Sequences from the ITS region were used to identify potential restriction enzyme sites *in silico*, followed by *in vitro* validation using the selected restriction enzymes. The obtained results were compared with species identification through morphological analyses and sequencing. The results demonstrated that the PCR-RFLP applied in this study accurately identified two major agents of chromoblastomycosis, *Fonsecaea pedrosoi* and *Fonsecaea monophora*, as well as *Cladophialophora bantiana* and *Exophiala dermatitidis*, both causative agents of phaeohyphomycosis. In this context, the proposed assay can complement current methods for identifying these species, aiding in diagnosis, and contributing to the proper management of these infections.

Keywords: chromoblastomycosis; phaeohyphomycosis; PCR-RFLP; *Herpotrichiellaceae*; sequencing; phenotypic; molecular

1. Introduction

Chromoblastomycosis (CBM) and phaeohyphomycosis (PHM) are fungal infections primarily instigated by dematiaceous fungi, predominantly classified under the *Herpotrichiellaceae* family [1,2]. CBM is categorized by the World Health Organization as a neglected tropical disease, primarily due to its higher prevalence in economically disadvantaged or developing countries situated within tropical or subtropical regions [3–5].

CBM prevails endemically in countries spanning Latin America, Central America, Africa, and Asia, with a notable predilection for rural laborers [2,3,6,7]. In contrast, PHM is reported in diverse nations globally and exhibits a similar affinity for tropical and subtropical climates, predominantly affecting immunocompromised individuals, thus denoting it as an opportunistic infection [1,8,9].

Chromoblastomycosis and phaeohyphomycosis are both mycoses primarily transmitted through direct contact with contaminated materials, often resulting from traumatic injuries. These materials may encompass plant thorns, branches, soil, and various other organic components [1,2].

The clinical manifestations of both infections are intricate, often mirroring other maladies, thereby complicating the process of differential diagnosis. Consequently, laboratory support is frequently imperative [8,10,11]. In this context, the prevailing diagnostic gold standard for both

infections are the identification of muriform cells in the case of chromoblastomycosis and dematiaceous hyphae in the case of phaeohyphomycosis through direct mycological examination [1,2,12].

In the presumptive species-level identification, morphological analyses are conducted based on the macrocolony's appearance and the arrangement and structure of conidial and hyphal elements [2,13,14]. However, due to the phenotypic similarities between the species responsible for CBM and PHM, such analyses become notably subjective [8,14].

Owing to this subjectivity, genetic sequencing of the Internal Transcribed Spacer (ITS) region has emerged as the most widely employed molecular technique for identifying agents of chromoblastomycosis and phaeohyphomycosis [1,2]. Nevertheless, this methodology is associated with high costs and is relatively inaccessible in regions where these infections are more endemic. Consequently, there is a pressing need for new, cost-effective molecular approaches suitable for resource-limited laboratories, enabling rapid and specific species identification [15].

In light of these considerations, the present study aimed to develop a molecular approach based on restriction fragment length polymorphism of the ITS region for the identification of species among the primary causative agents of chromoblastomycosis and phaeohyphomycosis in the Northern region of Brazil.

2. Materials and Methods

2.1. Clinical strains

The isolates from chromoblastomycosis (N=22) and phaeohyphomycosis (n=2) examined in the present study are maintained in the mycoteca of superficial and systemic mycoses at the Instituto Evandro Chagas (IEC) in the state of Pará, Brazil. The representative fungal agents were originally obtained from clinical samples sourced from healthcare units and hospitals within the state of Pará, as evidenced in Table 1. These agents have already been molecularly identified, and their sequences have been deposited in the GenBank platform of NCBI.

Table 1. Isolates causing chromoblastomycosis and phaeohyphomycosis used in this study.

Isolate	Genbank	Anatomical Site	Host	Geographic Origin
IEC-CBM02	KY312523	Thigh	Human	Pará/Brazil
IEC-CBM03	KY312524	Foot	Human	Pará/Brazil
IEC-CBM04	KY312525	Leg	Human	Pará/Brazil
IEC-CBM05	KY312526	Leg	Human	Pará/Brazil
IEC-CBM06	KY312527	CNS+	Human	Pará/Brazil
IEC-CBM07	KY312528	Hand	Human	Pará/Brazil
IEC-CBM08	KY312529	Leg	Human	Pará/Brazil
IEC-CBM09	KY312530	*	Human	Pará/Brazil
IEC-CBM10	KY312531	Arm	Human	Pará/Brazil
IEC-CBM11	KY312532	Leg	Human	Pará/Brazil
IEC-CBM12	KY312533	Foot	Human	Pará/Brazil

IEC-CBM13	KY312534	Leg	Human	Pará/Brazil
IEC-CBM14	KY312535	*	Human	Pará/Brazil
IEC-CBM15	KY312536	*	Human	Pará/Brazil
IEC-CBM16	KY312537	Thigh	Human	Pará/Brazil
IEC-CBM18	MF416919	Thigh	Human	Pará/Brazil
IEC-CBM19	MF416920	Forearm	Human	Pará/Brazil
IEC-CBM21	MF416922	Fist	Human	Pará/Brazil
IEC-CBM22	MF416923	Ankle	Human	Pará/Brazil
IEC-CBM23	MF416924	*	Human	Pará/Brazil

* Denotes unknown information; † Central Nervous System.

2.2. Morphological Identification

The morphological characteristics of the isolates were assessed through microcultivation on slides, employing lactrimel agar medium, and incubated at 37°C for 14 days. Subsequently, the slides were scrutinized with lactophenol cotton blue staining, utilizing 40x objectives, for the documentation of conidial arrangement organization [14,16].

2.3. DNA Extraction

The cultures were subcultured in a tube containing YPD Agar and incubated at 30°C for 14 days for DNA extraction. The extraction process involved collecting approximately 400 mg of fungal mass, which was then added to a 2 mL containing a 450 µL solution composed of 150 µL lysis buffer (SDS), 150 µL homogenization buffer, and 150 µL TE buffer. Glass beads were added, and the microtube was vortexed for 30 minutes. Subsequently, 15 µL of proteinase K were added, and the microtube was incubated in a water bath at 57°C for one hour. Following this, 200 µL of 5 mol/L sodium chloride were added, and the microtube was incubated again at 67°C for 10 minutes.

After incubation, 600 µL of the liquid was transferred to a new microtube and purified using the phenol-chloroform-isoamyl alcohol protocol described by Campos in 2017 [17]. To achieve enhanced DNA purity, the Bioflux DNA purification kit (Hangzhou Bioer Technology Co. Ltd, Hangzhou, China) was employed in conjunction with the product obtained through the phenol-chloroform methodology, adhering to the procedures delineated in the manufacturer's protocol.

The extracted DNA was quantified using the NanoDrop 2000© spectrophotometer (Thermo Fisher Scientific Inc.®, Waltham, MA, USA). We used the standard value of 1 OD = 50 µg/ml to determine the double-stranded DNA concentration. Only samples with an OD_{260/280} ratio between 1.7 and 2.0 were included in the study.

2.4. Molecular Identification through Sequencing

Molecular sequencing of the isolates was performed to confirm the previously identified species, focusing on the ITS region, which includes ITS1, 5.8S, and ITS2, using ITS1(F) TCCGTAGGTGAACCTGCGG and ITS4(R) TCCTCCGCTTATTGATATGC primers. The PCR conditions consisted of an initial denaturation at 95°C for 5 minutes, followed by 35 cycles of denaturation at 94°C for 1 minute, primer annealing at 55.5°C for 2 minutes, and extension at 72°C for 2 minutes. Finally, a 10-minute extension phase at 72°C was conducted [18].

The amplification reaction was carried out with 4 mM MgCl₂, 0.4 mM of each dNTP (deoxynucleotide triphosphate), 1 mM of the primers, 0.1 µL of Taq DNA polymerase (Thermo Fisher

Scientific Inc.®, Waltham, MA, USA), 2.5 µL of 10mM/L BSA, and 2 µL of DNA, in a final volume of 25 µL. PCR was performed using a PX2 Thermo Hybaid thermocycler (Artisan Technology Group, Champaign, IL, USA). The amplification product was visualized by electrophoresis on a 1.5% agarose gel.

Following electrophoresis, amplicon purification was carried out using the ExoSAP-IT™ PCR Product Cleanup Reagent (Thermo Fisher Scientific Inc.®, Waltham, MA, USA), following the manufacturer's recommendations. Sequencing was performed using the BigDye™ Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific Inc.®, Waltham, MA, USA), following the manufacturer's instructions, and the samples were sequenced on an ABI 3130/3130XL Automatic Genetic Analyzer using the dideoxynucleotide chain termination method (Thermo Fisher Scientific Inc.®, Waltham, MA, USA).

Identification was accomplished by aligning the sequences with those deposited in the GenBank and ISHAM databases, referencing type strains and considering a similarity value greater than 99% for species determination.

2.5. *In silico* ITS-RFLP

In order to identify enzymes with the potential for species differentiation among the etiological agents of chromoblastomycosis and phaeohyphomycosis, an *in silico* screening was conducted using sequences from the ITS region of the study's target species (N=60) available in the GenBank database. These sequences were downloaded and compiled into a MultiFasta file and aligned with the online software Mafft 7 (<https://mafft.cbrc.jp/alignment/server/index.html>) [19].

Subsequently, the presence of restriction sites for commercially available enzymes was analyzed using the CLC Sequence Viewer software 8.0® [20], with the goal of generating fragments of distinct sizes suitable for species differentiation while ensuring the consistency of restriction sites within the same species. Following the screening and identification of potential enzymes, the online tool NEBcutter v.3® from New England Biolabs (<https://nc3.neb.com/NEBcutter/>) was employed for visualizing the fragmentation pattern in a virtual agarose gel [21].

2.6. ITS-RFLP

The Internal Transcribed Spacer (ITS) fragments were amplified using the primer pair ITS1F (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4R (5'TCCTCCGCTTATTGATATGC -3'), as previously described. The negative control for the reaction contained all PCR components except the DNA of the agents responsible for chromoblastomycosis and phaeohyphomycosis. PCR products were visualized through electrophoresis on a 1.5% agarose gel using a UV transilluminator. Subsequently, restriction enzyme digestion was conducted in a final volume of 40 µl, following the reaction conditions recommended by the manufacturer (Thermo Fisher Scientific Inc.®, Waltham, MA, USA).

In the single enzymatic digestion, a mixture of 3 µl of Red Buffer (10x) and 1 µl of *HhaI* enzyme (10U/µl) was added for every 20 µl of PCR product. In the double digestion reaction, 1 µl of *HaeIII* enzyme (10U/µl), 1 µl of *HhaI* enzyme (10U/µl), and 3 µl of Red Buffer (10x) were added for every 20 µl of PCR product. A negative control reaction was also performed for each digestion, containing all reagents except the PCR product.

Both reactions were incubated at 37°C for 1 hour. The digestion products were then subjected to electrophoresis on a 3% (w/v) agarose gel for 120 minutes, and the fragments were visualized under a UV transilluminator.

3. Results

3.1. Morphological Identification and Sequencing of the Isolates

In this study, a total of 24 isolates originating from chromoblastomycosis and phaeohyphomycosis were utilized. Morphologic identification through micro cultivation analysis allowed for the determination of their genus-level classification, resulting in 21 isolates belonging to the genus *Fonsecaea* sp., 1 to the genus *Exophiala* sp., and 1 to the genus *Cladophialophora* sp.

Molecular identification through sequencing of the ITS region revealed that out of the 24 isolates, 18 were identified as *Fonsecaea pedrosoi*, 4 as *Fonsecaea monophora*, 1 as *Exophiala dermatitidis*, and 1 as *Cladophialophora bantiana*.

3.2. In Silico Analyses

Through in silico analyses, we have concluded that the enzymes *HhaI* and *HaeIII* are capable of distinguishing between the species in this study through the restriction of the ITS region. These enzymes can be used in two different assays, the first being a single digestion with *HhaI* and the second being a double digestion with *HhaI* and *HaeIII*. The two proposed assays differ in fragmentation patterns and their ability to distinguish between the species, as shown in Table 2.

Table 2. Comparison of the fragments obtained in in silico restrictions for the identification of Species *Fonsecaea pedrosoi*; *Fonsecaea monophora*; *Cladophialophora bantiana*; *Exophiala dermatitidis*.

Genbank Access	Specie	HhaI	HhaI and HaeIII
NR_130652.1	<i>Fonsecaea pedrosoi</i>	352, 219 bp	202, 130, 89, 52, 50, 48 bp
NR_131280.1	<i>Fonsecaea monophora</i>	330, 188, 53 bp	203, 130, 58, 53, 52, 48, 27 bp
KY312527.1	<i>Cladophialophora bantiana</i>	343, 186, 42 bp	182, 127, 95, 59, 49, 42, 17 bp
KY312530.1	<i>Exophiala dermatitidis</i>	193, 193, 135, 98 bp	193, 121, 97, 48, 32, 30, 28, 16, 6 bp

3.3. ITS-RFLP with *HhaI*

The in silico results of the single digestion with the *HhaI* enzyme were compared with the fragmentation patterns obtained in vitro (Figure 1), where the results were confirmed, allowing for the differentiation between the species *Fonsecaea pedrosoi* (352, 219 bp) and *Fonsecaea monophora* (330, 188, 53 bp). Fragmentation patterns also differed for the species *Exophiala dermatitidis* (193, 193, 135, 98 bp). However, it was not possible to distinguish between the species *Fonsecaea monophora* and *Cladophialophora bantiana* in the single digestion.




Figure 1. Comparison of fragmentation patterns obtained with enzymatic restriction using *HhaI* in silico and in vitro. A: *Fonsecaea pedrosoi*; B: *Fonsecaea monophora*; C: *Cladophialophora bantiana*; D: *Exophiala dermatitidis*.

3.4. ITS-RFLP com *HhaI* e *HaeIII*

A comparison was made between the double digestion fragmentation patterns obtained *in silico* and *in vitro* (Figure 2), where a similar results to that obtained with the single digestion was achieved for all species, but with a greater power of distinction between the species *Fonsecaea monophora* (203, 130, 58, 53, 52, 48, 27 bp) and *Cladophialophora bantiana* (182, 127, 95, 59, 49, 42, 17 bp).




Figure 2. Comparison of fragmentation patterns obtained with enzymatic restriction using *HhaI* and *HaeIII*. A: *Fonsecaea monophora*; B: *Fonsecaea pedrosoi*; C: *Cladophialophora bantiana*; D: *Exophiala dermatitidis*.

3.5. Comparative Analysis of Identification Methods

The results obtained in both enzyme restriction assays were compared to other identification methods employed in this study (Morphologic and Sequencing). It was observed that the ITS-RFLP method exhibited a higher species identification capability compared to the phenotypic method. Furthermore, the ITS-RFLP method demonstrated a species identification capacity similar to sequencing, as described in Table 3.

Table 3. Comparison of Identification Results between Morphological Analysis, ITS-RFLP (*HhaI*), ITS-RFLP (*HhaI* + *HaeIII*), and Sequencing of the ITS Region.

Sample	Morphological Identification	ITS-RFLP (<i>HhaI</i>)	ITS-RFLP (<i>HhaI</i> + <i>HaeIII</i>)	Sequencing Identification
IEC-CBM02	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM03	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM04	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM05	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM06	<i>Cladophialophora</i> sp.	<i>C. bantiana</i> / <i>F. monophora</i> *	<i>C. bantiana</i>	<i>C. bantiana</i>
IEC-CBM07	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM08	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM09	<i>Exophiala</i> sp.	<i>E. dermatitidis</i>	<i>E. dermatitidis</i>	<i>E. dermatitidis</i>

IEC-CBM10	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM11	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM12	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM13	<i>Fonsecaea</i> sp.	<i>F. monophora</i>	<i>F. monophora</i>	<i>F. monophora</i>
IEC-CBM14	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM15	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM16	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM17	<i>Fonsecaea</i> sp.	<i>F. monophora</i>	<i>F. monophora</i>	<i>F. monophora</i>
IEC-CBM18	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM19	<i>Fonsecaea</i> sp.	<i>F. monophora</i>	<i>F. monophora</i>	<i>F. monophora</i>
IEC-CBM21	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM22	<i>Fonsecaea</i> sp.	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>	<i>F. pedrosoi</i>
IEC-CBM23	<i>Fonsecaea</i> sp.	<i>F. monophora</i>	<i>F. monophora</i>	<i>F. monophora</i>

* Inconclusive Result due to similar patterns in two different species.

4. Discussion

Infections caused by melanized fungi can significantly impact people's quality of life, leading to discomfort, pain, and, in severe cases, disfigurement. Both chromoblastomycosis (CBM) and phaeohyphomycosis (FEO), when affecting the skin and subcutaneous tissues, can be persistent and challenging to treat, often requiring prolonged therapies and, in some cases, surgical interventions [1,8,22–24]. In immunocompromised patients, such as those with HIV/AIDS, uncontrolled diabetes, or transplant recipients, agents causing phaeohyphomycosis can disseminate to internal organs and the central nervous system, posing a severe life-threatening risk [1,25,26].

Diagnosing both infections is notoriously challenging and difficult, with chromoblastomycosis typically taking an average of nine years to be correctly identified. In many cases, the causative species remains unidentified [3], a crucial concern since recent studies have highlighted variations in pathogenicity, virulence, tropism, and susceptibility to antifungals among these species [1,2,27]. Therefore, identifying the causative agent is essential for a more accurate prognosis [28].

The lack of identification of these agents in many cases occurs due to the phenotypic and morphological similarity of the species associated with these infections, rendering conventional morphological analysis methods nonspecific. For that reason, applying sequencing and other molecular biology techniques is necessary in most cases [1,8,14,28].

It is evident that the identification of species causing CBM and FEO poses a challenge to overcome, and the use of molecular techniques can help overcome this barrier. In this context, RFLP emerges as a simple and promising technique for the identification of CBM and FEO agents. Recent studies have shown that this technique serves as a valuable tool for identifying a variety of pathogenic

fungal species, such as *Cryptococcus* sp., Dermatophytes, *Paracoccidioides* sp., and *Candida* sp. [20,29–31].

Currently, there is no existing literature on assays for the identification of CBM and FEO agents using RFLP, with only isolated studies of specific primers either do not encompass all currently identified agents or lack sensitivity and specificity [32].

Due to this gap, the methodology proposed in this study aimed to contribute to the standardization of a species-specific diagnostic method using RFLP of the ribosomal DNA ITS region, known for its high conservation in fungi and currently considered a pan-fungal barcode [33–35]. This region is also frequently used for the identification of fungi from the order Chaetothyriales, where the predominant agents of chromoblastomycosis and phaeohyphomycosis are found [1,2,36,37].

Furthermore, a recent study published by Fonseca in 2022 [30], highlighted the efficiency of the ITS region in a simple digestion RFLP assay for identifying species of dermatophytes of the genera *Trichophyton* sp. and *Microsporum* sp. A comparison between the results obtained through enzymatic restriction and morphological identification showed the greater precision of the molecular technique compared to macroscopic and microscopic analyses, emphasizing the importance of this technique in distinguishing cryptic species of filamentous fungi.

In this context, the molecular identification assays by ITS-RFLP conducted in this study have proven to accurately identify the main species responsible for CBM and FEO, showing greater specificity when compared to morphological analyzes methods. When we compare the RFLP results with those obtained by sequencing, we observe that simple digestion was able to accurately identify 95.8% of the species used in this study, while double digestion accurately identified 100% of the agents, making this technique promising for the molecular identification of these species without the need for sequencing.

Therefore, the use of multiple enzymes resulted in more precise outcomes, attributed to increased fragmentation of the target region, providing better differentiation of genetically close species such as *Fonsecaea monophora*, *Fonsecaea pedrosoi*, and *Cladophialophora bantiana*, all belonging to the same phylogenetic clade [2].

In this context, ITS-RFLP accurately identified two significant species responsible for chromoblastomycosis: *F. pedrosoi* and *F. monophora*, both belonging to the genus *Fonsecaea* [2,38]. *Fonsecaea* sp. is responsible for over 80% of chromoblastomycosis cases worldwide³. Specifically, *Fonsecaea pedrosoi* is responsible for 84.1% of registered cases in Latin America and the Caribbean [7].

Besides the agents of chromoblastomycosis, it is possible to identify two species responsible for phaeohyphomycosis. The first is *Exophiala dermatitidis*, a causative agent of cutaneous, subcutaneous, and invasive infections [26,39,40], which, in severe cases, can affect the central nervous system [41]. The second identifiable species is *Cladophialophora bantiana*, a causative agent of PHM with high lethality due to invasive infections, mainly affecting the central nervous system of immunocompromised and post-transplant patients [25,40–42].

5. Conclusions

Our results introduce a novel molecular identification method based on ITS-RFLP for the two major causative agents of chromoblastomycosis worldwide and for two significant agents causing opportunistic and invasive phaeohyphomycosis.

As the causative agents of these infections differ in terms of virulence, pathogenicity, tropism, and susceptibility to antifungals, necessitating their identification for a more accurate prognosis of the infection and guide appropriate therapy, preventing serious complications such as infection dissemination or an inadequate treatment response.

In this context, the identification of these species through ITS-RFLP expands the range of molecular identification possibilities for these agents, providing rapid results at a lower cost compared to sequencing. This enables the improvement of accuracy and speed in diagnosis, ensuring proper management of these fungal infections.

Author Contributions: Conceptualization, G.S.M.S., R.O., R.C.S. and S.H.M.S.; data curation E.P.T.E.S and G.S.M.S.; formal analysis, G.S.M.S. and A.B.S.; supervision, S.H.M.S.; writing - original draft, G.S.M.S.; writing - review & editing, R.O., L.F.F. and S.H.M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by Evandro Chagas Institute.

Institutional Review Board Statement: Not applicable.

Data Availability Statement: No data were reported in this study.

Acknowledgments: We express gratitude to the Evandro Chagas Institute, Ananindeua, PA, Brazil, for providing the necessary supplies, laboratory infrastructure, and strains of interest for the execution of this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Arcobello, J. T.; Revankar, S. G. Phaeohyphomycosis. *Semin Respir Crit Care Med* **2020**, *41* (1), 131–140. <https://doi.org/10.1055/s-0039-3400957>.
2. Queiroz-Telles, F.; de Hoog, S.; Santos, D. W. C. L.; Salgado, C. G.; Vicente, V. A.; Bonifaz, A.; Roilides, E.; Xi, L.; Azevedo, C. de M. P. E. S.; Da Silva, M. B.; Pana, Z. D.; Colombo, A. L.; Walsh, T. J. Chromoblastomycosis. *Clin Microbiol Rev* **2017**, *30* (1), 233–276. <https://doi.org/10.1128/CMR.00032-16>.
3. Santos, D. W. C. L.; de Azevedo, C. de M. P. e. S.; Vicente, V. A.; Queiroz-Telles, F.; Rodrigues, A. M.; de Hoog, G. S.; Denning, D. W.; Colombo, A. L. The Global Burden of Chromoblastomycosis. *PLoS Negl Trop Dis* **2021**, *15* (8). <https://doi.org/10.1371/journal.pntd.0009611>.
4. Enbiale, W.; Bekele, A.; Manaye, N.; Seife, F.; Kebede, Z.; Gebremeskel, F.; van Griensven, J. Subcutaneous Mycoses: Endemic but Neglected among the Neglected Tropical Diseases in Ethiopia. *PLoS Negl Trop Dis* **2023**, *17* (9). <https://doi.org/10.1371/journal.pntd.0011363>.
5. Who. Ending the neglect to attain the Sustainable Development Goals: a strategic framework for integrated control and management of skin-related neglected tropical diseases. Geneva: World Health Organization, **2022**.
6. Agarwal, R.; Singh, G.; Ghosh, A.; Verma, K. K.; Pandey, M.; Xess, I. Chromoblastomycosis in India: Review of 169 Cases. *PLoS Neglected Tropical Diseases*. Public Library of Science August 1, **2017**. <https://doi.org/10.1371/journal.pntd.0005534>.
7. Guevara, A.; Siqueira, N. P.; Nery, A. F.; Cavalcante, L. R. D. S.; Hagen, F.; Hahn, R. C. Chromoblastomycosis in Latin America and the Caribbean: Epidemiology over the Past 50 Years. *Medical Mycology*. Oxford University Press December 1, **2021**. <https://doi.org/10.1093/mmy/myab062>.
8. He, Y.; Zheng, H. L.; Mei, H.; Lv, G. X.; Liu, W. Da; Li, X. F. Phaeohyphomycosis in China. *Frontiers in Cellular and Infection Microbiology*. Frontiers Media S.A. June 13, **2022**. <https://doi.org/10.3389/fcimb.2022.895329>.
9. Thomas, E.; Bertolotti, A.; Barreau, A.; Klisnick, J.; Tournebize, P.; Borgherini, G.; Zemali, N.; Jaubert, J.; Jouvion, G.; Bretagne, S.; Picot, S. From Phaeohyphomycosis to Disseminated Chromoblastomycosis: A Retrospective Study of Infections Caused by Dematiaceous Fungi. *Med Mal Infect* **2018**, *48* (4), 278–285. <https://doi.org/10.1016/j.medmal.2017.09.011>.
10. Liu, S.; Zhi, H.; Shen, H.; Lv, W.; Sang, B.; Li, Q.; Zhong, Y.; Liu, Z.; Xia, X. Chromoblastomycosis: A Case Series from Eastern China. *PLoS Negl Trop Dis* **2022**, *16* (9). <https://doi.org/10.1371/JOURNAL.PNTD.0010800>.
11. Passero, L. F. D.; Cavallone, I. N.; Belda, W. Reviewing the Etiologic Agents, Microbe-Host Relationship, Immune Response, Diagnosis, and Treatment in Chromoblastomycosis. *Journal of Immunology Research*. Hindawi Limited **2021**. <https://doi.org/10.1155/2021/9742832>.
12. de Brito, A. C.; Bittencourt, M. de J. S. Chromoblastomycosis: An Etiological, Epidemiological, Clinical, Diagnostic, and Treatment Update. *An Bras Dermatol* **2018**, *93* (4), 495–506. <https://doi.org/10.1590/abd1806-4841.20187321>.
13. Revankar, S. G.; Sutton, D. A. Melanized Fungi in Human Disease. *Clinical Microbiology Reviews*. October **2010**, pp 884–928. <https://doi.org/10.1128/CMR.00019-10>.
14. Carolina Rojas, O.; León-Cachón, R. B. R.; Pérez-Maya, A. A.; Aguirre-Garza, M.; Moreno-Treviño, M. G.; González, G. M. Phenotypic and Molecular Identification of *Fonsecaea pedrosoi* Strains Isolated from

- Chromoblastomycosis Patients in Mexico and Venezuela. *Mycoses* **2015**, *58* (5), 267–272. <https://doi.org/10.1111/myc.12308>.
15. Bienvenu, A. L.; Picot, S. Mycetoma and Chromoblastomycosis: Perspective for Diagnosis Improvement Using Biomarkers. *Molecules*. MDPI AG June 1, **2020**. <https://doi.org/10.3390/molecules25112594>.
 16. De Hoog, G.S.; Guarro, J.; Gené, J.; Figueras, M.J. Atlas of clinical fungi, 2^{ed.}; Utrecht, The Netherlands and Universitat Rovira i Virgili Reus, Spain, **2000**, pp. 560- 680.
 17. Monteiro, RC. Epidemiologia Molecular e Caracterização fenotípica de Fungos Demáceos Agentes de Cromoblastomicose no Estado do Pará, com Relato de Fungos Neurotrópicos na Região Amazônica. Dissertação de Mestrado, Universidade Federal do Pará, Belém, Pará, Brasil, **2017**.
 18. Sun, J.; Najafzadeh, M. J.; Gerrits van den Ende, A. H. G.; Vicente, V. A.; Feng, P.; Xi, L.; de Hoog, G. S. Molecular Characterization of Pathogenic Members of the Genus *Fonsecaea* Using Multilocus Analysis. *PLoS One* **2012**, *7* (8). <https://doi.org/10.1371/journal.pone.0041512>.
 19. Katoh, K.; Rozewicki, J.; Yamada, K. D. MAFFT Online Service: Multiple Sequence Alignment, Interactive Sequence Choice and Visualization. *Brief Bioinform* **2018**, *20* (4), 1160–1166. <https://doi.org/10.1093/bib/bbx108>.
 20. Roberto, T. N.; Rodrigues, A. M.; Hahn, R. C.; De Camargo, Z. P. Identifying *Paracoccidioides* Phylogenetic Species by PCR-RFLP of the Alpha-Tubulin Gene. *Med Mycol* **2016**, *54* (3), 240–247. <https://doi.org/10.1093/mmy/myv083>.
 21. Vincze, T.; Posfai, J.; Roberts, R. J. NEBcutter: A Program to Cleave DNA with Restriction Enzymes. *Nucleic Acids Res* **2003**, *31* (13), 3688–3691. <https://doi.org/10.1093/nar/gkg526>.
 22. Borman, A. M.; Fraser, M.; Patterson, Z.; Linton, C. J.; Palmer, M.; Johnson, E. M. Fungal Infections of Implantation: More Than Five Years of Cases of Subcutaneous Fungal Infections Seen at the UK Mycology Reference Laboratory. *Journal of Fungi* **2022**, *8* (4). <https://doi.org/10.3390/jof8040343>.
 23. Sendrasoa, F. A.; Razanakoto, N. H.; Rakotoarisaona, M. F.; Andrianarison, M.; Raharolahy, O.; Rasamoelina, T.; Ranaivo, I. M.; Sata, M.; Ratovonjanahary, V.; Maubon, D.; Rakoto Andrianarivelo, M.; Cornet, M.; Ramarozatovo, L. S.; Rapelanoro Rabenja, F. Clinical Aspects of Previously Treated Chromoblastomycosis: A Case Series from Madagascar. *International Journal of Infectious Diseases* **2020**, *101*, 228–232. <https://doi.org/10.1016/j.ijid.2020.09.1479>.
 24. Ariani, T.; Rizal, Y.; Ledika Veroci, R. *Clinical and Mycological Spectrum of Chromoblastomycosis*; 2023; Vol. 33.
 25. Revankar, S. G. Phaeohyphomycosis in Transplant Patients. *Journal of Fungi*. MDPI AG March 1, 2016. <https://doi.org/10.3390/jof2010002>.
 26. Gu, J.; Xu, J.; Su, Q.; Chen, Y. Exophiala Dermatitis and Exacerbation of Chronic Obstructive Pulmonary Disease. *QJM* **2019**, *112* (11), 869–871. <https://doi.org/10.1093/qjmed/hcz190>.
 27. An, L.; Jia, G.; Tan, J.; Yang, L.; Wang, Y.; Li, L. Analysis of the Synergistic Antifungal Activity of Everolimus and Antifungal Drugs against Dematiaceous Fungi. *Front Cell Infect Microbiol* **2023**, *13*. <https://doi.org/10.3389/fcimb.2023.1131416>.
 28. Coelho, R. A.; Brito-Santos, F.; Figueiredo-Carvalho, M. H. G.; Silva, J. V. dos S.; Gutierrez-Galhardo, M. C.; do Valle, A. C. F.; Zancopé-Oliveira, R. M.; Trilles, L.; Meyer, W.; Freitas, D. F. S.; Almeida-Paes, R. Molecular Identification and Antifungal Susceptibility Profiles of Clinical Strains of *Fonsecaea* Spp. Isolated from Patients with Chromoblastomycosis in Rio de Janeiro, Brazil. *PLoS Negl Trop Dis* **2018**, *12* (7). <https://doi.org/10.1371/journal.pntd.0006675>.
 29. Nascimento, E.; Barião, P. H. G.; Kress, M. R. von Z.; Vilar, F. C.; Santana, R. de C.; Gaspar, G. G.; Martinez, R. Cryptococcosis by *Cryptococcus Neoformans*/*Cryptococcus Gattii* Species Complexes in Non-Hiv-Infected Patients in Southeastern Brazil. *Rev Soc Bras Med Trop* **2021**, *54*. <https://doi.org/10.1590/0037-8682-0169-2021>.
 30. da Fonseca, L. A. V.; Araújo, M. A. S.; Silva, D. M. W.; de Albuquerque Maranhão, F. C. ITS-RFLP Optimization for Dermatophyte Identification from Clinical Sources in Alagoas (Brazil) versus Phenotypic Methods. *J Infect Dev Ctries* **2022**, *16* (11), 1773–1777. <https://doi.org/10.3855/jidc.17077>.
 31. Fontecha, G.; Montes, K.; Ortiz, B.; Galindo, C.; Braham, S. Identification of Cryptic Species of Four *Candida* Complexes in a Culture Collection. *Journal of Fungi* **2019**, *5* (4). <https://doi.org/10.3390/jof5040117>.
 32. Maubon, D.; Garnaud, C.; Ramarozatovo, L. S.; Fahafahantsoa, R. R.; Cornet, M.; Rasamoelina, T. Molecular Diagnosis of Two Major Implantation Mycoses: Chromoblastomycosis and Sporotrichosis. *Journal of Fungi*. MDPI April 1, **2022**. <https://doi.org/10.3390/jof8040382>.

33. Fajarningsih, N. D. Internal Transcribed Spacer (ITS) as Dna Barcoding to Identify Fungal Species: A Review. *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology* **2016**, *11* (2), 37. <https://doi.org/10.15578/squalen.v11i2.213>.
34. Hoang, M. T. V.; Irinyi, L.; Chen, S. C. A.; Sorrell, T. C.; Meyer, W. Dual DNA Barcoding for the Molecular Identification of the Agents of Invasive Fungal Infections. *Front Microbiol* **2019**, *10* (JULY). <https://doi.org/10.3389/fmicb.2019.01647>.
35. Xu, J. Fungal DNA Barcoding1. *Genome*. Canadian Science Publishing **2016**, pp 913–932. <https://doi.org/10.1139/gen-2016-0046>.
36. Réblová, M.; Untereiner, W. A.; Réblová, K. Novel Evolutionary Lineages Revealed in the Chaetothyriales (Fungi) Based on Multigene Phylogenetic Analyses and Comparison of ITS Secondary Structure. *PLoS One* **2013**, *8* (5). <https://doi.org/10.1371/journal.pone.0063547>.
37. Teixeira, M. M.; Moreno, L. F.; Stielow, B. J.; Muszewska, A.; Hainaut, M.; Gonzaga, L.; Abouelleil, A.; Patané, J. S. L.; Priest, M.; Souza, R.; Young, S.; Ferreira, K. S.; Zeng, Q.; da Cunha, M. M. L.; Gladki, A.; Barker, B.; Vicente, V. A.; de Souza, E. M.; Almeida, S.; Henrissat, B.; Vasconcelos, A. T. R.; Deng, S.; Voglmayr, H.; Moussa, T. A. A.; Gorbushina, A.; Felipe, M. S. S.; Cuomo, C. A.; de Hoog, G. S. Exploring the Genomic Diversity of Black Yeasts and Relatives (Chaetothyriales, Ascomycota). *Stud Mycol* **2017**, *86*, 1–28. <https://doi.org/10.1016/j.simyco.2017.01.001>.
38. Najafzadeh, M. J.; Gueidan, C.; Badali, H.; Van Den Ende, A. H. G.; Xi, L.; Hoog, G. S. Genetic Diversity and Species Delimitation in the Opportunistic Genus *Fonsecaea*. *Med Mycol* **2009**, *47* (1), 17–25. <https://doi.org/10.1080/13693780802527178>.
39. Usuda, D.; Higashikawa, T.; Hotchi, Y.; Usami, K.; Shimosawa, S.; Tokunaga, S.; Osugi, I.; Katou, R.; Ito, S.; Yoshizawa, T.; Asako, S.; Mishima, K.; Kondo, A.; Mizuno, K.; Takami, H.; Komatsu, T.; Oba, J.; Nomura, T.; Sugita, M. *Exophiala* Dermatitis. *World J Clin Cases* **2021**, *9* (27), 7963–7972. <https://doi.org/10.12998/wjcc.v9.i27.7963>.
40. Castillo Bejarano, J. I.; de los Santos, A. M.; Saldaña, D. C.; Zambrano Lucio, M.; Pérez Cavazos, S.; Espinosa Villaseñor, F.; de la O Cavazos, M. E.; Vaquera Aparicio, D. N. Pediatric Phaeohyphomycosis: A 44-Year Systematic Review of Reported Cases. *J Pediatric Infect Dis Soc* **2022**. <https://doi.org/10.1093/jpids/piac106>.
41. Le Govic, Y.; Demey, B.; Cassereau, J.; Bahn, Y. S.; Papon, N. Pathogens Infecting the Central Nervous System. *PLoS Pathog* **2022**, *18* (2). <https://doi.org/10.1371/journal.ppat.1010234>.
42. Ganesan, V.; Hallur, V.; Velvizhi, S.; Rajendran, T. Cerebral Phaeohyphomycosis Due to *Cladophialophora Bantiana*: Case Report and Systematic Review of Cases. *Infection* **2023**. <https://doi.org/10.1007/s15010-023-02126-4>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.