

Article

Not peer-reviewed version

Effect of Chlorhexidine Digluconate on Oral Bacteria Adhesion to Surfaces of Orthodontic Appliance Alloys

Doria Gergeta , Matea Badnjević , [Ljerka Karleuša](#) , [Željka Maglica](#) , [Stjepan Spalj](#) ^{*} , [Ivana Gobin](#)

Posted Date: 19 January 2024

doi: 10.20944/preprints202401.1495.v1

Keywords: Biofilm; Dental plaque; Orthodontic appliances; Antiseptics



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Effect of Chlorhexidine Digluconate on Oral Bacteria Adhesion to Surfaces of Orthodontic Appliance Alloys

Gergeta D ^{1,†}, Badnjevic M ^{2,†}, Karleusa Lj ³, Maglica Z ⁴ and Spalj S ^{2,5,*} Gobin I ⁶

¹ Community Health Center Porec, Croatia; doriagergeta@outlook.com

² University of Rijeka, Faculty of Dental Medicine, Department of Orthodontics; matea.badnjevic@fdmri.uniri.hr, stjegan.spalj@uniri.hr

³ Department of Physiology and Immunology, Faculty of Medicine, University of Rijeka, Brace Banchetta 20, 51000 Rijeka, Croatia; ljerka.karleusa@medri.uniri.hr

⁴ Department of Biotechnology, University of Rijeka, 51000 Rijeka, Croatia; zeljka.maglica@biotech.uniri.hr

⁵ Department of Dental Medicine, J. J. Strossmayer University of Osijek, Faculty of Dental Medicine and Health, 31000 Osijek, Croatia

⁶ Department of Microbiology and Parasitology, Faculty of Medicine, University of Rijeka, 51000 Rijeka, Croatia; ivana.gobin@uniri.hr

* Correspondence: Stjepan Spalj, University of Rijeka, Faculty of Dental Medicine, Department of Orthodontics; stjegan.spalj@uniri.hr

† First two authors equally contributed sharing first authorship.

Abstract: This study aimed to analyse the effect of chlorhexidine digluconate (CHX DG) mouthwash on the adhesion of oral bacteria to orthodontic appliances. The interactions of four bacteria (*S. mutans*, *A. actinomycetemcomitans*, *S. oralis*, and *V. parvula*) with two alloys (stainless steel [SS] and nickel-titanium [NiTi]) and three CHX DG solutions (commercial products Curasept and Perio Plus, and pure CHX DG, all with 0.12% active substance) were tested. The adhesive effect on the orthodontic wires was evaluated after 24 h for *S. oralis* and after 72 h for the other bacteria. The minimum bactericidal concentration of the solution for each bacterial strain was determined using the dilution method to test the antibacterial action. Salivary-pretreated orthodontic arch wires were exposed to minimal bactericidal concentrations of solution and bacteria. Commercial antiseptic products, especially Perio Plus, showed better inhibition of bacterial adhesion to both alloys than pure CHX DG solution ($p < 0.05$). *A. actinomycetemcomitans* was most inhibited in the adhesion of all bacteria by the CHX DG products. Greater inhibition of streptococci adherence was observed on SS, while *A. actinomycetemcomitans* on NiTi. Inhibition of *V. parvula* was product dependent. In conclusion bacterial adhesion depends on the alloys, antiseptic products, and the bacteria. CHX DG is effective at inhibiting bacterial adhesion to orthodontic elements and is recommended for the chemical control of biofilms during orthodontic treatment with fixed appliances.

Keywords: biofilm; dental plaque; orthodontic appliances; antiseptics

1. Introduction

A biofilm is a sessile community of microbial cells immersed in a matrix composed of extracellular polymeric substances (EPS) attached to the surface. EPS consists of exopolysaccharides, lipids, proteins, and nucleic acids (eDNA and eRNA) and offers a functional environment and structural stability. Tolerance to immune cells and antimicrobials is enhanced by EPS [1]. Dental plaque is a complex biofilm whose formation begins with the formation of the salivary pellicle and changes progressively; therefore, we can differentiate early and mature plaque, as well as subgingival and supragingival plaque [2]. Supragingival biofilm bacteria like *S. mutans*, *Actinomyces spp.*, and *Lactobacillus* have the greatest influence on enamel and dentin demineralisation and ultimately caries

formation [3]. Some subgingival biofilm bacteria include *Actinomyces spp.*, *Tannerella forsythia*, and *Fusobacterium nucleatum* which are associated with gingivitis and periodontitis [4].

Malocclusion therapy in adolescents and adults commonly includes fixed orthodontic appliances containing wires, brackets, and ligatures [5]. Materials in these appliances are alloys such as stainless steel (SS), nickel-titanium (NiTi), titanium-molybdenum, and cobalt-chromium [6]. The difficulty in maintaining good oral hygiene is related to the additional retention places of the fixed appliance for plaque accumulation. Some studies suggest that plaque accumulation is especially increased in metal brackets with elastic ligatures in the maxilla, the maxillary canines, and lateral incisors [7], unlike in non-orthodontic patients, where the mandibula and maxillary molars are the primary sites of accumulation [8]. Areas around the brackets (mesial, distal, and gingival) accumulate more plaque than occlusal areas because they interfere most with archwires and ligatures; thus, natural self-cleaning is not possible [9]. Difficult and inadequate oral hygiene by brushing and a lower ability to use dental floss lead to plaque accumulation and oral flora changes [10]. It can cause gingivitis, enamel white spot lesions (WSL), and caries [11]. The progression of gingivitis can lead to periodontitis and ultimately cause irreversible damage, such as the loss of supporting tissue. Studies claim that some orthodontic patients do not complete treatment because of poor oral hygiene. Orthodontic patients need to have proper oral hygiene because it is very challenging, and patient motivation is crucial [12]. Therefore, patients should be aware of the consequences of improper hygiene. The clinician may use tools other than informed consent to motivate the patient, such as animations, leaflets, and other verbal information [13]. Appliance placement alters the oral microbiota, and we can observe high concentrations of acidogenic bacteria such as *Streptococcus* and *Lactobacillus* which result in WSL progression and caries development in patients with fixed appliances within a month after placement [14], unlike caries development in mouth without braces that can take up to 6 months [15]. Etiological factors associated with periodontitis include an increase in gram-positive supragingival bacteria, such as *Streptococcus* and *Actinomyces*. Regarding gingivitis, etiologies include an increase in gram-negative *Fusobacterium* and *Bacterioides* [16]. In addition to mechanical plaque control, such as brushing and flossing, chemical plaque control can be performed. The gold standard is chlorhexidine digluconate (CHX DG), which has side effects such as staining and altered taste after long-term use [17,18]. Short-term usage is recommended for people with difficulty in maintaining oral hygiene, such as those with fixed orthodontic appliances, mental and physical disabilities, intermaxillary fixations, and postsurgical or systematic diseases with oral manifestations [19]. Orthodontic wires provide a suitable environment for the growth of oral microorganisms because they are constantly present throughout orthodontic treatment and can play a significant role in enamel demineralization [20]. The contact areas between the wires and brackets make cleaning difficult by preventing proper access to the tooth surface. Some researchers claim that SS wires increase plaque retention capacity because of their high surface tension and energy [21].

This study aimed to analyze the effect of CHX DG on the adhesion of oral bacteria to SS and NiTi alloys, which are often used in orthodontic treatment. The hypothesis was that CHX DG with the addition of alcohol would reduce adhesion to steel alloys, especially *S. mutans* bacteria.

2. Materials and Methods

2.1. Cultivation of bacteria

The bacteria used were *Streptococcus mutans* ATCC 25175, *Streptococcus oralis* ATCC 6249, *Aggregatibacter actinomycetemcomitans* ATCC 29522, and *Veillonella parvula* ATCC 10790 (Microbiology, St Cloud, MN, USA). Streptococci were cultivated on Mutans-Sanguis agar (HIMEDIA, Mumbai, India) and incubated at 37° for 24 h with a candle to simulate anaerobic conditions. *A. actinomycetemcomitans* and *V. parvula* were cultivated on blood agar (Biolife, Milan, Italy) supplemented with 5% sheep blood (Biognost, Zagreb, Croatia) and incubated at 37° for 3-5 days in anaerobic conditions using AnaeroGen bags (GENbox anaerobe, bioMerieux SA, Marcy-l'Étoile, France). Brain Heart Infusion Broth (BHI-Brain Heart Infusion) (Becton, Dickinson and Company, Sparks, USA) was used to resuspend bacterial cultures supplemented with 1.0 mg/mL

Vitamin K and 5.0 mg/mL haemin (Sigma-Aldrich, Burlington, USA). Preparation of suspensions. The optical density at 600 nm (OD₆₀₀) was determined using a spectrophotometre and was set to OD₆₀₀=1, which was interpreted to mean that there were 1×10^9 colonies formed per millilitre (colony forming unit, CFU) in the suspension. Previously prepared suspensions were diluted and a suspension with a concentration of 1×10^7 CFU/mL was used.

2.2. Adhesion of bacteria to orthodontic archwires

Bacteria were transferred to the wells with 200 μ L of bacterial suspension and the microtitre plate was placed in the incubator for 4 h with stirring. Sterile orthodontic archwires of 0.019x0.025 " SS (Stainless steel, American Orthodontics, Sheboygan, USA) and 0.018x0.025 " NiTi (Neo Sentalloy, Dentsply Sirona, Charlotte, USA) were used. Before being placed in 500 μ L of sterile saline and exposed to sonication for 1 min (BathoSonic ultrasonic bath-Badelin, Berlin, Germany) at an intensity of 40 kHz, the wires were rinsed twice with saline. After homogenization, tenfold dilutions were made on the microtitre plates, and after planting on the substrate, the bacteria were incubated under anaerobic conditions or with a candle. After 24 h, *S. oralis* was counted, and after 72 h, *A. actinomycetemcomitans*, *S. mutans* and *V. parvula*.

2.3. Antibacterial action of antiseptics

The mouthwashes used were Curasept ADS 212 (Curasept, Italy), Perio Plus (Curasept, Switzerland), and a solution of pure 0.12% CHX DG in distilled water (Sigma-Aldrich, USA) all containing 0.12% CHX DG. A 200 μ L-mouthwash solution was applied to a microtitre plate that was diluted twice. The microtitre plate was incubated for 24 h at 37°. *V. parvula* was incubated on a shaker for 48 h. A suspension of 100 μ L of bacteria was applied to the wells. Sediment or turbidity is a sign of bacterial growth on microtitre plates. Turbidity in the wells was compared to that in the first column which was a control containing only bacteria. Because of the turbidity of the sample, it was not possible to determine the lowest concentration of the antimicrobial agent, the minimum inhibitory concentration (MIC). Therefore, the lowest concentration that inhibited the growth of bacteria on the nutrient medium, the minimum bactericidal concentration (MBC), was determined. Samples from wells without turbidity were plated on Mutans-Sanguis and blood agar and incubated at 37° for 24–72 h. The minimum bactericidal concentration was indicated as the sample without bacterial growth from the sample with the lowest concentration of the active component.

2.4. Anti-adhesive effects of antiseptics

Orthodontic arches were treated with saliva and exposed to 200 μ L half-subinhibitory concentration of oral antiseptic and 200 μ L bacterial suspension. The mixture was then stirred under anaerobic conditions for 4 h. After the incubation, the wires were washed twice with saline and centrifuged for 1 min. Bacteria were cultivated and incubated under anaerobic conditions (*Velionella*) or by candles (*Streptococcus*) from dilutions on microtitre plates. After 24 h, an increase was observed in the substrate for *S. oralis*, and after 72 h for *A. actinomycetemcomitans*, *V. parvula*, and *S. mutans*.

2.5. Visualization of adherent bacteria on archwires by fluorescence microscopy

The LIVE/DEAD BacLight Bacterial Viability Kit L-7012 (Invitrogen, Waltham, MA, USA) was used to stain the adherent bacteria on the archwires according to the manufacturer's instructions. SYTO 9 stained all cells in a green fluorescent colour, whereas propidium iodide (PI) stained cells with a damaged red membrane. Epifluorescence microscopy was used to visualise the adhered bacteria with GFP/FITC (excitation:480 nm and emission:500 nm) and rhodamine (excitation:490 nm and emission:635 nm) filters. Simultaneous dual-channel imaging was used to visualise green and red fluorescence. The obtained images were saved in the TIFF format and further processed using ImageJ 1.47. A minimum of three images per term were analysed.

2.6. Statistical methods

A logarithmic value of 10 CFU/mL was used for the statistical analysis. Factorial analysis of variance (ANOVA) was used to analyse the effects of time, alloy, bacterial interaction, and bacterial type on adhesion. ANOVA with the Student-Newman-Keuls post-hoc test was used to compare the adhesion between the four bacteria and the four periods. The proportion of inhibition of adhesion by preparations with CHX DG compared to the control was calculated according to formula 1 (number of treatments/number of controls). ANOVA was used to analyse the interaction of the agents, alloys, and bacteria with adhesion inhibition. An independent samples t-test was used to compare the adhesion inhibition and adhesion between the alloys. ANOVA with Student-Newman-Keuls post-hoc test was used to compare the types of oral antiseptics and bacteria. The t-test was calculated using the formula $r = \sqrt{t^2 / (t^2 + df)}$, and the effect size was quantified via η^2 for ANOVA. Cohen's r criteria were used in the interpretation: 0.1-0.3=small effect size, 0.3-0.5=moderate, 0.5-0.7=large, >0.7 very large. The squared values of r were used for the interpretation of η^2 . All analyses were performed using the commercial software SPSS 22 (IBM SPSS, Version 22.0, IBM, Armonk, NY, USA).

3. Results

3.1. Minimum bactericidal concentrations (MBC)

The MBCs are listed in Table 1. The CHX DG solution is presented in $\mu\text{g/mL}$ because pure 0.12% CHX DG was prepared in the laboratory, while Perio Plus and Curasept as factory solutions are shown in percentages.

Table 1. Minimum bactericidal concentrations for individual bacteria.

Bacteria	CHX DG solution ($\mu\text{g/mL}$)	Curasept (%)	Perio Plus (%)
<i>S. mutans</i>	12.5	1.56	$9.98 \times 10^{(-2)}$
<i>S. oralis</i>	25	3.13	$9.98 \times 10^{(-2)}$
<i>V. parvula</i>	12.5	0.78	0.195
<i>A. actinomycetemcomitans</i>	6.25	3.13	0.195

3.2. Adherence to alloys

Adherence to alloys was the highest in *V. parvula*, followed by *A. actinomycetemcomitans*, *S. oralis*, and *S. mutans* (Figure 1). The magnitude of the effect was greater for SS than for NiTi ($p < 0.001$; $\eta^2 = 0.930$ and 0.909 , respectively). Two-factor ANOVA pointed to the interaction of alloy and type of bacteria in adherence ($p < 0.001$; $\eta^2 = 0.184$). *S. mutans* showed greater adhesion to SS than to NiTi (3.5 ± 0.1 vs. 3.2 ± 0.1 ; $p < 0.001$; $r = 0.678$) as well as *V. parvula* (5.4 ± 0.3 vs. 5.0 ± 0.3 ; $p < 0.001$; $r = 0.519$), although the effect was greater on *S. mutans*. *A. actinomycetemcomitans* (NiTi 4.7 ± 0.1 and SS 4.6 ± 0.1) and *S. oralis* (NiTi 4.3 ± 0.2 and SS 4.2 ± 0.3) showed equal adhesion to both alloys.




Figure 1. Comparison of bacterial adhesion to alloys. Columns show mean values with 95% confidence intervals. Asterisks indicate significant differences between alloys for the same bacterium, and horizontal lines connect bacteria with a significant difference within the same alloy.

3.3. Inhibition of adhesion

Three-factor ANOVA indicated a significant interaction of the alloy, bacteria, and agent on adhesion reduction with a smaller effect size ($p=0.001$; $\eta^2=0.084$). Two-factor ANOVA showed that the interaction of bacteria and alloy was significant and with a greater effect with Perio Plus ($p<0.001$; $\eta^2=0.452$) as well as with pure CHX DG solution ($p<0.001$; $\eta^2=0.194$). Curasept showed no significant interactions. Compared to other bacteria, *A. actinomycetemcomitans* showed the greatest inhibition of adhesion to CHX DG solutions (Figure 2). The inhibition of adhesion on *S. mutans* was the lowest and the highest on *A. actinomycetemcomitans* on both alloys. The effect of CHX DG solution on inhibition of adhesion was greater with NiTi than with SS ($\eta^2=0.829$ and 0.626) and is significantly different for bacteria. On SS, there was less inhibition of adhesion for *V. parvula* than for *S. oralis*, and after the application of Curasept, there was less inhibition of adhesion of *V. parvula* on NiTi (84% vs. 99%–100%) ($p<0.001$; $\eta^2=0.757$) and SS (78 vs. 98%–100%; $p<0.001$; $\eta^2=0.713$) than other bacteria. Perio Plus had a weaker effect on the inhibition of *V. parvula* on NiTi (99.8% 100%) ($p<0.001$; $\eta^2=0.654$) with a greater effect size on SS (99.3% vs. 100%; $p<0.001$; $\eta^2=0.779$) than it acted on other bacteria with both alloys.




Figure 2. Comparison of control adherence and antiseptic treatment. Average column values are with 95% confidence intervals. Significant differences between alloys for the same bacteria and antiseptic are marked with an asterisk. Horizontal lines connect antiseptics that show a significant difference between the same alloy and bacteria.

Viability depended on the type of bacteria, exposure, and alloy combination ($p=0.047$; $\eta^2=0.217$; Figures 3 and 4). It was smaller with SS wires than with NiTi wires and after CHX DG therapy. Viability was the highest in *V. parvula* and lowest in *S. mutans*. CHX DG induced lower viability on both alloys in *V. parvula* and *S. mutans*, whereas in *A. actinomycetemcomitans* only on NiTi, with a large effect size ($p\leq 0.010$; $r=0.841-0.958$). There were significant differences in the viability of the control on SS in the order *V. parvula* > *S. oralis* > *A. actinomycetemcomitans* = *S. mutans* ($p<0.001$). In the NiTi control, there were significant differences in viability in the order of *V. parvula* > *A. actinomycetemcomitans* > *S. oralis* > *S. mutans* ($p<0.001$). In both alloys, significant differences in the viability of the CHX DG were observed in the following order: *V. parvula* > *A. actinomycetemcomitans* > *S. mutans* = *S. oralis* ($p<0.001$).




Figure 3. Representative images of bacteria adhering within 4 h to NiTi and SS wires with and without CHX treatment. Biofilms were stained with the LIVE/DEAD Sustainability Kit (FilmTracer™ LIVE/DEAD® Biofilm Viability Kit, Invitrogen). Green cells represent viable, live bacteria, while red cells represent dead bacteria.




Figure 4. Comparison of viability by fluorescence on confocal microscope. Total mean fluorescence is measured for propidium iodide and SYTO® 9 stained adherent bacteria before and after treatment with CHX DG. The experiment is repeated 2x and a minimum of three images are analysed and the mean fluorescence \pm SD is shown.

3.4. Differences between alloys

Two-factor ANOVA showed that the interaction of bacteria and agent was significant and with a greater effect for NiTi ($p < 0.001$; $\eta^2 = 0.609$) than for SS ($p < 0.001$; $\eta^2 = 0.380$). The inhibition of *V. parvula* depended on the solution. On SS, there was greater inhibition of adherence in streptococci, whereas,

on NiTi, there was greater inhibition of adherence in *A. actinomycetemcomitans*. After CHX DG therapy, the inhibition of adherence to SS was greater than that to NiTi in *S. oralis* (96% vs. 89%; $p < 0.001$; $r = 0.303$) and *V. parvula* (81% vs. 69%; $p < 0.001$; $r = 0.532$). The inhibition of adherence to SS after CHX DG in *S. mutans* was higher, but not significant, owing to the large scattering of the data. The inhibition of *A. actinomycetemcomitans* was equal for both metals after the application of CHX DG. Excellent inhibition of adherence was observed for both metals after Curasept therapy in *A. actinomycetemcomitans* (99% vs. 98%; $p = 0.001$; $r = 0.486$), although it was higher for NiTi than for SS. Greater inhibition was observed for *V. parvula* (84% vs. 78%), although the difference was not statistically significant because of the scattering of the data. After therapy with Curasept for streptococci, no differences were observed between the alloys. Perio Plus therapy was the least dependent on the type of alloy used. *V. parvula* showed greater inhibition of adherence to NiTi than to SS alloy after Perio Plus therapy (99.8% vs. 99.3%; $p < 0.001$; $r = 0.811$), and there was no difference between the alloys for other bacteria (100% inhibition).

3.5. Differences between solutions

V. parvula ($p < 0.001$; $\eta^2 = 0.208$) and *S. oralis* ($p < 0.001$; $\eta^2 = 0.675$) showed a significant interaction between the agents and alloys, with a larger effect in *S. oralis*, as shown by two-factor ANOVA. *S. mutans* and *A. actinomycetemcomitans* did not show significant interactions. CHX DG produced the lowest inhibition of adherence and Perio Plus produced the highest inhibition among all bacteria. CHX DG caused less adherence inhibition than Perio Plus and Curasept (100%; $p < 0.001$) on both alloys (66%–89%) in *S. oralis* with a larger effect size on NiTi than SS ($\eta^2 = 0.939$ and 0.708), no difference the last two solutions. A smaller inhibition of adherence on both alloys (44–59%) was shown by CHX DG than Perio Plus and Curasept (100%) in *S. mutans* with a larger effect size on NiTi ($p < 0.001$; $\eta^2 = 0.871$) than SS ($p = 0.003$; $\eta^2 = 0.871$). $\eta^2 = 0.532$, with no difference between the two last solutions. Adherence inhibition was high with all three agents (97–100%) in *A. actinomycetemcomitans*. Perio Plus had the greatest inhibition, and CHX DG had the smallest effect size on SS ($p = 0.003$; $\eta^2 = 0.311$) than NiTi ($p < 0.001$; $\eta^2 = 0.618$). After Perio Plus therapy, *V. parvula* had the highest inhibition on NiTi alloy, and the lowest after using CHX DG (100% and 69%; $p < 0.001$; $\eta^2 = 0.671$). On SS, Perio Plus showed the highest inhibition of adherence but the lowest inhibition after Curasept therapy (99% and 78%; $p < 0.001$; $\eta^2 = 0.604$).

4. Discussion

This study has shown that the adhesion of oral bacteria to orthodontic appliances depends on complex interactions between the type of alloy, antiseptic product used, and bacteria. The MIC could not be determined in the present study because of the presence of sediment in the wells. Other studies report that MIC in different strains of *S. mutans* treated with CHX DG varied from 0.03–0.2 $\mu\text{g/mL}$, and MBC 0.24–0.9 $\mu\text{g/mL}$. Sub-inhibitory concentrations of CHX DG have also been reported to influence the growth of planktonic *S. mutans* cells and biofilm [22].

Bacterial adhesion is the initial step in biofilm formation. Orthodontic appliances create new adhesion surfaces, and multiple bacterial species colonize the oral cavity and adhere to these surfaces to prevent swallowing. According to the present study, adhesion to alloys was the highest in *V. parvula* and lowest in *S. mutans* with *A. actinomycetemcomitans* and *S. oralis*. *S. mutans* and *V. parvula* showed greater adhesion to the SS wires, whereas *A. actinomycetemcomitans* and *S. oralis* had similar effects on both wires.

Bacterial hydrophobicity is one of the factors influencing the adhesion of oral bacteria to orthodontic alloys during the first 4 h while alloy surface characteristics have little influence. In the same study, *V. parvula* was found to have extremely hydrophobic properties, *S. oralis* and *S. mutans* were moderately hydrophobic, and *A. actinomycetemcomitans* exhibited hydrophilic properties. Both NiTi and SS exhibit hydrophilic characteristics; however, NiTi is more hydrophilic. In addition, NiTi has a higher roughness, polarity, and surface free energy than SS which makes NiTi more prone to bacterial adhesion. However, the difference between the surface free energies of the alloy and bacteria is inversely proportional to bacterial adhesion [23]. In addition, some biological processes regulate

bacterial adhesion, such as the production of extracellular polysaccharides, cell viability, charge and wall stiffness, bacterial surface protuberances, metabolic activity, hydrophobicity, and adhesin-mediated receptor-ligand binding [24].

The process of tooth preparation for bracket bonding by enamel etching alters the surface properties, and this initial step influences adhesion and biofilm formation [25]. Rough areas increase the space for biofilm accumulation and adhesion, whereas higher wettability makes it easier for biofilms to adhere to dental materials [26]. The correlation between the bonding procedure and the accumulation of *S. mutans* and *P. gingivalis* has been confirmed [27].

The market offers aesthetic archwires, in addition to traditional archwires made of iron, titanium, or cobalt alloys [28]. The wires are coated with epoxy resin and polytetrafluoroethylene [29]. In vitro research with *S. mutans* showed that there was no major difference in biofilm formation or bacterial accumulation between the coated and uncoated archwires when colony counting [30]. When comparing Cu-NiTi archwires and NiTi wires, it appears that *S. mutans* adheres better to the Cu-NiTi wires [31].

The main method of oral hygiene is the mechanical removal of plaque by tooth brushing (manual or electric). In addition to manual techniques, the gold standard for chemical plaque control is bisbiguanide CHX mouthwash. It was developed in the UK in the forties as a general disinfectant. In the seventies, it was being used as a mouthwash because its anti-plaque effects were discovered [32]. In dentistry 0.12%–0.2% CHX can be used preoperatively, postoperatively (if brushing is enabled), or long-term (CHX chips). The recommended dosage for maximum effect per use is 18–20 mg [33].

The bacterial wall is negatively charged and the CHX wall is positively charged; therefore, they are attracted to each other. On the surface of the bacterial wall, phosphate-containing molecules strongly adsorb onto CHX. CHX penetrates the bacterial wall, damaging the bacterial cell and inhibiting enzymes related to the cytoplasmic membrane, and causes an overflow of low-molecular-weight cytoplasmic components, such as potassium ions [34]. This is the bacteriostatic stage which can be reversed if CHX is removed, but if it persists, the antimicrobial action continues to the bactericidal stage, where cytoplasmic coagulation and precipitation occur, and complexes such as adenosine triphosphate and nucleic acids are formed. Oral surfaces are negatively charged; therefore, CHX, a cationic molecule, adheres well to these surfaces and exhibits good interference with bacterial adhesion [35]. Good binding to the oral surface allows it to last longer in the mouth. CHX exhibits a prolonged action and can be released through a period of 8–12 h in the mouth [19,36]. CHX has antifungal, antimicrobial (anaerobes and aerobes), and antiviral (RNA, DNA, and lipophilic-enveloped viruses) activities [37].

This study showed that the anti-adhesive effect is mostly related to the type of mouthwash used. CHX DG showed the lowest inhibition of adhesion, whereas Perio Plus showed the highest inhibition. The Curasept acted similarly to Perio Plus. *A. actinomycetemcomitans* was the most inhibited, whereas *V. parvula* was the least. Some differences were observed but not with commercial products. There is unclear evidence if mouthwash with CHX DG concentrations below 0.1% has an antiplaque effect. Some authors claim that CHX DG 0.05% with herbal extract has a similar antiplaque effect as CHX DG at concentrations of 0.1%, while others claim that CHX DG at concentrations below 0.1% has an unclear antiplaque effect [38,39]. It is generally recommended to rinse twice daily with 20 mg of CHX DG mouthwash at concentrations of 0.12%–0.2% [40]. In addition to the DG concentrations of CHX, it is necessary to be aware of toothpaste components. The effect of CHX DG can be reduced with ingredients such as sodium lauryl sulfate, cocamidopropyl betaine (CAPB), calcium and anionic surfactants, and sodium dodecyl sulfate in toothpaste. To avoid a reduction in the effect, the recommendation is to wait 30 min after brushing before using the mouthwash [41]. Fluoride toothpaste should be switched with CAPB paste because it provides better enamel remineralization as a surfactant with a 30-minute delayed rinse [42]. In addition to mouthwash, CHX DG can be used as a gel or dentifrice, although mouthwash should be the first choice, especially in non-brushing models. Researchers have shown that CHX mouthwash reduces *S. mutans* better than CHX gels and varnishes, even at higher concentrations, but when used alone, it does not have a clinical impact on significant caries reduction [43]. This study found that the cell viability was

dependent on the bacteria, alloys, and exposure. The viability of the wires was greatest in *V. parvula* and smallest in *S. mutans*. CHX DG decreased the viability of *S. mutans* and *V. parvula* on SS and NiTi and decreased the viability of *A. actinomycetemcomitans* only on NiTi.

Perio Plus is the only alcohol solution used in this study. When comparing mouthwash 0.2% CHX DG with a non-alcohol base and essential oils containing alcohol, both showed similar results on gingival health; however, because of the side effects of alcohol-based mouthwash, such as reduced salivation and burning mouth, it is recommended to use the non-alcohol-based one [44]. In contrast, a 7-day study using CHX DG showed a decrease in the pH of the saliva which is associated with enamel demineralisation and caries. CHX DG showed a shift to an acidic environment; therefore, mouthwash application should be cautiously considered [45].

The advantage of this research was the controlled environment for experiments in a microbiological laboratory, so it was possible to analyse each bacterium and product separately. The disadvantage is that bacteria work together in the oral cavity and coexist in dynamic media, making it difficult to imitate the oral cavity under laboratory conditions. The pH of the mouth was not considered in this study. Various factors influence biofilm formation on orthodontic appliances in the oral cavity; however, this connection is not fully understood. Additional research is required to clarify these gaps. The difference between the results of clinical and laboratory tests is large, which shows the complexity of dental biofilms.

5. Conclusions

The inhibition of bacterial adhesion depends on the alloy, antiseptic products, and biofilm bacteria. CHX DG is effective at inhibiting bacterial adhesion to orthodontic elements and is recommended for the chemical control of biofilms during orthodontic treatment with fixed appliances.

Author Contributions: Conceptualization I.G., D.G. and S.S.; methodology S.S.; validation L.J.K., Z.M., I.G., S.S., formal analysis, S.S.; investigation D.G.; resources S.S.; data curation, S.S.; writing—original draft preparation M.B, D.G.; writing—review and editing L.J.K., Z.M., I.G., S.S., M.B. and D.G.; visualization S.S., D.G.; supervision S.S. and I.G.; project administration S.S.; funding acquisition S.S. All authors have read and agreed to the published version of the manuscript.

Funding: Croatian Science Foundation IP-2020-02- 4027 (Environmental factors and microbial interactions in the structure of the dental biofilm) and DOK-2021-02-6438 (Career development of young researchers – training of new PhDs). Preliminary data was presented as a poster at Italian Orthodontic Society International Spring Meeting in Genova 18-19.3.2022.

Institutional Review Board Statement: This study was approved by authors' institution ethics committee and was conducted in accordance with the ethical standards of the 1964 Declaration of Helsinki and its subsequent amendments.

Informed Consent Statement: Written informed consent was obtained from all adult participants and the parents/guardians of all participants aged < 18 years before participation in the study.

Data Availability Statement: The datasets used and/or analyzed during the current study are available on Digital Academic Repository of the University of Rijeka, Faculty of Dental Medicine from the corresponding author on reasonable request. (Špalj S. Okolišni čimbenici i mikrobiološke interakcije u strukturi dentalnog biofilma: istraživački podaci. [Internet]. Fakultet dentalne medicine; 2023, [cited 2024 January 04] Available from: <https://urn.nsk.hr/urn:nbn:hr:271:843444>.)

Acknowledgments: We thank Gabrijela Begic for help in laboratory procedures.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S. Biofilms: an emergent form of bacterial life. *Nat. Rev. Microbiol.* **2016**, *14*, 563-575. doi: 10.1038/nrmicro.2016.94.
2. Kwon T, Lamster IB, Levin L. Current concepts in the management of periodontitis. *Int. Dent. J.* **2021**, *71*, 462-476. doi: 10.1111/idj.12630.

3. De Freitas AOA, Marquezan M, Nojima MDCG, Alviano DS, Maia LC. The influence of orthodontic fixed appliances on the oral microbiota: A systematic review. *Dent. Press J. Orthod.* **2014**, *19*, 46–55. doi: 10.1590/2176-9451.19.2.046-055.oar.
4. Zijngje V, van Leeuwen MBM, Degener JE, Abbas F, Thurnheer T, Gmür R, et al. Oral biofilm architecture on natural teeth. *Plos One.* **2010**, *5*, e9321. doi: 10.1371/journal.pone.0009321.
5. Liu H, Sun J, Dong Y, Lu H, Zhou H, Hansen BF, et al. Periodontal health and relative quantity of subgingival porphyromonas gingivalis during orthodontic treatment. *Angle Orthod.* **2011**, *81*, 609–615. doi: 10.2319/082310-352.1.
6. Kapila S, Sachdeva R. Mechanical properties and clinical applications of orthodontic wires. *Am. J. Orthod. Dentofacial Orthop.* **1989**, *96*, 100-9. doi: 10.1016/0889-5406(89)90251-5.
7. Li M, Chieng J, Wong C, Benic G, Farella M. Factors affecting dental biofilm in patients wearing fixed orthodontic appliances. *Prog. Orthod.* **2017**, *18*, 4. doi: 10.1186/s40510-016-0158-5.
8. Furuichi Y, Lindhe J, Ramberg P, Volpe AR. Patterns of de novo plaque formation in the human dentition. *J. Clin. Periodontol.* **1992**, *19*, 423–433. doi: 10.1111/j.1600-051X.1992.tb00673.x.
9. Garcez AS, Suzuki SS, Ribeiro MS, Mada EY, Freitas AZ, Suzuki H. Biofilm retention by 3 methods of ligation on orthodontic brackets: a microbiologic and optical coherence tomography analysis. *Am. J. Orthod. Dentofacial Orthop.* **2011**, *140*, e193–8. doi: 10.1016/j.ajodo.2011.04.019.
10. Goes P, Dutra CS, Lisboa MR, Gondim DV, Leitão R, Brito GA, et al. Clinical efficacy of a 1% Matricaria chamomile L. mouthwash and 0.12% chlorhexidine for gingivitis control in patients undergoing orthodontic treatment with fixed appliances. *J. Oral Sci.* **2016**, *58*, 569–574. doi: 10.2334/josnusd.16-0280.
11. Kleter GA. Discoloration of dental carious lesions (a review) *Arch. Oral Biol.* **1998**, *43*, 629–632. doi: 10.1016/s0003-9969(98)00048-x.
12. Huang J, Yunxia Y, Jiuhui, Cuiying L. Effects of motivational methods on oral hygiene of orthodontic patients A systematic review and meta-analysis. *Medicine (Baltimore).* **2018**, *97*, e13182. doi: 10.1097/MD.00000000000013182.
13. Shqaidef AJ, Saleh MYN, Kussad J, Khambay BS. Do parents of adolescent patients undergoing fixed appliance treatment recall more information using written material or an animated video? A randomized controlled trial. *Saudi Dent. J.* **2023**, *35*, 95-102. doi: 10.1016/j.sdentj.2022.11.011.
14. Lundström F, Krasse B. Streptococcus mutans and lactobacilli frequency in orthodontic patients; the effect of chlorhexidine treatments. *Eur. J. Orthod.* **1987**, *9*, 109–116. doi: 10.1093/ejo/9.2.109.
15. Mattousch TJ, van der Veen MH, Zentner A. Caries lesions after orthodontic treatment followed by quantitative light-induced fluorescence: A 2-year follow-up. *Eur. J. Orthod.* **2007**, *29*, 294–298. doi: 10.1093/ejo/cjm008.
16. Sinclair M, Berry CW, Bennet CL, Israelson H. Changes in gingival and gingival flora with bonding and banding. *Angle Orthod.* **1987**, *57*, 271–278. doi: 10.1043/0003-3219(1987)057<0271:CIGAGF>2.0.CO;2.
17. Martin BJ, Campbell PM, Rees TD, Buschang, PH. A randomized controlled trial evaluating antioxidant-essential oil gel as a treatment for gingivitis in orthodontic patients. *Angle Orthod.* **2016**, *86*, 407–412. doi: 10.2319/041515-251.1.
18. Yeturu SK, Acharya S, Urala AS, Pentapati KC. Effect of Aloe vera, chlorine dioxide, and chlorhexidine mouth rinses on plaque and gingivitis: A randomized controlled trial. *J. Oral Biol. Craniofac. Res.* **2016**, *6*, 54-58. doi: 10.1016/j.jobcr.2015.08.008.
19. Addy M. Chlorhexidine compared with other locally delivered anti-microbials. A short review. *J. Clin. Periodontol.* **1986**, *13*, 957-964. doi: 10.1111/j.1600-051x.1986.tb01434.x.
20. Balenseifen JW, Madonia JV. Study of dental plaque in orthodontic patients. *J. Dent. Res.* **1970**, *49*, 320–324. doi: 10.1177/00220345700490022101.
21. Eliades T, Eliades G, Brantley WA. Microbial attachment on orthodontic appliances. I. Wettability and early pellicle formation on bracket materials. *Am. J. Orthod. Dentofacial Orthop.* **1995**, *108*, 351–360. doi: 10.1016/S0889-5406(95)70032-3.
22. Gazola Filho J, Leal Vizoto N, de Aguiar Loesch ML, Dias de Sena M, Mendes da Camara D, Sampaio Caiaffa K, de Oliveira Mattos-Graner R, Duque C. Genetic and physiological effects of subinhibitory concentrations of oral antimicrobial agents on Streptococcus mutans biofilms. *Microb. Pathog.* **2021**, *150*, 104669. doi: 10.1016/j.micpath.2020.104669.
23. Radovic R, Begic G, Lucic Blagojevic S, Karleusa LJ, Spalj S, Gobin I. Temporal dynamics of adhesion of oral bacteria to orthodontic appliances. *Dent. Mater. J.* **2023**, *42*, 469-477. doi: 10.4012/dmj.2022-235.
24. Alam F, Balani K. Adhesion force of staphylococcus aureus on various biomaterial surfaces. *J. Mech. Behav. Biomed. Mater.* **2017**, *65*, 872–880. doi: 10.1016/j.jmbbm.2016.10.009.
25. Nandhra SS, Littlewood SJ, Houghton N, Luther F, Prabhu J, Munyombwe T, et al. Do we need primer for orthodontic bonding? A randomized controlled trial. *Eur. J. Orthod.* **2015**, *37*, 147–155. doi: 10.1093/ejo/cju024.

26. An JS, Kim K, Cho S, Lim BS, Ahn SJ. Compositional differences in multi-species biofilms formed on various orthodontic adhesives. *Eur. J. Orthod.* **2017**, *39*, 528–533. doi: 10.1093/ejo/cjw089.
27. Jeon DM, An JS, Lim BS, Ahn SJ. Orthodontic bonding procedure influence biofilm composition. *Prog. Orthod.* **2020**, *21*, 14. doi: 10.1186/s40510-020-00314-8.
28. Mikulewicz M, Gronostajski Z, Wielgus A, Chojnacka K. Transparent orthodontic archwires: a systematic literature review. *Arch. Civ. Mech. Eng.* **2017**, *17*, 651–657. doi: 10.1016/j.acme.2016.12.012.
29. Dittmer MP, Helleman CF, Grade S, Heuer W, Stiesch M, Schwestka-Polly R, et al. Comparative three-dimensional analysis of initial biofilm formation on three orthodontic bracket materials. *Head Face Med.* **2015**, *11*, 1–6. doi: 10.1186/s13005-015-0062-0.
30. Oliveira DC, Thomson JJ, Alhabeil JA, Toma JM, Plecha SC, Pacheco RR, et al. In vitro *Streptococcus mutans* adhesion and biofilm formation on different esthetic orthodontic archwires. *Angle Orthod.* **2021**, *91*, 786–793. doi: 10.2319/121220-998.1
31. Abraham KS, Jagdish N, Kailasam V, Padmanabhan S. *Streptococcus mutans* adhesion on nickel titanium (NiTi) and copper-NiTi archwires: A comparative prospective clinical study. *Angle Orthod.* **2017**, *87*, 448–454. doi: 10.2319/040516-270.1
32. Raszewski Z, Nowakowska-Toporowska A, Wezgowiec J, Nowakowska D. Design and characteristics of new experimental chlorhexidine dental gels with anti-staining properties. *Adv. Clin. Exp. Med.* **2019**, *28*, 885–890. doi: 10.17219/acem/94152.
33. Poppolo Deus F, Ouanounou A. Chlorhexidine in dentistry: pharmacology, uses, and adverse effect. *Int. Dent. J.* **2022**, *72*, 269–277. doi: 10.1016/j.identj.2022.01.005.
34. Łukomska-Szymańska M, Sokołowski J, Łapińska B. Chlorhexidine – mechanism of action and its application to dentistry. *J. Stomatol.* **2017**, *70*, 405–417. doi: 10.5604/01.3001.0010.5698.
35. Solderer A, Kaufmann M, Hofer D, Wiedemeier D, Attin T, Schmidlin PR. Efficacy of chlorhexidine rinses after periodontal or implant surgery: a systematic review. *Clin. Oral Investig.* **2019**, *23*, 21–32. doi: 10.1007/s00784-018-2761-y.
36. Bonesvoll P, Lökken P, Rølla G, Paus PN. Retention of chlorhexidine in the human oral cavity after mouth rinses. *Arch. Oral Biol.* **1974**, *19*, 209–212. doi: 10.1016/0003-9969(74)90263-5.
37. Poppolo Deus F, Ouanounou A. Mouthwashes and their use in dentistry: a review. *Oral Health.* **2021**. Available online: <https://www.oralhealthgroup.com/features/mouthwashes-and-their-use-in-dentistry-a-review/> (accessed on 14 January 2024).
38. Laugisch O, Ramseier CA, Salvi GE, Hägi TT, Bürgin W, Eick S, et al. Effects of two different post-surgical protocols including either 0.05 % chlorhexidine herbal extract or 0.1 % chlorhexidine on post-surgical plaque control, early wound healing and patient acceptance following standard periodontal surgery and implant placement. *Clin. Oral Invest.* **2016**, *20*, 2175–2183. doi: 10.1007/s00784-016-1713-7.
39. Duss C, Lang NP, Cosyn J, Persson GR. A randomized, controlled clinical trial on the clinical, microbiological, and staining effects of a novel 0.05% chlorhexidine/herbal extract and a 0.1% chlorhexidine mouthrinse adjunct to periodontal surgery. *J. Clin. Periodontol.* **2010**, *37*, 988–997. doi: 10.1111/j.1600-051X.2010.01609.x
40. Cortellini P, Labriola A, Zambelli R, Prato GP, Nieri M, Tonetti MS. Chlorhexidine with an anti-discoloration system after periodontal flap surgery: A cross-over, randomized, triple-blind clinical trial. *J. Clin. Periodontol.* **2008**, *35*, 614–620. doi: 10.1111/j.1600-051X.2008.01238.x
41. Thangavelu A, Kaspar S, Kathirvelu R, Srinivasan B, Srinivasan S, Sundram R. Chlorhexidine: an elixir for periodontics. *J. Pharm. Bioallied. Sci.* **2020**, *12*, 57. doi: 10.4103/jpbs.JPBS_162_20.
42. Almohefer SA, Levon JA, Gregory RL, Eckert GJ, Lippert F. Caries lesion remineralization with fluoride toothpastes and chlorhexidine - effects of application timing and toothpaste surfactant. *J. Appl. Oral. Sci.* **2018**, *26*, e20170499. doi: 10.1590/1678-7757-2017-0499.
43. Brookes ZLS, Bescos R, Belfield LA, Ali K, Roberts A. Current uses of chlorhexidine for management of oral disease: a narrative review. *J. Dent.* **2020**, *103*, 103497. doi: 10.1016/j.jdent.2020.103497.
44. Kulkarni PJ, Singh DK, Jalaluddin M, Mandal A. Comparative evaluation of antiplaque efficacy between essential oils with alcohol-based and chlorhexidine with nonalcohol-based mouthrinses. *Int. Soc. Prev. Community Dent.* **2017**, *7*(Suppl 1), S36-S41. doi: 10.4103/jispcd.JISPCD_131_17.
45. Bescos R, Ashworth A, Cutler C, Brookes ZL, Belfield L, Rodiles A et al. Effects of Chlorhexidine mouthwash on the oral microbiome. *Sci. Rep.* **2020**, *10*, 5254. doi: 10.1038/s41598-020-61912-4.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.