

Review

Not peer-reviewed version

Functionality and Usability of mHealth Apps in Patients with Peritoneal Dialysis: A Scoping Review

[Shu-Mei Chao](#) , Chao-Kuei Pan , Ming-Ling Wang , Yu-Wen Fang * , [Shu-Fen Chen](#) *

Posted Date: 23 February 2024

doi: 10.20944/preprints202402.1295.v1

Keywords: mHealth; peritoneal dialysis; functionality; usability



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

Functionality and Usability of mHealth Apps in Patients with Peritoneal Dialysis: A Systematic Review

Shu-Mei Chao ¹, Chao-Kuei Pan ¹, Ming-Ling Wang ², Yu-Wen Fang ^{1,*} and Shu-Fen Chen ^{3,*}

¹ Department of Nursing, School of Nursing, Tzu Chi University of Science and Technology, Hualien, Taiwan. R.O.C.

² Researcher, School of Nursing, International Ph. D Program in Gerontology and Long-term Care, Taipei Medical University, Taipei, Taiwan

³ Department of Nursing, Shuang Ho Hospital, Taipei Medical University, New Taipei City, Taiwan

* Corresponding author: ^{1st} Y.-W. F, yonned1998@gmail.com; ^{2nd} S.-F. C, moon2014100@gmail.com; Tel.: +886-958-959-318

Abstract: mHealth has been utilized in the care of patients with chronic kidney disease, allowing the collection of patient health-related data, offering disease-related information, enabling the tracking and recording of biochemical parameters, and enabling communication to healthcare providers in real-time through applications. mHealth may improve the health outcomes in patients with peritoneal dialysis. This systematic review aimed to summarize evidence regarding the functionality and usability of mHealth apps in patients with peritoneal dialysis. We conducted a comprehensive literature review, searching five databases, including CINAHL, Cochrane, PsycINFO, PubMed, and Web of Science, to retrieve titles and abstracts related to peritoneal dialysis and mHealth applications for PRISMA guidelines from January 2013 to December 2023. Overall, 11 studies met all the inclusion criteria. The functionality of mHealth apps included inform, instruct, record, display, guide, remind/alert, and communicate. Most of the apps have multifunctionality. The outcomes (usability) were categorized into three aspects: efficiency (self-efficacy and usability), satisfaction, and effectiveness (underwent kidney transplantation and switched to hemodialysis, rehospitalization, peritonitis rate, infection rates at exit sites, mortality, fluid overload, inadequate solute clearance, biochemical values, quality of life, consumer quality index, and technology readiness). Generally, outcomes in the intervention group had better effects compared to those in the control group. Multifunctional mHealth apps show good potential in improving the efficiency, satisfaction, and effectiveness for patients compared to traditional care. Future research should include more studies and participants to explore and verify the long-term effectiveness of mHealth apps.

Keywords: mHealth; peritoneal dialysis; functionality; usability

1. Introduction

Unlike hemodialysis (HD), peritoneal dialysis (PD) does not require patients to visit hospitals for dialysis two to three times a week, allowing them to maintain their regular work and social activities, preserving their independence and freedom [1]. Additionally, the cost of hemodialysis was 1.37 times compared to peritoneal dialysis in Taiwan, and opting for PD can reduce healthcare costs. Despite these advantages, most patients continue to prefer HD [2]. The reluctance toward PD may stem from the fact that it involves self-administration or caregiver assistance, requiring patients to manage the dialysis process, including attaching the dialysis bag and manually recording related data on paper [3], which can be challenging for some patients.

While patients can communicate with healthcare providers (HCP) over the phone to address any issues during daytime dialysis [4], seeking advice from them at night may pose problems due to their unavailability. Furthermore, for patients without a medical background, HCP must impart additional disease-related knowledge and provide opportunities for them to practice dialysis techniques. The lack of a medical background and sudden complications can cause patients to feel

overwhelmed, resulting in them choosing HD over PD [1]. Mobile health (mHealth) applications (apps) have the potential to increase patients' willingness to adopt PD. These apps can record PD conditions, enhance patients' self-practice of PD techniques at home, alleviate concerns about unmanageable health conditions, help patients with chronic kidney diseases understand their condition, strengthen disease self-management [5-7], facilitate two-way communication and data sharing between patients and HCP, assist patients in gaining necessary knowledge and skills, and support patients in problem-solving and decision-making [1, 8-11], ultimately increasing patients' sense of security and satisfaction.

mHealth provides healthcare services through portable devices such as smartphones, tablets, computers, and personal digital assistants [12, 13]. mHealth delivers pre-established health services and provides guidance regarding the PD procedure and disease-related information in real-time through features such as browsing, text messaging, communication, and videos [11, 14, 15]. mHealth apps enable the collection of patient health-related data, including access to personal medical records, obtaining nutritional or dietary information, receiving treatment-related details, offering disease-related information, understanding health status, tracking exercise activities, weight, and dietary habits, monitoring medication dosage, recording blood pressure, and providing convenience and efficiency for patients with PD [3, 4, 6, 9, 14, 16-19]. Additionally, some mHealth apps have interactive features that allow patients to communicate immediately with physicians or nurses when difficulties arise [6]. On identifying problems in patient data transmissions, HCP can proactively contact the patients. These features contribute to patients feeling reassured and maintaining their independence [3].

However, concerns such as unsuitable design, data reliability [13, 20], inadequate user technological literacy, and privacy and security concerns [3, 19] may affect patients' willingness to use mHealth services, and the efficacy of such services remains unclear. If mHealth apps lack scientific basis, certification, or regulatory requirements, they may be unsuitable or potentially harmful for patients [8]. A systematic literature review of the effectiveness of eHealth for patients with PD showed inconsistent results regarding infection rates, hospitalization outcomes, knowledge, skills, satisfaction, and quality of life [8]. However, van Eck van der Sluijs et al. (2021) suggested that eHealth can reduce the incidence of peritonitis [16]. Therefore, to determine the overall impact of mHealth apps on patients with PD, a comprehensive analysis through systematic evaluation is necessary. Thus, this study aimed to conduct a systematic review to explore the features, functions, and usability of mHealth apps used by patients with PD.

In terms of usage, each mobile app emphasizes different things, making it challenging to establish a standard for systematic evaluation [13]. To clarify the scope of assistance provided by mHealth apps for patients with PD, this study focused on the usability and functionality of mobile apps. Regarding functionality, we referred to the seven functions proposed by the Institute for Healthcare Informatics (IMS) for mHealth apps: inform, instruct, record, display, guide, remind/alert, and communicate [21]. These functions have been widely evaluated in mHealth apps, such as those for asthma [22], chronic kidney diseases [17], and heart failure [23]. In terms of usability assessment, Harrison et al. (2013) [24] and Weichbroth (2020) [25] advocated that an ideal mHealth app should be simple and user-friendly. According to Weichbroth (2020), an app's usability can be assessed by exploring three aspects: efficiency, satisfaction, and effectiveness [25]. Therefore, in this study, we explored the aforementioned aspects to evaluate and analyze the usability of mHealth apps.

mHealth apps have been widely used to enhance the care of patients with PD. However, whether they effectively improve health requires a comprehensive and concrete analysis. Therefore, this systematic review aimed to investigate studies on the use of mHealth apps in patients with PD in the following aspects: (1) features, (2) functionalities, and (3) usability.

2. Materials and Methods

This systematic review was reported in accordance with **PRISMA guidelines**, was conducted by two reviewers following the systematic review and meta-analysis expansion methodology [26, 27]. This review aimed to describe literature retrieval and study selection using the PRISMA checklist

for reporting systematic reviews. In case of inconsistencies in the results, a third reviewer was consulted. This review involved mapping the entirety of the relevant literature to identify and map features, functionalities, and usability related to mHealth apps for PD. The findings contribute not only to identifying empirical patterns and resources for clinical practice but also to recognizing the existing limitations of apps for future enhancement.

2.1. Search Strategy

Between January 2013 and December 2023, five databases—CINAHL, Cochrane, PsycINFO, PubMed, and Web of Science—were searched to retrieve titles and abstracts related to PD and mHealth apps. The search strategy incorporated MeSH terms and keywords such as “mHealth,” “mobile health,” “mobile app,” “mobile application,” “smartphone application,” “app,” “apps,” “telemedicine,” “peritoneal,” “dialysis,” and “peritoneal dialysis.” The following criteria were applied for selecting research articles: (1) full text; (2) English language; (3) studies classified according to Joanna Briggs Institute (JBI) levels of evidence, including experimental designs, quasi-experimental designs, observational – analytic designs, observational – descriptive studies, expert opinion, and bench research [28]; and (4) exclusion of study protocol, scoping literature, literature reviews, systematic reviews, study protocols, systematic reviews, integrative analysis reviews, conference abstracts, letters to the editor, and editorials and similar articles.

2.2. Inclusion and Exclusion Criteria

A review of the use of mHealth apps in PD was conducted. The inclusion criteria were as follows: (1) targeted patients with PD; (2) mobile app interventions delivered through smartphones, tablets, or web portals accessible via mobile devices; (3) aimed at enhancing patient care and management of the disease. The exclusion criteria were as follows: (1) not a mobile app (e.g., eHealth remote healthcare, electronic health records, and electronic medical records); and (2) designed and developed without adoption in patients.

2.3. Screening and Data Extraction

Title, abstract, and full-text screening was independently conducted by two reviewers (CS and FY). All identified articles were uploaded to EndNote 20, and duplicate articles were removed. The two reviewers used three predesigned forms to extract data from the eligible articles. The first form gathered the following information: author/year/country, research objectives, **app name/ device /platform/ objectives**, design/ sample size/ duration, intervention content, **and outcome variables/ results**. The second form assessed the functionality of mHealth apps by utilizing the seven functions proposed by IMS: inform, instruct, record, display, guide, remind/alert, and communicate. The third form focused on evaluating the apps’ usability by exploring the aspects of efficiency, satisfaction, and effectiveness. The extracted data were synthesized and analyzed using the three aforementioned forms.

2.4. Selection Process

A total of 184 articles were extracted from the five databases. After removing duplicates (n=10) as well as abstracts and articles that did not meet the inclusion criteria, 153 articles were excluded, leaving 21 articles for consideration. Two independent reviewers conducted the screening based on the inclusion and exclusion criteria and resolved any discrepancies through discussions. Based on the inclusion and exclusion criteria, 10 articles were subsequently excluded, resulting in a final set of 11 articles for the systematic review. The search results and study inclusion process are presented in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses **flowchart** (Figure 1). (Figure 1 insert here)




Figure 1. Flow diagram of study selection.

2.5. Functionality Assessment

The app's functionality was evaluated using the guidelines developed by the IMS [21] as follows: (1) inform: provide information in the form of text, photos, videos, etc.; (2) instruct: offer instructional guidance; (3) record: gather physiological data and other relevant patient information; (4) display: data entered by users and recipients in a graphical format; (5) guide: analyze user-inputted data and suggest relevant treatments; (6) remind/alert: send reminders or alerts to notify patients; (7) communicate: patients interact with HCP, peers, or community forums. These functionalities serve various purposes and enhance the overall capabilities of mHealth apps.

2.6. Usability Assessment

The usability of mHealth apps can be assessed by exploring their efficiency, satisfaction, and effectiveness [25]. (1) Efficiency refers to the user's ability to achieve goals quickly and accurately as well as the error rate during use. (2) Satisfaction relates to the user's comfort, happiness, meeting expectations, and fulfilling needs. (3) Effectiveness refers to the ability to accomplish tasks within a certain time frame. The two reviewers categorized the research results based on the aforementioned definitions of the components of usability. In case of any inconsistencies in the classification, a third reviewer reassessed the cases until a consensus was reached. This systematic approach ensured a thorough and standardized evaluation of the usability of mHealth apps.

3. Results

3.1. Characteristics of the Included Studies

Eleven articles published between 2013 and 2023 were included in the final review. Table 1 presents the characteristics of each mHealth app. Three studies were conducted in Mexico [29-31], two in Canada [32, 33], two in China [10, 34], and one each in Korea [35], the United Kingdom [11], Thailand [7], and Italy [36] (Table 1). The study designs varied, with three using experimental designs (Randomized Controlled Trials [RCTs]) [10, 32, 35], six using observational-analytic designs [7, 29, 31, 33, 34, 36], and two being observational – descriptive studies [11, 30]. mHealth apps were used on various devices: 27.2% (3/11) on smartphones, 18.2% (2/11) on tablets, 9.1% (1/11) on computers, and 72.7% (8/11) on unspecified devices. In terms of operating systems, 45.5% (5/11) were Android-based, 36.4% (4/11) were iOS-based, 36.4% (4/11) were Internet-based, and 45.5% (5/11) were unspecified. The measurement durations of these studies ranged from 10 weeks to 285 months. (Table 1 insert here)

Table 1. Characteristics of the included studies on patients with peritoneal dialysis (PD).

Authors/Year/Country ^{Ref no}	App name/ Device /Platform/ Objectives	Design/ Sample Size / Duration	Intervention Content	Outcome variables/ Results
Cao et al./2018 / China ¹⁰	<ul style="list-style-type: none"> ·The QQ Application · Smartphone & tablet · Android & <p>·To investigate the effectiveness of the QQ application.</p>	<ul style="list-style-type: none"> ·A randomized controlled trial design ·the QQ (experimental) group (n=80) the traditional follow-up group (n=80) ·11.4 months 	Nurses provide health education and disease-related information to patients through the QQ application, engaging in online conversations to address their health concerns.	<ul style="list-style-type: none"> ·Satisfaction, underwent kidney transplantation and switched to hemodialysis, rehospitalisation, infection rates, serum values ·(1) Patients in the QQ intervention group showed significantly higher levels of serum albumin, hemoglobin, and satisfaction; as well as lower levels of phosphorus and calcium-phosphorus products compared to those in the control group. (2) There was no difference in underwent kidney transplantation and switched to hemodialysis, as well serum calcium levels between the two groups.
Chae & Kim/2023 / Korea ³⁵	<ul style="list-style-type: none"> · PD With You · N/A · Android 9.0 <p>·Developing a mobile application for enhancing self-management and assessing its effectiveness for patients with PD.</p>	<ul style="list-style-type: none"> ·A randomized controlled trial ·the experimental group (n =27) the control group (n =26) ·10 weeks 	Patients used the self-management mobile application to record PD dialysis (i.e., replacement time and amount of removed fluid) and observe the physical indicators (i.e., body weight and blood pressure).	<ul style="list-style-type: none"> ·self-efficacy, serum values, knowledge, health behavior, QOL ·(1) The experimental group showed a significant improvement in PD-related knowledge, PD-related health behavior, serum albumin, hemoglobin level, and the domain symptoms/problems of kidney disease and disease impact on daily activity of HRQOL, compared to the control group. (2) There were no significant differences between the two groups in PD-related self-efficacy, serum K, serum P, the domains of physical, psychological, and burden of kidney disease of HRQoL.
Dey et al./2016/ UK ¹¹	<ul style="list-style-type: none"> · N/A · Computer tablets · N/A <p>· (1) The application can provide information related to diseases and</p>	<ul style="list-style-type: none"> ·Observational–Descriptive Studies(A cross-sectional study) ·22 participants ·Over 15 months 	Each patient is provided with a weight scale, blood pressure monitor, and computer tablets equipment. The patients were required to record data, such as	<ul style="list-style-type: none"> ·satisfaction, QOL ·(1) No significant improved in QOL. (2) No significant improved in satisfaction with assistive Technology.

Authors/Year/Country ^{Ref no}	App name/ Device /Platform/ Objectives	Design/ Sample Size / Duration	Intervention Content	Outcome variables/ Results
	<p>dietary considerations. (2) Patients can modify the dialysis plan or perform other actions, such as manual exchanges when dialysis resulted in problems. (3) Providing dietary advice through phone consultations by HCP. (4) Recording and accessing personal physiological data and details of dialysis sessions.</p>		<p>weight, blood pressure, dialysis exchanges, and ultrafiltration volume. Additionally, patients need to report on the presence or absence of symptoms, such as swelling, shortness of breath, fever, abdominal pain, tenderness around the catheter site, and other relevant conditions. When the patients' condition is not well, HCP will provide interventions or treatments.</p>	
Kiberd et al./ 2018/ Canda ³³	<ul style="list-style-type: none"> · McKesson · eHealth portal applications · N/A <p>·To investigate whether eHealth portal applications can effectively enhance the home dialysis care experience for patients.</p>	<ul style="list-style-type: none"> ·Observational–Analytic Designs, without a control group (A Single-Arm Pilot)Study ·participants (n =27) ·12 months 	<p>Patients and HCP can communicate through a web portal application. HCP can provide suggestions such as medication changes, post-clinic visit explanations, new appointment scheduling, and more.</p>	<ul style="list-style-type: none"> · satisfaction, acceptance, QOL, consumer quality index <p>· (1) Most participants were satisfied. (2) QOL did not improve, and it was difficult to ascertain the user acceptance due to the small sample size. (3) No significance in the consumer quality index</p>
Lukkanalikitkul et al /2022/Thailand ⁷	<ul style="list-style-type: none"> ·CKD-PD app ·Smart phone & near-field communication & optical character recognition ·Android & iOS <p>·To identify Chronic Kidney Disease - Peritoneal Dialysis Application with near-field communication and optical character recognition functions, which can automatically gather the hydration status to enhance care for patients with PD, and improve it.</p>	<ul style="list-style-type: none"> ·Observational– Analytic Designs (User- Centered Design Study) ·participants (n=10) ·12 months 	<p>Provide an application to collect the peritoneal dialysis fluid and upload the data to HCP.</p>	<ul style="list-style-type: none"> ·usability <p>·In the end, participants decreased interest in using this app. Especially, the entering data in NFC and OCR system of the app was difficult for participants compared to the manual data.</p>

Authors/Year/Country ^{Ref no}	App name/ Device /Platform/ Objectives	Design/ Sample Size / Duration	Intervention Content	Outcome variables/ Results
Martínez García et al./2018 /Mexico ²⁹	<ul style="list-style-type: none"> ·N/A ·A mobile Web application (for HCP) & Android application(for patients) ·Android ·Evaluating the usability of a remote monitoring system for patients undergoing peritoneal dialysis treatment. 	<ul style="list-style-type: none"> ·Quasi-experimental design (A case study) ·participants (n=24) · 9 months 	Providing applications to patients to monitor the patient' s condition and assess the usability of the app.	<ul style="list-style-type: none"> ·satisfaction, acceptance ·(1) 94.5% of participants were satisfied with the app among patients with APD and 92.3% among CAPD. (2) 89.5% of participants accepted.
Olivares-Gandy et al./2019/ Mexico ³⁰	<ul style="list-style-type: none"> ·N/A ·Android & iOS · N/A ·To Analyze, design, and develop a mobile health application for the dietary requirements of patients with PD, and then assessing its usability and satisfaction. 	<ul style="list-style-type: none"> ·Observational–Descriptive (A case study) ·one patient & one nutritionist ·N/A 	Patients' experiences with the use of mobile applications.	<ul style="list-style-type: none"> ·usability ·Patients find the application usable.
Polanco et al./2021/ Mexico ³¹	<ul style="list-style-type: none"> ·WhatsApp ·Computer, smartphone, tablet · N/A ·The healthcare team utilizes remote healthcare via WhatsApp to reduce hospital visits and the risk of infection during the COVID-19 pandemic. 	<ul style="list-style-type: none"> ·Observational– Analytic Designs (An observational prospective-longitudinal study) ·participants (n=946) ·3 months 	The patients send daily dialysis records and photos of their lower limbs to HCP by using WhatsApp. The medical team members utilize WhatsApp for communication and tracking the patient's condition.	<ul style="list-style-type: none"> ·switched to hemodialysis, rehospitalisation, peritonitis rate ·The incidence rate of peritonitis, switched to hemodialysis, and hospitalization rates showed no difference.
Viglino et al./2020/ Italy ³⁶	<ul style="list-style-type: none"> ·N/A ·N/A ·N/A ·To explore the reliability, safety, and effectiveness of videodialysis assisted PD patients. To investigate the possibilities of the videodialysis and whether it reduced family care burden and recourse to nurses at home. 	<ul style="list-style-type: none"> ·Observational–Analytic Designs ·the intervention group (n=15) ·the control group (n=62) ·Follow-up 285 months 	The experimental group was provided with videodialysis, a device consisting of two parts. One part included the equipment that patients should have at home, including a camera, monitor, microphone, and a technical connection box. The other part comprised the equipment required	<ul style="list-style-type: none"> ·satisfaction, switched to hemodialysis, Peritonitis rate ·(1) All the patients expressed a satisfaction with the app for enhance confidence. (2) There is no difference in the incidence of peritonitis. (3) Three out of the 15 participants transitioned to hemodialysis.

Authors/Year/Country ^{Ref no}	App name/ Device /Platform/ Objectives	Design/ Sample Size / Duration	Intervention Content	Outcome variables/ Results
			on the healthcare personnel's end, including a high-resolution display, a network camera, a computer with speakers, and software capable of managing six audio streams simultaneously.	
Xu et al./2022/China ³⁴	<ul style="list-style-type: none"> ·Manburs ·N/A ·N/A ·To explore the long-term impact of telemedicine (Manburs app) on patients in terms of mortality and technical performance failure.	<ul style="list-style-type: none"> ·Observational–Analytic Designs (A Propensity-Matched Study) ·Participants (n=7539) ·From June 2016 to December 2020 (4 years) 	Telemedicine, through the application (Manburs), involves self-monitoring records, online educational materials, and real-time doctor-patient communication.	<ul style="list-style-type: none"> ·switched to hemodialysis, infection rates, mortality, fluid overload, inadequate solute clearance, · The intervention group were observed to have significantly lower risks of all-cause mortality, CVD mortality, all-cause transfer to hemodialysis, transfer to hemodialysis from PD-related infection, severe fluid overload, inadequate solute clearance, and catheter-related noninfectious complications compared with that in the control group.

Notes: PD: Peritoneal dialysis; N/A: Not applicable; HCP: health care providers; QOL: quality of life; APD: automatic peritoneal dialysis; CAPD: continuously automatic peritoneal dialysis; CVD; cardiovascular diseases

3.2. Interventions Description

Regarding the functionality of mHealth apps, each app included “Inform,” “Instruct,” and “Communicate” features with a 100% inclusion rate (11/11). Regarding other features, “Display” was in 90.9% (10/11) of the apps, “Guide” in 90.9% (10/11), “Record” in 81.8% (9/11), and “Remind/Alert” —the least common feature—in 72.7% (8/11) of the apps (Table 2). The intervention content of these apps included disease-related information. Patients could record and transmit details such as the fluid volume during dialysis or the condition of dialysis wounds. Additionally, in case of symptoms or problems during dialysis, patients could communicate and interact with HCP or peers. (Table 2 insert here).

Table 2. IMS Functionality of mHealth Apps for Patients with Peritoneal Dialysis.

Authors(Year)	Functionality						
	Inform	Instruct	Record	Display	Guide	Remind/Alert	Communicate
Cao et al.(2018)	✓	✓	✓	✓	✓	✓	✓
Chae & Kim(2023)	✓	✓	✓	✓	✓	✓	✓
Dey et al.(2016)	✓	✓	✓	✓	✓	✓	✓
Farfan-Ruiz et al.(2021)	✓	✓		✓	✓		✓
Kiberd et al.(2018)	✓	✓	✓	✓	✓	✓	✓
Lukkanalikitkul et al.(2022)	✓	✓	✓	✓	✓	✓	✓
Martínez García et al. (2018)	✓	✓	✓	✓	✓	✓	✓
Olivares-Gandy et al.(2019)	✓	✓	✓	✓	✓	✓	✓
Polanco et al.(2021)	✓	✓	✓	✓	✓	✓	✓
Viglino et al.(2020)	✓	✓		✓	✓		✓
Xu et al.,(2022)	✓	✓	✓				✓

3.3. Outcomes

The outcomes were categorized into three main types: efficiency, satisfaction, and effectiveness (Table 3). In one study that investigated PD-related self-efficacy, no statistically significant differences were found [35]. Two studies that explored usability drew conflicting conclusions, with one study reporting positive [30] and the other negative usability [7]. All five studies that assessed satisfaction reported positive outcomes [10, 11, 29, 33, 36]. For acceptance, which was studied in one study, positive results were reported [29]. Regarding effectiveness, a lesser number of patients from the intervention group switched from PD to HD or kidney transplantation compared with those from the control group [34]; however, three studies reported no significant differences [10, 31, 36]. In two studies, re-hospitalization rates showed no significant differences [10, 31]. The peritonitis occurrence rates showed no significant differences in the two studies [31, 36]. Infection rates at the exit site and mortality rates were examined in two studies, with one showing better outcomes in relation to fluid overload and inadequate solute clearance in the experimental group [34] and the other reporting no significant differences [10]. Inconsistent results for biochemical parameters were found across studies. Serum albumin and hemoglobin levels were higher in the experimental group in two studies [10, 35]. However, changes in serum calcium levels did not differ significantly in two studies [10, 32]. Mixed results were obtained for serum phosphorus levels, with two studies reporting no significant differences [32, 35] and one showing better outcomes in the experimental group [10]. While no significant differences were found in the changes in serum potassium levels in a study [35], the calcium-phosphorus level was better in the experimental group in another study [10]. Taking calcium carbonate yielded no significant differences [32]. PD-related knowledge and health behaviors were better in the experimental group in one study [35]. Quality of life results varied, with one study showing better outcomes in the experimental group [35] and two studies reporting no significant

differences [11,33]. No significant differences were observed in the Consumer Quality Index [33] and Technology Readiness [32]. (Table 3 Insert here).

Table 3. Usability of mHealth Apps for Patients with Peritoneal Dialysis.

	Outcomes	Statistical significance		
		Positive	Negative	No difference
Efficiency	PD-related self-efficacy			Chae & Kim(2023)
	Usability	Olivares-Gandy et al.(2019)	Lukkanalikitkul et al. (2022)	
Satisfaction	Patients' satisfaction	Cao et al.(2018) Dey, et al.(2016) Kiberd et al.(2018) Martínez García et al. (2018) Viglino et al.(2020)		
	Acceptance	Martínez García et al. (2018)		Kiberd et al.(2018)
Effectiveness	Underwent kidney transplantation and switched to hemodialysis	Xu et al.(2022)		Cao et al.(2018) Polanco et al.(2021) Viglino et al.(2020) ^b
	Rehospitalisation			Cao et al.(2018) Polanco et al.(2021)
	Peritonitis rate			Polanco et al.(2021) Viglino et al.(2020)
	The infection rates at the exit site	Xu et al.(2022)		Cao et al.(2018)
	Mortality	Xu et al.(2022)		Cao et al.(2018)
	Fluid overload	Xu et al.(2022)		
	Inadequate solute clearance	Xu et al.(2022)		
	Serum albumin	Cao et al.(2018) Chae & Kim/2023		
	Serum Hemoglobin	Cao et al.(2018) Chae & Kim(2023)		
	Serum calcium			Cao et al.(2018) Farfan-Ruiz et al.(2021)
	Serum phosphorus	Cao et al.(2018) Farfan-Ruiz et al.(2021) ^a		Chae & Kim(2023)
	Serum potassium			Chae & Kim(2023)
	Calcium-phosphorus product	Cao et al./2018		
	Taking calcium carbonate			Farfan-Ruiz et al.(2021)
	PD-related knowledge	Chae & Kim(2023)		
	PD-related health behavior	Chae & Kim(2023)		
Quality of life	Chae & Kim(2023) ^a		Dey et al.(2016) Kiberd et al.(2018)	

	Consumer quality index (CQI)			Kiberd et al.(2018)
	Technology readiness			Farfan-Ruiz et al.(2021)

Notes: Positive: The intervention group was better than the control group; Negative: The control group was better than the intervention group; ^a : mixed effects; ^b : Not clear ; N/A: Not applicable.

4. Discussion

In this study, a systematic review was conducted to identify and map the existing evidence regarding the features, functionalities, and usability of mHealth apps for patients with PD. The findings revealed that the apps primarily focused on self-efficacy in performing PD procedures, user satisfaction, and monitoring and improving health conditions. Regarding functionality, mHealth apps mostly encompassed features listed by the IMS. Regarding usability, the findings are similar to those found in Cartwright et al.'s (2021) systematic review on eHealth [8]; however, the results pertaining to the outcomes are inconsistent.

The most successful feature for mHealth apps should be multifunctionality [22]. In this systematic review, most apps were found to have multiple functions. Functions such as record, display, guide, remind/alert, communicate [7, 35], and facilitating remote monitoring [8]. They allow patients to upload data, such as fluid volumes during the PD process, enabling communication and interaction between HCP and patients to ensure treatment outcome. HCP can also track patient treatment results [10]. These results are consistent with those of Eberle et al. (2021) who found that apps improved laboratory data and facilitated health promotion [37]. Moreover, these apps provide disease-related information for patients with PD, resulting in increased user satisfaction.

The usability of an app can be assessed by exploring three aspects: efficiency, satisfaction, and effectiveness. Some results obtained in this review seem to differ from those in previous studies. In terms of efficiency, the apps did not seem to enhance patients' self-efficacy. These results are inconsistent with the findings of Eberle et al. (2021) who indicated that patients with diabetes who used mHealth apps strengthened their self-efficacy [37]. This inconsistency may be attributed to the means of communicating or interacting with HCP, which are limited to phone calls, text messages, or monthly interviews, contributing to patients feeling less confident about dealing with disease-related situations. This could have influenced their lack of self-efficacy [35]. Furthermore, there was only one study on self-efficacy, and the intervention duration was only ten weeks; therefore, confirming whether apps can enhance self-efficacy is difficult. Similar findings were observed regarding usability. Lukkanalikitkul et al. (2022) believed that poor usability is due to unsuitable software development and limited Internet speeds, which do not meet user needs [7]. However, Olivares-Gandy et al. (2019) who focused on the app-development process, finding shortcomings for improvement, and allowing for timely adjustments to the app based on patient needs found the usability to be effective [30].

All study findings reported patient satisfaction with app usage [10, 11, 29, 33, 36]. The primary advantage of mHealth apps is that they are not bound by time or space constraints, allowing users to access disease-related information, record personal data (e.g., blood pressure, weight, and PD fluid), and monitor their health status at any time. Patients instantly communicated with HCP on encountering any problems or difficulties. Patient satisfaction significantly improved if the app features were comprehensive, practical, user-friendly, and suitable for patients [29, 30].

The effectiveness of apps in improving biochemical parameters revealed inconsistencies; while Eberle et al. (2021) showed some improvements in patients with diabetes, no statistically significant differences were found in others [37]. Continuous monitoring of nutritional management can help maintain normal hemoglobin and albumin levels in patients [35]. Learning fatigue, resulting in a low willingness to use apps, may occur when patients need to learn how to use apps correctly, which could be a contributing factor in the lack of improvement in physiological health indicators. In addition, the mHealth apps design, not aligning with the needs of patients and being difficult to operate, may contribute to increased frustration and discouragement among patients. It could lead to no ideally effective outcomes [32]. However, in a four-year study about the usage of a PD app, Xu

et al. (2022) found improvements in all physiological indicators, suggesting that the consistent and long-term use of apps to record daily life activities may yield effective outcomes [34].

App usage can enhance patients' quality of life, particularly in areas where the symptoms and disease impact daily living [22, 35]. However, no significant differences were observed in users' physiological and psychological domains of quality of life. Kiberd et al. (2018) indicated that the lower use of apps might be due to high rate of patient dropout [33]. Farfan-Ruiz et al. (2021) indicate that improvements in users' technological readiness might be associated with gender, educational level, and race [32].

5. Conclusions

Although the functional features of mHealth apps for patients with PD are mostly complete, their effectiveness varies. This inconsistency may be due to the fact that only three RCT studies were included in this review, of which two study samples, both in the experimental and control groups, comprised fewer than 30 participants, which might have led to poor-quality evidence for intervention outcomes. Comparing the usability of apps between the experimental and control groups, some studies showed no significant differences in effectiveness, whereas others did. Health outcomes with mobile apps may be better than or equivalent to those of traditional healthcare. Future research should include more studies with larger sample sizes to explore and verify the long-term effectiveness of mHealth apps. In this systematic review, user satisfaction with apps was found to be generally high. Thus, using these apps in the care of patients with PD to promote self-management of diseases and enhance health outcomes is recommended.

The ideal mHealth apps should have the functionality to record and monitor patients' physiological conditions and provide reminders or alerts for abnormal situations to patients and HCP. More importantly, it must feature interactive functions with HCP to increase patients' willingness to use it. Additionally, designing user-friendly software from the perspective of the patients is crucial. Only through user-friendly design can the mHealth apps fully function as an auxiliary tool for promoting healthcare.

Strengths and limitations

This study adheres to a systematic literature review approach by conducting a thorough and comprehensive search of the most common academic literature databases. Although the study findings suggest potential progress in the health of patients with PD as a result of using mHealth apps, this study has some limitations.

First, in eleven studies, only three studies were RCTs. This may have resulted in weak evidence for the study outcomes. However, owing to the limited research on mHealth apps, researchers have considered various study designs to broaden the scope of the systematic review. Nevertheless, this approach may introduce selection bias, and the overall quality of the evidence may be low. Second, grey literature, non-full-text articles, non-English publications, and conference reports in the field were excluded from this review. This omission may have led to valuable research in this field being excluded; therefore, the results should be interpreted cautiously. Third, this review only included studies published within the last 10 years, potentially overlooking valuable research published earlier. Finally, studies from different countries and regions were within the scope of the review, making it challenging to generalize the findings.

Author Contributions: S.-M. C. and Y.-W. Fang -Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Original draft preparation, Visualization, Supervision, and Project administration. S.-M. C. and S.-F. C. -Visualization, Supervision, Project administration. Validation, Investigation, and Data Curation. M.-L. W.- Software, Validation, and Formal analysis. S.-F. C. -Investigation. C.-K. P.-Data curation, Visualization. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by the National Science and Technology Council, Taiwan: NSTC 111-2314-B227-001. Tzu Chi University of Science and Technology, Taiwan: TCCT-1111A26.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available from the first author and availability Statements are available in section “MDPI Research Data Policies” at <https://www.mdpi.com/ethics>.

Acknowledgments: The authors wish to acknowledge the Department of Nursing, Tsu-Chi Hospital, Hualian.

Conflicts of Interest: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

1. Baumgart, A., et al., Meaning of empowerment in peritoneal dialysis: focus groups with patients and caregivers. *Nephrology Dialysis Transplantation*, 2020. 35(11): p. 1949-1958.
2. Tang, C. H., et al., Out-of-pocket costs and productivity losses in hemodialysis and peritoneal dialysis from a patient interview survey in Taiwan. *BMJ open*, 2019. 9(3): p. e023062.
3. Talbot, B., et al., Patient and Clinician Perspectives on the use of Remote Patient Monitoring in Peritoneal Dialysis. *Canadian Journal of Kidney Health and Disease*, 2022. 9: p. 20543581221084499.
4. Biebuyck, G.K., et al., Impact of telehealth interventions added to peritoneal dialysis-care: a systematic review. *BMC nephrology*, 2022. 23(1): p. 292.
5. He, T., et al., Remote home management for chronic kidney disease: a systematic review. *Journal of telemedicine and telecare*, 2017. 23(1): p. 3-13.
6. Markossian, T.W., et al., A mobile app to support self-management of chronic kidney disease: development study. *JMIR human factors*, 2021. 8(4): p. e29197.
7. Lukkanalikitkul, E., et al., Optimization of the Chronic Kidney Disease–Peritoneal Dialysis App to Improve Care for Patients on Peritoneal Dialysis in Northeast Thailand: User-Centered Design Study. *JMIR Formative Re-search*, 2022. 6(7): p. e37291.
8. Cartwright, E.J., et al., eHealth interventions to support patients in delivering and managing peritoneal dialysis at home: a systematic review. *Peritoneal Dialysis International*, 2021. 41(1): p. 32-41.
9. Schrauben, S.J., et al., Mobile health (mHealth) technology: assessment of availability, acceptability, and use in CKD. *American Journal of Kidney Diseases*, 2021. 77(6): p. 941-950. e1.
10. Cao, F., et al., Application of instant messaging software in the follow-up of patients using peritoneal dialysis, a randomized controlled trial. *Journal of Clinical Nursing*, 2018. 27(15-16): p. 3001-3007.
11. Dey, V., A. Jones, and E.M. Spalding, Telehealth: Acceptability, clinical interventions and quality of life in peritoneal dialysis. *SAGE open medicine*, 2016. 4: p. 2050312116670188.
12. Moss, R.J., A. Süle, and S. Kohl, eHealth and mHealth. *European Journal of Hospital Pharmacy*, 2019. 26(1): p. 57-58.
13. Lewis, R.A., et al., Identifying Mobile Applications Aimed at Self-Management in People With Chronic Kidney Disease. *Can J Kidney Health Dis*, 2019. 6: p. 2054358119834283.
14. Quach, S., et al., Features and characteristics of publicly available mHealth apps for self-management in chronic obstructive pulmonary disease. *Digital Health*, 2023. 9: p. 20552076231167007.
15. World Health Organization, mhealth Use of appropriate digital technologies for public health. 2018
16. van Eck van der Sluijs, A., et al., Good practices for dialysis education, treatment, and eHealth: A scoping review. *Plos one*, 2021. 16(8): p. e0255734.
17. Shen, H., et al., Electronic Health Self-Management Interventions for Patients with Chronic Kidney Disease: Systematic Review of Quantitative and Qualitative Evidence. *J Med Internet Res*, 2019. 21(11): p. e12384.
18. Ryan, S., et al., mHealth Apps for musculoskeletal rehabilitation: systematic search in APP stores and content analysis. *JMIR rehabilitation and assistive technologies*, 2022. 9(3): p. e34355.
19. Wilson, J., et al., Barriers and facilitators to the use of e-health by older adults: a scoping review. *BMC Public Health*, 2021. 21: p. 1-12.
20. Wang, C.-s. and E. Ku, eHealth in kidney care. *Nature Reviews Nephrology*, 2020. 16(7): p. 368-370.
21. IMS Institute for Healthcare Informatics. Patient Apps for Improved Healthcare. 2013.
22. Farzandipour, M., et al., Patient self-management of asthma using mobile health applications: a systematic re-view of the functionalities and effects. *Applied clinical informatics*, 2017. 8(04): p. 1068-1081.
23. Creber, R.M.M., et al., Review and analysis of existing mobile phone apps to support heart failure symptom monitoring and self-care management using the Mobile Application Rating Scale (MARS). *JMIR mHealth and uHealth*, 2016. 4(2): p. e5882.
24. Harrison, R., D. Flood, and D. Duce, Usability of mobile applications: literature review and rationale for a new usability model. *Journal of Interaction Science*, 2013. 1: p. 1-16.

25. Weichbroth, P., Usability of mobile applications: a systematic literature study. *Ieee Access*, 2020. 8: p. 55563-55577.
26. Moher, David, et al., Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine* 151.4 (2009): 264-269.
27. Moher, David, et al., Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *International journal of surgery* 8.5 (2010): 336-341.
28. Joanna Briggs Institute, JBI levels of evidence. 2014.
29. Martínez García, M.A., et al., Telemonitoring system for patients with chronic kidney disease undergoing peritoneal dialysis: usability assessment based on a case study. *PloS one*, 2018. 13(11): p. e0206600.
30. Olivares-Gandy, H.J., et al., A telemonitoring system for nutritional intake in patients with chronic kidney disease receiving peritoneal dialysis therapy. *Computers in Biology and Medicine*, 2019. 109: p. 1-13.
31. Polanco, E., et al., A COVID-19 pandemic-specific, structured care process for peritoneal dialysis patients facilitated by telemedicine: Therapy continuity, prevention, and complications management. *Therapeutic Apheresis and Dialysis*, 2021. 25(6): p. 970-978.
32. Farfan-Ruiz, A.C., et al., Multidisciplinary team versus a "Phosphate-Counting" app for serum phosphate control: a randomized controlled trial. *Kidney360*, 2021. 2(2): p. 290.
33. Kiberd, J., et al., Effectiveness of a web-based eHealth portal for delivery of care to home dialysis patients: a single-arm pilot study. *Canadian Journal of Kidney Health and Disease*, 2018. 5: p. 2054358118794415.
34. Xu, X., et al., Telemedicine and Clinical Outcomes in Peritoneal Dialysis: A Propensity-Matched Study. *American Journal of Nephrology*, 2022. 53(8-9): p. 663-674.
35. Chae, Y.J. and H.S. Kim, Effects of a mobile application on improving self-management of adult patients receiving peritoneal dialysis: A randomized controlled trial. *Japan Journal of Nursing Science*, 2023: p. e12555.
36. Viglino, G., et al., Videodialysis: a pilot experience of telecare for assisted peritoneal dialysis. *Journal of Nephrology*, 2020. 33: p. 177-182.
37. Eberle, C., M. Löhnert, and S. Stichling, Effectiveness of disease-specific mHealth apps in patients with diabetes mellitus: scoping review. *JMIR mHealth and uHealth*, 2021. 9(2): p. e23477. [CrossRef](#)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.