

Review

Not peer-reviewed version

Gene-Edited Medicinal 'Superplants': Advancements for Prolonged Space Travel

[Shreya Nair](#)*

Posted Date: 28 February 2024

doi: 10.20944/preprints202402.1643.v1

Keywords: Medicinal plants, long-duration space flight, gene-editing, radiation-induced cardiovascular disease, gut microbiome, microgravity



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

Gene-Edited Medicinal 'Superplants': Advancements for Prolonged Space Travel

Shreya Nair

University of Texas at Dallas, sxn220133@utdallas.edu

Abstract: Space exploration is an arduous journey, with future explorations extending in duration, increasing the risk for common space-related health issues such as bone density loss due to microgravity, impaired gut microbiota, cardiovascular ailments, and exposure to cosmic radiation. These ailments can be addressed using medicinal plants. Enhancing their efficacy entails employing gene editing to target specific secondary metabolite pathways associated with therapeutic properties. Moreover, gene editing can combine multiple medicinal properties from various plants, culminating in a singular 'superplant' with enhanced therapeutic benefits. This review discusses the current usage of medicinal plants in promoting human health during space travel, the application of gene editing technology in modifying medicinal plants and presents a vision for the utilization of gene-edited medicinal plants in space to mitigate the risk of various space-related health issues.

Keywords: Medicinal plants; long-duration space flight; gene-editing; radiation-induced cardiovascular disease; gut microbiome; microgravity

Introduction

Space travel presents an exciting opportunity to investigate celestial bodies and enhance our understanding of the cosmos as we know it. Alternatively, exploring outer space could become imperative if resources on Earth are depleted or a catastrophe occurs.

Researchers have spent decades exploring the potential of space exploration, but future missions will require even longer journeys through space. Humanity has made significant strides in space travel, from landing humans on the Moon and launching probes to explore distant planets. However, technological advancements and international collaborations have motivated us to pursue more ambitious space exploration goals. Space missions are constrained on time, storage, water, and food prep, even more so during future missions to the Moon and Mars, which will require extended stays for over twelve months. Space travel is one of the most challenging situations for the human body to endure. The adverse physiological effects of extended space travel are highly variable and continue to be an obstacle in carrying out missions (Bhuyan *et al.*, 2023). There is a plethora of health issues involved, including microgravity, bone density loss, impaired gut microbiota, and central fluid shift (Simanonok and Charles, 1994; Turrone *et al.*, 2020; Bhuyan *et al.*, 2023)

Longer space travel invariably leads to rising health concerns that can be addressed using medications. However, storing copious amounts of pills and vitamins for unpredictable health concerns is not practical due to limited storage space. Additionally, for extended space travel that may last many years, medications are likely to expire, and resupplying may not be possible (Wotring, 2015). A new system is urgently needed to address these issues.

Long-term stays in space require a sustainable life support mechanism that can provide fresh food, maintain a clean atmospheric balance, and regenerate resources - all of which can be supplied by plants (Wolff *et al.*, 2014). Additionally, plants can be used for medicinal purposes.

This review summarizes the potential applications of medicinal plants in addressing common space travel ailments, including microgravity, bone density loss, impaired gut microbiota, and central fluid shift. Additionally, I propose the development of a medicinal 'superplant' - a multipurpose medicinal plant capable of alleviating multiple symptoms associated with space travel.

Medicinal Plants

A medicinal plant contains beneficial nutrients which may serve as precursors to drugs or have therapeutic properties (Sofowora, Ogunbodede and Onayade, 2013). Several conditions must be met to accurately classify a medicinal plant. These conditions include its traditional use in formulations like herbal remedies, the extraction of its active compounds for medicinal purposes, its culinary uses as a spice, its role as a fragrant plant with medicinal benefits (e.g., ginger), and its utilization in the production of materials for medical dressings, such as cotton, jute, and flax (Sofowora, Ogunbodede and Onayade, 2013). In this article, we primarily focus on medicinal plants in which extracted substances are used for medicinal purposes.

Medicinal plants have been around for years, and various methods have already been used to cultivate and enhance their medicinal properties.

Indigenous knowledge

The curation of medicinal plants began through indigenous practices, which contributed to a variety of modern drugs (Gyllenhaal *et al.*, 2012). Rural communities have long used indigenous knowledge to guide their healthcare, energy, and natural resource practices. In some African tribes, medicinal plants have long been conserved by enforcing certain 'taboos', which restricted the harvesting of plants deemed to have medicinal qualities. Conditions included seasonality, timeframe, and cutting in specific areas (Kambizi and Bvenura, 2023). Ethnobotanical expertise is obtained by personal experience or through oral history over generations. (Aziz *et al.*, 2018). The preparation of medicinal plants varies in cultures and regions of the world. Additionally, different parts of the plant are used to treat different ailments. For instance, *Brassica Rapa* seeds are used for seasonal allergies, the bulb of *Allium cepa* is used to treat digestive issues, and the leaves of *Calotropis procera* alleviate joint pain (Aziz *et al.*, 2018). Various techniques are employed to extract the medicinal qualities of the plant parts. In a study conducted in Woreda Wukro klete Awlalo located in northern Ethiopia, common preparation methods of medicinal plants include, grinding, crushing, chewing, squeezing, and drinking. Grinding was the most common method of preparation (Mesfin, Tekle and Tesfay, 2013). Furthermore, Vaidyas in Uttaranchal, India typically crushes the medicinal herb into a powder and incorporate it with other ingredients through either cooking, dissolving in liquid, or forming pills for regular supplementation (Kala, Farooquee and Majila, 2005, p. 200).

Traditional breeding

Breeding medicinal plants is far more time-intensive than breeding crops. Medicinal plants are more prone to inhibit growth in response to various environmental stresses. Secondary metabolites aid plants in overcoming environmental stresses, and medicinal plants release secondary metabolites such as alkaloids and flavonoids, marking an important distinction between medicinal and crop plant breeding (Wang *et al.*, 2020). The initial domestication of plants was centered around staple crop plants such as wheat, rice, potato, and maize, and neglecting medicinal plants in the process (Niazian, 2019a). Medicinal plants are naturally highly variable and is one of the reasons traditional breeding methods were used until now, in addition to the low cost and return on investments [Carlen 2011].

Gene editing

The application of gene editing on medicinal plants is relatively new.

CRISPR-Cas technology has been applied to a few plants notably *Salvia miltiorrhiza* and *Cannabis sativa* to produce knockouts and highlight key genes that produce medicinal compounds (Zhou *et al.*, 2018; Guo *et al.*, 2022).

TILLING (Targeting-Induced Local Lesions IN Genomes) is a reverse genetic screening technique used to identify mutant genotypes, and while it has not been employed on medicinal plants specifically, it could be used to optimize detection of genes associated with medicinal properties (Niazian, 2019a).

TALENS (transcription activator-like effector nucleases) are engineered nucleases that induce targeted DNA modifications and achieve more than 10% DNA modification efficiency (Khan *et al.*, 2017; Niazi, 2019b; Niazi and Niedbała, 2020). Many crop plants have used TALENS for yield and quality improvement (Khan *et al.*, 2017).

Transgenes

Medicinal plant transgenes need to be able to withstand its important medicinal compounds without damaging them. Most methods for developing medicinal plant transgenes involve agrobacterium-mediated transformation (Trivedi *et al.*, 2016). In one such study, transgenic *Gentiana macrophylla* roots showed over 30-fold increased growth and produced a higher number of alkaloids for pharmaceutical use because of the larger root biomass in the T-DNA of agropine type of Ri plasmids (Tiwari *et al.*, 2007). Thus, the incorporation of transgenes can enhance the overall ability for a medicinal plant to generate therapeutic compounds.

Machine Learning-Assisted Methods

In medicinal plant breeding, machine learning can be used to classify molecular markers, which enable the study of genetic diversity. Convolutional neural networks (CNNs) can be trained to read DNA/RNA sequences and classify genotypes using molecular markers (Niazi and Niedbała, 2020). In essence, machine learning and deep learning techniques can be used as phenotype prediction tools using genome data, allowing for more efficient medicinal plant breeding outcomes.

Microgravity

Sustained microgravity for long periods of time could lead to severe physiological impairments in the human body. Most notably, a microgravity-induced bone loss trend suggests that more bone density and calcium is lost in longer missions. During spaceflight, microgravity conditions “unload” weight bearing bones, triggering increased bone mineral loss leading to secondary osteoporosis (Grimm *et al.*, 2016). Furthermore, calcium deficiency is common among astronauts, with nearly 50% decrease in calcium intake and absorption during a 3-month long term spaceflight (Smith *et al.*, 1999). Space travel results in increased bone density in the skull area, while causing significant bone depletion in the upper limbs, lumbar spine, thorax, and lower extremities (Stavnichuk *et al.*, 2020).

Microgravity conditions also impair cytoskeletal function. Human multipotent stem cells (hMSCs), which are found in adult bone marrow, are important for cytoskeleton integrity. In microgravity conditions, osteoblastic expression in hMSCs are inhibited, reducing osteoblastic differentiation and disrupting the cytoskeleton (Zayzafoon, Meyers and McDonald, 2005). A similar pattern has been observed in endothelial and glial cells upon exposure to space flight, suggesting that impaired cytoskeletons may contribute to cardiovascular, musculoskeletal, and immune system compromise (Uva *et al.*, 2002; Carlsson *et al.*, 2003; Zayzafoon, Meyers and McDonald, 2005).

Though there are no definite treatments to reverse bone osteoporosis, there are some management techniques. Firstly, exercise leads to the decrease of bone mineral loss, resulting in positive effects of bone loss. This has been determined in several animal models, but current human trials have shown exercise only provides a partial protection against bone mineral loss and requires supplementation with an anti-resorptive (Sibonga *et al.*, 2019). Pharmacologically, protein, calcium, vitamin D, and high intake of alkaline precursors are shown to reduce bone loss (Hajisadeghi *et al.*, 2021; Fischer *et al.*, 2023).

There have been no successful therapeutic strategies that have prevented osteocyte apoptosis, and future exploration may entail using a CRISPR-Cas 9 knockout for Cx43 or PGE2 biomarkers in apoptotic signaling (Ru and Wang, 2020, p. 20). Microgravity causes unloading and disuse of the bones speculates the dysfunction of Src/MEK/ERK, autophagy, Wnt/Beta-catenin, AMPK-PGC1A-ROS axis, NO-sGc-cGMP, PTHrP/PTH1R; VEGF/VEGFR2 pathway, though it is not clear what is the direct cause of osteocyte apoptosis (Ru and Wang, 2020).

Green tea polyphenol supplementation with drinking water has been shown to increase protection of bone mass by reducing inflammation, and downregulating COX-2 and TNF- α could reduce the risk of osteoporosis (Shen *et al.*, 2010).

Future studies should decipher the molecular mechanism by which osteocyte apoptosis occurs in microgravity conditions using animal models and gene editing of pathways.

Gut Microbiota

The gut microbiome plays a key role in immune functions and homeostasis. Impaired balance of microbiota could lead to a myriad of health concerns, including metabolic disorders, increased susceptibility to infectious pathogens, diseases of gut and body organs, and even cancer (Mosca, Leclerc and Hugot, 2016; Sommariva *et al.*, 2020; Shama *et al.*, 2024). Hind-limbed unloaded mice are well-established terrestrial models to simulate microgravity effects, mimicking various physiological changes akin to space travel (Shama *et al.*, 2022). In a hind-limbed unloaded mice model, microgravity conditions were mimicked over a three-week period; the study reported a decrease in intestinal microbial diversity and increase in pathogenic bacteria (Shama *et al.*, 2022). As established before, space-induced microgravity environments promote bone density loss and weakened muscles. Taking probiotics has been associated with regaining strength and muscle mass, suggesting that maintaining a healthy gut microbiome leads to improved overall health and maintained muscle strength - combatting common space travel ailments (Giron *et al.*, 2022).

Ceratonia Siliqua, or Carob, extract contains pectin, gums, and polyphenols and has been shown to flourish gut microbiome with its anti-inflammatory properties (Requena *et al.*, 2010; Micheletti *et al.*, 2023). Polyphenols are secondary metabolites of plants and act as antioxidants, neutralizing reactive oxygen species and improving gut health (Zhang and Tsao, 2016). Epigallocatechin-3-gallate (EGCG) in green tea has been shown to stimulate production of *Bifidobacterium adolescentis* when metabolized, creating a healthier gut bacteria profile and promoting human health (Liu *et al.*, 2020).

Cardiovascular Disease and Cosmic Radiation

In recent years, radiation-induced cardiovascular disease (RICVD) has been a rising concern for those pursuing long-distance space travel. Radiation can interact with genetics and lifestyle factors, potentially exacerbating the risk of RICVD during long-distance space travel (Huff *et al.*, 2022). Compared to Earth, exposure to high energy protons due to solar particle events and galactic cosmic rays are much higher in space. This leads to concentrated exposure to radiation particles which damage DNA, RNA, and proteins, which put astronauts at a higher risk for RICVD (Moeller *et al.*, 2017; Huff *et al.*, 2022). The use of antioxidants against radiation-induced tissue damage has long been studied, particularly the use of the Rosmarinic Acid antioxidant has been shown to protect the heart and decrease receive oxidative species by stimulating antioxidant enzymes and protect the heart (Goudarzi, Fatahi Asl and Shoghi, 2023). Though this was only studied in rats, human models are still yet to be studied to elucidate the effects of antioxidants in humans.

Supplementing antioxidants in a diet could enhance protection against cosmic radiation in space. One study described how the use of a 2% strawberry extract diet could significantly protect against heavy ion particles common in cosmic radiation than a 2% blueberry extract diet (Rabin, Joseph and Shukitt-Hale, 2005). Additionally, a single exposure to UV radiation stimulates the release of ROS. Topical application of green tea polyphenol extract epigallocatechin-3-gallate before exposure has been shown to inhibit hydrogen peroxide and nitric oxide production associated with ROS and reverses depletion of antioxidant enzymes such as catalase (Katiyar *et al.*, 2001).

However, the exact treatment and dosage of these remedies against cosmic radiation is highly variable and should be personalized to each astronaut upon further clinical study (Mitreia *et al.*, 2018). Finally, ferulic acid is typically found in wheat, oats, and rice (Kumar and Pruthi, 2014). Ferulic acid also preserves antioxidant enzymes and inhibits apoptotic activity in the spleen if administered before radiation exposure by reducing TBARS formation and enhancing catalase activity (Das *et al.*, 2016).

Additionally, cardiac arrhythmias are common among long duration spaceflight astronauts, and it has been shown that mitochondrial-associated antioxidants such as resveratrol activate the PI3K/AKT/eNOS pathway and reduce the occurrence of cardiac arrhythmias (Chong *et al.*, 2015; Joseph *et al.*, 2016).

Future Directions

Given the harsh environment and storage constraints of space travel, introducing a medicinal 'superplant' would be an ideal way to improve astronaut health. We introduce the novel idea of a 'superplant,' which would be genetically modified to contain multiple medicinal properties for different space-related ailments in one plant. This novel approach to space health provides a sustainable, regenerative supply of medicinal supplements to promote a more natural approach to controlling space-induced ailments than traditional chemical pharmaceuticals. Rather than relying on reactive treatments, continuous intake of these medicinal substances could serve as a key preventative measure for numerous diseases, especially during long-duration space flights.

In the previous discussion of space-related ailments, a common agent that caused microgravity-induced osteoporosis, impaired gut microbiota, and cardiovascular disease from cosmic radiation was the activation of ROS. In the human body, the Nrf2-ARE pathway regulates the detoxification process using antioxidants and rids of ROS (Nguyen, Nioi and Pickett, 2009). Thus, the primary focus of the 'superplant' is to contain high amounts of ROS antagonists which upregulate the NRF2-KEAP1 pathway in the body. NRF2 is a key gene involved in activating cytoprotective genes against ROS species, and KEAP1 is a key negative regulator of NRF2, degrading NRF2 in unstressed conditions and stabilizing it during defense responses (Yamamoto, Kensler and Motohashi, 2018). However, NRF2 is stabilized upon KEAP1 degradation (Wasik *et al.*, 2017). Additionally, the medicinal plant should over-express secondary metabolites that are able to facilitate nuclear translocation of NRF2 to upregulate the cytoprotective genes that provide the beneficial effect. Green tea catechins are one class of polyphenols that can induce the NRF2-KEAP1 defense mechanism via nuclear translocation of NRF2, transcribing cytoprotective genes such as glutathione S-transferases (GSTs) and glutathione reductases (GRs), to name a few (Yamamoto, Kensler and Motohashi, 2018; Talebi *et al.*, 2021; Liu *et al.*, 2022).

Additionally, phenolic acids like ferulic acid and chlorogenic acid, commonly present in fruits, along with flavonoids such as catechins and epicatechins, are notable for their potent antioxidant activity. These compounds have the capability to translocate NRF2 and enhance the expression of antioxidant enzymes like heme oxygenase-1 (HO-1) and catalase (CAT) (Kumar and Goel, 2019; Qader *et al.*, 2020). There is a rising interest in identifying botanical NRF2 activators due to its extreme promise in treatment of several human diseases.

Because each medicinal plant has unique compounds that target specific genes, employing a medicinal 'superplant' would be an ideal solution for space travel constraints. In order to pursue this, gene editing technology should be applied to consolidate multiple different medicinal plant properties into one 'superplant.'

However, before gene editing technology can be applied, the medicinal plant needs to be fully sequenced. Currently, medicinal plant genome sequencing is severely lacking compared to crop plants and organisms with mostly fully sequenced genomes, preventing many pathways and genes that control secondary metabolite production from being identified (Alami *et al.*, 2022).

The use of bioinformatics coupled with functional genomics approaches could radically speed up advances in medicinal plant sequencing and gene targeting to further elucidate secondary metabolite biosynthesis pathways. Modern plant metabolomics is a newly emerging tool used to uncover biosynthetic pathways (Sumner *et al.*, 2015). Using such screening tools, it would be ideal to identify key metabolic pathways and genes involved in secondary metabolite production and allow for a targeted gene editing approach. TILLING (Targeting-Induced Local Lesions IN Genomes) is a reverse genetic screening technique used to identify mutant genotypes, and while it has not been employed on medicinal plants specifically, it could be used to optimize detection of genes associated with medicinal properties (Niazian, 2019a).

The approach to engineer a medicinal 'superplant' suitable for space conditions is to employ gene editing and breeding tools to increase secondary metabolite production in the plant. Tandem duplication (TD) and whole-genome duplication (WGD) could be used to enhance secondary metabolite and antioxidant function and TD has been demonstrated successfully in poppy and tea plants (Alami *et al.*, 2022).

Cis-regulatory sequences such as transcriptional enhancers are able to increase transcription rate and expression rate with transcriptional factors (Schmitz, Grotewold and Stam, 2021). In Pea plants, the PetE enhancer upregulates genes of interest through induced histone acetylation. This hyperacetylation increases transcription rates by recruiting more transcription activators (Chua, Watson and Gray, 2003).

CRISPR-Cas 9 systems could be employed to target gene activation and improve nutritional content using sgRNA and Cas9, which creates a double stranded break whereby natural reannealing would induce mutations and gene knockouts (Sathee *et al.*, 2022). This approach could be used to downregulate the ROS signaling pathway. In rice, OsSRT1 overexpression leads to increased stress tolerance and decreased ROS expression (Huang *et al.*, 2019). Similarly, by identifying key ROS regulators in medicinal plants, appropriate gene regulation would allow the 'superplant' to better tolerate space environment stress.

Biofortification is the process of enhancing the nutritional quality of food crops by increasing the content of vitamins, minerals, and amino acids. Biofortification could be employed to increase the plant's nutritional quality without compromising its crucial characteristics. Selenium is a crucial trace element in boosting the immune system, and its deficiency is known to cause weakened immune systems and even cancer (Brown and Arthur, 2001). Deficiency is largely attributed to the lack of Selenium in plant soil and consequently in the plant itself. Attempts to restore selenium concentration in plants has been attempted with biofortification. This has been tested on a medicinal plant *Plantago ovata* with selenium (Se) biofortification, and results demonstrated an increase of production of polyphenols and overall antioxidant content (Dey and Raychaudhuri, 2023). Similar results have been shown in other medicinal plants such as *Plantago asiatica* and Iranian Borage (Hosseinzadeh Rostam Kalaei, Abdossi and Danaee, 2022; Liao and Zhu, 2022). This is largely explained due to the increase of ROS under Se-stress signaling. To date, biofortification has not been tested on medicinal plants under space conditions such as microgravity and extended radiation exposure. If testing Se-biofortification, it should be noted how space-induced stressors and genetic mutations would interact with Se-stress signaling, and if this combination may change expression levels of polyphenols.

Plants with existing rich levels of antioxidants may not require gene editing for metabolite production but may need modification to support stressors in space. Thus, the second approach would be to edit the plant to better withstand stressors and create space environments that enable the growth of the plants while maintaining its nutrient content.

There are several concerns of growing plants in space, and there is much still unknown. Unique concerns to growing plants in space is exposure to ionizing radiation, which could alter gene expression, and microgravity, which severely alters plant physiology (Vandenbrink and Kiss, 2016; De Micco *et al.*, 2023). A study found that secondary metabolite production of Cannabis was enhanced in a simulated microgravity setting (Darigh *et al.*, 2022). High temperature stress and concentrated ultraviolet-B light environments stimulate an increase of secondary metabolites and phenols (Naghiloo *et al.*, 2012; Li *et al.*, 2020). Further testing of medicinal plants rich in antioxidants such as green tea containing epigallocatechin-3-gallate, should be tested in a microgravity environment to corroborate the finding of enhanced secondary metabolite function.

Additionally, very few plants are being grown in space, only leafy vegetables, and little to no medicinal plants have been tested. A third approach would be to gene-edit plants that have already been grown in space to express more polyphenols and antioxidants. Overexpression of transcription factors in tomato plants have been used to increase expression of anthocyanin, a pigment found to be a protective agent against human diseases (Butelli *et al.*, 2008; Nguyen *et al.*, 2023, p. 20). Strawberries, an antioxidant rich fruit, have been cultivated in space with great potential for survival in a harsh space environment. As mentioned earlier, a strawberry extract supplement diet aids in protection

against cosmic radiation and associated heavy ion buildup. Overexpression of FaMAPK5 and FaMAPK10 in strawberries leads to increased anthocyanin enzyme activity (Zhang *et al.*, 2021; Zhang and Zhang, 2022). Utilizing such a modification would be essential for space travel, especially in situations of limited cultivation opportunities where each fruit should be optimized for nutrition.

Conclusion

Human space travel is a complicated venture, especially so for long-duration space flights. Given storage constraints and extended time periods, the use of chemical medications may not be the most effective tool to manage space travel ailments. In this review, we have outlined the various types of space ailments such as microgravity, bone-density loss, impaired gut bacteria biome, and cardiovascular health and possible use of medicinal plant treatment for each. We presented the possibility of developing a medicinal 'superplant' using gene-editing mechanisms, creating a viable medicinal supplement suitable for space travel. To our knowledge, there has been no research study to date that introduces this possibility.

Acknowledgements: The author thanks Dr. Christy Gault for invaluable guidance and support in editing and mentorship.

References

- Alami, M.M. *et al.* (2022) 'The Current Developments in Medicinal Plant Genomics Enabled the Diversification of Secondary Metabolites' Biosynthesis', *International Journal of Molecular Sciences*, 23(24), p. 15932. Available at: <https://doi.org/10.3390/ijms232415932>.
- Aziz, M.A. *et al.* (2018) 'Traditional uses of medicinal plants used by Indigenous communities for veterinary practices at Bajaur Agency, Pakistan', *Journal of Ethnobiology and Ethnomedicine*, 14, p. 11. Available at: <https://doi.org/10.1186/s13002-018-0212-0>.
- Bhuyan, N. *et al.* (2023) 'Prolonged space flight: Adverse health effects and treatment options with medicinal plants and natural products', *Sciences of Phytochemistry*, 2(1), pp. 82–97. Available at: <https://doi.org/10.58920/sciphy02010082>.
- Brown, K.M. and Arthur, J.R. (2001) 'Selenium, selenoproteins and human health: a review', *Public Health Nutrition*, 4(2B), pp. 593–599. Available at: <https://doi.org/10.1079/phn2001143>.
- Butelli, E. *et al.* (2008) 'Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors', *Nature Biotechnology*, 26(11), pp. 1301–1308. Available at: <https://doi.org/10.1038/nbt.1506>.
- Carlsson, S.I.M. *et al.* (2003) 'Endothelial stress by gravitational unloading: effects on cell growth and cytoskeletal organization', *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1642(3), pp. 173–179. Available at: <https://doi.org/10.1016/j.bbamcr.2003.08.003>.
- Chong, E. *et al.* (2015) 'Resveratrol, a red wine antioxidant, reduces atrial fibrillation susceptibility in the failing heart by PI3K/AKT/eNOS signaling pathway activation', *Heart Rhythm*, 12(5), pp. 1046–1056. Available at: <https://doi.org/10.1016/j.hrthm.2015.01.044>.
- Chua, Y.L., Watson, L.A. and Gray, J.C. (2003) 'The Transcriptional Enhancer of the Pea Plastocyanin Gene Associates with the Nuclear Matrix and Regulates Gene Expression through Histone Acetylation', *The Plant Cell*, 15(6), pp. 1468–1479. Available at: <https://doi.org/10.1105/tpc.011825>.
- Darigh, F. *et al.* (2022) 'Simulated microgravity contributed to modification of callogenesis performance and secondary metabolite production in *Cannabis Indica*', *Plant Physiology and Biochemistry*, 186, pp. 157–168. Available at: <https://doi.org/10.1016/j.plaphy.2022.07.012>.
- Das, U. *et al.* (2016) 'Ferulic acid (FA) abrogates ionizing radiation-induced oxidative damage in murine spleen', *International Journal of Radiation Biology*, 92(12), pp. 806–818. Available at: <https://doi.org/10.1080/09553002.2016.1230241>.
- De Micco, V. *et al.* (2023) 'Perspectives for plant biology in space and analogue environments', *npj Microgravity*, 9(1), pp. 1–10. Available at: <https://doi.org/10.1038/s41526-023-00315-x>.
- Dey, S. and Raychaudhuri, S.S. (2023) 'Selenium biofortification improves bioactive composition and antioxidant status in *Plantago ovata* Forsk., a medicinal plant', *Genes and Environment*, 45(1), p. 38. Available at: <https://doi.org/10.1186/s41021-023-00293-2>.
- Fischer, C. *et al.* (2023) 'Additive Effects of Exercise and Vitamin D Supplementation (with and without Calcium) on Bone Mineral Density in Older Adults: A Systematic Review and Meta-Analysis', *Journal of Osteoporosis*, 2023, p. 5570030. Available at: <https://doi.org/10.1155/2023/5570030>.
- Giron, M. *et al.* (2022) 'Gut microbes and muscle function: can probiotics make our muscles stronger?', *Journal of Cachexia, Sarcopenia and Muscle*, 13(3), pp. 1460–1476. Available at: <https://doi.org/10.1002/jcsm.12964>.

- Goudarzi, M., Fatahi Asl, J. and Shoghi, H. (2023) 'Comparison of the Effects of Rosmarinic Acid and Electromagnetic Radiation-Induced Cardiotoxicity on Rats', *The Journal of Tehran University Heart Center*, 18(3), pp. 207–213. Available at: <https://doi.org/10.18502/jthc.v18i3.14115>.
- Grimm, D. *et al.* (2016) 'The impact of microgravity on bone in humans', *Bone*, 87, pp. 44–56. Available at: <https://doi.org/10.1016/j.bone.2015.12.057>.
- Gyllenhaal, C. *et al.* (2012) 'Ethnobotanical approach versus random approach in the search for new bioactive compounds: Support of a hypothesis', *Pharmaceutical biology*, 50(1), pp. 30–41. Available at: <https://doi.org/10.3109/13880209.2011.634424>.
- Hajisadeghi, H. *et al.* (2021) 'Effect of regular resistance exercise, vitamin D, and calcium supplements on the bone mineral content and density in postmenopausal model of rats: An experimental study', *International Journal of Reproductive Biomedicine*, 19(1), pp. 63–74. Available at: <https://doi.org/10.18502/ijrm.v19i1.8181>.
- Hosseinzadeh Rostam Kalaei, M., Abdossi, V. and Danaee, E. (2022) 'Evaluation of foliar application of selenium and flowering stages on selected properties of Iranian Borage as a medicinal plant', *Scientific Reports*, 12(1), p. 12568. Available at: <https://doi.org/10.1038/s41598-022-16241-z>.
- Huang, H. *et al.* (2019) 'Mechanisms of ROS Regulation of Plant Development and Stress Responses', *Frontiers in Plant Science*, 10. Available at: <https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2019.00800> (Accessed: 20 February 2024).
- Huff, J.L. *et al.* (2022) 'Cardiovascular Disease Risk Modeling for Astronauts: Making the Leap From Earth to Space', *Frontiers in Cardiovascular Medicine*, 9, p. 873597. Available at: <https://doi.org/10.3389/fcvm.2022.873597>.
- Joseph, L.C. *et al.* (2016) 'Mitochondrial oxidative stress during cardiac lipid overload causes intracellular calcium leak and arrhythmia', *Heart Rhythm*, 13(8), pp. 1699–1706. Available at: <https://doi.org/10.1016/j.hrthm.2016.05.002>.
- Kala, C.P., Farooquee, N.A. and Majila, B.S. (2005) '**Indigenous knowledge and medicinal plants used by Vaidyas in Uttaranchal, India**', *NPR Vol.4(3) [May-June 2005]* [Preprint]. Available at: <http://nopr.niscpr.res.in/handle/123456789/8090> (Accessed: 20 February 2024).
- Kambizi, L. and Bvenura, C. (2023) *Sustainable Uses and Prospects of Medicinal Plants*. CRC Press.
- Katiyar, S.K. *et al.* (2001) 'Green tea polyphenol (–)-epigallocatechin-3-gallate treatment of human skin inhibits ultraviolet radiation-induced oxidative stress', *Carcinogenesis*, 22(2), pp. 287–294. Available at: <https://doi.org/10.1093/carcin/22.2.287>.
- Khan, Z. *et al.* (2017) 'Use of TALEs and TALEN Technology for Genetic Improvement of Plants', *Plant Molecular Biology Reporter*, 35(1), pp. 1–19. Available at: <https://doi.org/10.1007/s11105-016-0997-8>.
- Kumar, N. and Goel, N. (2019) 'Phenolic acids: Natural versatile molecules with promising therapeutic applications', *Biotechnology Reports*, 24, p. e00370. Available at: <https://doi.org/10.1016/j.btre.2019.e00370>.
- Kumar, N. and Pruthi, V. (2014) 'Potential applications of ferulic acid from natural sources', *Biotechnology Reports*, 4, pp. 86–93. Available at: <https://doi.org/10.1016/j.btre.2014.09.002>.
- Li, Y. *et al.* (2020) 'The effect of developmental and environmental factors on secondary metabolites in medicinal plants', *Plant Physiology and Biochemistry*, 148, pp. 80–89. Available at: <https://doi.org/10.1016/j.plaphy.2020.01.006>.
- Liao, R. and Zhu, J. (2022) 'Amino acid promotes selenium uptake in medicinal plant *Plantago asiatica*', *Physiology and Molecular Biology of Plants*, 28(5), pp. 1005–1012. Available at: <https://doi.org/10.1007/s12298-022-01196-2>.
- Liu, C. *et al.* (2022) 'The Gut Microbial Metabolite Pyrogallol Is a More Potent Inducer of Nrf2-Associated Gene Expression Than Its Parent Compound Green Tea (–)-Epigallocatechin Gallate', *Nutrients*, 14(16), p. 3392. Available at: <https://doi.org/10.3390/nu14163392>.
- Liu, Z. *et al.* (2020) 'Reciprocal Interactions between Epigallocatechin-3-gallate (EGCG) and Human Gut Microbiota In Vitro', *Journal of Agricultural and Food Chemistry*, 68(36), pp. 9804–9815. Available at: <https://doi.org/10.1021/acs.jafc.0c03587>.
- Mesfin, K., Tekle, G. and Tesfay, T. (2013) '*Journal of Medicinal Plants Studies*', 1(4).
- Micheletti, C. *et al.* (2023) 'Effects of Carob Extract on the Intestinal Microbiome and Glucose Metabolism: A Systematic Review and Meta-Analysis', *La Clinica Terapeutica*, 174(Suppl 2(6)), pp. 169–172. Available at: <https://doi.org/10.7417/CT.2023.2484>.
- Mitrea, D.R. *et al.* (2018) 'Antioxidant protection against cosmic radiation-induced oxidative stress at commercial flight altitude', *Journal of Physiology and Pharmacology: An Official Journal of the Polish Physiological Society*, 69(4). Available at: <https://doi.org/10.26402/jpp.2018.4.03>.
- Moeller, R. *et al.* (2017) 'STARLIFE-An International Campaign to Study the Role of Galactic Cosmic Radiation in Astrobiological Model Systems', *Astrobiology*, 17(2), pp. 101–109. Available at: <https://doi.org/10.1089/ast.2016.1571>.
- Mosca, A., Leclerc, M. and Hugot, J.P. (2016) 'Gut Microbiota Diversity and Human Diseases: Should We Reintroduce Key Predators in Our Ecosystem?', *Frontiers in Microbiology*, 7. Available at: <https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2016.00455> (Accessed: 20 February 2024).

- Naghiloo, S. *et al.* (2012) 'Ontogenetic Variation of Total Phenolics and Antioxidant Activity in Roots, Leaves and Flowers of *Astragalus compactus* Lam. (Fabaceae)', *BioImpacts: BI*, 2(2), pp. 105–109. Available at: <https://doi.org/10.5681/bi.2012.015>.
- Nguyen, M.T.P. *et al.* (2023) 'Space farming: Horticulture systems on spacecraft and outlook to planetary space exploration', *Plant Physiology and Biochemistry*, 194, pp. 708–721. Available at: <https://doi.org/10.1016/j.plaphy.2022.12.017>.
- Nguyen, T., Nioi, P. and Pickett, C.B. (2009) 'The Nrf2-Antioxidant Response Element Signaling Pathway and Its Activation by Oxidative Stress', *The Journal of Biological Chemistry*, 284(20), pp. 13291–13295. Available at: <https://doi.org/10.1074/jbc.R900010200>.
- Niazian, M. (2019a) 'Application of genetics and biotechnology for improving medicinal plants', *Planta*, 249(4), pp. 953–973. Available at: <https://doi.org/10.1007/s00425-019-03099-1>.
- Niazian, M. (2019b) 'Application of genetics and biotechnology for improving medicinal plants', *Planta*, 249(4), pp. 953–973. Available at: <https://doi.org/10.1007/s00425-019-03099-1>.
- Niazian, M. and Niedbala, G. (2020) 'Machine Learning for Plant Breeding and Biotechnology', *Agriculture*, 10(10), p. 436. Available at: <https://doi.org/10.3390/agriculture10100436>.
- Qader, M. *et al.* (2020) 'Natural Nrf2 Activators from Juices, Wines, Coffee, and Cocoa', *Beverages*, 6(4), p. 68. Available at: <https://doi.org/10.3390/beverages6040068>.
- Rabin, B.M., Joseph, J.A. and Shukitt-Hale, B. (2005) 'Effects of age and diet on the heavy particle-induced disruption of operant responding produced by a ground-based model for exposure to cosmic rays', *Brain Research*, 1036(1–2), pp. 122–129. Available at: <https://doi.org/10.1016/j.brainres.2004.12.041>.
- Requena, T. *et al.* (2010) 'Perspectives of the potential implications of wine polyphenols on human oral and gut microbiota', *Trends in Food Science & Technology*, 21(7), pp. 332–344. Available at: <https://doi.org/10.1016/j.tifs.2010.04.004>.
- Ru, J. and Wang, Y. (2020) 'Osteocyte apoptosis: the roles and key molecular mechanisms in resorption-related bone diseases', *Cell Death & Disease*, 11(10), pp. 1–24. Available at: <https://doi.org/10.1038/s41419-020-03059-8>.
- Sathee, L. *et al.* (2022) 'Genome Editing Targets for Improving Nutrient Use Efficiency and Nutrient Stress Adaptation', *Frontiers in Genetics*, 13. Available at: <https://www.frontiersin.org/journals/genetics/articles/10.3389/fgene.2022.900897> (Accessed: 20 February 2024).
- Schmitz, R.J., Grotewold, E. and Stam, M. (2021) 'Cis-regulatory sequences in plants: Their importance, discovery, and future challenges', *The Plant Cell*, 34(2), pp. 718–741. Available at: <https://doi.org/10.1093/plcell/koab281>.
- Shama, S. *et al.* (2022) 'The Role of 4-Phenylbutyric Acid in Gut Microbial Dysbiosis in a Mouse Model of Simulated Microgravity', *Life*, 12(9), p. 1301. Available at: <https://doi.org/10.3390/life12091301>.
- Shama, S. *et al.* (2024) 'Enhancing microbial diversity as well as multi-organ health in hind-limb unloaded mice', *Life Sciences in Space Research*, 40, pp. 62–71. Available at: <https://doi.org/10.1016/j.lssr.2023.08.006>.
- Shen, C.-L. *et al.* (2010) 'Green tea polyphenols mitigate bone loss of female rats in a chronic inflammation-induced bone loss model', *The Journal of Nutritional Biochemistry*, 21(10), pp. 968–974. Available at: <https://doi.org/10.1016/j.jnutbio.2009.08.002>.
- Sibonga, J. *et al.* (2019) 'Resistive exercise in astronauts on prolonged spaceflights provides partial protection against spaceflight-induced bone loss', *Bone*, 128, p. 112037. Available at: <https://doi.org/10.1016/j.bone.2019.07.013>.
- Simanonok, K.E. and Charles, J.B. (1994) 'Space sickness and fluid shifts: a hypothesis', *Journal of Clinical Pharmacology*, 34(6), pp. 652–663. Available at: <https://doi.org/10.1002/j.1552-4604.1994.tb02020.x>.
- Smith, S.M. *et al.* (1999) 'Calcium metabolism before, during, and after a 3-mo spaceflight: kinetic and biochemical changes', *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 277(1), pp. R1–R10. Available at: <https://doi.org/10.1152/ajpregu.1999.277.1.R1>.
- Sofowora, A., Ogunbodede, E. and Onayade, A. (2013) 'The role and place of medicinal plants in the strategies for disease prevention', *African Journal of Traditional, Complementary and Alternative Medicines*, 10(5), pp. 210–229. Available at: <https://doi.org/10.4314/ajtcam.v10i5.2>.
- Sommariva, M. *et al.* (2020) 'The lung microbiota: role in maintaining pulmonary immune homeostasis and its implications in cancer development and therapy', *Cellular and molecular life sciences: CMLS*, 77(14), pp. 2739–2749. Available at: <https://doi.org/10.1007/s00018-020-03452-8>.
- Stavnichuk, M. *et al.* (2020) 'A systematic review and meta-analysis of bone loss in space travelers', *npj Microgravity*, 6(1), pp. 1–9. Available at: <https://doi.org/10.1038/s41526-020-0103-2>.
- Sumner, L.W. *et al.* (2015) 'Modern plant metabolomics: advanced natural product gene discoveries, improved technologies, and future prospects', *Natural Product Reports*, 32(2), pp. 212–229. Available at: <https://doi.org/10.1039/c4np00072b>.
- Talebi, Marjan *et al.* (2021) 'New insights into the role of the Nrf2 signaling pathway in green tea catechin applications', *Phytotherapy research: PTR*, 35(6), pp. 3078–3112. Available at: <https://doi.org/10.1002/ptr.7033>.

- Tiwari, R.K. *et al.* (2007) 'Genetic transformation of *Gentiana macrophylla* with *Agrobacterium rhizogenes*: growth and production of secoiridoid glucoside gentiopicroside in transformed hairy root cultures', *Plant Cell Reports*, 26(2), pp. 199–210. Available at: <https://doi.org/10.1007/s00299-006-0236-0>.
- Trivedi, M. *et al.* (2016) 'Chapter 23 - GMO and Food Security', in Omkar (ed.) *Ecofriendly Pest Management for Food Security*. San Diego: Academic Press, pp. 703–726. Available at: <https://doi.org/10.1016/B978-0-12-803265-7.00023-3>.
- Turroni, S. *et al.* (2020) 'Gut Microbiome and Space Travelers' Health: State of the Art and Possible Pro/Prebiotic Strategies for Long-Term Space Missions', *Frontiers in Physiology*, 11. Available at: <https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2020.553929> (Accessed: 20 February 2024).
- Uva, B.M. *et al.* (2002) 'Clinorotation-induced weightlessness influences the cytoskeleton of glial cells in culture', *Brain Research*, 934(2), pp. 132–139. Available at: [https://doi.org/10.1016/S0006-8993\(02\)02415-0](https://doi.org/10.1016/S0006-8993(02)02415-0).
- Vandenbrink, J.P. and Kiss, J.Z. (2016) 'Space, the final frontier: A critical review of recent experiments performed in microgravity', *Plant Science: An International Journal of Experimental Plant Biology*, 243, pp. 115–119. Available at: <https://doi.org/10.1016/j.plantsci.2015.11.004>.
- Wang, W. *et al.* (2020) 'Advances and challenges in medicinal plant breeding', *Plant Science*, 298, p. 110573. Available at: <https://doi.org/10.1016/j.plantsci.2020.110573>.
- Wasik, U. *et al.* (2017) 'Protection against oxidative stress mediated by the Nrf2/Keap1 axis is impaired in Primary Biliary Cholangitis', *Scientific Reports*, 7(1), p. 44769. Available at: <https://doi.org/10.1038/srep44769>.
- Wolff, S.A. *et al.* (2014) 'Effects of the Extraterrestrial Environment on Plants: Recommendations for Future Space Experiments for the MELISSA Higher Plant Compartment', *Life*, 4(2), pp. 189–204. Available at: <https://doi.org/10.3390/life4020189>.
- Wotring, V.E. (2015) 'Chemical Potency and Degradation Products of Medications Stored Over 550 Earth Days at the International Space Station', *The AAPS Journal*, 18(1), pp. 210–216. Available at: <https://doi.org/10.1208/s12248-015-9834-5>.
- Yamamoto, M., Kensler, T.W. and Motohashi, H. (2018) 'The KEAP1-NRF2 System: a Thiol-Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis', *Physiological Reviews*, 98(3), p. 1169. Available at: <https://doi.org/10.1152/physrev.00023.2017>.
- Zayzafoon, M., Meyers, V.E. and McDonald, J.M. (2005) 'Microgravity: the immune response and bone', *Immunological Reviews*, 208(1), pp. 267–280. Available at: <https://doi.org/10.1111/j.0105-2896.2005.00330.x>.
- Zhang, H. and Tsao, R. (2016) 'Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects', *Current Opinion in Food Science*, 8, pp. 33–42. Available at: <https://doi.org/10.1016/j.cofs.2016.02.002>.
- Zhang, M. and Zhang, S. (2022) 'Mitogen-activated protein kinase cascades in plant signaling', *Journal of Integrative Plant Biology*, 64(2), pp. 301–341. Available at: <https://doi.org/10.1111/jipb.13215>.
- Zhang, Yunting *et al.* (2021) 'MAPK5 and MAPK10 overexpression influences strawberry fruit ripening, antioxidant capacity and resistance to *Botrytis cinerea*', *Planta*, 255(1), p. 19. Available at: <https://doi.org/10.1007/s00425-021-03804-z>.
- Zhou, Z. *et al.* (2018) 'CRISPR/Cas9-mediated efficient targeted mutagenesis of RAS in *Salvia miltiorrhiza*', *Phytochemistry*, 148, pp. 63–70. Available at: <https://doi.org/10.1016/j.phytochem.2018.01.015>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.