

Review

Not peer-reviewed version

---

# Lipid Oxidation Products and the Risk of Cardiovascular Diseases: Role of Lipoprotein Transport

---

[Markku Ahotupa](#)\*

Posted Date: 13 March 2024

doi: 10.20944/preprints202403.0781.v1

Keywords: Atherosclerosis; Cardiovascular diseases; Cholesterol; High-density lipoprotein; Lipid oxidation; Lipoprotein functions; Low-density lipoprotein; Risk factors



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

# Lipid Oxidation Products and the Risk of Cardiovascular Diseases: Role of Lipoprotein Transport

Markku Ahotupa <sup>1,2</sup>

<sup>1</sup> Centre for Population Health Research, University of Turku and Turku University Hospital, Turku, Finland.

<sup>2</sup> Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland.

**Abstract:** Cholesterol has ruled the history of atherosclerotic cardiovascular diseases (CVD) for decades, and the transport of cholesterol by lipoproteins has become a cornerstone of our view of etiology of the disease. Discovery of the lipoprotein-specific transport of lipid oxidation products (LOP) has opened a new and exciting area in research of CVD by revealing strong connections between atherogenic LOP, lipoprotein transport functions and CVD. This review goes through the evidence in support of the lipoprotein-specific LOP transport and speculates of possible consequences of the proposed role of lipoprotein LOP transport as a risk factor. Association of lipoprotein LOP with known risk factors of CVD are largely parallel with those of cholesterol, and due to the common transport and cellular intake mechanisms the elucidation of independent effects of either cholesterol or LOP is difficult. Yet, unlike cholesterol, LOP can directly initiate and boost atherogenic processes. It therefore appears inevitable that increasing knowledge of the lipoprotein-specific LOP transport will affect the present view of the role of cholesterol in aetiology and as a risk factor of CVD.

**Keywords:** atherosclerosis; cardiovascular diseases; cholesterol; high-density lipoprotein; lipid oxidation; lipoprotein functions; low-density lipoprotein; risk factors

---

## 1. Introduction

Plasma lipoproteins play a significant role in steering of lipid metabolism, both as the conveyors of lipophilic substances in blood, and as mediators of the receptor-mediated cellular uptake of lipids [1]. It is today well known that the two main lipoprotein classes, low-density lipoprotein (LDL) and high-density lipoprotein (HDL), transport plasma lipids to opposite directions. LDL is the main carrier of cholesterol to peripheral tissues, while the reverse transport by HDL returns excess tissue cholesterol to the liver. Likewise, it is also known that LDL and HDL have independent and opposing roles as risk factors for cardiovascular diseases (CVD). While LDL associates with elevated risk of CVD, high HDL concentration appears to be protective. The different roles of LDL and HDL as risk factors are explained by lipoprotein specificity of the transport and receptor binding functions, which have been worked out in studies over the past 60 years [1].

The formation of atheromatous plaques in arterial walls is a key element in development of atherosclerosis. The plaques have a complex histology, the core of which consist of lipids, chiefly cholesteryl esters, accumulated in the arterial intima. High level of plasma cholesterol, on the other hand, is undoubtedly a significant risk factor of atherosclerosis. These facts, recognized early in the history of atherosclerosis research, have been for decades the cornerstones for the undisputed dominant role of cholesterol in the science and management of CVD.

After “discovery” of the oxidized LDL, it soon became a target for investigations of molecular mechanisms of atherosclerosis, and the various ways by which oxidized LDL may contribute to

atherogenesis became evident. Based on this development, the idea of “the oxidation theory of atherosclerosis” was presented in 1989 [2].

The presence of lipid oxidation products (LOP) in plasma lipoproteins was first detected several decades ago [3–8]. Yet, physiological significance of the lipoprotein-specificity of LOP transport was recognized much later, as it turned out that plasma lipoproteins are active carriers of LOP. While LDL and triglyceride-rich lipoproteins (TRL) direct the transport towards peripheral tissues, HDL is active in the reverse transport of LOP [8–11]. LOP, formed during peroxidation of polyunsaturated fatty acids, can impair cellular functions in many ways and are well known from their multiple pro-atherogenic effects [12]. Therefore, emerging new information of lipoprotein-specific LOP transport likely affects conception of the etiology of atherosclerosis and CVD.

## 2. Oxidized LDL and Atherosclerosis

The oxidation theory of atherosclerosis is based on the fact that oxidative modification in the structure of LDL leads to inability to bind to the LDL receptor (LDLR), and thereby hampers normal metabolism of the lipoprotein. Scavenger receptors (like scavenger receptor A-1 and other scavenger receptors that were found later), on the other hand, bind avidly oxidized LDL, and this deviation of LDL metabolism leads to progressive transformation of scavenger receptor –bearing cells (macrophages, smooth muscle cells) into foam cells [13].

The mechanism whereby native LDL becomes a substrate for the scavenger receptor pathway is explained by initial oxidation of LDL lipids, and subsequent modification of apolipoprotein B by reactive LOP [12]. The reaction of LOP with the lysine groups in apolipoprotein B is postulated to be the reason for the altered receptor binding, since lysine residues are known to be required for LDL recognition by the LDLR [13].

Since the presentation of the idea of oxidation theory of atherosclerosis, it has been generally assumed that oxidation of LDL takes place within the vessel wall and is caused by endogenous free radical reactions induced e.g. by metal ions, superoxide anion, nitric oxide, lipoxygenase, or myeloperoxidase [12].

LDL oxidation contributes to atherogenesis not only by steering the metabolism towards scavenger receptor –bearing cells. Indeed, oxidized LDL has a wide spectrum of biological effects that may substantially increase its total proatherogenic impact [12,14]. Oxidized LDL is known to be cytotoxic and can cause endothelial injury or dysfunction allowing increased LDL deposition in the subendothelial space. Moreover, oxidized LDL is known to stimulate the production of monocyte chemoattractant protein-1 in vascular cells, which leads to recruitment of inflammatory cells; this is generally regarded as one key point in the pathogenesis of atherosclerosis. The injurious effects of recruited inflammatory cells may even be amplified by oxidized LDL, since it is known to be chemostatic for tissue macrophages and can induce the expression of scavenger receptors of macrophages. Yet another way through which oxidized LDL may be involved in atherogenesis is the mitogenic effect for smooth muscle cells. By this mechanism oxidized LDL can directly influence the formation of plaques and thickening of the vessel wall.

It is important to note that oxidized lipids in LDL appear to be responsible for most, if not all, of the atherogenic effects, since LOP have been shown to cause the toxic, inflammatory and mitogenic effects described for oxidized LDL (Table 1). Oxidatively modified apolipoprotein B, in turn, is merely known as the cause for disturbed LDLR-mediated cellular uptake of LDL.

**Table 1.** Adverse effects of LOP.

- 
- macrophage foam cell formation
  - chemotactic for monocytes, chemostatic for tissue macrophages
  - cytotoxic, induction of apoptosis
  - mitogenic for smooth muscle cells and macrophages
  - platelet aggregation and thrombosis
  - pathologic angiogenesis
- 

(Adapted from information provided in [12] and [14]).

### 3. The Protective Role of HDL

With advancement of the study of HDL it has come out that besides the lipid transport HDL appears to have several other physiological functions (Table 2). These include numerous examples of communication with cells as part of vital cellular functions, and the inactivation and removal of biohazards like bacterial toxins (for a recent review, see [15]).

**Table 2.** HDL functions.

---

•	Transport
○	cholesterol
○	fat-soluble vitamins
○	hormones
○	lipid oxidation products
○	xenobiotics
○	micro-RNAs
•	Cell communication
○	cholesterol efflux
○	signal transduction
○	endocytosis
•	Elimination of biohazards
○	bacterial toxins
○	lipid oxidation products

---

(Adapted from information provided in [15]).

Rather than on cholesterol carried by HDL, the various HDL functions are based on activities of entire HDL particles or distinct proteins or lipids on HDL [15]. Diversity of the physiological functions of HDL has been explained to result from heterogeneity of composition of the HDL particles. The prominent heterogeneity of HDL, which has come into daylight within the last two decades, has been somewhat surprising even to scientists working with this subject: the number of different HDL-associated proteins is estimated to be over 300, and the number of associated lipids may be even higher [15,16].

Concentrations of the distinct proteins and lipids on HDL may differ markedly from each other ranging from sub-micromolar ("low-abundant") to millimolar ("high-abundant"). The low-abundant HDL proteins, in turn, can be categorized either as lipoprotein-specific (e.g. lecithin:cholesterol acyltransferase, cholesteryl ester transfer protein, paraoxonase-1) or ancillary proteins. Despite their low abundance, ancillary proteins and low-abundant lipids are also known to have important biological functions [15,17].

The concentration of plasma HDL-cholesterol has for decades been regarded as an indicator of the risk of atherosclerosis and CVD. In addition to CVD, low level of HDL-cholesterol is known to be related to diabetes, chronic kidney disease, sepsis, and the risk of infectious diseases. It has turned out, however, that in case of CVD the HDL-cholesterol value may not be the best measure for estimation of risk, while new evidence has questioned a causal association between HDL-cholesterol and CVD [15,17]. In line with this, cholesterol efflux capacity, the first step of the "reverse cholesterol transport" (RCT) pathway and indicative of HDL functionality, has strong inverse association with indicators of atherosclerosis, and these associations are independent of the total level of HDL-cholesterol [18]. These findings underline significance of the HDL-mediated macrophage-cholesterol efflux, and have prompted a need for clinically usable biomarkers for estimation of efficacy of the efflux capacity. In addition to cholesterol efflux capacity also other functional measures of HDL, such as numbers of HDL particles or distinct HDL proteins, have proved to be better predictors of CVD than HDL-cholesterol [15]. It is obvious, as concluded by von Eckardstein et al. [15], that in clinical studies of the role of HDL only biomarkers directly related to essential pathological mechanisms should be used.


#### 4. Transport of Lipophilic Substances Other Than Plasma Lipids

In addition to plasma lipids and LOP, lipoproteins act as carriers for a wide variety of other lipophilic compounds in blood. For example, fat-soluble vitamins and non-nutrient lipophilic substances (phytochemicals, drugs, xenobiotics) that enter the alimentary tract can be absorbed in the intestine and incorporate into lipoproteins along with the dietary fat. These compounds, too, can be taken up by cells via receptor-mediated uptake mechanisms [19–25]. The physiological significance of the lipoprotein-dependent transport and uptake of substances other than dietary lipids has received little attention. It has been reported, however, that oxidized lipids in LDL are incorporated into the cells and detected in cell extracts [26]. Moreover, biologically active phytochemicals, incorporated into LDL and internalized by cultured cells via the LDLR pathway, have been shown to retain their biological activity [22].

Experimental studies with LDLR-overexpressing and –knockdown cells showed that LDLR regulates the behavior of drugs associated to VLDL/LDL [25]. The same report further showed that association with lipoproteins can affect the transport, metabolism, and efficacy of lipophilic drugs in humans. Consequently, the lipoprotein-mediated drug transport is a potential contributor to the transport and metabolism of lipophilic drugs in the body, the role of which should be considered when developing new pharmacotherapies [25]. In line with this, the evidence of the lipoprotein transport of fat-soluble xenobiotics would mean that lipoproteins are involved in distribution, and thereby toxic effects, of these chemicals in the body [27].

#### 5. Lipid Oxidation Products in Human Plasma

LOP arise during progression of the peroxidation of polyunsaturated fatty acids (Figure 1). Peroxidation reactions are induced by reactive free radicals and oxidant stress and can generate a wide variety of LOP of different structures. In the first phase of the stepwise chain of reactions, intermediary derivatives (e.g. hydroxides, hydroperoxides and epoxides) are formed. With advancement of the oxidative breakdown, these intermediary derivatives can give rise to formation of reactive aldehydes and less reactive products like ketones and alkanes (Figure 1).



**Figure 1.** Formation of LOP during peroxidation of polyunsaturated fatty acids.

Reactive aldehydes, in turn, can react and form adducts with cellular macromolecules, notably with proteins, giving rise to advanced lipid peroxidation end products that are known to cause protein modifications and disturb cellular functions [28,29]. Peroxidation of PUFA can also lead to formation of oxidatively modified lipids with biological activity, such as oxidized phospholipids and isoprostanes [28].

LOP in human plasma can have both endogenous and exogenous origins. Endogenously formed LOP result from oxidative stress in the body, which may be due e.g. to physical exercise, inflammation, irradiation, or exposure to chemicals [30,31]. Physical exercise has even proved to be a useful physiological model to investigate the formation and distribution of endogenous LOP [8,10,32]. Exogenous LOP, in turn, enter the body through the alimentary tract along with ingested food. It has been shown that LOP in food are absorbed in the intestine and incorporated into plasma lipoproteins together with the dietary lipids [5–7,10,33]. The average daily intake of LOP associated with dietary fat has been estimated to be more than 1 mmol per day [34].

LOP in food may arise from enzymatic oxidation, photo-oxidation, autoxidation, or thermal oxidation of food lipids, which can take place during storage and processing of food (Suomela, 2006). In addition, it has come out that peroxidation of food lipids occurs also in the alimentary tract during digestion [35]. Food LOP come from various lipid classes (e.g. triglycerides, phospholipids, sterol esters), and can be present attached to the original lipid (glycerol or sterol) backbone, or as free molecules. Food-contained LOP are shown to be absorbed and incorporated into plasma lipoproteins, including LDL and HDL [5–7,10,33]. In all, this indicates that food LOP appear as another type of dietary risk factors, emphasizing the importance of pro/antioxidant effects of nutrients.

The potential significance of food as a source of LOP was clearly shown in a study, where young healthy male volunteers consumed pan-fried and sous-vide thermally processed beef rump steaks. The experimental meals were prepared replicating real cooking conditions, which gave tender or medium rare steaks. Twenty-three oxidation-modified lipid structures were detected from the beef, and 15 of these were also found in postprandial plasma [36].


## 6. Lipoprotein-Specific Transport of LOP

The transport of LOP by plasma lipoproteins has received little attention, and the presence of LOP in circulating lipoproteins has mainly been regarded to result from endogenous oxidative stress. Accumulating evidence suggests, however, that the presence of LOP in plasma lipoproteins is not a random consequence of oxidative stress, but rather indicative of specific LOP transport functions. It appears that plasma lipoproteins are active carriers of LOP, LDL directing transport toward peripheral tissues and HDL being active in the reverse transport. Biological mechanisms responsible for the lipoprotein-specific transport and cellular uptake of the native (unoxidized) lipids may at the same time lead to lipoprotein-specificity in transport and cellular uptake of LOP.

Evidence of the lipoprotein-specific LOP transport comes mainly from studies of lipoprotein LOP transport during physiological oxidative stress and is supported by studies of the functionality apolipoprotein A-1 (Apo A-1) mimetic peptides.

### 6.1. Studies on Physiological Oxidative Stress in Human Volunteers

Acute increases in plasma LPO concentrations are known to be caused e.g. by LOP-containing meals [5–7,10,33] or by strenuous physical activity [31,32]. Both these physiological circumstances have been used as experimental models to investigate the lipoprotein LOP transport (Figure 2)[10]. Food LOP, typically found in fatty meals, represents an exogenous source of LOP for the human body, while LOP generated as a result of physical exercise-induced oxidative stress is an example of endogenously formed LOP.



**Figure 2.** Effects of high-fat meal and strenuous physical exercise on lipoprotein LOP.

High-fat meal (left panel): Blood samples were taken before, and 2, 4 and 6 h after consumption of the meal (a standard hamburger meal) and 4 dL of fruit juice. Physical exercise: The subjects performed on a treadmill a 40min tempo run at the velocity corresponding to 80%  $VO_{2max}$ . Blood samples were taken before, in the middle, and immediately after the exercise, as well as after a 90 min recovery period. Based on data from [10].

The roles of lipoproteins as carriers of exogenous LOP in human body were investigated in a model with a common hamburger meal (available in fast-food restaurants worldwide) as the “high-fat meal” [10]. Altogether over 100 volunteers participated these studies, in which the concentrations of LOP in plasma lipoproteins were monitored during postprandial period following the hamburger meal [8,10,37].

The results consistently showed that concentrations of LOP in triglyceride-rich lipoproteins (TRL and LDL) started to rise within 1 hour after the meal. LOP levels in TRL and LDL reached maximal levels within 2 – 4 hours, after which they gradually returned to pre-meal values. It is worth noting that even though the concentration of LOP in LDL did increase, the concentration of LDL-cholesterol was not changed. Moreover, detailed studies showed that the fatty meal –induced elevation of lipoprotein LOP was not due to increased oxidative stress, nor to altered antioxidant functions [10]. Unlike in the case of TRL and LDL, there was only a small and delayed increase in HDL-LOP concentration, 6 hours after the meal [10]. The uneven (lipoprotein-specific) postprandial distribution of LOP between the lipoprotein classes showed that food-derived LOP are not randomly incorporated into various lipoproteins.

In contrast to the substantial rise in LDL-LOP, the level of oxidatively modified apolipoprotein B has been reported to decrease after a high-fat meal [38]. This finding supports of the view that the fatty meal –induced increase of LDL-LOP is not due to oxidation of lipids within the LDL particle.

Among several other indicators of oxidative stress, strenuous physical exercise is well known to increase the level of LOP in human plasma [31,32], and follow-up of concentrations of LOP in lipoproteins during and after the exercise has allowed investigation of the transport of endogenously formed LOP [10].

Opposite to the effect of a fatty meal, strenuous physical exercise has little, if any, effect on concentration of LOP in LDL (Figure 2). Contrary to this, physical exercise results in a substantial increase in concentrations of LOP in HDL. This effect may last for hours after a single bout of aerobic exercise, and it appears to depend on intensity of the physical activity [8,10]. In a study with well-trained athletes (middle-distance vs. marathon runners) it was found that the exercise-induced increase of LOP in HDL is related to training history of the athletes [39].

Increase of LOP in HDL particles is not due to oxidation of lipids carried by HDL, since no such increase is seen in LDL despite the lower oxidation resistance of LDL [40]. In line with this, as reasoned by Shao and Heinecke [9], oxidation of either HDL or LDL in plasma is regarded highly

unlikely. Moreover, it seems unlikely that LOP would be transferred directly from LDL to HDL in plasma, since the transfer of LOP between lipoproteins appears to be too slow to affect their distribution in lipoproteins [3].

Taken together, the studies on physiological oxidative stress point to lipoprotein-specificity of the transport of LOP in circulation: LOP in food are absorbed and incorporated into serum TRL and LDL, directing the flow of LOP towards peripheral tissues; HDL appears to have an opposite transport function, and can respond to oxidative stress by substantially increasing the reverse transport of LOP.

### 6.2. Experimental Evidence of the Role of HDL

More evidence supporting the idea of LOP scavenging, transport and clearance function of HDL comes from studies with apo A-1 mimetic peptides and apolipoprotein M (apo M). Similarly to the atheroprotective functions of HDL, apo A-1 mimetic peptides are known to have positive effects on lipid metabolism, oxidative stress, inflammation, and insulin resistance [41–43]. Studies by Van Lenten et al. [41] showed that apo A-1 mimetics have a remarkable binding affinity for LOP. It is of special interest that the LOP scavenging and removing capacity of apo A-1 mimetics was found to be strongly related to the positive health effects [41].

Studies with the “4F peptide” as a representative of the apo A-1 mimetics has given additional evidence regarding the postulated link between the clearance of LOP and atheroprotection. While decreasing hepatic and plasma concentrations of LOP, the “4F peptide” also causes a significant reduction in atherosclerotic lesions [42,43]. In line with this, the ability of the HDL-associated apo M to trap oxidized phospholipids appears to be associated with its atheroprotective effects [44].

Studies in a perfused rat liver model have shed light on the role of HDL in clearance of LOP from the body. Early studies by Stocker and coworkers [45] showed that the liver removes LOP from HDL, but not from LDL particles. In support of the role of HDL in clearance of LOP, Fluiter et al. [46] further reported that there is a preferential liver uptake, coupled to a rapid biliary excretion pathway, of oxidized cholesteryl esters in HDL as compared with the unoxidized cholesteryl esters.

A recent study showed that HDLs can utilize their capacity of loading themselves with lipophilic compounds, similarly to their ability to extract cellular cholesterol, to reduce the cell content of lipophilic drugs and xenobiotics. ABC transporters were not required for the process but, however, could favor it [27].

Together with the evidence of intervention studies in human volunteers, results of the experimental studies suggest that the transport of LOP by HDL is part of a protective transport function, where HDL protects peripheral vessels and tissues from LOP by transporting these potentially toxic substances to the liver, as originally proposed for cholesterol [8–11].

## 7. Potential Significance of Lipoprotein LOP Transport for Oxidative Stress

Studies of LOP transport during physiological oxidative stress show that LDL carries LOP to peripheral tissues, and HDL is active in the reverse transport. Knowing the pro-oxidant potential of LOP, this would advocate a pro-oxidant role for the LDL transport, whereas transport by HDL would be part of antioxidant defence of the body.

At present there is little direct evidence of a causal connection between LDL and oxidative stress, but indirect evidence has suggested a link between these two. The level of oxidative stress is reported to be elevated among patients with FH [47,48], and suppressing of oxidative stress has been speculated as a potential therapeutic means to reduce atherogenesis in patients with FH [48]. In line with this, statin treatment has been reported to be associated with decreased oxidative stress [49,50], prompting speculations of the contribution of antioxidant properties of statins in their beneficial effects [50].

Besides the reverse cholesterol transport, HDL is known to have several other functions potentially affecting to atherogenesis, and antioxidant activity has frequently been mentioned as one of these. Inactivation of lipid hydroperoxides and hydrolysis of oxidized phospholipids are known as mechanisms of the LDL-specific antioxidant action of HDL. Apo A-1 and enzymes

associated with HDL particles are known to be important components of HDL particles contributing to these mechanisms [11,51].

Like in the case of LDL there is little direct evidence of a causal connection between HDL and oxidative stress in human body. Yet, as part of the studies on physiological oxidative stress and LOP transport, a positive association was found to exist between plasma HDL level and clearance of the fatty meal-induced LOP load, which was suggested to indicate antioxidant function of HDL in vivo [10,37].

## 8. Measurement of Lipoprotein LOP Transport

The common assays for “oxidized LDL” are based on (i) measurement of oxidatively modified apolipoprotein B, (ii) autoantibodies to oxidized LDL, or (iii) LOP in LDL. It is obvious that focusing either on the protein part of the LDL particle, or on autoantibodies developed against oxidatively modified LDL, are not indicative of the LOP transporting function of LDL. Instead, LOP found in the lipid part of LDL may be used for estimation of the LOP transport by LDL. The present analytical techniques for estimation of LOP in LDL are based on determination of oxidatively modified lipids or end products of lipid oxidation in LDL isolated from plasma. Common examples of these are conjugated dienes, fatty acid hydroperoxides, oxidized phospholipids, and malondialdehyde. Interest for LOP in HDL appeared much later than in the case of LDL. Therefore, there are far less studies conducted and data available of LOP in HDL.

Recent studies of the lipoprotein-specific transport of LOP are by large based on measurement of conjugated dienes in lipid fraction of LDL. LOP, as measured by the diene conjugation assay, are distributed to all main lipid classes of LDL. This assay has strong correlation with assays for oxidized apolipoprotein B and autoantibodies against oxidized LDL [52,53].

Based on the direction-specificity of the lipoprotein transport, it has been proposed that the level of LOP in LDL (LDL-LOP) would be indicative of potential exposure of peripheral vessels and tissues to LOP, and the concentration of LOP in HDL (HDL-LOP) would indicate reverse transport of LOP toward the liver. Accordingly, the ratio of LDL-LOP/HDL-LOP would be indicative of the direction of the net movement of LOP in the body [8,10].

## 9. Lipoprotein LOP and the Risk of Atherosclerosis

Cholesterol is known to be involved in many physiological functions, while LOP appear to be metabolic residues with no physiological role. Cholesterol per se is not regarded as the progenitor of chain of events leading to development of atherosclerosis. On the other hand, oxidized lipids in LDL are involved in all major pathophysiological processes leading to thickening and stagnation of arteries (Table 1). Therefore, the fact that LOP are transported and taken up by cells together with cholesterol has raised the question of a possible role of the lipoprotein LOP transport in development of atherosclerosis and CVD. As an attempt to answer this question, connections of lipoprotein LOP with known atherosclerosis risk factors, metabolic disorders related to atherosclerosis, subclinical atherosclerosis, and clinically verified atherosclerosis have been investigated in studies with human volunteers (Table 3).

**Table 3.** Findings of atherosclerosis –related factors and lipoprotein LOP.

<b>Influence of / association with</b>	<b>Oxidized LDL lipids</b>	<b>Oxidized HDL lipids</b>
Age	Increases with age	Decreases with age
Gender	Male > female	Female > male
Body mass index	Positive association	Negative association
Weight reduction	Substantial decrease	Not studied
Tobacco smoking	Elevated among smokers	Not studied
Statin treatment	Substantial decrease	
Acute physical exercise	No change / decrease	Increase

Physically active lifestyle	Substantial decrease	Slight increase
Fatty meal	Substantial increase within 2 hours	Slight increase after several hours
Saturated fat	Positive association	Not studied
Polyunsaturated fat	Negative association with n-6 PUFA	Not studied
Atherosclerosis	Positive association with brachial, carotid, and coronary atherosclerosis	Negative association with coronary atherosclerosis
Insulin resistance	Positive association	Not studied
Metabolic syndrome	Positive association	Not studied
Development of fatty liver	Positive association	Negative association

### 9.1. Common Risk Factors

**Age** and **gender** are well known factors related to the risk of atherosclerosis and CVD. The incidence of diseases associated with atherosclerosis typically increases with age and is higher in men than in women. There are only few studies where the dependence of lipoprotein LOP on either age or gender have been investigated. A substudy of The Cardiovascular Risk in Young Finns Study, with a total of 1395 subjects (629 males and 766 females), is by far the most comprehensive of these [53]. In addition to age and gender, associations of lipoprotein LOP with several other atherosclerosis risk factors were investigated in this multicenter follow-up study.

A significant sex difference was found to exist in both LDL-LOP and HDL-LOP concentrations: compared to men, the values of LDL-LOP were lower and HDL-LOP higher in female subjects [53]. Despite the relatively narrow age-range of participants (24–39 years old at time of blood sampling), age was associated in multivariate models with both LDL-LOP (direct association) and HDL-LOP (inverse association) [53]. It is worth mention that the age-dependence of HDL-LOP appears to be opposite to that for HDL-cholesterol, which has been reported to increase with age [54].

Of the common **lipid risk factors** of atherosclerosis, total cholesterol, LDL-cholesterol, and triglycerides showed strong correlation with LDL-LOP, but not with HDL-LOP [53]. Positive correlations of LDL-LOP with plasma lipids have been reported also in several previously published studies [52,55–57].

**Blood pressure** is another factor strongly related to the risk of atherosclerosis. As an indication of possible association between lipoprotein LOP and blood pressure, an increased level of LDL-LOP has been reported in young men with borderline hypertension. In this study ambulatory recording of blood pressure over a 24 h period was employed [58]. In line with this, a bivariate association between diastolic blood pressure and LDL-LOP was found in The Cardiovascular Risk in Young Finns Study [53].

The effect of **tobacco smoking** on lipoprotein LOP has been investigated in two studies. Participants of the first study (n=164) were obtained from a population cohort of Finnish men aged 40-70 years. In this study smokers had significantly higher levels of LDL-LOP (21%) than non-smokers [59]. In The Cardiovascular Risk in Young Finns Study daily smoking was found to be inversely associated with HDL-LOP in women [53].

### 9.2. Physical Activity

Physically active lifestyle is well known have beneficial effects on cardiovascular health reducing the overall risk of coronary heart disease and stroke [60,61]. During the past three decades several studies have been carried out to investigate effects of physical activity on lipoprotein LOP. The investigations range from cross-sectional studies of effects of physically active lifestyle to intervention studies on various forms of acute or long-term physical activity. Since the assay for HDL-LOP was taken in use 15 years later than that for LDL-LOP, the data concerning HDL-LOP is missing from studies published prior 2010.

The first study reporting of effects of physically active lifestyle was performed with veteran endurance athletes participating competitively in orienteering, running, cycling or triathlon. The

athletes were compared with matched subjects of the same age and socioeconomic status. It was found that the athletes had significantly lower LDL-LOP values than control subjects, and that LDL-LOP correlated inversely with leisure intensive activity score ( $MET_{int}$ ; an indication of intensive physical exercise energy expenditure) [55]. In another study on effects of physically active lifestyle, the study subjects consisted of young (9-15 years old) female athletes and nonathletic controls. The results of this study were well in line with the previous study with veteran athletes: physically active girls had lower LDL-LOP values, and LDL-LOP was inversely associated with the MET-index [62].

More evidence of the decrease of LDL-LOP by physical activity was obtained from an intervention study where sedentary middle-aged men and women ( $n=104$ ), recruited by the local occupational health service, participated in a 10-month exercise program. The exercise intervention produced favourable changes in lipid profiles, and reduced LDL-LOP by 23% and 26% in men and women, respectively. The ratio of LDL-LOP/LDL-cholesterol was also decreased, 14% in men and 18% in women [63]. Another long-term follow-up study was performed with marathon runners ( $n=127$ ) who were preparing for a marathon race [64]. Venous blood samples were taken at 4 different time points during the study: 3 months (training period) and 6 days (final preparation period) prior to marathon run, and on the marathon day 2-4 h before and immediately after the race. During the 6-days preparation period, subjects decreased training and increased carbohydrate intake. Interestingly, the training period did not affect LDL-LOP values, but after the preparation period the values were substantially increased. Then again, the marathon run restored LDL-LOP values towards the level before the preparation period [64].

The effect of prolonged, low intensity exercise on LDL-LOP was investigated in an experimental setting, where healthy well-trained men performed a walk exercise on two consecutive days (6 h per day) [65]. Like in the above studies, prolonged physical activity resulted in a decrease (-25 %) in the level of plasma LDL-LOP. Favourable changes were seen also in conventional plasma lipids, while serum LDL-cholesterol (-14 %) and triglycerides (-22 %) decreased, and HDL-cholesterol (+9 %) increased [65].

Thus far, only one published study has reported effects of physically active lifestyle on HDL-LOP. In this study 6 months supervised physical activity program increased the concentration of HDL-LOP and the ratio of HDL-LOP/HDL-cholesterol among elderly women. The concentration of LDL-LOP remained unchanged [66].

The effects of acute strenuous physical activity have been investigated in long distance runs and treadmill exercises. During long distance (31 km) run the total amount of LOP in whole serum increased but the concentration of LDL-LOP remained unchanged [67,68].

The finding that long distance run increased the amount of LOP in whole serum but not in LDL particles raised a question of the carrier of LOP generated during physical activity. This topic was later studied in more detail, and like in previous studies physical exercise did not affect LDL-LOP level. Contrary to this, HDL-LOP was substantially increased, and remained elevated for hours after the exercise. In addition to the total amount of HDL-LOP, also the ratio of HDL-LOP/ HDL-cholesterol was increased [10]. This was a significant novel discovery pointing at LOP clearing function of HDL. <sup>61,69,70</sup>

### 9.3. Obesity and Weight Reduction

Overweight and obesity are associated with significantly increased morbidity and mortality from CVD [61,69,70]. Consequently, weight reduction is among the first line of defence for patients with increased risk of developing atherosclerosis and resulting CVD.

The relationships of LOP with body weight and waist circumference were investigated in the comprehensive study material of The Cardiovascular Risk in Young Finns Study. It was found that LDL-LOP has significant bivariate associations with both BMI and waist circumference [53]. In accordance with this, Sorokin et al. [71] reported associations between lipoprotein LOP and BMI (positive association with LDL-LOP and negative association with HDL-LOP).

The effect of weight reduction has been investigated in three independent intervention studies. In the first study, effects of a 12-week weight-reduction and a subsequent 9-month weight-

maintenance program on LDL-LOP were investigated in a group (n=77) of obese premenopausal females [72]. A 13 kg decrease in body weight was associated with a 40% decrease in LDL-LOP concentration. The weight reduction correlated with the decrease of LDL-LOP. The concentration of LDL-cholesterol was also decreased, but to a smaller extent (11%). Altogether, this was the first study to investigate the effects of obesity and weight reduction on oxidized LDL.

In another study, performed with obese men (n=68), a 2-month weight reduction period was followed by 6-month weight maintenance program, after which participants of the study were followed up for another 2 years period [73]. The mean weight loss at the end of the weight reduction period was 14,5 kg (14 %), which associated with a 22 % decrease in LDL-LOP concentration. At the end of the 2-year follow-up, the weight regain from the end of the weight reduction period in whole study group was 11% and regain in concentration of LDL-LOP was 30 %. To study the effect of successful weight maintenance, the study population was divided into two subpopulations regarding success in maintaining the weight loss. It was found that the regain of weight was firmly associated with an increase of LDL-LOP concentration [73].

Relationship between the change of body weight and the level of LDL-LOP has been confirmed in yet another study on caloric restriction of obese individuals [74]. A 6-week weight reduction by very-low-calorie diet resulted in 11 kg (12%) weight loss. Serum LDL-LOP concentration was decreased (31%), and the decrease correlated with weight loss [74].

Cardiorespiratory fitness has been shown to protect obese individuals from cardiorespiratory mortality [75]. In accordance with this, overweight men with good aerobic or muscular fitness were found to have lower levels of LDL-LOP than unfit men of the same weight [76]. This finding further strengthened the view of a connection between LDL-LOP and the risk of atherosclerotic diseases.

#### 9.4. Dietary Factors


Despite the fact that nutritional aspects are well-known contributors to the risk of atherosclerosis and CVD, the number of studies focused on nutrition and lipoprotein LOP has remained low.

In a cross-sectional study setting, it was found that serum polyunsaturated fatty acids (PUFAs), and in particular the n6 PUFA proportion, were negatively associated with LDL-LOP, whereas the associations of saturated (SFAs) and monounsaturated fatty acids (MUFAs) were positive [77]. These findings at the population level are in line with observations linking PUFAs with lower and SFAs with increased risk of CDV [78].

The finding of the positive effect of PUFA on LDL-LOP level is supported by results from an intervention study with modified Mediterranean-type diet rich in n3 PUFA. A 12-week intervention resulted in an 8 % decrease in serum LDL-LOP level, and the concentrations of total and LDL-cholesterol decreased 8 % and 11 %, respectively [79].

The effects of separate components of nutrition on lipoprotein LOP were investigated in two early intervention studies. The first of these focused on effects of lycopene, a carotenoid available from tomatoes. In this study dietary supplementations doubled serum lycopene levels and significantly decreased LDL-LOP, but had no effect on total, LDL- or HDL-cholesterol concentrations [80]. In another study hyperlipidemic subjects consumed whole almonds as snacks for 1 month. Almond consumption resulted in significant reductions in LDL-LOP (14%) and LDL-cholesterol (9%) concentrations [81].

Marniemi et al. [82] studied effects of 8 weeks antioxidant vitamin supplementation (100 mg  $\alpha$ -tocopherol + 500 mg ascorbic acid per day) and found that even though antioxidant supplementation significantly increased the antioxidant capacity of LDL particles, the concentration of LDL-LOP did not change (Figure 3). The same result was obtained also in another antioxidant intervention study with different antioxidant dosing and treatment period. Daily doses of  $\alpha$ -tocopherol (294 mg), ascorbic acid (1000 mg) and ubiquinone (60 mg) for 4 weeks raised substantially the antioxidant capacity of LDL and the levels of antioxidant vitamins in blood but did not affect the level of LOP either in serum or LDL [68](Figure 3). The fact that antioxidant supplementation does not affect serum LDL-LOP levels despite the substantial increase in LDL antioxidant capacity indicates that LOP carried by circulating LDL do not come from peroxidation of lipids within the LDL particles.



**Figure 3.** Effects of antioxidant intervention on LDL-LOP and LDL antioxidant capacity (LDL-AOC) in healthy volunteers. Study 1: daily doses of 294 mg 100 mg d- $\alpha$ -tocopherol acetate, 1000 mg of ascorbic acid and 60 mg ubiquinone for 4-weeks. Study 2: daily doses of 100 mg d- $\alpha$ -tocopherol acetate and 500 mg of ascorbic acid for 8-weeks. Based on data from [68] (Study 1) and [82] (Study 2).

Similarly to effects of antioxidant vitamins, flavonoid supplements in food failed to influence serum LDL-LOP levels. Healthy male volunteers consumed flavonols (isorhamnetin, quercetin and kaempferol) daily for 4 weeks at a dose level that increased flavonols concentrations in plasma. Yet, serum LDL-LOP remained at the control level [83].

The studies described above have provided information of long-term dietary effects. In addition to these, acute postprandial effects of dietary fat on lipoprotein LOP have been investigated (described in *Studies on physiological oxidative stress in human volunteers*). In short, these studies substantiate that LOP in food are absorbed and incorporated into plasma lipoproteins and transported in plasma similarly to the native (unoxidized) lipids. This underlines the potential importance of lipoprotein transport of food-derived molecular species with respect to the risk of atherosclerosis.

#### 9.5. Metabolic Disorders Related to Atherosclerosis

Three clinical studies have focused on the co-existence of LDL-LOP and metabolic disorders related to atherosclerosis. In these studies, lipoprotein LOP was investigated in relation of insulin sensitivity, metabolic syndrome, and the development of fatty liver.

A 32-month lifestyle intervention study investigated whether changes in LDL-LOP levels are related to changes in insulin sensitivity [84]. The intervention consisted of two months weight reduction followed by 6-month dietary and physical exercise counseling, and a 2-year follow-up period. The experimental protocol suited well for the purpose, since it caused substantial changes in insulin sensitivity (determined by the homeostasis model assessment of insulin resistance, HOMA-IR) of participants. The detected changes in HOMA-IR were clearly parallel to those in LDL-LOP and LDL-LOP/HDL-cholesterol ratio, suggesting a connection between insulin metabolism and LDL-LOP [84].

Metabolic syndrome and lipoprotein LOP were investigated in a nationally representative sample of Finnish young men with and without metabolic syndrome (MetS) [85]. Participants were divided according to the IDF 2007 criteria into MetS (n=54) and non-MetS (n=790), and age, smoking and leisure-time physical activity were used as covariates. The results showed that a high LDL-LOP level predicted MetS, and it was concluded that elevated concentration of LDL-LOP is a significant predisposing factor in the development of MetS [85].

Fatty liver is another metabolic disorder that is a risk factor for CVD, and it is apparent that there are common progenitors for these pathological conditions [86]. Kaikkonen et al. [87] have studied associations of lipoprotein LOP with fatty liver assessed by ultrasonography. Participants of the study were 1286 middle-aged subjects with normal liver, and 288 subjects with fatty liver. After adjustment for age, sex, leisure-time physical activity, body mass index, alcohol intake, smoking, serum LDL- and HDL-cholesterol as well as particle concentrations, participants with elevated LDL-LOP were found to have an increased risk for fatty liver. In particular, the high ratio of LDL-LOP/HDL-LOP indicated increased risk. This relation was independent of LDL and HDL particle concentrations and cholesterol content, other common risk factors for fatty liver, and inflammation. The LDL/HDL-cholesterol ratio was not associated with future fatty liver [87]. Based on these data, lipoprotein LOP appeared to be implicated in yet another pathophysiological process, the development of fatty liver in middle-aged adults.

#### 9.6. Subclinical Atherosclerosis

Clinical complications of atherosclerosis usually occur in middle and late age. Yet, the development of atherosclerosis may begin early in life, and the asymptomatic subclinical phase may last for decades. Early atherosclerotic changes, such as thickening of vessel wall and alterations in endothelial functions, are indicators of subclinical atherosclerosis that can be detected by various non-invasive imaging methods. The use of these methodologies has given information of connections between subclinical atherosclerosis and LDL-LOP.

Toikka et al. [58] investigated indicators subclinical atherosclerosis and LDL oxidation among young healthy men with borderline hypertension. High-resolution ultrasound was used to measure intima-media thickness (IMT) of the carotid and brachial arteries, cardiac dimensions, and brachial artery endothelial function. In this study LDL-LOP was found to correlate positively with both carotid and brachial IMTs. In multivariate analyses, the only predictors of carotid IMT were 24-hour systolic blood pressure and LDL-LOP [58].

Another study from the same group investigated relationships between myocardial flow reserve, carotid IMT and LDL-LOP in young men free from coronary heart disease [88]. Basal and stimulated coronary blood flow was measured using positron emission tomography (PET) in 55 healthy men, and the mean carotid artery IMT was measured using high-resolution ultrasound. In this study LDL-LOP was found to correlate inversely with flow reserve, and directly with carotid IMT. These data suggested that increased LDL-LOP is directly related to early structural and functional atherosclerotic vascular changes.

In a study on surrogate markers of asymptomatic atherosclerosis, Toikka et al. [89] investigated arterial elasticity, serum lipoproteins and LDL-LOP in young healthy men and age-matched asymptomatic men with FH. As a marker of arterial elasticity, compliance in the thoracic aorta was measured by using magnetic resonance imaging, and in the common carotid artery by using ultrasound. It was found that elasticity of both the carotid and brachial arteries was negatively associated with LDL-LOP [89]. In all, this was the first study to demonstrate an *in vivo* association between oxidized LDL and arterial elasticity.

#### 9.7. Clinically Verified Atherosclerosis

The relation between LDL-LOP and coronary atherosclerosis was investigated in a sample of 62 men who underwent diagnostic coronary angiography and sonography to measure the carotid IMT [57]. It was found that the LDL-LOP/LDL-cholesterol ratio and carotid IMT were the only factors associated independently with the severity of coronary atherosclerosis. Moreover, patients with

multi-vessel disease who did not use lipid-lowering therapy had 41% higher LDL-LOP/LDL-cholesterol ratio than patients with normal vessels.

The finding of association between LDL-LOP and coronary atherosclerosis was confirmed and extended to also apply HDL-LOP in a study based on measurement of coronary plaque parameters by CT angiography [71]. While LDL-LOP had a significant positive and HDL-LOP negative association with non-calcified plaque burden, the traditional lipid risk factors showed no associations.


### 9.8. Mortality

The predictive role of LDL-LOP in all-cause mortality was investigated in a population of 1260 elderly (aged 64 years or more) inhabitants [90]. They participated the study in 1998-1999, and the medical records were re-examined one decade later. During the ten-year period 467 (37%) participants had died, and in 36% of cases the cause of death was atherosclerotic and/or ischemic CVD. Comparisons between the survivors and deceased subjects showed that total levels of serum LDL-LOP did not differ between the groups. However, when proportioned to LDL-cholesterol, HDL-cholesterol or apo A-1, LDL-LOP stood out as a predictor of all-cause mortality. This finding was independent of age, sex, BMI, smoking, blood pressure and diabetes.

### 9.9. Statin Treatment

Cholesterol lowering therapy is a major remedy in treatment of atherosclerotic vascular diseases, and the favorable effects of statins in primary and secondary prevention are well documented [91,92].

The effects of statins on LDL-LOP have been investigated in three different studies (Figure 4). The first study was a comparison between the effects of a Mediterranean-type diet and simvastatin treatment [79]. The trial was conducted in previously untreated hypercholesterolemic men aged 35 to 64 years. Simvastatin was given for 12 weeks at a dose of 20 mg/day. This treatment decreased total and LDL-cholesterol concentrations by 21% and 30%, respectively. The concentration of serum LDL-LOP was also decreased (by 16%). However, when serum LDL-LOP was proportioned to LDL-cholesterol, it appeared that the LDL-LOP/LDL-cholesterol ratio in fact was increased by 13% [79]. This means that despite the fact that the total amount of LDL-LOP in serum decreased due to simvastatin-treatment, concentrations of LDL-LOP in the remaining LDL particles increased.



**Figure 4.** Effect of statin treatment on LDL-cholesterol, LDL-LOP and the ratio of LDL-LOP/LDL-cholesterol. Statin treatments were investigated with three different statins in three separate studies. In the first study [79] simvastatin was given for 12 weeks at a dose of 20 mg/day ("Simva 1"). In the second study [93] simvastatin ("Simva 2") and atorvastatin ("Atorva") were given for 12 months at a dose that was initially 20 mg/day. The dose was doubled at 12 weeks if the subjects had not achieved predefined lipid goals. Subjects in the third study [94] were given pravastatin ("Prava") for 4 months at a dose of 40 mg/day.

Another study investigated the effects of 12-month simvastatin and atorvastatin therapies on LDL-LOP in patients with hypercholesterolemia and CVD [93]. Initial daily dose for both statins was 20 mg. The dose was doubled at 12 weeks if the subjects had not achieved predefined lipid goals (LDL-cholesterol <2.4 mmol/L; serum triglycerides <1.5 mmol/L). After 52 weeks' treatment LDL-cholesterol had decreased 47% by simvastatin and 50% by atorvastatin. Transient reductions in serum LDL-LOP concentrations were observed by both statins during the first 12 weeks' treatment, but after the 52 weeks' treatment period LDL-LOP did not differ from the baseline level in either statin group. This study confirmed results of the previous study by Jula et al. [79] regarding increase of the LDL-LOP/LDL-cholesterol ratio. After 52 weeks the ratio was substantially increased by both simvastatin (52%) and atorvastatin (59%) treatments [93].

The effects of pravastatin have been investigated in a study with 42 young normocholesterolemic subjects with type-1 diabetes [94]. Pravastatin was given to study subjects at a dose of 40 mg/day. Glycaemic control and insulin sensitivity remained unchanged during the 4 months' treatment period. Pravastatin decreased serum LDL-LOP concentration by 18%. Similarly to treatments with other statins, pravastatin, too, increased the ratio of LDL-LOP/LDL-cholesterol (by 32%).

## 10. LOP and Cholesterol: Commonalities and Disparities

Studies on LOP and the risk of atherosclerosis, as described in the previous chapter, indicate that attributes of lipoprotein LOP as a potential risk factor largely parallel those of cholesterol. As in the case of cholesterol, high LDL-LOP associates with increased risk and high HDL-LOP with protection. Lipoprotein LOP is related to blood pressure and BMI like cholesterol, and physically active lifestyle and caloric restriction affect similarly on both lipoprotein LOP and cholesterol levels. Due to the common transport and cellular intake mechanisms, however, the available data do not allow elucidation of independent effects of either cholesterol or LOP.

The age-dependency of lipoprotein-contained LOP and cholesterol, in turn, was found to be different. In case of LOP, the concentration in LDL increases and in HDL decreases with age. While there appears to be no clear trend for a change of LDL-cholesterol, the concentration of HDL-cholesterol is reported to increase with age [54]. Increase of the LDL-LOP/HDL-LOP ratio may mean increase of pro-oxidant burden and elevated risk for atherosclerotic changes with increasing age.

The most prominent differences between lipoprotein LOP and cholesterol are seen in their acute responses to high-fat meal and physical exercise. Both of these have notable influences on lipoprotein LOP levels: a fatty meal increases for several hours the concentration of LDL-LOP, and strenuous physical exercise elevates substantially the concentration of HDL-LOP. The effects of high-fat meal and exercise on lipoprotein cholesterol levels are negligible.

The different responses of LDL-LOP and LDL-cholesterol to fatty meal demonstrates that unlike cholesterol, LOP in food may rapidly affect the amount of atherogenic substances in LDL. It should be kept in mind that LOP comprise of molecules which have no physiological role and are capable of directly initiating and boosting atherogenic processes. The exercise-induced increase in HDL-LOP, on the other hand, may be indicative of protective LOP clearing function of HDL, but also of the ability of HDL to respond rapidly to increased LOP levels in the body.

Yet another disparity between lipoprotein LOP and cholesterol is seen in their response to statin treatment. Three independent studies on statins and LDL-LOP have uniformly shown that statin treatment is less effective in reducing the concentration of LDL-LOP than that of LDL-cholesterol, which results in elevation of the LDL-LOP/LDL-cholesterol ratio. The apparent discrepancy may

indicate that LOP in LDL impair the LDL-receptor –mediated cellular intake of LDL particles. Another possible explanation is that statin accelerates the LDL-receptor –mediated removal of native LDL particles from circulation, but at the same time the rate of the scavenger receptor –mediated removal of oxidized LDL is not changed [93]. Nevertheless, a higher amount of LOP in remaining LDL particles could mean increased atherogenicity of circulating LDL. It is well known that CVD events may remain prevalent also during high-intensity statin therapy [95]. This so called “residual risk” appears to depend on lipid and/or lipoprotein factors [96], and it is tempting to speculate that enrichment of LOP in LDL, caused by statin treatment, could in part explain the residual risk.

## 11. Lipoprotein LOP as a Potential Risk Factor

LOP in lipoproteins have usually been regarded as a consequence of endogenous oxidative stress, but the presented evidence of lipoprotein-specific LOP transport is suggesting a role for lipoprotein LOP as a risk factor of atherosclerosis. It is therefore worth considering how this would affect the conception and management of the disease.

### 11.1. LOP as Dietary Risk Factor

Oxidative modification of LDL has long been assumed to result from endogenous free radical reactions [12]. It has come out, however, that a fatty meal increases the amount of LOP in LDL, capable of causing atherogenic LDL modifications. These findings draw attention to another type of dietary risk factors, LOP formed e.g. during food storage, processing, or digestion, emphasizing the importance of pro/antioxidant effects of nutrients. In support of this view, experimental evidence shows that postprandial LDL is more atherogenic than LDL in the post absorptive state. The more pronounced atherogenicity was postulated to be due to lipid peroxides in postprandial LDL [97].

A hamburger meal, the LOP content of which was reported to be close to 1 mmol, was sufficient to double the concentration of LOP in postprandial LDL [10]. In UK the average daily intake of lipid hydroperoxides associated with fats and oils was estimated to be approximately of 1.5 mmol per day [34]. It is therefore possible that at the average level of fat consumption a substantial part of LDL-LOP could be of dietary origin. This applies particularly to the postprandial situation, and except for the early morning, most individuals are in the nonfasting state most of the time. In all, if the conclusion of the dietary origin of LOP is tenable, then LOP in food should be considered as a notable risk factor for atherosclerosis. It would also mean that LDL is an important mediator of the risk.

### 11.2. HDL and the Clearance of LOP

According to the conception of lipoprotein-specific LOP transport, HDL-assisted reverse transport carries LOP from peripheral veins and tissues towards the liver. This conclusion is based on studies showing that physical exercise affects the LDL-LOP/HDL-LOP balance in such a way that the net flow of LOP shifts towards the liver. Together with previous observations of capability of the liver to remove and eliminate LOP in HDL, and results from studies with apoA1-mimetic peptides, these findings suggest a protective LOP clearing function for HDL. This is an interesting new perspective for the discussion of functionality of HDL. HDL-LOP is also reported to be associated with a wide spectrum of atherosclerosis risk factors. In most cases these associations are opposite to those of LDL-LOP, giving further support for the protective role of HDL.

## 12. Significance for Other Oxidative Stress-Dependent Diseases

As carriers of LOP in circulation, lipoproteins likely contribute to the pro/antioxidant balance in the body. This raises the question whether the lipoprotein –specific transport of LOP would be implicated in oxidative stress –dependent pathological processes other than atherosclerosis. Yet, it must be kept in mind that there are no common criteria for “oxidative stress –dependent” diseases and/or pathological processes, and it may sometimes be difficult to differentiate whether oxidative stress is a cause or consequence of the disease process.

In a previous chapter (*Metabolic disorders related to atherosclerosis*) data was presented of associations of lipoprotein LOP with the development of fatty liver, which also associates with oxidative stress [98]. Similarly, data was presented from connection between lipoprotein LOP and insulin sensitivity. Lipoprotein LOP was found to be involved in both these pathophysiological situations. Besides these reported studies, there appear to be no investigations on lipoprotein LOP in diseases or pathological processes in which oxidative stress is implicated.

Type 2 diabetes, which is a major worldwide public health problem today, is an example of diseases in which the contribution of lipoprotein LOP transport would be worth investigating. The involvement of oxidative stress in type 2 diabetes has been demonstrated by biomarkers of oxidative lipid, protein, and nucleic acid damage [99]. Another disease of emerging worldwide concern, chronic kidney disease (CKD), could also be of interest in this respect. Oxidative stress and inflammation are important contributors in the pathogenesis and progression of CKD [100–102]. In both these diseases dyslipemia, too, appears to play an important role [103–105]. It is possible that together with oxidative stress, unfavorable lipoprotein levels could enhance development of these pathophysiological conditions.

### 13. Perspective Regarding Prevention and Treatment

If the lipoprotein-specific transport of LOP would play a role in development of atherosclerosis, then strategies for prevention and treatment should aim at low LDL-LOP/HDL-LOP ratio by lowering LDL-LOP and/or elevating HDL-LOP levels. As discussed in a previous chapter, caloric restriction, physically active lifestyle, and statin treatment have been shown to improve LDL-LOP effectively. Therefore, since these current strategies for a healthier lipid profile also decrease the potential risk due to LOP, the role of LOP as a risk factor would hardly radically change the management of atherosclerosis and CVD.

On the other hand, as it has turned out that the amount and composition of circulating LOP, especially during the postprandial period, is largely determined by ingested food, LOP in food could be regarded as a new type of dietary risk factor. Shifting the focus from cholesterol toward LOP could therefore give additional tools for prevention and therapy. This could mean in the future e.g. reporting of the presence of LOP in foodstuffs, similarly to the present practice for the content and quality of fat.

In line with the postulated role of LOP as a risk factor, epidemiological studies show that low levels of dietary antioxidants are associated with increased risk for cardiovascular disease and that increased intakes are protective [106]. Yet, antioxidant supplementation is not an answer since, as discussed earlier (in *Dietary factors*), it does not affect the concentration of LDL-LOP even though it substantially increases the antioxidant capacity of LDL. This is well in agreement with the poor success of antioxidant strategies in limiting atherosclerosis and CVD [12], and supports the idea of dietary origin of LOP in LDL.

In recent years the development of new strategies for prevention and treatment of atherosclerosis has increasingly focused on HDL and its functions [15,16]. Despite huge investments on research and drug development projects, attempts to reinforce and exploit protective effects of HDL as clinical applications have turned out to be challenging. As for the LOP transport by HDL, there is little information concerning possibilities to enhance the function. It has been reported that 6 months supervised physical activity program increased the concentration of HDL-LOP and the HDL-LOP/HDL -cholesterol ratio among elderly women [66]. No doubt, a deeper understanding of factors specifically determining the transport capacity is needed before it is possible to modulate this function.

Since the studies of LOP transport function are based on measurement of LOP in lipoproteins, it would be worth considering whether assays for lipoprotein LOP could form basis for new diagnostic applications. Circulating LDL-LOP have been measured over the past 25 years in a variety of cross-sectional, intervention, and population-based studies, and it appeared early that LDL-LOP is an applicable indicator of the risk of atherosclerosis [107]. In these studies, high LDL-LOP has

consistently pointed to increased risk of the disease (See *Lipoprotein-specific transport of LOP and the risk of atherosclerosis*).

The assay for HDL-LOP was introduced much later, but in studies thus far low HDL-LOP level has been found to associate with an elevated risk. The ratio of LDL-LOP/HDL-LOP, as a putative indicator of the direction of the net movement of circulating LOP, has proved to be potentially interesting and worth further investigation. Even though available evidence supports the usability of lipoprotein LOP as an indicator of the risk of atherosclerosis and CVD, large-scale epidemiological studies are needed for verification. In addition, the present methodologies need to be developed to meet the requirements of modern clinical laboratories. Malondialdehyde (MDA) and other reactive LOP are known to form adducts on proteins or lipids of lipoproteins [12,108], which can serve as indicators of oxidative stress. Yet, due to firm attachment on structural components of lipoproteins they are not indicative of the LOP transport function. Similarly, the common assay for oxidized LDL, which is a measure of oxidation –dependent aldehyde substitution of lysine residues in apolipoprotein B, is indicative of oxidative stress but not of the LOP transport function.

#### 14. Consistence with Present Theories

How does the lipoprotein-specific transport of LOP comport with present view of the etiology of atherosclerosis? Various theories have been developed to explain etiology of the disease, each of them emphasizing a specific part in the processes of atherogenesis [12]. Since the discoveries by Goldstein and Brown [109], the role of lipoproteins as risk factors for atherosclerosis has been explained by cholesterol transport function. More recently, the oxidation theory of atherosclerosis and the oxidative response to inflammation –hypothesis [12] stressing the role of monocytes in subendothelial space and the ensuing vascular inflammation, have further shaped the view.

The proposed role of the LOP transport function does not conflict with any of these theories, and rather underlines the significance of both lipoprotein transport and oxidation. Moreover, since LOP are implicated in accumulation of monocytes in the subendothelial space, the hypothesis of the role of lipoprotein LOP transport is compatible also with the oxidative response to inflammation – hypothesis. Taken together, the proposed role of the lipoprotein-specific LOP transport by large agrees with previously presented hypotheses, and, in fact, not only expand but also combine previous theories in a rational way. The only deviation from the previously presented theories concerns origin of the “oxidized LDL”. Thus far it has been assumed that “oxidized LDL” is generated within the vessel wall in reactions caused by endogenous free radicals [12]. As an addition to this, the theory emphasizing role of lipoprotein LOP transport proposes that part of lipoprotein LOP would be of dietary origin. This argument is reasoned by evidence from the dietary studies with fatty meals.

#### 15. Open Questions and Future Directions

The fact that life-style dependent factors affect in a parallel way on both cholesterol and LOP levels suggests that the lipoprotein-specific LOP transport at least in part works under the same regulatory control as cholesterol transport. Unlike in case of cholesterol, there is at present no information regarding influence of LOP on regulation of lipoprotein metabolism.

The molecular mechanisms of cholesterol transfer from cells and tissues to HDL, between lipoproteins, and from lipoproteins to liver are today well known, but is not known whether the same mechanisms apply also for LOP. To increase credibility of the lipoprotein-specific LOP transport – hypothesis, it would be important to know the molecular mechanisms of LOP transfer at various steps of the transportation. In the current situation it is not known whether drug development for modulation of lipoprotein LOP levels could follow the same paths as in case of cholesterol. Increasing the knowledge would help to find out whether there are ways to improve the situation beyond strategies that are in use in cholesterol-targeted treatment protocols.

The idea of the lipoprotein-specific LOP transport is rather young, and therefore more evidence based on research is needed. An advancement that could help to boost the development would be availability of methodology that is suitable for large-scale epidemiological studies and applicable in modern clinical laboratories.

The multiplicity of LOP, which are an assortment of chemicals with various structures, forms a challenge for attempts to control possible risks. Diverse constituents of LOP presumably differ from each other regarding their biological effects, and research should therefore be directed to characterization of the role of individual chemicals and chemical groups and within the LOP.

Another place where chemical analytics will be needed is monitoring of LOP in foodstuffs. Reduction of dietary intake of LOP will not be possible without knowledge of the presence of LOP in dietary components. Thus far, data concerning LOP in food have come out randomly with no aspiration to a systematical survey.

## 16. Conclusions

The evidence connecting lipoprotein LOP transport to atherosclerosis is versatile and consistent revealing surprisingly strong connections between LOP transport and the risk of atherosclerotic CVD. If confirmed, the proposed LOP transport function would open a new and interesting viewpoint to study of atherosclerosis. Importantly, the theory of lipoprotein-specific LOP transport is in full agreement with the existing theories of etiology of atherosclerosis and combines these in a rational way.

The data presented in this review show that effects of lipoprotein LOP as a potential risk factor are largely parallel with those of cholesterol. However, due to common transport and cellular intake mechanisms, elucidation of independent effects of cholesterol and LOP is difficult. Keeping in mind that LOP, unlike cholesterol, can directly initiate and boost atherogenic processes, it appears likely that increasing knowledge of the lipoprotein-specific LOP transport will affect the present view of the role of cholesterol in etiology and as a risk factor of atherosclerosis.

If confirmed, the proposed lipoprotein LOP transport function would open new perspectives regarding the risk of CVD, such as the dietary origin of LOP, and the protective function of HDL in clearance of LOP. Focusing on LOP could give additional tools especially for prevention and diagnosis but would hardly radically change the management of atherosclerosis and CVD.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Feingold, K. R. Introduction to lipids and lipoproteins. In *Endotext*; Feingold, K. R., Anawalt, B., Blackman, M. R., Boyce, A., Chrousos, G., Corpas, E., de Herder, W. W., Dhatariya, K., Dungan, K., Hofland, J., Kalra, S., Kalsas, G., Kapoor, N., Koch, C., Kopp, P., Korbonits, M., Kovacs, C. S., Kuohung, W., Laferrère, B., Levy, M., McGee, E. A., McLachlan, R., New, M., Purnell, J., Sahay, R., Shah, A. S., Singer, F., Sperling, M. A., Stratakis, C. A., Trencé, D. L., Wilson, D. P., Eds.; MDText.com, Inc.: South Dartmouth (MA), 2000.
2. Steinberg, D.; Parthasarathy, S.; Carew, T. E.; Khoo, J. C.; Witztum, J. L. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med* 1989, 320 (14), 915–924. <https://doi.org/10.1056/NEJM198904063201407>.
3. Bowry, V. W.; Stanley, K. K.; Stocker, R. High density lipoprotein is the major carrier of lipid hydroperoxides in human blood plasma from fasting donors. *Proc. Natl. Acad. Sci. U.S.A.* 1992, 89 (21), 10316–10320. <https://doi.org/10.1073/pnas.89.21.10316>.
4. Esterbauer, H.; Gebicki, J.; Puhl, H.; Jürgens, G. The role of lipid peroxidation and antioxidants in oxidative modification of LDL. *Free Radical Biology and Medicine* 1992, 13 (4), 341–390. [https://doi.org/10.1016/0891-5849\(92\)90181-F](https://doi.org/10.1016/0891-5849(92)90181-F).
5. Staprans, I.; Rapp, J. H.; Pan, X. M.; Feingold, K. R. The effect of oxidized lipids in the diet on serum lipoprotein peroxides in control and diabetic rats. *J. Clin. Invest.* 1993, 92 (2), 638–643. <https://doi.org/10.1172/JCI116632>.
6. Staprāns, I.; Rapp, J. H.; Pan, X. M.; Kim, K. Y.; Feingold, K. R. Oxidized lipids in the diet are a source of oxidized lipid in chylomicrons of human serum. *Arterioscler Thromb* 1994, 14 (12), 1900–1905. <https://doi.org/10.1161/01.ATV.14.12.1900>.
7. Staprans, I.; Rapp, J. H.; Pan, X. M.; Feingold, K. R. Oxidized lipids in the diet are incorporated by the liver into very low density lipoprotein in rats. *J Lipid Res* 1996, 37 (2), 420–430.

8. Ahotupa, M. Oxidized lipoprotein lipids and atherosclerosis. *Free Radical Research* 2017, 51 (4), 439–447. <https://doi.org/10.1080/10715762.2017.1319944>.
9. Shao, B.; Heinecke, J. W. HDL, lipid peroxidation, and atherosclerosis. *Journal of Lipid Research* 2009, 50 (4), 599–601. <https://doi.org/10.1194/jlr.E900001-JLR200>.
10. Ahotupa, M.; Suomela, J.-P.; Vuorimaa, T.; Vasankari, T. Lipoprotein-specific transport of circulating lipid peroxides. *Annals of Medicine* 2010, 42 (7), 521–529. <https://doi.org/10.3109/07853890.2010.510932>.
11. Soran, H.; Schofield, J. D.; Durrington, P. N. Antioxidant properties of HDL. *Front. Pharmacol.* 2015, 6. <https://doi.org/10.3389/fphar.2015.00222>.
12. Stocker, R.; Keaney, J. F. Role of oxidative modifications in atherosclerosis. *Physiological Reviews* 2004, 84 (4), 1381–1478. <https://doi.org/10.1152/physrev.00047.2003>.
13. Steinbrecher, U. Receptors for oxidized low density lipoprotein. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* 1999, 1436 (3), 279–298. [https://doi.org/10.1016/S0005-2760\(98\)00127-1](https://doi.org/10.1016/S0005-2760(98)00127-1).
14. Miller, Y. I.; Shyy, J. Y.-J. Context-dependent role of oxidized lipids and lipoproteins in inflammation. *Trends in Endocrinology & Metabolism* 2017, 28 (2), 143–152. <https://doi.org/10.1016/j.tem.2016.11.002>.
15. Von Eckardstein, A.; Nordestgaard, B. G.; Remaley, A. T.; Catapano, A. L. High-density lipoprotein revisited: biological functions and clinical relevance. *European Heart Journal* 2023, 44 (16), 1394–1407. <https://doi.org/10.1093/eurheartj/ehac605>.
16. Asztalos, B. F.; Tani, M.; Schaefer, E. J. Metabolic and functional relevance of HDL subspecies. *Current Opinion in Lipidology* 2011, 22 (3), 176–185. <https://doi.org/10.1097/MOL.0b013e3283468061>.
17. Ben-Aicha, S.; Badimon, L.; Vilahur, G. Advances in HDL: much more than lipid transporters. *IJMS* 2020, 21 (3), 732. <https://doi.org/10.3390/ijms21030732>.
18. Khera, A. V.; Cuchel, M.; De La Llera-Moya, M.; Rodrigues, A.; Burke, M. F.; Jafri, K.; French, B. C.; Phillips, J. A.; Mucksavage, M. L.; Wilensky, R. L.; Mohler, E. R.; Rothblat, G. H.; Rader, D. J. Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. *N Engl J Med* 2011, 364 (2), 127–135. <https://doi.org/10.1056/NEJMoa1001689>.
19. Spindler-Vomachka, M.; Vodcnik, M. J.; Lech, J. J. Transport of 2,4,5,2',4',5'-Hexachlorobiphenyl by lipoproteins in vivo. *Toxicology and Applied Pharmacology* 1984, 74 (1), 70–77. [https://doi.org/10.1016/0041-008X\(84\)90271-0](https://doi.org/10.1016/0041-008X(84)90271-0).
20. Borlakoglu, J. T.; Welch, V. A.; Wilkins, J. P. G.; Dils, R. R. Transport and cellular uptake of polychlorinated biphenyls (PCBs)—I. *Biochemical Pharmacology* 1990, 40 (2), 265–272. [https://doi.org/10.1016/0006-2952\(90\)90687-G](https://doi.org/10.1016/0006-2952(90)90687-G).
21. Lamon-Fava, S.; Sadowski, J.; Davidson, K.; O'Brien, M.; McNamara, J.; Schaefer, E. Plasma lipoproteins as carriers of phyloquinone (vitamin K1) in humans. *The American Journal of Clinical Nutrition* 1998, 67 (6), 1226–1231. <https://doi.org/10.1093/ajcn/67.6.1226>.
22. Meng, Q.-H.; Wähälä, K.; Adlercreutz, H.; Tikkanen, M. J. Antiproliferative efficacy of lipophilic soy isoflavone phytoestrogens delivered by low density lipoprotein particles into cultured U937 cells. *Life Sciences* 1999, 65 (16), 1695–1705. [https://doi.org/10.1016/S0024-3205\(99\)00418-X](https://doi.org/10.1016/S0024-3205(99)00418-X).
23. Gershkovich, P.; Hoffman, A. Effect of a high-fat meal on absorption and disposition of lipophilic compounds: the importance of degree of association with triglyceride-rich lipoproteins. *European Journal of Pharmaceutical Sciences* 2007, 32 (1), 24–32. <https://doi.org/10.1016/j.ejps.2007.05.109>.
24. Li, Y.; Wongsiriroj, N.; Blaner, W. S. The multifaceted nature of retinoid transport and metabolism. *Hepatobiliary Surg Nutr* 2014, 3 (3), 126–139. <https://doi.org/10.3978/j.issn.2304-3881.2014.05.04>.
25. Yamamoto, H.; Takada, T.; Yamanashi, Y.; Ogura, M.; Masuo, Y.; Harada-Shiba, M.; Suzuki, H. VLDL/LDL acts as a drug carrier and regulates the transport and metabolism of drugs in the body. *Sci Rep* 2017, 7 (1), 633. <https://doi.org/10.1038/s41598-017-00685-9>.
26. Saito, Y.; Noguchi, N. Oxidized lipoprotein as a major vessel cell proliferator in oxidized human serum. *PLoS ONE* 2016, 11 (8), e0160530. <https://doi.org/10.1371/journal.pone.0160530>.
27. Zheng, A.; Dubuis, G.; Georgieva, M.; Mendes Ferreira, C. S.; Serulla, M.; Del Carmen Conde Rubio, M.; Trofimenko, E.; Mercier, T.; Decosterd, L.; Widmann, C. HDLs extract lipophilic drugs from cells. *Journal of Cell Science* 2022, 135 (5), jcs258644. <https://doi.org/10.1242/jcs.258644>.
28. Negre-Salvayre, A.; Coatrieux, C.; Ingueneau, C.; Salvayre, R. Advanced lipid peroxidation end products in oxidative damage to proteins. Potential role in diseases and therapeutic prospects for the inhibitors. *British J Pharmacology* 2008, 153 (1), 6–20. <https://doi.org/10.1038/sj.bjp.0707395>.
29. Vistoli, G.; De Maddis, D.; Cipak, A.; Zarkovic, N.; Carini, M.; Aldini, G. Advanced glycooxidation and lipoxidation end products (AGEs and ALEs): An overview of their mechanisms of formation. *Free Radical Research* 2013, 47 (sup1), 3–27. <https://doi.org/10.3109/10715762.2013.815348>.
30. Halliwell, B.; Gutteridge, J. M. C. *Free Radicals in Biology and Medicine*, 3rd ed.; Oxford science publications; Clarendon Press ; Oxford University Press: Oxford : New York, 1999.

31. Fisher-Wellman, K.; Bloomer, R. J. Acute exercise and oxidative stress: A 30 year history. *Dyn Med* 2009, 8 (1), 1. <https://doi.org/10.1186/1476-5918-8-1>.
32. Vasankari, T.; Kujala, U.; Heinonen, O.; Kapanen, J.; Ahotupa, M. Measurement of serum lipid peroxidation during exercise using three different methods: diene conjugation, thiobarbituric acid reactive material and fluorescent chromolipids. *Clin Chim Acta* 1995, 234 (1–2), 63–69. [https://doi.org/10.1016/0009-8981\(94\)05976-y](https://doi.org/10.1016/0009-8981(94)05976-y).
33. Suomela, J.; Ahotupa, M.; Kallio, H. Triacylglycerol oxidation in pig lipoproteins after a diet rich in oxidized sunflower seed oil. *Lipids* 2005, 40 (5), 437–444. <https://doi.org/10.1007/s11745-005-1402-4>.
34. Wolff, S. P.; Nourooz-Zadeh, J. Hypothesis: UK consumption of dietary lipid hydroperoxides — a possible contributory factor to atherosclerosis. *Atherosclerosis* 1996, 119 (2), 261–263. [https://doi.org/10.1016/0021-9150\(95\)05640-8](https://doi.org/10.1016/0021-9150(95)05640-8).
35. Tarvainen, M.; Phuphusit, A.; Suomela, J.-P.; Kuksis, A.; Kallio, H. Effects of antioxidants on rapeseed oil oxidation in an artificial digestion model analyzed by UHPLC-ESI-MS. *J Agric Food Chem* 2012, 60 (14), 3564–3579. <https://doi.org/10.1021/jf2050944>.
36. Nuora, A.; Chiang, V. S.-C.; Milan, A. M.; Tarvainen, M.; Pundir, S.; Quek, S.-Y.; Smith, G. C.; Markworth, J. F.; Ahotupa, M.; Cameron-Smith, D.; Linderborg, K. M. The impact of beef steak thermal processing on lipid oxidation and postprandial inflammation related responses. *Food Chemistry* 2015, 184, 57–64. <https://doi.org/10.1016/j.foodchem.2015.03.059>.
37. Tiainen, S.; Ahotupa, M.; Ylinen, P.; Vasankari, T. High density lipoprotein level is negatively associated with the increase of oxidized low density lipoprotein lipids after a fatty meal. *Lipids* 2014, 49 (12), 1225–1232. <https://doi.org/10.1007/s11745-014-3963-y>.
38. Cortés, B.; Núñez, I.; Cofán, M.; Gilabert, R.; Pérez-Heras, A.; Casals, E.; Deulofeu, R.; Ros, E. Acute effects of high-fat meals enriched with walnuts or olive oil on postprandial endothelial function. *Journal of the American College of Cardiology* 2006, 48 (8), 1666–1671. <https://doi.org/10.1016/j.jacc.2006.06.057>.
39. Välimäki, I.; Vuorimaa, T.; Ahotupa, M.; Vasankari, T. Effect of continuous and intermittent exercises on oxidised HDL and LDL lipids in runners. *Int J Sports Med* 2016, 37 (14), 1103–1109. <https://doi.org/10.1055/s-0042-114703>.
40. Välimäki, I. A.; Vuorimaa, T.; Ahotupa, M.; Vasankari, T. J. Strenuous physical exercise accelerates the lipid peroxide clearing transport by HDL. *Eur J Appl Physiol* 2016, 116 (9), 1683–1691. <https://doi.org/10.1007/s00421-016-3422-y>.
41. Van Lenten, B. J.; Wagner, A. C.; Anantharamaiah, G. M.; Navab, M.; Reddy, S. T.; Buga, G. M.; Fogelman, A. M. Apolipoprotein A-I mimetic peptides. *Curr Atheroscler Rep* 2009, 11 (1), 52–57. <https://doi.org/10.1007/s11883-009-0008-8>.
42. Imaizumi, S.; Grijalva, V.; Navab, M.; J. Van Lenten, B.; C. Wagner, A.; Anantharamaiah, G. M.; M. Fogelman, A.; T. Reddy, S. L-4F differentially alters plasma levels of oxidized fatty acids resulting in more anti-inflammatory HDL in mice. *DML* 2010, 4 (3), 139–148. <https://doi.org/10.2174/187231210791698438>.
43. Morgantini, C.; Imaizumi, S.; Grijalva, V.; Navab, M.; Fogelman, A. M.; Reddy, S. T. Apolipoprotein A-I mimetic peptides prevent atherosclerosis development and reduce plaque inflammation in a murine model of diabetes. *Diabetes* 2010, 59 (12), 3223–3228. <https://doi.org/10.2337/db10-0844>.
44. Elsäe, S.; Ahnström, J.; Christoffersen, C.; Hoofnagle, A. N.; Plomgaard, P.; Heinecke, J. W.; Binder, C. J.; Björkbacka, H.; Dahlbäck, B.; Nielsen, L. B. Apolipoprotein M binds oxidized phospholipids and increases the antioxidant effect of HDL. *Atherosclerosis* 2012, 221 (1), 91–97. <https://doi.org/10.1016/j.atherosclerosis.2011.11.031>.
45. Christison, J.; Karjalainen, A.; Brauman, J.; Bygrave, F.; Stocker, R. Rapid reduction and removal of HDL-but not LDL-associated cholesteryl ester hydroperoxides by rat liver perfused *in situ*. *Biochemical Journal* 1996, 314 (3), 739–742. <https://doi.org/10.1042/bj3140739>.
46. Fluiter, K.; Vietsch, H.; Biessen, E. A.; Kostner, G. M.; van Berkel, T. J.; Sattler, W. Increased selective uptake in vivo and in vitro of oxidized cholesteryl esters from high-density lipoprotein by rat liver parenchymal cells. *Biochem J* 1996, 319 ( Pt 2) (Pt 2), 471–476. <https://doi.org/10.1042/bj3190471>.
47. Nourooz-Zadeh, J. Measures of oxidative stress in heterozygous familial hypercholesterolaemia. *Atherosclerosis* 2001, 156 (2), 435–441. [https://doi.org/10.1016/S0021-9150\(00\)00677-8](https://doi.org/10.1016/S0021-9150(00)00677-8).
48. Mollazadeh, H.; Carbone, F.; Montecucco, F.; Pirro, M.; Sahebkar, A. Oxidative burden in familial hypercholesterolemia. *Journal Cellular Physiology* 2018, 233 (8), 5716–5725. <https://doi.org/10.1002/jcp.26466>.
49. Shahsavari, G.; Raoufi, A.; Toolabi, A.; Hosseninejadmir, N.; Ahmadvand, H.; Safariebrahimsarabie, M. The Effect of atorvastatin treatment duration on oxidative stress markers and lipid profile in patients with coronary artery diseases: a case series study. *ARYA Atheroscler* 2017, 13 (6), 282–287.

50. Sørensen, A. L.; Hasselbalch, H. C.; Nielsen, C. H.; Poulsen, H. E.; Ellervik, C. Statin treatment, oxidative stress and inflammation in a danish population. *Redox Biology* 2019, 21, 101088. <https://doi.org/10.1016/j.redox.2018.101088>.
51. Rosenson, R. S.; Brewer, H. B.; Ansell, B.; Barter, P.; Chapman, M. J.; Heinecke, J. W.; Kontush, A.; Tall, A. R.; Webb, N. R. Translation of high-density lipoprotein function into clinical practice: current prospects and future challenges. *Circulation* 2013, 128 (11), 1256–1267. <https://doi.org/10.1161/CIRCULATIONAHA.113.000962>.
52. Ahotupa, M.; Marniemi, J.; Lehtimäki, T.; Talvinen, K.; Raitakari, O. T.; Vasankari, T.; Viikari, J.; Luoma, J.; Ylä-Herttuala, S. Baseline diene conjugation in LDL lipids as a direct measure of in vivo LDL oxidation. *Clinical Biochemistry* 1998, 31 (4), 257–261. [https://doi.org/10.1016/S0009-9120\(98\)00018-6](https://doi.org/10.1016/S0009-9120(98)00018-6).
53. Kresanov, P.; Ahotupa, M.; Vasankari, T.; Kaikkonen, J.; Kähönen, M.; Lehtimäki, T.; Viikari, J.; Raitakari, O. T. The associations of oxidized high-density lipoprotein lipids with risk factors for atherosclerosis: The Cardiovascular Risk in Young Finns Study. *Free Radical Biology and Medicine* 2013, 65, 1284–1290. <https://doi.org/10.1016/j.freeradbiomed.2013.09.023>.
54. Sniderman, A. D.; Islam, S.; McQueen, M.; Pencina, M.; Furberg, C. D.; Thanassoulis, G.; Yusuf, S. Age and cardiovascular risk attributable to apolipoprotein B, low-density lipoprotein cholesterol or non-high-density lipoprotein cholesterol. *J Am Heart Assoc* 2016, 5 (10), e003665. <https://doi.org/10.1161/JAHA.116.003665>.
55. Kujala, U. M.; Ahotupa, M.; Vasankari, T.; Kaprio, J.; Tikkanen, M. J. Low LDL oxidation in veteran endurance athletes. *Scandinavian Med Sci Sports* 1996, 6 (5), 303–308. <https://doi.org/10.1111/j.1600-0838.1996.tb00475.x>.
56. Kujala, U. M.; Ahotupa, M.; Vasankari, T. J.; Kaprio, J.; Tikkanen, M. J. Familial aggregation of LDL oxidation. *Scandinavian Journal of Clinical and Laboratory Investigation* 1997, 57 (2), 141–146. <https://doi.org/10.1080/00365519709056382>.
57. Vasankari, T.; Ahotupa, M.; Toikka, J.; Mikkola, J.; Irjala, K.; Pasanen, P.; Neuvonen, K.; Raitakari, O.; Viikari, J. Oxidized LDL and thickness of carotid intima-media are associated with coronary atherosclerosis in middle-aged men: lower levels of oxidized LDL with statin therapy. *Atherosclerosis* 2001, 155 (2), 403–412. [https://doi.org/10.1016/S0021-9150\(00\)00573-6](https://doi.org/10.1016/S0021-9150(00)00573-6).
58. Toikka, J. O.; Laine, H.; Ahotupa, M.; Haapanen, A.; Viikari, J. S. A.; Hartiala, J. J.; Raitakari, O. T. Increased arterial intima-media thickness and in vivo LDL oxidation in young men with borderline hypertension. *Hypertension* 2000, 36 (6), 929–933. <https://doi.org/10.1161/01.HYP.36.6.929>.
59. Linna, M. S.; Ahotupa, M.; Irjala, K.; Pöllänen, P.; Huhtaniemi, I.; Mäkinen, J.; Perheentupa, A.; Vasankari, T. J. Smoking and low serum testosterone associates with high concentration of oxidized LDL. *Annals of Medicine* 2008, 40 (8), 634–640. <https://doi.org/10.1080/07853890802161007>.
60. Li, J.; Siegrist, J. Physical activity and risk of cardiovascular disease—a meta-analysis of prospective cohort studies. *IJERPH* 2012, 9 (2), 391–407. <https://doi.org/10.3390/ijerph9020391>.
61. Martin, S. S.; Aday, A. W.; Almarzooq, Z. I.; Anderson, C. A. M.; Arora, P.; Avery, C. L.; Baker-Smith, C. M.; Barone Gibbs, B.; Beaton, A. Z.; Boehme, A. K.; Commodore-Mensah, Y.; Currie, M. E.; Elkind, M. S. V.; Evenson, K. R.; Generoso, G.; Heard, D. G.; Hiremath, S.; Johansen, M. C.; Kalani, R.; Kazi, D. S.; Ko, D.; Liu, J.; Magnani, J. W.; Michos, E. D.; Mussolino, M. E.; Navaneethan, S. D.; Parikh, N. I.; Perman, S. M.; Poudel, R.; Rezk-Hanna, M.; Roth, G. A.; Shah, N. S.; St-Onge, M.-P.; Thacker, E. L.; Tsao, C. W.; Urbut, S. M.; Van Spall, H. G. C.; Voeks, J. H.; Wang, N.-Y.; Wong, N. D.; Wong, S. S.; Yaffe, K.; Palaniappan, L. P.; on behalf of the American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. 2024 Heart disease and stroke statistics: a report of US and global data from the American Heart Association. *Circulation* 2024, CIR.0000000000001209. <https://doi.org/10.1161/CIR.0000000000001209>.
62. Vasankari, T.; Lehtonen-Veromaa, M.; Möttönen, T.; Ahotupa, M.; Irjala, K.; Heinonen, O.; Leino, A.; Viikari, J. Reduced mildly oxidized LDL in young female athletes. *Atherosclerosis* 2000, 151 (2), 399–405. [https://doi.org/10.1016/S0021-9150\(99\)00401-3](https://doi.org/10.1016/S0021-9150(99)00401-3).
63. Vasankari, T. J.; Kujala, U. M.; Vasankari, T. M.; Ahotupa, M. Reduced oxidized LDL levels after a 10-month exercise program. *Medicine & Science in Sports & Exercise* 1998, 30 (10), 1496–1501. <https://doi.org/10.1097/00005768-199810000-00005>.
64. Välimäki, I. A.; Vuorimaa, T.; Ahotupa, M.; Kekkonen, R.; Korpela, R.; Vasankari, T. Decreased training volume and increased carbohydrate intake increases oxidized LDL levels. *Int J Sports Med* 2012, 33 (4), 291–296. <https://doi.org/10.1055/s-0031-1291223>.
65. Vuorimaa, T.; Ahotupa, M.; Irjala, K.; Vasankari, T. Acute prolonged exercise reduces moderately oxidized LDL in healthy men. *Int J Sports Med* 2005, 26 (6), 420–425. <https://doi.org/10.1055/s-2004-821142>.

66. Tiainen, S.; Luoto, R.; Ahotupa, M.; Raitanen, J.; Vasankari, T. 6-mo aerobic exercise intervention enhances the lipid peroxide transport function of HDL. *Free Radical Research* 2016, *50* (11), 1279–1285. <https://doi.org/10.1080/10715762.2016.1252040>.
67. Vasankari, T. J.; Kujala, U. M.; Vasankari, T. M.; Vuorimaa, T.; Ahotupa, M. Effects of acute prolonged exercise on serum and LDL oxidation and antioxidant defences. *Free Radical Biology and Medicine* 1997, *22* (3), 509–513. [https://doi.org/10.1016/S0891-5849\(96\)00373-5](https://doi.org/10.1016/S0891-5849(96)00373-5).
68. Vasankari, T.; Kujala, U.; Vasankari, T.; Vuorimaa, T.; Ahotupa, M. Increased serum and low-density lipoprotein antioxidant potential after antioxidant supplementation in endurance athletes. *The American Journal of Clinical Nutrition* 1997, *65* (4), 1052–1056. <https://doi.org/10.1093/ajcn/65.4.1052>.
69. Hubert, H. B.; Feinleib, M.; McNamara, P. M.; Castelli, W. P. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 1983, *67* (5), 968–977. <https://doi.org/10.1161/01.CIR.67.5.968>.
70. Calle, E. E.; Thun, M. J.; Petrelli, J. M.; Rodriguez, C.; Heath, C. W. Body mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med* 1999, *341* (15), 1097–1105. <https://doi.org/10.1056/NEJM199910073411501>.
71. Sorokin, A. V.; Kotani, K.; Elnabawi, Y. A.; Dey, A. K.; Sajja, A. P.; Yamada, S.; Ueda, M.; Harrington, C. L.; Baumer, Y.; Rodante, J. A.; Gelfand, J. M.; Chen, M. Y.; Joshi, A. A.; Playford, M. P.; Remaley, A. T.; Mehta, N. N. Association between oxidation-modified lipoproteins and coronary plaque in psoriasis: an observational cohort study. *Circ Res* 2018, *123* (11), 1244–1254. <https://doi.org/10.1161/CIRCRESAHA.118.313608>.
72. Vasankari, T.; Fogelholm, M.; Kukkonen-Harjula, K.; Nenonen, A.; Kujala, U.; Oja, P.; Vuori, I.; Pasanen, P.; Neuvonen, K.; Ahotupa, M. Reduced oxidized low-density lipoprotein after weight reduction in obese premenopausal women. *Int J Obes* 2001, *25* (2), 205–211. <https://doi.org/10.1038/sj.ijo.0801533>.
73. Linna, M. S.; Borg, P.; Kukkonen-Harjula, K.; Fogelholm, M.; Nenonen, A.; Ahotupa, M.; Vasankari, T. J. Successful weight maintenance preserves lower levels of oxidized LDL achieved by weight reduction in obese men. *Int J Obes* 2007, *31* (2), 245–253. <https://doi.org/10.1038/sj.ijo.0803413>.
74. Raitakari, M.; Ilvonen, T.; Ahotupa, M.; Lehtimäki, T.; Harmoinen, A.; Suominen, P.; Elo, J.; Hartiala, J.; Raitakari, O. T. Weight reduction with very-low-caloric diet and endothelial function in overweight adults: role of plasma glucose. *Arterioscler Thromb Vasc Biol* 2004, *24* (1), 124–128. <https://doi.org/10.1161/01.ATV.0000109749.11042.7c>.
75. Lee, C. D.; Blair, S. N.; Jackson, A. S. Cardiorespiratory fitness, body composition, and all-cause and cardiovascular disease mortality in men. *The American Journal of Clinical Nutrition* 1999, *69* (3), 373–380. <https://doi.org/10.1093/ajcn/69.3.373>.
76. Kosola, J.; Ahotupa, M.; Kyröläinen, H.; Santtila, M.; Vasankari, T. Good aerobic or muscular fitness protects overweight men from elevated oxidized LDL. *Medicine & Science in Sports & Exercise* 2012, *44* (4), 563–568. <https://doi.org/10.1249/MSS.0b013e31823822cc>.
77. Kaikkonen, J. E.; Kresanov, P.; Ahotupa, M.; Jula, A.; Mikkilä, V.; Viikari, J. S. A.; Kähönen, M.; Lehtimäki, T.; Raitakari, O. T. High serum n6 fatty acid proportion is associated with lowered LDL oxidation and inflammation: The Cardiovascular Risk in Young Finns Study. *Free Radical Research* 2014, *48* (4), 420–426. <https://doi.org/10.3109/10715762.2014.883071>.
78. Harris, W. S.; Mozaffarian, D.; Rimm, E.; Kris-Etherton, P.; Rudel, L. L.; Appel, L. J.; Engler, M. M.; Engler, M. B.; Sacks, F. Omega-6 fatty acids and risk for cardiovascular disease: a science advisory from the American Heart Association Nutrition Subcommittee of the Council on Nutrition, Physical Activity, and Metabolism; Council on Cardiovascular Nursing; and Council on Epidemiology and Prevention. *Circulation* 2009, *119* (6), 902–907. <https://doi.org/10.1161/CIRCULATIONAHA.108.191627>.
79. Jula, A.; Marniemi, J.; Huupponen, R.; Virtanen, A.; Rastas, M.; Rönnemaa, T. Effects of diet and simvastatin on serum lipids, insulin, and antioxidants in hypercholesterolemic men: a randomized controlled trial. *JAMA* 2002, *287* (5), 598. <https://doi.org/10.1001/jama.287.5.598>.
80. Agarwal, S.; Rao, A. V. Tomato lycopene and low density lipoprotein oxidation: a human dietary intervention study. *Lipids* 1998, *33* (10), 981–984. <https://doi.org/10.1007/s11745-998-0295-6>.
81. Jenkins, D. J. A.; Kendall, C. W. C.; Marchie, A.; Parker, T. L.; Connelly, P. W.; Qian, W.; Haight, J. S.; Faulkner, D.; Vidgen, E.; Lapsley, K. G.; Spiller, G. A. Dose response of almonds on coronary heart disease risk factors: blood lipids, oxidized low-density lipoproteins, lipoprotein(a), homocysteine, and pulmonary nitric oxide: a randomized, controlled, crossover trial. *Circulation* 2002, *106* (11), 1327–1332. <https://doi.org/10.1161/01.CIR.0000028421.91733.20>.
82. Marniemi, J.; Hakala, P.; Mäki, J.; Ahotupa, M. Partial resistance of low density lipoprotein to oxidation in vivo after increased intake of berries. *Nutr Metab Cardiovasc Dis* 2000, *10* (6), 331–337.

83. Suomela, J.-P.; Ahotupa, M.; Yang, B.; Vasankari, T.; Kallio, H. Absorption of flavonols derived from sea buckthorn (*Hippophaë Rhamnoides* L.) and their effect on emerging risk factors for cardiovascular disease in humans. *J. Agric. Food Chem.* 2006, *54* (19), 7364–7369. <https://doi.org/10.1021/jf061889r>.
84. Linna, M. S.; Ahotupa, M.; Kukkonen-Harjula, K.; Fogelholm, M.; Vasankari, T. J. Co-Existence of insulin resistance and high concentrations of circulating oxidized LDL lipids. *Annals of Medicine* 2015, *47* (5), 394–398. <https://doi.org/10.3109/07853890.2015.1043939>.
85. Kosola, J.; Vaara, J. P.; Ahotupa, M.; Kyröläinen, H.; Santtila, M.; Oksala, N.; Atalay, M.; Vasankari, T. Elevated concentration of oxidized LDL together with poor cardiorespiratory and abdominal muscle fitness predicts metabolic syndrome in young men. *Metabolism* 2013, *62* (7), 992–999. <https://doi.org/10.1016/j.metabol.2013.01.013>.
86. Targher, G.; Day, C. P.; Bonora, E. Risk of cardiovascular disease in patients with nonalcoholic fatty liver disease. *N Engl J Med* 2010, *363* (14), 1341–1350. <https://doi.org/10.1056/NEJMra0912063>.
87. Kaikkonen, J. E.; Kresanov, P.; Ahotupa, M.; Jula, A.; Mikkilä, V.; Viikari, J. S. A.; Juonala, M.; Hutri-Kähönen, N.; Kähönen, M.; Lehtimäki, T.; Kangas, A. J.; Soininen, P.; Ala-Korpela, M.; Raitakari, O. T. Longitudinal study of circulating oxidized LDL and HDL and fatty liver: The Cardiovascular Risk in Young Finns Study. *Free Radical Research* 2016, *50* (4), 396–404. <https://doi.org/10.3109/10715762.2015.1133906>.
88. Raitakari, O. T.; Toikka, J. O.; Laine, H.; Ahotupa, M.; Iida, H.; Viikari, J. S. A.; Hartiala, J.; Knuuti, J. Reduced myocardial flow reserve relates to increased carotid intima-media thickness in healthy young men. *Atherosclerosis* 2001, *156* (2), 469–475. [https://doi.org/10.1016/S0021-9150\(00\)00689-4](https://doi.org/10.1016/S0021-9150(00)00689-4).
89. Toikka, J. O.; Niemi, P.; Ahotupa, M.; Niinikoski, H.; Viikari, J. S. A.; Rönnemaa, T.; Hartiala, J. J.; Raitakari, O. T. Large-artery elastic properties in young men: relationships to serum lipoproteins and oxidized low-density lipoproteins. *ATVB* 1999, *19* (2), 436–441. <https://doi.org/10.1161/01.ATV.19.2.436>.
90. Linna, M.; Ahotupa, M.; Lopponen, M. K.; Irjala, K.; Vasankari, T. Circulating oxidised LDL lipids, when proportioned to HDL-c, emerged as a risk factor of all-cause mortality in a population-based survival study. *Age and Ageing* 2013, *42* (1), 110–113. <https://doi.org/10.1093/ageing/afs074>.
91. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: The Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994, *344* (8934), 1383–1389.
92. Shepherd, J.; Cobbe, S. M.; Ford, I.; Isles, C. G.; Lorimer, A. R.; Macfarlane, P. W.; McKillop, J. H.; Packard, C. J. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med* 1995, *333* (20), 1301–1308. <https://doi.org/10.1056/NEJM199511163332001>.
93. Vasankari, T.; Ahotupa, M.; Viikari, J.; Nuotio, I.; Vuorenmaa, T.; Strandberg, T.; Vanhanen, H.; Tikkanen, M. J. Effects of statin therapy on circulating conjugated dienes, a measure of LDL Oxidation. *Atherosclerosis* 2005, *179* (1), 207–209. <https://doi.org/10.1016/j.atherosclerosis.2004.11.008>.
94. Janatuinen, T.; Knuuti, J.; Toikka, J. O.; Ahotupa, M.; Nuutila, P.; Rönnemaa, T.; Raitakari, O. T. Effect of pravastatin on low-density lipoprotein oxidation and myocardial perfusion in young adults with type 1 diabetes. *ATVB* 2004, *24* (7), 1303–1308. <https://doi.org/10.1161/01.ATV.0000132409.87124.60>.
95. Baigent et al. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90 056 participants in 14 randomised trials of statins. *The Lancet* 2005, *366* (9493), 1267–1278. [https://doi.org/10.1016/S0140-6736\(05\)67394-1](https://doi.org/10.1016/S0140-6736(05)67394-1).
96. Lawler, P. R.; Akinkuolie, A. O.; Chu, A. Y.; Shah, S. H.; Kraus, W. E.; Craig, D.; Padmanabhan, L.; Glynn, R. J.; Ridker, P. M.; Chasman, D. I.; Mora, S. Atherogenic lipoprotein determinants of cardiovascular disease and residual risk among individuals with low low-density lipoprotein cholesterol. *JAMA* 2017, *6* (7), e005549. <https://doi.org/10.1161/JAHA.117.005549>.
97. Marschang, P.; Götsch, C.; Kirchmair, R.; Kaser, S.; Kähler, C. M.; Patsch, J. R. Postprandial, but not postabsorptive low-density lipoproteins increase the expression of intercellular adhesion molecule-1 in human aortic endothelial cells. *Atherosclerosis* 2006, *186* (1), 101–106. <https://doi.org/10.1016/j.atherosclerosis.2005.07.014>.
98. Sumida, Y.; Niki, E.; Naito, Y.; Yoshikawa, T. Involvement of free radicals and oxidative stress in NAFLD/NASH. *Free Radical Research* 2013, *47* (11), 869–880. <https://doi.org/10.3109/10715762.2013.837577>.
99. Bigagli, E.; Lodovici, M. Circulating oxidative stress biomarkers in clinical studies on type 2 diabetes and its complications. *Oxidative Medicine and Cellular Longevity* 2019, *2019*, 1–17. <https://doi.org/10.1155/2019/5953685>.
100. Oberg, B. P.; McMenamin, E.; Lucas, F. L.; McMonagle, E.; Morrow, J.; Ikizler, T. A. L. P.; Himmelfarb, J. Increased prevalence of oxidant stress and inflammation in patients with moderate to severe chronic kidney disease. *Kidney International* 2004, *65* (3), 1009–1016. <https://doi.org/10.1111/j.1523-1755.2004.00465.x>.
101. Tucker, P. S.; Scanlan, A. T.; Dalbo, V. J. Chronic kidney disease influences multiple systems: describing the relationship between oxidative stress, inflammation, kidney damage, and concomitant disease. *Oxidative Medicine and Cellular Longevity* 2015, *2015*, 1–8. <https://doi.org/10.1155/2015/806358>.

102. Ho, H.-J.; Shirakawa, H. Oxidative stress and mitochondrial dysfunction in chronic kidney disease. *Cells* 2022, 12 (1), 88. <https://doi.org/10.3390/cells12010088>.
103. Ruan, X. Z.; Varghese, Z.; Moorhead, J. F. An Update on the lipid nephrotoxicity hypothesis. *Nat Rev Nephrol* 2009, 5 (12), 713–721. <https://doi.org/10.1038/nrneph.2009.184>.
104. De Ferranti, S. D.; De Boer, I. H.; Fonseca, V.; Fox, C. S.; Golden, S. H.; Lavie, C. J.; Magge, S. N.; Marx, N.; McGuire, D. K.; Orchard, T. J.; Zinman, B.; Eckel, R. H. Type 1 diabetes mellitus and cardiovascular disease: a scientific statement from the American Heart Association and American Diabetes Association. *Circulation* 2014, 130 (13), 1110–1130. <https://doi.org/10.1161/CIR.0000000000000034>.
105. Bae, J. C.; Han, J. M.; Kwon, S.; Jee, J. H.; Yu, T. Y.; Lee, M. K.; Kim, J. H. LDL-C/apoB and HDL-C/apoA-1 ratios predict incident chronic kidney disease in a large apparently healthy cohort. *Atherosclerosis* 2016, 251, 170–176. <https://doi.org/10.1016/j.atherosclerosis.2016.06.029>.
106. Jialal, I.; Devaraj, S. Antioxidants and atherosclerosis: don't throw out the baby with the bath water. *Circulation* 2003, 107 (7), 926–928. <https://doi.org/10.1161/01.CIR.0000048966.26216.4C>.
107. Ahotupa, M.; Vasankari, T. J. Baseline diene conjugation in LDL lipids: *Free Radical Biology and Medicine* 1999, 27 (11–12), 1141–1150. [https://doi.org/10.1016/S0891-5849\(99\)00201-4](https://doi.org/10.1016/S0891-5849(99)00201-4).
108. Bacot, S.; Bernoud-Hubac, N.; Chantegrel, B.; Deshayes, C.; Doutheau, A.; Ponsin, G.; Lagarde, M.; Guichardant, M. Evidence for in situ ethanolamine phospholipid adducts with hydroxyalkenals. *Journal of Lipid Research* 2007, 48 (4), 816–825. <https://doi.org/10.1194/jlr.M600340-JLR200>.
109. Goldstein, J. L.; Brown, M. S. The LDL receptor. *ATVB* 2009, 29 (4), 431–438. <https://doi.org/10.1161/ATVBAHA.108.179564>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.