

Article

Not peer-reviewed version

Trichoderma virens and Pseudomonas chlororaphis Differentially Regulate Maize Resistance to Anthracnose Leaf Blight and Insect Herbivores When Grown in Sterile versus Non-sterile Soils

[Pei-Cheng Huang](#) , Peiguo Yuan , [John M. Grunseich](#) , [James Taylor](#) , [Eric-Olivier Tiénébo](#) , Elizabeth A. Pierson , [Julio S. Bernal](#) , [Charles M. Kenerley](#) , [Michael V. Kolomiets](#) *

Posted Date: 22 March 2024

doi: 10.20944/preprints202403.1285.v1

Keywords: Trichoderma virens; Pseudomonas chlororaphis; induced systemic resistance; jasmonic acid; Spodoptera frugiperda; Diabrotica virgifera virgifera



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Trichoderma virens and *Pseudomonas chlororaphis* Differentially Regulate Maize Resistance to Anthracnose Leaf Blight and Insect Herbivores When Grown in Sterile Versus Non-Sterile Soils

Pei-Cheng Huang ¹, Peiguo Yuan ¹, John M. Grunseich ², James Taylor ¹, Eric-Olivier Tiénébo ^{3,4}, Elizabeth A. Pierson ^{1,3}, Julio S. Bernal ², Charles M. Kenerley ¹ and Michael V. Kolomiets ^{1,*}

¹ Department of Plant Pathology and Microbiology, Texas A&M University, College Station, TX 77843-2132, USA; pei-cheng.huang@ag.tamu.edu (P.-C.H.); peiguo.yuan@ag.tamu.edu (P.Y.); jimtaylor011@gmail.com (J.T.); elizabeth.pierson@ag.tamu.edu (E.A.P.); mike.kolomiets@ag.tamu.edu (M.V.K.)

² Department of Entomology, Texas A&M University, College Station, TX 77843-2475, USA; johngrunseich@tamu.edu (J.M.G.); julio.bernal@ag.tamu.edu (J.S.B.)

³ Department of Horticultural Sciences, Texas A&M University, College Station, TX 77843-2133, USA; elizabeth.pierson@ag.tamu.edu (E.A.P.)

⁴ Agronomic Sciences and Transformation Processes Joint Research and Innovation Unit, Institut National Polytechnique Félix Houphouët-Boigny, Yamoussoukro, PO BOX 1093 Yamoussoukro, Côte d'Ivoire; eric.tienebo@inphb.ci (E.-O.T.)

* Correspondence: mike.kolomiets@ag.tamu.edu (M.V.K.)

Abstract: Soil-borne *Trichoderma* spp. have been extensively studied for their biocontrol activities against pathogens and growth promotion ability with plants. However, the beneficial effect of *Trichoderma* on inducing resistance against insect herbivores has been considered only recently. Among diverse *Trichoderma* species, consistent with previous reports, we showed that root colonization by *T. virens* triggered induced systemic resistance (ISR) to leaf-infecting fungal pathogens. Whether *T. virens* induces ISR to insect pests has not been tested before. In this study, we investigated whether *T. virens* affects jasmonic acid (JA) biosynthesis and defense against fall armyworm (FAW) and western corn rootworm (WCR). Unexpectedly, the results showed that *T. virens* colonization of maize seedlings grown in autoclaved soil suppressed wound-induced production of JA, resulting in reduced resistance to FAW. Similarly, the bacterial endophyte *Pseudomonas chlororaphis* 30-84 suppressed systemic resistance to FAW due to reduced JA. Further comparative analyses of the systemic effects of these endophytes when applied into sterile or non-sterile field soil showed that both *T. virens* and *P. chlororaphis* 30-84 triggered ISR against hemibiotrophic fungal pathogen *Colletotrichum graminicola* in both soil conditions, but only suppressed resistance to FAW in sterile soil. In contrast to the effect on FAW defense, *T. virens* colonization of maize roots suppressed WCR larvae survival and weight gain. This is the first report suggesting the potential role of *T. virens* as a biocontrol agent against WCR.

Keywords: *Trichoderma virens*; *Pseudomonas chlororaphis*; induced systemic resistance; jasmonic acid; *Spodoptera frugiperda*; *Diabrotica virgifera virgifera*

1. Introduction

Maize is the top-yielding cereal crop produced globally and a major source of livestock feed, biofuel, and staple food for many regions worldwide [1]. Pathogen infection and insect herbivory are constant threats to crop production causing over 20% yield losses of maize annually at a global level, resulting in substantial economic impact and food insecurity [2]. The hemibiotrophic fungal pathogen *Colletotrichum graminicola* is the causal agent of anthracnose leaf blight and stalk rot of maize and

accounts for millions of dollars in yield losses annually in the United States [3-5]. Fall armyworm (FAW; *Spodoptera frugiperda*) originates in the Americas and has recently invaded Europe, Africa, Asia, and Australia causing serious damage [6, 7]. FAW larvae feed on all the aboveground tissue of maize, resulting in several billions of dollars in yield losses globally [6]. Western corn rootworm (WCR; *Diabrotica virgifera virgifera*) is the most economically important maize pest in the United States and causes losses in excess of US \$1 billion annually in the United States and Europe [8-10]. WCR larvae cause the most significant damage through feeding on the corn root while WCR adult beetles damage the silks [8, 11]. Published evidence indicates that the plant phytohormone salicylic acid (SA) plays a critical role in maize defense against *C. graminicola* [12, 13], while resistance to chewing insects, including FAW, is mainly mediated by the lipid-derived phytohormone jasmonic acid (JA) [14-16]. However, resistance mechanisms against root-feeding WCR remain largely unknown [17].

Microbial biological control agents, including plant growth-promoting rhizobacteria and fungi (PGPR and PGPF, respectively), have been widely used to increase resistance in plants against pathogens and pests [18-20]. Isolates of soil-borne *Trichoderma* spp. and bacterium *Pseudomonas chlororaphis* are well-known biocontrol agents and have been extensively studied for their beneficial traits, including growth promotion and enhanced disease resistance of the host plants [21-24]. However, using biocontrol agents against pests has only recently been considered, and little is known about their impact on defense against chewing insects [25, 26]. Several studies have shown that *T. virens* induces systemic resistance (ISR) against leaf-infecting pathogens, the physiological process tightly regulated by fungal small secreted proteins, Sm1 and Sir1, which play pivotal but contrasting roles in triggering ISR responses [27-30]. Analyses of the *T. virens sm1* deletion mutants revealed that they are unable to induce ISR against fungal pathogens in above-ground organs, suggesting that Sm1 peptide is a positive regulator of ISR [27, 28]. In sharp contrast, deletion of *SIR1* resulted in much stronger ISR response [29]. Wang et al. (2020)[30] showed that *T. virens* colonization in maize roots upregulated genes involved in biosynthesis of JA precursors, including 12-OPDA (12-oxo-10(Z),15(Z)-phytyldienoic acid) and OPC-4:0 ((Z)-4-[3-oxo-2-(pent-2-en-1-yl)cyclopentyl]butanoic acid), yet downregulated genes downstream of 12-OPDA for JA biosynthesis. Whether *T. virens* colonization affects wound-induced JA production in leaves and confers resistance to leaf-feeding FAW and root-feeding WCR larvae remains unexplored.

The principal objective of this study was to investigate the effects of *T. virens* and its secreted peptide signals on the production of wound-induced JA and other defensive oxylipin metabolites and to test whether the beneficial effects of this endophyte extend to increasing resistance against FAW and WCR. Contrary to our expectation, the results demonstrated that maize seedlings grown in sterile soil amended with wild-type *T. virens*, the *sm1* or *sir1* single or the double mutants were found to suppress wound-induced JA biosynthesis, resulting in reduced resistance to FAW. We followed up by testing whether such a detrimental effect on defense against herbivory could be ascribed to another well-studied growth-promoting bacterial endophyte, *P. chlororaphis*. Colonization of roots by the *P. chlororaphis* 30-84 also reduced production of JA and defense against FAW. Further analysis showed that while both *T. virens* and *P. chlororaphis* induced ISR against *C. graminicola* when added in sterile or nonsterile soil, suppression of defense against FAW occurred only when maize seedlings were grown in sterile soil amended with these biocontrol agents. In addition, we demonstrated that *T. virens* reduced survival and growth (weight gain) of WCR larvae, making our study the first showing the potential of *T. virens* as a biocontrol agent against this devastating pest of maize.

2. Results

2.1. Colonization of Maize Roots by either $\Delta sir1$ or $\Delta sm1\Delta sir1$ Enhanced Greater Levels of ISR against *C. Graminicola*

Previously, we have shown that *T. virens* colonization triggered ISR against *C. graminicola* in a Sm1 peptide-dependent manner as the $\Delta sm1$ mutant lost its ISR-inducing capability [28, 30]. In contrast, plants treated with the deletion mutant of Sir1 peptide, $\Delta sir1$, exhibited enhanced systemic

resistance [29, 30]. Moreover, Wang et al. (2020)[30] showed that the expression of *SIR1* in $\Delta sm1$ remained similar to the level in wild-type *T. vires* strain (TvWT), while $\Delta sir1$ had elevated expression of *SM1*. These results suggested that one potential reason for increased ISR by $\Delta sir1$ is due to elevated *SM1* expression in this strain. To test this possibility, we created $\Delta sm1\Delta sir1$ double mutant and tested its capability to induce resistance to *C. graminicola*. The result showed that colonization of both $\Delta sir1$ and $\Delta sm1\Delta sir1$ mutants triggered greater level of ISR against *C. graminicola* compared to TvWT (32%, 59%, and 56% reduction in lesion areas on TvWT-, $\Delta sir1$ -, and $\Delta sm1\Delta sir1$ -treated plants, respectively, compared to control) (Figure 1). These results suggest that a functional Sir1 suppresses disease resistance to *C. graminicola* by suppressing not only *SM1*, but other as yet unknown mechanisms.




Figure 1. Both *T. vires* $\Delta sir1$ and $\Delta sm1\Delta sir1$ treatments enhanced greater level of ISR against *C. graminicola* than TvWT. Lesion areas of maize seedlings colonized by different *T. vires* strains after inoculation with *C. graminicola* for 7 days. Bars are means \pm SE. Different letters indicate statistically significant differences (Tukey's HSD test, $P < 0.05$).

2.2. *T. vires* Colonization Suppressed Insect Defense

Numerous studies have shown that beneficial *Trichoderma* spp. induce resistance against pathogens [31-33]. However, their effect on inducing insect resistance remains largely unexplored. Here, we tested whether different *T. vires* stains, including $\Delta sm1$, $\Delta sir1$, and $\Delta sm1\Delta sir1$, induce differential resistance to the foliar-feeding FAW. Surprisingly, the results showed that FAW larvae gained significantly more weight after feeding on *T. vires*-colonized plants for 7 days (60% to 95% increase) (Figure 2A) and 3rd-instar larvae consumed more leaf area on *T. vires*-colonized plants (64% to 83% increase) (Figure 2B). Treatment with *T. vires* reduced resistance to FAW, regardless of the mutation of these two signaling peptides.




Figure 2. *T. vires* colonization suppressed insect defense. One-week old maize seedlings were colonized by *T. vires* strains for 14 days before FAW infestation. (A) FAW larval weight was determined after neonates feeding on maize plants for 7 days. (B) Maize leaf consumed area by 3rd-

instar FAW larvae for 6 hours. Bars are means \pm SE. Different letters indicate statistically significant differences (Tukey's HSD test, $P < 0.05$).

2.3. *T. vires* Treatment Reduced Wound-Induced JA Accumulation by Suppressing Biosynthesis and Enhancing Catabolism

A previous study showed that treatment with *T. vires* downregulated JA biosynthesis genes in colonized maize roots [30] and that JA dominantly modulates defense responses against chewing insects [14]. In the present study, we found that *T. vires* treatment decreased insect defense, which prompted us to test whether *T. vires* colonization regulates wound-induced JA production in leaves. Phytohormone and oxylipin profiling showed that significantly lower amounts of jasmonates, including JA precursors, 12-OPDA, OPC-4:0, JA, and the bioactive JA derivatives, JA-Isoleucine (JA-Ile), JA-leucine (JA-Leu), and JA-Valine (JA-Val) conjugates, were accumulated in the *T. vires*-treated plants at 1 and 2 hours post wounding (hpw) (Figure 3A). Specifically, JA accumulation was reduced by 23% to 37% and 33% to 38% and JA-Ile content was reduced by 8% to 37% and 44% to 50% compared to the amounts accumulated in the control plants at 1 and 2 hpw, respectively (Figure 3B,C). Notably, we also found that one of the JA catabolites, 12OH-JA, accumulated to a significantly higher level in the *T. vires*-treated plants in response to mechanical wounding (83% to 171% and 23% to 82% increase compared to the contents in the control plants at 1 and 2 hpw, respectively) (Figure 3A and 3D). This latter result suggests that one potential reason behind decreased JA content may be due to increased catabolism of JA. To confirm the reduced insect defense by *T. vires* colonization was modulated by JA signaling, we tested the effect of *T. vires* on the JA-deficient *opr7opr8* mutant [14] to FAW. The result showed that *T. vires* colonization did not increase susceptibility to FAW on *opr7opr8* double mutants, suggesting that attenuated JA signaling plays an essential role in *T. vires*-reduced insect defense (Figure 4). Interestingly, the levels of the α -ketol, 9,10-KODA (9-hydroxy-10-oxo-12(Z),15(Z)-octadecadienoic acid), have been significantly reduced at 1 hpw (24% to 37% reduction compared to the contents in the control plants) by all strains of *T. vires* (Figure 3E). This molecule has been recently shown to play a major JA-dependent signaling role in defense against FAW [34]. Together, these results indicate that *T. vires* colonization suppressed defense against FAW via downregulating JA biosynthesis and/or enhancing JA catabolism.




Figure 3. Cont.




Figure 3. *T. vires* colonization suppressed wound-induced JA biosynthesis and enhanced JA catabolism. (A) Heatmap showing relative accumulation of oxylipins and phytohormones in leaves of B73 seedlings colonized by different *T. vires* strains or untreated control at 0, 1, and 2 hours post wounding. Contents of (B) JA, (C) JA-Ile, (D) 12OH-JA, and (E) 9,10-KODA in different *T. vires* strains colonized or control plants at 0, 1, and 2 hours post wounding. Abbreviation for compounds were listed in Supplemental Table 1. Bars are means \pm SE. Different letters indicate statistically significant differences within the same time points (Tukey's HSD test, $P < 0.05$).




Figure 4. Reduced defense against FAW by *T. vires* colonization is modulated by JA signaling. Maize B73 wild-type or *opr7opr8* double mutant seedlings were colonized by TvWT for 14 days or left as untreated control (CK) before FAW infestation. FAW larval weight was determined after neonates feeding on maize plants for 7 days. Bars are means \pm SE. Different letters indicate statistically significant differences on log-transformed data (Two-way ANOVA with Tukey's HSD test, $P < 0.05$).

2.4. *T. vires* and *P. chlororaphis* Colonization Enhanced Growth but Suppressed Insect Defense against FAW in Sterile Soil While Increasing Resistance to *C. graminicola* in both Sterile and Nonsterile Soil

Bacterial biocontrol agent *P. chlororaphis* 30-84 was previously reported to promote growth and suppress growth of fungal pathogens [35, 36], yet its role in insect defense against foliar-feeding insects remains mostly unexplored. Therefore, we also tested the impact of *P. chlororaphis* 30-84 on

plant growth promotion, defense against FAW, and pathogen resistance and the effect of combining the two beneficial microorganisms, *T. virens* and *P. chlororaphis* 30-84, on these physiological processes. Because the beneficial microorganisms were commonly tested in an agricultural setting, we further tested their effect in both sterile and nonsterile soil mixtures of field soil and potting mix. Similar to *T. virens* treatment, colonization by *P. chlororaphis* 30-84 promoted plant growth (12% to 15% increase in plant height) but suppressed defense against FAW (41% to 51% increase in weight gain) in the sterile soil mixture (Figure 5A,B). Plant growth and insect defense were not significantly affected when seedlings were grown in a nonsterile soil mixture upon treatment with *T. virens* and *P. chlororaphis* 30-84. However, both *T. virens* and *P. chlororaphis* 30-84 were able to increase resistance to *C. graminicola* in both soil types (53% to 66% reduction and 44% to 55% reduction in lesion areas in nonsterile and sterile soils, respectively) (Figure 5C).




Figure 5. *T. virens* and *P. chlororaphis* 30-84 colonization enhanced growth but suppressed defense against FAW only in sterile soil while increased resistance to *C. graminicola* in both soil types. Maize seedlings were colonized by TvWT, *P. chlororaphis* 30-84, or both for 14 days before plant height measurement and *C. graminicola* infection or FAW infestation. (A) Plant height measured at 14 days after colonization. (B) FAW larval weight and (C) lesion areas were determined at 7 days after infection or infestation. Bars are means \pm SE. Different letters indicate statistically significant differences on log-transformed data within the same treatment (Tukey's HSD test, $P < 0.05$). Asterisks represent statistically significant differences between sterile and nonsterile soils (Student's *t*-test, ** $P < 0.01$, *** $P < 0.001$).

2.5. *T. vires* and *P. chlororaphis* 30-84 Colonization Reduced Wound-Induced JA Accumulation by Suppressing Biosynthesis and Enhancing Catabolism in Plants Grown in Sterile Soil

Decreased FAW resistance found with the amendment of *P. chlororaphis* 30-84 in this study prompted us to investigate whether this biocontrol agent impacts JA homeostasis in response to wounding in sterile soil. Phytohormone and oxylipin profiling revealed that *P. chlororaphis* 30-84 treatment reduced wound-induced accumulation of jasmonates, including 12-OPDA, JA, JA-Ile, and JA-Leu in sterile soil (Figure 6A). More specifically, treatments with *T. vires*, *P. chlororaphis* 30-84, and the mixture of *T. vires* and *P. chlororaphis* 30-84 resulted in 25%, 21%, and 19% reduction in JA accumulation and 27%, 32%, and 27% reduction in JA-Ile level at 1 hpw, respectively (Figure 6B,C). Moreover, JA catabolism was increased by 50%, 13%, and 21% in the plants treated with *T. vires*, *P. chlororaphis* 30-84, and the mixture of *T. vires* and *P. chlororaphis* 30-84, respectively (Figure 6D). Interestingly, suppression of wound-induced JA biosynthesis (Figure 6) and FAW insect defense (Figure 5) by either *T. vires* or *P. chlororaphis* 30-84 was only observed in the sterile soil condition. A similar accumulation pattern was observed for 9,10-KODA (17% to 34% reduction) in sterile soil (Figure 6E). Together, these results suggest that *T. vires* and *P. chlororaphis* 30-84 suppress foliar-feeding insect defense via regulating JA homeostasis in response to wounding only in plants grown in sterile soil amended with either of these biocontrol agents.




Figure 6. *T. virens* and *P. chlororaphis* 30-84 colonization suppressed wound-induced JA biosynthesis and enhanced JA catabolism in sterile soil. (A) Heatmap showing relative accumulation of oxylipins and phytohormones. Contents of (B) JA, (C) JA-Ile, (D) 12OH-JA, and (E) 9,10-KODA in plants colonized by *T. virens*, *P. chlororaphis* 30-84, and both or control in response to wounding at 0 and 1 hpw. Abbreviations for samples: NS, nonsterile soil; SS, sterile soil. Abbreviation for compounds were listed in Supplemental Table 1. Bars are means \pm SE. Different letters indicate statistically significant differences within the same treatment per time point (Tukey's HSD test, $P < 0.05$).

2.6. *T. virens* Colonization Suppressed WCR Larval Survival and Weight Gain

We also tested whether *T. virens* plays a role in conferring resistance to root-feeding WCR larvae. The results showed that significantly fewer larvae were recovered (50% fewer) (Figure 7A), and WCR larvae gained significantly less weight after feeding on *T. virens*-colonized plants (57% less) (Figure 7B). However, *T. virens* treatment did not reduce tissue consumption (Figure 7C,D). These results suggest that *T. virens* may be a potent biocontrol agent against WCR by suppressing WCR larval survival and weight gain.




Figure 7. *T. virens* colonization suppressed WCR larval survival and weight gain. (A) Larvae recovery, (B) larval mass, (C) root, and (D) shoot ratios were recorded 10 days post-infestation. Bars are means \pm SE. Asterisks represent statistically significant differences (Student's *t*-test, * $P < 0.05$, *** $P < 0.001$). Different letters indicate statistically significant differences (Two-way ANOVA with Tukey's HSD test, $P < 0.05$).

3. Discussion

Soil-borne *Trichoderma* spp. are ubiquitous in soil worldwide, and have been used as biocontrol agents against pathogens for decades [23, 31, 37-39]. However, the potential utility of *Trichoderma* spp. as biocontrol agents against insect pests has just begun to be addressed [26], and information on their effects against foliar-feeding and root-feeding insects is scarce. Accumulating evidence suggests that root colonization by *Trichoderma* spp. triggered ISR against insects, such as aphids [40-43], whiteflies [44], thrips [45], and Lepidoptera [41, 46-50], and also enhanced defense against spider mites [51] and nematodes [52-54]. Previous reports [27, 28, 30] and this study showed that *T. virens* confers resistance to *C. graminicola*. In addition, these studies have identified a fungal-secreted peptide, Sm1, as a signal necessary for inducing systemic resistance by showing that the $\Delta sm1$ deletion mutant of *T. virens* is unable to induce ISR [27, 28, 30]. In contrast to $\Delta sm1$, another signaling peptide mutant of *T. virens*, $\Delta sir1$ [29, 30], triggered enhanced ISR against *C. graminicola*. As this

mutant overexpressed *SM1* transcripts [30], we hypothesized that the increased ISR following colonization by this mutant is due to elevated *SM1* expression. We tested this hypothesis by deleting *SM1* in the $\Delta sir1$ mutant background and found that the $\Delta sm1\Delta sir1$ double mutant still displayed stronger resistance as compared to the wild-type strain. We are suggesting that Sir1 peptide acts to suppress ISR not only by downregulating *SM1* expression, but also by employing other as-yet-unknown mechanisms of ISR suppression.

Several reports showed that several *Trichoderma* spp. directly impact Lepidoptera spp. by producing antifeedant compounds and toxic secondary metabolites [26, 55-59] or indirectly impact them via activation of JA-mediated systemic defense, production of parasitoid-attracting volatile compounds, or disruption of insect gut microbiota and metabolome [41, 42, 46-48, 50, 60]. Here, we demonstrated that the treatment with *T. virens* suppressed defense against FAW by reducing wound-induced JA production and increased JA catabolism in plants grown in sterile soil. Consistent with our findings, Lelio et al. (2021)[47] showed that when grown at cool temperatures (20°C as compared to 25°C), tomato plants colonized with *T. afroharzianum* T22 had less negative impact on *Spodoptera littoralis* larvae and this was accompanied with suppressed JA biosynthesis and signaling pathways. Also consistent with our results, Kinyungu et al. (2023)[49] showed that the treatment with *T. atroviride* F2S21 strain resulted in a higher leaf defoliation rate by FAW larvae. The fact that some studies showed increased resistance to certain chewing insects due to *Trichoderma* spp. treatment [41, 46, 50] while others, including the current study, show opposite effect of *Trichoderma* treatment, suggest *Trichoderma* species- and strain-specific effects as well an environment-dependent effect on insect defense. Interestingly, our results indicate that the reduced insect defense is independent of the fungal secreted peptides, Sm1 and Sir1, suggesting that fungal signals regulating ISR against pathogens and increased induced susceptibility against chewing insects involve different molecular signals to be identified in the future.

Previous reports showed that root colonization by *T. virens* induced biosynthesis of the JA precursor, 12-OPDA, and ketols [30, 61]. Both 12-OPDA and ketols were shown to act as xylem-mobile long-distance signals that prime the above-ground organs for greater resistance against pathogens in maize [30, 61]. In contrast to these oxylipins, JA-Ile levels in xylem sap were not significantly increased in response to *T. virens* treatment and transfusion of maize seedlings with xylem sap supplemented with JA-Ile increased susceptibility to *C. graminicola* [30]. The role of JA as a susceptibility factor against anthracnose leaf blight caused by *C. graminicola* was further supported by the studies of Gorman et al. (2020)[12] and Huang et al. (2023)[13]. Moreover, transcriptome analysis showed that maize roots responded to colonization by *T. virens* by increased expression of 12-OPDA biosynthesis genes but suppressed expression of the genes required for 12-OPDA conversion to JA-Ile [30]. Because JA is widely reported to suppress growth in numerous plant species [62-64], it is tempting to speculate that attenuated synthesis of JAs in roots colonized by *T. virens* is a potential mechanism accounting for the strong growth promotion effect by this symbiont observed in this study and previous reports. Similar to the *T. virens*-mediated induction of ISR against pathogens and growth promotion, we showed that root colonization by the bacterial symbiont, *P. chlororaphis* 30-84, resulted in increased seedling growth and systemic resistance against *C. graminicola*, suggesting a similar mechanism underlying these two beneficial responses. However, as in the case of *T. virens*, *P. chlororaphis* 30-84 treatment reduced JA production and decreased resistance to herbivory by FAW when treated plants were grown in sterile soil. Therefore, it is likely that growth promotion by both beneficial microorganisms, *T. virens* and *P. chlororaphis* 30-84, and ISR against the hemibiotrophic pathogen *C. graminicola* is linked to their ability to suppress JA synthesis and JA-mediated signaling. Interestingly, wound-induced JA accumulation and insect defense were not affected when maize seedlings were treated with *T. virens* or *P. chlororaphis* 30-84 in nonsterile soil, suggesting that natural microbial communities dampened JA suppression activities by both endophytes.

WCR root feeding causes substantial economic damage to corn production in the United States and Europe [65, 66] and the insect has adapted to overcome most of the available management techniques, including chemical insecticides and *Bacillus thuringiensis* (Bt) toxins [66, 67]. So far, there

is a lack of efficient management of this notorious pest and, thus, there is an urgent need for a sustainable and economical management approach, such as utilizing efficient biocontrol agents. In this study, we showed that *T. virens* colonization significantly enhanced mortality and suppressed weight gain of WCR larvae. Our previous report showed that increased WCR mortality was correlated with greater levels of herbivory-induced production of insecticidal death acids and ketols [16]. Therefore, one of the potential mechanisms of *T. virens*-mediated WCR defense may be to activate the production of insecticidal defensive compounds. The other potential mechanism behind the detrimental effect on larvae survival and growth conferred by *T. virens* may be the production of some fungus-derived insecticidal metabolites as reported in several studies [26, 55-59]. Further experiments are needed to explore whether *T. virens* has a direct insecticidal impact on the WCR root-feeding larvae or has an indirect effect due to the induction of the host-mediated synthesis of toxic compounds.

4. Materials and Methods

4.1. Plant, Growth Medium, and Plant-Beneficial Inoculants

The maize (*Zea mays*) inbred line B73 and *opr7opr8* double mutant (Yan et al., 2012)[14] were used in this study. Plants were grown in conical pots (20.5 × 4 cm) filled with autoclaved (121°C for 60 minutes, 2 cycles) commercial potting mix (Jolly Gardener Pro Line C/20) on light shelves at room temperature (22–24°C) with a 16-h light period. To test the efficacy of the beneficial microorganisms in an agricultural setting, maize B73 seeds were grown in a mixture of field soil (collected from Texas A&M University Research Farm in the fall of 2022 after maize was harvested) and potting mix (Jolly Gardener Pro Line C/20) at the ratio of 1 to 3 that was either autoclaved (121°C for 60 minutes, 2 cycles; henceforth referred to as sterile) or untreated (hence referred to as nonsterile) after mixing. Chlamydospores of *T. virens* strains TvWT (Gv29-8; Baek and Kenerly, 1998)[68], $\Delta sm1$ (Djonovic' et al., 2006; Djonovic et al., 2007)[27, 28], $\Delta sir1$ (formerly D77560; Lamdan et al., 2015)[29], and $\Delta sm1\Delta sir1$ (this study) were obtained from liquid cultures grown in molasses-yeast extract medium as described in Wang et al. (2020)[30]. *P. chlororaphis* 30-84 was grown in Luria-Bertani (LB) medium containing 5 g of NaCl per liter at pH 7 at 28°C with agitation at 200 rpm, as described in Yuan et al. (2020)[36]. Seven days after sowing, seedlings were treated with 0.05 g of the *T. virens* chlamydospores, or 300 μ l of *P. chlororaphis* 30-84 suspension at OD₆₂₀ = 0.8, or a mixture of both *T. virens* and *P. chlororaphis* with the same amount described above (added to the soil at a depth of approximately 2 cm around the seeds), or left untreated.

4.2. Anthracnose Leaf Blight Assay

C. graminicola stain 1.001 was grown on half strength potato dextrose agar (PDA) plates for 2-3 weeks to allow sporulation. Two weeks after being colonized by beneficial microorganisms, the plants were inoculated with 10 μ l of conidial suspension (10⁶ conidia/ml) with six droplets of spore suspensions per leaf collected from hemibiotrophic fungal pathogen *C. graminicola* plates as described in Huang et al. (2023a)[13]. The leaves were scanned and the lesion areas measured using ImageJ software [69] seven days post-inoculation.

4.3. Fall Armyworm Assay

The laboratory strain of fall armyworm (FAW; *Spodoptera frugiperda*) was purchased from Benzon Research (Carlisle, PA, USA). Two weeks after incubation with the beneficial microorganisms, six FAW neonates were placed to the whorls of maize seedlings contained in individual plastic jars and allowed to move and feed on the plants freely for 7 days, as described in Huang et al. (2023b)[16]. FAW larvae were removed from the plants, and the total weight was determined 7 days post-infestation. To evaluate insect resistance and leaf damaged area, one maize leaf was individually caged in a handmade clip-cage and infested with one 3rd-instar FAW larva per spot for approximately one hour, then moved toward the base of the plant. Leaves were then scanned, and eaten areas were measured with ImageJ software[69].

4.4. Oxylipin Profiling of Wounded Leaf Tissue

For the wounding treatment, the third fully expanded leaf of seedlings grown in sterile or nonsterile soil conditions that were previously colonized with beneficial microorganisms for two weeks or left untreated was wounded seven times using a hemostat, with three wound sites on one side and four on the other side of the midvein with the wound sites approximately 1 cm apart. The wounded regions were then harvested and placed in 2 mL screw-cap Fast-Prep tubes (Qbiogene, Carlsbad, CA, U. S. A.) in liquid nitrogen and stored in a -80 °C freezer. Phytohormone and oxylipin extraction and profiling of wounded leaf tissue were performed as described previously [13].

4.5. Western Corn Rootworm Bioassays

Western corn rootworm (WCR; *Diabrotica virgifera virgifera*) eggs were provided by USDA-ARS North Central Agriculture Research Laboratory (Brookings, SD, USA). Two weeks after being colonized by beneficial microorganisms, ten WCR neonates were placed on the soil surface in individual cone-tainers with a maize seedling and allowed to burrow and feed for 10 days. WCR larvae were removed from the soil, and the total weight was determined 10 days post-infestation. The fresh weights of shoot and root tissues was measured after the soil was removed and the ratio was analyzed as larvae damaged over control root mass, as described in Huang et al., (2023)[16].

4.6. Statistical Methods

Statistical analyses were performed using the software programs JMP Pro 17 (SAS Institute Inc., Cary, NC, USA). Lesion areas, FAW larval weight, leaf eaten areas, wound-induced accumulation of metabolites, and plant heights were analyzed using One-way Analysis of Variance (ANOVA) with post-hoc Tukey HSD (Honestly Significant Difference). FAW larval weight after feeding on *opr7opr8* double mutants and wild-type seedlings treated with *T. virens* or left untreated and root and shoot ratios after WCR infestation were analyzed using Two-way ANOVA with Tukey HSD. Effect of sterile and nonsterile mixture was analyzed using Student's *t*-test.

5. Conclusions

The elevated level of ISR against *C. graminicola* triggered by both *T. virens* $\Delta sir1$ and $\Delta sm1\Delta sir1$ double mutants revealed that a functional Sir1 signal peptide reduces disease resistance by downregulating not only *SM1* (as demonstrated previously by Wang et al. (2020)) but also involves other unknown mechanisms as well. Colonization of plants grown in sterile soil by different *T. virens* mutants suppressed wound-induced JA accumulation, resulting in reduced defense against FAW, which is independent of both signal peptides, Sm1 and Sir1. When testing both beneficial microorganisms, *T. virens* and *P. chlororaphis* 30-84, using both sterile and nonsterile mixtures of field soil and potting mix, both biocontrol agents triggered ISR against *C. graminicola* in both soil types. However, these agents reduced defense against FAW only when the treated plants were grown in sterile soil due to reduced JA. In contrast to the effect on leaf-feeding FAW, *T. virens* colonization suppressed root-feeding WCR larvae survival and weight gain, the first report revealing the potential of *T. virens* as a biocontrol agent against WCR.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Figure S1: title; Table S1: title; Video S1: title.

Author Contributions: Designed the research and analyzed the data, P.-C.H. and M.V.K.; performed most of the experiments and drafted the article, P.-C.H.; treatment with *P. chlororaphis* 30-84, P.Y.; conducted the WCR bioassays, J.M.G and P.-C.H.; created $\Delta sm1\Delta sir1$ double mutant, J.T.; helped design the experiments and wrote the article with input from and revisions by J.M.G., J.T., E.-O.T., E.A.P., J.S.B., and C.M.K. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported in part by United States Department of Agriculture (USDA)-National Institute of Food and Agriculture (NIFA) 2017-67013- 26524 to M.V.K and 2021-67013-33568 grants awarded to M.V.K and J.S.B.

Data Availability Statement: The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

Acknowledgments: We thank Andie Miller and Dr. Greg Sword, Department of Entomology, Texas A&M University, for kindly sharing FAW eggs/neonates.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Erenstein, O.; Jaleta, M.; Sonder, K.; Mottaleb, K.; Prasanna, B. M., Global maize production, consumption and trade: trends and R&D implications. *Food Security* **2022**, *14* (5), 1295-1319.
- Savary, S.; Willocquet, L.; Pethybridge, S. J.; Esker, P.; McRoberts, N.; Nelson, A., The global burden of pathogens and pests on major food crops. *Nature ecology & evolution* **2019**, *3* (3), 430-439.
- Belisário, R.; Robertson, A. E.; Vaillancourt, L. J., Maize Anthracnose Stalk Rot in the Genomic Era. *Plant Disease* **2022**, *106* (9), 2281-2298.
- Bergstrom, G. C.; Nicholson, R. L., The Biology of Corn Anthracnose: Knowledge to Exploit for Improved Management. *Plant Disease* **1999**, *83* (7), 596-608.
- Dean, R.; Van Kan, J. A. L.; Pretorius, Z. A.; Hammond-Kosack, K. E.; Di Pietro, A.; Spanu, P. D.; Rudd, J. J.; Dickman, M.; Kahmann, R.; Ellis, J.; Foster, G. D., The Top 10 fungal pathogens in molecular plant pathology. *Molecular plant pathology* **2012**, *13* (4), 414-430.
- Kenis, M. B.; Giovanni; Biondi, Antonio; Calatayud, Paul-André; Day, Roger; Desneux, Nicolas; Harrison, Rhett D; Kriticos, Darren; Rwomushana, Ivan; van den Berg, Johnnie; Verheggen, François; Zhang, Yong-Jun; Agboyi, Lakpo Koku; Ahissou, Régis Besmer; Ba, Malick N; Wu, Kongming, Invasiveness, biology, ecology, and management of the fall armyworm, Spodoptera frugiperda. *Entomologia Generalis* **2023**, *43* (2), 55.
- Overton, K.; Maino, J. L.; Day, R.; Umina, P. A.; Bett, B.; Carnovale, D.; Ekesi, S.; Meagher, R.; Reynolds, O. L., Global crop impacts, yield losses and action thresholds for fall armyworm (Spodoptera frugiperda): A review. *Crop Protection* **2021**, *145*, 105641.
- Gray, M. E.; Sappington, T. W.; Miller, N. J.; Moeser, J.; Bohn, M. O., Adaptation and invasiveness of western corn rootworm: intensifying research on a worsening pest. *Annu Rev Entomol* **2009**, *54*, 303-21.
- Darlington, M.; Reinders, J. D.; Sethi, A.; Lu, A. L.; Ramaseshadri, P.; Fischer, J. R.; Boeckman, C. J.; Petrick, J. S.; Roper, J. M.; Narva, K. E.; Vélez, A. M., RNAi for Western Corn Rootworm Management: Lessons Learned, Challenges, and Future Directions. *Insects* **2022**, *13* (1).
- Wechsler, S.; Smith, D., Has resistance taken root in US corn fields? Demand for insect control. *American Journal of Agricultural Economics* **2018**, *100* (4), 1136-1150.
- Gyeraj, A.; Szalai, M.; Pálkás, Z.; Edwards, C. R.; Kiss, J., Effects of adult western corn rootworm (*Diabrotica virgifera virgifera* LeConte, Coleoptera: Chrysomelidae) silk feeding on yield parameters of sweet maize. *Crop Protection* **2021**, *140*, 105447.
- Gorman, Z.; Christensen, S. A.; Yan, Y.; He, Y.; Borrego, E.; Kolomiets, M. V., Green leaf volatiles and jasmonic acid enhance susceptibility to anthracnose diseases caused by *Colletotrichum graminicola* in maize. *Molecular plant pathology* **2020**, *21* (5), 702-715.
- Huang, P. C.; Tate, M.; Berg-Falloure, K. M.; Christensen, S. A.; Zhang, J.; Schirawski, J.; Meeley, R.; Kolomiets, M. V., A non-JA producing oxophytodiene reductase functions in salicylic acid-mediated antagonism with jasmonic acid during pathogen attack. *Molecular plant pathology* **2023**, *24* (7), 725-741.
- Yan, Y.; Christensen, S.; Isakeit, T.; Engelberth, J.; Meeley, R.; Hayward, A.; Emery, R. J.; Kolomiets, M. V., Disruption of OPR7 and OPR8 reveals the versatile functions of jasmonic acid in maize development and defense. *Plant Cell* **2012**, *24* (4), 1420-36.
- Christensen, S. A.; Nemchenko, A.; Borrego, E.; Murray, I.; Sobhy, I. S.; Bosak, L.; DeBlasio, S.; Erb, M.; Robert, C. A.; Vaughn, K. A.; Herrfurth, C.; Tumlinson, J.; Feussner, I.; Jackson, D.; Turlings, T. C.; Engelberth, J.; Nansen, C.; Meeley, R.; Kolomiets, M. V., The maize lipoxygenase, ZmLOX10, mediates green leaf volatile, jasmonate and herbivore-induced plant volatile production for defense against insect attack. *The Plant journal : for cell and molecular biology* **2013**, *74* (1), 59-73.
- Huang, P. C.; Grunseich, J. M.; Berg-Falloure, K. M.; Tolley, J. P.; Koiwa, H.; Bernal, J. S.; Kolomiets, M. V., Maize OPR2 and LOX10 Mediate Defense against Fall Armyworm and Western Corn Rootworm by Tissue-Specific Regulation of Jasmonic Acid and Ketol Metabolism. *Genes* **2023**, *14* (9).
- Castano-Duque, L.; Loades, K. W.; Tooker, J. F.; Brown, K. M.; Paul Williams, W.; Luthe, D. S., A Maize Inbred Exhibits Resistance Against Western Corn Rootworm, *Diabrotica virgifera virgifera*. *Journal of chemical ecology* **2017**, *43* (11-12), 1109-1123.
- Köhl, J.; Kolnaar, R.; Ravensberg, W. J., Mode of Action of Microbial Biological Control Agents Against Plant Diseases: Relevance Beyond Efficacy. *Frontiers in plant science* **2019**, *10*.

19. Tiénébo, E. O.; Harrison, K.; Abo, K.; Brou, Y. C.; Pierson, L. S., 3rd; Tamborindeguy, C.; Pierson, E. A.; Levy, J. G., Mycorrhization Mitigates Disease Caused by "Candidatus Liberibacter solanacearum" in Tomato. *Plants (Basel, Switzerland)* **2019**, *8* (11).
20. El-Saadony, M. T.; Saad, A. M.; Soliman, S. M.; Salem, H. M.; Ahmed, A. I.; Mahmood, M.; El-Tahan, A. M.; Ebrahim, A. A. M.; Abd El-Mageed, T. A.; Negm, S. H.; Selim, S.; Babalghith, A. O.; Elrys, A. S.; El-Tarabily, K. A.; AbuQamar, S. F., Plant growth-promoting microorganisms as biocontrol agents of plant diseases: Mechanisms, challenges and future perspectives. *Frontiers in plant science* **2022**, *13*, 923880.
21. Zin, N. A.; Badaluddin, N. A., Biological functions of Trichoderma spp. for agriculture applications. *Annals of Agricultural Sciences* **2020**, *65* (2), 168-178.
22. Tyśkiewicz, R.; Nowak, A.; Ozimek, E.; Jaroszuk-Ściśeł, J., Trichoderma: The Current Status of Its Application in Agriculture for the Biocontrol of Fungal Phytopathogens and Stimulation of Plant Growth. *International journal of molecular sciences* **2022**, *23* (4).
23. Yao, X.; Guo, H.; Zhang, K.; Zhao, M.; Ruan, J.; Chen, J., Trichoderma and its role in biological control of plant fungal and nematode disease. *Frontiers in microbiology* **2023**, *14*, 1160551.
24. Anderson, A. J.; Kim, Y. C., Insights into plant-beneficial traits of probiotic Pseudomonas chlororaphis isolates. *Journal of medical microbiology* **2020**, *69* (3), 361-371.
25. Contreras-Cornejo, H. A.; Macías-Rodríguez, L.; del-Val, E.; Larsen, J., The root endophytic fungus Trichoderma atroviride induces foliar herbivory resistance in maize plants. *Applied Soil Ecology* **2018**, *124*, 45-53.
26. Poveda, J., Trichoderma as biocontrol agent against pests: New uses for a mycoparasite. *Biological Control* **2021**, *159*, 104634.
27. Djonović, S.; Pozo, M. J.; Dangott, L. J.; Howell, C. R.; Kenerley, C. M., Sm1, a proteinaceous elicitor secreted by the biocontrol fungus Trichoderma virens induces plant defense responses and systemic resistance. *Molecular plant-microbe interactions : MPMI* **2006**, *19* (8), 838-53.
28. Djonovic, S.; Vargas, W. A.; Kolomiets, M. V.; Horndeski, M.; Wiest, A.; Kenerley, C. M., A proteinaceous elicitor Sm1 from the beneficial fungus Trichoderma virens is required for induced systemic resistance in maize. *Plant physiology* **2007**, *145* (3), 875-89.
29. Lamdan, N. L.; Shalaby, S.; Ziv, T.; Kenerley, C. M.; Horwitz, B. A., Secretome of Trichoderma interacting with maize roots: role in induced systemic resistance. *Molecular & cellular proteomics : MCP* **2015**, *14* (4), 1054-63.
30. Wang, K. D.; Borrego, E. J.; Kenerley, C. M.; Kolomiets, M. V., Oxylinins Other Than Jasmonic Acid Are Xylem-Resident Signals Regulating Systemic Resistance Induced by Trichoderma virens in Maize. *Plant Cell* **2020**, *32* (1), 166-185.
31. Khan, R. A. A.; Najeeb, S.; Chen, J.; Wang, R.; Zhang, J.; Hou, J.; Liu, T., Insights into the molecular mechanism of Trichoderma stimulating plant growth and immunity against phytopathogens. *Physiologia plantarum* **2023**, *175* (6), e14133.
32. Dehbi, I.; Achemrk, O.; Ezzouggar, R.; El Jarroudi, M.; Mokrini, F.; Legrifi, I.; Belabess, Z.; Laasli, S. E.; Mazouz, H.; Lahlali, R., Beneficial Microorganisms as Bioprotectants against Foliar Diseases of Cereals: A Review. *Plants (Basel, Switzerland)* **2023**, *12* (24).
33. Saldaña-Mendoza, S. A.; Pacios-Michelena, S.; Palacios-Ponce, A. S.; Chávez-González, M. L.; Aguilar, C. N., Trichoderma as a biological control agent: mechanisms of action, benefits for crops and development of formulations. *World journal of microbiology & biotechnology* **2023**, *39* (10), 269.
34. Yuan, P.; Borrego, E.; Park, Y. S.; Gorman, Z.; Huang, P. C.; Tolley, J.; Christensen, S. A.; Blanford, J.; Kilaru, A.; Meeley, R.; Koiwa, H.; Vidal, S.; Huffaker, A.; Schmelz, E.; Kolomiets, M. V., 9,10-KODA, an α -ketol produced by the tonoplast-localized 9-lipoxygenase ZmLOX5, plays a signaling role in maize defense against insect herbivory. *Molecular plant* **2023**.
35. Yu, J. M.; Wang, D.; Pierson, L. S., 3rd; Pierson, E. A., Effect of Producing Different Phenazines on Bacterial Fitness and Biological Control in Pseudomonas chlororaphis 30-84. *The plant pathology journal* **2018**, *34* (1), 44-58.
36. Yuan, P.; Pan, H.; Boak, E. N.; Pierson, L. S., 3rd; Pierson, E. A., Phenazine-Producing Rhizobacteria Promote Plant Growth and Reduce Redox and Osmotic Stress in Wheat Seedlings Under Saline Conditions. *Frontiers in plant science* **2020**, *11*, 575314.
37. Harman, G. E.; Howell, C. R.; Viterbo, A.; Chet, I.; Lorito, M., Trichoderma species — opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology* **2004**, *2* (1), 43-56.
38. Lorito, M.; Woo, S. L.; Harman, G. E.; Monte, E., Translational research on Trichoderma: from 'omics to the field. *Annual review of phytopathology* **2010**, *48*, 395-417.
39. Woo, S. L.; Hermosa, R.; Lorito, M.; Monte, E., Trichoderma: a multipurpose, plant-beneficial microorganism for eco-sustainable agriculture. *Nature Reviews Microbiology* **2023**, *21* (5), 312-326.
40. Coppola, M.; Cascone, P.; Chiusano, M. L.; Colantuono, C.; Lorito, M.; Pennacchio, F.; Rao, R.; Woo, S. L.; Guerrieri, E.; Digilio, M. C., Trichoderma harzianum enhances tomato indirect defense against aphids. *Insect science* **2017**, *24* (6), 1025-1033.

41. Coppola, M.; Cascone, P.; Lelio, I. D.; Woo, S. L.; Lorito, M.; Rao, R.; Pennacchio, F.; Guerrieri, E.; Digilio, M. C., Trichoderma atroviride P1 Colonization of Tomato Plants Enhances Both Direct and Indirect Defense Barriers Against Insects. *Frontiers in physiology* **2019**, *10*.
42. Coppola, M.; Diretto, G.; Digilio, M. C.; Woo, S. L.; Giuliano, G.; Molisso, D.; Pennacchio, F.; Lorito, M.; Rao, R., Transcriptome and Metabolome Reprogramming in Tomato Plants by Trichoderma harzianum strain T22 Primes and Enhances Defense Responses Against Aphids. *Frontiers in physiology* **2019**, *10*.
43. Islam, M. S.; Subbiah, V. K.; Siddiquee, S., Efficacy of Entomopathogenic Trichoderma Isolates against Sugarcane Woolly Aphid, Ceratovacuna lanigera Zehntner (Hemiptera: Aphididae). *Horticulturae* **2022**, *8* (1), 2.
44. Jafarbeigi, F.; Samih, M. A.; Alaei, H.; Shirani, H., Induced Tomato Resistance Against Bemisia tabaci Triggered by Salicylic Acid, β -Aminobutyric Acid, and Trichoderma. *Neotropical entomology* **2020**, *49* (3), 456-467.
45. Muvea, A. M.; Meyhöfer, R.; Subramanian, S.; Poehling, H. M.; Ekesi, S.; Maniania, N. K., Colonization of onions by endophytic fungi and their impacts on the biology of Thrips tabaci. *PLoS One* **2014**, *9* (9), e108242.
46. Contreras-Cornejo, H. A.; del-Val, E.; Macías-Rodríguez, L.; Alarcón, A.; González-Esquivel, C. E.; Larsen, J., Trichoderma atroviride, a maize root associated fungus, increases the parasitism rate of the fall armyworm Spodoptera frugiperda by its natural enemy Campoletis sonorensis. *Soil Biology and Biochemistry* **2018**, *122*, 196-202.
47. Di Lelio, I.; Coppola, M.; Comite, E.; Molisso, D.; Lorito, M.; Woo, S. L.; Pennacchio, F.; Rao, R.; Digilio, M. C., Temperature Differentially Influences the Capacity of Trichoderma Species to Induce Plant Defense Responses in Tomato Against Insect Pests. *Frontiers in plant science* **2021**, *12*, 678830.
48. Papantoniou, D.; Vergara, F.; Weinhold, A.; Quijano, T.; Khakimov, B.; Pattison, D. I.; Bak, S.; van Dam, N. M.; Martínez-Medina, A., Cascading Effects of Root Microbial Symbiosis on the Development and Metabolome of the Insect Herbivore Manduca sexta L. *Metabolites* **2021**, *11* (11).
49. Kinyungu, S. W.; Agbessenou, A.; Subramanian, S.; Khamis, F. M.; Akutse, K. S., One stone for two birds: Endophytic fungi promote maize seedlings growth and negatively impact the life history parameters of the fall armyworm, Spodoptera frugiperda. *Frontiers in physiology* **2023**, *14*, 1253305.
50. Di Lelio, I.; Forni, G.; Magoga, G.; Brunetti, M.; Bruno, D.; Becchimanzi, A.; De Luca, M. G.; Sinno, M.; Barra, E.; Bonelli, M.; Frusciante, S.; Diretto, G.; Digilio, M. C.; Woo, S. L.; Tettamanti, G.; Rao, R.; Lorito, M.; Casartelli, M.; Montagna, M.; Pennacchio, F., A soil fungus confers plant resistance against a phytophagous insect by disrupting the symbiotic role of its gut microbiota. *Proceedings of the National Academy of Sciences of the United States of America* **2023**, *120* (10), e2216922120.
51. Pappas, M. L.; Samaras, K.; Koufakis, I.; Broufas, G. D., Beneficial Soil Microbes Negatively Affect Spider Mites and Aphids in Pepper. *Agronomy* **2021**, *11* (9), 1831.
52. Medeiros, H. A. d.; Araújo Filho, J. V. d.; Freitas, L. G. d.; Castillo, P.; Rubio, M. B.; Hermosa, R.; Monte, E., Tomato progeny inherit resistance to the nematode Meloidogyne javanica linked to plant growth induced by the biocontrol fungus Trichoderma atroviride. *Scientific Reports* **2017**, *7* (1), 40216.
53. Fan, H.; Yao, M.; Wang, H.; Zhao, D.; Zhu, X.; Wang, Y.; Liu, X.; Duan, Y.; Chen, L., Isolation and effect of Trichoderma citrinoviride Snf1910 for the biological control of root-knot nematode, Meloidogyne incognita. *BMC Microbiology* **2020**, *20* (1), 299.
54. Saharan, R.; Patil, J. A.; Yadav, S.; Kumar, A.; Goyal, V., The nematicidal potential of novel fungus, Trichoderma asperellum FbMi6 against Meloidogyne incognita. *Scientific Reports* **2023**, *13* (1), 6603.
55. Berini, F.; Caccia, S.; Franzetti, E.; Congiu, T.; Marinelli, F.; Casartelli, M.; Tettamanti, G., Effects of Trichoderma viride chitinases on the peritrophic matrix of Lepidoptera. *Pest Management Science* **2016**, *72* (5), 980-989.
56. Vijayakumar, N.; Alagar, S.; Madanagopal, N., Effects of chitinase from Trichoderma viride on feeding, growth and biochemical parameters of the rice moth, Corcyra cephalonica Stainton. *Journal of Entomology and Zoology Studies* **2016**, *4* (4), 520-523.
57. Massry, S.; Shokry, H. M.; Hegab, M. In *EFFICIENCY OF Trichoderma harzianum AND SOME ORGANIC ACIDS ON THE COTTON BOLLWORMS, Earias insulana AND Pectinophora gossypiella*, 2016.
58. Chinnaperumal, K.; Govindasamy, B.; Paramasivam, D.; Dilipkumar, A.; Dhayalan, A.; Vadivel, A.; Sengodan, K.; Pachiappan, P., Bio-pesticidal effects of Trichoderma viride formulated titanium dioxide nanoparticle and their physiological and biochemical changes on Helicoverpa armigera (Hub.). *Pesticide Biochemistry and Physiology* **2018**, *149*, 26-36.
59. Ghosh, S. K.; Pal, S., Entomopathogenic potential of Trichoderma longibrachiatum and its comparative evaluation with malathion against the insect pest Leucinodes orbonalis. *Environmental Monitoring and Assessment* **2015**, *188* (1), 37.
60. Zhou, D.; Huang, X.-F.; Guo, J.; dos-Santos, M. L.; Vivanco, J. M., Trichoderma gamsii affected herbivore feeding behaviour on Arabidopsis thaliana by modifying the leaf metabolome and phytohormones. *Microbial Biotechnology* **2018**, *11* (6), 1195-1206.

61. Wang, K. D.; Gorman, Z.; Huang, P. C.; Kenerley, C. M.; Kolomiets, M. V., Trichoderma virens colonization of maize roots triggers rapid accumulation of 12-oxophytodienoate and two γ -ketols in leaves as priming agents of induced systemic resistance. *Plant signaling & behavior* **2020**, *15* (9), 1792187.
62. Yan, Y.; Huang, P.-C.; Borrego, E.; Kolomiets, M., New perspectives into jasmonate roles in maize. *Plant signaling & behavior* **2014**, *9* (10), e970442-e970442.
63. Huang, H.; Liu, B.; Liu, L.; Song, S., Jasmonate action in plant growth and development. *Journal of Experimental Botany* **2017**, *68* (6), 1349-1359.
64. Wang, J.; Song, L.; Gong, X.; Xu, J.; Li, M., Functions of Jasmonic Acid in Plant Regulation and Response to Abiotic Stress. *International journal of molecular sciences* **2020**, *21* (4).
65. Bažok, R.; Lemić, D.; Chiarini, F.; Furlan, L., Western Corn Rootworm (*Diabrotica virgifera virgifera* LeConte) in Europe: Current Status and Sustainable Pest Management. *Insects* **2021**, *12* (3).
66. Paddock, K. J.; Robert, C. A. M.; Erb, M.; Hibbard, B. E., Western Corn Rootworm, Plant and Microbe Interactions: A Review and Prospects for New Management Tools. *Insects* **2021**, *12* (2).
67. Carrière, Y.; Brown, Z.; Aglasan, S.; Dutilleul, P.; Carroll, M.; Head, G.; Tabashnik, B. E.; Jørgensen, P. S.; Carroll, S. P., Crop rotation mitigates impacts of corn rootworm resistance to transgenic Bt corn. *Proceedings of the National Academy of Sciences* **2020**, *117* (31), 18385-18392.
68. Baek, J.; Kenerley, C., Detection and enumeration of a genetically modified fungus in soil environments by quantitative competitive polymerase chain reaction. *FEMS Microbiology Ecology* **2006**, *25*, 419-428.
69. Schneider, C. A.; Rasband, W. S.; Eliceiri, K. W., NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* **2012**, *9* (7), 671-675.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.