

Article

Not peer-reviewed version

---

# Finding All Efficient Solutions of an Interval Multiple Objective Linear Programming Problem

---

[Ta Van Tu](#) \*

Posted Date: 21 March 2024

doi: 10.20944/preprints202403.1314.v1

Keywords: Interval multiple objective linear programming; the efficient basic set; the efficient set



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# Finding All Efficient Solutions of an Interval Multiple Objective Linear Programming Problem

Ta Van Tu

Department of Operations Research, Corvinus University of Budapest, H-1093 Budapest, Hungary;  
t\_vantu@yahoo.com

**Abstract:** In this paper, we propose a method to find all efficient solutions of an interval multiple objective linear programming (IMOLP) problem with interval coefficients in the objective functions, the constraint matrix and the right-hand side vector. The set of all efficient solutions of an IMOLP problem can completely supply information to the decision makers in finding their most preferred solutions but, up to now, not all solutions of an IMOLP problem can be found by known methods. A numerical example is given to illustrate the working of the proposed method.

**Keywords:** interval multiple objective linear programming; the efficient basic set; the efficient set

**MSC:** 90C29; 90C50; 90C90

## 1. Introduction

Multiple objective linear programming (MOLP) models play an important role in solving and investigating real-life practical problems. There is a practical fact that the exact values of the coefficients of an MOLP model are very difficult to determine but intervals containing them can be easily determined. Thus, practical problems can be described more correctly and more easily by using interval multiple objective linear programming (IMOLP) models than using MOLP models. For brevity of presentation, we shall use the following notation: For two vectors  $a = (a_1, \dots, a_n)^T$  and  $b = (b_1, \dots, b_n)^T$ ,  $a \leq b$  if and only if  $a_i \leq b_i$  for all  $i = 1, \dots, n$ .

An interval multiple objective linear programming (IMOLP) problem, denoted by  $P(IA, IC, ib)$ , can be stated as follows:

$$\text{"maximize"} \quad Cx, \quad (1)$$

$$Ax \leq b, \quad (2)$$

$$C \in IC, \quad A \in IA, \quad b \in ib, \quad (3)$$

where  $IA = \{A | \underline{A} \leq A \leq \bar{A}\}$  is an interval matrix in  $R^{m \times n}$ ,  $IC = \{C | \underline{C} \leq C \leq \bar{C}\}$  is an interval matrix in  $R^{k \times n}$ ,  $ib = \{b | \underline{b} \leq b \leq \bar{b}\}$  is an interval column vector in  $R^{m \times 1}$ ,  $\underline{A}$ ,  $\bar{A}$ ,  $\underline{C}$ ,  $\bar{C}$ ,  $\underline{b}$  and  $\bar{b}$  are determined.

For every  $A \in IA$ ,  $C \in IC$  and  $b \in ib$  a multiple objective linear programming (MOLP) problem, denoted by  $P(A, C, b)$ , is obtained from problem (1)-(3). Let  $L(A, b)$  be the feasible set of  $P(A, C, b)$ . A point  $x \in L(A, b)$  is called *efficient* for  $P(A, C, b)$  if there is no  $y \in L(A, b)$  such that  $Cx \leq Cy$  and  $Cx \neq Cy$ . A feasible point of problem (1)-(3) is called *efficient* for it if there are  $A \in IA$ ,  $C \in IC$  and  $b \in ib$  such that it is efficient for  $P(A, C, b)$ .

Let  $L(IA, ib)$  be the feasible set of problem (1)-(3),  $E(A, C, b)$  and  $E(IA, IC, ib)$  be the efficient sets of  $P(A, C, b)$  and  $P(IA, IC, ib)$ , respectively. It is easily seen that

$$L(IA, ib) = \left\{ x \in R^m \mid \begin{array}{l} Ax \leq b, \\ A \in IA, b \in ib \end{array} \right\}, \quad (4)$$

$$L(IA, ib) = \cup \{ L(A, b) \mid A \in IA, b \in ib \}, \quad (5)$$

$$E(IA, IC, ib) = \cup \{ E(A, C, b) \mid A \in IA, C \in IC, b \in ib \}. \quad (6)$$

The notion of efficient solutions of IMOLP problem (1)-(3) can be found in, for example, Tu [34], Allahdadi and Batamiz [1]. Another name of an efficient solution of IMOLP problem (1)-(3) is *possibly efficient* used in, for example, Bitran [6], Inuiguchi and Kume [17], Oliveira and Antunes [24]. There are many known methods for finding the efficient set of an MOLP problem, see, for example, Yu and Zeleny [40], Isermann [19], Ecker et al. [10], Dauer and Liu [9], Armand and Malivert [3], Armand [2], Sayin [32], Dauer and Gallagher [8], Benson [5], Tu ([34–38]), Yan et al. [39], Foroughi and Jafari [12], Pourkarimi et al. [25], Krichen et al. [21], Rudloff et al. [31], Tohidi and Hassasi [33]. It can be easily seen that all the known methods for finding the efficient set of an MOLP problem must recomputed the efficient set when the data of the MOLP problem is changed. Therefore, they cannot find the efficient set of an IMOLP problem. Solving an IMOLP problem is much more difficult than solving an MOLP problem. Methods to solve and investigate an IMOLP problem are increasingly developed. Chanas and Kuchta [7], Ishibuchi and Tanaka [20] study interval coefficients only in objective functions, Tu [34] investigate interval coefficients only in the right-hand side vector. Some efficient solutions can be found by methods given by Urli and Nadeau [41] based on an interactive method, by Inuiguchi and Kume [18] based on a goal programming method, by Rivaz and Yaghoobi [27] based on a weighted sum of maximum regrets, Hajiagha et al. [14], Rivaza and Saeidib [26] based on fuzzy programming methods, etc.. Efficient extreme points of an IMOLP problem are dealt with in Inuiguchi and Kume [17]. If the coefficients of an IMOLP problem are satisfied probability distributions, then stochastic programming methods can be used to study this problem, see, for example, Batamiz and Allahdadi [4]. Theoretically, an IMOLP problem can be stated based on fuzzy numbers and can be solved by fuzzy programming methods. However, huge difficulties in this way lie in constructing adequate membership functions and finding all efficient solutions of an IMOLP problem.

In this paper, we propose a method to determine all efficient solutions of an IMOLP problem. This method is developed based on the methods of Tu [34,35] and can be easily implemented.

The paper is organized as follows: A method for determining the efficient set of an IMOLP problem is presented in Section 2 and an example is given in Section 3.

## 2. A Method for Determining the Efficient Set of an IMOLP Problem

### 2.1. A General Case

Corresponding to IMOLP problem (1)-(3), we consider the following set:

$$\tilde{L}(IA, IC) = \left\{ p = (y, z)^T \in R^{m+k} \mid \begin{array}{l} y^T A - z^T C = e^T C, y \geq 0, z \geq 0, \\ A \in IA, C \in IC \end{array} \right\}, \quad (7)$$

where  $e = (1, \dots, 1)^T \in R^k$ . For every  $A \in IA$  and  $C \in IC$ , the following convex polyhedron is directly obtained from (7):

$$\tilde{L}(A, C) = \left\{ p = (y, z)^T \in R^{m+k} \mid y^T A - z^T C = e^T C, y \geq 0, z \geq 0 \right\}$$

Thus, it is clear that

$$\tilde{L}(IA, IC) = \cup \{ \tilde{L}(A, C) \mid A \in IA, C \in IC \}. \quad (8)$$

A formula presented in the following property to describe the solution set of an interval linear equation is given by Oettli-Prager [23]:

**Property 2.1.**  $X = \{x \mid |A_c x - b_c| \leq \Delta |x| + \delta\}$ ,

where  $X = \{x \in R^n \mid Ax = b, A \in IA, b \in ib\}$ ,  $A_c = (\underline{A} + \bar{A})/2$ ,  $b_c = (\underline{b} + \bar{b})/2$ ,  $\Delta = (\bar{A} - \underline{A})/2$ ,

$\delta = (\bar{b} - \underline{b})/2$ ,  $|x| = (|x_1|, \dots, |x_n|)^T$ ,  $IA$  and  $ib$  are defined in problem (1)-(3).

This property is proven by Rohn ([29], Theorem 2.1, p.43) in a special case when  $n = m$  but his proof can be easily modified to prove Property 2.1. In the case when  $x \geq 0$ , from Property 2.1 the following property is easily obtained:

**Property 2.2.** If  $x \geq 0$ , then  $X = \left\{ x \in R^n \mid \begin{array}{l} \underline{A}x \leq \bar{b}, \\ -\underline{A}x \leq -\underline{b} \end{array} \right\}$ .

The solution set of an interval linear inequality with positive variables is given in the following property whose proof is obvious:

**Property 2.3.**  $IS(IA, ib) = \{x \in R^n \mid \underline{A}x \leq \bar{b}, x \geq 0\}$ ,

where

$$IS(IA, ib) = \left\{ x \in R^n \mid \begin{array}{l} Ax \leq b, x \geq 0, \\ A \in IA, b \in ib \end{array} \right\}.$$

Noting that the variables in (7) are non-negative, based on Property 2.2 the following property can be easily obtained:

**Property 2.4.**  $\tilde{L}(IA, IC) = \bar{L}$ ,

where

$$\bar{L} = \left\{ p = (y, z)^T \in R^{m+k} \mid \begin{array}{l} y^T \underline{A} - z^T \bar{C} \leq e^T \bar{C}, \\ -y^T \bar{A} + z^T \underline{C} \leq -e^T \underline{C} \\ y \geq 0, z \geq 0 \end{array} \right\}.$$

This property is also presented in Li et al ([22], Theorem 2.5), Rohn [30].

**Remark 2.1.** Since  $\bar{L}$  is a convex polyhedron described by a system of linear inequalities with non-negative variables,  $\bar{L}$  has an extreme point if and only if it is not empty.

In order to find extreme points of  $\bar{L}$  based on the simplex method,  $\bar{L}$  is stated in the following form:

$$\bar{\bar{L}} = \left\{ p = (y, y^1, y^2, z)^T \in R^{m+2n+k} \mid \begin{array}{l} y^T \underline{A} + I_n y^1 - z^T \bar{C} = e^T \bar{C}, \\ -y^T \bar{A} + I_n y^2 + z^T \underline{C} = -e^T \underline{C}, \\ y \geq 0, y^1 \geq 0, y^2 \geq 0, z \geq 0 \end{array} \right\}.$$

Let

$$T_0(IA, IC) = \left\{ p \mid p = (y, y^1, y^2, z)^T \text{ is an extreme point of } \bar{\bar{L}} \right\},$$

$$T_1(IA, IC) = \{I_+(p) \mid p \in T_0(IA, IC)\},$$

where  $I_+(p) = \{i \in \{1, \dots, m\} | p_i > 0\}$  and  $p_i$  is the  $i$ -th component of  $p$ ;

$T_2(IA, IC)$  is a set consisting of all minimal elements of  $T_1(IA, IC)$  by inclusion.

Let  $T_2(A, C)$  be a set established based on  $\tilde{L}(A, C)$  by a way similar to that used to establish  $T_2(IA, IC)$  based on  $\bar{L}$ .

It can be easily seen that the set  $T_2(IA, IC)$  can be found by the method given in Tu [34] without determining all extreme points of  $\bar{L}$ .

A relation between  $T_2(IA, IC)$  and  $T_2(A, C)$  is considered in the following property:

**Property 2.5.** For every  $I \in T_2(A, C)$  there is  $J \in T_2(IA, IC)$  such that  $J \subseteq I$ .

**Proof.** There is an extreme point  $p^0$  of  $\tilde{L}(A, C)$  such that  $I_+(p^0) = I$ . Noting that  $p^0 \in \tilde{L}(IA, IC)$ , from (8) and Property 2.4 it follows that  $p^0 \in \bar{L}$ . Based on a proof similar to that of Property 2.4 in [34], it can be easily seen that there is an extreme point  $p^1$  of  $\bar{L}$  such that  $I_+(p^1) \subseteq I_+(p^0)$ . From the definition of  $T_2(IA, IC)$  it follows that there is  $J \in T_2(IA, IC)$  such that  $J \subseteq I_+(p^1)$ . Therefore,  $J \subseteq I$ .  $\square$

Let

$$S(I) = \{x \in L(IA, ib) | a_i x = b_i, i \in I\},$$

$$S(I, A, b) = \{x \in L(A, b) | a_i x = b_i, i \in I\},$$

where  $(a_i, b_i)$  is the  $i$ -th row of a matrix  $(A, b)$  defined in (4).

A formula to compute the efficient set of IMOLP problem (1)-(3) is shown in the following property:

**Property 2.6.**  $E = \bigcup_{I \in T_2(IA, IC)} S(I)$ .

**Proof.** For every element  $x^0 \in E$  there are  $A \in IA$ ,  $C \in IC$  and  $b \in ib$  such that  $x^0 \in E(A, C, b)$ . Based on Property 2.4 in [34], there is  $J \in T_2(A, C)$  such that  $J \subseteq ID(x^0, A, b)$ , where

$$ID(x^0, A, b) = \{i \in \{1, \dots, m\} | a_i x^0 = b_i\}.$$

Therefore,  $x^0 \in S(J, A, b)$ . From Property 2.5 it follows that there is  $I \in T_2(IA, IC)$  such that  $I \subseteq J$ . Thus,  $x^0 \in S(J, A, b) \subseteq S(I, A, b) \subseteq S(I)$ . Therefore,  $E \subseteq \bigcup_{I \in T_2(IA, IC)} S(I)$ .

Conversely, for every element  $x^0 \in \bigcup_{I \in T_2(IA, IC)} S(I)$  there is  $I \in T_2(IA, IC)$  such that  $x^0 \in S(I)$ .

Based on Property 2.4 and (8), from  $I \in T_2(IA, IC)$  it follows that there are  $A \in IA$ ,  $C \in IC$  and an extreme point  $p^0 = (y^0, z^0)^T \in \tilde{L}(A, C)$  such that  $I_+(p^0) = I$ . It is clear that  $x^0 \in S(I, A, b^0)$ , where  $b^0 = Ax^0$ . From  $x^0 \in S(I, A, b^0)$  it follows that  $I \subseteq ID(x^0, A, b^0)$ . Noting that  $I_+(p^0) = I$  and  $p^0 = (y^0, z^0)^T$ , it can be easily seen that  $y^0 \in \tilde{F}(ID(x^0, A, b^0), x^0)$ ,

where

$$\tilde{F}\left(ID(x^0, A, b^0), x^0\right) = \left\{ y \in R^m \left| \begin{array}{l} y^T A = (e + z^0)^T C, \\ y_i = 0 \text{ for all } i \in \{1, \dots, m\} \setminus ID(x^0, A, b^0), \\ y_i \geq 0 \text{ for all } i \in ID(x^0, A, b^0) \end{array} \right. \right\}.$$

Thus, based on the complementary theorem of linear programming,  $x^0$  is an optimal solution of the linear programming problem  $\max \left\{ (e + z^0)^T Cx \mid x \in L(A, b^0) \right\}$ . Therefore,  $x^0 \in E(A, C, b^0)$ . It is clear that  $b^0 \in ib$ . Therefore,  $x^0 \in E$ . The proof is complete.  $\square$

Since the solution set of an interval linear equation, in general, is not convex, see, for example, Hensen [15], Fiedler et al. [11], Rohn [28,30]. Therefore, the sets  $S(I)$  defined in Property 2.6 can be not convex polyhedrons. This can cause difficulties in finding most preferred solutions from the efficient set of IMOLP problem (1)-(3).

Now we consider an IMOLP problem of which the efficient set can be computed by a union of a finite number of convex polyhedrons.

## 2.2. A Special Case

We consider the following IMOLP problem:

$$\text{“maximize” } Cx, \quad (9)$$

$$A^1 x \leq b^1, \quad x \geq 0, \quad (10)$$

$$C \in IC, \quad A^1 \in IA^1, \quad b^1 \in ib^1, \quad (11)$$

where  $IA^1 = \left\{ A^1 \mid \underline{A}^1 \leq A^1 \leq \bar{A}^1 \right\}$  is an interval matrix in  $R^{m_2 \times n}$ ,  $IC = \left\{ C \mid \underline{C} \leq C \leq \bar{C} \right\}$  is an interval matrix in  $R^{k \times n}$ ,  $ib^1 = \left\{ b^1 \mid \underline{b}^1 \leq b^1 \leq \bar{b}^1 \right\}$  is an interval column vector in  $R^{m_2 \times 1}$ . Problem (9)-(11) is a special case of IMOLP problem (1)-(3) because its variables are restricted in sign. IMOLP problem (9)-(11) can be easily solved by the above presented method for solving problem (1)-(3). To do this, we restate problem (9)-(11) in the form of problem (1)-(3) by defining  $A = \begin{pmatrix} A^1 \\ -I_{n \times n} \end{pmatrix} \in R^{(m_2+n) \times n}$ ,  $\underline{A} = \begin{pmatrix} \underline{A}^1 \\ -I_{n \times n} \end{pmatrix} \in R^{(m_2+n) \times n}$ ,  $\bar{A} = \begin{pmatrix} \bar{A}^1 \\ -I_{n \times n} \end{pmatrix} \in R^{(m_2+n) \times n}$ ,  $b = \begin{pmatrix} b^1 \\ O_{n \times 1} \end{pmatrix} \in R^{(m_2+n) \times 1}$ ,  $\underline{b} = \begin{pmatrix} \underline{b}^1 \\ O_{n \times 1} \end{pmatrix} \in R^{(m_2+n) \times 1}$ ,  $\bar{b} = \begin{pmatrix} \bar{b}^1 \\ O_{n \times 1} \end{pmatrix} \in R^{(m_2+n) \times 1}$  and  $m = m_2 + n$ , where  $I_{n \times n}$  is the unit matrix in  $R^n$  and  $O_{n \times 1}$  is the  $n$  column vector with components being 0. Thus, the efficient set of problem (9)-(11), denoted by  $E_+$ , can be computed by the formula given in Property 2.6. Now we represent this formula with using the data of problem (9)-(11).

**Property 2.7.**  $E_+ = \bigcup_{I \in T_2(IA, IC)} S_+(I)$ ,

where  $(\bar{a}_i, \underline{b}_i)$  is the  $i$ -th row of the matrix  $(\bar{A}^1, \underline{b}^1)$  and

$$S_+(I) = \left\{ x \in R^n \left| \begin{array}{l} -\bar{a}_i x \leq -\underline{b}_i, i \in I \cap \{1, \dots, m_2\}, \\ x_i = 0, \text{ if } (i + m_2) \in I \text{ and } i \in \{1, \dots, n\}, \\ \underline{A}^1 x \leq \bar{b}^1, \\ x \geq 0 \end{array} \right. \right\}$$

**Proof.** Let  $A(I), IA(I), b(I)$  and  $ib(I)$  be the matrices obtained from the matrices  $A, IA, b$  and  $ib$  by dropping rows whose indices are not in  $I$ , respectively. Based on Properties 2.2 and 2.3, it

can be easily seen that  $S(I) = \{x \in L(IA, ib) | a_i x = b_i, i \in I\} = \left\{ x \in R^n \begin{cases} a_i x = b_i, i \in I \\ Ax \leq b, x \geq 0, \\ A \in IA, b \in ib \end{cases} \right. =$

$$\left\{ x \in R^n \begin{cases} A(I)x = b(I), x \geq 0, \\ A(I) \in IA(I), b(I) \in ib(I), \\ Ax \leq \bar{b} \end{cases} \right\} = \left\{ x \in R^n \begin{cases} -\bar{a}_i x \leq -\bar{b}_i, i \in I, \\ a_i x \leq \bar{b}_i, i \in I, \\ Ax \leq \bar{b}, \\ x \geq 0 \end{cases} \right\} \left( (a_i, \bar{b}_i) \text{ and } (\bar{a}_i, \bar{b}_i) \text{ are the } i\text{-th rows} \right.$$

of the matrices  $(\underline{A}, \bar{b})$  and  $(\bar{A}, \underline{b})$ , respectively)  $= \left\{ x \in R^n \begin{cases} -\bar{a}_i x \leq -\bar{b}_i, i \in I, \\ Ax \leq \bar{b}, \\ x \geq 0 \end{cases} \right. =$

$$\left\{ x \in R^n \begin{cases} -\bar{a}_i x \leq -\bar{b}_i, i \in I \cap \{1, \dots, m_2\}, \\ x_i = 0, \text{ if } (i + m_2) \in I \text{ and } i \in \{1, \dots, n\}, \\ \underline{A}x \leq \bar{b}^1, \\ x \geq 0 \end{cases} \right\}. \text{ Based on Property 2.6, the proof is complete.}$$

□

**Remark 2.2.** IMOLP problem (9)-(11) is a popular one used in investigating practical problems because the condition of the variables is natural. Since its efficient set can be computed by the union of convex polyhedrons, finding most preferred solutions based on IMOLP problem (9)-(11) has many advantages.

**Remark 2.3.** Interval linear programming (ILP) problems are extensively investigated by many researchers, for example, Garajova and Hladik [13], Hladik [16]. Since ILP problems are a special case of IMOLP problems, the above presented results also validate for ILP problems.

### 3. Examples

We consider problem (9)-(11) when

$$IC = \begin{pmatrix} [0.5, 1.8] & [-0.5, 0.5] \\ [0.3, 0.8] & [1, 1.2] \end{pmatrix}, \quad IA^1 = \begin{pmatrix} [1.5, 2.5] & [0.5, 1] \\ [0.5, 2] & [3, 6] \end{pmatrix} \text{ and } ib^1 = \begin{pmatrix} [6, 10] \\ [14, 16] \end{pmatrix}.$$

This example is recalled from Oliveira and Antunes [24]. It is clear that

$$\bar{L} = \left\{ p = (y, z)^T \in R^{10} \begin{cases} 1.5y_1 + .5y_2 - y_3 + y_1^1 - 1.8z_1 - .8z_2 = 2.6, \\ .5y_1 + 3y_2 - y_4 + y_2^1 - .5z_1 - 1.2z_2 = 1.7, \\ -2.5y_1 - 2y_2 + y_3 + y_1^2 + .5z_1 + .3z_2 = -.8, \\ -y_1 - 6y_2 + y_4 + y_2^2 - .5z_1 + z_2 = -.5, \\ y \geq 0, y^1 \geq 0, y^2 \geq 0, z \geq 0 \end{cases} \right\}.$$

Using Lips-1.9.4 software in MATLAB, based on the method given in Tu [35], we can easily obtain  $T_2(IA, IC) = \{\{1\}, \{2\}\}$  after solving 4 linear programming problems. Based on Properties 2.7,

we have  $E_+ = S_+(\{1\}) \cup S_+(\{2\})$  , where  $S_+(\{1\}) = \left\{ x \in R^2 \begin{cases} -2.5x_1 + x_2 \leq -6, \\ 1.5x_1 + .5x_2 \leq 10, \\ .5x_1 + 3x_2 \leq 16, \\ x_1 \geq 0, x_2 \geq 0 \end{cases} \right\}$  ,

$$S_+(\{2\}) = \left\{ x \in R^2 \begin{cases} -2x_1 + 6x_2 \leq -14, \\ 1.5x_1 + .5x_2 \leq 10, \\ .5x_1 + 3x_2 \leq 16, \\ x_1 \geq 0, x_2 \geq 0 \end{cases} \right\}.$$

#### 4. Conclusions

We propose a method to find all efficient solutions of an IMOLP problem with interval coefficients in objective functions, the constraint matrix and the right-hand side vector. The set of all efficient solutions of an IMOLP problem can completely supply information to the decision makers in finding their most preferred solutions but, up to now, only some solutions of an IMOLP problem can be found by known methods. The proposed method is simple, is easy to implement and illustrated by a numerical example.

#### References

1. M. Allahdadi and A. Batamiz, Generation of some methods for solving interval multi-objective linear programming models. *OPSEARCH* **2021**, 58, 1077-1115.
2. P. Armand, Finding all maximal efficient faces in multiobjective linear programming. *Math. Program.* **1993**, 61, 357-375.
3. P. Armand and C. Malivert, Determination of the efficient set in multiobjective linear programming. *J. Optim. Theory and Appl.* **1991**, 70, 467-489.
4. A. Batamiz and M. Allahdadi, Finding efficient solutions in the interval multi-objective linear programming models, *Yugosl. J. Oper. Res.* **2021**, 31, 95-119.
5. H.P. Benson, Hybrid approach for solving multi-objective linear programs in outcome space. *J. Optim. Theory Appl.* **1998**, 98, 17-35.
6. G.R. Bitran, Linear multiobjective problems with interval coefficients, *Management Science* **1980**, 26, 694 – 706.
7. S. Chanas and D. Kuchta, Multiobjective programming in optimization of interval objective functions – A generalized approach, *European J. Oper. Res.* **1996**, 94, 594 – 598.
8. J.P. Dauer and R.J. Gallagher, A combined constraint-space, objective-space approach for determining high-dimensional maximal efficient faces of multiple objective linear programs. *European J. Oper. Res.* **1996**, 88, 368-381.
9. J.P. Dauer and Y.H. Liu, Solving multiple objective linear programs in objective space. *European J. Oper. Res.* **1990**, 46, 350-357.
10. J.G. Ecker, N.S. Hegner and I.A. Kouada, Generating all maximal efficient faces for multiple objective linear Programs. *J. Optim. Theory and Appl.* **1980**, 30, 353-381.
11. M. Fiedler, J. Nedoma, J. Ramík, J. Rohn and K. Zimmermann, Linear optimization problems with inexact data. Springer, New York, **2006**.
12. A.A. Foroughi and Y. Jafari, A modified method for constructing efficient solutions structure of molp. *Appl. Math. Model.* **2009**, 3, 2403-2410.
13. E. Garajova and M. Hladik, On the optimal solution set in interval linear programming. *Comput. Optim. Appl.* **2019**, 72, 269-292.
14. S.H.R. Hajiagha, H.A. Mahdiraji and S.S. Hashemi, Multi-objective linear programming with interval coefficients: A fuzzy set based approach. *Kybernetes* **2013**, 42, 482-496.
15. E. Hansen, On linear algebraic equations with interval coefficients, in Topics in Interval Analysis (E. Hansen, Ed.), Oxford U.P., Oxford, **1969**.
16. M. Hladik, Interval linear programming: A survey. Nova Science Publishers, New York, **2012**.
17. M. Inuiguchi and Y. Kume, A discrimination method of possibly efficient extreme points for interval multiobjective linear programming. *Transactions of the Society of Instrument Control Engineers* **1989**, 25, 823– 825.
18. M. Inuiguchi and Y. Kume, Goal programming problems with interval coefficients and target intervals. *European J. Oper. Res.* **1991**, 52, 345-360.

19. H. Isermann, The enumeration of the set of all efficient solutions for a linear multiple objective program. *Oper. Res. Q.* **1977**, 28, 711-725.
20. H. Ishibuchi and H. Tanaka, Multiobjective programming in optimization of the interval objective function. *European J. Oper. Res.* **1990**, 48, 219-225.
21. S. Krichen, H. Masri and A. Guitouni, Adjacency based method for generating maximal efficient faces in molp. *Appl. Math. Model.* **2012**, 36, 6301-6311.
22. H. Li, J. Luo and Q. Wang, Solvability and feasibility of interval linear equations and inequalities, *Linear Algebra and its Appl.* **2014**, 463, 78-94.
23. W. Oettli and W. Prager, Compatibility of approximate solution of linear equations with given error bounds for coefficients and right-hand sides, *Numerische Mathematik* **1964**, 6, 405-409.
24. C. Oliveira and C.H. Antunes, Multiple objective linear programming models with interval coefficients—an illustrative overview. *European J. Oper. Res.* **2007**, 181, 1434-1463.
25. L. Pourkarimi, M.A. Yaghoobi and M. Mashinchi, Determining maximal efficient faces in multiobjective linear programming problem. *J. Math. Anal. Appl.* **2009**, 354, 234-248.
26. S. Rivaza and Z. Saeidib, Solving Multiobjective Linear Programming Problems with Interval Parameters, *Fuzzy Information and Engineering* **2021**, 13, 497-504.
27. S. Rivaz and M.A. Yaghoobi, Weighted sum of maximum regrets in an interval MOLP problem. *Int Trans Oper Res.* **2018**, 25, 1659-1676.
28. J. Rohn, Solvability of systems of linear interval equations, *SIAM J. Matrix Anal. Appl.* **2003**, 25, 237-245.
29. J. Rohn, Systems of linear interval equations, *Linear Algebra Appl.* **1989**, 126, 39-78.
30. J. Rohn, Solvability of systems of interval linear equations and inequalities, in: M. Fiedler, J. Ne-doma, J. Ramík, J. Rohn, K. Zimmermann (Eds.), *Linear Optimization Problems with Inexact Data*, 35-77 (Chapter 2). Springer, New York, **2006**.
31. B. Rudloff, F. Ulus and R.J. Vanderbei, A parametric simplex algorithm to solve linear vector optimization problems, *Math. Programming*, vol. 163, pp. 213-242, 2017.
32. S. Sayin, An algorithm based on facial decomposition for finding the efficient set in multiple objective linear Programming. *Oper. Res. Lett.* **1996**, 19, 87-94.
33. G. Tohidi and H. Hassasi, An adjacency based local top-down search method for finding maximal efficient faces in multiple objective linear programming. *Naval Res. Logist.* **2018**, 65, 203-217.
34. T.V. Tu, Optimization over the efficient set of a parametric multiple objective linear programming Problem. *European J. Oper. Res.* **2000**, 122, 570-583.
35. T.V. Tu, A common formula to compute the efficient sets of a class of multiple objective linear programming problems. *Optimization* **2015**, 64, 2065-2092.
36. T.V. Tu, The maximal descriptor index set for a face of a convex polyhedral set and some applications. *J. Math. Anal. Appl.* **2015**, 429, 395-414.
37. T.V. Tu, A new method for determining all maximal efficient faces in multiple objective linear programming. *Acta Math. Vietnam.* **2017**, 42, 1-25.
38. T.V. Tu, A combined top-down and bottom-up search method for determining all maximal efficient faces in multiple objective linear programming. **2023**. <https://doi.org/10.21203/rs.3.rs-2934665/v1>.
39. H. Yan, Q. Wei and J. Wang Constructing efficient solutions structure of multiobjective linear programming. *J. Math. Anal. Appl.* **2005**, 307, 504-523.
40. P.L. Yu and M. Zeleny, The set of all nondominated solutions in linear cases and a multicriteria simplex method. *J. Math. Anal. Appl.* **1975**, 49, 430-468.
41. B. Urli and R. Nadeau, An interactive method to multiobjective linear programming problems with interval coefficients, *INFOR* **1992**, 30, 127 - 137.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.