

Article

Not peer-reviewed version

---

# *In vitro* embryos of Romosinuano and Tropical Milking cattle during three seasons in Veracruz, Mexico

---

[Froylan Rosales-Martínez](#), [Carlos Miguel Becerril-Pérez](#)<sup>\*</sup>, Adalberto Rosendo-Ponce, Alberto Riaño-Gaya, [César Cortez-Romero](#), [Jaime Gallegos-Sánchez](#), Salvador Romo-García

Posted Date: 5 April 2024

doi: 10.20944/preprints202404.0398.v1

Keywords: Adapted breeds; climatic change; genetic resources; hot climates; OPU-IVF; tropical region



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# *In vitro* embryos of Romosinuano and Tropical Milking Cattle during Three Seasons in Veracruz, Mexico

Froylan Rosales-Martínez <sup>1</sup>, Carlos Miguel Becerril-Pérez <sup>2,3,\*</sup>, Adalberto Rosendo-Ponce <sup>2</sup>, Alberto Riaño-Gaya <sup>4</sup>, César Cortez-Romero <sup>5</sup>, Jaime Gallegos-Sánchez <sup>3</sup> and Salvador Romo-García <sup>6</sup>

<sup>1</sup> Universidad Autónoma de Chiapas, Facultad Maya de Estudios Agropecuarios, Catazajá, Chiapas, México. C.P. 29980, froylan.rosales@unach.mx

<sup>2</sup> Colegio de Postgraduados, Programa de Agroecosistemas Tropicales, Campus Veracruz, Manlio Fabio Altamirano, Veracruz, México. C.P. 94251, arosendo@colpos.mx

<sup>3</sup> Colegio de Postgraduados, Programa de Ganadería, Campus Montecillo, Texcoco, Estado de México, México. C.P. 56230, gallegos@colpos.mx

<sup>4</sup> Reproducción Genética Avanzada RGA *In vitro*, Boca del Río, Veracruz, México. C.P. 94299, rga\_in\_vitro@hotmail.com

<sup>5</sup> Colegio de Postgraduados, Campus San Luis, Salinas de Hidalgo, San Luis Potosí, México. C.P. 78622; ccortez@colpos.mx

<sup>6</sup> Universidad Nacional Autónoma de México, Facultad de Estudios Superiores de Cuautitlán, Estado de México, México. C.P. 54714, sromo\_99@yahoo.com

\* Correspondence: color@colpos.mx

**Simple Summary:** Heat stress in bovine females from hot tropical regions is one of the main factors that negatively affect the success of ovum pick-up and *in vitro* fertilization programs (OPU-IVF). Seasonal climatological variation allows us to differentiate significant physiological effects in donors that affect the quantity and quality of embryos. The results of this study indicated a significant effect of season and breed on cumulus-oocyte complexes (COCs) and embryos, as well as the influence of weight and age, hormonal and physiological conditions of criollo Romosinuano (RM) and Tropical Milking (TM) donors. The greatest quantity and quality of COCs and code one blastocysts was observed in the hot humid and fresh dry seasons, so it is concluded that these seasons are the most suitable to perform OPU-IVF in RM and TM donors.

**Abstract:** One of the main factors that influence the fertility of cattle in grazing systems in hot tropical climates is heat stress. The objective of this study was to evaluate the effect of season, breed, hormonal and physiological condition on the quantity and quality of cumulus-oocyte complexes (COCs) and embryos produced *in vitro*, from Romosinuano (RM) and Tropical Milking (TM) donors. Three ovum pick-up and *in vitro* fertilization (OPU-IVF) were performed, one per season, hot dry (HD, 10, 10), hot humid (HH, 9, 9) and fresh dry (FD, 7, 10) in RM and LT donors. Serum levels of cortisol, insulin and glucose were measured, in addition to heart rate (HR), respiratory rate (RR) and rectal temperature (RT). Effect of season x genotype interaction ( $p \leq 0.05$ ) was observed in all COCs variables and only in cleavage embryos (CLI) ( $p \leq 0.05$ ). Body weight (BW) affected all COCs variables ( $p \leq 0.01$ ), except unviable (UNV) although affected degenerated embryos (DEG) ( $p \leq 0.01$ ) and total blastocysts (BLAST) ( $p \leq 0.01$ ). Cow age only affected viable COCs (VIAB) ( $p \leq 0.05$ ), code one blastocysts (BC1) and BLAST ( $p \leq 0.01$ ). Cortisol affected total COCs (COCsT), VIAB and total matured *in vitro* (TMIV) ( $p \leq 0.01$ ), as well as CLI, BC1 ( $p \leq 0.01$ ) and BLAST ( $p \leq 0.05$ ). Insulin affected COCsT ( $p \leq 0.01$ ), UNV ( $p \leq 0.05$ ), denuded oocytes (DE) ( $p \leq 0.01$ ), BC1 and code two blastocysts (BC2) ( $p \leq 0.01$ ). Glucose affected all COCs variables ( $p \leq 0.01$ ), except UNV and all embryo variables except BC2. HR affected COCsT, DE, TMIV ( $p \leq 0.01$ ), CLI, BLAST and DEG ( $p \leq 0.05$ ). RR affected COCsT, UNV, VIAB, CLI ( $p \leq 0.05$ ), BC1, BLAST and DEG ( $p \leq 0.01$ ). RT only affected DE, VIAB ( $p \leq 0.01$ ) and BLAST ( $p \leq 0.05$ ). The seasonal climatic year variation of Veracruz and changes in physiological and

hormonal variables have diverse effects on the cumulus-oocyte complexes and embryos produced by RM and TM donors.

**Keywords:** Adapted breeds; climatic change; genetic resources; hot climates; OPU-IVF; tropical region

---

## 1. Introduction

Ovum pick-up (OPU) and in vitro fertilization (IVF) are used in cattle to accelerate the diaspora of genotypes and genetically superior females [1], furthermore, they enable the multiplication of small populations adapted to adverse environments and even in danger of extinction [2]. OPU-IVF has been successful for *Bos taurus* breeds in the temperate climatic region and for *Bos indicus* breeds in the hot climatic tropical region [3,4]. However, for *Bos taurus* adapted breeds to the hot climatic tropical region studies are scarce and less successful [5,6], among other possible factors, the adverse hot humid climate affects directly and indirectly the reproductive performance of females [7]. The low fertility of cattle in hot climates is related to heat stress caused by high maximum temperatures (Tmax) and relative humidity (RH) [8]. Animals under heat stress increase the concentrations of cortisol, insulin and glucose in blood [9,10]. In addition, they respond physiologically by increasing heart rate, respiratory rate and rectal temperature, among other acclimatization mechanisms, with the aim of dissipating body heat [11,12], these changes negatively affect reproductive behavior [13].

Although OPU-IVF are reproductive technologies that contribute to rapidly increasing herds of pure animals, inter annual season variability with Tmax and RH higher than 30 °C and 84% [14,15] can decrease their success. In recent years, 90% of the annual pluvial precipitation has occurred in the hot humid season, while in the fresh dry and hot dry only around 5% in each, with a total of seven months of drought [16–18]. In the female bovine, the increase in body temperature reduces the follicular population and decreases the size of the preovulatory follicles, in addition, a higher degree Celsius in the temperature of the follicles in the ovaries inhibits ovulation [19]. In the oocyte, high temperatures cause damage to cellular organelles and DNA [20] and in the first stages of embryo development, cellular and epigenetic changes are generated that limit its development and quality [21].

The criollo breeds Romosinuano and Tropical Milking are *Bos taurus* naturalized to hot climates of the tropical region, with populations established in different countries of the Americas, although relatively small of thousands of individuals [22–24]. Studies have been carried out to determine the effect of Tmax and RH on the fertility of TM heifers and cows, it was found that the lowest pregnancy, less than 50%, occur in the hot dry season with Tmax greater than 27 °C and RH of 85% for heifers [18], but no season effect was observed in cows [25]. When comparing, Brahman and Angus breeds to determine the effect of heat shock on oocytes cultured in vitro at 41°C, oocytes had greater cleavage, with embryos of more than eight cells four days after IVF [26]. The objective of this study was to evaluate the effect of the season of the year, the breed, the hormonal and physiological condition of cows on the quantity and quality of the cumulus-oocyte complexes and embryos produced in vitro of Romosinuano and Tropical Milking females.

## 2. Materials and Methods

### 2.1. Characterization of the Seasons of the Year

The study was carried out in three climatic seasons (EST): hot dry (HD, March-May, with Tmax and RH of  $31.6 \pm 1.0$  °C and  $77.2 \pm 1.1$ %), hot humid (HH, June-October, with Tmax and RH of  $31.4 \pm 0.4$  °C and  $84.7 \pm 1.0$ %), and fresh dry (FD, November-February, with Tmax and RH of  $26.8 \pm 0.5$  °C and  $84.6 \pm 0.3$ %) [18] in Veracruz, Mexico, at 19° 11'N and 96° 20'W, at 23 m amsl, with average annual temperature and precipitation of 24.8 °C and 1,525.8 mm [15]. The climate of the region is AW0(w)(i)gw'', hot sub humid with rains in summer [14].

## 2.2. Management of Donors

The study was carried out in accordance with the Official Mexican Standard (NOM-062-ZOO-1999) of technical specifications for the production, care and use of laboratory animals [27], as well as the regulations for the use and care of research animals from the Colegio de Postgraduados [28]. Twenty-six Romosinuano (RM) and 29 Tropical Milking (TM) donors, nulliparous and up to three parturitions were used, selected by ultrasonography to rule out reproductive anomalies. The RM and TM cows had body weight (BW) of  $429.3 \pm 7.2$  and  $368.7 \pm 17.8$  kg, mean age of  $95.9 \pm 5.5$  and  $54.2 \pm 4.9$  months, and body condition score (BC) of  $2.6 \pm 0.1$  and  $2.9 \pm 0.1$  (scale 1 emaciated – 5 obese [29]), respectively. The BC in HD, HH and FD was  $2.6 \pm 0.3$ ,  $2.9 \pm 0.5$  and  $2.7 \pm 0.4$ . In each climatic season, an OPU-IVF session was performed, with 10, 9 and 7 RM and 10, 9 and 10 TM cows. The donors were rotationally grazed fed with para (*Brachiaria mutica* Forssk) and native grass (*Paspalum spp*), were dewormed with Ivermectin (Virbamec® Platinum, Virbac, Mexico) and received an application of phosphorus (Phospho® 20, Virbac, Mexico) monthly for four occasions, in addition to free mineral salt (Vimifos®, Mexico).

## 2.3. Follicular Aspiration and In Vitro Embryo Production

The most used reagents were from Sigma-Aldrich (St Louis, Mo, USA). For OPU, an ultrasound was used with a 7.5 MHz micro-convex transducer (Mindray®, DP-50Vet, Shenzhen, China) placed in an aspiration guide (WTA, Cravinhos, SP, Brazil). To aspirate the follicles, a 1 mm diameter x 120 cm plastic aspiration line and 20 G x 50 mm OPU needles were used. The follicular fluid was collected in a 150 mL conical tube with collection medium (Dulbecco's Phosphate buffer, DPBS), supplemented with 1% fetal bovine serum (FBS) and 125 IU mL<sup>-1</sup> of heparin at 37 °C.

The collected cumulus-oocyte complexes (COCs) were washed with DPBS and observed in a stereoscope (EMZ-8TR, Meiji Techno., Saitama, Japan). COCs were classified into four classes according to the number of cumulus cell layers and the characteristics of the ooplasm [30]. Class 1, more than three complete layers of cumulus cells and uniform granulation, the ooplasm fills the zona pellucid uniformly (ZP), class 2, less than three layers thick, ooplasm fills the ZP, class 3, shrunken and degenerated ooplasm, far from the ZP with partial filling, and class 4, naked oocytes, only enclosed by the ZP, in addition to presenting damage and remains of ooplasm.

Once classified, the COCs were grouped into viable (VIAB) ones, which included classes 1 and 2, unviable (UNV) class 3 and bare class 4. Only the VIAB COCs were selected for subsequent maturation. The COCs were transported to the laboratory in 1.5 mL cryotubes with in vitro maturation medium (IVM), TCM-199, L-glutamine and phenol red supplemented with 2.2 mg mL<sup>-1</sup> of sodium bicarbonate, 1% (v/v) of FBS with 2 µL of heparin, 0.01 µg mL<sup>-1</sup> of streptomycin, 0.01 µg mL<sup>-1</sup> of penicillin-G, in a lab mix portable incubator (WTA, Cravinhos, SP, Brazil). In the laboratory, 10 COCs were placed in 50 µL drops of IVM, covered with mineral oil and transferred to the fixed bench incubator (WTA, 12419, Cravinhos, SP, Brazil) to complete the 22 h period from IVM to 38 °C, 5% CO<sub>2</sub> and saturation humidity. Once maturation was completed, the COCs were transferred to 90 µL drops of in vitro fertilization medium (IVFM), supplemented with 20 µL mL<sup>-1</sup> of heparin and 6 mg mL<sup>-1</sup> of essential fatty acids, free of bovine serum albumin fatty acids and covered with mineral oil.

IVF was carried out in the three seasons of the year, with the use of two high fertility straws from RM and LT sires. The semen was prepared using a discontinuous Percoll gradient and  $1 \times 10^6$  sperm mL<sup>-1</sup> for each of the drops. After the 18 h IVF period, the presumptive zygotes were partially denuded with hyaluronidase, in a vortex for 5 min. In vitro culture medium (IVCM), KSOM, MR-107-D, supplemented with 10% FBS, 0.25 µg mL<sup>-1</sup> gentamicin and 0.5% non-essential amino acids was used for seven days after IVF, at which time the evaluation of the number and quality of the blastocysts was carried out. Only code 1 embryos were cryopreserved by vitrification.

#### 2.4. COCs Response Variables

Total COCs (COCsT) were observed in the field, with a stereoscope (EMZ-8TR, Meiji Techno., Saitama, Japan) in DPBS medium, when performing follicular aspiration, including VIAB, UNV, denuded (DE) and total matured in vitro (TMIV) with the cumulus cells expanded after the maturation period and the maintenance of cytoplasmic integrity.

#### 2.5. Embryo Response Variables

In the embryos, those cleavage (CLI) with at least two cells after fertilization were observed. Code one blastocysts (BC1) with symmetrical and spherical internal cell mass with individual blastomeres similar in size, color and density with at least 85% of the cellular material intact, code two blastocysts (BC2) with minor irregularities in the inner cell mass or in size, color and density of individual cells and at least 50% of their cell mass intact, total blastocysts (BLAST) all embryos at the blastocyst stage, including codes one and two, and degenerate blastocysts (DEG) with unviable cell mass [31].

#### 2.6. Hormonal Covariates and Glucose

Two hormones were measured in each donor, basal insulin and serum cortisol, as well as serum glucose concentration. A blood sample was collected by puncture of the coccygeal vein, one day before each OPU at 6:00 h, with 21G x 38 mm needles and Vacutainer tubes (Vacutainer®, Becton Dickinson, Franklin Lakes, NJ, USA). Blood samples were collected and centrifuged for 15 min at 750 x g at 4 °C. Serum samples were frozen and kept at -20 °C until further analysis. Basal insulin and serum cortisol concentrations were determined by chemiluminescence immunoassay (CLIA) (Maglumi 800, Shenzhen New Industries Biomedical Engineering Co., Ltd. (Snibe), Shenzhen, China) and serum glucose concentration was determined by automated spectrophotometry with a clinical chemistry analyzer (Architect c4000, Abbott Diagnostics, Abbott Park, IL, USA).

#### 2.7. Physiological Covariates

The heart rate (HR) was measured in each donor, one day before the OPU at 15:00 h (hottest of the day), with a double bell stethoscope (BLESMED®, CDMX, Mexico), for one minute, the respiratory rate (RR), by observing the thoracic abdominal expansion on the left flank, for one minute and the rectal temperature (RT) with a veterinary digital thermometer (FlashCheck®, Delta Trak Inc., International, Mexico).

#### 2.8. Statistical Analysis

Linear fixed effects statistical models were used. The models included the factors of EST, breed and their interaction, body weight (BW) and age onset covariates were also included, the hormonal and glucose ones and the physiological ones of the hottest time. A  $P(\lambda)$  distribution of the experimental error was considered. Data were processed using generalized linear models and the GLM and GENMOD procedures of SAS [32].

### 3. Results

#### 3.1. COCs

The EST x breed interaction affected all COCs response variables ( $p \leq 0.001$ , Table 1). The highest number of COCsT was observed in the HD and FD seasons and RM breed, higher for more than 10 in the lowest FD season and LT breed, with similar numbers in the other season and 9% more for RM. The highest number of VIABs occurred in FD and RM with more than 18 the lowest in FD and TM, VIAB in both breeds were between 12 and 16 ( $p \leq 0.001$ ) and between seasons 12 to 15 ( $p > 0.05$ ). The number of UNV was close to zero in all seasons and breeds, except in the HH and TM, which was more than 8 times higher than in HD and TM. The lowest number of DE occurred in the TM, close to four, although in the HD, with four times less than in FD, which had more DE than the HD

and HH. The highest number of TMIV with more than 27 occurred in HD and RM and with less than 14 for TM in the same season, values close to 20 occurred in all other EST x breed combinations. RM had 8.5 more TMIV than TM.

**Table 1.** Cumulus-oocyte complexes (COCs) collected by ovum pick-up in Romosinuano (RM) and Tropical Milking (TM) cows in three seasons in Veracruz, Mexico.

Variable Breed	Season			
	Hot dry	Hot humid	Fresh dry	Global
<i>Total COCs</i>				
RM	29.6 ± 2.6a	22.6 ± 1.7b	29.1 ± 2.3a	26.9 ± 1.3
TM	18.9 ± 2.0b	28.4 ± 2.2a	26.6 ± 2.1a	24.2 ± 1.3
Global	23.6 ± 1.5	25.4 ± 1.4	27.8 ± 1.6	
<i>Viable</i>				
RM	17.3 ± 2.1a	12.8 ± 1.3b	18.7 ± 1.9a	16.1 ± 1.0A
TM	13.7 ± 1.8ab	17.4 ± 1.8a	8.0 ± 1.0c	12.4 ± 0.9B
Global	15.4 ± 1.2	14.9 ± 1.1	12.2 ± 1.0	
<i>Unviable</i>				
RM	1.7 ± 0.6b	1.7 ± 0.5b	1.8 ± 0.5b	1.8 ± 0.3
TM	0.4 ± 0.3a	3.0 ± 0.8b	1.5 ± 0.5b	1.2 ± 0.3
Global	0.9 ± 0.3Y	2.3 ± 0.4XZ	1.7 ± 0.4YZ	
<i>Denuded</i>				
RM	9.4 ± 1.4b	7.5 ± 1.0b	8.3 ± 1.1b	8.4 ± 0.7
TM	4.2 ± 0.8a	6.0 ± 1.0ab	17.1 ± 2.2c	7.6 ± 0.7
Global	6.3 ± 0.7X	6.7 ± 0.7X	11.9 ± 1.1Y	
<i>Total matured in vitro</i>				
RM	27.8 ± 2.7a	24.6 ± 2.0a	22.9 ± 2.1a	25.0 ± 1.3A
TM	13.3 ± 1.5c	16.7 ± 1.5c	20.0 ± 1.8abc	16.5 ± 1.0B
Global	19.3 ± 1.3	20.3 ± 1.2	21.4 ± 1.4	

<sup>A,B</sup>. Different literal per row indicates statistical difference ( $p \leq 0.01$ ). <sup>X,Y,Z</sup>. Different literal per column indicates statistical difference ( $p \leq 0.006$ ). <sup>a,b,c</sup>. Different literal per column for season and row for breed indicates statistical difference ( $p \leq 0.01$ ).

A main effect of EST was observed on UNV ( $p \leq 0.05$ ) and DE ( $p \leq 0.01$ ), and of breed on VIAB ( $p \leq 0.05$ ) and TMIV ( $p \leq 0.01$ ). The lowest number of UNV was observed in HD, almost three times less than the highest number observed in HH. The number of DE was similar to six in HD and HH, but in FD more than eleven were observed. Regarding breed, the VIAB and TMIV were higher in RM, with four and eight more compared to TM.

### 3.2. Effect of Hormonal and Physiological Covariates on COCs

Covariates affected COCs. BW affected all variables ( $p \leq 0.01$ ) except UNV, the betas were positive and similar,  $\beta = 0.002 \pm 0.001$  for COCsT, VIAB and DE and  $\beta = 0.004 \pm 0.001$  for TMIV. Age only had effect on VIAB with  $\beta = 0.005 \pm 0.002$ . Regarding the hormonal and glucose covariates, cortisol affected COCsT, VIAB and TMIV ( $p \leq 0.01$ ), the betas were all negative with the lowest  $\hat{\beta} = -0.098 \pm 0.014$  for TMIV and the highest  $\hat{\beta} = -0.045 \pm 0.013$  for COCsT. Insulin only affected COCsT ( $p \leq 0.01$ ), UNV ( $p \leq 0.05$ ) and DE ( $p \leq 0.01$ ), the betas were all negative, the lowest  $\hat{\beta} = -0.251 \pm 0.109$  was observed in UNV and the highest  $\hat{\beta} = -0.159 \pm 0.032$  in COCsT. Glucose significantly affected all the response variables of the COCs ( $p \leq 0.01$ ), except UNV. Glucose betas were all positive.

Physiological covariates affected COCs. HR affected COCsT, DE and TMIV ( $p \leq 0.01$ ), the lowest  $\hat{\beta} = -0.020 \pm 0.004$ , for TMIV and the highest  $\hat{\beta} = -0.008 \pm 0.003$  for DE. Effect of RR was observed in COCsT, UNV and VIAB ( $p \leq 0.01$ ), the lowest  $\hat{\beta} = -0.070 \pm 0.021$  for UNV and the highest  $\hat{\beta} = -0.017 \pm 0.004$  for COCsT. RT only affected DE and VIAB ( $p \leq 0.01$ ), with beta negative for DE and positive for VIAB.

### 3.3. Embryos

The response in the embryonic variables due to the study factors is presented in Table 2. The EST x breed interaction only affected CLI ( $p \leq 0.05$ ), the greatest amount of CLI was observed in HD and RM, with more of nine than in the minor FD and TM, RM had twice as many CLI as TM in all seasons. Main effect of EST was observed in BLAST ( $p \leq 0.01$ ) and DEG ( $p \leq 0.05$ ), the BLAST means were similar in HD and HH with values above three and lower in FD, with less than half compared to HD and HH. Regarding DEG, the lowest mean was observed in FD, similar to HH and nine times lower than HD, the highest. Breed only affected CLI ( $p \leq 0.01$ ), more than eleven were observed in RM, double the five observed in TM.

**Table 2.** Embryos produced in vitro (IVF) from Romosinuano (RM) and Tropical Milking (TM) cows in three seasons in Veracruz, México.

Variable Breed	Season			
	Hot dry	Hot humid	Fresh dry	Global
<i>Cleavage</i>				
RM	10.8 ± 1.6 <sup>ab</sup>	13.3 ± 1.7 <sup>a</sup>	10.8 ± 1.5 <sup>ab</sup>	11.6 ± 1.0 <sup>A</sup>
TM	3.9 ± 0.7 <sup>c</sup>	5.4 ± 0.9 <sup>bc</sup>	7.8 ± 1.2 <sup>b</sup>	5.5 ± 0.6 <sup>B</sup>
Global	6.5 ± 0.7	8.5 ± 0.8	9.2 ± 0.9	
<i>Blastocysts code one</i>				
RM	1.5 ± 0.5	1.6 ± 0.5	1.1 ± 0.4	1.4 ± 0.3
TM	1.3 ± 0.6	1.5 ± 0.4	0.6 ± 0.2	1.1 ± 0.3
Global	1.4 ± 0.4	1.5 ± 0.3	0.9 ± 0.2	
<i>Blastocysts code two</i>				
RM	1.3 ± 0.7	0.7 ± 0.4	0.4 ± 0.2	0.7 ± 0.2
LT	0.5 ± 0.3	0.5 ± 0.3	0.3 ± 0.2	0.4 ± 0.2
Global	0.8 ± 0.3	0.6 ± 0.2	0.4 ± 0.2	
<i>Total blastocysts</i>				
RM	5.0 ± 1.2	3.4 ± 0.8	1.8 ± 0.5	3.1 ± 0.5
TM	2.6 ± 0.7	3.0 ± 0.6	1.1 ± 0.3	2.1 ± 0.4
Global	3.6 ± 0.6 <sup>x</sup>	3.2 ± 0.5 <sup>x</sup>	1.4 ± 0.3 <sup>y</sup>	
<i>Degenerate</i>				
RM	2.0 ± 0.9	0.3 ± 0.1	0.2 ± 0.1	0.5 ± 0.2
TM	0.6 ± 0.3	0.7 ± 0.3	0.2 ± 0.1	0.4 ± 0.2
Global	1.1 ± 0.3 <sup>y</sup>	0.4 ± 0.1 <sup>xy</sup>	0.2 ± 0.1 <sup>x</sup>	

<sup>A,B</sup>. Different literal per row indicates statistical difference ( $p \leq 0.01$ ). <sup>x,y,z</sup>. Different literal per column indicates statistical difference ( $p \leq 0.03$ ). <sup>a,b</sup>. Different literal per column for season and row for breed indicates statistical difference ( $p \leq 0.02$ ).

### 3.4. Effect of Hormonal and Physiological COVARIATES on embryos

The covariates affected the embryos. BW affected DEG and BLAST ( $p \leq 0.01$ ) and age BC1 and BLAST ( $p \leq 0.01$ ). The hormones affected the embryos. Cortisol affected CLI ( $p \leq 0.01$ ), BC1 ( $p \leq 0.01$ ) and BLAST ( $p \leq 0.05$ ), negatively with the lowest  $\hat{\beta} = -0.124 \pm 0.053$  for BC1 and the highest  $\hat{\beta} = -0.081 \pm 0.035$  for BLAST. Insulin affected BC1 and BC2 ( $p \leq 0.01$ ), with  $\hat{\beta} = -0.381 \pm 0.132$  for BC1 and  $\hat{\beta} = -0.543 \pm 0.232$  for BC2. Glucose affected all embryo variables except BC2, with the lowest  $\hat{\beta} = 0.201 \pm 0.037$  for BC1 and the highest  $\hat{\beta} = 0.057 \pm 0.014$  for CLI. Physiological covariates also affected embryos. The HR affected CLI, BLAST and DEG ( $p \leq 0.05$ ), the lowest and positive  $\hat{\beta} = 0.029 \pm 0.014$  was observed in DEG and the highest and negative  $\hat{\beta} = -0.010 \pm 0.005$  in CLI. The RR affected CLI ( $p \leq 0.05$ ), BC1, BLAST and DEG ( $p \leq 0.01$ ), the betas were all negative except for CLI, with the lowest  $\hat{\beta} = 0.154 \pm 0.038$  for DEG and the highest  $\hat{\beta} = 0.015 \pm 0.007$  for CLI. RT only affected BLAST ( $p \leq 0.05$ ) with  $\hat{\beta} = -0.885 \pm 0.427$ .

#### 4. Discussion

It has been observed that, as the rectal temperature of cows subjected to heat stress increases, the uterine temperature increases, which affects the number of small and medium-sized follicles [33]. In this study the lowest amount of COCsT, less than 20, was observed in the TM breed in the HD season, when the least precipitation occurs and the highest maximum temperatures are observed, however, the RM breed observed higher and similar numbers of COCsT. At the FD season, medium numbers of COCsT were observed with no difference between breeds ( $p \leq 0.05$ , Table 1). In Brahman and Simmental  $\times$  Brahman crossbreed heifers, a greater amount of COCsT and medium follicles ( $4.0 \pm 0.2$ ) useful in IVF were obtained in the fresh humid season and a lower in the hot dry ( $2.9 \pm 0.2$ ) [34].

In Black Japanese cows in the hot ( $28.3 \pm 0.3$  °C, 66.5% RH) and cool ( $17.5$  °C, 62.9% RH) seasons,  $11.2 \pm 0.8$  and  $23.2 \pm 1.9$  COCsT were obtained by OPU [35] and in Gyr cows for 14 days in controlled environments at 25 and 38 °C,  $11.2 \pm 2.8$  and  $14.3 \pm 2.5$  COCsT were obtained [36]. In addition to climatological and environmental factors, genotype, aspiration frequency and semen used, among others, affect COCsT in an OPU-IVF program, [37]. In this study, for each kg of increase in BW of the donors, the COCsT increased with  $\hat{\beta} = 0.002 \pm 0.001$ , well fed donors have greater energy reserves that are used in reproductive events [38]. On the other hand, as cortisol and insulin increased, COCsT decreased with  $\hat{\beta} = -0.045 \pm 0.013$  and  $\hat{\beta} = -0.159 \pm 0.032$ . Heat stress in females activates the hypothalamic-pituitary-adrenal axis, which excites the pituitary gland to release the adrenocorticotrophic hormone, which stimulates the release of glucocorticoids, such as cortisol, which when increased causes an inhibitory effect on the reproductive axis, which causes a delay in follicular development [39]. Furthermore, high concentrations of cortisol cause a decrease in glucose entry into cells, which result in high amounts of glucose and insulin in the blood [40]. Similarly, as the HR and RR increased, the COCsT decreased, increases in physiological covariates are presented as a mechanism generated by animals to release heat [41] and their higher frequencies indicate greater stress in the animal, which negatively affects its reproductive performance.

It is important to incubate only viable COCs, since correct IVM and subsequent IVF depend on it, as well as their development to the blastocyst stage [42,43]. The highest VIAB were observed in FD for RM and HH for TM (Table 1). The lowest temperatures in the region occur in FD and although in HH the Tmax and RH are high, also the greatest precipitation occurs, which allow the females to reduce body heat, with positive impact on the recovered VIAB. Increasing Tmax in HD directly affect follicular development and oocyte competence [44]. Our results differ from those by Torres-Júnior et al. [36] in Gyr cows, who observed no effect on VIAB (27.7 and 31.5%) from 25 °C and 38 °C environments. In this study, BW and age positively affected the COCsT, which may be related to the amount of VIAB, the highest COCsT occurred in FD and HH, the highest precipitation also occurs in HH, so the availability of forage is greater in this season, donors with excellent body condition and BW correctly generate hormonal processes, allowing the development of follicles and oocytes of excellent quality [45]. On the other hand, although in other breeds the donor's age affects the VIAB, the donors in this study had an age of  $95.9 \pm 5.5$  in RM and  $54.2 \pm 4.9$  months in LT, criollo breeds are characterized by their longevity, with cows that can remain productive for more than 15 years [23].

The greatest number of UNV was observed in HD and FD in TM, however, in these seasons the greatest amount of COCsT was recovered, which may be related to the increase in UNV observed in TM (Table 1). In cows under heat stress, a lower quality of the COCs for IVF is observed. It has been found that oocytes recovered in hot seasons have a lower capacity to reach the blastocyst stage [46]. In cows under heat stress, the components of the cytoskeleton and mitochondrial functions and the correct expansion in cumulus cells are altered during maturation, which induces cell apoptosis [47] and requires two to three estrous cycles to recover good quality COCs damaged in the HD season [48]. In vitro matured studies have been carried out at 41 °C, to evaluate the damage caused by thermal shock in COCs and it has been observed that the damage in the ZP, with a greater number of pores, as well as abnormalities increase after 22 h of exposure to high temperatures [49]. The cows in this study were kept exposed to grazing weather conditions throughout the year and the highest Tmax occurred in HD [18]. It has been observed that as heat stress increases, animals activate

physiological mechanisms to release body heat [50]. In this study, as RR increased, UNV decreased with  $\hat{\beta} = -0.070 \pm 0.021$ , as product of the heat release that occurred in the donors as respiration increased.

Factors that are related to the increase in DE oocytes include the frequency and speed in the OPU and the gauge of the needle, among others [51,52]. However, there are no previous studies showing the effect of season on the number of DE oocytes. In this study, a greater amount of DE was observed in HH and FD in TM and, although without statistical difference, greater amounts of DE in HD and HH in RM (Table 1). In Gyr cows, 15.8 and 23.0% DE oocytes were observed at low and high temperatures.

Hormonal covariates, insulin and glucose affected DE (Table 1), at greater increases in glucose, a greater amount of DE was observed. It is important to put at IVM those COCs with the greatest number of cumulus cells, since they allow their communication with the oocyte, influencing protein synthesis and gene expression for the subsequent expansion and maturation of the oocyte [53].

The highest TMIV means were observed in HD and HH for RM, but also the lowest in the same seasons for TM (Table 1). In zebu and crossbred cows, oocytes had nuclear maturation of 76.9, 52.2 and 39.1% in the HH, HD and FD seasons [54]. In COCs subjected to heat shock of 41 °C in the first 12 h of maturation *in vitro*, oxidative stress and apoptosis were increased, the distribution of cortical granules was altered, the formation and potential of the mitochondrial membrane was reduced, and damage was presented. In the cytoskeleton, reducing the cortical organization of actin [55] and the expansion of cumulus cells was reduced by 20% [47]. Andrew-Vázquez et al. [56] observed 14.5 times more abnormalities in the morphology of metaphase II oocytes subjected to heat shock of 41.5 °C in the first 3 h of *in vitro* maturation. As BW increased in the donors in this study, TMIV increased, which is related to the increase in VIAB as the females gained weight. It is advisable to maintain excellent nutrition in donors when undergoing reproductive biotechnologies, good nutrition generates better results [57]. In the donors of this study, as cortisol increased, TMIV decreased, high levels of cortisol occur as a response to heat stress [50], which in addition, caused an increase in HR and as a result a lower amount of COCsT and VIAB. Therefore, it is advisable to expose only VIAB to *in vitro* maturation, as they have the highest probability of completing the process [43].

Among the factors that affect the division of the presumed zygotes are the quality of the semen used [58], the culture medium [59] and the quality of the oocyte collected, which can come from donors under heat stress [60]. In this study, the highest means of CLI embryos were observed in the HH for RM and FD for TM and the lowest in HD also for TM with almost 70% less in relation to the highest (Table 2). The highest temperatures occur in HD, it has been observed that more than six hours of exposure to extreme temperatures generate heat stress in females that decreases the cleavage of embryos, with a smaller number of embryos reaching at least eight cells [61] and a low percentage of embryos in the blastocyst stage [55].

Jersey embryos matured *in vitro* at 38.5 and 41 °C observed 11.4% less CLI at the highest temperature [62]. In Gyr cows, 74.6 and 80% of CLI were obtained at low and high temperatures [36]. Cortisol and glucose affected CLI in this study. As cortisol increased, CLI decreased. Large concentrations of cortisol indicate stress in the donors, which is reflected in lower quality oocytes that, although they complete the IVM, do not have the ability to cleavage. On the other hand, HR and RR affected CLI, the increases in heart and respiratory rates are used by animals as a heat release mechanism [41], so by increasing the flow of oxygenated blood, heat decreases, that in turn benefits the quality of the COCs and that affects the increase in CLI, as occurred in this study.

Efforts *in vitro* embryo production focus on increasing BC1, since they have the greatest capacity to generate a pregnancy after being transferred [63] or survive cryopreservation [64]. In this study, the highest mean BC1 was observed in HH for both breeds, although without statistical differences (Table 2). The greatest precipitation occurs in the HH season, with greater availability and quality of forage [65]. However, the means of  $1.4 \pm 0.3$  and  $1.1 \pm 0.3$  for RM and TM were not different between breeds adapted to warm environments [26]. These *Bos taurus* breeds in adverse environments over the centuries developed adaptation characteristics such as slick hair coat [66] and mechanisms of metabolic heat reduction and increases in its dissipation through sweat glands and respiratory rate

[67]. The age of the donor influenced the BC1, the females had a maximum of three births, so as the age increased better development has been observed in the OPU that allows recovering COCs of better quality, without damage to the ZP and the greater number of cumulus cells, which allow the increase of BC1. The observed effect of the hormonal covariates and glucose is related to the increase in the heat stress of the donors [68]. As the hormones increased, the recovered COCs presented greater damage that was reflected in a decrease in BC1.

No effect of season or breed was observed on BC2 (Table 2), however, the means were very low, less than  $0.8 \pm 0.2$ . It is preferable to obtain BC1 and a smaller amount of BC2, since pregnancies are influenced by the quality of the blastocyst [63]. The increases in insulin in the donors decreased BC2, the high concentrations in the blood are related to the stress of the donors, since by reducing the entry of glucose into the cells, insulin concentrations increase as a mechanism protection and adaptation [69].

The quality of the oocyte is an important indicator of its development [42], however, in hot climate tropical regions, exposure of females to more than 12 h of heat stress reduces the ability of embryos to reach the blastocyst stage [61]. The highest mean BLAST was observed in HD ( $3.6 \pm 0.6, 45.5\%$ ) greater than HH ( $3.2 \pm 0.5, 40.4\%$ ) and FD ( $1.4 \pm 0.3, 24.9\%$ ). Criollo breeds are adapted to seasonal changes [23], since stressful conditions over decades generate genetic and metabolic adaptations that contribute to their physiological performance [70]. In Gyr cows [36] averages of  $3.0 \pm 0.5$  and  $1.7 \pm 0.5$  embryos were observed per donor at low and high temperatures. In RM cows [26], 30% and 18% BLAST were obtained in oocytes subjected to heat shock of 41 and 38.5 °C during IVM. On the other hand, Al-Katanani et al. [44] observed in BLAST derived from ovaries of Holstein cows, collected in winter and summer, 34.3 and 11.4% respectively, with a decrease of 22.9%. The global BLAST means obtained by season (Table 2) are similar to the three and four BLAST obtained by OPU-IVF in heifers and criollo cows from Ecuador [71,72].

Hormonal covariates, with the exception of insulin and physiological covariates, affected BLAST, these effects were also observed in VIAB. It is known that their quality has an impact on the development of embryos [43]. Small populations need the adoption of new technologies that facilitate the increase of pure and genetically superior animals [73]. In Colombia, a germplasm bank of criollo cattle has been created with embryos obtained in vivo and in vitro [74], this activity can be of great support for the establishment of new pure cattle herds and the exchange between countries of criollo bovine germplasm of high genetic quality [75].

The greatest amount of DEG was observed in HD ( $1.1 \pm 0.3$ ), compared to HH ( $0.4 \pm 0.1$ ) and FD ( $0.2 \pm 0.1$ ), with no difference between breeds (Table 2). During the Veracruz HD season, there is less precipitation and greater solar radiation, with Tmax of  $31.6 \pm 1.0$  °C. High temperatures have been observed to induce DNA fragmentation and affect RNA levels [20,76]. Therefore, if COCs are exposed to high temperatures in hot seasons, the number of cells is reduced in the few embryos that reach the blastocyst stage, in addition to greater damage to the blastomeres [55] than increases the number of DEG embryos. The increase in BW of the donors in this study decreased the DEG, well-fed females allocate energy to reproductive events with better quality COCs. On the other hand, as glucose and HR increased, DEG increased, which is an indication of stress in donors [69].

## 5. Conclusions

The exposure of criollo cows to the seasonal climatic variations of Veracruz and the physiological and hormonal changes had adverse effects on the cumulus-oocyte complexes and embryos. Although no effect of season and breed was observed in code one blastocysts, the number of degenerated embryos was greater in the hot dry season, so it can be concluded that the hot humid and cool dry seasons are the most favorable to perform OPU-IVF in Romosinuano and Tropical Milking cows.

**Author Contributions:** Conceptualization, F.R.M. and C.M.B.P., methodology, F.R.M. and C.M.B.P., formal analysis, F.R.M. and C.M.B.P., investigation, C.M.B.P., resources, A.R.P., C.M.B.P. and A.R.G., data curation, F.R.M. and C.M.B.P., writing—original draft preparation, F.R.M. and C.M.B.P., writing—review and editing, A.R.P., C.C.R., J.G.S., and S.R.G., visualization, A.R.P., C.C.R., J.G.S., and S.R.G., supervision, C.M.B.P., project

administration, A.R.P. and C.M.B.P., funding acquisition, A.R.P., C.M.B.P. and A.R.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** The Colegio de Postgraduados gave financial support for this research. In the Reproducción Genética Avanzada RGA In vitro company, the OPU-IVF processes were carried out. Carlos Miguel Becerril-Pérez and Adalberto Rosendo-Ponce provided the Romosinuano cows for the study.

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Institutional Review Board of Colegio de Postgraduados (Reglamento para el uso y cuidado de animales destinados a la investigación en el Colegio de Postgraduados).

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We thank CONACyT, Mexico for the doctoral scholarship awarded to FRM.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Ferré, B.L., Alvarez-Gallardo, H., Romo, S., Fresno, C., Stroud, T., Stroud, B., Lindsey, B., Kjelland, M. Transvaginal ultrasound-guided oocyte retrieval in cattle: State-of-the-art and its impact on the in vitro fertilization embryo production outcome. *Reprod. Domest. Anim* **2022**, 00, 1-16. Doi: 10.1111/rda.1403.
2. Herrick, J.R. Assisted reproductive technologies for endangered species conservation: developing sophisticated protocols with limited access to animals with unique reproductive mechanisms. *Biol. Reprod* **2019**, 1, 1158-1170. Doi: 10.1093/biolre/ioz025.
3. Sartori, R., Prata, A.B., Figueredo, A.C.S., Sanches, B.V., Pontes, G.C.S., Viana, J.H.M., Pontes, J.H., Vasconcelos, J.L.M., Pereira, M.H.C., Dode, M.A.N., Monteiro Jr. P.L.J., Baruselli, P.S. Update and overview on assisted reproductive technologies (ARTs) in Brazil. *Anim. Reprod* **2016**, 13, 300-312. Doi: 10.21451/1984-3143-AR873.
4. Viana, J.H.M., Siqueira, L.G.B., Palhao, M.P., Camargo, S.A., Use of in vitro fertilization technique in the last decade and its effect on Brazilian embryo industry and animal production. *Acta Sci. Vet* **2010**, 38(Supp 2), s661-s674. Available on line: [https://www.ufrgs.br/actavet/38-suple-2/23\\_SBTE\\_JHENRIQUE.pdf](https://www.ufrgs.br/actavet/38-suple-2/23_SBTE_JHENRIQUE.pdf) (accessed on 28 June 2022).
5. Goicochea, V.J., Rondón, J.W., Acosta, P.F., Gómez, M.Y., Montalvo, M.M., Salvatierra, A.M., Martel, F.J., Ballarte, Z.O., Díaz, Z.J., Ratto, F.M. Efecto de dos medios de fertilización en el desarrollo in vitro de embriones bovinos criollos. *Rev. Inv. Vet. Perú* **2021**, 32, e18709. Doi: 10.15381/rivep.v32i5.18709.
6. Rocha, A., Randel, D., Broussard, J.R., Lim, J.M., Blair, R.M., Roussel, J.D., Godke, R.A., Hansel, W. High environmental temperature and humidity decrease oocyte quality in *Bos taurus* but not in *Bos indicus* cows. *Theriogenology* **1997**, 49, 657-665. Doi: 10.1016/s0093-691x(98)00016-8.
7. Breuel, K.F., Lewis, P.E., Schrick, F.N., Lishman, A.W., Inskip, E.K., Butcher, R.L. Factors affecting fertility in the postpartum cow: role of the oocyte and follicle in conception rate. *Biol. Reprod* **1993**, 48, 655-661. Doi: 10.1095/biolreprod48.3.655.
8. Boni, R. Heat stress, a serious threat to reproductive function in animals and humans. *Mol. Reprod. Dev* **2019**, 86, 1307-1323. Doi: 10.1002/mrd.23123.
9. Gonçalves, T.C., Negrão, J.A., de Souza, C.T., Martins, T.R., da Cunha, L.S.T.M., Luis, H.F., Ferrari, C.R., Franco, P.A.M., Lencioni, T.E.A. Heat stress and ACTH administration on cortisol and insulin-like growth factor I (IGF-I) levels in lactating Holstein cows. *J. App. Anim. Res* **2017**, 45, 1-7. Doi: 10.1080/09712119.2015.1091326.
10. Itoh, F., Obara, Y., Rose, M.T., Fuse, H., Hashimoto, H. Insulin and glucagon secretion in lactating cows during heat exposure. *J. Anim. Sci* **1998**, 76, 2182-2189. Doi: 10.2527/1998.7682182x.
11. Rashamol, V.P., Sejian, V., Bagath, M., Krishnan, G., Archana, P.R., Bhatta, R. Physiological adaptability of livestock to heat stress: an updated review. *J. Anim. Behav. Biometeorol* **2018**, 6, 62-71. Doi: 10.31893/2318-1265jabb.v6n3p62-71.
12. Cartwright, S., Schmied, J., Livernois, A., Mallard, B.A. Physiological response to heat stress in immune phenotyped Canadian Holstein dairy cattle in free-stall and tie-stall management systems. *Front. Anim. Sci* **2022**, 3, 852958. Doi: 10.3389/fanim.2022.852958.
13. Fernandez-Novio, A., Pérez-Garnelo, S.S., Villagrà, A., Pérez-Villalobos, N., Astiz, S. The effect of stress on reproduction and reproductive technologies in beef cattle- a review. *Animals* **2020**, 10, 2096. Doi: 10.3390/ani10112096.
14. García, E. *Modificaciones al Sistema de Clasificación Climática de Köppen*, 5ta ed., Universidad Nacional Autónoma de México, Instituto de Geografía: Ciudad de México, México, **2004**. Available online: <http://www.publicaciones.igg.unam.mx/index.php/ig/catalog/view/83/82/251-1> (accessed on 23 April 2020).

15. SMN. *Sistema de información climatológica y geográfica del estado de Veracruz, México*. Consulta de base de datos 2000-2013. **2020**. Available online: <https://smn.conagua.gob.mx/es/>. (Accessed on 20 April 2020). (In Spanish).
16. Gutiérrez-García, G., Ricker, M. Climate and climate change in the region of Los Tuxtlas (Veracruz, México): A statistical analysis. *Atmósfera* **2011**, *24*, 347-373. Available on line: [https://www.scielo.org.mx/scielo.php?script=sci\\_arttext&pid=S0187-62362011000400001](https://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0187-62362011000400001).
17. Ortega-Gaucin, D., Cruz, B.J., Castellano, B.H.V. Drought vulnerability indices in Mexico. *Water* **2018**, *10*, 1671. Doi:10.3390/w10111671.
18. Rosales-Martínez, F., Rosendo-Ponce, A., Cortez-Romero, C., Gallegos-Sánchez, J., Cuca-García, J.M., Becerril-Pérez, C.M. Relation of the maximum temperature and relative humidity close to the insemination with the tropical milking criollo heifer's gestation in three seasons. *Trop. Anim. Health Prod* **2020**, *23*, 53. 27. Doi: 10.1007/s11250-020-02430-3.
19. López-Gatius, F., Hunter, R.H.F. Pre-ovulatory follicular cooling correlates positively with the potential for pregnancy in dairy cows: Implications for human IVF. *J. Gynecol. Obstet. Human. Reprod* **2019**, *48*, 419-422. Doi: 10.1016/j.jogoh.2019.03.005.
20. Sakatani, M. Effects of heat stress on bovine preimplantation embryos produced *in vitro*. *J. Reprod. Dev* **2017**, *63*, 347-352. Doi: 10.1262/jrd.2017-045.
21. Barros, R.O.F., Paula-Lopez, F.F. Cellular and epigenetic changes induced by heat stress in bovine preimplantation embryos. *Mol. Reprod. Dev* **2018**, *85*, 810-820. Doi.org/10.1002/mrd.23040.
22. AMCROLET (Asociación Mexicana de Criadores de Ganado Romosinuano y Lechero Tropical, A.C). Libro de registro de raza Criollo Lechero Tropical **2015**, Veracruz, México.
23. de Alba, J. *El libro de los Bovinos Criollos de América*. Ed. Colegio de Postgraduados. Biblioteca Básica de Agricultura, México **2011**.
24. Rosendo-Ponce, A., Becerril-Pérez, C.M. Avance en el conocimiento del bovino criollo Lechero Tropical de México. *Ecosistemas Rec. Agrop* **2015**, *2*, 233-243, Available on line: [https://www.scielo.org.mx/scielo.php?script=sci\\_arttext&pid=S2007-90282015000200012](https://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S2007-90282015000200012).
25. Rosales-Martínez, F., Becerril-Pérez, C.M., Rosendo-Ponce, A., Cortez-Romero, C., Torres-Hernández, G., Gallegos-Sánchez, J. Effects of season, maximum temperature and relative humidity on the gestation success of tropical milking criollo cows. *Agrociencia* **2023**, *57*, 5, 860-881. Doi: 10.47163/agrociencia.v57i5.2802.
26. Hernández-Cerón, J., Chase Jr, C.C., Hansen, J.P. Differences in heat tolerance between preimplantation embryos from Brahman, Romosinuano, and Angus breeds. *J. Dairy Sci* **2004**, *87*, 53-58. Doi: 10.3168/jds.S0022-0302(04)73141-0.
27. SAGARPA. Norma Oficial Mexicana, Especificaciones Técnicas para la Producción, Cuidado y uso de los Animales de Laboratorio. Diario Oficial de la Federación **2001**. Available online in pdf: [https://www.gob.mx/cms/uploads/attachment/file/203498/NOM-062-ZOO-1999\\_220801.pdf](https://www.gob.mx/cms/uploads/attachment/file/203498/NOM-062-ZOO-1999_220801.pdf).
28. Colegio de Postgraduados (COLPOS). Reglamento Para el Uso y Cuidado de Animales Destinados a la Investigación en el Colegio de Postgraduados. **2016**. Available online: [www.colpos.mx](http://www.colpos.mx) (accessed on 18 May 2021).
29. Wildman, E.E., Jones, G.M., Wagner, P.E., Boman, R.L., Trout, J.R., Lesch, T.N. A dairy cow body condition scoring system and its relationship to selected production characteristics. *J. Dairy Sci* **1982**, *65*, 495-512. Doi: 10.3168/jds.S0022-0302(82)82223-6.
30. Leibfried, L., First, N.L., Characterization of bovine follicular oocytes and their ability to mature *in vitro*. *J. Anim. Sci* **1979**, *48*, 76-86. Doi: 10.2527/jas1979.48176x.
31. Bó, A.G., Mapletoft, J.R. Evaluation and classification of bovine embryos. *Anim. Reprod* **2013**, *10*, 344-348, Available on line: <https://www.animal-reproduction.org/article/5b5a604cf7783717068b46a2/pdf/animreprod-10-3-344.pdf>.
32. Statistical Analysis System (SAS). *User's Guide: Statistic*, Version 9.4 for Windows. SAS Inst. Inc. Cary, NC, USA, **2010**.
33. Wolfenson, D., Thatcher, W.W., Badinga, L., Savio, J.D., Meidan, R., Lew, B.J., Braw-Tal, R., Berman, A. Effect of heat stress on follicular development during the estrous cycle in lactating dairy cattle. *Biol. Reprod* **1995**, *52*, 1106-1113. Doi: 10.1095/biolreprod52.5.1106.
34. Peralta-Torres, J.A., Aké-López, J.R., Centurión-Castro, F.G., Segura-Correa, J.C. Effect of season and breed group on the follicular population and cyclicity of heifers under tropical conditions. *Trop. Anim. Health Prod* **2017a**, *49*, 207-2011. Doi: 10.1007/s11250-016-1182-7.
35. Takuma, T., Sakai, S., Ezoe, D., Ichimaru, H., Jinnouchi, T., Kaedei, Y., Nagai, T., Otoi, T. Effect of season and reproductive phase on the quality, quantity and developmental competence of oocytes aspirated from Japanese Black cows. *J. Reprod. Dev* **2010**, *56*, 55-59. Doi: 10.1262/jrd.09-071h.
36. Torres-Júnior, S.J.R., Pires, F.A.M., Sá, F.W., Ferreira, M.A., Viana, M.J.H., Camargo, A.L.S., Ramos, A.A., Folhadella, M.I., Polisseni, J., Freitas, C., Clemente, A.C.A., Sá Filho, M.F., Paula-Lopes, F.F., Baruselli,

- S.P. Effect of maternal heat-stress on follicular growth competence in *Bos indicus* cattle. *Theriogenology*, **2008**, 69, 155-166. Doi: 10.1016/j.theriogenology.2007.06.023.
37. Nogueira, B.G.R.,Souza, L.F.A.de.,Puelker, R.Z.,Giometti, I.C.,Firretti, S.M.G.,Dias, T.S. dos S.B.,Castilho, C. Factors affecting the in vitro production of embryos in a commercial program. *Res. Soc. Dev* **2021**, 10, 2, e16110212264. Doi: 10.33448/rsd-v10i2.12264.
  38. Ponsart, C.,Gamarra, G.,Lacaze, S.,Ponter, A.A. Nutritional status of donor cows: insulin related strategies to enhance embryo development. *Anim. Reprod* **2014**, 11, 3, 195-198, Available on line: <https://www.animal-reproduction.org/article/5b5a603ef7783717068b4655/pdf/animreprod-11-3-195.pdf>.
  39. Krishnan, G.,Bagath, M.,Pragna, P.,Kusha, V.M.,Aleena, J.,Ravindranathan, A. P.,Sejian, V.,and Bhatta, R. *Mitigation of the heat stress impact in livestock reproduction*. In: Carreira, P.R, editor. *Theriogenology*, 1. Portugal: InTech,**2017**.
  40. Romero, P.M.H.,Uribe-Velásquez, L.F.,Sánchez, V.J.A. Biomarcadores de estrés como indicadores de bienestar animal en ganado de carne. *Biosalud*, **2011**, 10, 71-87, Available on line: [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S1657-95502011000100007](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S1657-95502011000100007).
  41. Idris, M.,Uddin, J.,Sullivan, M.,McNeill, M.D.,Phillips, J.C.C. Non-invasive physiological indicators of heat stress in cattle. *Animals* **2021**, 11, 71. Doi: 10.3390/ani11010071.
  42. Krisher, R.L. The effect of oocyte quality on development. *J. Anim. Sci* **2004**, 82(E. Suppl.), E14-E23. Doi: 10.2527/2004.8213\_supplE14x.
  43. Aguila, L.,Treulen, F.,Therrien, J.,Felmer, R.,Valdivia, M.,Smith, C.L. Oocyte selection for in vitro embryo production in bovine species: noninvasive approaches for new challenges of oocyte competence. *Animals* **2020**, 10, 2196. Doi: 10.3390/ani10122196.
  44. Al-Katanani, Y.M.,Paula-Lopes, F.F.,Hansen, J.P. Effect of season and exposure to heat stress on oocyte competence in Holstein cows. *J. Dairy Sci* **2002**, 85, 390-396. Doi: 10.3168/jds.s0022-0302(02)74086-1.
  45. Nazhat, S.A.,Aziz, A.,Zabuli, J.,Rahmati, S. Importance of body condition scoring in reproductive performance of dairy cows: A review. *Open J. Vet. Med* **2021**, 11, 272-288. Doi: 10.4236/ojvm.2021.117018.
  46. Ferreira, M.R.,Ayres, H.,Chiaratti, R.M.,Ferraz, L.M.,Araujo, B.A.,Rodrigues, A.C.,Watanabe, F.Y.,Vireque, A.A.,Joaquim, C.D.,Smith, C.L.,Meirelles, V.F.,Baruselli, S.P. The low fertility of repeated-breeder cows during summer heat stress is related to a low oocyte competence to develop into blastocysts. *J. Dairy Sci* **2011**, 94, 2383-2392. Doi: 10.3168/jds.2010-3904.
  47. Ahmed, J.A.,Nashiruddullah, N.,Dutta, D.,Biswas, R.K.,Borah, P. Cumulus cell expansion and ultrastructural changes in matured bovine oocytes under heat stress. *Iran J. Vet. Res*, **2017**, 18, 203-207. Available on: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5674444/pdf/ijvr-18-203.pdf>.
  48. Gendelman, M.,Roth, Z.,Incorporation of coenzyme Q10 into bovine oocytes improves mitochondrial features and alleviates effects of summer thermal stress on developmental competence. *Biol. Reprod* **2012**, 87, 118. Doi: 10.1095/biolreprod.112.101881.
  49. Báez, F.,Camargo, A.,Reyes, A.L.,Márquez, A.,Paula-Lopes, F.,Viñoles, C. Time-dependent effects of heat shock on the zona pellucida ultrastructure and in vitro developmental competence of bovine oocytes. *Reprod. Biol* **2019**, 19, 195-203. Doi: 10.1016/j.repbio.2019.06.002.
  50. Kim, W.S.,Lee, J.S.,Jeon, S.W.,Peng, D.Q.,Kim, Y.S.,Bae, M.H.,Jo, Y.H.,Lee, H.G. Correlation between blood, physiological and behavioral parameters in beef calves under heat stress. *Asian-Australas J. Anim. Sci* **2018**, 31, 6, 919-925. Doi: 10.5713/ajas.17.0545.
  51. Harkal, S.B.,Pawshé, C.H.,Ingawale, M.V.,Deshmukh, S.G.,Ambalkar, M.B.,Kale, S.D. Effect of frequency of follicular aspiration on recovery of oocytes and follicular development. *J. Entomol. Zool. Stud* **2019**, 7, 568-571. Doi: 10.1016/s0093-691x(98)00162-9.
  52. Ward, F.A.,Lonergan, P.,Enright, B.P.,Boland, M.P. Factors affecting recovery and quality of oocytes for bovine embryo production in vitro using ovum pick-up technology. *Theriogenology*, **2000**, 54, 433-446. Doi: 10.1016/s0093-691x(00)00360-5.
  53. Turathum, B.,Gao, E.-M.,Chian, R.C. The function of cumulus cells in oocyte growth and maturation and in subsequent ovulation and fertilization. *Cells* **2021**, 10, 2292. Doi: 10.3390/cells10092292.
  54. Peralta-Torres, J.A.,Aké-López, J.R.,Segura-Correa, J.C.,Aké-Villanueva, J.R. Effect of season on follicular population, quality and nuclear maturation of bovine oocytes under tropical conditions. *Anim. Reprod. Sci* **2017b**, 187, 47-53. Doi: 10.1016/j.anireprosci.2017.10.004.
  55. Rodrigues, A.T.,Ispada, J.,Risolia, H.B.P.,Rodrigues, T.M.,Lima, S.R.,Assumpcao, F.O.A.M.,Visintin, A.J.,Paula-Lopes, F.F. Thermoprotective effect of insulin-like growth factor 1 on in vitro matured bovine oocyte exposed to heat shock. *Theriogenology*, **2016**, 86, 2028-2039. Doi: 10.1016/j.theriogenology.2016.06.023.
  56. Andrew-Vázquez, C.,López-Gatius, F.,García-Ispuerto, I.,Maya-Soriano, M.J.,Hunter, R.H.F.,López-Béjar, M. Does heat stress provoke the loss of continuous layer of cortical granules beneath the plasma membrane during oocyte maturation? *Zygote*, **2010**, 18, 293-299. Doi: 10.1017/S0967199410000043.
  57. Santos, P.J.E.,Cerri, A.R.L.,Sartori, R. Nutritional management of donor cow. *Theriogenology* **2008**, 69, 1, 88-97. Doi: 10.1016/j.theriogenology.2007.09.010.

58. Chaveiro, A., Santos, P., Carvalhais, I., Agrícola, R., Faheem, M., Habibi, A., Silva, M.F. Sire effect and sperm apoptosis on bovine embryonic cleavage and subsequent in vitro embryo development. *Vet. Sci* **2010**. Doi:10.3814/2010/324529.
59. Lee, Y.J., Jung, G.Y., Seo, B.B. Effects of culture media conditions on production of eggs fertilized in vitro of embryos derived from ovary of high grade Hanwoo. *J. Anim. Sci. Technol* **2016**, 1, 58, 11. Doi: 10.1186/s40781-016-0093-5.
60. Ealy, D.A., Drost, M., Hansen, J.P. Developmental changes in embryonic resistance to adverse effect of maternal heat stress in cows. *J. Dairy Sci* **1993**, 76, 2899-2905. Doi: 10.3168/jds.S0022-0302(93)77629-8.
61. Payton, R.R., Romar, R., Coy, P., Saxton, A.M., Lawrence, J.L., Edwards, J.L. Susceptibility of bovine germinal vesicle-stage oocytes from antral follicles to direct effects of stress in vitro. *Biol. Reprod* **2004**, 71, 1303-1308. Doi: 10.1095/biolreprod.104.029892.
62. Silva, F.C., Sartorelli, S.E., Castilho, C.S.A., Satrapa, A.R., Puelker, Z.R., Razza, M.E., Ticianelli, S.J., Eduardo, P.H., Loureiro, B., Barros, M.C. Effects of heat stress on development, quality and survival of *Bos indicus* and *Bos taurus* embryos produced in vitro. *Theriogenology*, **2013**, 79, 351-357. Doi: 10.1016/j.theriogenology.2012.10.003.
63. Erdem, H., Karasahin, T., Alkan, H., Dursun, S., Satilmis, F., Guler, M. Effect of embryo quality and developmental stages on pregnancy rates during fresh embryo transfer in beef heifers. *Trop. Anim. Health Prod* **2020**, 52, 2541-2547. Doi: 10.1007/s11250-020-02287-6.
64. Marsico, V.T., Camargo, J., Valente, S.R., Sudano, J.M. Embryo competence and cryosurvival: molecular and cellular features. *Anim. Reprod* **2019**, 16, 423-439. Doi: 10.21451/1984-3143-AR2019-0072.
65. Jarrillo-Rodríguez, J., Castillo-Gallegos, E., Flores-Garrido, A.F., Valles de la Mora, B., Ramírez, A.L., Escobar-Hernández, R., Ocaña-Zavaleta, E. Forage yield, quality and utilization efficiency on native pastures under different stocking rates and seasons of the year in the mexicans humid tropics. *Trop. Subtrop. Agroecosystems* **2011**, 13, 417-427. Available on line: <https://www.revista.ccba.uady.mx/ojs/index.php/TSA/article/view/1379/680>.
66. Baena, M.M., Costa, A.C., Viera, G.C., Rocha, R.F.B., Ribeiro, A.R.B., Ibelli, A.M.G., Meirelles, S.L.C. Heat tolerance responses in a *Bos taurus* cattle raised in a Brazilian climate. *J. Therm. Biol* **2019**, 81, 162-169. Doi: 10.1016/j.jtherbio.2019.02.017.
67. Seijian, V., Bhatta, R., Gaughan, J.B., Dunshea, F.R., Lacetera, N. Review: Adaptation of animals to heat stress. *Animal* **2018**, 12, S2, s431-s444. Doi: 10.1017/S1751731118001945.
68. O'Brien, M.D., Rhoads, R.P., Sander, S.R., Duff, G.C., Baumgard, L.H. Metabolic adaptations to heat stress in growing cattle. *Domest. Anim. Endocrinol* **2010**, 38, 86-94. Doi: 10.1016/j.domaniend.2009.08.005.
69. Abbas, Z., Sammad, A., Hu, L., Fang, H., Xu, Q., Wang, Y. Glucose metabolism and dynamics of facilitative glucose transporters (GLUTs) under the influence of heat stress in dairy cattle. *Metabolites*, **2020**, 10, 312. Doi: 10.3390/metabo10080312.
70. Collier, R.J., Baumgard, L.H., Zimelman, R.B., Xiao, Y. Heat stress: physiology of acclimation and adaptation. *Anim. Front* **2019**, 9, 1, 1-19. Doi: 10.1093/af/vfy031.
71. Méndez, S.M., Soria, E.M., Galarza, R.L., Perea, P.F., Argudo, E.D. Effect of fetal calf serum on production and cryotolerance of in vitro bovine embryos from Ecuadorian creole heifers. *Reprod. Fertil. Dev* **2018**, 31, 141-142. Doi: 10.1071/RDv31n1Ab31.
72. Méndez, S.M., Argudo, E.D., Soria, E.M., Galarza, R.L., Perea, P.F. Efecto de la adición de melatonina en el medio de maduración y/o vitrificación de ovocitos sobre la producción in vitro de embriones bovinos. *Rev. Investig. Vet. Perú*, **2020**, 31. Doi: 10.15381/rivep.v31i1.17557.
73. Biscarini, F., Nicolazzi, E.L., Stella, A., Boettcher, P.J., Gandini, G. Challenges and opportunities in genetic improvement of local livestock breeds. *Front. Genet* **2015**, 6. Doi: 10.3389/fgene.2015.00033.
74. Martínez, R., Ávila, O., Pérez, J., Gallego, J., Onofre, H. Estructura y función del banco de germoplasma in vitro en Colombia. *Arch. Zoot* **2005**, 54, 545-550. Available on line: <https://www.redalyc.org/pdf/495/49520767.pdf>.
75. Chase Jr, C.C., Hammond, A.C., Olson, T.A., Murphy, C.N., Tewolde, A., Griffin, J.L. Introduction and evaluation of Romosinuano in the USA. *Latin American Arch. Anim. Prod*, **1997**, 5(Suppl. 2), 57-70. Available on line: [https://ojs.alpa.uy/index.php/ojs\\_files/article/view/223/216](https://ojs.alpa.uy/index.php/ojs_files/article/view/223/216).
76. Lima, S.R., Assumpcao, O.D.M.E., Visintin, J.A., Paula-Lopes, F.F. Alteracoes celulares inducidas pelo estresse térmico em embriões bovinos. *Braz. J. Vet. Res. Anim. Sci* **2013**, 50, 257-264. Doi: 10.11606/issn.2318-3659.v50i4p257-264.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.