

Article

Not peer-reviewed version

Effect of Menthol Treatment on the Sprouting and Quality of Potato Tuber

[Ye Xu](#), [Yang Qin](#), [Qiangjan Hou](#), [Defu Niu](#), [Qingmin Chen](#)*

Posted Date: 22 April 2024

doi: 10.20944/preprints202404.1391.v1

Keywords: potato tuber; menthol; sprouting inhibition; quality; glucoside alkaloids



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Effect of Menthol Treatment on the Sprouting and Quality of Potato Tuber

Ye Xu, Yang Qin, Qianqian Hou, Defu Niu and Qingmin Chen *

College of Food Science and Engineering, Shandong Agriculture and Engineering University, Jinan, 250100, PR China

* Correspondence: qmchen1207@163.com (Q.M. Chen)

Abstract: This study used Dutch potatoes at the end of dormancy as the material to explore the impact of menthol (0.2 and 0.5 g/kg based on potato mass) treatment on sprouting inhibition and potato quality. The findings revealed that a menthol concentration of 0.5 g/kg effectively inhibited potato tuber sprouting and significantly reduced glucoside alkaloids production. After a storage period of 15 days, the sprouting percentage and glucoside alkaloids content of potatoes treated with 0.5 g/kg menthol were observed to be significantly lower at 4.17% and 68.63 mg/kg, respectively, compared to the control group which exhibited values of 100% and 282.01 mg/kg, respectively ($P < 0.05$). Throughout the storage period, 0.5 g/kg of menthol promoted respiration, reduced malondialdehyde production in potatoes, inhibited polyphenol oxidase activity, and slowed down tissue browning. Additionally, it mitigated the decline in starch and soluble protein content, inhibiting the accumulation of reducing sugars.

Keywords: potato tuber; menthol; sprouting inhibition; quality; glucoside alkaloids

1. Introduction

Potato (*Solanum tuberosum* L.) is an annual plant belonging to the Solanaceae family. With a global production exceeding 300 million tonnes each year [1,2], potatoes hold considerable economic value due to their high yield and nutritional quality, playing a crucial role in ensuring food security, driving industrial poverty alleviation, and fostering regional economic revitalisation across all nations, especially in developing countries [3]. However, potatoes are highly susceptible to sprouting during storage and transport, accompanied by the production of toxins such as glucoside alkaloids [4]. According to statistics, the annual loss of potatoes due to sprouting accounts for about 20–25% of the total potato production [5], resulting in substantial economic losses and posing food safety hazards.

Currently, various methods are employed for inhibiting potato sprouting during storage, including chemical treatment, irradiation, phytohormone treatment, and plant essential oil treatment. Common chemicals used for this purpose include chlorpropham (CIPC) and maleic hydrazide. CIPC, the most widely utilised commercial potato sprout inhibitor, hinders tuber sprouting by affecting spindle formation and interfering with cell division in meristematic tissues [6]. However, the safety concerns related to pharmaceutical residues in CIPC pose a significant threat to human health [7], leading to its complete ban in the EU since 2020. Maleic hydrazide, also known as 3,6-dihydroxypyridazine, acts as a growth regulator with various functions, including inhibiting potato sprouting. Despite its common use in potato cultivation, many countries lack established maximum residue limits for maleic hydrazide in potatoes, limiting its application. Radiation treatments, though effective, are irreversible, expensive, and have low market acceptance.

In phytohormone treatment, ethylene is a primary agent used to inhibit potato sprouting by influencing carbohydrate metabolism [8]. However, continuous introduction of ethylene into the potato storage environment requires specialised equipment for sealing and introduction, leading to higher costs. Additionally, ethylene can enhance the activity of potato polyphenol oxidase (PPO), accelerating potato browning and adversely affecting both potato quality and subsequent processing quality [9].

Recently, the inhibitory effect of plant essential oils on potato sprouting has gained attention due to their status as pure natural extracts. Essential oils, such as carvone, lemongrass oil, jasmine oil, and menthol, have been studied. Oosterhaven et al. [10] found that carvone effectively inhibits sprouting in a potato bud growth model, and its inhibitory effect is reversible. Yuan et al. [11] demonstrated that lemongrass essential oil effectively inhibits potato sprouting while significantly impacting potato sugar metabolism, delaying the decline of starch and soluble protein content during storage and preserving the nutritional components of potato tubers. Huang et al. [12] investigated the effects of jasmine essential oil and menthol on sprouting inhibition and quality in fresh potatoes. However, the effectiveness of menthol in inhibiting potato sprouting at the end of dormancy and its impact on quality remain unexplored and require further study.

The present study aims to investigate the impact of different concentrations of menthol treatment on Dutch potatoes at the end of dormancy. The effects of two concentrations of menthol on potato sprouting rate, glucoside alkaloid content, and quality parameters during storage at room temperature were investigated. Physiological changes, browning, starch and reducing sugar content, as well as amino acid composition were analyzed in order to provide a potential solution for sprout control.

2. Materials and Methods

2.1. Materials

The potatoes utilised in this study were Dutch potatoes procured from the Guanzhuang Vegetable Wholesale Market in Zhoucun District, Zibo City. Harvested in July 2022 after a dormant period, these potatoes were subsequently transported to the Fruit and Vegetable Storage and Preservation Laboratory of Shandong Agriculture and Engineering University. Fresh potatoes, characterised by uniform size, devoid of mechanical damage or diseases, were meticulously chosen for the experimentation. Menthol (C₁₀H₂₀O, CAS 89-78-1, purity: 99%) was obtained from the Wuhan Jixin Yibang Biotechnology Co., Ltd.

2.2. Treatments

Potato tubers were segregated into plastic baskets, with each group comprising six baskets, each containing 10 potatoes (approximately 5 kg per basket). Three distinct sets were arranged: Control CK (no menthol application), Group 1 treated with an effective concentration of 0.2 g/kg menthol, and Group 2 treated with 0.5 g/kg menthol. The menthol and an inert material mixture were uniformly applied to the potato surfaces.

All groups were stored at room temperature (25 °C) for 15 days, with samples collected at 3-day intervals. For analysis, three baskets from each group were allocated to assess sprouting percentage, sprouting index, weight loss, and respiration intensity. The remaining three baskets were utilised for determining additional indices, including moisture content, browning degree, malondialdehyde (MDA) content, PPO activity, starch content, and soluble protein content.

To ensure sampling uniformity, the four-quadrant method was employed, and the collected samples were promptly frozen using liquid nitrogen and stored at -80°C for future analysis.

2.3. Determination of Sprouting Percentage and Sprouting Index

The sprouting percentage and sprouting index were assessed at 3-day intervals. Refer to method of Yuan et al. [11] for the determination of sprouting percentage. Ten potatoes were randomly selected from each treatment, and the length of the longest sprout was measured using a vernier caliper. Sprouts with a length less than 2 mm were classified as non-sprouting or in an initial sprouting state. Refer to method of Şanlı et al. [13] for the determination of sprouting index. Ten potatoes were randomly chosen for each treatment. The number of buds on each tuber was documented, and the length of the longest bud was measured with a vernier caliper, then averaged. The bud lengths were categorised into eight grades (Table 1), and the sprouting percentage (%) was computed as the numbers of sprouted tubers / total number of tubers ×100%; and the sprouting index (%) was was

computed as the sum of the length of sprouted tubers / (total number of tubers × highest length) ×100%.

Table 1. Sprouting grading criteria.

Bud length/mm	sprouting grade	Bud length/mm	sprouting grade
0~2	0	15~20	4
2~5	1	20~25	5
5~10	2	25~30	6
10~15	3	> 30	7

2.4. Determination of Glucoside Alkaloids Content

The extraction method of glucoside alkaloids followed slight modifications of method of [Oishi et al.](#) [14] procedure: 5.0 g of sample were mixed with 30 mL of acidified acetonitrile ((acetonitrile: formic acid = 99:1): water = 5:1, v:v), vortexed and oscillated for 5 min. Then, sodium acetate (1.0 g) and anhydrous magnesium sulfate (2.0 g) were added, followed by another vortexing and oscillation for 5 min before centrifugation at a rate of 1.0×10^4 g for 10 min. The supernatant was filtered through a 0.45 μm organic membrane filter. The detection of α -solanine and α -chaconine was performed using HPLC (Shimadzu Corporation, Tokyo, Japan).

Separation occurred on a C_{18} column (Waters Corporation; dimensions: 4.6 mm * 250 mm; particle size: 5 μm), employing a mixture consisting of dipotassium hydrogen phosphate solution (pH adjusted to 7.6 with potassium dihydrogen phosphate; concentration: 0.1 mol L^{-1}), acetonitrile, and water in a ratio of 1: 6: 3 as mobile phase. A stepwise gradient program was used for isocratic elution. The injection volume for all samples was 10 μL . The flowrate was set at 1.2 mol L^{-1} , the temperature of the column incubator was set at 40 $^{\circ}\text{C}$, and the detector wavelength was set at a wavelength of 208 nm. Standard solutions in the concentration range from 10 to 200 mgL^{-1} were dissolved in acetonitrile and used for quantification purposes. Standard curves were constructed with concentration on the abscissa and peak area on the ordinate.

The glucoside alkaloids content (x) was calculated as $c \times v \times f / m$. Where, x-The content of the target in the sample, in grams (mg kg^{-1}); c-To determination of target concentration in sample solution from calibration curve, in grams (mg kg^{-1}); v-The final constant volume of the sample solution to be measured, in grams (mL); m-The mass of the sample is measured, in grams (g); f -Diluted multiples.

2.5. Determination of Moisture Content and Weight Loss

2.5.1. Determination of Moisture Content

The direct drying method, as outlined in GB5009.3-2016 [15] was employed for assessing the moisture content in potato tubers during storage. The moisture content (x) was calculated as $(m_1 - m_2) / (m_1 - m_3) \times 100$. Where, x-the content of moisture in the sample, in grams ($\text{g}/100 \text{ g}$); m_1 - The mass of the weighing bottle (plus sea sand, glass rod) and the sample, in grams (g); m_2 - The mass of the weighing bottle (plus sea sand and glass rod) and the sample after drying, in grams (g); m_3 - The mass of the weighing bottle (plus sea sand and glass rod), in grams (g); 100-Unit conversion factor.

2.5.2. Determination of Weight Loss

The mass of each tuber before and after storage were individually determined. The rate of weight loss (x) was calculated as $(m_1 - m_2) / m_1 \times 100$. Where, x -Weight loss rate in sample, in grams (%); m_1 - Mass of tuber before storage, in grams (g); m_2 -Mass of tubers after storage, in grams (g); 100-Unit conversion factor.

2.6. Determination of Browning

2.6.1. Determination of Browning Index

The browning index determination was done according to the method of Wang et al. [16] with minor modifications. The supernatant was obtained by collecting a 5 g sample of frozen and ground

potato tubers. Subsequently, 3 ml of pre-cooled (4 °C) 95% ethanol solution was added, followed by incubation at 4 °C for 6 hours and centrifugation at 12,000 g for 20 min. The absorbance at a wavelength of 410 nm was measured using the control group containing only 95% ethanol. The calculation formula used was $BD = \Delta OD_{410} \times 6$ (where ΔOD_{410} represents the absorbance value at a wavelength of 410 nm).

2.6.2. Determination of Total Phenolic Content

Folin phenol colorimetry was employed to determine the total phenolic content in extracts of potato samples as described by [Joly et al.](#) [17]. A frozen potato sample was pulverized into a fine powder using a grinder and liquid nitrogen. Take 3.0 g of the potato powder and grind it thoroughly in a mortar, then add 10 mL of methanol and refrigerate at 4 °C for 12 h. Subsequently, mix 300 μ L of the extract with 1.7 mL of distilled water, followed by adding 1 mL each of folin phenol solution and a 7.5% anhydrous sodium carbonate solution. The mixture was kept warm in a water bath at 25 °C for 2 h, after which the absorbance was measured at a wavelength of 765 nm. Notably, the standard curve equation is $y = 0.0111x + 0.0203$ with an R^2 value of 0.9954 (where x represents gallic acid concentration and y represents light absorption).

2.6.3. Determination of PPO Activity

PPO activity in potato tubers was determined using a plant PPO ELISA kit. After allowing the aluminum foil bag to equilibrate at room temperature for 20 min, remove the required slats and seal the remaining ones in a self-sealing bag before storing them at 4 °C. Set up standard product holes and sample holes, and add different concentrations of standard product (50 μ L) to each standard hole. Then add 10 μ L of sample to the sample hole, followed by adding 40 μ L of sample diluent; Blank holes are not added. In addition to blank holes, add horseradish peroxidase (HRP)-labeled detection antibody (100 μ L) to each well containing standards or samples, seal with a sealing plate film, and incubate at 37 °C in a water bath or incubator for 1 h. Discard liquid contents, pat dry on absorbent paper, fill each well with washing solution for one minute before shaking off excess liquid and patting dry on absorbent paper again, repeat this process five times. Add substrate A and B (50 μ L each) into every well then incubate without light at 37 °C for 15 min. Adding termination solution (50 μ L) into every well then measure OD value within fifteen minutes using a wavelength of 450 nm. The standard curve equation is $y = 0.001x + 0.1139$ with an R^2 value of 0.9923 (where x represents PPO activity and y represents the OD value at a wavelength of 450 nm).

2.7. Determination of Respiration Intensity and MDA Content

2.7.1. Determination of Respiration Intensity

The respiration intensity of potato tubers was measured using a gas analyser, following the method of [Yuan et al.](#) [11]. The respiration intensity (Q) was calculated as $(V \times N \times 1.894 \times 1000) / (m \times t)$. Where Q is the respiration intensity [mg/(kg·h)]; V is the volume of the container (plastic basket volume with potato volume, L); N is the percentage of CO₂ (%); 1.894 is the density of carbon dioxide at 10°C under normal pressure; m is the mass of the sample (kg); and t is the airtight time (h).

2.7.2. Determination of MDA Content

The MDA content of the samples was determined following the method of [Palamutoğlu et al.](#) [18] with minor modifications. The potato sample (1 g) was homogenized with 10 mL of pre-cooled trichloroacetic acid solution (1 g/L) and centrifuged at 4000rpm for 15 min at 4°C. The resulting supernatants were used for MDA analysis. A mixture of 0.6 mL supernatant and 1 mL thiobarbituric acid solution (6.7 g/L) was prepared in glass tubes and incubated at a temperature of 95 °C for a duration of 20 min. Subsequently, the mixture was cooled using running water, followed by measuring its absorbance at wavelengths of 450 nm, 532 nm, and 600 nm. Finally, equation (6) was employed to calculate the MDA content (C) was calculated as $6.45 \times (OD_{532} - OD_{600}) - 0.56 \times OD_{450}$. In the formula, OD_{450} , OD_{532} , and OD_{600} represent the absorbance at 450, 532, and 600 nm, respectively. C denotes the concentration of MDA in μ mol/kg.

2.8. Determination of Reducing Sugar and Starch Content

2.8.1. Determination of Reducing Sugar Content

The concentration of reducing sugar was determined using the 3,5-dinitrosalicylic acid method as described by Hu et al. [19]. A total of 1.0 g of potato powder was weighed and washed into a conical flask with 50 mL of distilled water. After thorough mixing, the mixture was heated in a water bath at 50 °C for 30 min. Once cooled, the homogenate was filtered and the filtrate transferred to a 100 mL volumetric flask for volume adjustment. The absorbance of the resulting solution was measured at a wavelength of 540 nm. The standard curve equation is $y=0.9817x-0.0412$ with an R^2 value of 0.9962 (where x represents the quality of sample and y represents the OD value at a wavelength of 540nm). The reducing sugar content was calculated as $\text{Reducing sugar content} = \frac{m' \times V \times N}{V_s \times m}$. Where, m' -Mass of glucose in the standard curve, in grams (μg); V -Mass of total volume of extract, in grams (mL); N -Mass of sample dilution fold; V_s - Mass of the volume of sample aspirated during determination, in grams (mL); m -Mass of the sample, in grams (g).

2.8.2. Determination of Starch Content

For the determination of starch content, the iodine-starch colorimetric method [11] was followed. The starch content was calculated using equation $\text{Starch content}/\% = \frac{m' \times V \times N}{V_s \times m \times 10^6}$. Where, m' is the mass of starch in the standard curve (μg); V is the total volume of extract (mL); N is the sample dilution fold; V_s is the volume of sample aspirated during determination (mL); and m is the sample mass (g).

2.9. Determination of Soluble Protein and Amino Acid Content

2.9.1. Determination of Soluble Protein Content

The soluble protein content in potatoes was determined following the method of [Wegener et al.](#) [20]. The calculation was performed using equation of soluble protein content $= \frac{m' \times V}{V_s \times m \times 1000}$. Where m' is the mass of soluble protein in the standard curve (μg); V is the total volume of the extract (mL); V_s is the volume of the sample aspirated during determination (mL); and m is the sample mass (g).

2.9.2. Determination of Amino Acid Content

The amino acid content was determined using an amino acid analyser according to the method outlined in GB 5009.124-2016 [21] in China.

2.10. Data Analysis

Three parallel trials were conducted for each treatment group, and the significance of differences in the test results was analysed using SPSS 23.0 software. After one-way analysis of variance, Duncan's multiple comparisons were employed to analyse differences between the data at a significance level of $P < 0.05$. Graphs were generated using Excel 2010 software, and the results were expressed as mean \pm standard deviation.

3. Results and Analysis

3.1. Effect of Menthol Treatment on Potato Tuber Sprouting

During the storage period, daily observations of potato tuber sprouting were made, and on the 15th d, a morphological picture of potato sprouting was taken. As depicted in Figure 1A, both the CK group and the 0.2 g/kg group exhibited sprouting, with budding observed on the lateral side of the tuber. The buds appeared yellow, while the apical buds were black. In contrast, the 0.5 g/kg group showed no signs of sprouting. The tuber eyes were completely black, the apical bud did not sprout, and other parts of the eyes were necrotic. This suggests that menthol treatment can damage the meristematic tissues of potato tubers, causing injury to the apical meristems, resulting in skin damage and bud necrosis, ultimately inhibiting potato sprouting.




Figure 1. Impact of menthol treatment on the sprouting state (A), sprouting rate (B), sprouting index (C), and solanine content (D) in potato (*Solanum tuberosum* L) tubers. Different letters on the same day indicate a significant ($P < 0.05$) difference. The data represent mean \pm SD of three replicates.

The impact of menthol on potato sprouting percentage and sprouting index is illustrated in Figure 1B-1C, respectively. In Figure 1B, it is evident that from day 0 to day 3, no sprouting occurred in any of the three treated groups. By day 6, both the control group (CK) and the 0.2 g/kg group initiated sprouting, with sprouting percentages of 33.3% and 20.8%, respectively, while the 0.5 g/kg group remained free from sprouting. On day 12, the sprouting percentages for CK and the 0.2 g/kg group were 100% and 91.67%, respectively, while the 0.5 g/kg group continued to show no signs of sprouting. After 15 days of storage, the sprouting percentage for the 0.5 g/kg group was 4.1%, significantly lower than that of CK and the 0.2 g/kg group ($P < 0.05$).

Turning to Figure 1C, it is observed that starting from day 6, the sprouting index of the CK group and the 0.2 g/kg group exhibited an increasing trend. Moreover, the sprouting index of the CK group was significantly higher than that of the 0.2 g/kg group ($P < 0.05$). At the conclusion of storage, the sprouting index for CK, 0.2 g/kg, and 0.5 g/kg groups was 70.8%, 56.6%, and 0.8%, respectively, indicating significant differences ($P < 0.05$) in the sprouting index among these three groups. These results affirm that menthol effectively inhibits potato tuber sprouting.

3.2. Impact of Menthol Treatment on Glucoside Alkaloids Content in Potatoes

Following storage, potatoes may exhibit elevated glucoside alkaloids levels due to tuber sprouting, particularly in the sprouting and greening sections of the potato. Figure 1D illustrates the effect of menthol on glucoside alkaloids content in potatoes after storage. After 15 days of storage, the glucoside alkaloids content in the 0.5 g/kg group was 68.63 mg/kg, significantly lower than that in the CK group (282.01 mg/kg) and the 0.2 g/kg group (208.91 mg/kg) ($P < 0.05$). The Chinese national limit for glucoside alkaloids is 200 mg/kg, confirming that the glucoside alkaloids content in the 0.5 g/kg group complies with national safety standards. Comparative analysis across different storage times and treatment groups suggests that high concentrations of menthol effectively inhibit glucoside alkaloids accumulation in potatoes by suppressing sprouting.

3.3. Impact of Menthol Treatment on Potato Moisture Content, Weight Loss, Respiration Intensity, and MDA Content

Figure 2A-2B showcase the effect of menthol on moisture content and weight loss of potatoes during storage. As observed in Figure 2A, the moisture content of potatoes significantly increased during the initial six days of storage. Subsequently, after the sixth day, the moisture content exhibited a gradual decline. This phenomenon can be attributed to the initial storage of potatoes at 4°C. The influence of enzymes, such as ADP-glucose pyrophosphorylase, UDP-glucose pyrophosphorylase,

and sucrose synthase, leads to starch conversion and breakdown into reducing sugars [22]. The accumulation of these reducing sugars results in low-temperature sweetening. During this process, the free water content becomes lower than the bound water content, contributing to an overall lower moisture content [23]. Conversely, in the early stages of potato storage, as the temperature rises, previously accumulated reducing sugars revert to starch, causing the free water content to exceed the bound water content and resulting in a higher moisture content. Comparative analysis among experimental groups indicates that the moisture content of the 0.5 g/kg group surpassed that of the 0.2 g/kg group and the CK group during storage, indicating that a high menthol concentration effectively reduces water loss and maintains higher moisture content.




Figure 2. Impact of menthol treatment on the moisture content (A) and weight loss (B) in potato (*Solanum tuberosum* L) tubers.

In Figure 2B, the weight loss rate of potatoes gradually increased during storage. The CK group exhibited a significantly higher weight loss rate than the 0.2 g/kg and 0.5 g/kg groups ($P < 0.05$). At the end of the storage period, the weight loss rates were 0.83% in the CK group, 0.74% in the 0.2 g/kg group, and 0.67% in the 0.5 g/kg group ($P < 0.05$). This indicates that menthol-treated potatoes effectively inhibit respiration, transpiration, and other processes, thereby reducing nutrient losses.

Respiration intensity reflects the strength of plant metabolism, as depicted in Figure 3A, showcasing changes in potato respiration intensity during storage. The three groups exhibited an initial increase followed by a decrease in respiration intensity throughout the storage period, with the most pronounced trend observed in the 0.2 g/kg group. Comparing different experimental groups, on day 3 of storage, the respiration intensity of the 0.5 g/kg group (9.78 mg/(kg·h)) was significantly higher than that of the CK group (4.38 mg/(kg·h)) and the 0.2 g/kg group (7.10 mg/(kg·h)) ($P < 0.05$). This may be attributed to the high menthol concentration triggering a stress response in the potatoes. On day 15 of storage, the respiration intensity in the 0.5 g/kg group (3.13 mg/(kg·h)) was significantly lower ($P < 0.05$) than that in the CK group (3.70 mg/(kg·h)) and the 0.2 g/kg group (4.11 mg/(kg·h)). This suggests that menthol-treated potatoes displayed robust early-stage metabolic activity and maintained a higher respiration intensity. In the middle and later stages, the treated potatoes exhibited a relatively stable metabolic state, indicating that menthol treatment can inhibit potato respiration intensity. This aligns with the results of sprouting percentage and sprouting index, indicating increased respiration intensity during potato tuber sprouting in the CK group.




Figure 3. Impact of menthol treatment on the respiration intensity (A) and MDA (malondialdehyde) content (B) in potato (*Solanum tuberosum* L) tubers. Different letters indicate a significant ($P < 0.05$) difference. Data are presented as the mean \pm SD of three replicates.

MDA is a major byproduct in the process of cell membrane lipid peroxidation, with a high concentration having a detrimental impact on the cell membrane [24]. MDA content serves as an indicator of membrane lipid peroxidation, reflecting damage to the membrane system and plant stress tolerance [25]. Figure 3B depicts the fluctuations in potato MDA content during storage. The MDA content across the three groups exhibited a general decreasing trend throughout storage. Comparative analysis indicated that the 0.5 g/kg group consistently maintained lower MDA content than the CK and 0.2 g/kg groups. At the end of the storage period (15 days), the MDA content was 0.18 $\mu\text{mol/g}$ in the 0.5 g/kg group, significantly lower than that in the CK group (0.32 $\mu\text{mol/g}$) and the 0.2 g/kg group (0.26 $\mu\text{mol/g}$) ($P < 0.05$). This underscores the substantial inhibitory effect of high menthol concentration on the accumulation of harmful substances in potatoes.

3.4. Impact of Menthol Treatment on Browning Index, Total Phenol Content, and PPO Activity in Potatoes

The browning index serves as an indirect gauge of soluble pigments generated in the browning reaction, reflecting the degree of browning in fruits and vegetables [26]. Figure 4A illustrates the changes in the browning index among the three potato groups during storage. Over time, the browning index in all groups exhibited an ascending and then descending trend. Comparative analysis indicated that the browning index of the 0.5 g/kg group remained consistently lower than that of the CK and 0.2 g/kg groups throughout the storage period. At the conclusion of the storage period (15 days), the browning index reached 0.42 in the 0.5 g/kg group, significantly lower ($P < 0.05$) than the values of 0.54 and 0.51 in the CK and 0.2 g/kg groups, respectively. This underscores the effective ability of 0.5g/kg menthol to delay and inhibit potato browning, preserving the quality of potatoes and achieving a preservation effect.




Figure 4. Impact of menthol treatment on the browning index (A), total phenol content (B) and PPO (polyphenol oxidase) activity in potato (*Solanum tuberosum* L) tubers.

Plants harbor a diverse array of phenolics, crucial substrates for the enzymatic coloring of fruits and vegetables [27]. In Figure 4B, the total phenol content in potato tubers exhibited erratic variations across all groups during the storage period. There was a declining trend in total phenol content from day 3 to day 6, suggesting phenolics were utilised by phenolases, contributing to enzymic browning and accelerating the tuber browning rate, consistent with the observed changes in browning degree. In the later stages, as the browning rate slowed, total phenol accumulation increased, resulting in a gradual rise in total phenol content after 12 days. Upon comparison of experimental groups, the total phenol content of the 0.5 g/kg group consistently lagged behind that of the CK and 0.2 g/kg groups. At the end of storage, the total phenol content in the 0.5 g/kg group was 0.23 mg/g, significantly lower than the CK group (0.27 mg/g) and the 0.2 g/kg group (0.31 mg/g) ($P < 0.05$). This indicates that menthol-treated potato tubers experienced a gradual change in total phenol content, exhibiting a slow browning rate and maintaining good storage quality.

PPO serves as the primary enzyme responsible for the enzymatic browning of potatoes, impacting their flavor and quality [28]. PPO activity is a crucial indicator for assessing the extent of browning. Additionally, PPO functions as a terminal oxidase system in plant tissues' respiration, influencing dormancy and sprouting in tubers [29]. Within 15 days of storage, the binding of PPO to the inner vesicle membrane may be compromised, leading to PPO activation [30]. As depicted in Figure 4C, PPO activity exhibited an initial increase (during the first six days) followed by a decrease during storage across all groups. Comparative analysis among the three groups on day 6 revealed that the PPO activity of the 0.5 g/kg group was 301.43 U/mL, significantly lower than that of the CK group (400.10 U/mL) and the 0.2 g/kg group (349.77 U/mL) ($P < 0.05$). Even at the end of the storage period, the PPO activity of the 0.2 g/kg group remained significantly lower than that of the CK group and the 0.2 g/kg group ($P < 0.05$). In conclusion, menthol treatment demonstrated effective inhibition of PPO activity, showcasing a positive correlation with the observed changes in browning degree in this study.

3.5. Effect of Menthol Treatment on Reducing Sugar and Starch Content of Potatoes

In Figure 5A, it is observed that during the initial storage phase, the potato exhibited the highest reducing sugar content of 173.47 mg/g. This could be attributed to the storage at 4°C before testing, where the activity of amylolytic enzyme in the tuber surpassed that of amylopectin synthase, resulting in the conversion of starch to reducing sugar [31]. In the early storage period (up to day 6), reducing sugar content in all groups declined due to increased respiration and the conversion of reducing sugars into starch after removal from cold storage. However, a temporary surge in reducing sugar content occurred during days 6~12, followed by a subsequent decline from days 12~15. Comparative analysis at the late storage stage revealed that the reducing sugar content of the 0.5 g/kg group was lower than that of both the CK and 0.2 g/kg groups. At the end of storage, the reducing sugar content of the 0.5 g/kg group was 80.30 mg/g, significantly lower than that of both the CK group (121.86 mg/g) and the 0.2 g/kg group (99.58 mg/g) ($P < 0.05$). These findings indicate that menthol markedly inhibited the accumulation of reducing sugars during the storage of potato tubers.




Figure 5. Impact of menthol treatment on the reducing sugar content (A) and starch content (B) in potato (*Solanum tuberosum* L) tubers.

Starch serves as the primary nutrient and energy source in potato tubers, reflecting changes in storage quality [32]. In Figure 5B, the starch content initially rises, falls, then rises again with prolonged storage. Comparisons reveal that the 0.5 g/kg group consistently maintains higher starch content than the CK group. At the 6th day, the 0.5 g/kg group reaches 14.92%, significantly surpassing the CK group (8.33%) and the 0.2 g/kg group (10.70%) ($P < 0.05$). At the storage end, starch content for the 0.5 g/kg, 0.2 g/kg, and CK groups is 14.84%, 10.86%, and 21.89%, respectively, compared to their initial values of 95.18%, 72.21%, and 76.27%. In summary, high-concentration menthol treatment effectively retards starch content decline, favoring quality maintenance during storage.

3.6. Effect of Menthol Treatment on Soluble Protein and Amino Acid Content of Potatoes

Soluble proteins serve as the foundation for physiological metabolism in fruit and vegetable cells, with intricate enzyme systems primarily comprising non-membrane-bound proteins [20]. Table 2 illustrates the impact of menthol on soluble protein content in potato tubers. Throughout storage, soluble protein content steadily increased. The CK group consistently exhibited lower soluble protein content, possibly due to untreated potato sprouting, leading to higher metabolic activity and energy consumption. At the 12th and 15th days of storage, the 0.5 g/kg group showed significantly higher soluble protein content (4.06 mg/g) than the CK and 0.2 g/kg groups during the same period ($P < 0.05$), indicating menthol's effective delay in soluble protein loss, showcasing superior preservation.

Table 2. Impact of menthol treatment on the soluble protein content in potato tubers.

soluble protein content (mg/g)	treatment groups		
	CK	0.2 g/kg Men	0.5 g/kg Men
0 d	3.18±0.07 ^a	3.18±0.07 ^a	3.18±0.07 ^a
3 d	3.21±0.05 ^a	3.62±0.05 ^b	3.69±0.04 ^b
6 d	2.99±0.12 ^a	3.38±0.11 ^b	3.43±0.11 ^b
9 d	3.55±0.17 ^a	3.78±0.09 ^a	3.63±0.10 ^a
12 d	3.73±0.07 ^a	3.71±0.03 ^a	4.06±0.08 ^b
15 d	3.87±0.05 ^a	3.86±0.06 ^a	4.06±0.08 ^b

The different letters indicate a significant ($P < 0.05$) difference. Data are presented as the mean ± SD of three replicates.

Amino acids, crucial osmotic regulators, can reduce potato browning by forming colourless adducts with quinones or acting as chelating agents [9]. Table 3 displays the menthol effect on amino acid content in potatoes before and after storage. At the storage end, the 0.5 g/kg group exhibited lower glycine, methionine, and proline content (0.39 mg/g, 0.021 mg/g, and 0.43 mg/g, respectively) compared to the 0.2 g/kg group and the CK group. This may be attributed to untreated and low-concentration menthol-treated potatoes containing higher glycine, methionine, and proline, forming coloured catechol-amino acid adducts and inducing browning [33]. However, further exploration is needed for a comprehensive understanding of this mechanism. The majority of amino acid contents in potatoes remained stable during storage, suggesting minimal impact on amino acid content due to menthol.

Table 3. Impact of menthol treatment on the amino acid content in potato tubers.

amino acid content (mg/g)	0 d		15 d	
	control value	CK	0.2 g/kg Men	0.5 g/kg Men
aspartic acid	4.7	4.6	4.3	4.2
threonine	0.47	0.5	0.49	0.51
threonine	0.42	0.43	0.4	0.42

glutamic acid	3.1	2.6	2.7	2.7
glycine	0.43	0.43	0.43	0.39
alanine	0.53	0.48	0.51	0.52
valine	0.77	0.77	0.76	0.75
methionine	0.03	0.026	0.025	0.021
isoleucine	0.49	0.48	0.47	0.48
leucine	0.78	0.78	0.79	0.8
tyrosine	0.24	0.22	0.28	0.21
phenylalanine	0.66	0.69	0.7	0.68
lysine	0.74	0.8	0.79	0.81
histidine	0.37	0.4	0.37	0.38
arginine	0.55	0.56	0.54	0.59
proline	0.39	0.49	0.45	0.43
total amino acid	14.67	14.256	14.005	13.891

4. Discussions

Potatoes are susceptible to various quality deteriorations during storage, including sprouting, browning, and nutrient loss [34]. This study investigated the impact of menthol on sprout inhibition and storage quality in potato tubers, utilising two menthol concentrations. The findings indicate that the postharvest application of 0.5 g/kg menthol effectively hinders tuber sprouting, leading to a decreased sprouting percentage and sprouting index, along with reduced glucoside alkaloids production. Notably, 0.5 g/kg menthol treatment induced blackening of potato tuber buds and disrupted bud tissue during sprouting, resulting in delayed sprouting.

In terms of quality, the treatment with 0.5 g/kg menthol influenced the respiration intensity of potatoes, slowing down browning, inhibiting PPO activity, and reducing the production of MDA. Furthermore, it contributed to the preservation of starch, reducing sugars, and soluble protein content, while inhibiting the accumulation of reducing sugars. In summary, the application of 0.5 g/kg menthol proved effective in inhibiting potato sprouting, minimising nutrient loss, and preserving overall storage quality.

This study observed that menthol treatment maintained higher respiration levels in potato tubers during the early storage stage, which declined after wound healing, and suppressed potato respiration intensity during the late storage stage [35]. PPO activity and total phenolic content in potatoes correlate with browning [36]. Results demonstrated that high menthol concentrations significantly reduced PPO activity and total phenol content at the end of storage compared to control and low concentration groups, positively correlating with the browning degree [37]. The high menthol treatment delayed the decline in potato starch content, inhibited reducing sugar production, and maintained low reducing sugar levels. After breaking dormancy, proteins broke down into amino acids to support bud development [38]. Liu et al. [39] noted differentially expressed proteins during secondary growth of meristematic tissues, leading to potato tuber dormancy break. Soluble protein content in the high-concentration menthol-treated group remained higher than the control and low-concentration groups at the end of storage, suggesting menthol slows soluble protein depletion and delays potato aging, with less obvious effects on amino acid content.

5. Conclusions

In conclusion, menthol, a natural plant ingredient, not only exhibits a significant sprout inhibition effect but also enhances the overall quality of potatoes during storage, preserving their commercial value. Moreover, menthol significantly affects the colour quality and carbohydrate content during potato sprouting, with a relatively minor influence on protein metabolism. Although the glucoside alkaloids content in the 0.5 g/kg menthol treatment group was found to be below the

national safety standard, its presence remained detectable. Further investigation is needed to unveil the specific physiological, biochemical, and molecular mechanisms underlying these observed effects.

Author Contributions: Ye Xu: Writing-original draft, Data curation, Investigation, Visualization. Yang Qin: Investigation, Methodology, Data curation. Qianqian Hou: Investigation, Methodology. Defu Niu: Methodology and Supervision, Data curation. Qingmin Chen: Methodology Establishment, Parameter Testing, Funding acquisition.

Acknowledgments: This project was supported by the Shandong Provincial Science and Technology of Innovation Ability Enhancement Project of Small and Medium Enterprises (2023TSGC0153), the Project of Innovation Team of Potato Storage and Processing of Shandong Agriculture and Engineering University (sgykcxtd2020-02), and the Agricultural Industrial Technology System of Shandong Province (SDAIT-05) in China.

Conflicts of Interest: The authors report there are no competing interests to declare.

References

- Gumbo, N.; Magwaza, L. S.; Ngobese, N. Z. Evaluating ecologically acceptable sprout suppressants for enhancing dormancy and potato storability: a review. *Plants*. **2021**, *10*(11), 2307.
- Sampaio, S. L.; Petropoulos, S. A.; Alexopoulos, A.; Heleno, S. A.; Santos-Buelga, C.; Barros, L.; Ferreira, I. C. Potato peels as sources of functional compounds for the food industry: A review. *Trends in Food Science & Technology*. **2020**, *103*, 118-129.
- Ahmadu, T.; Abdullahi, A.; Ahmad, K.; Ahmadu, T.; Abdullahi, A.; Ahmad, K. The role of crop protection in sustainable potato (*Solanum tuberosum* L.) production to alleviate global starvation problem: An overview. *Solanum tuberosum-A Promising Crop for Starvation Problem*. **2021**, 19-51.
- Liu J. M.; Wang S. S.; Zheng X. Antimicrobial activity against phytopathogens and inhibitory activity on glucoside alkaloids in potatoes of the endophytic bacteria isolated from potato tubers. *Frontiers in microbiology*. **2020**, *11*: 570926.
- Zhao, S.; Fu, M. R.; Liu, X. H. Research progress of potato sprout inhibition technology. *Food Science*. **2013**, *17*, 338-343.
- Alamar, M. C.; Tosetti, R.; Landahl, S.; Bermejo, A.; Terry, L. A. Assuring potato tuber quality during storage: A future perspective. *Frontiers in Plant Science*. **2017**, *8*, 2034.
- Paul, V.; Ezekiel, R.; Pandey, R. Sprout suppression on potato: need to look beyond CIPC for more effective and safer alternatives. *Journal of Food Science and Technology*. **2016**, *53*, 1-18.
- Dai, H.; Fu, M.; Yang, X.; Chen, Q. Ethylene inhibited sprouting of potato tubers by influencing the carbohydrate metabolism pathway. *Journal of Food Science and Technology*. **2016**, *53*, 3166-3174.
- Levaj, B.; Pelaić, Z.; Galić, K.; Kurek, M.; Šćetar, M.; Poljak, M.; Repajić, M. Maintaining the quality and safety of fresh-cut potatoes (*Solanum tuberosum*): Overview of recent findings and approaches. *Agronomy*. **2023**, *13*(8), 2002.
- Oosterhaven, K.; Hartmans, K. J.; Scheffer, J. J. C. Inhibition of potato sprout growth by carvone enantiomers and their bioconversion in sprouts. *Potato Research*. **1995**, *38*, 219-230.
- Yuan, L.; Wang, J.; Guan, Z.; Yue, F.; Wang, S.; Chen, Q.; Fu, M. Optimized preparation of methyl salicylate hydrogel and its inhibition effect on potato tuber sprouting. *Horticulturae*. **2022**, *8*(10), 866.
- Huang, T.; Ye, X.; Huang, X.; Li, X.; Mei, M.; Yu, L.; Wang, X. Effect of menthol and Jasmine essential oil on potato bud inhibition. *Journal of Sichuan Agricultural University*. **2018**, *(5)*, 618-625.
- Şanlı, A.; Karadoğan, T.; Tonguc, M.; Baydar, H. Effects of caraway (*Carum carvi* L.) seed on sprouting of potato (*Solanum tuberosum* L.) tubers under different temperature conditions. *Turkish Journal of Field Crops*. **2010**, *15*(1), 54-58.
- Oishi, A.; Nagatomi, Y.; Suzuki, K. Simultaneous LC-MS/MS Determination of 18 Plant Toxins in Beverages. *Food Hygiene and Safety Science*. **2019**, *60*(4), 108-112.
- GB 5009.3-2016. Determination of moisture in food. *Beijing: National standards for food safety*. **2016**.
- Wang, T.; Yan, T.; Shi, J.; Sun, Y.; Wang, Q.; Li, Q. The stability of cell structure and antioxidant enzymes are essential for fresh-cut potato browning. *Food Research International*. **2023**, *164*, 112449.
- Joly, N.; Souidi, K.; Depraetere, D.; Wils, D.; Martin, P. Potato by-products as a source of natural chlorogenic acids and phenolic compounds: extraction, characterization, and antioxidant capacity. *Molecules*. **2020**, *26*(1), 177.
- Palamutoğlu, R. Antibrowning effect of commercial and acid-heat coagulated whey on potatoes during refrigerated storage. *Journal of Food Science*. **2020**, *85*(11), 3858-3865.
- Hu, R.; Lin, L.; Liu, T.; Ouyang, P.; He, B.; Liu, S. Reducing sugar content in hemicellulose hydrolysate by DNS method: a revisit. *Journal of Biobased Materials and Bioenergy*. **2008**, *2*(2), 156-161.

20. Wegener, C. B.; Jansen, G.; Jurgens, H. U. Influence of drought and wounding stress on soluble phenols and proteins in potato tubers. *Sustainable Agriculture Research*. **2014**, 3(3), 1-15.
21. GB5009.124-2016. Determination of amino acids in food under National Food Safety standards. *Beijing: National standards for food safety*. **2016**.
22. Lee, S. K.; Jeon, J. S. Crucial role of inorganic pyrophosphate in integrating carbon metabolism from sucrose breakdown to starch synthesis in rice endosperm. *Plant Science*. **2020**, 298, 110572.
23. Khan, M.I.H.; Wellard, R.M.; Nagy, S.A.; Joardder, M.U.H.; Karim, M.A. Experimental investigation of bound and free water transport process during drying of hygroscopic food material. *International Journal of Thermal Sciences*. **2017**, 117, 266-273.
24. Xu, Y.; Wang, D.; Zhao, W.; Zheng, Y.; Wang, Y.; Wang, P.; Ma, Y.; Zhao, X. Low frequency ultrasound treatment enhances antibrowning effect of ascorbic acid in fresh-cut potato slices. *Food Chemistry*. **2022**, 380, 132190.
25. Hu, W.; Guan, Y.; Ji, Y.; Yang, X. Effect of cutting styles on quality, antioxidant activity, membrane lipid peroxidation, and browning in fresh-cut potatoes. *Food Bioscience*. **2021**, 44, 101435.
26. He, X.; Pu, Y.; Chen, L.; Jiang, H.; Xu, Y.; Cao, J.; Jiang, W. A comprehensive review of intelligent packaging for fruits and vegetables: Target responders, classification, applications, and future challenges. *Comprehensive Reviews in Food Science and Food Safety*. **2023**, 22(2), 842-881.
27. Basheer, S. M.; Chellappan, S.; Sabu, A. Enzymes in fruit and vegetable processing. *Value-Addition in Food Products and Processing Through Enzyme Technology*. **2022**, 101-110.
28. Nath, P.; Pandey, N.; Samota, M.; Sharma, K.; Kale, S.; Kannaujia, P.; Chauhan, O. P. Browning reactions in foods. *In Advances in Food Chemistry*. **2022**, 117-159.
29. Wang, L.; Wang, W. X.; Zeng, L.; Suo, H. C.; Li, C. C.; Shan, J. W.; Xiong, X. Y. Characteristics and differences of polyphenol oxidase, peroxidase activities and polyphenol content in different potato (*solanum tuberosum*) tubers. *Applied Ecology & Environmental Research*. **2020**, 18(6), 8171-8187.
30. Dobránszki, J.; Hidvégi, N.; Gulyás, A.; Tóth, B.; Teixeira da Silva, J. A. Transcription profile of potato (*Solanum tuberosum* L.) growing *in vitro*. *Journal of Plant Growth Regulation*. **2021**, 40, 749-760.
31. Cui, P.; Li, Y.; Cui, C.; Huo, Y.; Lu, G.; Yang, H. Proteomic and metabolic profile analysis of low-temperature storage responses in *Ipomoea batata* Lam. tuberous roots. *BMC Plant Biology*. **2020**, 20, 1-15.
32. Gikundi, E. N. Physico-chemical properties and storability of selected Irish potato varieties grown in Kenya (Doctoral dissertation, JKUAT-CoANRE). **2021**.
33. Ali, H. M.; El-Gizawy, A. M.; El-Bassiouny, R. E.; Saleh, M. A. The role of various amino acids in enzymatic browning process in potato tubers, and identifying the browning products. *Food Chemistry*. **2016**, 192, 879-885.
34. Sharma, N.; Sucheta, Dangi, S.; Yadav, S. K. Long-term storability of potato tubers in aspect of biochemical changes and overall quality index affected by different packaging materials in refrigerated and non-refrigerated storage. *Potato Research*. **2020**, 63(3), 303-321.
35. Aroiee, H.; Babaei, M.; Ajdanian, L.; Javdani, M.; Azizi, M.; Asgari Lajayer, B.; Dell, B. Effect of essential oil of seven medicinal plants on longevity, nongermination, qualitative and quantitative traits of *Solanum tuberosum* cv. Agria. *Journal of Food Processing and Preservation*. **2022**, 46(8), e16754.
36. Kasnak, C. Evaluation of the anti-browning effect of quercetin on cut potatoes during storage. *Food Packaging and Shelf Life*. **2022**, 31, 100816.
37. Hu, W.; Guan, Y.; Ji, Y.; Yang, X. Effect of cutting styles on quality, antioxidant activity, membrane lipid peroxidation, and browning in fresh-cut potatoes. *Food Bioscience*. **2021**, 44, 101435.
38. Bernal, J.; Mouzo, D.; López-Pedrouso, M.; Franco, D.; García, L.; Zapata, C. The major storage protein in potato tuber is mobilized by a mechanism dependent on its phosphorylation status. *International Journal of Molecular Sciences*. **2019**, 20(8), 1889.
39. Liu, P.; Xu, N.; Liu, R.; Liu, J.; Peng, Y.; Wang, Q. Exogenous proline treatment inhibiting enzymatic browning of fresh-cut potatoes during cold storage. *Postharvest Biology and Technology*. **2022**, 184, 111754.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.