

---

# High Burden of Non-Clonal Chromosome Aberrations Before Onset of Detectable Neoplasia in Fanconi Anemia Bone Marrow

---

Silvia Sánchez , [Benilde García-de-Teresa](#) , Marco A. Mejía-Barrera , [Pedro Reyes](#) , [Antonio Paz-Martínez](#) , Miguel A. Martínez , [Moisés O. Fiesco-Roa](#) , Angélica Monsiváis-Orozco , [Bertha Molina](#) , [Leda Torres](#) , Alfredo Rodríguez , [Sara Frias](#) \*

Posted Date: 17 March 2025

doi: 10.20944/preprints202503.1158.v1

Keywords: fanconi anemia; non clonal chromosome aberrations; clonal chromosome aberrations; complex karyotypes; cancer evolution; bone marrow failure; AML; MDS



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

# High Burden of Non-Clonal Chromosome Aberrations Before Onset of Detectable Neoplasia in Fanconi Anemia Bone Marrow

Silvia Sánchez <sup>1†</sup>, Benilde García-de-Teresa <sup>1†</sup>, Marco A. Mejía-Barrera <sup>1,2</sup>, Pedro Reyes-Jiménez, Antonio Paz-Martínez <sup>1,2</sup>, Miguel A. Martínez <sup>1,2</sup>, Moisés O. Fiesco-Roa <sup>1</sup>, Angélica Monsiváis-Orozco <sup>3</sup>, Bertha Molina <sup>1</sup>, Leda Torres <sup>1</sup>, Alfredo Rodríguez <sup>4,5</sup> and Sara Frias <sup>1,5,\*</sup>

<sup>1</sup> Laboratorio de Citogenética, Instituto Nacional de Pediatría, Insurgentes Sur 3700-C, CP04530, Ciudad de México, México

<sup>2</sup> Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Ciudad Universitaria, CP04510, Ciudad de México, México

<sup>3</sup> Servicio de Hematología, Instituto Nacional de Pediatría, Insurgentes Sur 3700-C, CP04530, Ciudad de México, México

<sup>4</sup> Laboratorio de Falla Medular y Carcinogénesis, Instituto Nacional de Pediatría, Insurgentes Sur 3700-C, CP04530, Ciudad de México, México

<sup>5</sup> Departamento de Medicina Genómica y Toxicología Ambiental, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Ciudad Universitaria, CP04510, Ciudad de México, México

\* Correspondence: sarafrias@iibiomedicas.unam.mx; Tel.: +52 550840900 ext. 1326, 1436

† Both authors contributing equally to this paper.

**Simple Summary:** Fanconi anemia (FA) is the most commonly inherited bone marrow failure syndrome, characterized by chromosomal instability and a high risk of cancer. Hematologic malignancy in FA, mainly myelodysplastic neoplasm (MDS) and acute myeloid leukemia (AML), associates with characteristic clonal chromosomal abnormalities in the bone marrow. Although clonal chromosome abnormalities (CCA) associated with malignant progression in FA involve chromosomes 1, 3 and 7, information on the broader preleukemic bone marrow cytogenetic landscape is scarce. In this study we report the type and frequency of every kind of non-clonal chromosomal abnormalities (NCCA) appearing in the bone marrow of a group of patients with FA, spanning from cancer-free patients to hematologic malignancy, where CCA appears. This study unveils the dynamism of emerging karyotypes in the FA bone marrow and its potential association with the patient's hematological state.

**Abstract:** Fanconi anemia (FA) is an inherited bone marrow failure syndrome characterized by chromosome instability and predisposition to develop myelodysplastic neoplasm (MDS) and acute myeloid leukemia (AML). Clonal chromosome aberrations (CCA) in chromosomes 1, 3, and 7 frequently appear in the bone marrow (BM) of patients with FA and associate to MDS/AML progression. Given the underlying DNA repair defect that characterizes FA, non-clonal chromosomal abnormalities (NCCA) are expected to be common events in the FA BM, however little is known about its presence and significance. Here, we transversally examined the BM karyotypes of 43 non-transplanted patients with FA, 41 with non-clinically detectable hematologic neoplasia and two with diagnosed MDS. We searched for the presence of NCCAs, complex karyotypes (CKs) and CCA, and their association with the natural history of the disease, including age, degree of BM failure, and neoplastic transformation. NCCAs were observed in the metaphase spreads of 41/43 FA patients; CKs were observed in 25/43 patients, whereas CCAs were found in 15/43 patients, CCA involving chromosomes 1, 3 and/or 7 in four patients, and other autosomes in the remaining 11 patients. Overall, we observed a baseline large karyotypic heterogeneity in the BM of FA patients, demonstrated by the ubiquitous presence of NCCA; such karyotypic heterogeneity

preceding the eventual emergence of CKs and selection of cells carrying fitness-improving CCAs. Finally, CCAs involving chromosomes 1, 3 and 7, well-known drivers of hematological malignancy in FA, become established. Overall, we observed that the frequency of NCCA and CCA increased with age, even though a significant correlation was not found. Preleukemic BM of patients with FA displays a baseline large karyotypic heterogeneity, evidenced by the ubiquitous presence of NCCA; such karyotypic heterogeneity preceding the eventual emergence of CKs and selection of cells carrying fitness-improving CCAs. Finally, CCAs involving chromosomes 1, 3 and 7, well-known drivers of hematological malignancy in FA, become established. These observations fit the model of evolution towards cancer that comprises a first phase of macroevolution represented by NCCA and karyotypic heterogeneity, followed by the establishment of clones with CCA, leading to microevolution and cancer.

**Keywords:** fanconi anemia; non clonal chromosome aberrations; clonal chromosome aberrations; complex karyotypes; cancer evolution; bone marrow failure; AML; MDS

---

## 1. Introduction

Fanconi anemia (FA) is a chromosome instability syndrome affecting 1 to 5 individuals per 1,000,000 individuals. FA is also the most frequently inherited bone marrow failure syndrome and originates from the inheritance of pathogenic variants (PV) in one of any 22 genes (FANCA-FANCW) that operate in the FA/BRCA pathway, responsible for recognizing and repairing DNA interstrand cross-links (ICL), through homologous recombination, an error-free pathway of DNA repair [1,2]. In absence of a functional FA/BRCA pathway, the ICLs can be repaired by low-fidelity DNA repair pathways, such as the non-homologous end joining (NHEJ) or the microhomology-mediated end joining (MMEJ) pathways [3,4] which rescue cell viability at the expense of increasing chromosome instability (CIN). The latter is a pathognomonic feature of FA, which allows its diagnosis due to the exquisite sensitivity of FA cells to ICLs inducing agents, like mitomycin C (MMC) and Diepoxybutane (DEB) [2,5–8].

The FA clinical phenotype is highly heterogeneous affecting multiple organs and systems. The phenotype includes, developmental abnormalities that can be detected since birth and are present in 80-96% of patients [9,10]; bone marrow (BM) failure, which appears in over 86% of patients with FA and consists of aplastic BM accompanied by peripheral blood cytopenias of different degrees mild, moderate or severe; patients with FA will develop BM failure around 7 years of age [10,11]. Also, FA patients are prone to develop cancer, both hematologic and solid tumors, with a cumulative incidence by 40 years of age of 30-33% and 20-28%, respectively [12,13]. The relative risk for a patient with FA to develop cancer, with respect to the general population is 19X and rises post HSC transplant to 55X; specifically, 527X for head and neck squamous cell carcinomas (HNSCC) and after transplant 933X; 582X for vulvar squamous cell carcinomas, after transplant 6298X; 213X for acute myeloid leukemia (AML), and 5669X for myelodysplastic neoplasia (MDS). The risk for hematological cancer, MDS and AML, increases by the age of 10 years old, plateauing at 20-30 years old [13,14].

BM failure in FA has been proposed to be the consequence of damage accumulation in the DNA of hematopoietic stem and progenitor cells (HSPCs) and hyperactivation of growth suppressing pathways, such as the TGF $\beta$  (Transforming growth factor  $\beta$ ) and the p53/p21 pathways, thus resulting in critical reductions in the number of HSPCs [15,16]. Despite this harsh BM microenvironment, certain HSPCs thrive, probably due to the inherent CIN in FA, and the acquisition of chromosome abnormalities that allow temporary survival, although not in the long-term.

CIN, one of the hallmarks of cancer, is characterized by aneuploidy and structural chromosomal abnormalities (CA), and can be observed in the early stages of cancer in the general population [17–19]. Importantly, CIN is a constant in FA patients, since zygote formation and embryonic development, it produces single cell genomic variation in the form of heterogeneous karyotypes and

has the potential to produce across the body of patients with FA, diverse altered genomes with short-term survival and elevated level of cell death, leading to the loss of somatic and HSPS cells, [20,21] however, this karyotypic heterogeneity and its inherent proapoptotic tendency, might also be the foundation for the well-known appearance of clonal hematopoiesis and eventual progression to cancer in FA [21,22]. The latter might be the reason why up to 40% of children and young patients with FA show signs of clonal evolution in their BM, and 15 to 60%, will develop MDS or leukemia, mainly AML, at an early age [2,23].

Multiple studies have described, and confirmed, the involvement of clonal chromosome abnormalities (CCA) in the emergence of MDS or AML in patients with FA [22–27]. Most of these studies screened for common CCA using microarrays or fluorescent in situ hybridization (FISH) directed to regions of interest in chromosomes 1q, 3q and 7q, since these are the most informative cytogenetic markers of clonal evolution towards MDS/AML [23,28,29]. Cytogenetic analysis, however constantly misses or discards non-clonal chromosome aberrations (NCCA), as they are considered "background noise", specially in patients with CIN, as is the case for FA [23].

Cancer evolution, however, is known to be a dynamic process, directed by large-scale chromosomal copy number variations that create karyotypic heterogeneity between the cells that compose a tissue, privileging the selection of the genomes with the greatest fitness [17], eventually leading to the selection of clones with CCA that will orchestrate cancer development and evolution [17,30]. FA inherent CIN is expected to create large levels of NCCA, generating a BM with high karyotypic heterogeneity years before full-blown MDS or AML. In this sense, NCCA could precede the formation of complex genomes and stable CCAs eventually transitioning to cancer.

The aim of this study was to cytogenetically analyze the BM of a group of non-transplanted patients with FA, to investigate whether the multiple steps of karyotypic evolution towards cancer can be appreciated, in relation to the age of the patients and their hematological condition. We did not find an association between the complexity of the BM karyotype and the age of the patients or their hematological condition, suggesting that age and stage of bone marrow failure are not the only drivers of leukemogenesis in FA. However, inspecting the group of FA patients as a whole, revealed transitions from a karyotypically diverse population of BM cells with NCCA, to the presence of complex karyotypes, and emergence of CCA involving CA of indeterminate potential in patients with FA [29] and CCA with the classic high risk chromosome alterations found in chromosomes 1, 3, and 7; the only two patients in the group who were identified with MDS presented with the classic alterations in chromosomes 3 and 7.

## 2. Materials and Methods

### 2.1. Patients

We performed a transversal study including non-transplanted patients with a confirmed diagnosis of FA with the DEB chromosome breakage test. The samples analyzed corresponded to the first cytogenetic BM evaluation of each patient within the framework of the recommended annual follow-up in search for MDS/AML progression markers. All samples were processed at the Cytogenetics Laboratory of the National Institute of Pediatrics (Mexico) between 2015 and 2024. In accordance with the Declaration of Helsinki, all patients or their relatives gave written informed consent for genetic analyses under project number INP 2014/041 and 2020/012; clinical and demographic data were obtained from the patient's medical records and from interviews conducted in the frame of the Fanconi Anemia Registry from Mexico (RAFMex) project INP 2020/053. These projects were approved by the institutional research and ethics committees at Instituto Nacional de Pediatría (México).

Age at diagnosis was calculated considering the date of positive DEB test. BMF severity at the time of cytogenetic study was classified according to Fanconi anemia guidelines for diagnosis and management [31], using the hemogram data (free of transfusion) nearest to the cytogenetic study. The use of androgens was recorded irrespective of the length of treatment or if it was being taken at

the time of sample collection. Transfusion dependency was considered when the patient could not maintain adequate blood counts without sequential transfusions at any timepoint.

## 2.2. Cytogenetics

Conventional karyotypes were performed on BM cells using the standard direct method. Briefly, 1-2 mL of heparinized BM samples were incubated in a 5% CO<sub>2</sub> incubator, cultured during 48 h in MarrowMax medium without cell division stimulant at 37°C, after which colcemid [10 µg/mL] was added for the final 2 h of culture. Harvesting, slides preparation and GTG banding were made according to the classic methodology [32]. On average, 20 metaphase spreads were analyzed per patient and reported according to the International System for Human Cytogenetic Nomenclature (ISCN) [33], less than 18 metaphases were analyzed in four patients with severe aplastic anemia due to hypocellular BM.

Chromosome abnormalities were classified as: a) NCCA, non-clonal chromosomal abnormality, both numerical and structural; b) Complex karyotype (CK), defined in this study as a cell with  $\geq 3$  independent cytogenetic abnormalities [34]; c) CCA, clonal chromosomal abnormalities, when at least two cells bear the same chromosome gains or structural rearrangements (including deletions and duplications), or when at least three cells present with the same whole chromosome loss [33]. CCA 1,3,7 refers to clonal abnormalities involving at least one of the known recurrent alterations: i.e. duplication of the long arm of chromosome 1 (commonly known as 1q+), duplication of the long arm of chromosome 3 (commonly known as 3q+), complete monosomy of chromosome 7 or deletion of the long arm of chromosome 7 (commonly known as -7/7q-), while CCA refers to clonal chromosome alterations in chromosomes other than 1q+,3q+ and -7/7q-. When aneuploidies of a whole chromosome were detected at a given metaphase, and to reduce potential technical errors, the periphery of such metaphase was analyzed with a 10x objective to increase the visibility margin by 10 fields and to detect whether the lost chromosome was located at the periphery of the metaphase, or whether the gained chromosome belonged to a nearby metaphase. Chromosome breaks and radial exchange figures were recorded during the screening but are not part of the analysis in this report.

## 2.3. Statistics

All statistical analyses were performed in Prism (version 10.4.0). Normal distribution of the data was assessed with the Shapiro Wilk test. Data with normal distribution were compared using one-way ANOVA with Tukey's post-test for multiple comparisons. Comparisons between two groups were performed with the unpaired t test. The Fisher's exact test was used to compare proportions between two groups. Pearson correlation test was used to probe correlation between age and frequency of chromosome abnormalities.

## 3. Results

### 3.1. Patients

Samples from 50 non transplanted patients with a positive diagnosis of FA, were sent to the cytogenetics Laboratory at Instituto Nacional de Pediatría (Mexico) for routine annual follow-up between 2015 and 2024; six of them had insufficient material to perform cytogenetic analysis, one patient was excluded since the provided sample was obtained once antineoplastic treatment for AML was started. Appropriate cells, in quality and number, for cytogenetic analysis were obtained from 43 patients (Table S1); in two of them (FANC031 and RAFMex015) the concurrence of cytopenia and a monosomy of chromosome 7 clone allowed the diagnosis of MDS in accordance with International Consensus Classification criteria (CITAR PMID: 35767897). During the duration of this study, some of the patients had more than one cytogenetic analysis on more than one sample, only the first cytogenetic study was considered for the purposes of this report.

Median age at diagnosis was 8.1 (1.4-15.6) years old, and median age at the cytogenetic study was 9.14 (4.7-28.6) years old, with a female-male relationship 1.4:1. A summary of the population

demographics is shown in **Table 1**, and main clinical data are shown in **Table 2**. Eleven patients died during the elapsed time between cytogenetic study and this report: due to BM failure or HSCT-related complications in eight patients, MDS in two patients and acute lymphoblastic leukemia (ALL) in the remaining one.

**Table 1. Clinical characteristics of study population**

<b>Gender and age of patients (n=43)</b>	<b>n</b>	<b>Percentage % or (range)</b>
Female	25	58
Male	18	42
Median age at diagnosis years	8.1	(1.4-15.6)
Median age (years) at cytogenetic study	9.19	(4.7-28.6)
<b>Affected gene (n=28)</b>		
<i>FANCA</i>	19	67.8
<i>FANCE</i>	2	7.1
<i>FANCD2/BRCA1</i>	1	3.6
<i>FANCF</i>	1	3.6
<i>FANCG</i>	2	7.1
<i>FANCI/BRIP1</i>	1	3.6
<i>FANCL</i>	2	7.1
<b>BMF Status (n=43)</b>		
No	5	11.6
Mild	8	18.6
Moderate	15	34.9
Severe	13	30.2
MDS	2	4.7
<b>Androgen treatment (n=42)</b>		
Yes	29	69
No	13	31
<b>Transfusion dependency (n=42)</b>		
Yes	13	31
No	29	69
<b>Transplant (n=42)</b>		
Yes	8	19
No	34	81
<b>Alive/Deceased (Cause of death) (n=41)</b>		
Alive	30	73.2
Deceased (ALL)	1	2.4
Deceased (BMF)	4	9.8
Deceased (BMF, Intracranial bleeding)	1	2.4
Deceased (HSPCT complication)	4	9.8
MDS	1	2.4

BMF-Bone Marrow Failure. MDS-Myelodysplastic Neoplasm.

ALL- Acute Lymphoid Leukemia.

HSPCT-hematopoietic stem and progenitor cells transplantation.


Table 2. Population and clinical data.

ID	Gender	Age at diagnosis (years)	Age at cytogenetic study (years)	Affected gene	BMF Status	Androgen treatment	Transfusion dependency	HSPCT	Alive/Deceased (Cause of death)
RAFMex057	Male	4.2	4.7	FANCE	Moderate	No	No	No	Alive
RAFMex058	Female	4.1	4.8	NI	Mild	No	No	No	Alive
RAFMex006	Male	3.5	5.5	FANCF	Severe	Yes	No	Yes	Alive
RAFMex029	Male	4.5	5.5	NI	Mild	Yes	No	No	Alive
RAFMex009	Female	3.3	6.2	FANCA	Moderate	Yes	No	No	Alive
RAFMex043	Male	4.9	6.6	FANCE	Severe	Yes	Yes	Yes	Deceased (HSPCT complication)
RAFMex010	Female	5.9	6.7	FANCA	Moderate	Yes	No	No	Alive
RAFMex042	Female	6.9	6.9	FANCL	Severe	Yes	Yes	Yes	Alive
RAFMex047	Female	7.1	6.9	FANCA	Severe	Yes	No	No	Alive
RAFMex002	Female	6.0	7.8	FANCA	No	No	No	No	Alive
RAFMex044	Female	1.4	7.9	FANCA	Mild	Yes	No	No	Alive
RAFMex033	Male	5.1	8.1	NI	Moderate	No	No	No	Alive
RAFMex025	Female	6.1	8.1	FANCA	No	No	No	No	Alive
RAFMex005	Female	6.1	8.2	FANCA	Moderate	Yes	No	No	Alive
FANC143	Female	3.6	8.3	FANCG	Severe	Unclear	Yes	Yes	Deceased (HSPCT complication)
RAFMex026	Female	8.5	8.5	FANCA	Severe	No	No	Yes	Deceased (HSPCT complication)
RAFMex011	Female	7.1	8.7	FANCA	Moderate	No	No	Yes	Alive
FANC161	Female	8.0	8.8	NI	Severe	Yes	Yes	NI	Deceased (BMF)
FANC157	Male	9.0	9.0	NI	Severe	Yes	Yes	No	Deceased (BMF)
FANC162	Male	9.1	9.12	NI	Severe	No	NI	No	NI
FANC148	Male	7.7	9.14	NI	Severe	No	Yes	No	Deceased (BMF, Intracranial bleeding)
FANC101	Female	8.2	9.2	FANCA	Severe	Yes	Yes	Yes	Deceased (HSPCT complication)
RAFMex07	Female	9.7	9.7	FANCJ/BRIP1	Moderate	No	No	No	Alive
RAFMex035	Male	9.7	10.2	FANCA	Mild	No	No	No	Alive
FANC032	Male	9.2	10.7	FANCG	Moderate	Yes	No	No	NI
RAFMex024	Female	8.8	10.9	FANCA	Moderate	Yes	Yes	No	Alive
RAFMex08	Male	9.5	11.7	FANCA	Mild	Yes	No	No	Deceased (ALL)
RAFMex032	Female	12.6	12.6	NI	Moderate	No	No	No	Alive
RAFMex031	Male	13.1	13.4	FANCA	Severe	Yes	Yes	No	Deceased (BMF)
RAFMex018	Female	9.8	14.7	FANCA	Severe	Yes	Yes	No	Alive
FANC024	Female	7.4	16.7	FANCA	Moderate	Yes	No	Yes	Alive
RAFMex051	Female	11.7	17.7	NI	Moderate	Yes	Yes	No	Alive
RAFMex036	Male	6.4	17.8	NI	Moderate	Yes	No	No	Alive
RAFMEX066	Male	9.5	17.9	FANCL	No	Yes	No	No	Alive
RAFMex038	Male	12.7	18.4	NI	Mild	Yes	No	No	Alive
RAFMex037	Female	14.4	20.2	NI	No	Yes	No	No	Alive
RAFMex059	Male	13.3	22.50	NI	Moderate	Yes	No	No	Alive
RAFMex027	Male	5.6	22.51	NI	No	No	No	No	Alive
RAFMex022	Female	15.6	25.0	FANCA	Mild	Yes	Yes	No	Alive
RAFMex034	Female	11.7	25.7	FANCA	Mild	Yes	No	No	Alive
RAFMex030	Female	12.0	28.6	NI	Moderate	Yes	No	No	Alive
FANC031	Female	10.3	15.5	FANCD2	MDS	Yes	Yes	No	Deceased (MDS)
RAFMex015	Male	9.7	17.2	FANCA	MDS	Yes	Yes	No	Deceased (MDS)

BMF-Bone Marrow Failure. HSPCT-hematopoietic stem and progenitor cells transplantation. ALL- Acute Lymphoid Leukemia. MDS- Myelodysplastic Neoplasm. NI-No Information.

### 3.2. BM Cytogenetics of Patients with FA

As expected for patients with FA, their BM showed enormous karyotypic heterogeneity, evidencing the underlying CIN. The cytogenetic analysis in the 43 included patients, showed that all but one patient displayed chromosomal abnormalities, that were classified in three types, NCCA, non recurrent CK, and CCA. Representative karyotypes per CA type are shown in **Figure 1**.



**Figure 1. Representative karyotypes showing the various types of chromosomal abnormalities found in the BM from patients with FA.** NCCA: Non-Clonal Chromosome Aberrations; CK:Complex Karyotype; CCA=Clonal Chromosome Aberrations. Red arrowheads indicate deletions; blue arrowheads indicate duplications.

NCCA were observed in 41/43 patients; karyotypes were NCCA were the only findings occurred in 26/43 patients, the presence of only CCA was observed in a single patient; and the combination of both NCCA and CCA were observed in 15/43 patients. In addition, 25 patients presented with non recurrent CK. Among the 15 patients with CCA, eight had clones with aneuploidy for one complete chromosome, six had clones with structural chromosomal aberrations and one patient had clones with both types of chromosomal aberration (**Figure 2**). Chromosome losses were more commonly observed in comparison to chromosome gains. The chromosomes more frequently involved in any kind of CA were chromosomes 3, 7, 8 and 18, followed by chromosomes 6, 16 12, 17, 20 and 22; also, we found additional material (add), marker chromosomes (mar) and neutral aberrations like inversions and balanced translocations (**Figure 3 and Table 3**).

### 3.2.1. Non Clonal Chromosomal Abnormalities

Patients with inherent CIN, such as FA, are expected to have enormous karyotypic heterogeneity, this was evidenced in their BM where stochastic chromosomal alterations were found (**Table 3 and Figure 2**). All 24 human chromosomes were involved in these CA, numerical aberrations were the most common (**Figure 3**).




Figure 2. Frequency of chromosomal aberrations found in BM of FA patients ordered by age and MDS diagnosis at cytogenetic study. Distribution of the frequencies of the different type of chromosomal aberrations, per patient, in ascending order of age. The age is indicated between parenthesis.


Table 3. Cytogenetic findings in the BM of 43 patients with FA.

n	ID	Total frequency of CA	Frequency of NCCA/cell (Frequency of cells with CK)	Frequency of CCA/cell	Involved chromosome in CCA
1	RAFMEK057	0.8	0.55 (0.10)	0.25	del(3q27)
2	RAFMEK058	0.26	0.26 (0.04)	0.00	
3	RAFMEK006	0.75	0.75 (0.04)	0.00	
4	RAFMEK029	0.65	0.65 (0.05)	0.00	
5	RAFMEK009	0.45	0.45 (0.00)	0.00	
6	RAFMEK043	0.54	0.33 (0.03)	0.21	del(6p22)
7	RAFMEK010	0.50	0.30 (0.05)	0.20	-21
8	RAFMEK042	0.36	0.36 (0.00)	0.00	
9	RAFMEK047	0.00	0.00 (0.00)	0.00	
10	RAFMEK002	0.23	0.00 (0.00)	0.23	add(2p)
11	RAFMEK044	0.50	0.50 (0.05)	0.00	
12	RAFMEK033	0.90	0.90 (0.21)	0.00	
13	RAFMEK025	0.85	0.85 (0.10)	0.00	
14	RAFMEK005	0.90	0.75 (0.10)	0.15	-20
15	FANCI43	0.60	0.60 (0.10)	0.00	
16	RAFMEK026	0.95	0.63 (0.11)	0.32	dup(3q26)
17	RAFMEK011	0.40	0.40 (0.00)	0.00	
18	FANCI61	0.35	0.35 (0.05)	0.00	
19	FANCI57	0.10	0.10 (0.00)	0.00	
20	FANCI62	0.10	0.10 (0.00)	0.00	
21	FANCI48	1.25	0.55 (0.15)	0.70	+8
22	FANCI01	0.40	0.40 (0.00)	0.00	
23	RAFMEK007	0.85	0.85 (0.10)	0.00	
24	RAFMEK035	0.25	0.25 (0.00)	0.00	
25	FANCI32	0.30	0.30 (0.00)	0.00	
26	RAFMEK024	0.15	0.15 (0.00)	0.00	
27	RAFMEK008	0.63	0.63 (0.05)	0.00	
28	RAFMEK032	0.70	0.40 (0.10)	0.30	-17, -22
29	RAFMEK031	1.20	1.20 (0.15)	0.00	
30	RAFMEK018	0.05	0.05 (0.00)	0.00	
31	FANCI24	0.40	0.25 (0.05)	0.15	-18
32	RAFMEK051	0.22	0.22 (0.00)	0.00	
33	RAFMEK036	0.68	0.68 (0.05)	0.00	
34	FANCI46	1.13	0.93 (0.07)	0.20	-16
35	RAFMEK038	0.81	0.67 (0.00)	0.14	add(3p25)
36	RAFMEK037	0.99	0.99 (0.06)	0.00	
37	RAFMEK059	0.75	0.50 (0.15)	0.25	der(6)t(1;6)
38	RAFMEK027	0.70	0.45 (0.10)	0.25	del(3p13)
39	RAFMEK022	0.30	0.30 (0.00)	0.00	
40	RAFMEK034	0.25	0.25 (0.00)	0.00	
41	RAFMEK030	0.35	0.35 (0.05)	0.00	
42	FANCI31	1.05	0.45 (0.1)	0.60	-7, -8
43	RAFMEK015	3.40	0.40 (1.00)	3.00	-7 der(18)t(3;18)
	SUM (percentage)	27 (100%)	20.05 (74.3%)	6.95 (25.7%)	
	Average	0.63	0.47 (0.07)	0.16	

Abbreviations: CA- Chromosome Abnormalities. NCCA- No Clonal Chromosome Abnormalities. CK- Complex Karyotype. CCA-Clonal Chromosome Abnormalities.

NCCA were observed in the vast majority of FA patients, 95% (41/43), including non recurrent CK. NCCA, without evidence of any additional chromosomal clones were found in 26 patients. NCCA were the most prevalent type of CA, accounting for 74.6% of the total abnormalities observed, and included CK in 25/43 patients, which most of them were non-recurrent. Patient RAFMex015, was exceptional, since clonal and complex karyotypes were found in 100% of the analyzed cells, concurrent with an MDS diagnosis (Table 3 and Figure 2).

In most patients, NCCA were the most frequent CA, these were highly diverse and included numerical and structural alterations. Among the structural CA, 11/78 were neutral in that they did not appear to condition a loss or gain of chromosome information; most were inversions but there was also one translocation (Figure 3, green cells).



**Figure 3. NCCA and CCA abnormalities in 43 FA patients.** In the upper panel, patients are ordered by age and MDS diagnosis. The rightmost column signals which chromosomes are involved in the CA. Complete aneuploidy, including loss and gain were observed for all 24 chromosomes. Structural aberrations, depicted in the lower section, involved all chromosomes except 16,19,20,21 and Y. Marker chromosomes refers to non-


identified abnormal chromosomes and add refers to additional chromosomal material of unknown origin. Neutral aberrations refers to inversions and balanced translocations. Patient RAFMex057 showed two structural alterations on chromosome 8, one deletion and one inversion, indicated by .

### 3.2.2. Clonal Chromosomal Abnormalities (CCA)

CCA accounted for 25.4% of the total CA found in the 43 patients with FA (**Table 3**), they were found in 35% (15/43) among these the presence of CCA involving monosomy of chromosome 7 in two of these patients led to a MDS diagnosis. Within the 15 patients which presented with clonal chromosome abnormalities (WC), seven presented clones consisting of whole-chromosome aneuploidy, losses involving chromosomes 7, 8, 16, 17, 18, 20, 21 and 22; gains involving chromosome 8, and/or a marker chromosome. Seven patients had clones with structural CA in chromosomes 1, 3, 6 and 18, and one patient presented a clone with both numerical (chromosomes 7 and +mar) and structural (chromosomes 3 and 18) CA. Among 15 patients WC, 11 presented CCA involving autosomes other than 1,3 and 7, and in 4/15 patients WC, the clone involved the regions on chromosomes 1, 3 and 7 associated with the evolution towards hematological malignancy; one of them had a duplication 1q21-qter, due to a translocation t(1;6)(q21;p21.3). Patients WC, in addition to having cytogenetic clones, revealed a variety of NCCA: 3/15 had whole-chromosome aneuploidies, 11 had numerical and structural NCCA, and only one patient did not present NCCA (**Figure 3 and Table 3**).

### 3.3. Clonicity and Chromosomal Damage


In this group of patients, we found that the average frequency of CCA (0.16) is only ~20% of the observed average frequency of NCCA (0.47) (**Table 3**), in addition, it is evident that when the total frequency of ab/cell is higher, CCA appears in a patient (**Figure 4**). In patients WC who have MDS, a large number of CA are observed in their BM, because the fittest clone is present in a large proportion (**Figures 1 and 4**). In patients WC, CCA alterations coexist with NCCA, although the latter are generally the most common alteration. However, it should be noted that in the two patients who already present MDS, CCA prevail over NCCA (**Figure 2**).



**Figure 4. Frequency of chromosomal aberrations per cell.** The population was divided in patients without clones (blue box), with clones (red box), and patients with -7/ MDS (myelodysplastic neoplasm). Statistical analyses were performed using one-way ANOVA with Tukey's post-test for multiple comparisons.

### 3.4. Complex Karyotypes and Clonicity

Several analyzed metaphases had more than one CA, we classified the patients that had at least one metaphase that had three or more CA as having a CK. Twenty five out of 43 patients had metaphases with these CK, which represent a greater complexity within the population of cells with NCCA. When comparing the frequency of CK in patients with and without clones, there is a higher proportion of patients with CK among those who have already developed clones, showcasing an increased complexity of the diverse karyotypes found in their BM (**Figure 5**).




**Figure 5. Patients with complex karyotypes (CK), according to clonicity.** In yellow, the proportion of patients who present at least one metaphase with CK ( $\geq 3$  CA). Patients with MDS were excluded.


### 3.5. Chromosomal Damage and Patient Demographic and Clinical Characteristics

#### 3.5.1. Chromosomal Abnormalities and the Age and Gender of Patients

Pearson correlation analysis revealed that as the patients grow older the total frequency of CA does not increase, ( $R=0.056$ ,  $p=0.73$ ) (**Figure 6a**). However, when patients were sub-divided into three groups of age: 4 -9 years old, 10-15 years old and >16 years old, a significantly higher frequency of CA was observed between the older patients WC (excluding MDS patients) and the younger patients without clones (one-way ANOVA with Tukey's post-test) (**Figure 6b**). Finally, There were no significant differences in AC frequency when comparing males and females. (**Figure 7**).




**Figure 6. Correlation between the total frequency of CA and the patients' age.** a) No correlation was found between the amount of CA and increasing age of patients WC and without clones (Pearson  $R=0.056$ ,  $p=0.73$ ). b) A significant increase in total CA was found only between patients 4-9 years old without clones and patients >16 years with clones.



**Figure 7.** No differences in the total frequency of CA were found between females vs males.

Finally, we search for an association between the frequency of chromosomal aberrations and the bone marrow failure severity: 5/43 did not have BMF, 28/43 had moderate or severe BMF. As a group, the patients with WC show slightly more CA than WOC, and although no statistical differences were found it would appear that in individuals WC, CA would increase as the BMF becomes more severe. (Figure 8a). Other parameters regarding to the hematological status, including transfusion dependency and androgen treatment, did not influence the total frequency of CA observed in patients with FA. (Figure 8b and 8c).



**Figure 8. Frequency of CA with respect to hematologic status.** a) The frequency of CA in FA does not correlate with the severity of bone marrow failure, determined in accordance with the FA clinical care guidelines [31]. b) CA per cell with respect to transfusion dependency. c) CA per cell in reference to androgen therapy. No differences in the frequency of CA were observed with respect to hematological conditions or patients' treatment.

#### 4. Discussion

In this study, we conducted a cytogenetic cross-sectional analysis of the BM of 43 patients with the chromosome instability and cancer predisposition syndrome FA. As anticipated, a high number of CA were observed in virtually all participants. NCCA were the most prevalent, reflecting an underlying karyotype heterogeneity caused by faulty DNA repair mechanisms in individuals with FA. CCAs, mostly abnormalities of indeterminate potential, were also observed, with high-risk CAs being detected in four patients, two of whom had MDS defining CAs. This comprehensive analysis of the type and frequency of CA in the BM of individuals with FA, stresses out the natural history of hematological cancer in FA, from heterogeneous NCCA towards CK, and finally CCA and cancer. This study complements the information on the genomic changes involved in the evolution towards hematological cancer, which has been extensively studied by other authors [21,22,29], given that we analyzed the chromosomes of pre-leukemic BM and at the individual cell level, revealing the presence of NCCA in addition to CCA.

As shown in **Figure 3**, the observed NCCAs included both structural alterations and aneuploidies (gains and losses of whole chromosomes). Structural alterations included gains of chromosome material and chromosome deletions. Specifically, chromosome gains may arise from the joining of non-homologous chromosome segments through the non-homologous end-joining (NHEJ) pathway, causing chromosome translocations. Additionally, the segregation of radial exchange figures during mitosis, could produce daughter cells harboring single derivative chromosomes (with the corresponding gain and/or loss of chromosome material). On the other hand, chromosome deletions, on the other hand, were six times more frequent than gains. Their origin could be chromosomal rearrangements, or non-repaired double-strand breaks (DSB) that reached mitosis, and whose broken segment was not retained in the same cell after exiting mitosis, leaving a deleted chromosome.

We saw an overrepresentation of loss-of-chromosome aneuploidies, raising the possibility of technical errors during the preparation of metaphase spreads, nonetheless given our strict scoring criteria during chromosome analysis (see methodology), we consider that most of these alterations are genuine chromosome numerical losses. In this context of genomic instability, we hypothesize that

chromosome losses are related to the mis-segregation of structurally altered chromosomes that became incorporated into micronuclei and further shattered, as others have shown to occur with lagged chromosomes [35].

Interestingly, we also identified neutral CA, i.e. balanced CA such as chromosome inversions and translocations. Traditionally, these alterations were thought to be less detrimental than deletions and duplications, but it is now recognized that they can have a significant impact in the cancer evolution, since the change in the 3D distribution pattern of genes and regulatory elements addresses is responsible for emerging network dynamics [36]. Inversions are not commonly reported in previous cytogenetic studies of patients with FA, most likely because cytogenomic analysis using microarrays is usually preferred over GTG banded chromosomes. are not typically analyzed [22]. These inversions could result from the presence of two DSBs in the same chromosome that were rejoined after a 180-degree rotation. Other mechanisms may also explain the presence of inversions, including chromothripsis-like rearrangements, but with fewer breakpoints than traditional chromothripsis, which usually involves extensive fragmentation and rearrangement of a single chromosome. It is known that an intact FA/BRCA pathway plays a crucial role in the micronucleus-related chromosome fragmentation that leads to chromothripsis, while a deficient FA/BRCA pathway generates significantly less chromosome fragmentation under similar conditions [35]. This type of alterations evidenced here by G banding, may represent just the tip of the iceberg of more complex ACs, which deserve further study.

Of note, NCCAs in FA are quite variable from cell to cell, and this is probably due to multiple factors, including: first, a continuous generation of DSBs, that are incorrectly and not homogeneously rejoined due to the reliance of the FA cells on error-prone DNA repair pathways [4]. Second, dividing cells harboring NCCAs will not parent chromosomally identical daughter cells, particularly when dicentric chromosomes and radial figures are present, as mitotic segregation will not equally distribute the CAs into the daughter cells [2]. Third, NCCA might confer differential surviving capacities, therefore alterations incompatible with life will not thrive, while fitness providing CAs will persist, in a constant cycle of cell replacement and cell attrition. During this cycle of depletion and emergence of cells with NCCAs, new genome compositions in the tissue will arise, producing a unique combination of CAs that will benefit the survival of cells harboring stable karyotypes with CCA.

Cells carrying a specific CCA can be considered the evidence of a successful genome arrangement. However, this specific clone could also be transient, as the continuous production of CAs does not cease, leading to the emergence of CK, most of which are NCCA. Some CK may appear in cells that already contain a specific CCA, generating genomes that may either disappear or adapt, eventually evolving towards a clone with higher fitness. In a cell population exhibiting CIN, NCCAs will create dynamic karyotypic diversity. This heterogeneity of cells, with stochastic CA, eventually gives rise to CCA, which may be transient, creating a dynamic cycle of NCCA/CCA. The NCCA/CCA cycle establishes a macro evolutive phase that may continue until emergency of stable clones that encompasses a phase transition. One surviving genome with CCA carrying high-risk CAs emerge, such as clones with dup(1q), dup(3q), del(7q), and -7 and the accompanying NCCAs decrease (as in patients with MDS). Clonal expansion is characteristic for the microevolutionary phase [37]. These specific karyotype alterations are known to facilitate the progression to cancer. The cells with these Cas will proliferate until they become a clone, thus completing the transition from NCCA to CCA. In patients with FA, the transition to cancer also involves specific genetic alterations such as RUNX1 and TP53, among others [22].

Importantly, progression towards cancer, which typically takes a long time in DNA repair proficient individuals, is accelerated in patients with FA. In this work, through an in-depth cytogenetic analysis of the BM of patients with FA, we recapitulate macroevolution patterns previously described [23,30,31], including an initial phase of high karyotypic heterogeneity leading to more stable "end products of evolution", i.e. CCA abnormalities that combine large-scale

chromosomal changes with genetic mutations and copy number alterations, that will ultimately drive micro-evolution steps towards cancer.

## 5. Conclusions

In this study, we observed that the frequency of ACs was not associated with gender, bone marrow failure, or treatment. The frequency of NCCA and CCA increased with age; although no significant correlation was found, a significant difference was observed between older patients with CCA and younger patients without CCA.

In general, the preleukemic bone marrow of patients with FA exhibit significant basal karyotypic heterogeneity, evidenced by the widespread presence of NCCA. This karyotypic heterogeneity precedes the eventual appearance of CK and the selection of CCA-bearing cells that enhance adaptation, which may be transient until the appearance of a stable CCA. The NCCA/CCA cycle establishes a macroevolutionary phase that could continue until the emergence of stable clones, leading to a phase transition. A surviving CCA genome harboring high-risk ACs emerges, such as clones with dup(1q), dup(3q), del(7q), and -7, and the associated NCCAs decline. These observations fit the model of evolution towards cancer that comprises a first phase of macroevolution represented by NCCA and karyotypic heterogeneity, followed by the establishment of clones with CCA, which leads to microevolution and cancer.

These observations warrant a longitudinal follow-up study of patients with FA to determine the macro- and microevolutionary phases and to detect potential cytogenetic biomarkers that precede clonal hematopoiesis.

**Supplementary Materials:** The following supporting information can be downloaded at: Preprints.org, Table S1. Data Base containing the Cytogenetics of 43 patients with FA. Complete data.

**Author Contributions:** Conceptualization, S.F.; S.S.; B.G.; methodology, S.F.; S.S.; B.G.; M.O.F.; A.R.; investigation, S.F.; S.S.; B.G.; P.R.; A.P.; M.A.M.; B.M. resources, B.G.; A.M.; M.O.F.; B.M.; U.J.; L.T.; data curation, S.S.; B.G. writing—original draft preparation, S.F.; S.S.; B.G.; writing—review and editing, S.F.; S.S.; B.G.; A.R.; visualization, S.S.; B.G.; B.M.; P.R.; A.R.; supervision, S.F.; S.S.; B.M.; U.J.; L.T.; B.G.; A.M.; project administration, S.F.; B.M.; funding acquisition, S.F.; B.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Dirección General de Asuntos del Personal Académico, Universidad Nacional Autónoma de México, UNAM-PAPIIT, grant number IN205120; Consejo Nacional de Humanidades, Ciencias y Tecnologías, Grant number CF-2023-G-800; Instituto Nacional de Pediatría, Recursos Fiscales, Projects 2020/012 and 2020/053. The APC was funded by XXX”.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Ethics Committee of Instituto Nacional de Pediatría (protocol code 2014/041 approved on July 14<sup>th</sup>, 2014, 2020/053 and 2020/012 approved on March 23<sup>rd</sup>, 2020).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The original contributions presented in this study are included in the article/supplementary material. Further inquiries can be directed to the corresponding author(s).

**Conflicts of Interest:** The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## References

1. Rodríguez, A.; D’Andrea, A. Fanconi anemia pathway. *Curr. Biol.* 2017.
2. García-De-teresa, B.; Rodríguez, A.; Frias, S. Chromosome instability in fanconi anemia: From breaks to

- phenotypic consequences. *Genes (Basel)*. **2020**, *11*, 1–35.
3. Ayala-Zambrano, C.; Yuste, M.; Frías, S.; García de Teresa, B.; Mendoza, L.; Azpeitia, E.; Rodríguez, A.; Torres, L. A Boolean network model of the double-strand break repair pathway choice. *J. Theor. Biol.* **2023**, *573*.
  4. Rogers, C.B.; Kram, R.E.; Lin, K.; Myers, C.L.; Sobek, A. dependent on POL  $\theta$  -mediated alternative end joining. **2023**, *42*.
  5. Auerbach, A.D. Fanconi anemia and its diagnosis. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* **2009**, *668*, 4–10.
  6. Moreno, O.M.; Paredes, A.C.; Suarez-Obando, F.; Rojas, A. An update on Fanconi anemia: Clinical, cytogenetic and molecular approaches (review). *Biomed. Reports* **2021**, *15*, 1–10.
  7. Esmer, C.; Sánchez, S.; Ramos, S.; Molina, B.; Frias, S.; Carnevale, A. DEB Test for Fanconi Anemia Detection in Patients with Atypical Phenotypes. *Am. J. Med. Genet.* **2004**.
  8. Merfort, L.W.; Lisboa, M. de O.; Cavalli, L.R.; Bonfim, C.M.S. Cytogenetics in Fanconi Anemia: The Importance of Follow-Up and the Search for New Biomarkers of Genomic Instability. *Int. J. Mol. Sci.* **2022**, *23*.
  9. Fiesco-Roa, M.O.; Giri, N.; McReynolds, L.J.; Best, A.F.; Alter, B.P. Genotype-phenotype associations in Fanconi anemia: A literature review. *Blood Rev.* 2019.
  10. Altintas, B.; Giri, N.; McReynolds, L.J.; Best, A.; Alter, B.P. Genotype-phenotype and outcome associations in patients with Fanconi anemia: the National Cancer Institute cohort. *Haematologica* **2023**, *108*, 69–82.
  11. Risitano, A.M.; Marotta, S.; Calzone, R.; Grimaldi, F.; Zatterale, A. Twenty years of the Italian Fanconi Anemia Registry: Where we stand and what remains to be learned. *Haematologica* **2016**, *101*, 319–327.
  12. Kutler, D.I.; Singh, B.; Satagopan, J.; Batish, S.D.; Berwick, M.; Giampietro, P.F.; Hanenberg, H.; Auerbach, A.D. A 20-year perspective on the International Fanconi Anemia Registry (IFAR). *Blood* **2003**, *101*, 1249–1256.
  13. Alter, B.P. Fanconi anemia and the development of leukemia. *Best Pract. Res. Clin. Haematol.* **2014**, *27*, 214–221.
  14. Alter, B.P.; Giri, N.; Savage, S.A.; Rosenberg, P.S. Cancer in the national cancer institute inherited bone marrow failure syndrome cohort after fifteen years of follow-up. *Haematologica* **2018**, *103*, 30–39.
  15. Ceccaldi, R.; Parmar, K.; Mouly, E.; Delord, M.; Kim, J.M.; Regairaz, M.; Pla, M.; Vasquez, N.; Zhang, Q.S.; Pondarre, C.; et al. Bone marrow failure in fanconi anemia is triggered by an exacerbated p53/p21 DNA damage response that impairs hematopoietic stem and progenitor cells. *Cell Stem Cell* **2012**, *11*, 36–49.
  16. Rodríguez, A.; Zhang, K.; Färkkilä, A.; Filiatrault, J.; Yang, C.; Velázquez, M.; Furutani, E.; Goldman, D.C.; García de Teresa, B.; Garza-Mayén, G.; et al. MYC Promotes Bone Marrow Stem Cell Dysfunction in Fanconi Anemia. *Cell Stem Cell* **2020**.
  17. Tjhuis, A.E.; Foijer, F. Characterizing chromosomal instability-driven cancer evolution and cell fitness at a glance. *J. Cell Sci.* **2024**, *137*.
  18. Abdallah, B.Y.; Horne, S.D.; Stevens, J.B.; Liu, G.; Ying, A.Y.; Vanderhyden, B.; Krawetz, S.A.; Gorelick, R.; Heng, H.H.Q. Single cell heterogeneity: Why unstable genomes are incompatible with average profiles. *Cell Cycle* **2013**, *12*, 3640–3649.
  19. Davoli, T.; Xu, A.W.; Mengwasser, K.E.; Sack, L.M.; Yoon, J.C.; Park, P.J.; Elledge, S.J. XChromosomal haploinsufficiency and triplosensitivity drive aneuploidy patterns and shape the cancer genome. *Cell* **2013**, *155*, 948.

20. Liu, G.; Stevens, J.B.; Horne, S.D.; Abdallah, B.Y.; Ye, K.J.; Bremer, S.W.; Ye, C.J.; Chen, D.J.; Heng, H.H. Genome chaos: Survival strategy during crisis. *Cell Cycle* **2014**, *13*, 528–537.
21. Cioc, A.M.; Wagner, J.E.; MacMillan, M.L.; DeFor, T.; Hirsch, B. Diagnosis of myelodysplastic syndrome among a cohort of 119 patients with fanconi anemia: Morphologic and cytogenetic characteristics. *Am. J. Clin. Pathol.* **2010**, *133*, 92–100.
22. Sebert, M.; Gachet, S.; Leblanc, T.; Rousseau, A.; Bluteau, O.; Kim, R.; Ben Abdelali, R.; Sicre de Fontbrune, F.; Maillard, L.; Fedronie, C.; et al. Clonal hematopoiesis driven by chromosome 1q/MDM4 trisomy defines a canonical route toward leukemia in Fanconi anemia. *Cell Stem Cell* **2023**, *30*, 153-170.e9.
23. Butturini, A.; Gale, R.P.; Verlander, P.C.; Adler-Brecher, B.; Gillio, A.P.; Auerbach, A.D. Hematologic abnormalities in Fanconi anemia: An International Fanconi Anemia Registry study. *Blood* **1994**, *84*, 1650–1655.
24. Berger, R.; Jonveaux, P. Clonal chromosome abnormalities in Fanconi anemia. *Hematol. Cell Ther.* **1996**, *38*, 291–296.
25. Quentin, S.; Cucchini, W.; Ceccaldi, R.; Nibourel, O.; Pondarre, C.; Pagès, M.P.; Vasquez, N.; D'Enghien, C.D.; Larghero, J.; De Latour, R.P.; et al. Myelodysplasia and leukemia of fanconi anemia are associated with a specific pattern of genomic abnormalities that includes cryptic RUNX1/AML1 lesions. *Blood* **2011**, *117*, 161–170.
26. Mehta, P.A.; Harris, R.E.; Davies, S.M.; Kim, M.O.; Mueller, R.; Lampkin, B.; Mo, J.; Myers, K.; Smolarek, T.A. Numerical chromosomal changes and risk of development of myelodysplastic syndrome-acute myeloid leukemia in patients with Fanconi anemia. *Cancer Genet. Cytogenet.* **2010**, *203*, 180–186.
27. Lisker, R.; Gutiérrez, A.C. de Cytogenetic studies in Fanconi's anemia. Description of a case with bone marrow clonal evolution. *Clin. Genet.* **1974**, *5*, 72–76.
28. Tönnies, H.; Huber, S.; Kühl, J.S.; Gerlach, A.; Ebell, W.; Neitzel, H. Clonal chromosomal aberrations in bone marrow cells of Fanconi anemia patients: Gains of the chromosomal segment 3q26q29 as an adverse risk factor. *Blood* **2003**, *101*, 3872–3874.
29. Behrens, Y.L.; Göhring, G.; Bawadi, R.; Cöktü, S.; Reimer, C.; Hoffmann, B.; Sängler, B.; Käfer, S.; Thol, F.; Erlacher, M.; et al. A novel classification of hematologic conditions in patients with fanconi anemia. *Haematologica* **2021**, *106*, 3000–3003.
30. Heng, J.; Heng, H.H. Genome Chaos, Information Creation, and Cancer Emergence: Searching for New Frameworks on the 50th Anniversary of the "War on Cancer." *Genes (Basel)*. **2022**, *13*.
31. Fanconi Anemia Research Fund *Fanconi anemia clinical care Guidelines*; Sroka, I., Frohnmayer, L., Van Ravenhorst, S., Wirkkula, L., Eds.; 5th ed.; Fanconi Anemia Research Fund: Oregon, 2020;
32. Sánchez, S.; Reyes, P.; Barrera, M.A.M.; Mar-Tínez, A.P.; Frias, S. Fanconi anemia, Part 3. Cytogenetic monitoring in the bone marrow of patients with Fanconi anemia. *Acta Pediatr. Mex.* **2024**, *45*, 343–368.
33. McGowan-Jordan J, Simons A, S.M. ISCN 2020: An International System for Human Cytogenomic Nomenclature. *Cytogenet. Genome Res.* **2020**, *160*, 503.
34. Nguyen-Khac, F.; Bidet, A.; Daudignon, A.; Lafage-Pochitaloff, M.; Ameye, G.; Bilhou-Nabéra, C.; Chapiro, E.; Collonge-Rame, M.A.; Cucchini, W.; Douet-Guilbert, N.; et al. The complex karyotype in hematological malignancies: a comprehensive overview by the Francophone Group of Hematological Cytogenetics (GFCH). *Leukemia* **2022**, *36*, 1451–1466.
35. Engel, J.L.; Zhang, X.; Wu, M.; Wang, Y.; Espejo Valle-Inclán, J.; Hu, Q.; Woldehawariat, K.S.; Sanders, M.A.; Smogorzewska, A.; Chen, J.; et al. The Fanconi anemia pathway induces chromothripsis and ecDNA-driven cancer drug resistance. *Cell* **2024**, 1–16.
36. Heng, J.; Heng, H.H. Genome chaos: Creating new genomic information essential for cancer

macroevolution. *Semin. Cancer Biol.* **2022**, *81*, 160–175.

37. Ye, J.C.; Editors, H.H.H. *Cancer Cytogenetics and Cytogenomics*; ISBN 9781071639450.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.