

Review

Not peer-reviewed version

Genome Agnostic Reprogramming of Acute Myelocytic Leukemia Hallmarks by Targeting Non-Oncogene Addictions with Azacitidine plus Pioglitazone and All-Trans Retinoic Acid

[Dennis Christoph Harrer](#), [Florian Lüke](#), [Tobias Pukrop](#), [Albrecht Reichle](#)^{*}, [Daniel Heudobler](#)

Posted Date: 25 December 2025

doi: 10.20944/preprints202512.2313.v1

Keywords: relapsed/refractory acute myelocytic leukemia (AML); leukemia hallmarks as therapeutic target; non-oncogene addictions; differentiation; immuno surveillance; inflammation; metabolism; anakoinosis; pioglitazone; all-trans retinoic acid; azacitidine



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Review

Genome Agnostic Reprogramming of Acute Myelocytic Leukemia Hallmarks by Targeting Non-Oncogene Addictions with Azacitidine Plus Pioglitazone and All-Trans Retinoic Acid

Dennis Christoph Harrer ¹, Florian Lücke ^{1,2}, Tobias Pukrop ^{1,3}, Albrecht Reichle ^{1*} and Daniel Heudobler ^{1,3}

¹ Department of Internal Medicine III, Hematology and Oncology, University Hospital Regensburg, Regensburg, Germany

² Division of Personalized Tumor Therapy, Fraunhofer Institute for Toxicology and Experimental Medicine, Regensburg, Germany

³ Bavarian Cancer Research Center (BZKF), University Hospital Regensburg, Regensburg, Germany

* Correspondence: albrecht.reichle@ukr.de

Abstract

The search for new therapeutic principles is essential for treating relapsed/refractory (r/r) acute myeloid leukemia (AML). Novel principles include genome-agnostic differentiation induction, controlling AML-triggering inflammation, potentiating the immune response and 'normalizing' AML metabolism. This review summarizes data from a phase I study (10 patients, pts) and three case reports (7pts) on the treatment of r/r AML by reprogramming AML hallmarks using APA, low dose azacitidine, pioglitazone (PPAR α/γ agonist) and all-trans retinoic acid. APA reprograms the r/r AML phenotype in patients with clinically and molecularly/genetically unfavorable risk profiles (17pts, 16 refractory, one relapsed) in a genome-agnostic manner, restoring the plasticity of AML hallmarks, thereby improving immune surveillance, attenuating inflammation-triggered promotion of AML and distant microbial inflammation (healing of fungal pneumonia during induction of complete remission (CR) with APA), while normalizing leukemia metabolism (restoring phagocytosis and ROS production in leukemic neutrophils). APA induces complete remission (CR) in 10pts (59%), with only modest hematotoxicity following CR induction. This allows treatment to be carried out in an outpatient setting, including for elderly and comorbid patients. Triple transcriptional modulation, facilitated by epigenetic modelling with azacitidine, targets reprogramming of non-oncogene addiction networks in AML, re-establishing functionally active, closely interrelated myeloid hallmarks and AML cell death genome-agnostically.

Keywords: Relapsed/refractory acute myelocytic leukemia (AML); leukemia hallmarks as therapeutic target; non-oncogene addictions; differentiation; immuno surveillance; inflammation; metabolism; anakoinosis; pioglitazone; all-trans retinoic acid; azacitidine

1. Introduction

Despite the steadily growing portfolio of approved drugs, including classic cytotoxic therapies such as CPX351 (liposomal encapsulation of cytarabine/daunorubicin at a molar ratio of 5:1), immunotherapies such as improved techniques of allogeneic hematopoietic stem cell transplantation (allo-HSCT) and CAR T cell therapy, as well as chemo-immunoconjugates such as gemtuzumab ozogamicin (a CD33 antibody–drug conjugate) and drugs targeting non-oncogene addiction targets for large-scale reprogramming AML (such as hypomethylating agents (HMAs), 5-azacytidine

(azacitidine) administered intravenously or orally or 2-deoxy-5-azacitidine (decitabine), remission induction and maintenance therapy in acute myeloid leukemia (AML) for relapsed or refractory (r/r) AML is still challenging. Besides HMAs, which initiate epigenetic modelling and derepression of tumor suppressor genes further drugs targeting non-oncogene addictions are venetoclax (a BCL2 inhibitor); glasdegib (a hedgehog inhibitor). A specific group of drugs directed towards oncogene addictions, whose clinical activity depends on the presence of recurrent aberrations, includes midostaurin, gilteritinib and quizartinib (FMS-like tyrosine kinase 3 inhibitors); ivosidenib, olutasidenib and enasidenib (isocitrate dehydrogenase 1 and 2 inhibitors); and others [1–17].

The search for novel therapeutic principles for treating r/r AML is imperative, including exploring mechanisms for inducing differentiation, controlling inflammation, potentiating the immune response, and normalizing AML metabolism to prevent the rapid repopulation of blasts [18–23].

At first glance, the unique functional defect acquired by AML blasts, the inability to differentiate into neutrophils under disease conditions, may be at the center of attention. However, the question of how systemic AML therapy may induce genome agnostic differentiation in non-acute promyelocytic leukemia (non-PML) AML remains widely unanswered. Furthermore, how are the differential molecular rationalizations for differentiation induction organized in commonly genetically heterogeneous r/r non-PML AML [17,24,25]?

Similar to ATRA or arsenic trioxide in PML, isocitrate dehydrogenase 1 and 2 inhibitors, menin inhibitors and FMS-like tyrosine kinase 3 inhibitors induce differentiation in AML in an oncogene-addicted way [12,26]. Differentiation in non-PML AML facilitates clinically valuable control of leukemia. [12,18,27–29]. The therapeutic challenge lies in demonstrating that the various rationalizations for differentiation can be reconstituted therapeutically with reduced toxicity in molecularly and genetically heterogeneous relapsed/refractory (r/r) non-PML AML.

Substituting the BCL-2 inhibitor venetoclax for a peroxisome proliferator-activated receptor alpha/gamma (PPAR α/γ) agonist (pioglitazone) and an all-trans retinoic acid (ATRA) agonist (APA schedule) within the established azacitidine/venetoclax (aza/ven) schedule may promote the differentiation of blasts to neutrophils in r/r non-PML AML and induces leukemia cell death, as well as comprehensively 'normalized' myeloid hallmarks. [18,25,30–33]. When applied at a reduced dose within the APA schedule, azacitidine acts as a facilitator for the activity of pioglitazone/ATRA. [18]. Otherwise, neither of the nuclear receptor agonists would have any clinically relevant activity in r/r AML, while all-trans retinoid acid (ATRA) alone may dramatically promote differentiation in promyelocytic leukemia (PML) [25,26].

Successful differentiation induction in r/r non-PML AML reveals still pathophysiologically preserved and therapeutically actionable pathways if azacitidine may pave the way for triple transcriptional modulation by generating stress conditions in non-PML AML bone marrow tissue or extramedullary disease [18,28,34]. In vitro data show that the well-known monoactivity and differentiation-inducing capacity of azacitidine may be enhanced to restore phagocytic capability and increase reactive oxygen species in non-PML AML neutrophils [18,25]. Even in cases of azacitidine failure, complete remission (CR) can be achieved, which highlights the novel synergistic activity profile of the APA drug cocktail [18,27]. CR may be established after just one cycle of APA therapy in patients with r/r non-PML AML [18].

This review summarizes the data from a phase I study of 10 patients with r/r non-PML AML who were treated with APA (azacitidine, pioglitazone and all-trans retinoic acid) together with published data on an additional seven cases of r/r non-PML AML treated with the APA protocol [18,27–29]. The treatment results for these 17 patients with r/r non-PML AML support the use of a drug combination that induces differentiation and leukemia cell death genome agnostically, even in patients with highly heterogeneous cytogenetic/molecular-genetic aberration patterns and complex pre-treatment conditions. For example, one patient had an early relapse following allogeneic hematopoietic stem cell transplantation (allo-HSCT) [29].

The data reveal that the differentiation associated with the improvement of immunosurveillance, as well as the attenuation of inflammation and the 'normalization of leukemia metabolism' by pioglitazone, may be followed by leukemia cell death and complete remission (CR), even in r/r non-PML AML, while synergistically targeting non-oncogene additions with three nuclear receptor agonists: the peroxisome proliferator-activated receptor (PPAR) α/γ agonist, the retinoid receptor RARA agonist all-trans retinoid acid (ATRA), and the epigenetic modulator, azacitidine [18,27–29,35,36].

2. APA Therapy

Patients selected for APA therapy were not eligible for dose-intensive therapy or allo-HSCT due to age or (treatment-related) comorbidity or due to insufficient induction response [18,27–29]. The time to progression (n = 16) or relapse (n = 1) was less than six months.

Between 2014 and 2020, a total of 17 patients with r/r non-PML AML received the APA schedule (azacitidine, pioglitazone and all-trans-retinoic acid). Ten of these patients were treated within a phase I trial of the German-Austrian AML Study Group (AMLSG) in Germany (the ViVA trial) [18], seven consecutive patients also received APA and their cases were published due to their significant responses [27–29].

Patients in the AML ViVA trial were treated with an ATRA dose of 45 mg/m²/day from days 1 to 28, followed by 15 mg/m² per os from day 29 onwards, in combination with subcutaneously administered azacitidine (75 mg fixed dose from days 1 to 7 of each 28-day treatment cycle) and pioglitazone (45 mg per os continuously from day 1). Non-study patients received APA with a fixed daily dose of ATRA (45 mg per os) continuously from day 1. Following APA induction, patients continued APA therapy for as long as was clinically appropriate, until AML relapsed or progressed. Cycle intervals beyond the first cycle were adapted to the time of hematopoietic regeneration.

3. Activity Profile of APA Therapy in Refractory or Relapsed non-PML AML

Patients received a median of three cycles of the study treatment (range 1–14). The rapid onset of complete remission (CR) in r/r non-PML AML, even after one or two cycles, highlights the effectiveness of APA rescue therapy in r/r non-PML AML. Responding patients showed improvement or normalization of peripheral blood cell counts. However, early neutrophil recovery may be due to blast differentiation [27].

The low toxicity of APA enabled outpatient treatment for patients with CR and PR, including those aged 70 years and over (n = 4).

The rescue therapy, APA works in r/r non-PML AML and can induce complete remission (CR), complete remission with residual thrombocytopenia (<100 G/L) or neutropenia (<1 G/L) (CRi), complete cytogenetic remission (CRc) or complete molecular remission (CRm). Ten of the seventeen patients (59%) who did not achieve CR in response to previous therapy experienced CR, CRi/c/m. APA maintained CR or PR for a long time (Table 1). Two of the four patients aged over 70 remained in continuous complete remission (cCR) for over six months.

Table 1. AML diagnostics/history of pre-treatment and response to APA therapy.

Table 1	AML diagnostics/history of pre -treatment and response to APA therapy			
	Variable	Best response	Cycles/ No. of days on APA treatment	N=17
Sex, N (%)	Male	-	-	17 (100)
	Female	-	-	1 (6)
Age in years, median (range)	Age >= 70 years	-	-	66 (55-76)
	Age < 70 years	3x CR, 1x SD (1xAML, 3x de novo)	-	4 (70-76)
Type of AML, N (%)	de novo AML	3x CR, 1x PR, 4x SD, 1x NA	CR 10/350, 4/240, 4/1405/175; PR 14/490; SD 1 to 3, NA-	9 (52)
	secondary AML	2x SD, 1x PR, 1x CR	SD 2x1/30; PR 3/110; CR 5/210	4 (24)
ECOG performance score, N (%)	0	-	-	3 (18)
	1	-	-	12 (70)
Cytogenetics at baseline, N (%)	Complex karyotype (+/- molecular genetic aberrations)	SD, PR, CR	-	14 (82)
	90 complex karyotype involving chromosomes 5, 7, 8, 12, 13, 18, 21, and 22	1 CR skin	6/210	1 (6)
Molecular genetics (mutations) at baseline, N (%)	TP53	2 CR	4/140; 5/210	2 (12)
	TP53, ASXL1	2x CR, 1x PR	2/74; 5/175; 3/110	3 (17)
ELN 2022 genetic risk category, N (%)	adverse	-	-	17 (100)
N and type of pretreatment, N (%)	1 cycle	CR, SD	-	9 (53)
	2 cycles	NA, SD	-	7 (41)
3 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
4 cycles	CR	-	-	6 (35)
	PR	-	-	3 (17)
5 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
6 cycles	CR	-	-	2 (12)
	PR	-	-	2 (12)
7 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
8 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
9 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
10 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
11 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
12 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
13 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
14 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
15 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
16 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
17 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
18 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
19 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
20 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
21 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
22 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
23 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
24 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
25 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
26 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
27 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
28 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
29 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
30 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
31 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
32 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
33 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
34 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
35 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
36 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
37 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
38 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
39 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
40 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
41 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
42 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
43 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
44 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
45 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
46 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
47 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
48 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
49 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
50 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
51 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
52 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
53 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
54 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
55 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
56 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
57 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
58 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
59 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
60 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
61 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
62 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
63 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
64 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
65 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
66 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
67 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
68 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
69 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
70 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
71 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
72 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
73 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
74 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
75 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
76 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
77 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
78 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
79 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
80 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
81 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
82 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
83 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
84 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
85 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
86 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
87 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
88 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
89 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
90 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
91 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
92 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
93 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
94 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
95 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
96 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
97 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
98 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
99 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)
100 cycles	CR	-	-	1 (6)
	PR	-	-	1 (6)

For patients with refractory non-PML AML, achieving eligibility for allo-HSCT is a high priority. The quality of remission prior to allo-HSCT is important for the long-term outcome [37]. Consolidation with allo-HSCT was facilitated by first-time CR with APA therapy in three patients with refractory non-PML AML (n = 16).

The pathogenesis of de novo, secondary or treatment-related non-PML AML seems to have no significant impact on the response to APA therapy within small subgroups, as all three groups achieved CR, CRi/c/m: in de novo AML, three out of nine patients (33%); in secondary AML, one out of four patients (25%); and in treatment-related AML, three out of four patients (75%).

According to the 3-tiered venetoclax prognostic risk score (VEN-PRS), extramedullary disease is a poor prognostic parameter for aza/ven treatment [38]. One patient with complex aberrant cytogenetic involving chromosomes 5, 7, 8, 12, 13, 18, 21, and 22 (Karyotype formula: 45,XX,der(5;7)(5pter->5p11::7q22->7q11::5p11->5q11::7q11->7pter)[4]; 45,XX,der(5;7)(5pter->5p11::7q22->7q11::5p11->5q11::7q11->7pter), der(12;18)t(12;18)(p13;q23)del(12)(p12p13)[2]; 45,XX,der(5;7)(5pter->5p11::7q22->7q11::5p11->5q11::7q11->7pter), +8,der(12;18)t(12;18)(p13;q23)del(12)(p12p13), ider(13) (q10)del(13)(q13q22), +21,+22[4]; 46,XX[3]; ISCN: nuc ish 7cen(D7Z1x2), 7q31(D7S486x1), 8cen(D8Z2x3), 12p13(3'ETV6x1,5'ETV6x2)(3'ETV6 con 5' ETV6x1)[74/100]) achieved continuous CR of extramedullary non-PML AML in the skin. The multilineage dysplasia with ringsideroblasts (MDS-RS-MLD) remained stable [28].

Due to the small patient population, the observation that patients with complex karyotypes and no recurrent aberration (n = 7; 41%) responded poorly (100% no CR) may be accidental. Further studies should confirm this observation. At diagnosis, the following recurrent aberrations were routinely analyzed in the standard panel: MLLT3/MLL, CEBPA, PML/RARA, RUNX1, FLT3-ITD and FLT3-TKD, as well as NPM1.

In contrast, patients with complex karyotypes and additional molecular-genetic aberrations (n = 7, 41%) and those with molecular-genetic aberrations only (n = 2) responded with CRi/c/m (n = 6, 86%) and CR (n = 2, 100%), respectively. Considering molecular aberrations irrespective of complex aberrant karyotypes, eight out of nine patients (89%) achieved a CR.

Moreover, three high-risk patients with induction failure and a TP53 mutation responded: two achieved a CR, and one achieved a prolonged partial response (PR). As recently shown, this result was also achieved with decitabine plus ATRA [39]. According to the 3-tiered venetoclax prognostic risk score (VEN-PRS) for aza/ven treatment, CR in TP53 mutated AML is generally difficult to achieve, as is the case with NF1 and FLT3 mutations [40,41] (Table 1). FLT3 mutations commonly

render cells resistant to aza/ven [39] One patient with CMML, transformed into non-PML AML, had no aberration and could be reversed to CMML-2 with APA therapy [27].

Even a single intensive treatment regimen, including allo-HSCT, did not affect the likelihood of a positive response [29]. Patients who underwent one, two or three intensive treatment schedules achieved a complete response (CR), including a patient who had relapsed after allo-HSCT.

A total of seven patients died during the follow-up period. The median overall survival (mOS) in the ViVA trial was 131 days (i.e., 4.3 months).

4. Adverse Events on Patient Basis

The mean treatment duration for the 17 patients was 136 days (range 27–426 days), following a median follow-up period of 142 days. Major grade 3/4 events were likely due to baseline cytopenias caused by preceding treatment failure, AML progression, and toxicity from dose-intensive induction therapies (Table 2).

Therefore, hematological toxicities and infections are primarily attributable to the adverse starting conditions for APA therapy in advanced treatment lines.

Table 2. Hematologic parameters before APA treatment.

Table 2

Hematologic parameters before APA treatment	
Variable	Readout
Hemoglobin g/dL, median (range)	8.8 (6.3-10.6)
Platelets x10 ⁹ /L, median (range)	48.3 (5-175)
White blood count x10 ⁹ /L, median (range)	0.93 (0.33- 4.8)
Neutrophils x10 ⁹ /L, median (range)	0.19 (0-2.7)
Bone marrow blasts %, median (range)	58 (10-90)
Peripheral blood blasts %, median (range)	0 (0-26)

The 3. infection-associated adverse events (AEs) for the 17 summarized patients is similar to that published for the AML ViVA trial, which included 10 patients. Fever occurred in three patients (18%), lung infections in six patients (35%), urinary tract infections in four patients (24%), sepsis in one patient (6%), and other infections in two patients (12%) [18].

No additional non-hematological toxicities of grades 3 or 4 occurred in comparison to the ViVA study population. However, hematological toxicities occurred in the additional seven patients: five cases of grade 4 neutropenia, three cases of grade 3 thrombocytopenia and one case of grade 4 thrombocytopenia [27–29].

A retrospective study of venetoclax-based HMA combinations in r/r AML shows grade ≥4 neutropenia in 100% of patients, grade ≥4 thrombocytopenia in 95.7%, and grade ≥4 febrile neutropenia in 45%. Compared to this toxicity data, APA treatment appears to be less myelosuppressive. Pioglitazone and ATRA do not increase toxicity [42].

Importantly, the concerted targeting of non-oncogene addictions predominantly with APA results in modest toxicity characterized by myelotoxicity, while promoting multiple favorable biological effects via the re-establishment of myeloid hallmarks, leading to AML cell death.

5. APA as Rescue Therapy for Azacitidine Failure

APA therapy successfully reversed the effects of azacitidine failure in three cases. Rescuing azacitidine failure with pioglitazone and ATRA illustrates the synergistic activity of the triple APA

combination. Azacitidine failure generally indicates a poor prognosis for overall survival (OS), according to the three-tiered venetoclax prognostic risk score (VEN-PRS) [38]. Nevertheless, azacitidine paves the way for successful triple transcriptional modulation via receptor-triggered nuclear receptors, even though it was clinically ineffective as monotherapy [43].

Of the two patients with azacitidine failure, one with transformed chronic myelomonocytic leukemia (CMML) responded with CMML-2 after two cycles of APA therapy, while the other patient with an MLL-PTD mutation achieved a complete remission (CR) and received a successful allogeneic hematopoietic stem cell transplantation (allo-HSCT) as consolidation [27]. In a further patient, APA therapy rescued azacitidine failure following early relapse after allo-HSCT in a patient with high-risk AML. APA led to complete donor chimerism within one cycle [29].

Recently published results suggesting that PPAR γ antagonists could be used to overcome decitabine/ven resistance contradict clinical results achieved with APA, as azacitidine resistance may be reversed by a PPAR α/γ agonist (pioglitazone) when administered alongside APA [32]. Data from APA are consistent with results from the same group showing that pioglitazone induces apoptosis in AML in vitro without coculturing AML blasts with stroma [44]. The specific impact of pioglitazone's agonistic activity on PPAR α cannot be determined [45–47].

Contradictory preclinical and clinical data on the targeting of nuclear transcription factors may be resolved by considering the systems context of adjacent stroma and AML blasts, and by focusing in vitro and in clinical trials on drug combinations including at least one nuclear transcription factor agonist. For example, there have been positive clinical results for the combination of decitabine and ATRA in non-PML AML, and in the case of APA, even after preceding azacitidine failure. [18,25,48,49] (Figure 1, 2).

Pioglitazone and/or ATRA alone are clinically ineffective in non-PML AML, although both drugs are physically interacting at their receptor sites [50]. Following the rescue of clinical azacitidine failure in r/r non-PML AML, the reciprocal biological activity of the two drug groups, pioglitazone/ATRA and azacitidine, becomes apparent [51]. Although azacitidine is clinically ineffective, it is found to be the decisive facilitator of pioglitazone/ATRA activity [18,25].

A recent experimental study shows that the resistance of AML blasts to pioglitazone may be mediated by stromal cells and can be overcome by an AXL inhibitor, bemcentinib [44]. These data support the simultaneous targeting of AML blasts and adjacent stroma cells to overcome resistance as shown with the APA schedule in clinical practice. Dual versus triple exposure of AML blasts to APA drug components supports the clinically observed azacitidine rescue approach, as triple activity is a prerequisite for establishing cell differentiation and regaining the neutrophil activity profile in leukemic neutrophils [25].

The re-establishment of novel tissue homeostasis by reprogramming the communication between transcriptional networks maintained by tumor and adjacent stromal cells is called 'anakinosis', a novel therapeutic principle [52,53]. Azacitidine serves as the initiating drug, implementing the prerequisites for the clinically efficacious application of triple transcriptional modulation, allowing differentiation induction and the re-establishment of immunosurveillance, as demonstrated both clinically and experimentally [18,25].

6. APA and re-Establishment of Immuno Surveillance

Of the patients in the series who qualified for APA therapy as a last-resort treatment for r/r non-PML AML, two suffered from additional autoimmune diseases: one had psoriasis involving the large joints, and the other had systemic lupus erythematosus [54,55]. However, pre-treatment with cytotoxic agents did not sufficiently control the autoimmune diseases. Both diseases were finally controlled after one cycle of APA therapy, revealing APA's strong immunoregulatory activity profile, as has been demonstrated for APA-like therapies, including metronomic low-dose chemotherapy, in non-hematological neoplasias [54,56,57].

Consistent with this is the finding that restored immunocompetence can treat severe fungal pneumonia acquired during intensive induction therapy and induction failure (Figure 1). The fungal

pneumonia was resolved after two cycles of APA therapy, after which the patient qualified for allo-HSCT, which was successfully performed [18] .




Figure 1. Simultaneously targeting non-oncogene addiction in both bone marrow stromal cells and leukemia cells using APA establishes myeloid hallmarks that impact the immune response in distant organs, such as the lungs.

Regaining immune surveillance during APA therapy may be due to the immunoregulatory effects of the drugs themselves, as well as the differentiation of blasts into neutrophils being associated with the restoration of neutrophil functions [25,58,59]. This has been demonstrated experimentally, with the reacquisition of phagocytic activity and increased reactive oxygen species (ROS). These are both prerequisites for processing microbial infections [18,25].

Targeting NRF2 with pioglitazone and attenuating the oxidative stress and stress conditions induced by azacitidine may ultimately enhance immune surveillance [20,47,60–62]. Experimental data show that T-cell specific response to breast cancer is possible in the combination metronomic chemotherapy plus pioglitazone and also following DNA methyltransferase inhibition [57,59].

Achieving complete donor chimerism with APA, but not azacitidine, reveals a synergistic immunological effect of APA with the graft-versus-leukemia (GvL) reaction. Differentiation of blasts may again play an important role here, as the overexpression of antigens such as proteinase 3 and other azurophil granule proteins may serve as targets for allogeneic and autologous T-cell responses [63–65]. Azacitidine-associated DNA methyltransferase inhibition may further enhance T-cell responses by inducing interferon and upregulating major histocompatibility class I (MHC-I) genes [59,66] .

7. Differential AML Hallmarks Targeted with Pioglitazone, ATRA

In addition to clinical and experimental observations regarding the re-establishment of myeloid hallmarks in AML blasts and the differentiation and reconstitution of immune surveillance, APA modulates the metabolism of leukemia cells (e.g. leukemic neutrophils) and the inflammation that promotes AML, as well as microbial inflammation at a distant site (e.g. resolution of fungal pneumonia) [18,25,38]. The combination of pioglitazone and ATRA targets both AML blasts and stromal cells, reestablishing hallmarks that are therapeutically reconstituted via concerted regulatory

activity in AML blasts and adjacent stroma [18,22,25,50]. Distant inflammation, manifested as fungal infection, may benefit from the therapeutic re-establishment of myeloid hallmarks.

Inflammation is considered to be at the forefront of AML pathogenesis, as has recently been demonstrated [38]. A blood-based leukemia inflammatory risk score (LIRS) is a significantly better predictor than the current European LeukemiaNet 2022 risk model, which considers genetic, molecular-genetic and clinical risk factors. LIRS also serves as independent prognostic factor for overall survival based on both known clinical risk profiles and inflammation-based markers in serum. Thus, inflammation control by pioglitazone for AML control resumes the originally approved activity profile in insulin-resistant diabetes mellitus, inflammation control [35,67–69]. Moreover, attenuation of pro-inflammatory drivers of AML pathogenesis might explain alone genome agnostic activity of APA. As recently shown, differential APA-like schedules designed for multiple r/r neoplasias reveal the correlation of serum C-reactive protein response and neoplasia control [19,56].




Figure 2. Therapy with APA (azacitidine, pioglitazone and all-trans retinoic acid): Remodelling non-oncogene addictions, biological and clinical results, and systems-relevant correlates in relapsed/refractory acute myeloid leukemia.

The Leukemia Inflammatory Risk Score (LIRS) uses blood-based pro-inflammatory biomarkers to predict AML survival. These markers provide a key rationale for substantial anti-inflammatory therapy strategies, e.g. pioglitazone for AML control [18,38] . Interestingly, the LIRS remains relevant regardless of the intensity of previous treatment.

APA-like schedules have principally demonstrated potent anti-inflammatory effects in various neoplasias, cancers, including hematological neoplasias, when treated with metronomic low-dose chemotherapy in conjunction with additional transcriptional modulators. These schedules can control relapsed/refractory neoplasms up to the induction of continuous CR (cCR) [56,70].

An excellent example of how pioglitazone attacks leukemia stem cells in chronic myeloid leukemia (CML) has been demonstrated through its interaction with imatinib in patients whose imatinib molecular response is failing. [35,71]. Once again, the addition of pioglitazone or rosiglitazone can revitalize the activity profile of monotherapies that have failed, such as imatinib, azacitidine, MEK inhibitors and gefitinib (an EGFR TKI) [18,27,28,71–73].

Broadly speaking, a variety of approaches to therapy that induce anakoinosis with APA-like therapies may focus on establishing distinct disease-relevant hallmarks that are lost during neoplastic transformation. These approaches can be used to control recurrent or refractory neoplasms

(hematological neoplasms, sarcomas and cancers), and presumably r/r non-PML AML as well [35,56,74].

Experimental designs considered the therapeutic use of PPAR γ agonists/antagonists. Resistance to decitabine/ven in AML may be induced by the upregulation of fatty acid metabolism, which is thought to be counter-regulated by PPAR γ antagonists [32]. The antileukemic activity of selenium may be mediated by activation of PPAR γ via the agonist prostaglandin J2. [75]

Drugs that target oncogenic aberrations in AML might enhance the biomodulatory effects of therapeutic approaches such as APA. Clinical trials have included the FLT3 inhibitor gilteritinib as an adjunct to aza/ven. In FLT3-positive relapsed/refractory AML, the CR/CRi rate was 27%, but impressive toxicity remains a striking problem [16]. ATRA may increase cell death induced by quizartinib [76].

8. APA and AML Microenvironment

The resolution of extensive leukemic skin infiltration in a patient with treatment-related MDS-RS-MLD, but no increased blast count in the bone marrow, impressively demonstrates the impact of environmental context (bone marrow versus skin) on leukemia growth control with APA therapy. The MDS-RS-MLD transformed into AML after 210 days of APA treatment, during which there were no skin relapses [28]. Despite repetitive therapies for bone marrow AML, the leukemic skin infiltration did not recur. These observations highlight the importance of simultaneously targeting AML blasts and adjacent stromal cells to effectively control AML.

Myeloid hallmarks may be reinstated by APA-mediated tissue reprogramming of AML. Myeloid hallmarks are collectively constituted via targeting of AML blasts and adjacent stromal cells [62,68]. The targeted nuclear receptors are typically distributed throughout the stroma and leukemia cells. PPAR γ expression may change depending on the context [35,44].

The adaptation of the transcriptionally active drug cocktail for use in r/r neoplasms may be tumor-agnostic, as demonstrated in r/r multisystem Langerhans cell histiocytosis (mLCH) and r/r renal clear cell carcinoma (RCCC). In patients with r/r mLCH who failed to respond to treatment with pioglitazone and dexamethasone, substitution of the transcriptionally active drug cocktail with interferon- α and pioglitazone — a schedule that had previously been successfully tested for r/r renal clear cell carcinoma — resulted in cCR. Each transcriptional cocktail was combined with metronomic low-dose chemotherapy [74,77–79].

As with the combination of a PPAR γ agonist and interferon- α for controlling r/r RCCC/mLCH, the combination of a PPAR γ agonist and a RAR agonist (ATRA) can also be successfully used in a tumor-agnostic manner for treating r/r non-PML AML or muscle-invasive bladder cancer (in a mouse model) [56,80]. Facilitators of efficacious combined transcriptional intervention include metronomic low-dose chemotherapy, azacitidine and MEK inhibitors, revealing the potential for further systems pharmacological exploitation [80].

9. APA Therapy Predominantly Targets Non-Oncogene Addictions

Clinically used azacitidine doses predominantly exert their activity through hypomethylating effects [14,81,82]. As AML blasts depend heavily on controlling DNA methylation, the epigenetic modifier azacitidine is a key inhibitor of the methylation process, notably in a non-oncogene-addicted manner [81,82]. Consequently, azacitidine neutralizes the suppression of tumor suppressor genes in parallel with pioglitazone [82,83]. RNA-mediated cytotoxic effects are not the main mechanisms of azacitidine action at clinically applicable doses [82].

Response to or primary resistance to azacitidine in non-PML AML cannot be predicted from specific mutational signatures, as azacitidine activity is not predominantly disruptive via cytotoxicity, but may reprogram AML cell fate by establishing myeloid hallmarks genome agnostically, such as differentiation [34,84].

Azacitidine makes full use of the plasticity of AML tissues, particularly by inducing novel stress responses in neoplastic cells. For example, it targets stress dependency on the antiapoptotic BCL-2 family proteins BCL-XL, BCL-2 and MCL-1 [85,86]. The clinical efficacy of azacitidine in combination with the BCL-2 inhibitor venetoclax indicates the therapeutic importance of the epigenetic modulation of non-oncogene addictions in the treatment of AML.

Compared to approved cytotoxic or aza/ven therapies, APA is a less toxic, synergistic, bioregulatory combination therapy [18,42]. The rationale behind APA is to induce AML cell death by re-establishing myeloid hallmarks in non-PML AML blasts, probably also in cases that are resistant to decitabine/ven [32,87].

Myeloid hallmarks may be reinstated by APA-mediated tissue reprogramming of AML. Myeloid hallmarks are collectively constituted via targeting of AML blasts and adjacent stromal cells [62,68]. The targeted nuclear receptors are typically distributed throughout the stroma and leukemia cells. PPAR γ expression may change depending on the context [35,44].

Azacitidine was administered at a lower dose than the approved one in the APA schedule. In the APA schedule, azacitidine predominantly serves as a facilitator for the re-establishment of AML plasticity (**Figure 1**). Unlike azacitidine in the aza/ven schedule, which has predominant cytotoxic effects, azacitidine in the APA schedule acts as a 'facilitator' to recover bioregulatory processes [18,25,88].

In non-PML AML, clinically non-functional transcriptional regulators such as pioglitazone and ATRA cannot unlock AML tissue plasticity independently of azacitidine's epigenetic regulation [18,25,29]. Preclinical data reveal a synergistic activity profile of pioglitazone and ATRA in a variety of neoplasias [25,88–90].

Inducing stress dependencies in the bone marrow tissue of AML patients through APA therapy leads to obvious genome-agnostic failure of the tissue to adapt to APA-induced stress, resulting in the development of resistance mechanisms [91]. Conversely, a targeted and coordinated reorganization of myeloid hallmarks occurs in AML blasts through the reciprocal activity of APA drugs. This enables APA to overcome AML addictions and establish the ability to differentiate into neutrophils under disease conditions [18,25]. Leukemia plasticity, which is often discussed as a prerequisite for the development of resistance, is now therapeutically exploited by involving the AML-silenced but still organizable networks of non-oncogene addictions that contribute to differentiation in AML [92] (**Figure 2**).

In summary, APA therapy promotes the plasticity of non-PML AML blasts by reprogramming the non-oncogene addiction networks in non-PML AML to reintegrate 'normalized' myeloid hallmarks.

However, there is variability in the rationalizations of the processes that guide differentiation and the restoration of immune competence, as demonstrated by the poor responses observed, particularly in AML patients with genetic aberrations, only. APA therapy successfully covers a wide range of chromosomal and molecular genetic aberrations due to the frequently occurring, uniquely organized non-oncogene addiction networks that facilitate the differentiation of blasts and the re-establishment of immune competence [59,64,93,94].

This principally demonstrates that non-oncogene addictions provide a significant means of achieving a clinically meaningful response in r/r non-PML AML [18,64]. The huge heterogeneity of chromosomal and molecular genetic aberrations in non-PML AML contrasts with the limited diversity of rationalizations that facilitate myeloid differentiation and the reconstitution of immune competence. Due to the genome-agnostic activity of the APA schedule in mediating differentiation induction, a limited number of schedules, such as APA or APA-like schedules, seem necessary to cover differentiation in AML blasts, despite the significant genetic heterogeneity.

10. APA, a Genome Agnostic Therapy

Descriptive statistics show that only one subgroup of refractory non-PML AMLs responds poorly to APA: those with chromosomal aberrations only. Despite the significant heterogeneity of

molecular and genetic aberrations, the unfavorable prognostic features of the 16 refractory patients and one relapsed did not preclude the possibility of an APA response with complete remission (CR), complete remission with incomplete blood count recovery (CRi), complete remission with partial blood count recovery (CRi) or complete remission with minor blood count recovery (CRm). CRs occurred in elderly patients (≥ 70 years), in de novo, secondary or treatment-related AML, and in patients with a high number of pre-treatments. This occurred independently of molecular-genetic aberrations, plus/minus complex aberrant karyotypes.

The wide range of genetic aberrations in AML blasts within the APA cohort, and the unique, therapeutically successful targeting of growth-promoting AML hallmarks with APA, reveal that APA activity is genome-agnostic (**Table 1, Figure 2**).

Novel targeted therapies that induce differentiation in AML blasts are predominantly active in the presence of a single recurrent mutation; therefore, they are not generally applicable. Isocitrate dehydrogenase 1 and 2 inhibitors, menin inhibitors and FMS-like tyrosine kinase 3 inhibitors can cause lysis syndromes similar to those seen with all-trans retinoid acid (ATRA) therapy for PML [95].

These recurrent mutations may be detected in 12–25% of non-PML AML cases and comprise key genetic drivers of AML (e.g., KMT2A). Their occurrence may also increase with age [96,97]. Targeting these mutations therapeutically to block AML differentiation may elicit a response in patients who have not responded to previous AML therapies. However, the APA cohort studied comprises a sizable proportion of genetically heterogeneous AML patients who may benefit from APA therapy inducing anakoinosis. The comparison of APA-mediated differentiation versus differentiation induction by inhibitors of recurrent mutations reveals that AML blasts may provide different rationalizations for initiating differentiation (**Figure 2**). In contrast to therapies targeting recurrent mutations, APA therapy is multi-level and restores a pattern of stroma-linked hallmarks, differentiation, immune response and inflammation control and AML metabolism. This has also been demonstrated in a variety of r/r neoplasias [51,56,98].

11. Second-Line aza/ven Versus APA in r/r non-PML AML

Selecting salvage therapies for patients with relapsed or refractory AML remains challenging and controversial, as response rates are poor [12,30,99,100].

Published second-line response and overall survival (OS) rates to AZA/VEN are compared to those achieved with APA. As aza/ven is frequently used in the first line of treatment for elderly or medically unfit patients, it is often not considered as a second-line treatment, so alternative approaches are welcome [101].

Importantly, aza/ven has changed the prognostic parameters for first- and second-line AML treatment compared to classic intensive induction therapies [38]. Aza/ven responses demonstrate a novel therapeutic approach to AML compared to chemotherapy [102].

A large retrospective study developed a three-tiered venetoclax prognostic risk score (VEN-PRS) for azacitidine/venetoclax-treated r/r AMLs. According to the novel VEN-PRS scoring system, the APA patient cohort comprises patients at adverse risk, as indicated in Table 1. The median overall survival (OS) for the ViVA trial cohort ($n = 10$ patients) was 131 days (i.e. 4.3 months), reflecting the OS of the adverse patient population with a VEN-PRS of 4.6 months (1.7–8.2 months) [38]. Further large-scale APA trials are necessary to fully demonstrate the impact of APA on survival.

The APA patient cohort ($n = 17$) comprises patients at adverse risk, 10 of whom (59%) have mutations in TP53 ($n = 3$), NF1 ($n = 1$) and FLT3 ($n = 2$). Furthermore, three patients experienced failure of azacitidine pretreatment, and one patient had extramedullary AML, according to the adverse risk parameters included in the VEN-PRS. Additionally, the APA cohort was characterized by multiple myelodysplasia-related gene mutations, which are associated with poor outcomes based on classic intensive AML therapies. Overall survival (OS) was significantly shorter for patients with RUNX1 ($n = 2$), SF3B1 ($n = 0$) and U2AF1 ($n = 1$) mutations compared to patients in the ELN intermediate risk group [39,103].

In summary, 11 patients (65%) harbor dismal prognostic genetic parameters. According to the 2022 ELN genetic risk category, all APA patients who were treated were at adverse risk [104].

The change in prognostic parameters with aza/ven therapy versus chemotherapy indicates a significantly different pathophysiological approach to AML blasts compared to dose-intensive chemotherapy. APA uses one cytotoxic agent (azacitidine at a reduced dose of 45 mg per day from day 1 to day 7 of a 28-day treatment cycle) and two non-hematotoxic transcriptional agents, instead of 400 mg of venetoclax per day from day 1 to day 7 of a 28-day treatment cycle.

These changes to the APA schedule do not merely indicate a substitution of the cytotoxic therapy element, venetoclax, for a targeted therapy. The triple combination of APA comprehensively uncovers molecular-biological changes as novel therapeutic principles via the triple combination's synergistic and biomodulatory interaction. Hidden myeloid functions in AML blasts that guide AML cell death can be re-established, even in patients who have been pre-treated with azacitidine [18,27–29,42,105,106]. Therefore, azacitidine-induced tRNA methylation, which may redirect the translational response to stress, does not appear to be sufficient for coordinating myeloid differentiation with clinical significance [34].

As neutropenic infections are a major cause of mortality in AML patients, they should be avoided through reducing treatment-related hematotoxicity and providing supportive care. [107]. Until now, the reconstitution of myeloid cell functions in acute myeloid leukemia (AML) blasts, and the improvement of immunosurveillance by the antileukemic agent (APA), seemed incompatible with the induction of AML cell death in parallel. Initially observed APA hematotoxicity is primarily related to the AML bone marrow infiltrate and the preceding toxicity of intensive induction treatments (Table 1). Beyond cycles one or two, all subsequent cycles can be administered in an outpatient setting.

12. No Long-Term Continuous Complete Remission Without allo-HSCT Consolidation

Long-term observation of patients treated with aza/ven in the first line, with a median follow-up of 43.2 months, shows a steadily declining survival curve, indicating that a cure may not be achievable for most patients, except those with favorable risk parameters such as NPM1 mutations [108–110].

The same continuous decline in survival curves can be expected for patients treated with APA if they do not qualify for allo-HSCT after treatment. A cure for r/r non-PLM AML was possible in cases of extramedullary leukemic manifestation (skin) or via consolidation with allo-HSCT (n = 3). The patient who received allo-HSCT after APA treatment relapsed before the start of the seventh APA cycle.

APA facilitates CR and, consequently, successful consolidation with allo-HSCT and long-term maintenance of CR or PR, even in patients with TP53 mutations and other poor-risk mutations, as indicated in Table 1.

The advantages of APA therapy are convincing, even though they are based on a small cohort. These advantages include APA activity after azacitidine failure; rescue of poor-risk cytogenetics and molecular genetics with APA; low toxicity of APA in consecutive cycles if CR or PR has been achieved; the ability to generate prerequisites for allo-HSCT in patients who are primarily ineligible; the ability to rescue allo-HSCT failure; and the ability to facilitate long-term stabilization of CR and PR. Last but not least, APA therapy can address the needs of elderly and frail patients.

13. Discussion

In the case of r/r AML, the introduction of low-dose cytotoxic facilitators such as azacitidine is a prerequisite for the successful implementation of targeted transcription modulation in AML blasts and adjacent tissue, and for the concerted reprogramming of AML hallmarks. As demonstrated in clinical trials, these facilitators also include low-dose metronomic chemotherapy and decitabine

[111,112]. As demonstrated in numerous clinical trials, facilitators that enable the targeted modulation of tissue-specific cancer hallmarks can induce stress responses in neoplastic tissue [61]. This creates the prerequisites for the reorganization of neoplasia-associated hallmarks, which is of particular importance in terms of therapy of r/r neoplasias [61,74,113–115]. As demonstrated in the context of APA-like schedules, metronomic low-dose chemotherapy can facilitate genome- and even neoplasia-agnostic schedules [56].

APA omits major non-hematological side effects in normal, non-addicted tissues, thereby highlighting the specificity of the pro-anakoinotic process. The induction of CR in non-PML AML, which is refractory in a genome-agnostic manner, reveals the therapeutic accessibility of a large, well-organized set of non-oncogene addiction targets for achieving pivotal, therapeutically relevant leukemia plasticity. This can be done by reactivating hidden myeloid hallmarks in AML tissue [18,111].

As clinically demonstrated, pioglitazone and ATRA redirected AML tissue-triggered hallmarks comprise differentiation and enhanced immuno surveillance, as well as inflammation control and 'normalized' metabolism. As AML inflammation scores demonstrate high predictive value for outcome, controlling inflammation with pioglitazone, as demonstrated for other r/r neoplasias treated with APA-like schedules, may also be pivotal in controlling r/r AML with APA-like schedules [38,68].

Considering the adverse side effects of current chemotherapies, most targeted therapies or the aza/ven schedule for controlling non-PML AML, differentiation induction combined with the regain of immune competence and inflammation control with APA are contradictory treatment experiences compared with chemotherapy [116].

Phase II/III trials on APA are necessary to establish the APA schedule clinically. However, the presented APA data exemplifies the novel, genome-agnostic therapy principle and the new, systems-relevant correlates that lead to AML cell death (**Figure 2**). Further studies must elucidate the precise molecular mechanisms of APA therapy within a systems context. Meanwhile, pre-clinical data on pioglitazone in AML are continuously adding to our understanding of its activity in neoplasias [22,35,61,117].

The fact that APA treatment has made previously ineligible patients eligible for allo-HSCT is remarkable, given their pre-APA infection history and the response quality induced by APA. [118]. Thus, APA made successful allo-HSCT possible and facilitated cures in high-risk patients, including the elderly.

Targeting non-oncogene addiction networks in a concerted manner using APA-like therapy strategies enables specific adaptation to the pathophysiological requirements of refractory neoplasias and offers novel therapeutic development possibilities [56]. (1) A phase II/III trial should be established to confirm the promising APA results, particularly in light of the novel data showing inflammation-triggered AML pathogenesis [19]. On the other hand, extended experience from clinical trials on inflammation control with pioglitazone in relapsed/refractory neoplasms treated with APA-like schedules reveals the therapeutic relevance of attenuating neoplasia-associated inflammation in order to achieve continuous CR [19,35,56,70]. (2) Furthermore, the differential combined targeting of receptor-triggered transcription factors enables the reprogramming of the pathogenetically most relevant AML hallmarks, with the aim of establishing the most effective genome-agnostic leukemia control. This is achieved by coordinating non-oncogene addiction networks (anakoinosis), which promote the restoration of myeloid hallmarks lost during malignant transformation to AML [18,25,27–29]. (3) Additionally, oncogene-addicted targets may be therapeutically addressed in the context of APA-like schedules. Even myelotoxic drugs can be added due to APA's modest toxicity profile after CR or PR induction [13,56]. (4) The diverse facilitators of effective combined receptor-triggered transcriptional modulation for the control of recurrent and refractory neoplasms reveal novel systems pharmacological activity profiles in a genome- and tumor-agnostic manner [18,56,80]. (5) Proteomic platforms may enable the testing of novel pharmacological

approaches for AML 'normalization' in vitro, as indicated by the therapeutic establishment of myeloid hallmarks [13,56,74,117,119].

Author Contributions: The authors D.H., F.L., and A.R. made substantial contributions to the conception of the review. D.C.H., D.H., F.L., and A.R. contributed to Writing – Review & Editing and to the interpretation of the reviewed data and drafted the work. T.P. revised the work critically. All authors approved the submitted paper and agreed to all aspects of the work, ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. and appropriately documented in the literature.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lancet, J.E.; Uy, G.L.; Cortes, J.E.; Newell, L.F.; Lin, T.L.; Ritchie, E.K.; Stuart, R.K.; Strickland, S.A.; Hogge, D.; Solomon, S.R.; et al. CPX-351 (cytarabine and daunorubicin) Liposome for Injection Versus Conventional Cytarabine Plus Daunorubicin in Older Patients With Newly Diagnosed Secondary Acute Myeloid Leukemia. *J. Clin. Oncol.* **2018**, *36*, 2684–2692, doi:10.1200/JCO.2017.77.6112.
2. Kawamura, K. Allogeneic Hematopoietic Stem Cell Transplantation for Older Patients with Hematological Malignancies. *Yonago Acta Med.* **2025**, *68*, 169–179, doi:10.33160/yam.2025.08.001.
3. Poiré, X.; Labopin, M.; Polge, E.; Blaise, D.; Chevallier, P.; Maertens, J.; Kröger, N.; Besley, C.; Nguyen, S.; Castilla-Llorente, C.; et al. Older matched sibling donor vs young haploidentical donor for older patients with acute myeloid leukemia. *Blood Adv.* **2025**, *9*, 5192–5200, doi:10.1182/bloodadvances.2024015582.
4. Cao, J.; Zhuang, X.; Luo, D.; Pei, R.; Lu, Y.; Chen, D.; Li, S.; Du, X.; Liu, X. Efficacy of Haploidentical Allogeneic Hematopoietic Cell Transplantation following Two Courses of Venetoclax and Azacytidine Therapy in Patients over 55 Years Old with Acute Myelogenous Leukemia. *Acta Haematol.* **2025**, *148*, 427–436, doi:10.1159/000542034.
5. Silva, H.J.; Martin, G.; Birocchi, F.; Wehrli, M.; Kann, M.C.; Supper, V.; Parker, A.; Graham, C.; Bratt, A.; Bouffard, A.; et al. CD70 CAR T cells secreting an anti-CD33/anti-CD3 dual-targeting antibody overcome antigen heterogeneity in AML. *Blood* **2025**, *145*, 720–731, doi:10.1182/blood.2023023210.
6. Haubner, S.; Subklewe, M.; Sadelain, M. Honing CAR T cells to tackle acute myeloid leukemia. *Blood* **2025**, *145*, 1113–1125, doi:10.1182/blood.2024024063.
7. Kantarjian, H.M.; DeAngelo, D.J.; Stelljes, M.; Martinelli, G.; Liedtke, M.; Stock, W.; Gökbuget, N.; O'Brien, S.; Wang, K.; Wang, T.; et al. Inotuzumab Ozogamicin versus Standard Therapy for Acute Lymphoblastic Leukemia. *N. Engl. J. Med.* **2016**, *375*, 740–753, doi:10.1056/NEJMoa1509277.
8. Castro-Muñoz, L.J.; Ulloa, E.V.; Sahlgren, C.; Lizano, M.; La Cruz-Hernández, E. de; Contreras-Paredes, A. Modulating epigenetic modifications for cancer therapy (Review). *Oncol. Rep.* **2023**, *49*, doi:10.3892/or.2023.8496.
9. Sekeres, M.A.; Montesinos, P.; Novak, J.; Wang, J.; Jeyakumar, D.; Tomlinson, B.; Mayer, J.; Jou, E.; Robak, T.; Taussig, D.C.; et al. Glasdegib plus intensive or non-intensive chemotherapy for untreated acute myeloid leukemia: results from the randomized, phase 3 BRIGHT AML 1019 trial. *Leukemia* **2023**, *37*, 2017–2026, doi:10.1038/s41375-023-02001-z.
10. Gao, S.; Wang, X.; Zhao, X.; Xiao, Z. Rational design of next-generation FLT3 inhibitors in acute myeloid leukemia: From laboratory to clinics. *Eur. J. Med. Chem.* **2025**, *301*, 118214, doi:10.1016/j.ejmech.2025.118214.
11. Maschmeyer, G.; Bullinger, L.; Garcia-Vidal, C.; Herbrecht, R.; Maertens, J.; Menna, P.; Pagano, L.; Thiebaut-Bertrand, A.; Calandra, T. Infectious complications of targeted drugs and biotherapies in acute leukemia. Clinical practice guidelines by the European Conference on Infections in Leukemia (ECIL), a joint

- venture of the European Group for Blood and Marrow Transplantation (EBMT), the European Organization for Research and Treatment of Cancer (EORTC), the International Immunocompromised Host Society (ICHS) and the European Leukemia Net (ELN). *Leukemia* **2022**, *36*, 1215–1226, doi:10.1038/s41375-022-01556-7.
12. Issa, G.C.; Stein, E.M.; DiNardo, C.D. How I treat acute myeloid leukemia with differentiation therapy. *Blood* **2025**, *145*, 1251–1259, doi:10.1182/blood.2024024008.
 13. Kantarjian, H.M.; DiNardo, C.D.; Kadia, T.M.; Daver, N.G.; Altman, J.K.; Stein, E.M.; Jabbour, E.; Schiffer, C.A.; Lang, A.; Ravandi, F. Acute myeloid leukemia management and research in 2025. *CA Cancer J. Clin.* **2025**, *75*, 46–67, doi:10.3322/caac.21873.
 14. Wenger, V.; Garcia-Manero, G.; Zeiser, R.; Lübbert, M. DNA methyltransferase inhibitors in hematological malignancies and solid tumors. *Int. J. Cancer* **2025**, doi:10.1002/ijc.70197.
 15. Haumschild, R.; Kennerly-Shah, J.; Barbarotta, L.; Zeidan, A.M. Clinical activity, pharmacokinetics, and pharmacodynamics of oral hypomethylating agents for myelodysplastic syndromes/neoplasms and acute myeloid leukemia: A multidisciplinary review. *J. Oncol. Pharm. Pract.* **2024**, *30*, 721–736, doi:10.1177/10781552241238979.
 16. Short, N.J.; Daver, N.; DiNardo, C.D.; Kadia, T.; Nasr, L.F.; Macaron, W.; Yilmaz, M.; Borthakur, G.; Montalban-Bravo, G.; Garcia-Manero, G.; et al. Azacitidine, Venetoclax, and Gilteritinib in Newly Diagnosed and Relapsed or Refractory FLT3-Mutated AML. *J. Clin. Oncol.* **2024**, *42*, 1499–1508, doi:10.1200/JCO.23.01911.
 17. Stein, E.M.; DiNardo, C.D.; Pollyea, D.A.; Fathi, A.T.; Roboz, G.J.; Altman, J.K.; Stone, R.M.; DeAngelo, D.J.; Levine, R.L.; Flinn, I.W.; et al. Enasidenib in mutant IDH2 relapsed or refractory acute myeloid leukemia. *Blood* **2017**, *130*, 722–731, doi:10.1182/blood-2017-04-779405.
 18. Heudobler, D.; Luke, F.; Hahn, J.; Grube, M.; Schlosser, P.; Kremers, S.; Sudhoff, T.; Westermann, J.; Hutter-Kronke, M.L.; Schlenk, R.F.; et al. Low-dose azacitidine, pioglitazone and all-trans retinoic acid is safe in patients aged ≥ 60 years with acute myeloid leukemia refractory to standard induction chemotherapy (AMLSG 26-16/AML-ViVA): results of the safety run-in phase. *Haematologica* **2024**, *109*, 1274–1278, doi:10.3324/haematol.2023.283864.
 19. Reville, P.K.; Wang, B.; Marvin-Peek, J.; Yuan, B.; Kuo, Y.-A.; Garza, A.; Root, J.; Qiao, W.; Arruda, A.; Veletic, I.; et al. Blood-based proteomic profiling identifies OSMR as a novel biomarker of AML outcomes. *Blood* **2025**, *145*, 3015–3029, doi:10.1182/blood.2024027244.
 20. Hegde, S.; Giotti, B.; Soong, B.Y.; Halasz, L.; Le Berichel, J.; Schaefer, M.M.; Kloeckner, B.; Mattiuz, R.; Park, M.D.; Magen, A.; et al. Myeloid progenitor dysregulation fuels immunosuppressive macrophages in tumours. *Nature* **2025**, *646*, 1214–1222, doi:10.1038/s41586-025-09493-y.
 21. Gozdecka, M.; Dudek, M.; Wen, S.; Gu, M.; Stopforth, R.J.; Rak, J.; Damaskou, A.; Grice, G.L.; McLoughlin, M.A.; Bond, L.; et al. Mitochondrial metabolism sustains DNMT3A-R882-mutant clonal haematopoiesis. *Nature* **2025**, *642*, 431–441, doi:10.1038/s41586-025-08980-6.
 22. Atas, E.; Berchtold, K.; Schleder, M.; Prodinger, S.; Sternberg, F.; Pucci, P.; Steel, C.; Matthews, J.D.; James, E.R.; Philippe, C.; et al. The anti-diabetic PPAR γ agonist Pioglitazone inhibits cell proliferation and induces metabolic reprogramming in prostate cancer. *Mol. Cancer* **2025**, *24*, 134, doi:10.1186/s12943-025-02320-y.
 23. Reichle, A.; Hildebrandt, G.C. Principles of Modular Tumor Therapy. *Cancer Microenvironment* **2009**, *2*, 227–237, doi:10.1007/s12307-009-0023-x.
 24. Meier, R.; Greve, G.; Zimmer, D.; Bresser, H.; Berberich, B.; Langova, R.; Stomper, J.; Rubarth, A.; Feuerbach, L.; Lipka, D.B.; et al. The antileukemic activity of decitabine upon PML/RARA-negative AML blasts is supported by all-trans retinoic acid: in vitro and in vivo evidence for cooperation. *Blood Cancer J.* **2022**, *12*, 122, doi:10.1038/s41408-022-00715-4.
 25. Klobuch, S.; Steinberg, T.; Bruni, E.; Mirbeth, C.; Heilmeier, B.; Ghibelli, L.; Herr, W.; Reichle, A.; Thomas, S. Biomodulatory Treatment With Azacitidine, All-trans Retinoic Acid and Pioglitazone Induces Differentiation of Primary AML Blasts Into Neutrophil Like Cells Capable of ROS Production and Phagocytosis. *Front. Pharmacol.* **2018**, *9*, 1380, doi:10.3389/fphar.2018.01380.
 26. Chen, Y.; Tong, X.; Lu, R.; Zhang, Z.; Ma, T. All-trans retinoic acid in hematologic disorders: not just acute promyelocytic leukemia. *Front. Pharmacol.* **2024**, *15*, 1404092, doi:10.3389/fphar.2024.1404092.

27. Thomas, S.; Schelker, R.; Klobuch, S.; Zaiss, S.; Troppmann, M.; Rehli, M.; Haferlach, T.; Herr, W.; Reichle, A. Biomodulatory therapy induces complete molecular remission in chemorefractory acute myeloid leukemia. *Haematologica* **2015**, *100*, e4–6, doi:10.3324/haematol.2014.115055.
28. Heudobler, D.; Klobuch, S.; Thomas, S.; Hahn, J.; Herr, W.; Reichle, A. Cutaneous Leukemic Infiltrates Successfully Treated With Biomodulatory Therapy in a Rare Case of Therapy-Related High Risk MDS/AML. *Front. Pharmacol.* **2018**, *9*, 1279, doi:10.3389/fphar.2018.01279.
29. Kattner, A.-S.; Holler, E.; Herr, W.; Reichle, A.; Wolff, D.; Heudobler, D. Successful Treatment of Early Relapsed High-Risk AML After Allogeneic Hematopoietic Stem Cell Transplantation With Biomodulatory Therapy. *Front. Oncol.* **2020**, *10*, 443, doi:10.3389/fonc.2020.00443.
30. Wei, A.H.; Loo, S.; Daver, N. How I treat patients with AML using azacitidine and venetoclax. *Blood* **2025**, *145*, 1237–1250, doi:10.1182/blood.2024024009.
31. Lehmann, S.; Paul, C.; Törmä, H. Retinoid receptor expression and its correlation to retinoid sensitivity in non-M3 acute myeloid leukemia blast cells. *Clin. Cancer Res.* **2001**, *7*, 367–373.
32. Yamatani, K.; Watanabe, T.; Saito, K.; Khasawneh, A.; Maiti, A.; Zeng, Z.; Hayes, K.; Kimura, S.; DiNardo, C.D.; Su, X.; et al. PPAR γ -induced upregulation of fatty acid metabolism confers resistance to venetoclax and decitabine therapy in AML. *Blood Neoplasia* **2025**, *2*, 100121, doi:10.1016/j.bneo.2025.100121.
33. Lindblad, K.E.; Goswami, M.; Hourigan, C.S.; Oetjen, K.A. Immunological effects of hypomethylating agents. *Expert Rev. Hematol.* **2017**, *10*, 745–752, doi:10.1080/17474086.2017.1346470.
34. Stein, A.; Platzbecker, U.; Cross, M. How Azanucleosides Affect Myeloid Cell Fate. *Cells* **2022**, *11*, doi:10.3390/cells11162589.
35. Harrer, D.C.; Lücke, F.; Pukrop, T.; Ghibelli, L.; Gerner, C.; Reichle, A.; Heudobler, D. Peroxisome proliferator-activated receptor α/γ agonist pioglitazone for rescuing relapsed or refractory neoplasias by unlocking phenotypic plasticity. *Front. Oncol.* **2023**, *13*, 1289222, doi:10.3389/fonc.2023.1289222.
36. Di Martino, O.; Welch, J.S. Retinoic Acid Receptors in Acute Myeloid Leukemia Therapy. *Cancers (Basel)* **2019**, *11*, doi:10.3390/cancers11121915.
37. Bornhäuser, M.; Schliemann, C.; Schetelig, J.; Röllig, C.; Kramer, M.; Glass, B.; Platzbecker, U.; Burchert, A.; Hänel, M.; Müller, L.P.; et al. Allogeneic Hematopoietic Cell Transplantation vs Standard Consolidation Chemotherapy in Patients With Intermediate-Risk Acute Myeloid Leukemia: A Randomized Clinical Trial. *JAMA Oncol.* **2023**, *9*, 519–526, doi:10.1001/jamaoncol.2022.7605.
38. Shahswar, R.; Gabdoulline, R.; Krueger, K.; Wichmann, M.; Götze, K.S.; Braitsch, K.; Meggendorfer, M.; Schmalbrock, L.; Bullinger, L.; Modemann, F.; et al. A novel prognostic risk model for patients with refractory/relapsed acute myeloid leukemia receiving venetoclax plus hypomethylating agents. *Leukemia* **2025**, *39*, 614–622, doi:10.1038/s41375-024-02501-6.
39. Döhner, H.; Pratz, K.W.; DiNardo, C.D.; Wei, A.H.; Jonas, B.A.; Pullarkat, V.A.; Thirman, M.J.; Récher, C.; Schuh, A.C.; Babu, S.; et al. Genetic risk stratification and outcomes among treatment-naïve patients with AML treated with venetoclax and azacitidine. *Blood* **2024**, *144*, 2211–2222, doi:10.1182/blood.2024024944.
40. Bresser, H.; Schmoor, C.; Grishina, O.; Pfeifer, D.; Thomas, J.; Rehman, U.-U.; Crysandt, M.; Jost, E.; Thol, F.; Heuser, M.; et al. Impact of TP53 Mutation Status in Elderly AML Patients When Adding All-Trans Retinoic Acid or Valproic Acid to Decitabine. *Eur. J. Haematol.* **2025**, *114*, 231–237, doi:10.1111/ejh.14304.
41. Daver, N.; Schlenk, R.F.; Russell, N.H.; Levis, M.J. Targeting FLT3 mutations in AML: review of current knowledge and evidence. *Leukemia* **2019**, *33*, 299–312, doi:10.1038/s41375-018-0357-9.
42. Piccini, M.; Pileri, S.; Merlini, M.; Grieco, P.; Scappini, B.; Bencini, S.; Peruzzi, B.; Caporale, R.; Signori, L.; Pancani, F.; et al. Venetoclax-Based Regimens for Relapsed/Refractory Acute Myeloid Leukemia in a Real-Life Setting: A Retrospective Single-Center Experience. *J. Clin. Med.* **2021**, *10*, doi:10.3390/jcm10081684.
43. Taylor, S.M.; Jones, P.A. Mechanism of action of eukaryotic DNA methyltransferase. Use of 5-azacytosine-containing DNA. *J. Mol. Biol.* **1982**, *162*, 679–692, doi:10.1016/0022-2836(82)90395-3.
44. Zhu, J.; Guérineau, H.; Lefebvre-Fortané, A.-M.; Largeaud, L.; Lambert, J.; Rousselot, P.; Boudouin, M.; Calvo, J.; Prost, S.; Clauser, S.; et al. The AXL inhibitor bemcentinib overcomes microenvironment-mediated resistance to pioglitazone in acute myeloid leukemia. *FEBS J.* **2025**, *292*, 115–128, doi:10.1111/febs.17263.

45. Luo, X.; Zhong, L.; Yu, L.; Xiong, L.; Dan, W.; Li, J.; Ye, J.; Chu, X.; Liu, C.; Liu, B. TRIB3 destabilizes tumor suppressor PPAR α expression through ubiquitin-mediated proteasome degradation in acute myeloid leukemia. *Life Sci.* **2020**, *257*, 118021, doi:10.1016/j.lfs.2020.118021.
46. Ricote, M.; Glass, C.K. PPARs and molecular mechanisms of transrepression. *Biochim. Biophys. Acta* **2007**, *1771*, 926–935, doi:10.1016/j.bbailip.2007.02.013.
47. Zhou, H.; Jiang, Y.; Huang, Y.; Zhong, M.; Qin, D.; Xie, C.; Pan, G.; Tan, J.; Deng, M.; Zhao, H.; et al. Therapeutic inhibition of PPAR α -HIF1 α -PGK1 signaling targets leukemia stem and progenitor cells in acute myeloid leukemia. *Cancer Lett.* **2023**, *554*, 215997, doi:10.1016/j.canlet.2022.215997.
48. Xie, C.; Zhou, H.; Qin, D.; Zheng, H.; Tang, Y.; Li, W.; Zhou, J.; Liu, L.; Yu, X.; Duan, H.; et al. Bcl-2 inhibition combined with PPAR α activation synergistically targets leukemic stem cell-like cells in acute myeloid leukemia. *Cell Death Dis.* **2023**, *14*, 573, doi:10.1038/s41419-023-06075-6.
49. Lübbert, M.; Grishina, O.; Schmoor, C.; Schlenk, R.F.; Jost, E.; Crysandt, M.; Heuser, M.; Thol, F.; Salih, H.R.; Schittenhelm, M.M.; et al. Valproate and Retinoic Acid in Combination With Decitabine in Elderly Nonfit Patients With Acute Myeloid Leukemia: Results of a Multicenter, Randomized, 2 $\times$ 2, Phase II Trial. *J. Clin. Oncol.* **2020**, *38*, 257–270, doi:10.1200/JCO.19.01053.
50. Kucharova, S.; Farkas, R. Hormone nuclear receptors and their ligands: role in programmed cell death (review). *Endocr. Regul.* **2002**, *36*, 37–60.
51. Contractor, R.; Samudio, I.J.; Estrov, Z.; Harris, D.; McCubrey, J.A.; Safe, S.H.; Andreeff, M.; Konopleva, M. A novel ring-substituted diindolylmethane, 1,1-bis(3'-(5-methoxyindolyl)-1-(p-t-butylphenyl) methane, inhibits extracellular signal-regulated kinase activation and induces apoptosis in acute myelogenous leukemia. *Cancer Res.* **2005**, *65*, 2890–2898, doi:10.1158/0008-5472.CAN-04-3781.
52. Heudobler, D.; Lüke, F.; Vogelhuber, M.; Klobuch, S.; Pukrop, T.; Herr, W.; Gerner, C.; Pantziarka, P.; Ghibelli, L.; Reichle, A. Anakoinosis: Correcting Aberrant Homeostasis of Cancer Tissue-Going Beyond Apoptosis Induction. *Front. Oncol.* **2019**, *9*, 1408, doi:10.3389/fonc.2019.01408.
53. Heudobler, D.; Rechenmacher, M.; Lüke, F.; Vogelhuber, M.; Klobuch, S.; Thomas, S.; Pukrop, T.; Hackl, C.; Herr, W.; Ghibelli, L.; et al. Clinical Efficacy of a Novel Therapeutic Principle, Anakoinosis. *Front. Pharmacol.* **2018**, *9*, 1357, doi:10.3389/fphar.2018.01357.
54. Vogt, T.; Coras, B.; Hafner, C.; Landthaler, M.; Reichle, A. Antiangiogenic therapy in metastatic prostate carcinoma complicated by cutaneous lupus erythematosus. *Lancet Oncol.* **2006**, *7*, 695–697, doi:10.1016/S1470-2045(06)70798-7.
55. Nasrabadi, M.E.; Al-Harrasi, A.; Mohammadi, S.; Zarif Azam Kardani, F.; Rahmati, M.; Memarian, A. Pioglitazone as a potential modulator in autoimmune diseases: a review on its effects in systemic lupus erythematosus, psoriasis, inflammatory bowel disease, and multiple sclerosis. *Expert Rev. Clin. Immunol.* **2025**, *21*, 5–15, doi:10.1080/1744666X.2024.2401614.
56. Harrer, D.C.; Lüke, F.; Pukrop, T.; Ghibelli, L.; Reichle, A.; Heudobler, D. Addressing Genetic Tumor Heterogeneity, Post-Therapy Metastatic Spread, Cancer Repopulation, and Development of Acquired Tumor Cell Resistance. *Cancers (Basel)* **2023**, *16*, doi:10.3390/cancers16010180.
57. Ge, Y.; Domschke, C.; Stoiber, N.; Schott, S.; Heil, J.; Rom, J.; Blumenstein, M.; Thum, J.; Sohn, C.; Schneeweiss, A.; et al. Metronomic cyclophosphamide treatment in metastasized breast cancer patients: immunological effects and clinical outcome. *Cancer Immunol. Immunother.* **2012**, *61*, 353–362, doi:10.1007/s00262-011-1106-3.
58. Singh, A.; Chaudhary, R. Potentials of peroxisome proliferator-activated receptor (PPAR) α , β/δ , and γ : An in-depth and comprehensive review of their molecular mechanisms, cellular Signalling, immune responses and therapeutic implications in multiple diseases. *Int. Immunopharmacol.* **2025**, *155*, 114616, doi:10.1016/j.intimp.2025.114616.
59. Luo, N.; Nixon, M.J.; Gonzalez-Ericsson, P.I.; Sanchez, V.; Opalenik, S.R.; Li, H.; Zahnow, C.A.; Nickels, M.L.; Liu, F.; Tantawy, M.N.; et al. DNA methyltransferase inhibition upregulates MHC-I to potentiate cytotoxic T lymphocyte responses in breast cancer. *Nat. Commun.* **2018**, *9*, 248, doi:10.1038/s41467-017-02630-w.

60. Kamel, G.A.M.; Elariny, H.A. Pioglitazone attenuates tamoxifen-induced liver damage in rats via modulating Keap1/Nrf2/HO-1 and SIRT1/Notch1 signaling pathways: In-vivo investigations, and molecular docking analysis. *Mol. Biol. Rep.* **2023**, *50*, 10219–10233, doi:10.1007/s11033-023-08847-x.
61. Vallée, A.; Lecarpentier, Y. Crosstalk Between Peroxisome Proliferator-Activated Receptor Gamma and the Canonical WNT/ β -Catenin Pathway in Chronic Inflammation and Oxidative Stress During Carcinogenesis. *Front. Immunol.* **2018**, *9*, 745, doi:10.3389/fimmu.2018.00745.
62. Zhu, W.; Yan, H.; Li, S.; Nie, W.; Fan, F.; Zhu, J. PPAR- γ agonist pioglitazone regulates dendritic cells immunogenicity mediated by DC-SIGN via the MAPK and NF- κ B pathways. *Int. Immunopharmacol.* **2016**, *41*, 24–34, doi:10.1016/j.intimp.2016.09.028.
63. Molldrem, J.J.; Komanduri, K.; Wieder, E. Overexpressed differentiation antigens as targets of graft-versus-leukemia reactions. *Curr. Opin. Hematol.* **2002**, *9*, 503–508, doi:10.1097/00062752-200211000-00006.
64. Scheibenbogen, C.; Letsch, A.; Thiel, E.; Schmitt, A.; Mailaender, V.; Baerwolf, S.; Nagorsen, D.; Keilholz, U. CD8 T-cell responses to Wilms tumor gene product WT1 and proteinase 3 in patients with acute myeloid leukemia. *Blood* **2002**, *100*, 2132–2137, doi:10.1182/blood-2002-01-0163.
65. Hu, L.; Zhang, X.; Zhang, W.; Jin, S.; Zhao, J.; Zheng, J.; Song, W.; Shen, Z. Targeting TCMR-associated cytokine genes for drug screening identifies PPAR γ agonists as novel immunomodulatory agents in transplantation. *Front. Immunol.* **2025**, *16*, 1539645, doi:10.3389/fimmu.2025.1539645.
66. Chiappinelli, K.B.; Strissel, P.L.; Desrichard, A.; Li, H.; Henke, C.; Akman, B.; Hein, A.; Rote, N.S.; Cope, L.M.; Snyder, A.; et al. Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* **2015**, *162*, 974–986, doi:10.1016/j.cell.2015.07.011.
67. Montaigne, D.; Butruille, L.; Staels, B. PPAR control of metabolism and cardiovascular functions. *Nat. Rev. Cardiol.* **2021**, *18*, 809–823, doi:10.1038/s41569-021-00569-6.
68. Lasry, A.; Nadorp, B.; Fornerod, M.; Nicolet, D.; Wu, H.; Walker, C.J.; Sun, Z.; Witkowski, M.T.; Tikhonova, A.N.; Guillaumot-Ruano, M.; et al. An inflammatory state remodels the immune microenvironment and improves risk stratification in acute myeloid leukemia. *Nat. Cancer* **2023**, *4*, 27–42, doi:10.1038/s43018-022-00480-0.
69. Goleij, P.; Khazeei Tabari, M.A.; Ahmed, A.R.D.; Mohamed, L.M.E.; Saleh, G.A.H.; Abdu Hassan, M.T.M.; Moahammednoor, A.G.M.; Khan, H. Molecular Secrets Revealed: How Diabetes may be Paving the Way for Leukemia. *Curr. Treat. Options Oncol.* **2024**, *25*, 1563–1579, doi:10.1007/s11864-024-01281-6.
70. Lüke, F.; Harrer, D.C.; Pantziarka, P.; Pukrop, T.; Ghibelli, L.; Gerner, C.; Reichle, A.; Heudobler, D. Drug Repurposing by Tumor Tissue Editing. *Front. Oncol.* **2022**, *12*, 900985, doi:10.3389/fonc.2022.900985.
71. Prost, S.; Relouzat, F.; Spentchian, M.; Ouzegdough, Y.; Saliba, J.; Massonnet, G.; Beressi, J.-P.; Verhoeven, E.; Raggueneau, V.; Maneglier, B.; et al. Erosion of the chronic myeloid leukaemia stem cell pool by PPAR γ agonists. *Nature* **2015**, *525*, 380–383, doi:10.1038/nature15248.
72. Ben-Yishay, R.; Globus, O.; Balint-Lahat, N.; Arbili-Yarhi, S.; Bar-Hai, N.; Bar, V.; Aharon, S.; Kosenko, A.; Zundevich, A.; Berger, R.; et al. Class Effect Unveiled: PPAR γ Agonists and MEK Inhibitors in Cancer Cell Differentiation. *Cells* **2024**, *13*, doi:10.3390/cells13171506.
73. To, K.K.W.; Wu, W.K.K.; Loong, H.H.F. PPARgamma agonists sensitize PTEN-deficient resistant lung cancer cells to EGFR tyrosine kinase inhibitors by inducing autophagy. *Eur. J. Pharmacol.* **2018**, *823*, 19–26, doi:10.1016/j.ejphar.2018.01.036.
74. Harrer, D.C.; Lüke, F.; Pukrop, T.; Ghibelli, L.; Reichle, A.; Heudobler, D. MEPED as salvage therapy for relapsed/refractory Hodgkin's lymphoma incorporating edited non-oncogene addiction: mTOR as a bottleneck. *Front. Pharmacol.* **2025**, *16*, 1553331, doi:10.3389/fphar.2025.1553331.
75. Vunta, H.; Davis, F.; Palempalli, U.D.; Bhat, D.; Arner, R.J.; Thompson, J.T.; Peterson, D.G.; Reddy, C.C.; Prabhu, K.S. The anti-inflammatory effects of selenium are mediated through 15-deoxy-Delta12,14-prostaglandin J2 in macrophages. *J. Biol. Chem.* **2007**, *282*, 17964–17973, doi:10.1074/jbc.M703075200.
76. Sánchez-Mendoza, S.E.; Deus-Wagatsuma, V.M. de; do Nascimento, M.C.; Lima, K.; Machado-Neto, J.A.; Djavaheri-Mergny, M.; Rego, E.M. All-trans retinoic acid potentiates cell death induced by quizartinib in acute myeloid leukemia with FLT3-ITD mutations. *Ann. Hematol.* **2024**, *103*, 5405–5416, doi:10.1007/s00277-024-06089-w.

77. Lüke, F.; Harrer, D.C.; Menhart, K.; Wolff, D.; Holler, E.; Hellwig, D.; Herr, W.; Grube, M.; Vogelhuber, M.; Reichle, A.; et al. Biomodulatory Treatment Regimen, MEPED, Rescues Relapsed and Refractory Classic Hodgkin's Disease. *Front. Pharmacol.* **2021**, *12*, 599561, doi:10.3389/fphar.2021.599561.
78. Harrer, D.C.; Jakob, M.; Vogelhuber, M.; Lüke, F.; Utpatel, K.; Corbacioglu, S.; Herr, W.; Reichle, A.; Heudobler, D. Biomodulatory therapy induces durable remissions in multi-system Langerhans cell histiocytosis. *Leuk. Lymphoma* **2022**, *63*, 2858–2868, doi:10.1080/10428194.2022.2095627.
79. Reichle, A.; Grassinger, J.; Bross, K.; Wilke, J.; Suedhoff, T.; Walter, B.; Wieland, W.-F.; Berand, A.; Andreesen, R. C-reactive Protein in Patients with Metastatic Clear Cell Renal Carcinoma: An Important Biomarker for Tumor-associated Inflammation. *Biomark. Insights* **2007**, *1*, 87–98.
80. Plumber, S.A.; Tate, T.; Al-Ahmadie, H.; Chen, X.; Choi, W.; Basar, M.; Lu, C.; Viny, A.; Batourina, E.; Li, J.; et al. Rosiglitazone and trametinib exhibit potent anti-tumor activity in a mouse model of muscle invasive bladder cancer. *Nat. Commun.* **2024**, *15*, 6538, doi:10.1038/s41467-024-50678-2.
81. Wouters, B.J.; Delwel, R. Epigenetics and approaches to targeted epigenetic therapy in acute myeloid leukemia. *Blood* **2016**, *127*, 42–52, doi:10.1182/blood-2015-07-604512.
82. Mulet-Lazaro, R.; van Herk, S.; Nuetzel, M.; Sijs-Szabo, A.; Díaz, N.; Kelly, K.; Erpelinck-Verschueren, C.; Schwarzfischer-Pfeilschifter, L.; Stanewsky, H.; Ackermann, U.; et al. Epigenetic alterations affecting hematopoietic regulatory networks as drivers of mixed myeloid/lymphoid leukemia. *Nat. Commun.* **2024**, *15*, 5693, doi:10.1038/s41467-024-49811-y.
83. Esmaeili, S.; Safaroghli-Azar, A.; Pourbagheri-Sigaroodi, A.; Salari, S.; Gharehbaghian, A.; Hamidpour, M.; Bashash, D. Stimulation of Peroxisome Proliferator-Activated Receptor-Gamma (PPAR γ) using Pioglitazone Decreases the Survival of Acute Promyelocytic Leukemia Cells through Up-Regulation of PTEN Expression. *Anticancer. Agents Med. Chem.* **2021**, *21*, 108–119, doi:10.2174/1871520620666200817101533.
84. Symeonidou, V.; Metzner, M.; Usukhbayar, B.; Jackson, A.E.; Fox, S.; Craddock, C.F.; Vyas, P. Heterogeneous genetic and non-genetic mechanisms contribute to response and resistance to azacitidine monotherapy. *EJHaem* **2022**, *3*, 794–803, doi:10.1002/jha2.527.
85. Bogenberger, J.M.; Kornblau, S.M.; Pierceall, W.E.; Lena, R.; Chow, D.; Shi, C.-X.; Mantei, J.; Ahmann, G.; Gonzales, I.M.; Choudhary, A.; et al. BCL-2 family proteins as 5-Azacitidine-sensitizing targets and determinants of response in myeloid malignancies. *Leukemia* **2014**, *28*, 1657–1665, doi:10.1038/leu.2014.44.
86. Jin, S.; Cojocari, D.; Purkal, J.J.; Popovic, R.; Talaty, N.N.; Xiao, Y.; Solomon, L.R.; Boghaert, E.R.; Levenson, J.D.; Phillips, D.C. 5-Azacitidine Induces NOXA to Prime AML Cells for Venetoclax-Mediated Apoptosis. *Clin. Cancer Res.* **2020**, *26*, 3371–3383, doi:10.1158/1078-0432.CCR-19-1900.
87. Esmaeili, S.; Salari, S.; Kaveh, V.; Ghaffari, S.H.; Bashash, D. Alteration of PPAR-GAMMA (PPARG; PPAR γ) and PTEN gene expression in acute myeloid leukemia patients and the promising anticancer effects of PPAR γ stimulation using pioglitazone on AML cells. *Mol. Genet. Genomic Med.* **2021**, *9*, e1818, doi:10.1002/mgg3.1818.
88. Yasugi, E.; Horiuchi, A.; Uemura, I.; Okuma, E.; Nakatsu, M.; Saeki, K.; Kamisaka, Y.; Kagechika, H.; Yasuda, K.; Yuo, A. Peroxisome proliferator-activated receptor gamma ligands stimulate myeloid differentiation and lipogenesis in human leukemia NB4 cells. *Dev. Growth Differ.* **2006**, *48*, 177–188, doi:10.1111/j.1440-169X.2006.00855.x.
89. Zang, C.; Wächter, M.; Liu, H.; Posch, M.G.; Fenner, M.H.; Stadelmann, C.; Deimling, A. von; Possinger, K.; Black, K.L.; Koeffler, H.P.; et al. Ligands for PPARgamma and RAR cause induction of growth inhibition and apoptosis in human glioblastomas. *J. Neurooncol.* **2003**, *65*, 107–118, doi:10.1023/b:neon.0000003728.80052.a8.
90. Liu, H.; Zang, C.; Fenner, M.H.; Possinger, K.; Elstner, E. PPARgamma ligands and ATRA inhibit the invasion of human breast cancer cells in vitro. *Breast Cancer Res. Treat.* **2003**, *79*, 63–74, doi:10.1023/a:1023366117157.
91. Liccardo, F.; Śniegocka, M.; Tito, C.; Iaiza, A.; Ottone, T.; Divona, M.; Travaglini, S.; Mattei, M.; Cicconi, R.; Miglietta, S.; et al. Retinoic acid and proteotoxic stress induce AML cell death overcoming stromal cell protection. *J. Exp. Clin. Cancer Res.* **2023**, *42*, 223, doi:10.1186/s13046-023-02793-z.

92. Li, B.; Zheng, L.; Yang, J.; Qu, L. Targeting oncogene-induced cellular plasticity for tumor therapy. *Adv. Biotechnol. (Singap)* **2024**, *2*, 24, doi:10.1007/s44307-024-00030-y.
93. Fu, S.; He, K.; Tian, C.; Sun, H.; Zhu, C.; Bai, S.; Liu, J.; Wu, Q.; Di Xie; Yue, T.; et al. Impaired lipid biosynthesis hinders anti-tumor efficacy of intratumoral iNKT cells. *Nat. Commun.* **2020**, *11*, 438, doi:10.1038/s41467-020-14332-x.
94. Reichle, A.; Vogt, T. Systems biology: a therapeutic target for tumor therapy. *Cancer Microenvironment* **2008**, *1*, 159–170, doi:10.1007/s12307-008-0012-5.
95. Huls, G.; Woolthuis, C.M.; Schuringa, J.J. Menin inhibitors in the treatment of acute myeloid leukemia. *Blood* **2025**, *145*, 561–566, doi:10.1182/blood.2024026232.
96. Issa, G.C.; Aldoss, I.; DiPersio, J.; Cuglievan, B.; Stone, R.; Arellano, M.; Thirman, M.J.; Patel, M.R.; Dickens, D.S.; Shenoy, S.; et al. The menin inhibitor revumenib in KMT2A-rearranged or NPM1-mutant leukaemia. *Nature* **2023**, *615*, 920–924, doi:10.1038/s41586-023-05812-3.
97. Halik, A.; Tilgner, M.; Silva, P.; Estrada, N.; Altwasser, R.; Jahn, E.; Heuser, M.; Hou, H.-A.; Pratcorona, M.; Hills, R.K.; et al. Genomic characterization of AML with aberrations of chromosome 7: a multinational cohort of 519 patients. *J. Hematol. Oncol.* **2024**, *17*, 70, doi:10.1186/s13045-024-01590-1.
98. Gottfried, E.; Rogenhofer, S.; Waibel, H.; Kunz-Schughart, L.A.; Reichle, A.; Wehrstein, M.; Peuker, A.; Peter, K.; Hartmannsgruber, G.; Andreesen, R.; et al. Pioglitazone modulates tumor cell metabolism and proliferation in multicellular tumor spheroids. *Cancer Chemother. Pharmacol.* **2011**, *67*, 117–126, doi:10.1007/s00280-010-1294-0.
99. Wattad, M.; Weber, D.; Döhner, K.; Krauter, J.; Gaidzik, V.I.; Paschka, P.; Heuser, M.; Thol, F.; Kindler, T.; Lübbert, M.; et al. Impact of salvage regimens on response and overall survival in acute myeloid leukemia with induction failure. *Leukemia* **2017**, *31*, 1306–1313, doi:10.1038/leu.2017.23.
100. Thol, F.; Schlenk, R.F.; Heuser, M.; Ganser, A. How I treat refractory and early relapsed acute myeloid leukemia. *Blood* **2015**, *126*, 319–327, doi:10.1182/blood-2014-10-551911.
101. Westermann, J.; Bullinger, L. Precision medicine in myeloid malignancies. *Semin. Cancer Biol.* **2022**, *84*, 153–169, doi:10.1016/j.semcancer.2021.03.034.
102. DiNardo, C.D.; Jonas, B.A.; Pullarkat, V.; Thirman, M.J.; Garcia, J.S.; Wei, A.H.; Konopleva, M.; Döhner, H.; Letai, A.; Fenaux, P.; et al. Azacitidine and Venetoclax in Previously Untreated Acute Myeloid Leukemia. *N. Engl. J. Med.* **2020**, *383*, 617–629, doi:10.1056/NEJMoa2012971.
103. Bill, M.; Eckardt, J.-N.; Döhner, K.; Röhnert, M.-A.; Rausch, C.; Metzeler, K.H.; Spiekermann, K.; Stasik, S.; Wurm, A.A.; Sauer, T.; et al. Differential prognostic impact of myelodysplasia-related gene mutations in a European cohort of 4978 intensively treated AML patients. *Leukemia* **2025**, doi:10.1038/s41375-025-02781-6.
104. Döhner, H.; Wei, A.H.; Appelbaum, F.R.; Craddock, C.; DiNardo, C.D.; Dombret, H.; Ebert, B.L.; Fenaux, P.; Godley, L.A.; Hasserjian, R.P.; et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood* **2022**, *140*, 1345–1377, doi:10.1182/blood.2022016867.
105. Labrador, J.; Saiz-Rodríguez, M.; Miguel, D. de; Laiglesia, A. de; Rodríguez-Medina, C.; Vidriales, M.B.; Pérez-Encinas, M.; Sánchez-Sánchez, M.J.; Cuello, R.; Roldán-Pérez, A.; et al. Use of Venetoclax in Patients with Relapsed or Refractory Acute Myeloid Leukemia: The PETHEMA Registry Experience. *Cancers (Basel)* **2022**, *14*, doi:10.3390/cancers14071734.
106. DiNardo, C.D.; Maiti, A.; Rausch, C.R.; Pemmaraju, N.; Naqvi, K.; Daver, N.G.; Kadia, T.M.; Borthakur, G.; Ohanian, M.; Alvarado, Y.; et al. 10-day decitabine with venetoclax for newly diagnosed intensive chemotherapy ineligible, and relapsed or refractory acute myeloid leukaemia: a single-centre, phase 2 trial. *Lancet Haematol.* **2020**, *7*, e724–e736, doi:10.1016/S2352-3026(20)30210-6.
107. DiNardo, C.D.; Jen, W.-Y.; Takahashi, K.; Kadia, T.M.; Loghavi, S.; Daver, N.G.; Xiao, L.; Reville, P.K.; Issa, G.C.; Short, N.J.; et al. Long term results of venetoclax combined with FLAG-IDA induction and consolidation for newly diagnosed and relapsed or refractory acute myeloid leukemia. *Leukemia* **2025**, *39*, 854–863, doi:10.1038/s41375-025-02531-8.
108. Maiti, A.; Rausch, C.R.; Cortes, J.E.; Pemmaraju, N.; Daver, N.G.; Ravandi, F.; Garcia-Manero, G.; Borthakur, G.; Naqvi, K.; Ohanian, M.; et al. Outcomes of relapsed or refractory acute myeloid leukemia after frontline

- hypomethylating agent and venetoclax regimens. *Haematologica* **2021**, *106*, 894–898, doi:10.3324/haematol.2020.252569.
109. DiNardo, C.D.; Tiong, I.S.; Quaglieri, A.; MacRaild, S.; Loghavi, S.; Brown, F.C.; Thijssen, R.; Pomilio, G.; Ivey, A.; Salmon, J.M.; et al. Molecular patterns of response and treatment failure after frontline venetoclax combinations in older patients with AML. *Blood* **2020**, *135*, 791–803, doi:10.1182/blood.2019003988.
 110. Müller-Tidow, C.; Tschanter, P.; Röllig, C.; Thiede, C.; Koschmieder, A.; Stelljes, M.; Koschmieder, S.; Dugas, M.; Gerss, J.; Butterfaß-Bahloul, T.; et al. Azacitidine in combination with intensive induction chemotherapy in older patients with acute myeloid leukemia: The AML-AZA trial of the Study Alliance Leukemia. *Leukemia* **2016**, *30*, 555–561, doi:10.1038/leu.2015.306.
 111. Thomas, J.; Rehman, U.-U.; Bresser, H.; Grishina, O.; Pfeifer, D.; Sollier, E.; Döhner, K.; Plass, C.; Becker, H.; Schmoor, C.; et al. Continued decitabine/all-trans retinoic acid treatment: extended complete remission in an elderly AML patient with multi-hit TP53 lesions and complex-monosomal karyotype. *Clin. Epigenetics* **2024**, *16*, 126, doi:10.1186/s13148-024-01737-4.
 112. Wang, L.; Zhang, Q.; Ye, L.; Ye, X.; Yang, W.; Zhang, H.; Zhou, X.; Ren, Y.; Ma, L.; Zhang, X.; et al. All-trans retinoic acid enhances the cytotoxic effect of decitabine on myelodysplastic syndromes and acute myeloid leukaemia by activating the RAR α -Nrf2 complex. *Br. J. Cancer* **2023**, *128*, 691–701, doi:10.1038/s41416-022-02074-0.
 113. Gehrmann, M.; Brunner, M.; Pfister, K.; Reichle, A.; Kremmer, E.; Multhoff, G. Differential up-regulation of cytosolic and membrane-bound heat shock protein 70 in tumor cells by anti-inflammatory drugs. *Clin. Cancer Res.* **2004**, *10*, 3354–3364, doi:10.1158/1078-0432.CCR-03-0382.
 114. Doloff, J.C.; Waxman, D.J. Transcriptional profiling provides insights into metronomic cyclophosphamide-activated, innate immune-dependent regression of brain tumor xenografts. *BMC Cancer* **2015**, *15*, 375, doi:10.1186/s12885-015-1358-y.
 115. Stelmach, P.; Trumpp, A. Leukemic stem cells and therapy resistance in acute myeloid leukemia. *Haematologica* **2023**, *108*, 353–366, doi:10.3324/haematol.2022.280800.
 116. Feld, J.; Tremblay, D.; Dougherty, M.; Czaplinska, T.; Sanchez, G.; Brady, C.; Kremyanskaya, M.; Bar-Natan, M.; Keyzner, A.; Marcellino, B.K.; et al. Safety and Efficacy: Clinical Experience of Venetoclax in Combination With Hypomethylating Agents in Both Newly Diagnosed and Relapsed/Refractory Advanced Myeloid Malignancies. *Hemasphere* **2021**, *5*, e549, doi:10.1097/HS9.0000000000000549.
 117. Heudobler, D.; Reichle, A.; Ghibelli, L. Editorial: Anakoinosis: An Innovative Anticancer Therapy Targeting the Aberrant Cancer Tissue Homeostasis. *Front. Pharmacol.* **2021**, *12*, 779021, doi:10.3389/fphar.2021.779021.
 118. Loke, J.; Buka, R.; Craddock, C. Allogeneic Stem Cell Transplantation for Acute Myeloid Leukemia: Who, When, and How? *Front. Immunol.* **2021**, *12*, 659595, doi:10.3389/fimmu.2021.659595.
 119. Meier-Menches, S.M.; Neuditschko, B.; Janker, L.; Gerner, M.C.; Schmetterer, K.G.; Reichle, A.; Gerner, C. A Proteomic Platform Enables to Test for AML Normalization In Vitro. *Front. Chem.* **2022**, *10*, 826346, doi:10.3389/fchem.2022.826346.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.