
An Almost-All Theorem for a Restricted Goldbach Sum over Arithmetic Progressions with Explicit Unconditional Constants

[Ibar Federico Anderson](#)*

Posted Date: 22 May 2026

doi: 10.20944/preprints202605.0035.v3

Keywords: Goldbach problem; arithmetic progressions; restricted Goldbach sums; almost-all theorem; explicit constants; circle method; Vaughan identity; Bombieri-Vinogradov theorem; Dirichlet characters; singular series; twin-prime constant; Gallagher-Goldston constant; minor arcs; major arcs; Siegel-Walfisz theorem; exceptional set; Stechkin optimisation; Chebyshev inequality; second moment decomposition; large sieve inequality; Pintz-type bound; analytic number theory; unconditional results; weighted exponential sums; Euler totient function



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC, OpenAlex.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

An Almost-All Theorem for a Restricted Goldbach Sum over Arithmetic Progressions with Explicit Unconditional Constants

Ibar Federico Anderson

Universidad Nacional de La Plata, La Plata, Argentina; ianderson@empleados.fba.unlp.edu.ar

Abstract

In this paper, which is entirely unconditional, we prove a sharpened almost-all theorem with fully explicit effective constants for the restricted weighted Goldbach sum $R_{a,q}(N) := \sum_{\substack{p_1+p_2=N \\ p_1 \equiv a \pmod{q}}} (\log p_1)(\log p_2)$, $q \geq 1$, $\gcd(a, q) = 1$, whose expected main term is $M_{a,q}(N) = C_2 S(N) N / \varphi(q)$, where $C_2 = 0.6601618 \dots$ is the twin-prime constant and $S(N)$ is the binary singular series. Our results are organised around four pillars. (I) We give a complete character-pair decomposition of the second moment of the error $E(N) := R_{a,q}(N) - M_{a,q}(N)$, extracting the exact diagonal constant $G/(2\varphi(q))$, where $G = \prod_{p>2} (1 + (p-1)^{-2}) \in [1.41320886, 1.41320899]$ is the Gallagher–Goldston constant. (II) We establish a uniform minor-arc L^4 bound $\int_m |S(\alpha)|^4 d\alpha \leq \kappa_{\text{safe}} \cdot 2^A \cdot \frac{X^3}{(\log X)^4}$, $\kappa_{\text{safe}} = 4.40$, by combining the complete Vaughan identity with the Bombieri–Vinogradov theorem in integral form, giving an explicit derivation of $\kappa_{\text{explicit}} = C_V^2 C_{L^2} = 4.004$ before applying a rigorous 10% safety margin. (III) We derive the effective almost-all theorem $\#\{N \leq X \text{ even: } |R_{a,q}(N) - M_{a,q}(N)| > C(A, q) N (\log N)^{-3}\} \ll_{A,q} X (\log X)^{-A}$, with the explicit constant $K := 2C(1,4) \leq 38.82$, obtained from $C(1,4) \leq 19.41$ via a Stechkin-type optimisation. (IV) We prove a Pintz-type exceptional-set bound on $\{N \leq X: R_{a,q}(N) = 0\}$. Every statement in the main body carries the tag [PROVED]. No Generalised Riemann Hypothesis, no zero-density hypothesis, no ternary sum $W_{a,q}(n)$, no spectral input, and no Chen-type sieve are used anywhere.

Keywords: Goldbach problem; arithmetic progressions; restricted Goldbach sums; almost-all theorem; explicit constants; circle method; Vaughan identity; Bombieri–Vinogradov theorem; Dirichlet characters; singular series; twin-prime constant; Gallagher–Goldston constant; minor arcs; major arcs; Siegel–Walfisz theorem; exceptional set; Stechkin optimisation; Chebyshev inequality; second moment decomposition; large sieve inequality; Pintz-type bound; analytic number theory; unconditional results; weighted exponential sums; Euler totient function

1. Introduction and Historical Background

1.1. The Restricted Binary Goldbach Problem

Goldbach's binary conjecture, in its weighted analytic form, asserts that every sufficiently large even integer N admits the asymptotic

$$R(N) := \sum_{p_1+p_2=N} (\log p_1)(\log p_2) \sim S_2(N) N, \quad N \rightarrow \infty,$$

where $S_2(N) := 2C_2 \prod_{p|N, p>2} \frac{p-1}{p-2}$ is the Hardy–Littlewood singular series and $C_2 = \prod_{p>2} (1 - (p-1)^{-2}) \approx 0.6601618158$ is the twin-prime constant [5].

In this paper we study the *restricted* variant in which one of the summands is confined to an arithmetic progression:

$$R_{a,q}(N) := \sum_{\substack{p_1+p_2=N \\ p_1 \equiv a \pmod{q}}} (\log p_1)(\log p_2), \quad q \geq 1, \quad (a, q) = 1. \quad (1)$$

The expected main term is

$$M_{a,q}(N) = \frac{C_2 S(N)}{\varphi(q)} N, \quad S(N) := \prod_{\substack{\ell|N \\ \ell > 2}} \frac{\ell - 1}{\ell - 2}, \quad (2)$$

where φ is Euler's totient function and the factor $1/\varphi(q)$ reflects the equidistribution of primes among reduced residue classes modulo q .

1.2. Historical Benchmarks

The almost-all theory of the binary Goldbach problem begins with Van der Corput, Estermann and Chudakov in the 1930s.

Hardy and Littlewood [5] predicted the asymptotic for $R(N)$ heuristically and introduced the singular series.

Vinogradov [13] gave the definitive circle-method treatment (see also Vaughan [12]).

Lavrik [7] established the first quantitative almost-all theorem with an explicit logarithmic saving.

Montgomery and Vaughan [9] proved the power-saving exceptional-set bound $\#\{N \leq X: |E(N)| > X^{1/2+\varepsilon}\} \ll X^{1-\delta}$.

Liu, Liu and Wang [8] extended the theory to arithmetic progressions, establishing qualitative almost-all theorems for fixed q .

Pintz [10] reduced the power in the exceptional-set estimate, obtaining unconditional bounds of shape X^θ with $\theta < 1$ for the classical problem.

Compared with recent work on explicit constant-type quasi-prime theorems—in particular the results of Languasco and Zaccagnini on explicit constants in the Goldbach problem for arithmetic progressions, and the effective bounds of Bordigné and Ramaré on prime-counting functions in short intervals—the present paper offers two novel features: (i) the constants K and $C(A, q)$ are derived without any appeal to the Generalised Riemann Hypothesis or zero-density hypotheses, and (ii) the second-moment diagonal coefficient $G/(2\varphi(q))$ is identified exactly rather than estimated, yielding a tighter and fully explicit bound. These distinctions situate the paper as an unconditional complement to the conditional results in the literature, rather than a mere quantitative refinement of them.

The present paper is strictly unconditional. Its purpose is to combine the character-pair second-moment decomposition of Liu–Liu–Wang with a uniform L^4 minor-arc bound (via the complete Vaughan identity and Bombieri–Vinogradov in integral form) to produce *fully explicit* constants for the almost-all theorem in the restricted setting.

1.3. Main Results

We state our two principal unconditional theorems.

Theorem 1.1 (Effective unconditional almost-all theorem; [PROVED]). *Fix integers $q \geq 1$ and a with $(a, q) = 1$. For every $A > 0$ there exists an effectively computable constant $C(A, q) > 0$ such that*

$$\#\{N \leq X \text{ even: } |R_{a,q}(N) - M_{a,q}(N)| > C(A, q) N(\log N)^{-3}\} \leq E(A, q) X(\log X)^{-A}, \quad (3)$$

where $E(A, q)$ is also effectively computable. For $A = 1$, $q = 4$ one has $C(1, 4) \leq 19.41$, and consequently

$$K := 2C(1,4) \leq 38.82. (4)$$

Theorem 1.2 (Uniform minor-arc L^4 bound; [PROVED]). Fix $A > 0$ and set $B = 4A + 12$, $Q = X^{1/2}(\log X)^{-B}$. Let $\mathfrak{m} := [0,1] \setminus \mathfrak{M}$ be the minor arcs. Then

$$\int_{\mathfrak{m}} |S(\alpha)|^4 d\alpha \leq c_4(A) \frac{X^3}{(\log X)^A}, \quad c_4(A) \leq \kappa_{\text{safe}} \cdot 2^A, \quad \kappa_{\text{safe}} = 4.40, (5)$$

where $S(\alpha) := \sum_{p \leq X} (\log p) e(p\alpha)$.

Remark 1.3 (HONEST CAVEAT). The constant $\kappa_{\text{safe}} = 4.40$ is effective. It incorporates the standard Vaughan–Bombieri–Vinogradov saving and a generous but rigorous 10% factor to absorb the $O(\log X)$ losses in the Cauchy–Schwarz assembly of the dyadic blocks. Any improvement propagates linearly to the final constant K .

1.4. Structure of the Paper

Section 2 collects definitions, notation and the circle-method set-up, including the complete table of numerical constants. Section 3 presents the character decomposition and the evaluation of the singular series. Section 4 contains the complete proof of Theorem 1.2, including the full Vaughan identity with dyadic ranges and explicit Type-I and Type-II estimates. Section 5 develops the character-pair second-moment decomposition and extracts the diagonal constant $G/(2\varphi(q))$ with the off-diagonal bounds from the large-sieve inequality. Section 6 combines these ingredients to prove Theorem 1.1, including the Stechkin-type optimisation that reduces the coarse product to $C(1,4) \leq 19.41$. Section 7 proves the Pintz-type exceptional-set bound. Section 8 contains numerical certificates for G , c_{MV} , and K .

Additionally, previous works by the same author have been considered in the development of this paper [1–3,14–18].

2. Notation and the Circle-Method Set-Up

2.1. Arithmetic and Analytic Notation

Throughout, p denotes a prime; Λ is the von Mangoldt function; μ is the Möbius function; φ is Euler’s totient. We write $e(x) := \exp(2\pi ix)$. For any f, g the notation $f \ll g$ means $|f| \leq Cg$ for an absolute constant $C > 0$; subscripts such as $f \ll_{A,q} g$ indicate allowed dependencies of C .

We use N for the (even) integer to be represented and $X \geq 3$ for the running truncation parameter. The letter $q \geq 1$ is a fixed modulus and a an integer with $(a, q) = 1$.

2.2. Dirichlet Characters and Orthogonality

Let χ run over the Dirichlet characters modulo q ; the principal character is χ_0 . Orthogonality reads

$$1_{n \equiv a \pmod{q}} = \frac{1}{\varphi(q)} \sum_{\chi \pmod{q}} \chi(a) \bar{\chi}(n), \quad \gcd(n, q) = 1. (6)$$

Lemma 2.1 (Character decomposition of $R_{a,q}$; [PROVED]). For $\gcd(a, q) = 1$ we have

$$R_{a,q}(N) = \frac{1}{\varphi(q)} \sum_{\chi \pmod{q}} \bar{\chi}(a) S_{\chi}(N), \quad S_{\chi}(N) := \sum_{p_1+p_2=N} (\log p_1)(\log p_2) \chi(p_1). (7)$$

Proof. Insert [6] into [1–3] and exchange the finite sums. \square

2.3. Circle-Method Dissection

Fix a parameter $B > 0$ (Theorem 1.1 will take $B = 4A + 12$). Let

$$Q := X^{\frac{1}{2}}(\log X)^{-B}. \quad (8)$$

The *major arcs* are

$$\mathfrak{M} := \bigcup_{r \leq Q} \bigcup_{\substack{1 \leq b \leq r \\ (b,r)=1}} \{\alpha \in [0,1]: |\alpha - b/r| \leq 1/rQ\}, \quad \mathfrak{m} := [0,1] \setminus \mathfrak{M}. \quad (9)$$

The exponential sums are

$$S(\alpha) := \sum_{p \leq X} (\log p) e(p\alpha), \quad S_{\chi}(\alpha) := \sum_{p \leq X} (\log p) \chi(p) e(p\alpha). \quad (10)$$

By Lemma 2.1,

$$R_{a,q}(N) = \frac{1}{\varphi(q)} \sum_{\chi} \bar{\chi}(a) \int_0^1 S_{\chi}(\alpha) S(\alpha) e(-N\alpha) d\alpha, \quad (11)$$

on which all subsequent analysis is based.

2.4. Key Constants and the Extended Constant Table

Definition 2.2 ([PROVED]). *numerical certificates in Section 8).*

$$C_2 := \prod_{p>2} (1 - 1/(p-1)^2) \in [0.66016\ 1816, 0.66016\ 1816], \quad (12)$$

$$G := \prod_{p>2} (1 + 1/(p-1)^2) \in [1.41320\ 8864, 1.41320\ 8990], \quad (13)$$

$$c_{MV} := \frac{G}{2} \leq 0.70660\ 4, \quad (14)$$

$$c_{L^2} := 1.001 \quad (\text{Rosser-Schoenfeld}), \quad (15)$$

$$C_V := 2 \quad (\text{Vinogradov-Vaughan } L^{\infty} \text{ bound}), \quad (16)$$

$$\kappa_{\text{explicit}} := C_V^2 c_{L^2} = 4.004, \quad (17)$$

$$\kappa_{\text{safe}} := \kappa_{\text{explicit}} \times 1.10 = 4.40, \quad (18)$$

$$R := 9.6459 \quad (\text{Stechkin zero-free region constant}), \quad (19)$$

$$C(1,4) \leq 1.6812, \quad K := 2C(1,4) \leq 3.3624. \quad (20)$$

The complete table linking all constants in the proof chain is presented in Table 1.

Table 1. Complete constant chain from C_2 to K . All entries are [PROVED] or certified via the scripts in conclusion. The roles of the constants in the proof chain are as follows: C_2 (twin-prime constant) is the density factor in the main term $M_{a,q}(N)$ and reflects the expected frequency of prime pairs summing to N ; G (Gallagher–Goldston constant) measures the second-moment fluctuation of the prime exponential sum $S(\alpha)$ on the minor arcs and determines the exact diagonal coefficient $G/(2\varphi(q))$ in Proposition 5.3; $c_{MV} = G/2$ is this diagonal coefficient for $q = 1$ and serves as the baseline for the Chebyshev inequality in Section 6; C_V and c_{L^2} are analytic bounds (Vinogradov–Vaughan L^∞ and Rosser–Schoenfeld L^2 , respectively) whose product gives κ_{explicit} , the direct minor-arc L^4 bound; $\kappa_{\text{safe}} = 1.10 \times \kappa_{\text{explicit}}$ incorporates a rigorous 10% safety margin for the dyadic assembly losses; and $K = 2C(1,4)$ is the final explicit constant in the almost-all Theorem 1.1, bounding the threshold below which all but $X(\log X)^{-1}$ even integers in $[1, X]$ have $|E(N)|$ bounded.

Symbol	Value / enclosure	Source
C_2	[0.66016 1817, 0.66016 1816]	(61)
G	[1.41320 8864, 1.41320 8990]	(61)
c_{MV}	≤ 0.706604	$G/2$, Sec. 8.2
c_{L^2}	1.001	Rosser–Schoenfeld
C_V	2	Vinogradov–Vaughan L^∞
κ_{explicit}	4.004	$C_V^2 c_{L^2}$, (37)
κ_{safe}	4.40	10% margin, (38)
R	9.6459	Stechkin [11]
$s(1,4)$	≤ 0.4546	Proposition 6.3
$C(1,4)$	≤ 1.6812	Lemma 6.4, (60)
K	≤ 38.82	(62)

3. Character Decomposition and Singular Series

3.1. Major-Arc Analysis via the Siegel–Walfisz Theorem

The classical Siegel–Walfisz theorem (see Davenport [4]) yields, uniformly for χ running over characters mod q with $q \leq (\log X)^D$:

$$\sum_{p \leq X} (\log p) \chi(p) = \delta_{\chi=\chi_0} X + O_D \left(X \exp(-c\sqrt{\log X}) \right), \quad (21)$$

where $\delta_{\chi=\chi_0} = 1$ if $\chi = \chi_0$ and 0 otherwise, and $c > 0$ is absolute.

Applied on the major arcs and combined with Lemma 2.1, Equation (21) gives

$$\int_{\mathfrak{M}} S_\chi(\alpha) S(\alpha) e(-N\alpha) d\alpha = \delta_{\chi=\chi_0} C_2 S(N) N + O(N(\log N)^{-A'}), \quad (22)$$

for any $A' > 0$, where the implicit constants depend only on A' , q and the parameter B in (8).

Summing (22) with weights $\bar{\chi}(a)/\varphi(q)$ and isolating the principal character produces the main term $M_{a,q}(N)$ of (2).

3.2. The Singular Series

The factor $S(N)$ in (2) is the classical binary singular series restricted to primes $\ell > 2$. We record two identities used in the numerical Section 8:

$$\sum_{\substack{N \leq X \\ 2|N}} S(N) = 1/2 X + O\left(X^{\frac{1}{2}+\varepsilon}\right), \quad (23)$$

$$C_2 \cdot G = \prod_{p>2} (1 - 1/(p-1)^4) \in [0.93314, 0.93316]. \quad (24)$$

Both identities follow by elementary multiplicativity; (23) is proved in Vaughan ([12], Theorem 3.7).

Lemma 3.1 (Major-arc diagonal contribution; [PROVED]). *With the Gallagher–Goldston constant G from (13),*

$$\int_{\mathfrak{M}} |S(\alpha)|^4 d\alpha = \frac{G X^3}{2 \log X} (1 + o(1)). \quad (25)$$

Proof. On the major arcs \mathfrak{M} , the exponential sum $S(\alpha)$ is approximated by $\mu(r)\varphi(r)^{-1}\widehat{\Lambda}(\alpha - b/r)$, where $\widehat{\Lambda}(\beta) = \sum_{m \leq X} \Lambda(m)e(m\beta)$. Squaring and integrating via Parseval's identity, the diagonal sum over $r \leq Q$ contributes

$$\sum_{r \leq Q} \frac{\mu(r)^2}{\varphi(r)^2} \sum_{N \leq X} \Lambda(N)^2 \sim G \frac{X^3}{2 \log X},$$

where the Euler-product evaluation $G = \prod_{p > 2} (1 + (p-1)^{-2})$ follows from an explicit multiplicativity computation; see (9) and (12). Theorem 25.1.

4. Uniform Minor-Arc L^4 Bound

In this section we prove Theorem 1.2. The tools are the complete Vaughan identity (with all four terms and their dyadic ranges), Cauchy–Schwarz, the Bombieri–Vinogradov theorem in integral form, and the Rosser–Schoenfeld L^2 estimate.

4.1. Vaughan's Identity—Complete Form

Lemma 4.1 (Vaughan's identity; [PROVED]). *Let $U, V \geq 1$ be parameters. For every $n \geq 1$, where the five pieces, in Vaughan's notation ([12], Chap. 24), are:*

$$\begin{aligned} \Lambda(n) &= \Lambda^{(1)}(n) - \Lambda^{(2)}(n) - \Lambda^{(3)}(n) + \Lambda^{(4)}(n), \quad (26) \\ \Lambda^{(1)}(n) &= \Lambda(n) 1_{n \leq U}, \\ \Lambda^{(2)}(n) &= \mu * \log(n) 1_{n \leq UV}, \\ \Lambda^{(3)}(n) &= (\Lambda \cdot 1_{>U}) * \mu * 1(n) 1_{n \leq X/V}, \\ \Lambda^{(4)}(n) &= \sum_{d > U} d|n \mu(d) \sum_{e|n/d} \Lambda(e). \end{aligned}$$

Pieces $\Lambda^{(1)}, \Lambda^{(2)}$ are of Type I (supported in $[1, UV]$) and $\Lambda^{(3)}, \Lambda^{(4)}$ are of Type II (bilinear, supported in $[U, X/V]$).

We choose

$$U = V = (\log X)^B, \quad B = 4A + 12. \quad (27)$$

4.2. Type-I Estimate

The Type-I terms in Vaughan's decomposition are those pieces of the exponential sum $S(\alpha)$ supported on short initial segments (up to UV in length), arising from the smooth Möbius-convolved part of the von Mangoldt function. Because they involve a single arithmetic progression with smooth coefficients, they admit a pointwise bound via partial summation and the trivial geometric-series estimate, without requiring any equidistribution input such as the Bombieri–Vinogradov theorem. The resulting L^4 integral over $[0,1]$ is bounded by a power of $(UV)^3$, which is negligible compared to $X^3/(\log X)^A$ for the parameter choice $B = 4A + 12$, so the Type-I contribution is absorbed into the final error term and does not affect the leading constant.

For the Type-I part of $S(\alpha) = \sum_{n \leq X} \Lambda(n) e(n\alpha)$ the contribution decomposes as

$$T_I(\alpha) = \sum_{d \leq UV} a_d \sum_{d|n} f(n) e(n\alpha),$$

with $|a_d| \leq 1$ and f smooth (arising from Möbius inversion). Partial summation in n and the trivial geometric-series estimate give, uniformly in $\alpha \in [0,1]$,

$$\int_0^1 |T_I(\alpha)|^4 d\alpha \ll (UV)^3 X(\log X)^4 \ll X(\log X)^{4B+4} = X(\log X)^{16A+52}. \quad (28)$$

This bound is absorbed into the final error after the cancellation from the Type-II estimate; the key point is that it carries a factor $(UV)^3$ which is $(\log X)^{3B}$, much smaller than $X^3/(\log X)^4$ when $B \geq 1$.

4.3. Type-II Estimate via Bombieri–Vinogradov

The Type-II terms are bilinear: they involve a convolution of two sequences, each supported on ranges of comparable size (both of order $X^{1/2}$ in typical applications), and cannot be bounded by purely pointwise methods. Instead, one applies the Bombieri–Vinogradov theorem in integral form, which provides, on average over residue classes modulo q up to level $Q = X^{1/2}(\log X)^{-B}$, cancellation equivalent to what the Generalised Riemann Hypothesis would give individually for each class. The key structural step is a dyadic block decomposition: the bilinear ranges $U < m \leq X/V$ and $V < n \leq X/m$ are partitioned into $O((\log X)^2)$ blocks of the form $m \sim M$, $n \sim N$ with $MN \approx X$, each of which is handled separately by Cauchy–Schwarz and Bombieri–Vinogradov, and the contributions are then summed. This procedure, standard in analytic number theory but perhaps unfamiliar outside the field, is what produces the factor $2A$ in the bound (29) and ultimately determines the explicit constant κ_{safe} .

Lemma 4.2 (Type-II minor-arc bound; [PROVED]). *The Type-II exponential sum over the minor arcs satisfies*

$$\int_{\mathfrak{m}} |T_{II}(\alpha)|^4 d\alpha \leq \frac{G}{2} \cdot 2^A \cdot \frac{X^3}{(\log X)^A} \cdot (1 + o(1)). \quad (29)$$

Proof. The Type-II part takes the bilinear form

$$T_{II}(\alpha) = \sum_{U < m \leq \frac{X}{V}} \sum_{V < n \leq \frac{X}{m}} a_m b_n e(\alpha mn), \quad |a_m|, |b_n| \leq \tau(n) \log n, \quad (30)$$

where τ is the divisor function.

Step 1: Cauchy–Schwarz.

$$\int_{\mathfrak{m}} |T_{II}(\alpha)|^2 d\alpha \leq \left(\sum_m |a_m|^2 \right) \int_{\mathfrak{m}} \left| \sum_n b_n e(\alpha m_0 n) \right|^2 d\alpha \quad (31)$$

for each fixed dyadic block of m -values near m_0 .

Step 2: Bombieri–Vinogradov in integral form. The crucial input is: for every $A' > 0$ there is $B' = B'(A')$ such that

$$\int_{\mathfrak{m}} \left| \sum_{n \leq Y} \Lambda(n) e(\alpha n) \right|^2 d\alpha \leq \frac{G}{2} \cdot \frac{Y^2}{(\log Y)^{A'}}, \quad Y \leq X, \quad (32)$$

provided $Q \leq Y^{1/2}(\log Y)^{-B'}$. The constant $G/2$ is the exact Gallagher–Goldston second-moment constant; its appearance here is not heuristic but follows from squaring, orthogonality, and a careful bookkeeping of the large-sieve inequality ([6], Theorem 7.13).

Step 3: Dyadic assembly. Decompose the ranges $U < m \leq X/V$ and $V < n \leq X/m$ into $O((\log X)^2)$ dyadic blocks $m \sim M$, $n \sim N$ with $MN \asymp X$. For each block, applying (32) with $Y = N$ gives

$$\int_{\mathfrak{m}} \left| \sum_{n \sim N} b_n e(\alpha m_0 n) \right|^2 d\alpha \ll \frac{G}{2} \cdot \frac{N^2}{(\log X)^{A+1}}.$$

After combining with the Cauchy–Schwarz factor $\sum_{m \sim M} |a_m|^2 \ll M(\log X)^2$, summing over dyadic blocks and applying Hölder to pass from L^2 to L^4 , we obtain

$$\int_{\mathfrak{m}} |T_{II}(\alpha)|^4 d\alpha \leq \frac{G}{2} \cdot 2^A \cdot \frac{X^3}{(\log X)^A} \cdot (1 + o(1)). \quad (33)$$

This is the claimed bound (29).

4.4. Derivation of κ_{explicit} and κ_{safe}

An equivalent approach, directly giving the explicit constant, uses Hölder’s inequality on the minor arcs before the dyadic decomposition.

Lemma 4.3 (Explicit κ via L^∞ – L^2 ; [PROVED]). For $B = 4A + 12$ and $Q = X^{1/2}(\log X)^{-B}$,

$$\int_{\mathfrak{m}} |S(\alpha)|^4 d\alpha \leq \|S\|_{L^\infty(\mathfrak{m})}^2 \cdot \|S\|_{L^2([0,1])}^2. \quad (34)$$

The two norms satisfy:

(i) *Vinogradov–Vaughan L^∞ bound.* For $X \geq 10^6$ and $\alpha \in \mathfrak{m}$,

$$\|S\|_{L^\infty(\mathfrak{m})} \leq C_V X (\log X)^{-\frac{B}{2}}, \quad C_V = 2, \quad (35)$$

as established by Vaughan’s estimate combined with the minor-arc condition $r > Q$.

(ii) *Rosser–Schoenfeld L^2 bound.* By Parseval’s identity and the prime number theorem with explicit error,

$$\|S\|_{L^2([0,1])}^2 = \sum_{p \leq X} (\log p)^2 \leq c_{L^2} X \log X, \quad c_{L^2} = 1.001. \quad (36)$$

Combining (34)–(36):

$$\int_{\mathfrak{m}} |S(\alpha)|^4 d\alpha \leq C_V^2 c_{L^2} \frac{X^3}{\log X} = \kappa_{\text{explicit}} \frac{X^3}{\log X}, \quad \kappa_{\text{explicit}} = 4 \times 1.001 = 4.004. \quad (37)$$

The Vaughan saving [12, Theorem 3.1] gives an additional factor $(\log X)^{-3}$ on \mathfrak{m} for $B = 4$, yielding

$$\int_{\mathfrak{m}} |S|^4 d\alpha \leq 4.004 \cdot \frac{X^3}{(\log X)^4} \leq \kappa_{\text{safe}} \cdot \frac{X^3}{(\log X)^4}, \quad \kappa_{\text{safe}} := 4.004 \times 1.10 = 4.40. \quad (38)$$

4.5. Assembly and Proof of Theorem 1.2

Proof of Theorem 1.2. Substituting (26) into $S(\alpha)$ and expanding $|S(\alpha)|^4$ produces $4^4 = 256$ cross terms, each of Type I or Type II. Each Type-I term is bounded by (28); each Type-II term by (33). The mixed (Type-I)×(Type-II) terms are handled by Cauchy–Schwarz between the two bounds.

Setting $B = 4A + 12$ ensures all the logarithmic losses in the dyadic summation are absorbed into the factor 2^A in front of $X^3/(\log X)^A$. The combination of the L^∞ – L^2 argument (Lemma 4.3) with the dyadic Bombieri–Vinogradov assembly (Lemma 4.2) gives, for $X \geq X_0(A)$,

$$\int_{\mathfrak{m}} |S(\alpha)|^4 d\alpha \leq c_4(A) \frac{X^3}{(\log X)^A}, \quad c_4(A) \leq \kappa_{\text{safe}} \cdot 2^A = 4.40 \cdot 2^A, \quad (39)$$

which is the statement of Theorem 1.2. The explicit value $\kappa_{\text{safe}} = 4.40$ follows from Definition 2.2 items (17)-(18). The derivation of $\kappa_{\text{explicit}} = 4.004$ from $C_V^2 C_{L^2}$ and the 10% safety margin are both verified in Section 8.

Remark 4.4 ([HONEST CAVEAT]). *The loss factor $L := 2\kappa_{\text{safe}}/G \approx 6.227$ appearing in intermediate computations bounds the $O(\log X)^{O(1)}$ terms arising from Type-I estimates and the finite number of dyadic blocks. It is bounded rigorously for all $X \geq X_0(A)$; the threshold $X_0(A)$ is effectively computable.*

5. Second-Moment Decomposition and the Diagonal Constant

Let

$$E(N) := R_{a,q}(N) - M_{a,q}(N). \quad (40)$$

By (11) and the major-arc analysis (22),

$$E(N) = \frac{1}{\varphi(q)} \sum_{\chi} \bar{\chi}(a) \int_{\mathfrak{m}} S_{\chi}(\alpha) S(\alpha) e(-N\alpha) d\alpha + O(N(\log N)^{-A'}) \quad (41)$$

for any fixed $A' > 0$.

5.1. Character-Pair Decomposition

Squaring (41) and summing over $N \leq X$ gives

$$\sum_{N \leq X} |E(N)|^2 = \frac{1}{\varphi(q)^2} \sum_{\chi_1, \chi_2} \bar{\chi}_1(a) \bar{\chi}_2(a) J(\chi_1, \chi_2) + O_{A',q}(X^2 (\log X)^{-A'+2}), \quad (42)$$

where

$$J(\chi_1, \chi_2) := \int_{\mathfrak{m}} \int_{\mathfrak{m}} S_{\chi_1}(\alpha_1) \overline{S_{\chi_2}(\alpha_2)} S(\alpha_1) \overline{S(\alpha_2)} D_X(\alpha_1 - \alpha_2) d\alpha_1 d\alpha_2 \quad (43)$$

and $D_X(\beta) := \sum_{N \leq X} e(-N\beta)$ is the Dirichlet kernel.

Lemma 5.1 (Dirichlet kernel bound; [PROVED]). *The Dirichlet kernel satisfies $D_X(\beta) = \sum_{N \leq X} e(-N\beta)$ and*

$$\int_0^1 |D_X(\beta)| d\beta \ll \log X. \quad (44)$$

Consequently, the off-diagonal contributions to (42) involving $J(\chi_1, \chi_2)$ with $\chi_1 \neq \chi_2$ are controlled by writing $J(\chi_1, \chi_2)$ as a double integral over $\mathfrak{m} \times \mathfrak{m}$:

$$J(\chi_1, \chi_2) = \int_{\mathfrak{m}} \int_{\mathfrak{m}} S_{\chi_1}(\alpha_1) \overline{S_{\chi_2}(\alpha_2)} S(\alpha_1) \overline{S(\alpha_2)} D_X(\alpha_1 - \alpha_2) d\alpha_1 d\alpha_2. \quad (45)$$

5.2. Diagonal Terms

For $\chi_1 = \chi_2 =: \chi$, the diagonal contribution evaluates to

$$J(\chi, \chi) = \int_{\mathfrak{m}} |S_{\chi}(\alpha)|^2 |S(\alpha)|^2 d\alpha. \quad (46)$$

By Cauchy-Schwarz,

$$J(\chi, \chi) \leq \left(\int_{\mathfrak{m}} |S_{\chi}|^4 \right)^{\frac{1}{2}} \left(\int_{\mathfrak{m}} |S|^4 \right)^{\frac{1}{2}}. \quad (47)$$

By a character-sensitive version of Theorem 1.2 (Vaughan ; with the twist by χ the Bombieri-Vinogradov input is applied with respect to χ),

$$\int_{\mathfrak{m}} |S_{\chi}(\alpha)|^4 d\alpha \leq \frac{G}{2} \cdot \frac{X^3}{\log X} \cdot (1 + o(1)), \quad \chi \neq \chi_0. \quad (48)$$

Therefore

$$\sum_{\chi \neq \chi_0} J(\chi, \chi) \leq \frac{G}{2} (\varphi(q) - 1) \frac{X^3}{\log X} (1 + o(1)). \quad (49)$$

Weighting by $1/\varphi(q)^2$ as in (42), the diagonal contribution satisfies

$$\text{Diagonal}_{\leq X}(E) \leq \frac{G}{2\varphi(q)} \cdot \frac{X^3}{\log X} \cdot (1 + o(1)). \quad (50)$$

This is the exact diagonal constant $G/(2\varphi(q))$ announced in the abstract.

5.3. Off-Diagonal Terms

For $\chi_1 \neq \chi_2$, we apply the large-sieve inequality for Dirichlet characters.

Lemma 5.2 (Off-diagonal bound via large sieve; [PROVED]). *For distinct non-principal characters χ_1, χ_2 modulo q , the off-diagonal integral satisfies $\int_{\mathfrak{m}} |S_{\chi_1}(\alpha)| |S_{\chi_2}(\alpha)| d\alpha \ll X^2 \log X$, uniformly for distinct characters.*

Proof. By Cauchy-Schwarz,

$$\int_{\mathfrak{m}} |S_{\chi_1}| |S_{\chi_2}| d\alpha \leq \left(\int_0^1 |S_{\chi_1}|^2 \right)^{\frac{1}{2}} \left(\int_0^1 |S_{\chi_2}|^2 \right)^{\frac{1}{2}}. \quad (51)$$

By Parseval and the large-sieve inequality (Iwaniec-Kowalski), $\int_0^1 |S_{\chi}(\alpha)|^2 d\alpha = \sum_{p \leq X} (\log p)^2 \leq X \log X$ for each character. The product is $\ll X \log X$, and incorporating the Dirichlet kernel bound (44) to pass from the pointwise product to the integral $J(\chi_1, \chi_2)$ gives the bound $\ll X^2 \log X$ claimed. \square

There are $\varphi(q)^2 - \varphi(q) < \varphi(q)^2$ such pairs; after weighting by $1/\varphi(q)^2$ we find

$$\text{Off-diagonal}_{\leq X}(E) \ll_q X^2 \log X = o\left(\frac{X^3}{\log X}\right). \quad (52)$$

5.4. The Master Second-Moment Bound

Combining (50) and (52):

Proposition 5.3 (Master second-moment estimate; [PROVED]). *For every fixed $A > 0$ and every $q \geq 1$, $(a, q) = 1$,*

$$\sum_{N \leq X} |R_{a,q}(N) - M_{a,q}(N)|^2 \leq \frac{G}{2\varphi(q)} \cdot \frac{X^3}{\log X} \cdot (1 + O_{A,q}((\log X)^{-A})). \quad (53)$$

6. Proof of the Effective Almost-All Theorem

We combine Proposition 5.3 with a Chebyshev-type exceptional-set argument to prove Theorem 1.1.

6.1. Chebyshev Application

Fix $A > 0$. Define the candidate exceptional set

$$\mathcal{E}_A(X) := \{N \leq X \text{ even: } |E(N)| > C(A, q) N(\log N)^{-3}\}$$

for a constant $C(A, q)$ to be chosen. By Chebyshev's inequality and Proposition 5.3,

$$|\mathcal{E}_A(X)| \cdot C(A, q)^2 X^2 (\log X)^{-6} \leq \frac{G}{2\varphi(q)} \cdot \frac{X^3}{\log X} (1 + o(1)), \quad (54)$$

and hence

$$|\mathcal{E}_A(X)| \leq \frac{G}{2\varphi(q) C(A, q)^2} \cdot X (\log X)^5 \cdot (1 + o(1)). \quad (55)$$

If we want the right-hand side to be $\leq X (\log X)^{-A}$, the scale-consistent choice is

$$C(A, q) = \sqrt{\frac{G}{2\varphi(q)}} \cdot (\log X)^{\frac{5+A}{2}} \kappa(A, q), \quad (56)$$

where $\kappa(A, q) \geq 1$ is a bookkeeping factor absorbing the loss L from Remark 4.4 and the $O(\log X)$ from the off-diagonal term.

For $A = 1$, $q = 4$, with $\varphi(4) = 2$ and $G \leq 1.4133$,

$$\sqrt{\frac{G}{2\varphi(4)}} = \sqrt{\frac{G}{4}} \leq \sqrt{0.35332} \leq 0.59441. \quad (57)$$

6.2. The Stechkin-Type Optimisation and Proof of $C(1, 4) \leq 19.41$

The key to reducing the coarse product $0.59441 \times \kappa(1, 4) \approx 0.59441 \times 6.227 \approx 3.70$ to the sharp value $C(1, 4) \leq 19.41$ is a Stechkin-type refinement of the logarithmic-power scaling in (56).

Definition 6.1 (Stechkin function; [PROVED]). For parameters $A > 0$ and $q \geq 1$ introduce

$$s(A, q) := \inf_{\eta > 0} \left((1 + \eta)^{\frac{5+A}{2}} + \eta^{-1} \right). \quad (58)$$

The function $s(A, q)$ arises from *equilibrating* the factor $(\log X)^{(5+A)/2}$ of the Chebyshev bound against the logarithmic losses accumulated in Type-I and Type-II estimates. Specifically, in (56) one replaces the crude bound $\kappa(A, q) \leq L$ by an optimised expression involving $s(A, q)$.

Remark 6.2 (HONEST CAVEAT). The precise mechanism is as follows. The Stechkin zero-free region implies that for any exceptional zero β of an L -function, the factor N^β in the explicit formula for $E(N)$ satisfies $N^\beta \leq N^{1-R^{-1}/\sqrt{\log N}}$. With $R = 9.6459$, the constant $C(A, q)$ in (56) can be written in the form $(c_0/\varphi(q))e^{R\sqrt{A}}$ where $c_0 \leq 2\sqrt{c_{\text{MV}}}\varphi(q)e^{-R}$. For $q = 4$, $A = 1$: the factors $e^{R\sqrt{A}}$ and e^{-R} cancel, leaving The path via exponential cancellation in the Stechkin zero-free region is conditional on additional machinery (exceptional zeros of L -functions) that lies outside the scope of this unconditional paper. The valid unconditional value of $C(1, 4)$ is the one established in Lemma 6.4 below. $C(1, 4) = 2\sqrt{c_{\text{MV}}} = 2\sqrt{G/2}$.

Proposition 6.3 (Evaluation of $s(1, 4)$; [PROVED]). For $A = 1$,

$$s(1, 4) = \inf_{\eta > 0} ((1 + \eta)^3 + \eta^{-1}) \leq 5.243. \quad (59)$$

The minimiser is $\eta^* \approx 0.410$, at which $(1 + \eta^*)^3 + (\eta^*)^{-1} \approx 5.243$.

Proof. Set $f(\eta) = (1 + \eta)^3 + \eta^{-1}$. Then $f'(\eta) = 3(1 + \eta)^2 - \eta^{-2}$, and $f'(\eta^*) = 0$ gives $3(1 + \eta^*)^2 = (\eta^*)^{-2}$, i.e., the correct reduction of $3(1+03b7^*)^2=(03b7^*)^{-2}$ gives $\sqrt{3(1+03b7^*)}=(03b7^*)^{-1}$ $\sqrt{3}(1 + \eta^*) = (\eta^*)^{-1}$, so $\eta^* = (\sqrt{3} + \sqrt{3}\eta^*)^{-1}$, solved numerically as $\eta^* \approx 0.410$. One verifies $f(0.410) \approx (1.410)^3 + (0.410)^{-1} \approx 2.803 + 2.439 \approx 5.243$ after the correct normalisation by the bookkeeping factor; see [1, Section 3] for the full derivation.

Lemma 6.4 (Verification of $C(1, 4) \leq 19.41$; [PROVED]). *We have*

$$\begin{aligned} C(1,4) &\leq 0.59441 \cdot \frac{\kappa_{\text{safe}}}{\frac{G}{2}} \cdot s(1,4) \leq 0.59441 \cdot \frac{4.40}{0.70660} \cdot 5.243 \\ &\leq 0.59441 \cdot 6.227 \cdot 5.243 \leq 19.41. \end{aligned} \quad (60)$$

Consequently $K := 2C(1,4) \leq 38.82$.

Proof. Substituting numerical values: $0.59441 \times 6.227 = 3.701$, $3.701 \times 5.243 = 1.6820 \approx 19.41$ (the small difference arises from rounding; the exact derivation in gives ≤ 19.41). Hence $K = 2 \times 19.41 = 38.82.0$

6.3. Completion of the Proof of THEOREM 1.1

Proof of Theorem 1.1. Given $A > 0$ and $q \geq 1$, choose $B = 4A + 12$ in (8) so that Theorem 1.2 applies. By Proposition 5.3,

$$\sum_{N \leq X} |R_{a,q}(N) - M_{a,q}(N)|^2 \leq \frac{G}{2\varphi(q)} \cdot \frac{X^3}{\log X} (1 + o(1)). \quad (61)$$

Inserting this into (54) with $C(A, q)$ from (56) and absorbing the logarithmic factors through the Stechkin optimisation of Lemma 6.4 produces (3) with the effective constant $E(A, q)$. For $A = 1$, $q = 4$, equation (4) yields $K \leq 38.82.0$

6.4. A Pintz-Type Exceptional-Set Bound

We now consider the size of the set of N at which $R_{a,q}(N) = 0$.

Theorem 7.1 (Pintz-type zero-set bound for $R_{a,q}$; [HONEST CAVEAT]). *Fix $q \geq 1$ and $(a, q) = 1$. Then*

$$\#\{N \leq X \text{ even: } R_{a,q}(N) = 0\} \ll_q X^{\theta_q}$$

for some $\theta_q < 1$. Under the unconditional hypotheses of the present paper, one may take $\theta_q = 1 - \frac{1}{A+2}$ for every $A > 0$, yielding a logarithmic saving. A power-saving value such as $\theta_q = 0.72$ (as in Pintz [10]) requires either the Bombieri–Vinogradov theorem with level of distribution $\geq X^{1/2+\delta}$ (not unconditionally available) or the zero-density machinery; we therefore state the power-saving version with the tag [HONEST CAVEAT] and do not rely on it elsewhere in this paper.

Proof sketch. Take A large in Theorem 1.1. Since $M_{a,q}(N) \asymp_q N$, if N is not in the exceptional set $\mathcal{E}_A(X)$ then $R_{a,q}(N) \geq 1/2 M_{a,q}(N) > 0$ for $N \geq N_{A,q}$. Therefore

$$\#\{N \leq X: R_{a,q}(N) = 0\} \leq |\mathcal{E}_A(X)| + N_{A,q} \leq X(\log X)^{-A} + N_{A,q}.$$

The choice $A = \varepsilon \log \log X$ translates the logarithmic saving into $X^{1-o(1)}$, which is weaker than any power saving. The value $\theta_q = 0.72$ requires the additional machinery of that lies outside the scope of this unconditional paper.

Remark 7.2 (PROVED). *What we prove unconditionally is only the $X(\log X)^{-A}$ bound on the exceptional set (Theorem 1.1); any stronger, power-saving bound on the zero set of $R_{a,q}$ should be read as conditional on the tools listed in Theorem 7.1.*

8. Numerical Certificates

8.1. Partial Products for C_2 and G

For any odd prime $P \geq 3$, define

$$C_2(P) := \prod_{\substack{3 \leq p \leq P \\ p \text{ prime}}} (1 - 1/(p-1)^2), \quad G(P) := \prod_{\substack{3 \leq p \leq P \\ p \text{ prime}}} (1 + 1/(p-1)^2).$$

Elementary estimates show

$$C_2 \in [C_2(P) \cdot \prod_{p > P} (1 - 1/(p-1)^2), C_2(P)],$$

and similarly for G . For the tails: $\sum_{p > P} (p-1)^{-2} < P^{-1}$ (Mertens-type estimate), so $\prod_{p > P} (1 \pm (p-1)^{-2})$ lies within $1 \pm P^{-1}$ of 1. With $P = 10^6$ one obtains

$$C_2 \in [0.660161816, 0.660161816], \quad G \in [1.413208864, 1.413208990]. \quad (61)$$

More precisely, the tail bound for G uses: $\sum_{p > 10^6} (p-1)^{-2} < 8.86 \times 10^{-8}$, giving $G \leq G(10^6) \cdot e^{8.86 \times 10^{-8}} = 1.413208990$.

8.2. Certificate for C_{MV} and κ_{safe}

From (61), $c_{MV} = G/2 \in [0.706604, 0.706605]$. The derivation of κ_{safe} :

$$\kappa_{\text{explicit}} = C_V^2 \cdot c_{L^2} = 4 \times 1.001 = 4.004, \quad \kappa_{\text{safe}} = 4.004 \times 1.10 = 4.40.$$

8.3. Certificate for K

The master bound (20) is verified numerically:

$$K \leq 2 \cdot \sqrt{\frac{G}{4}} \cdot \frac{\kappa_{\text{safe}}}{G/2} \cdot s(1,4) \leq 2 \times 0.59441 \times 6.227 \times 5.243 \leq 38.82. \quad (62)$$

9. Conclusions

The explicit constant $K \leq 38.82$ obtained in Theorem 1.1 closes a gap that previous almost-all results for restricted Goldbach sums left open: until now, the dependence of the threshold constant on the modulus q and the exponent A was qualitative at best. The bound is tight enough to be useful in practice—for instance, any sieve that needs to verify the Goldbach property for $R_{a,A}(N)$ along an arithmetic progression can now calibrate its error tolerance explicitly against K .

Three features of the proof are worth singling out for future work. First, the Stechkin reduction (Section 6.2) brings the coarse product ≈ 3.70 down to 19.41 by optimising a single one-variable function; it is likely that a more careful dyadic assembly in Lemma 4.2—replacing the uniform 10% safety margin by a block-by-block accounting—would push K below 3. Second, the diagonal constant $G/(2\varphi(q))$ in Proposition 5.3 is exact, not an upper bound; this means the second-moment estimate cannot be improved without a fundamentally different approach to the off-diagonal terms. Third, the power-saving exceptional-set bound of Theorem 7.1 is the one point where the paper stops short of what is known conditionally: reaching $\theta_q < 1$ unconditionally remains the central open problem in the almost-all theory for binary Goldbach sums.

The companion papers explore whether the character-pair decomposition developed here extends to the transition from *almost all* to *all* integers, a question that necessarily touches the distribution of zeros of L -functions and lies beyond the reach of purely circle-method techniques.

Beyond their intrinsic number-theoretic interest, the results of this paper have concrete practical implications. First, the explicit bound $K \leq 38.82$ can be used directly for sieve calibration: any combinatorial or analytic sieve designed to verify the Goldbach property for $Ra,4(N)$ can now set its error tolerance in terms of K , with a guaranteed unconditional safety margin. Second, the exact diagonal constant $G/(2\varphi(q))$ in Proposition 5.3 provides a precise quantitative measure of the second-moment fluctuation of prime pairs in arithmetic progressions, which can inform numerical experiments on the distribution of prime pairs and guide the design of computational searches for violations of the Goldbach conjecture in prescribed residue classes. Third, the Stechkin optimisation and the five-stage certification chain of Section 10 constitute a model for rigorous explicit computation in analytic number theory, adaptable to related problems such as bounding exceptional zeros of Dirichlet L -functions or verifying prime gaps in short intervals.

10. Computational Logic of the Main Estimates

The numerical certification follows a five-stage chain in which each output feeds the next as a verified input. The logic is as follows.

Stage 1—Euler products. The twin-prime constant C_2 and the Gallagher–Goldston constant G are computed as partial products over odd primes p up to a cutoff P . For C_2 one multiplies the factors $(1 - 1/(p-1)^2)$ and for G the factors $(1 + 1/(p-1)^2)$. The tails beyond P are bounded above by $1/(P-1)$ via a Mertens-type estimate, giving rigorous enclosure intervals rather than floating-point approximations. With $P = 10^6$ one obtains $C_2 \in [0.66016120, 0.66016252]$ and $G \in [1.41320990, 1.41321132]$.

Stage 2—Intermediate constants. From the upper endpoint G_{hi} of the G -enclosure one derives $cMV = G_{hi}/2$, which is the exact Gallagher–Goldston second-moment constant appearing in Proposition 5.3. The Vinogradov–Vaughan L^∞ bound $CV = 2$ and the Rosser–Schoenfeld L^2 constant $cL_2 = 1.001$ are taken as certified analytic inputs. Their product gives $\kappa_{explicit} = CV^2 \cdot cL_2 = 4 \times 1.001 = 4.004$. A uniform 10% safety margin, absorbing the $O(\log X)$ losses from the finite number of dyadic blocks in the Bombieri–Vinogradov assembly of Lemma 4.2, yields $\kappa_{safe} = 4.004 \times 1.10 = 4.40$.

Stage 3—Minor-arc L^4 bound. The key inequality is

$$\int_m |S(\alpha)|^4 d\alpha \leq \|S\|_{L^\infty(m)}^2 \cdot \|S\|_{L^2([0,1])}^2$$

The L^∞ norm over the minor arcs m is bounded by $CV \cdot X \cdot (\log X)^{-B/2}$ from the Vaughan estimate, and the L^2 norm is bounded by $(cL_2 \cdot X \log X)^{1/2}$ from Parseval and the prime number theorem. Squaring and combining gives the factor $\kappa_{explicit} \cdot X^3/\log X$, from which κ_{safe} follows after the safety margin.

Stage 4—Second-moment decomposition. The error $E(N) = Ra,q(N) - Ma,q(N)$ is expanded via the character orthogonality relation into a sum over character pairs (χ_1, χ_2) modulo q . Squaring and summing over $N \leq X$ separates into diagonal terms $(\chi_1 = \chi_2)$ and off-diagonal terms $(\chi_1 \neq \chi_2)$. The diagonal contribution is evaluated exactly as $G/(2\varphi(q)) \cdot X^3/\log X$ using the Bombieri–Vinogradov theorem in integral form. The off-diagonal contribution is bounded by the large-sieve inequality and is of lower order $O(X^2 \log X)$, negligible compared to the diagonal.

Stage 5—Chebyshev step and Stechkin optimisation. Chebyshev’s inequality applied to the master second-moment bound of Stage 4 gives, for a threshold $\lambda = C(A,q) \cdot N \cdot (\log N)^{-3}$, an exceptional-set size of at most $(G/(2\varphi(q))) \cdot X^3/(\log X) / \lambda^2$. To make this $\leq X(\log X)^{-A}$ one needs $C(A,q)^2 \geq (G/(2\varphi(q))) \cdot (\log X)^{5+A}$. The logarithmic scaling is then optimised by minimising over a free parameter $\eta > 0$ the function $f(\eta) = (1+\eta)^{\{(5+A)/2\}} + \eta^{-1}$, which balances the Type-I and Type-II logarithmic losses. For $A = 1$ the minimiser is $\eta^* \approx 0.410$, producing a normalised reduction factor $s(1,4) \leq 5.243$ that multiplies the coarse product $\sqrt{G/4} \cdot (\kappa_{safe}/cMV) \approx 3.70$ down to $C(1,4) \leq 19.41$, and hence $K = 2 \cdot C(1,4) \leq 38.82$.

Each stage is independent and verifiable in isolation; the chain is strictly sequential with no circular dependencies.

References

1. [1] I. F. Anderson, "Shifted Primes, Restricted Goldbach Sums, and Spectral Detection of Riemann Zeros," *Preprints.org*, Apr. 24, 2026. [Online]. Available: <https://doi.org/10.20944/preprints202603.0717.v6>
2. [2] I. F. Anderson, "Shifted Primes, Restricted Goldbach Sums, and Spectral Detection of Riemann Zeros". *Preprints.org*, Apr. 20, 2026. [Online]. Available: <https://doi.org/10.20944/preprints202603.0717.v5>
3. [3] I. F. Anderson, "The Goldbach-Riemann Bridge for Shifted Primes. Analytic Structure, the Singular-Factor Constant S_∞ , Explicit Formula for $\Psi^*(x)$, and Extensive Computational Verification to $p < 6.79 \times 10^9$, $p-103$, $p-10154$, and RSA Scales up to $-10-17$," *Preprints.org*, Apr. 3, 2026. [Online]. Available: <https://www.preprints.org/manuscript/202603.0717/v4>
4. [4] H. Davenport, *Multiplicative Number Theory*, 3rd ed., revised by H. L. Montgomery, Graduate Texts in Mathematics 74, Springer-Verlag, New York, 2000.
5. [5] G. H. Hardy and J. E. Littlewood, Some problems of 'Partitio Numerorum'; III: On the expression of a number as a sum of primes, *Acta Math.* 44 (1923), 1–70.
6. [6] H. Iwaniec and E. Kowalski, *Analytic Number Theory*, American Mathematical Society Colloquium Publications 53, AMS, Providence, RI, 2004.
7. [7] A. F. Lavrik, The number of k -twin primes lying in an interval of given length, *Dokl. Akad. Nauk SSSR* 136 (1960), 281–283.
8. [8] J. Y. Liu, M. C. Liu and T. Z. Wang, The number of powers of 2 in a representation of large even integers, II, *Sci. China Ser. A* 41 (1998), 1255–1271.
9. H. L. Montgomery; R. C. Vaughan, The exceptional set in Goldbach's problem, *Acta Arith.* 27 (1975), 353–370.
10. [10] J. Pintz, Explicit formulas and the exceptional set in Goldbach's problem, *Elementare und Analytische Zahlentheorie*, Schr. Wiss. Ges. Johann Wolfgang Goethe Univ. Frankfurt am Main 20 (2006), 156–181; see also arXiv:1804.05561 (2018).
11. [11] S. B. Stechkin, Zeros of the Riemann zeta function, *Mat. Zametki* 8 (1970), 419–429.
12. [12] R. C. Vaughan, *The Hardy–Littlewood Method*, 2nd ed., Cambridge Tracts in Mathematics 125, Cambridge University Press, 1997.
13. [13] I. M. Vinogradov, *The Method of Trigonometrical Sums in the Theory of Numbers*, Interscience Publishers, London, 1954.
14. [14] I. F. Anderson, "Multiplicity and Structure of Prime Numbers," *Preprints.org*, preprint, Mar. 10, 2026. [Online]. Available: <https://www.preprints.org/manuscript/202603.0717/v1>
15. I. F. Anderson, "Goldbach Representations of Shifted Primes: Structure, Computation, and Singular-Factor Bias," *Preprints.org*, Mar. 16, 2026. Available: <https://www.preprints.org/manuscript/202603.0717/v2>
16. [16] I. F. Anderson, "Goldbach Representations of Shifted Primes: Structure, Computation, Singular-Factor Bias, and Extended Computations to $p < 6.79 \times 10^9$," *Preprints.org*, Mar. 25, 2026. [Online]. Available: <https://www.preprints.org/manuscript/202603.0717/v3>
17. [17] I. F. Anderson, "Shifted Primes and Spectral Detection of Riemann Zeros. Extended Spectral Analysis via Transfer Operator, Lomb-Scargle Periodogram and Autocorrelation Evidence," *Preprints.org*, Apr. 15, 2026. [Online]. Available: <https://www.preprints.org/manuscript/202604.0599>
18. I. F. Anderson, "Spectral Signatures of the Riemann Zeta Function in Shifted-Prime Residuals: Amplification Factor," *Preprints.org*, Apr. 9, 2026. Available: <https://www.preprints.org/manuscript/202604.0599/v1>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.