

Article

Not peer-reviewed version

First Computational Characterization of HTR5A-AS1: a Schizophrenia-Linked Antisense RNA with Synaptic Functions

[Grant H. Ruttenberg](#)*

Posted Date: 25 September 2025

doi: 10.20944/preprints202509.2129.v1

Keywords: schizophrenia; long noncoding RNA; HTR5A-AS1; antisense RNA; serotonin system; transcriptome-wide association study; BrainSpan; synaptic signaling; GABAergic signaling; neurodevelopment



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

First Computational Characterization of *HTR5A-AS1*: A Schizophrenia-Linked Antisense RNA with Synaptic Functions

Grant H. Ruttenberg ^{1,2} 

¹ Independent Researcher, Los Angeles, CA 90024, USA; ghruttenberg@gmail.com

² Department of Pancreas and Liver Transplant Surgery, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90095, USA

Abstract

Background/Objectives: Schizophrenia is a highly heritable psychiatric disorder that affects approximately 1% of the global population. Genome-wide association studies (GWAS) have mapped most schizophrenia risk variants to noncoding regions, highlighting the role of regulatory processes and noncoding RNAs in schizophrenia pathology. Despite this, and schizophrenia's association with 5-hydroxytryptamine (serotonin) system dysfunction, *HTR5A-AS1*, a long noncoding RNA (lncRNA) antisense to the serotonin receptor (HTR, 5-hydroxytryptamine receptor) gene *HTR5A*, remains virtually unstudied. This study provides the first systematic characterization of *HTR5A-AS1*, validating its transcript structure and investigating its genetic associations, expression dynamics, developmental regulation, and potential synaptic and GABAergic functions in schizophrenia. **Methods:** Transcriptome-wide association study (TWAS) summary statistics were integrated with postmortem RNA sequencing (RNA-seq), BrainSpan developmental transcriptomes, UCSC Genome Browser annotations, and functional prediction tools. These complementary approaches enabled validation of the transcript's structure, quantification of regional and developmental expression, and assessment of potential molecular functions. **Results:** *HTR5A-AS1* showed significant TWAS associations with schizophrenia in the hippocampus and dorsolateral prefrontal cortex (dlPFC). In postmortem schizophrenia donor tissue, expression was significantly reduced in the hippocampus, with a non-significant but directionally similar decrease in the dlPFC. Developmental transcriptomes revealed region-specific developmental trajectories, with steep increases during adolescence, aligning with the age range of typical schizophrenia onset. *HTR5A-AS1* was strongly co-expressed with *HTR5A*, and functional predictions implicated involvement in synaptic and GABAergic signaling, consistent with cortico-hippocampal circuit disruption in schizophrenia. **Conclusions:** These findings provide the first evidence that *HTR5A-AS1* is a bona fide antisense transcript with developmental and synaptic roles that may contribute to schizophrenia risk. Future single-cell and functional perturbation studies are needed to test causality and define mechanisms of regulation.

Keywords: schizophrenia; long noncoding RNA; *HTR5A-AS1*; antisense RNA; serotonin system; transcriptome-wide association study; BrainSpan; synaptic signaling; GABAergic signaling; neurodevelopment

1. Introduction

Schizophrenia is a severe psychiatric disorder that impacts approximately 0.3–0.7% of people during their lifetime, corresponding to about 21 million individuals worldwide [1]. Age-adjusted prevalence has remained relatively stable, but incidence rates have risen modestly over recent decades, with more new diagnoses per year [2]. Schizophrenia is characterized by hallucinations, delusions, cognitive deficits, and social withdrawal [3]. Environmental factors such as maternal stress and especially adolescent cannabis use may contribute to a person's likelihood of getting schizophrenia.

However, twin, family, and genome-wide association studies (GWAS) firmly establish genetics as the central risk component, with genetic factors explaining approximately 80% [4] of the risk for schizophrenia. GWAS have identified 287 genetic variants and 120 genes linked to schizophrenia. Interestingly, of the 287 genome-wide significant risk loci, only 106 are in protein-coding genes [5,6]. These findings highlight the importance of regulatory processes and noncoding RNAs as a risk factor for schizophrenia.

In addition, schizophrenia is widely considered a neurodevelopmental disorder, with symptoms typically emerging in adolescence or early adulthood. Converging evidence suggests that genetic risk factors act, in part, by altering neurodevelopmental trajectories: large-scale transcriptomic studies show that schizophrenia-associated genes are preferentially expressed prenatally and during adolescence, when synaptic pruning and cortical maturation occur [7]. This framework emphasizes the importance of evaluating potential risk genes across developmental stages and brain regions.

Schizophrenia is consistently linked to structural and functional abnormalities across brain regions. Meta-analyses and longitudinal magnetic resonance imaging (MRI) studies reveal reduced gray matter volumes and altered connectivity in areas including the prefrontal cortex, hippocampus, superior temporal cortex, thalamus, and anterior cingulate cortex [8]. These structures are critical for cognition, memory, and emotional regulation, and their dysfunction highlights schizophrenia's characterization as a dysconnection syndrome, defined by disrupted cognition and impaired high-order neural integration.

The 5-hydroxytryptamine (serotonin) system has been consistently implicated in the pathology of schizophrenia, especially through its involvement in mood, cognition, and sensory processing [9]. The serotonin receptor family (HTR, 5-hydroxytryptamine receptor) consists of seven main classes from *HTR1A* to *HTR7* and 14 subtypes. Antipsychotic drugs used to treat schizophrenia primarily target dopamine receptors [10], although many have also been shown to act on serotonin receptors [11], particularly *HTR1A* and *HTR2A*. Interestingly, variants in *HTR2A*, *HTR1A*, *HTR2C*, and *HTR3A* have been implicated in schizophrenia [12].

By contrast, *HTR5A* is far less studied despite its clear physiological relevance. *HTR5A* encodes the serotonin 5-HT_{5A} receptor, a G protein-coupled receptor (GPCR) that inhibits adenylate cyclase and modulates cyclic adenosine monophosphate (cAMP) signaling [13]. *HTR5A* is highly expressed in the cortex and hippocampus [13], areas that are consistently disrupted in schizophrenia. Nevertheless, unlike *HTR2A* and *HTR1A*, which are highly characterized, *HTR5A*'s role in psychiatric disease remains largely unexplored. This gap makes it especially intriguing to investigate *HTR5A* as a potential contributor to the pathophysiology of schizophrenia [14].

Transcriptome-wide association studies (TWAS) present a means of addressing gene-disease associations by integrating GWAS summary statistics with expression quantitative trait loci (eQTL) data to score a gene based on an association between its predicted expression and a trait [15]. A recent TWAS study identified the expression of a long noncoding RNA (lncRNA) transcribed antisense to *HTR5A*, *HTR5A-AS1*, to be significantly associated with schizophrenia risk [16]. Collado-Torres et al. profiled 900 postmortem brain tissue samples from the hippocampus ($n = 447$) and dorsolateral prefrontal cortex (dlPFC; $n = 453$) from 286 individuals diagnosed with schizophrenia and 265 not diagnosed with schizophrenia. By integrating the most recent schizophrenia GWAS and their own eQTL results, the authors performed brain region-specific TWAS [16] and reported over 1,140 significant associations spanning 333 genes, including two independent associations with *HTR5A-AS1*: a junction-level hippocampal signal ($Z = -3.99$, $p = 6.52 \times 10^{-5}$, false discovery rate [FDR] = 2.93×10^{-3}) and an exon-level signal in the dlPFC ($Z = -3.54$, $p = 4.06 \times 10^{-4}$, FDR = 1.04×10^{-2}). The significant association of *HTR5A-AS1* in the hippocampus is linked to schizophrenia GWAS variants that are predicted to act as eQTLs regulating its expression. Nevertheless, *HTR5A-AS1* remains largely unexplored in the literature, with PubMed [17] and GeneCards [18] searches showing no PubMed-indexed publications explicitly mentioning it to date.

2. Materials and Methods

2.1. Transcriptome-Wide Association Study (TWAS)

R [22] version 4.4.3 with the readr [23], dplyr [24], ggplot2 [25], and stringr [26] packages was used to analyze Collado-Torres et al. TWAS results, which were downloaded from <http://eqtl.brainseq.org/phase2/>. The TWAS data were imported, and gene identifiers were cleaned by removing Ensembl version suffixes to allow for consistent matching across datasets. The data were then filtered to include only rows containing *HTR5A-AS1* or its Ensembl ID: ENSG00000220575. TWAS p -values were converted to a numeric format, and the lowest p -values across all brain regions were identified and tested to determine the most significant gene–trait association for *HTR5A-AS1*. This step was performed to establish whether *HTR5A-AS1* showed robust statistical evidence for involvement in schizophrenia across the transcriptome.

2.2. Linkage Disequilibrium Analysis

LD statistics were obtained using the LD1inkR [27] package in R, which provides programmatic access to the NIH LDlink suite [28]. The LDpair() function was used to calculate pairwise LD between the schizophrenia GWAS sentinel SNP (rs1583830) and the lead exon-level eQTL SNP (rs1881691). Analyses were restricted to the European (EUR) reference population from the 1000 Genomes Project Phase 3 under the GRCh38 genome build. LD metrics reported include D' , r^2 , and p -values. This analysis was performed to evaluate whether the schizophrenia risk SNP and the eQTL regulating *HTR5A-AS1* are linked on the same haplotype, suggesting a shared genetic mechanism.

2.3. Postmortem Expression Analysis (BrainSeq)

Bulk RNA-seq expression data containing raw count data and sample metadata from postmortem brain tissue were loaded into R. The SummarizedExperiment [29], dplyr, ggplot2, forcats [30], and ggbeeswarm [31] packages were used to extract expression values for *HTR5A-AS1*, which were converted to counts per million (CPM) using sample-specific library sizes. The samples were filtered to include only adult donors (at least 18 years old) from the hippocampus and dlPFC. Expression data were \log_2 -transformed, and group differences between control and schizophrenia samples were assessed using a Wilcoxon rank-sum test. This analysis directly tested whether *HTR5A-AS1* dysregulation is present in disease-relevant cortical and limbic regions.

2.4. Developmental Expression Trajectories (BrainSpan)

Expression trajectories of *HTR5A-AS1* were derived from BrainSpan bulk RNA-seq data. Transcript abundance values were extracted from the expression matrix using gene annotations from the accompanying metadata. Reported sample ages were parsed into post-conception days (PCD) by converting gestational weeks, months, and years into a continuous day scale, with birth defined at 280 days. Expression values were plotted against \log_{10} -scaled PCD, with vertical reference lines marking birth and selected developmental timepoints.

A linear regression model ($\text{TPM} \sim \text{PCD}$) was fit to quantify the global developmental trend. The regression slope (TPM/day), coefficient of determination (R^2), and p -value were calculated using the broom [32] package and annotated directly onto the plot. The slope serves as a quantitative measure of whether *HTR5A-AS1* expression systematically increased or decreased across human brain development.

Sample metadata were mapped from BrainSpan structure acronyms into broad anatomical classes: NCX, hippocampus, STR, MD, AMY, and CBC. Regions with the six highest median TPM values were retained for visualization. For each of the six regions, linear regression models ($\text{TPM} \sim \text{PCD}$) were fit independently. Slopes (TPM change per post-conception day) were extracted using the broom package, and slope values were embedded into the figure legend for interpretability. Points were overlaid on regression lines, with a \log -scaled x -axis and vertical markers at birth and selected guide marks, enabling identification of the most developmentally dynamic brain regions for *HTR5A-AS1* expression.

2.5. UCSC Genome Browser Validation

The UCSC Genome Browser (hg38 assembly, accessed 2025) was used to confirm the presence and transcriptional structure of *HTR5A-AS1*. The locus was examined with the following tracks enabled: Base Position, database of single nucleotide polymorphisms (dbSNP) build 155 (to visualize genome-wide significant GWAS SNP rs1583830 and the lead eQTL SNP rs1881691), GENCODE v48 annotation, CLS transcript models, Adult Blood/Brain transcript models, Cell Line and Embryonic Brain transcript models, ENCODE4 long-read transcripts, and GTEx tissue expression. A PDF export of the genome browser view was generated to document transcript models and SNP placement relative to *HTR5A-AS1*.

2.6. Functional Predictions (lncHUB)

Functional predictions for *HTR5A-AS1* were obtained from the lncHUB platform [21], including Kyoto Encyclopedia of Genes and Genomics (KEGG) pathway Z-scores, Mammalian/MGI mouse phenotype Z-scores, Gene Ontology (GO) biological process Z-scores, and co-expressed gene Z-scores. The exported result tables were read into R and processed in `dp1yr`; display names were wrapped for readability and converted to title case while preserving acronyms (e.g., GPCR, GABA). For each category, the top entries by Z-score were selected without transformation (KEGG top 15; Mouse Phenotype top 15; GO Biological Process top 20; Co-expression top 25). Ranked horizontal bar charts were generated in `ggplot2`, with bars representing the reported Z-scores and numeric labels printed just beyond bar ends. Axis text was brought closer to bars to improve legibility, and figure margins were adjusted to avoid clipping of labels. No statistical re-analysis or rescaling of Z-scores was performed; ordering and formatting were applied for visualization. All plots were produced in R (packages: `tibble` [33], `dp1yr`, `ggplot2`, `stringr`, `forcats`, `rlang` [34], `grid` [35]) using fixed fill colors and identical geometries across panels to facilitate comparison.

3. Results

3.1. TWAS Identifies *HTR5A-AS1* Association with Schizophrenia

The first step of this study was to determine whether *HTR5A-AS1* is associated with schizophrenia on a genetic and expression level, as establishing such links is critical before pursuing functional interpretation. In the hippocampus, the strongest TWAS association with *HTR5A-AS1* was detected at the junction level ($Z = -3.99$, $p = 6.52 \times 10^{-5}$, $FDR = 2.93 \times 10^{-3}$; Figure 1A) [16]. Junction-level hits indicate that genetic regulation influences exon–exon splicing events, implicating *HTR5A-AS1* isoform usage in disease susceptibility. The lead GWAS variant in this locus, rs1583830, showed robust association with schizophrenia (GWAS $Z = 4.73$, $p = 2.25 \times 10^{-6}$). The corresponding hippocampal eQTL, rs1881691, exerted a strong effect on *HTR5A-AS1* expression (eQTL $Z = -5.71$, $p = 9.15 \times 10^{-5}$; Figure 1B). In the dlPFC, an exon-level feature of *HTR5A-AS1* also reached statistical significance (TWAS $Z = -3.54$, $p = 4.06 \times 10^{-4}$, $FDR = 1.04 \times 10^{-2}$; Figure 1A). Exon-level hits capture expression changes, reflecting the regulatory effects of transcript expression level rather than isoform usage. The same GWAS sentinel single-nucleotide polymorphism (SNP), rs1583830, again showed strong association with schizophrenia, while the lead exon eQTL, rs1881691, exhibited an even stronger effect in this region (eQTL $Z = -6.92$, $p = 9.15 \times 10^{-5}$; Figure 1B). Across both regions, the GWAS sentinel SNP (rs1583830) and the exon-level eQTL SNP (rs1881691) were found to be in moderate-to-strong linkage disequilibrium (LD; $D' = 0.82$, $r^2 = 0.60$, $p = 1.0 \times 10^{-4}$), suggesting that the same haplotype block drives the schizophrenia risk association and *HTR5A-AS1* regulation. Taken together, these results indicate that the haplotype containing rs1583830 and rs1881691 regulates *HTR5A-AS1* expression and splicing, supporting its role as a schizophrenia risk gene.




Figure 1. *HTR5A-AS1* TWAS results and eQTL associations with schizophrenia. (A) Bar plot showing $-\log_{10}(p\text{-value})$ for *HTR5A-AS1* in dorsolateral prefrontal cortex (dlPFC) and hippocampus (HIPPO) from transcriptome-wide association studies (TWAS) analysis [16]. *HTR5A-AS1* showed its strongest association in the HIPPO ($p = 6.52 \times 10^{-5}$, FDR-adjusted $p = 2.93 \times 10^{-3}$), exceeding the false discovery rate (FDR) threshold for significance, and a weaker but notable association in dlPFC ($p = 4.06 \times 10^{-4}$, FDR-adjusted $p = 2.93 \times 10^{-2}$). (B) Expression quantitative trait loci (eQTL) association for *HTR5A-AS1* with lead SNP rs1881691. Scatterplot showing eQTL Z-scores (x-axis) versus $-\log_{10}(p\text{-values})$ (y-axis) across all tested variants. The lead variant rs1881691 (highlighted in red) shows a strong association with *HTR5A-AS1* expression ($Z = -6.92$, $p = 9.15 \times 10^{-5}$) in the HIPPO samples.

3.2. Reduced *HTR5A-AS1* Expression in the Hippocampus

The next step was to test whether *HTR5A-AS1* is differentially expressed in schizophrenia. Analyses of postmortem raw count RNA sequencing (RNA-seq) data from 900 brain samples across hippocampus ($n = 447$) and dlPFC ($n = 453$), from both control and schizophrenia donors, were performed [16]. In the hippocampus, *HTR5A-AS1* was expressed significantly lower ($p = 0.0193$, Wilcoxon rank-sum test; Figure 2A) in schizophrenia than in the control samples, consistent with its negative TWAS Z-score, -3.99 . In the dlPFC, expression was lower in schizophrenia but did not reach statistical significance ($p = 0.0954$, Wilcoxon rank-sum test; Figure 2B). These findings indicate that *HTR5A-AS1* is significantly downregulated in the hippocampus of schizophrenia donors, consistent with genetic predictions of disease risk.




Figure 2. Violin plots of regional *HTR5A-AS1* expression using Collado-Torres et al. (2019) TWAS data [16]. (A) Violin plot showing $\log_2(\text{counts per million [CPM]} + 1)$ *HTR5A-AS1* expression in adult hippocampus (HIPPO) from postmortem schizophrenia and control donors. Data were normalized to CPM using sample-specific library sizes. A Wilcoxon rank-sum test revealed significantly lower expression in schizophrenia ($p = 0.0193$). (B) Violin plot showing $\log_2(\text{CPM} + 1)$ *HTR5A-AS1* expression in adult dorsolateral prefrontal cortex (dlPFC) from postmortem schizophrenia and control donors ($p = 0.0954$, Wilcoxon rank-sum test).

3.3. Region-Specific Expression in the Human Brain

To evaluate whether *HTR5A-AS1* is expressed in regions most consistently disrupted in schizophrenia, its expression patterns across brain regions were analyzed using RNA-seq data from the BrainSpan Atlas of the Developing Human Brain [19]. Expression was quantified as transcripts-per-million (TPM), a normalized metric that controls for sequencing depth and gene length. Expression was highest in the cerebellar cortex (CBC; mean = 5.17 TPM, median = 5.03, $n = 29$), followed by the primary somatosensory cortex (S1C; mean = 3.38 TPM, median = 2.45, $n = 26$), the posterior superior temporal cortex (STC; mean = 3.30 TPM, median = 1.52, $n = 36$), and the inferior parietal cortex (IPC; mean = 3.22 TPM, median = 1.24, $n = 33$). Other notable regions with moderate expression included the inferolateral temporal cortex (ITC; mean = 2.95 TPM, $n = 34$) and the primary motor cortex (M1C; mean = 2.75 TPM, $n = 26$), whereas expression was comparatively lower in the amygdaloid complex (AMY; mean = 1.62 TPM, $n = 33$) and especially the cerebellum (CB; mean = 0.73 TPM,

$n = 3$) (Figure 3A). This analysis confirmed that *HTR5A-AS1* shows strong region-specific enrichment, particularly in cortical and cerebellar regions.

3.4. Developmental Trajectory of *HTR5A-AS1*

To determine whether *HTR5A-AS1* expression, like schizophrenia risk genes, is developmentally regulated, brain RNA-seq data spanning prenatal to adult stages were analyzed from BrainSpan [19]. Developmental analyses of *HTR5A-AS1* showed dynamic changes in expression across the lifespan. Linear regression across all samples revealed an overall positive developmental trajectory ($\beta \approx 0.0003$ TPM/day; Figure 3B). Region-stratified linear fits showed that *HTR5A-AS1* expression rises at distinct rates across brain regions (Figure 3C). The fastest rate of *HTR5A-AS1* expression increase was observed in the CBC ($\beta = 0.000639$ TPM/day), followed by the neocortex (NCX; $\beta = 0.000484$) and striatum (STR; $\beta = 0.000362$). Moderate increases were seen in the AMY ($\beta = 0.000233$) and hippocampus ($\beta = 0.000208$), while the mediodorsal nucleus of the thalamus (MD) showed the lowest rates ($\beta = 0.000045$), suggesting relative developmental stability. Together, these results demonstrate that *HTR5A-AS1* expression is developmentally regulated, with the steepest increases occurring in cortical and cerebellar regions relevant to schizophrenia.




Figure 3. BrainSpan RNA-seq *HTR5A-AS1* expression by brain region, across development [19]. (A) Mean TPM values are shown for selected brain regions, ordered by decreasing expression. (B) Expression of *HTR5A-AS1* increases steadily from prenatal to postnatal stages in BrainSpan RNA-seq data. A linear regression shows a significant positive trajectory (slope = 0.0004 TPM/day; $R^2 = 0.44$; $p = 5.7 \times 10^{-68}$), indicating strong developmental regulation. (C) Region-stratified regression shows distinct developmental patterns.

3.5. Transcript Validation via UCSC Genome Browser and Co-Expression Analysis

To verify that *HTR5A-AS1* is a genuine transcript rather than an artifact of transcriptional noise, its annotation and expression were examined using independent datasets. In the UCSC Genome Browser analysis (hg38 assembly; accessed 2025) [20], long-read RNA-seq tracks confirmed multi-exonic transcripts in brain tissues overlapping the GENCODE-annotated locus. The schizophrenia-associated SNPs appeared upstream of the *HTR5A-AS1* locus, consistent with potential regulatory positioning. These data support *HTR5A-AS1* as a distinct multi-exonic antisense transcript to *HTR5A* (Figure 4B). To test whether *HTR5A-AS1* expression is related to its sense strand partner, *HTR5A*, the same BrainSpan dataset was examined [19]. Across 524 samples, *HTR5A-AS1* expression correlated strongly with *HTR5A* (Pearson's $r = 0.943$, $p < 2.2 \times 10^{-16}$; Figure 4A). These findings confirm that *HTR5A-AS1* is a bona fide transcript and suggest coordinated regulation with its coding partner *HTR5A*.




Figure 4. Co-expression with *HTR5A* and UCSC Genome Browser analysis. (A) Scatterplot showing TPM for *HTR5A-AS1* (y-axis) versus *HTR5A* (x-axis) across samples in the BrainSpan dataset [19], colored by brain region. Pearson's $r = 0.943$, $p < 2.2 \times 10^{-16}$. (B) UCSC Genome Browser (hg38) snapshot of the *HTR5A-AS1* locus on chromosome 7 [20]. Displayed tracks include genome-wide association studies (GWAS) and eQTL sentinel SNPs (rs1583830, red; rs1881691, blue), GENCODE v48 annotation, transcript models, ENCODE4 long-read RNA-seq, and GTEx RNA-seq.

3.6. Predicted Functional Associations of *HTR5A-AS1*

To generate functional hypotheses, pathway enrichment and co-expression analyses were performed using IncHUB [21]. These analyses indicated strong enrichment for pathways related to synaptic transmission, learning, and addiction (Figure 5A). The top KEGG pathway was amyotrophic lateral sclerosis (ALS; $Z = 6.354$), followed by nicotine addiction ($Z = 5.917$) and long-term potentiation ($Z = 5.653$). Predicted mouse phenotypes highlighted associations with abnormal short-term spatial reference memory ($Z = 7.54$), abnormal hearing electrophysiology ($Z = 7.20$), and abnormal peripheral synaptic transmission ($Z = 6.93$; Figure 5B). Gene Ontology (GO) enrichment identified GABAergic synaptic transmission ($Z = 7.943$), regulation of secretion by cell ($Z = 6.449$), and gamma-

aminobutyric acid signaling pathway ($Z = 6.396$) among the top biological processes (Figure 5C). Analysis of the top 500 co-expressed genes revealed many known synaptic and neuronal regulators, including *HTR5A* ($Z = 0.948$), *CAMK2A* ($Z = 0.896$), *SNAP25* ($Z = 0.880$), and *GRIN1* ($Z = 0.871$; Figure 5D). Collectively, these functional predictions implicate *HTR5A-AS1* in synaptic transmission, inhibitory signaling, and cognitive processes.




Figure 5. IncHUB predictions [21]. (A) Horizontal bar chart showing the top 20 Kyoto Encyclopedia of Genes and Genomics (KEGG) pathways predicted by IncHUB based on co-expression patterns. The highest-scoring pathway was amyotrophic lateral sclerosis (ALS) ($Z = 6.354$), followed by nicotine addiction ($Z = 5.917$) and long-term potentiation ($Z = 5.653$). (B) Horizontal bar chart showing the top 22 mouse phenotypes predicted by IncHUB. The top phenotype was abnormal short-term spatial reference memory ($Z = 7.54$), followed by abnormal hearing electrophysiology ($Z = 7.20$) and abnormal PNS synaptic transmission ($Z = 6.93$). (C) Horizontal bar chart showing the top 35 Gene Ontology (GO) biological process predictions from IncHUB. The highest-scoring process was GABAergic synaptic transmission ($Z = 7.943$), followed by regulation of secretion by cell ($Z = 6.449$) and gamma-aminobutyric acid signaling pathway ($Z = 6.396$). (D) Horizontal bar chart showing the top 25 protein-coding genes co-expressed with *HTR5A-AS1* from IncHUB analysis. The highest-ranking gene was *HTR5A* ($Z = 0.948$), followed by *CAMK2A* ($Z = 0.896$), *SNAP25* ($Z = 0.880$), and *GRIN1* ($Z = 0.871$).

4. Discussion

This study provides the first detailed characterization of a lncRNA transcribed antisense to the serotonin receptor gene *HTR5A*, *HTR5A-AS1*, in the context of schizophrenia. TWAS analyses [16] identified two significant associations between *HTR5A-AS1* expression and schizophrenia: a junction-level signal in the hippocampus and an exon-level signal in the dlPFC. Beyond genetic associations, expression analyses revealed significantly reduced *HTR5A-AS1* expression in the hippocampus of

schizophrenia brain donors, aligning with TWAS predictions [16]. While the reduction in the dlPFC did not reach statistical significance, the trend aligns with the negative TWAS association and should be interpreted cautiously. The Wilcoxon rank-sum test, which compares observed expression distributions between diagnostic groups, does not account for technical factors or genetically mediated effects. By contrast, TWAS integrates eQTL and GWAS data, making it more sensitive to genetic associations that may be obscured in bulk RNA-seq analyses. Notably, the TWAS associations were identified at the junction and exon level, suggesting that schizophrenia risk may involve altered isoform usage of *HTR5A-AS1*. Such splicing-dependent effects would be diluted in bulk gene-level differential expression analyses, underscoring the need for isoform-aware approaches.

Using BrainSpan data [19], analyses revealed that *HTR5A-AS1* is developmentally regulated, with distinct region-specific trajectories. Consistent with schizophrenia's developmental origins, *HTR5A-AS1* displayed steep increases in expression in the cerebellar cortex (CBC) and neocortex (NCX), moderate increases in the hippocampus and amygdala, and relative stability in the thalamus. These trajectories suggest that *HTR5A-AS1* may contribute to the maturation of cortical–cerebellar and cortico-limbic circuits, which are strongly implicated in schizophrenia's cognitive and affective dysfunction. Thus, the developmental regulation of *HTR5A-AS1* aligns it with the temporal and regional vulnerability windows most relevant to schizophrenia risk.

The possibility that *HTR5A-AS1* is simply transcriptional noise arising from *HTR5A* was also refuted. This study provides multiple independent lines of evidence to validate the transcript's authenticity: multi-exonic structure validated by long-read RNA-seq, consistent annotation in GENCODE, and strong independent expression signals across multiple datasets [20]. Its co-expression with *HTR5A* is expected for antisense pairs but does not negate potential regulatory function. Instead, it raises the possibility of cis-regulation of *HTR5A*, as well as coordinated involvement in common pathways.

Functional predictions and co-expression analyses consistently implicated synaptic and cognitive processes, particularly GABAergic signaling and long-term potentiation [21]. These findings align with the developmental regulation and cortical–hippocampal enrichment of *HTR5A-AS1*, regions in which disruptions of inhibitory–excitatory balance are strongly implicated in schizophrenia. Together, these results suggest that *HTR5A-AS1* may modulate neuronal excitability or plasticity, potentially by regulating *HTR5A* expression or by interacting with other synaptic genes such as *CAMK2A*, *SNAP25*, and *GRIN1*.

These findings support a model in which *HTR5A-AS1* plays a role in shaping typical brain development and synaptic function. Its developmental trajectory suggests a role in tuning serotonergic and inhibitory circuits during adolescence, consistent with the developmental onset of schizophrenia. In pathology, reduced hippocampal expression of *HTR5A-AS1*, coupled with its genetic associations, raises the possibility that its dysregulation contributes to schizophrenia disease risk by disrupting serotonergic signaling, inhibitory balance, or developmental trajectories in vulnerable brain regions.

There are several implications to be acknowledged. First, most predicted associations obtained from lncHUB are correlative based on co-expression, and functional experiments are required to establish their causality [21]. Second, *HTR5A-AS1*'s high co-expression with *HTR5A* complicates interpretation, as distinguishing cis (local effects on the paired sense transcript) from trans (distal effects on other genes) effects will require CRISPR-mediated perturbation, allele-specific expression assays, or other targeted experiments. Third, sample sizes, while larger than prior work, are still few relative to the polygenic architecture of schizophrenia, and expression variability across cell types is obscured in bulk tissue. Future work should emphasize single-cell and spatial transcriptomic profiling to define cellular specificity, along with experimental manipulations in neuronal models to test regulatory effects on *HTR5A* and synaptic physiology.

Author Contributions: Conceptualization, methodology, software, validation, formal analysis, investigation, resources, data curation, writing—original draft, writing—review and editing, visualization, project administration: Grant H. Rutenber. The author has read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Publicly available datasets were analyzed in this study. BrainSpan data are available at <https://www.brainspan.org/>, GTEx data at <https://gtexportal.org/>, and Collado-Torres et al. (2019) data at <https://doi.org/10.1016/j.neuron.2019.05.013>. Custom scripts used for processing and plotting are available from the author upon request.

Conflicts of Interest: The author declares no conflicts of interest.

References

1. Schizophrenia. Available online: <https://www.who.int/news-room/fact-sheets/detail/schizophrenia> (accessed on 8 September 2025).
2. Solmi, M.; Seitidis, G.; Mavridis, D.; Correll, C.U.; Dragioti, E.; Guimond, S.; Tuominen, L.; Dargél, A.; Carvalho, A.F.; Fornaro, M.; et al. Incidence, Prevalence, and Global Burden of Schizophrenia—Data, with Critical Appraisal, from the Global Burden of Disease (GBD) 2019. *Mol. Psychiatry* **2023**, *28*, 5319–5327. <https://doi.org/10.1038/s41380-023-02138-4>.
3. Hany, M.; Rizvi, A. Schizophrenia. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2025.
4. Marder, S.R.; Cannon, T.D. Schizophrenia. *N. Engl. J. Med.* **2019**. <https://doi.org/10.1056/NEJMra1808803>.
5. Chick, S.L.; Holmans, P.; Cameron, D.; Grozeva, D.; Sims, R.; Williams, J.; Bray, N.J.; Owen, M.J.; O'Donovan, M.C.; Walters, J.T.R.; Rees, E. Whole-Exome Sequencing Analysis Identifies Risk Genes for Schizophrenia. *Nat. Commun.* **2025**, *16*, 1–9. <https://doi.org/10.1038/s41467-025-62429-y>.
6. Functional Genomics Reveal Gene Regulatory Mechanisms Underlying Schizophrenia Risk. *Nat. Commun.* **2019**. Available online: <https://www.nature.com/articles/s41467-019-08666-4> (accessed on 16 August 2025).
7. Birnbaum, R.; Weinberger, D.R. Genetic Insights into the Neurodevelopmental Origins of Schizophrenia. *Nat. Rev. Neurosci.* **2017**, *18*, 727–740. <https://doi.org/10.1038/nrn.2017.125>.
8. Karlsgodt, K.H.; Sun, D.; Cannon, T.D. Structural and Functional Brain Abnormalities in Schizophrenia. *Curr. Dir. Psychol. Sci.* **2010**, *19*, 226–231. <https://doi.org/10.1177/0963721410377601>.
9. Quednow, B.B.; Geyer, M.A.; Halberstadt, A.L. Serotonin and Schizophrenia. In *Handbook of Behavioral Neuroscience*; Müller, C.P., Cunningham, K.A., Eds.; Elsevier: Amsterdam, The Netherlands, 2020; Volume 31, pp. 711–743. <https://doi.org/10.1016/B978-0-444-64125-0.00039-6>.
10. Li, P.; Snyder, G.L.; Vanover, K.E. Dopamine Targeting Drugs for the Treatment of Schizophrenia: Past, Present and Future. *Curr. Top. Med. Chem.* **2016**, *16*, 3385–3403. <https://doi.org/10.2174/1568026616666160608084834>.
11. Meltzer, H.Y.; Li, Z.; Kaneda, Y.; Ichikawa, J. Serotonin Receptors: Their Key Role in Drugs to Treat Schizophrenia. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **2003**, *27*, 1159–1172. <https://doi.org/10.1016/j.pnpbp.2003.09.010>.
12. Grubor, M.; Zivkovic, M.; Sagud, M.; Nikolac Perkovic, M.; Mihaljevic-Peles, A.; Pivac, N.; Muck-Seler, D.; Svob Strac, D. HTR1A, HTR1B, HTR2A, HTR2C and HTR6 Gene Polymorphisms and Extrapyramidal Side Effects in Haloperidol-Treated Patients with Schizophrenia. *Int. J. Mol. Sci.* **2020**, *21*, 2345. <https://doi.org/10.3390/ijms21072345>.
13. Thomas, D.R. 5-HT_{5A} Receptors as a Therapeutic Target. *Pharmacol. Ther.* **2006**, *111*, 707–714. <https://doi.org/10.1016/j.pharmthera.2005.12.006>.
14. Zhang, Y.; Smith, E.M.; Baye, T.M.; Eckert, J.V.; Abraham, L.J.; Moses, E.K.; Kissebah, A.H.; Martin, L.J.; Olivier, M. Serotonin (5-HT) Receptor 5A Sequence Variants Affect Human Plasma Triglyceride Levels. *Physiol. Genomics* **2010**, *42*, 168–176. <https://doi.org/10.1152/physiolgenomics.00038.2010>.
15. Gusev, A.; Ko, A.; Shi, H.; Bhatia, G.; Chung, W.; Penninx, B.W.J.H.; Jansen, R.; de Geus, E.J.; Boomsma, D.I.; Wright, F.A.; et al. Integrative Approaches for Large-Scale Transcriptome-Wide Association Studies. *Nat. Genet.* **2016**, *48*, 245–252. <https://doi.org/10.1038/ng.3506>.
16. Collado-Torres, L.; Burke, E.E.; Peterson, A.; Shin, J.; Straub, R.E.; Rajpurohit, A.; Semick, S.A.; Ulrich, W.S.; Price, A.J.; Valencia, C.; et al. Regional Heterogeneity in Gene Expression, Regulation, and Coherence in the Frontal Cortex and Hippocampus across Development and Schizophrenia. *Neuron* **2019**, *103*, 203–216.e8. <https://doi.org/10.1016/j.neuron.2019.05.013>.

17. National Center for Biotechnology Information. Available online: <https://www.ncbi.nlm.nih.gov/> (accessed on 17 September 2025).
18. GeneCards. Available online: <https://www.genecards.org/> (accessed on 17 September 2025).
19. BrainSpan Atlas of the Developing Human Brain. Available online: <https://www.brainspan.org/> (accessed on 8 September 2025).
20. UCSC Genome Browser. Available online: <https://genome.ucsc.edu/> (accessed on 17 September 2025).
21. IncHUB. Available online: <https://maayanlab.cloud/Inchub/> (accessed on 17 September 2025).
22. R: The R Project for Statistical Computing. Available online: <https://www.r-project.org/> (accessed on 17 September 2025).
23. readr: Read Rectangular Text Data. Available online: <https://CRAN.R-project.org/package=readr> (accessed on 17 September 2025).
24. dplyr: A Grammar of Data Manipulation. Available online: <https://CRAN.R-project.org/package=dplyr> (accessed on 17 September 2025).
25. Wickham, H. *ggplot2*; Springer International Publishing: Cham, Switzerland, 2016. <https://doi.org/10.1007/978-3-319-24277-4>.
26. stringr: Simple, Consistent Wrappers for Common String Operations. Available online: <https://CRAN.R-project.org/package=stringr> (accessed on 17 September 2025).
27. Myers, T.A.; Chanock, S.J.; Machiela, M.J. LDlinkR: An R Package for Rapidly Calculating Linkage Disequilibrium Statistics in Diverse Populations. *Front. Genet.* **2020**, *11*, 157. <https://doi.org/10.3389/fgene.2020.00157>.
28. Lin, S.-H.; Thakur, R.; Machiela, M.J. LDexpress: An Online Tool for Integrating Population-Specific Linkage Disequilibrium Patterns with Tissue-Specific Expression Data. *BMC Bioinformatics* **2021**, *22*, 608. <https://doi.org/10.1186/s12859-021-04531-8>.
29. SummarizedExperiment: Summarized Experiment Container. Available online: <https://bioconductor.org/packages/SummarizedExperiment> (accessed on 17 September 2025).
30. forcats: Tools for Working with Categorical Variables (Factors). Available online: <https://CRAN.R-project.org/package=forcats> (accessed on 17 September 2025).
31. ggbeeswarm: Categorical Scatter (Violin Point) Plots. Available online: <https://CRAN.R-project.org/package=ggbeeswarm> (accessed on 17 September 2025).
32. broom: Convert Statistical Objects into Tidy Tibbles. Available online: <https://CRAN.R-project.org/package=broom> (accessed on 17 September 2025).
33. tibble: Simple Data Frames. Available online: <https://CRAN.R-project.org/package=tibble> (accessed on 17 September 2025).
34. rlang: Functions for Base Types and Core R and Tidyverse Features. Available online: <https://CRAN.R-project.org/package=rlang> (accessed on 17 September 2025).
35. R Core Team. grid: The Grid Graphics Package. R package version bundled with R. Available online: <https://stat.ethz.ch/R-manual/R-devel/library/grid/doc/grid.pdf> (accessed on 17 September 2025).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.