

Article

Not peer-reviewed version

---

# Carbon Saving Potential of Urban Parks in the Central Plains City: A High Spatial Resolution Study Using a Forest City, Shangqiu, China as a Lens

---

[Jianwei Gao](#) <sup>\*,†</sup>, [Haiting Han](#) <sup>+</sup>, [Shidong Ge](#) <sup>\*</sup>

Posted Date: 2 June 2023

doi: [10.20944/preprints202306.0137.v1](https://doi.org/10.20944/preprints202306.0137.v1)

Keywords: urban park; carbon saving potential; high spatial resolution; Shangqiu



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# Carbon Saving Potential of Urban Parks in the Central Plains City: A High Spatial Resolution Study Using a Forest City, Shangqiu, China as a Lens

Jianwei Gao <sup>1,\*†</sup>, Haiting Han <sup>2,†</sup> and Shidong Ge <sup>3,\*</sup>

<sup>1</sup> School of Economics, Tianjin University of Commerce, Tianjin 300134, China; gjw0923@tjcu.edu.cn

<sup>2</sup> Københavns Universitet, Rolighedsvej 23, Copenhagen, 1953 FC, Denmark; hanht@pku.edu.cn

<sup>3</sup> College of Landscape Architecture and Art and International Union Laboratory of Landscape Architecture, Henan Agricultural University, Zhengzhou 450002, China; shidongge@henau.edu.cn

\* Correspondence: gjw0923@126.com; shidongge@henau.edu.cn

† These authors contributed equally to this work.

**Abstract:** This article investigates the potential for carbon reduction in urban parks in Shangqiu City using high-resolution remote sensing imagery. The aim is to guide modern urban carbon neutrality strategies. The carbon reduction potential is estimated based on the mitigation of the urban heat island effect by park greenery, which reduces energy consumption. The parks are regarded as cool island centers, and 100 cooling gradients are gradually formed outward until the temperature reaches that of impermeable urban surfaces. The energy saved by the parks in mitigating the heat island effect is statistically calculated as the carbon reduction potential of urban park greenery. Additionally, this article classifies the sample parks into different categories and selects 26 landscape metrics to analyze their relationship with carbon-saving potential and driving factors.

**Keywords:** urban park; carbon saving potential; high spatial resolution; Shangqiu

## 1. Introduction

Changes in urban land cover have caused dramatic alterations in local urban microclimates, leading to a series of ecological issues. As urbanization continues to accelerate, extreme weather events and the urban heat island (UHI) effect have become hot topics among academics. The UHI effect refers to the phenomenon in which urban areas have significantly higher temperatures than adjacent rural areas. Although the theory was proposed early on, it remains a topic of great interest in the academic community, with many unresolved issues even after extensive research into its causes and mitigation strategies [1]. Some studies suggest that the UHI effect leads to the formation of local air circulation within cities, carrying air pollutants to higher altitudes and posing serious threats to human health [2]. In addition, a strong UHI effect can greatly impact the comfort of urban living conditions [3]. Especially in summer, the UHI effect is particularly pronounced, increasing the use of air conditioning and leading to higher energy consumption and carbon emissions. Therefore, reducing the UHI effect is a crucial component in achieving urban carbon neutrality.

Urban park green spaces have a significant impact on mitigating the urban heat island effect [4,5]. Many scholars have recognized the positive role of urban parks in offsetting this effect and creating "cool islands" within cities [6]. There are several reasons for this. First, vegetation can provide shade, reducing direct solar radiation on the ground and decreasing the absorption of heat radiation by the ground. Second, through the transpiration of plants and their low albedo, some of the solar radiation can be absorbed. Finally, blue-green infrastructure, including urban green spaces, can effectively reduce the surface temperature of adjacent areas by directing airflow within the city. Therefore, enhancing the "cool island" effect within urban areas is considered an effective way to mitigate the heat effect [7].


In the past, most studies on urban green spaces have focused on their direct carbon sequestration capacity, which increases as vegetation biomass increases [8]. For example, in Xi'an, urban green spaces have an annual fixed carbon amount of  $0.19 \text{ t C ha}^{-1}$  [9], while in Beijing, Changchun, and Harbin the average carbon densities are  $7.8 \text{ t C ha}^{-1}$  [10],  $8.7 \text{ t C ha}^{-1}$  [11], and  $7.7 \text{ t C ha}^{-1}$  [12], respectively. However, other studies have pointed out that the role of vegetation in reducing energy consumption in urban areas and bringing about carbon emission reductions cannot be ignored [13,14]. An empirical study showed that park green space has great potential to reduce carbon emissions and could save  $23.7 \pm 1.6 \text{ t CO}_2$  in the Yangtze River Economic Belt region [13]. Therefore, it is critical to understand the potential carbon emission reduction capacity through reducing thermal environmental pressure. However, the potential carbon emission reduction capacity of urban green spaces has received limited attention, especially in central China.

This study examines the potential for reducing carbon emissions in urban parks in Shangqiu, a city in Central China. Our objective is to provide guidance for urban development and construction, from the perspective of urban ecology and environmental economics. Specifically, we aim to answer the following research questions: 1) What is the magnitude and spatial distribution of carbon saving potential in Shangqiu's urban parks? 2) What are the landscape factors driving spatial heterogeneity in the carbon saving potential across different urban parks in Shangqiu?

## 2. Materials and Methods

### 2.1. Study Area

Shangqiu is located in the eastern Henan province of China between longitudes  $114^{\circ}49' \sim 116^{\circ}39'$  E and latitudes  $33^{\circ}43' \sim 34^{\circ}52'$  N (Figure 1). It has a total area of 10,704 square kilometers and a population of 7.723 million. It is a national civilized city, national health city, national garden city, and national forest city, with tourist attractions like the ancient city of Shangqiu. The city's plains account for 99.2% of its total area, with only 0.8% being hilly areas. Shangqiu has a warm temperate semi-humid continental monsoon climate with distinct four seasons, ample sunshine, and abundant rainfall. The city has an annual average of 1944 hours of sunshine, an average annual temperature of  $14.6^{\circ}\text{C}$ , an average annual rainfall of 736.2 millimeters, and a frost-free period of about 211 days. Since its establishment in 1997, Shangqiu has undergone rapid urbanization, with significant expansion of construction and transportation land use. From 2000 to 2015, the city's construction land use expanded rapidly, increasing by as much as 696.57 square kilometers, which accounted for 37.57% of the total construction land use in 2000. This rapid urbanization has led to significant changes in the city's landscape pattern, making it an ideal area to study the potential for carbon reduction in urban areas.



**Figure 1.** Location and land cover classification of Shangqiu City, Henan, China.

## 2.2. Remote-Sensed Urban Park

The Split Window algorithm is a commonly used method in the field of land surface temperature (LST) remote sensing retrieval. It uses the ratio of reflectance in two spectral bands to remove atmospheric effects and obtain the land surface radiation temperature, which is then converted to LST using thermal radiation principles [15,16]. This algorithm applies to remote sensing image data of different resolutions and spectral bands and has the advantages of high accuracy and strong operability [17]. In this study, the original data was obtained from Landsat 8 image data set with a spatial resolution of 30 meters. All remote sensing images of cloudless and sunny days were selected from the summers of 2020-2021 (Table 1).

**Table 1.** Information on remotely sensed data for Shangqiu.

Satellite	Path/Row	Period(Year-Month-Day)
Landsat	122/36	2020-08-28
Landsat	122/36	2020-09-04
Landsat	122/36	2021-06-28
Landsat	122/36	2021-07-30
Landsat	122/36	2021-09-16
GF-2	--	2021-08-10

The radiative transfer equation is expressed as follows:

$$L_\lambda = [\varepsilon \cdot B(T_s) + (1 - \varepsilon) \cdot L_\downarrow] \cdot \tau + L_\uparrow \quad (1)$$

The radiative transfer equation used in the Split-Window algorithm for calculating land surface temperature (LST).  $L_\lambda$  represents the thermal radiation intensity of wavelength  $\lambda$  received by the satellite sensor,  $\varepsilon$  is the land surface emissivity,  $B(T_s)$  is the radiation brightness received by a blackbody with temperature  $T_s$ , and its unit is  $\text{W} \cdot \text{m}^{-2} \cdot \text{sr}^{-1} \cdot \mu\text{m}^{-1}$ .  $\tau$  is the atmospheric transmissivity, and  $L_\uparrow$  and  $L_\downarrow$  are the upwelling and downwelling atmospheric radiances obtained from NASA [18]. This equation, based on the principles of thermal radiation, considers the effects of the atmosphere on radiation and can be used to retrieve land surface temperature from satellite data. It is a crucial component of the Split-Window algorithm.

Later, we used the GF-2 image (Table 1) and the method of Object-based Image classification to interpret the land use in the study area. To improve the accuracy of interpretation, we optimized the samples based on auxiliary data and combined manual corrections by researchers to modify the parts with significant deviation, maximizing the accuracy of land use data. We divided the land use into five categories: forest (including trees and shrubs), grass, bare land (including sandy and soil surfaces), water bodies (including ponds, streams, etc.), and impervious surfaces (including buildings, squares, and other facilities).

Based on high-resolution remote sensing images, the 118 parks in Shangqiu City can be classified according to different features and classification standards. Firstly, according to the nature of the park and the urban land classification standard (CJJ/T85-2017), parks can be classified into community parks, small parks, specific parks, and comprehensive parks. Secondly, according to the size of the park's area, it can be divided into miniature parks (less than 2 hectares), small parks (2-5 hectares), medium parks (5-10 hectares), medium-large parks (10-50 hectares) and large parks (more than 50 hectares). Thirdly, the presence of water bodies within the park is also a classification criterion that can divide parks into those with and without water bodies. Through these classification criteria, we can have a more comprehensive understanding of the distribution and characteristics of parks in Shangqiu City.

## 2.3. Landscape Metrics

Landscape indices are statistical measures used to describe and quantify landscape patterns. They are commonly used to analyze the impact of different land use types and landscape

characteristics on ecosystems, including biodiversity, ecosystem services, and landscape functionality. In this study, a set of landscape indices was selected based on the classification criteria of Fragstats 4.2 software, including:

Patch Density (PD), Number of Patches (NP), Patch Area Fraction (PAFRAC), Mean Patch Area (PARA\_MN), Mean Fragment Shape Index (FRAC\_MN), Mean Patch Shape Index (SHAPE\_MN), Mean Patch Area (AREA\_MN), Largest Patch Index (LSI), Aggregation Index (AI), Cohesion Index (COHESION), Interspersion and Juxtaposition Index (IJI), Simpson's Evenness Index (SHEI), Simpson's Diversity Index (SHDI), Simpson's Dominance Index (SIDI), Percentage of Forest, Grass, Impervious surfaces and water(PLAND\_Forest, PLAND\_Grass, PLAND\_Imper, and PLAND\_Water).

These indices were selected to investigate the effect of landscape patterns on land surface temperature. They represent the fragmentation, connectivity, and diversity of the landscape within the study area.

#### 2.4. Quantification of UPSs' Carbon Saving Potential

This section aims to estimate the carbon emission reduction potential of urban green spaces by calculating the carbon emission reduction resulting from the alleviation of the urban heat island effect in city parks. Following the previous study [13,14], we calculate the carbon saving intensity (CSI) and the carbon emission reduction efficiency (CSE):

$$CSI = k \cdot \rho \cdot a \cdot \int_0^H \sum_{i=0}^N \frac{1}{3} (S_i + S_{i+1} + \sqrt{S_i S_{i+1}}) \Delta T dh \quad (2)$$

$$CSE = CSI \div S \quad (3)$$

Among them,  $k$  represents the specific heat at constant pressure, with a value of  $1004.68 \text{ J kg}^{-1} \text{ }^{\circ}\text{C}^{-1}$ ,  $\rho$  represents the air density ( $1.2923 \text{ kg} \cdot \text{m}^{-3}$ ),  $a$  is the conversion coefficient of energy consumption into carbon emissions from coal-fired power generation (841 g/3.6 MJ).  $h$  represents the vertical influence range ( $H=70\text{m}$ ).  $\Delta T$  is the temperature difference between adjacent buffer zones,  $S_i$  and  $S_{i+1}$  are the base areas and  $S$  is the park area.

#### 2.5. Correlation between UPSs' Carbon Saving Potential and Landscape Metrics


This article proposes a method for studying the correlation between the carbon-saving intensity of urban parks and their influencing factors. The method uses various statistical analysis methods, including the Shapiro-Wilk test, analysis of variance (ANOVA), independent sample t-test, Pearson correlation analysis, and stepwise multiple linear regression. First, the Shapiro-Wilk test is used to ensure that the sampled data set conforms to a normal distribution. Then, ANOVA and independent sample t-tests are used to analyze whether there are significant differences in the carbon-saving intensity of urban parks under different park classifications. In this process, the average carbon-saving intensity of different parks is taken as the dependent variable, and park classification calibration is taken as the factor. Pearson correlation analysis is used to determine whether there is a statistically significant correlation between various landscape indices and carbon saving intensity, as well as if the correlation is obvious. Stepwise multiple linear regression is used to establish the relationship between carbon saving intensity and landscape patterns, the best-fitting model is selected, and the significance of the coefficients is determined based on regression statistics ( $R^2$ , p-value). This study analyzes landscape indices that can explain the intensity of carbon saving and proposes strategies for addressing urban heat island effects based on these indices. To eliminate the influence of collinearity between independent factors, the variance inflation factor (VIF) is used to assess the model, and independent variables with collinearity are removed to obtain the final regression model. In conclusion, this method provides a comprehensive way to analyze factors that

influence carbon-saving intensity in urban parks and can be used to propose strategies for achieving carbon neutralization.

### 3. Results

#### 3.1. Spatial Heterogeneity of Carbon Saving Potential in Different Urban Parks

For CSI, Shangqiu City Park Green Space (538 ha) can reduce a total of 300.57 t CO<sub>2</sub> emissions. Specifically, in the parks of Shangqiu, the CSI distribution range is 0.04~18.93 t, and on average, each park can save  $2.55 \pm 0.31$  t CO<sub>2</sub> emissions for the city because of its contribution to the mitigation of surface heat effects, of which the canal ribbon park has the highest CSI, reaching 18.93 t CO<sub>2</sub>. Among the parks divided according to different classification criteria, special parks (3.42 t), parks larger than 10 ha (6.94 t), and parks with water bodies (4.71 t) had the largest CSI average in their respective classifications. From the efficiency perspective, the CSE distribution in Shangqiu City was 16.34~0.04 t CO<sub>2</sub> ha<sup>-1</sup>, and the average CSE was  $1.79 \pm 0.29$  t CO<sub>2</sub> ha<sup>-1</sup>. Among them, the average CSE of amusement parks (2.03 t CO<sub>2</sub> ha<sup>-1</sup>), parks less than 2ha (2.83 t CO<sub>2</sub> ha<sup>-1</sup>), and anhydrous parks (2.04 t CO<sub>2</sub> ha<sup>-1</sup>) had larger CSE, while smaller parks showed larger CSE and the CSE of Chinese Fatong Square (0.10 ha) was the highest (16.34 t CO<sub>2</sub> ha<sup>-1</sup>). Fig.2 shows the results of the analysis of the difference between CSI and CSE in different classifications of parks, and there are certain differences in the performance of CSI in different types of parks, but the CSE of each park does not show significant differences, and this difference is not reflected in all park classifications. According to the classification of different areas, the CSI of parks less than 2 ha is significantly lower than that of parks with an area of 5-10 ha and parks larger than 10 ha, the CSI of parks with an area of 2-5 ha is significantly lower than that of parks with an area greater than 5ha, and the CSI of parks with an area greater than 10 ha is significantly larger. Among the parks classified according to whether the park contains water bodies, the CSI of water-bearing parks is significantly higher than that of non-water-bearing parks.




**Figure 2.** Park category differences between CSI and CSE. a, b, c, d, represent significant differences determined by Fisher's least significant difference (LSD) tests ( $p < 0.05$ ) on different seasons for different park types.

#### 3.2. Spatial Changes in Landscape Metrics

In the study area, the park area was distributed in 0.03~357.00 ha, and the proportion of various landscapes in the park was different, from large to small, they were woodland (59.42%), grassland (23.09%), impervious surface (22.20%) and water body (12.34%). We selected 15 landscape pattern indexes to explore the differences in surface landscape patterns of various parks, and on the whole, Shangqiu City Park has a high degree of fragmentation, a high degree of aggregation, and landscape diversity, but the shape of various landscape patches is more complex. In different types of parks, there are certain differences in landscape indicators (Figure 3). The analysis of the differences in the landscape pattern indices of different parks shows that the landscape indices do not show significant

differences when the parks are divided according to the park category, and among the parks with different areas, the AI, SHDI, and COHESION of parks with an area of less than 2ha are significantly larger than those with an area greater than 2ha, and COHESION is the opposite. Parks with an area greater than 5 ha SHAPE\_MN are significantly larger than parks with an area of less than 2ha, LSI and NP are significantly larger than parks with an area of less than 5ha in parks with an area of 5-10ha, and LSI, NP, and PLAND\_Imper are significantly smaller in parks with an area of less than 2ha. LSI, NP, SPLIT, AREA\_MN increased significantly with the increase of park area when the park area was less than 10ha, and on the contrary, PD decreased significantly with the increase of area when the area was greater than 2ha. In waterless parks and water-bearing parks, the values of AI, LPI, COHESION, PD, and PLAND\_Forest in waterless parks were significantly greater than those in waterless parks, while SHDI, SPLIT, AREA\_MN, LSI, and NP were the opposite, and the values in waterless parks were significantly larger.




**Figure 3.** Park category differences of landscape patterns. a, b, c, d, represent significant differences determined by Fisher's least significant difference (LSD) tests ( $p < 0.05$ ) on different seasons for different park types.

### 3.3. The Relationship between the Carbon Saving Potential and Landscape Driving Factors in Different Urban Parks

There were similarities and differences in the correlation results between various park landscape indexes and CSI (Figure 4). On the whole, CSI was significantly positively correlated with SPLIT, IJI, AREA\_MN, LSI, NP, and PLAND\_Water, and significantly negatively correlated with LPI and PLAND\_Imper. Among the different types of parks, the CSI of AREA\_MN and general parks (-0.952) and community parks (-0.869) showed a strong and significant positive correlation, while


PLAND\_imper was the opposite. The CSI of the garden was positively correlated with NP, IJI, and SPLIT, but negatively correlated with LPI. The CSI of special parks showed a significantly strong positive correlation with AREA\_MN (0.952), NP (0.92), and PLAND\_water (0.905), while it showed a significantly strong negative correlation with FRAC\_MN (-0.989), and the CSI of comprehensive parks showed a correlation with the most landscape index, among which there was a strong positive correlation with AREA\_MN (0.952) and CONTAG (0.856) and a strong positive correlation with PLAND\_Imper (-0.957), PLAND\_Grass (-0.818) showed a strong negative correlation. Among parks of different sizes, the number of landscape indices associated with CSI was slightly smaller than in the other two categories, but the correlation between them was strongest. The CSI of 2-5ha parks was significantly positively correlated with SHEI (0.995), SHDI (0.995), and PLAND\_Grass (0.998), the CSI and AREA\_MN of parks with an area greater than 2ha showed a significant positive correlation, and the CSI of parks with an area greater than 10ha showed a different correlation from other parks, and the CSI was significantly positively correlated with NP (0.717) and PLAND\_Water (0.813). It was significantly negatively correlated with PLAND\_imper (-0.591). In the classification according to whether the park has a water body, there are more landscape pattern indices related to CSI, and the correlation between anhydrous parks and waterless parks and landscape pattern indexes is similar, and there are significant positive correlations with SPLIT, AREA\_MN, LSI, and NP, among which the positive correlation between CSI and NP in waterless parks is the strongest (0.733), while the negative correlation between anhydrous parks and LPI is the strongest (-0.385). The difference is that the significant correlation between IJI (0.37), PLAND\_Water (0.644), and PLAND\_Imper (-0.348) and waterless parks is not reflected in waterless parks.



**Figure 4.** Heatmaps of correlation coefficients between landscape patterns and CSI in different types of parks.

Figure 5 shows the correlation between CSE and landscape pattern index of various parks in Shangqiu City, and the results show that the correlation between CSE and landscape pattern index is similar to the CSI trend overall, both in parks classified by category, the most relevant landscape pattern index, followed by parks classified according to water bodies, and the correlation between CSE and landscape pattern in parks classified by area is weak. The difference is that the landscape pattern index related to CSE of various parks is completely different from CSI, and AI, COHESION,

and park CSE are significantly negatively correlated, while PARA\_MN and PD are significantly positively correlated with park CSE. Among different types of parks, the landscape pattern indexes related to CSE of community parks were SHEI (0.809), SHDI (0.809), PARA\_MN (0.831), and PLAND\_Grass (0.945), respectively, and all showed positive correlations. In contrast, CSE in integrated parks was only negatively correlated with PLAND\_Grass significantly (-0.836). CSE WAS SIGNIFICANTLY NEGATIVELY CORRELATED WITH AI AND COHESION IN AMUSEMENT PARKS AND SPECIAL PARKS, AND POSITIVELY CORRELATED WITH PARA\_MN AND PD, and this correlation was stronger in special parks. In addition, SHAPE\_MN (-0.458), AREA\_MN (-0.466), and amusement park CSE were significantly negatively correlated, while special park CSE and LSI were negatively correlated and positively correlated with PLAND\_Imper (0.986). Among the parks with different areas, park CSEs less than 2 ha were positively correlated with NP (0.85), and park CSEs with 5-10 ha also showed a positive correlation with AREA\_MN (0.734). Among watered parks and waterless parks, the correlation between landscape pattern index and CSE was more consistent, significantly negatively correlated with AI and COHESION, and this correlation was stronger in water-bearing parks, and the significant positive correlation with PD was stronger in waterless parks, and the difference was reflected in the significant positive correlation between CSE and PARA\_MN (0.406) in waterless parks, while CSE in waterless parks was positively correlated with PLAND\_Grass (0.292).




**Figure 5.** Heatmaps of correlation coefficients between landscape patterns and CSE in different types of parks.


### 3.4. Identify the Landscape-Driving Factors

For all the landscape pattern variables examined in our study, the direction and magnitude of their impact on park CSI and CSE were generally different (Figures 6 and 7). Moreover, the specific influencing factors are different from the direction and degree of their influence on CSI and CSE to different park categories, and the overall changes of CSI and CSE in different parks have a high degree of explanation. The fragmentation index has a more common effect on the changes of CSI and CSE in various parks, while the degree of explanatory degree of diversity index on CSI and CSE changes is significantly higher than that of other factors. The overall interpretation rate of CSI of various landscape pattern indexes can reach 83%, of which NP contributed the largest 52% of the

interpretation rate, followed by AREA\_MN (17.39%), PD (9.41), and FRAC\_MN (4.2%). Among the different types of parks, the regression model of CSI for comprehensive parks was the best, with a goodness-of-fit of 99%, and the fitting degree of the model for special parks was also high, 98.03%. Among them PLAND\_Imper contributed the highest explanation rate (88.7%) for the CSI of the comprehensive park, PARA\_MN contributed 10.3% of the explanation rate, the largest contributor in the regression model of the special park was FRAC\_MN (94.7%), and SPLIT also explained a small part of the CSI change. The CSI model fit was slightly weak, but the influencing factors were relatively balanced, and SPLIT contributed 51.8% of the explanation rate and PLAND\_Forest contributed 11.8%. In parks with different areas, except for parks with an area of less than 2ha, the park model with a smaller area has a higher fit and clearer influencing factors. The park regression model of 2-5ha had the highest goodness-of-fit (98.6%), and the vast majority of explanatory degrees were contributed by PLAND\_Grass (97.8%), while SHDI explained the model only 0.8%. For 5-10ha parks, two variables, AREA\_MN and LSI, were introduced into the regression model, with explanatory rates of 75.5% and 22.2%, respectively. Among the parks with an area greater than 10 ha, the interpretation of CSI was PLAND\_Water (63.7%), AREA\_MN (13.82%), and NP (5.36%), respectively. Compared with the above classification, the CSI driving factors of the anhydrous park and watery park are scattered, and the four influencing factors are introduced into the waterless park regression model, which is FRAC\_MN (4.16%), AREA\_MN (17.39%), PD (9.41%), and NP (52%), and the overall model goodness-of-fit reached 83%. The goodness-of-fit of the CSI regression model for anhydrous parks was 50.5%, of which SPLIT contributed 40.4% of the explanatory degrees.



**Figure 6.** Stepwise regression diagram of landscape pattern and CSI in different parks.



**Figure 7.** Stepwise regression diagram of landscape pattern and CSE in different parks.

The results of CSE stepwise regression in various parks (Figure 7) showed that compared with the CSI regression model, the CSE model introduced fewer variables and the overall explanatory rate was not high. All parks were included in the model, and the model introduced COHESION as the only explanatory variable, with an explanatory rate of 44.2%. Among the different types of parks, community parks also introduced PLAND\_Grass as the only explanatory variable, but the explanatory degree reached 85.7%, and PD in amusement parks and special parks were the most explanatory landscape pattern indicators, with 93.1% and 96.8% explanatory degrees, respectively. The goodness-of-fit of the amusement park CSE regression model was 97.7%, and in addition to PD AREA\_MN and PLAND\_imper also contributed 3.99% and 0.78% of the explanatory degrees, respectively. IJI was also introduced into the special park CSE model to provide an explanatory rate of 0.2%. Classified by area, none of the factors were introduced into the regression model due to confounding factors. COHESION and PD were introduced as the only explanatory variables in the regression model of the anhydrous park and watery park, and the goodness-of-fit was 44.2% and 43.1%, respectively.

#### 4. Discussion

##### 4.1. The Carbon-Saving Potential of Urban Parks

The results showed that the average CSE of Shangqiu City Park was  $1.79 \pm 0.29 \text{ t CO}_2 \text{ ha}^{-1}$ , which was higher than the previous CSE study of the Yangtze River Economic Belt City Park in China ( $1.08 \pm 0.03 \text{ t CO}_2 \text{ ha}^{-1}$ ). First, studying the size difference of the region may be one of the reasons for the difference in CSE. The study area is a city-level city in Henan Province, with a total of 118 parks. The previous study area included 1,510 parks in 26 cities, spanning 11 provinces in China. A higher number of parks studied directly means greater differences in size between parks, many types of parks, and more complex basic information about parks. So it's understandable that the average CSE for 1510 parks is lower than the average CSE for 118 parks. On the other hand, the size of the study area will cause differences in the results of urban LST results, and may also indirectly affect the calculation results of CSI and CSE. For example, a study of Bangkok, Jakarta, and Manila showed

that LST studies presented different results at different spatial resolutions, and proposed that 210 m×210 m is an optimal characteristic area or land climate footprint that can be used for examining any meteorological, climatic, or environmental issues in urban areas or for landscape and urban planning [19]. The results for the 960 m scale and the 240 m scale are completely different [20]. It is worth noting that the previous research results also show that the average CSE of small urban parks is generally high, which is partly consistent with the results of this study.

Second, the climate type of the study area is also responsible for the difference in CSE. Previous results on the Yangtze River Economic Belt showed that the average CSE of parks with humid subtropical climates was higher than that of parks located in humid subtropical monsoon climates. The climate of Shangqiu City is a temperate monsoon climate, and the difference in climatic characteristics is more obvious. Different climatic characteristics lead to different LST research results and indirectly lead to different CSEs [21,22].

Third, studying differences in regional economic development, population density, and urbanization levels will also lead to differences in the average CSE of parks. Several studies have pointed out that economic development, population growth, and urbanization are important causes of rising surface temperature [23]. Higher levels of urbanization mean more natural features such as vegetation and water bodies that have been replaced by impervious materials and buildings [24]. Economic activity and population agglomeration consume large amounts of fossil fuel resources and exacerbate the urban thermal environment [25]. Compared to Shangqiu City, the Yangtze River Economic Belt, the former study area, has the most prosperous and dense urbanization performance in China, which also means a denser population [26]. In contrast, Shangqiu City, Henan Province, located in central China, is lower than the Yangtze River Economic Belt in terms of economy, population density, and urbanization level. This directly leads to the different urban heat island effects between the two, and also indirectly leads to the difference in the average CSE of the park.

Fourth, Shangqiu City's emphasis on park construction has also greatly increased the average CSE of the park. Shangqiu City was awarded the honorary title of "National Garden City" as early as 2010. Shangqiu City takes the "14th Five-Year Plan for Urban Renewal and Urban and Rural Habitat Environment Construction of Shangqiu City" as the program, intending to build a green city and ecological city. Continuously strengthen the urban landscaping construction in Shangqiu City, continue to pay attention to the incremental quality improvement of park green space, and improve the carbon emission reduction capacity of Shangqiu City Park.

In addition, this study shows that the annual CSE of the park green space in Shangqiu City is 154.8 t CO<sub>2</sub> ha<sup>-1</sup> year<sup>-1</sup>. Another result of our ongoing study shows that the annual average Carbon Serum in Zhengzhou Parkland is 19.07 t CO<sub>2</sub> ha<sup>-1</sup> year<sup>-1</sup>. Another study based on the rate of major Chinese cities showed that the average annual carbon sequestration of green infrastructure in Zhengzhou was 10.52 t CO<sub>2</sub> ha<sup>-1</sup> year<sup>-1</sup> [27]. This study shows that the carbon emission reduction efficiency of Shangqiu City is 8~14 times that of carbon sequestration in the green space of the two Zhengzhou parks. This strongly indicates that the cooling effect of the park's green space leads to significant carbon reduction.

#### 4.2. Effects of Landscape Patterns on the Carbon Saving Potential

In general, the impact of the park's landscape pattern on the park's carbon reduction potential is significant. The park's carbon reduction capacity is equivalent to the park's cooling and energy-saving effects. This result is therefore consistent with previous studies confirming the cooling effect of landscape patterns on parks [22,28~31]. Specifically, the main drivers of CSI in Shangqiu City Park are NP, AREA\_MN, and PD. In other words, the number of park patches and the degree of landscape fragmentation (characterized by AREA\_MN and PD) significantly affected the park CSI. For the number of patches (NP), it can be interpreted that a smaller, more plaque in the park can play a role in cooling energy saving and carbon reduction. A study of 197 water bodies in Beijing showed that tiny lakes and ponds play an important role in cooling. Therefore, it is suggested that decomposing parks with large water areas into smaller ones can improve the cooling effect of parks, and the same applies to park carbon reduction [32]. The impact on landscape fragmentation depends on the type

of land use. For example, the AREA\_MN of impervious surface panel blocks has an increasing effect on LST, while plant plaques have the opposite effect. Today's increased urbanization has led to the fragmentation of impervious surfaces and patches of green space, resulting in the fragmentation of the urban landscape pattern. By reducing the fragmentation of the urban landscape, urban LST can be lowered and the carbon reduction potential of parks can be increased [31].

In addition, the impact of the PLAND index on the CSI and CSE of Shangqiu City parks was shown in different categories of parks, including PLAND\_Grass, PLAND\_Water, PLAND\_Imper, and PLAND\_Forest. This result is supported by many LST studies [20,33,34]. Even if the study area is different from the meteorological environment, the enhancement effect of PLAND\_Imper on park LST and the cooling effect of PLAND\_Grass, LAND\_Forest, and PLAND\_Water on LST due to high reflectivity, transpiration, providing shade to reduce cooling and its specific heat capacity have been widely demonstrated [35–38]. As a park's carbon reduction potential, which is closely related to the park's cooling and energy-saving effects, it will also be affected by the PLAND index.

But contrary to Hao Hou's index that Shape\_MN is the best-performing cooling effect [29], Shape\_MN does not affect Shangqiu City's CSI. This inconsistency is normal in LST studies. Different study areas and different climatic conditions significantly affect the performance of landscape indicators [20,39,40]. Therefore, when exploring the relationship between landscape patterns and the effects of park cooling energy saving and carbon emission reduction, it is necessary to refer to the natural environmental factors of the study area [39–41].

#### 4.3. Implications for Urban Planning and Management

This study explores the impact of landscape patterns on CSI and CSE of different types of urban park green space, the specific relationship, and its potential value in reducing carbon emissions from the perspective of carbon emission reduction caused by the cooling effect of green space. Rapid urbanization has caused an imbalance in land resource distribution, leading to the need for rational allocation of the internal landscape pattern of parks. This is important in maximizing the cooling effect of urban green spaces and achieving the strongest carbon emission reduction efficiency in the context of carbon neutrality in China. The study focuses on several parks and green spaces in the study area to understand the impact of the internal landscape pattern on the efficiency and intensity of urban carbon emission reduction at the regional scale. This will provide guidance and suggestions for upgrading and renovating urban parks. Previous research shows that landscape pattern change greatly affects the surface thermal environment of urban parks and reduces carbon emissions. Urban planners can explore the strongest mode of urban park green space to reduce carbon emissions by updating and transforming the urban park landscape pattern through design.

The results showed that the CSI and CSE of the park differed with the change in landscape pattern, and this difference was manifested in different types of parks. Overall, FRAC\_MN, PLAND\_Forest, PLAND\_Grass, and NP were all significant influencing factors with positive effects on CSI and CSE. This indicates that parks with a larger proportion of trees and ground cover plants, as well as more complex patch shapes, have a better potential for carbon saving. These findings are consistent with previous research in the field of the thermal environment [42–44]. The proportion of water in parks has a significant positive impact on CSI, as water bodies have a high specific heat capacity. Small lakes or ponds, in particular, play an important role in carbon emission reduction [45]. Parks with water, special parks, and parks larger than 10 hectares show significant carbon reduction effects. Therefore, water design and transformation should be the focus of upgrading and renovation in the aforementioned types of parks. Our research results indicate that changes in various landscape patterns in different types of parks have varying degrees of impact on carbon emission reduction. This paper proposes targeted park improvement and update suggestions for park categories with relatively low carbon emission reduction in the study area. These include reorganizing landscapes and changing the area and location of different landscape types. Suggestions for transforming different types of park green spaces with weaker carbon emission reduction capacity were also proposed to adjust the density of patches and the shape of parks, to accurately promote the carbon emission reduction of park green spaces in Shangqiu.

The diversity index has a significant positive impact on the CSE of the integrated park, and the increase in PLAND\_Imper can bring about a significant decrease in CSI. Therefore, when transforming comprehensive parks, the focus should be on reducing the proportion of impervious surface area in the park and increasing the diversity of landscape patches. This can include introducing a variety of plant communities into a single green space and adding small water bodies, such as fountains and pools, to waterless parks. The CSI of LSI amusement park has a significant negative impact, and the complexity of the park boundary provides an opportunity for energy exchange between the park and the surrounding area, thereby increasing the cooling effect of the park to a certain extent [46], so the design and transformation of the park should focus on the change of the shape of the park boundary. The increase of AREA\_MN and PLAND\_Grass can significantly improve the carbon emission reduction capacity of parks and community parks with an area of 2-5 ha [47], so attention should be paid to improving the cover of surface grassland in the renovation of such parks, to enhance the carbon storage of vegetation and thus improve the carbon emission reduction capacity of parks. For parks with an area of 2-5 ha, it is also possible to improve the carbon-saving capacity of the park by increasing the type of landscape patch. In the waterless park, PD, SPLIT, and AREA\_MN all showed a significant positive relationship with carbon emission reduction capacity, which indicates that the degree of plate fragmentation has the most significant impact on the sewage park, so attention should be paid to dividing the internal patches of the park to make them as dispersed as possible to achieve greater carbon emission reduction efficiency.

#### 4.4. Limitations and Future Research Directions

In this paper, 19 landscape pattern indicators were selected to comprehensively describe the morphology, patch characteristics, fragmentation degree, and aggregation degree of the park, and the selected variables explained the changes in CSI (83%) and CSE (44.2%). However, our study has some limitations in some aspects, and the influence mechanism of landscape patterns on CSI and CSE of urban parks and green spaces needs further study. First of all, it should be recognized that this study is based on a 2D plane, and the impact of landscape changes in the vertical 3D range, including green amount, water depth, and other factors on CSI and CSE, needs to be refined and improved in future studies, and the specific configuration of plant communities in the park, the physiological and ecological indicators of vegetation and the vertical and planar structure of surrounding features and other factors that have a potential impact on the cooling effect of urban park green space are included in the research of carbon emission reduction and park enhancement strategy of urban parks. Comprehensively consider the impact of influencing factors on carbon emission reduction from various perspectives.

Secondly, the research on the carbon emission reduction effect of urban parks in this paper is based on surface temperature. Due to the difficulty in acquiring high-quality continuous meteorological data, this paper relied only on remote sensing images of cloudless and sunny summer days in 2020-2021 to measure the carbon emission reduction of urban park green spaces. Actual results may be biased due to this limitation. Future research should explore higher precision surface temperature data combined with field survey data to supplement remote sensing images. It should also introduce multi-source data from seasonal changes in surface temperature, daily dynamic changes, interannual changes, urban and rural changes, and other multi-temporal and spatial perspectives to explore the changes in land surface temperature and its impact on carbon emission reduction efficiency in a comprehensive manner. This approach will clarify the specific impact of landscape pattern index, urban park CSI and CSE, and pay attention to the practical significance and practical effect of research. It will also promote the combination of research results and urban construction strategies, and conduct in-depth research on the carbon emission reduction effect of urban green spaces in practice.

### 5. Conclusions

Currently, there is extensive research on mitigating the surface heat island effect in urban parks in the context of carbon neutrality. However, it is of significant importance to quantify the carbon

emission reduction resulting from the mitigation of urban park thermal effects, as it contributes to achieving urban carbon neutrality goals. Additionally, it is crucial to quantitatively analyze the influence of urban park landscape patterns on carbon emission reduction intensity and efficiency. In this study, we estimated the carbon emission reduction intensity and efficiency of 118 urban parks located in Shangqiu City, Henan Province. The average CSI was found to be  $2.55 \pm 0.31$  t CO<sub>2</sub>, and the CSE was  $1.79 \pm 0.29$  t CO<sub>2</sub> per hectare across all studied parks. Consequently, a total carbon emission reduction of 300.57 t CO<sub>2</sub> was achieved. Parks larger than 10 hectares and parks with water features exhibited higher carbon emission reduction intensity, indicating that the landscape layout of these parks is more conducive to carbon reduction compared to other park types. Among all park categories, the proportion of trees, herbs, and water bodies significantly influenced carbon conservation. Furthermore, the concentration of landscape patches, including cohesion, split, and aggregation index (AI), played a crucial role in the CSI and CSE of urban parks. Increased fragmentation also led to a stronger carbon emission reduction effect. Based on these findings, we propose a series of strategic suggestions for the renovation and improvement of different park types, aiming to enhance the carbon emission reduction intensity of urban park green spaces through landscape pattern transformations at the regional scale. These suggestions provide theoretical support and practical guidance for urban planning, renovation, and renewal efforts, contributing to carbon emission reduction, mitigation of the urban thermal environment, and enhancement of ecological benefits in the Central Plains region. Ultimately, this research promotes the acceleration of the carbon neutrality process in the Central Plains.

**Author Contributions:** Conceptualization, J.G. and S.G.; methodology, J.G., H.H. and S.G.; software, H.H. and S.G.; validation, J.G. and H.H.; formal analysis, S.G. and H.H.; investigation, S.G. and H.H.; resources, J.G. and S.G.; data curation, J.G., H.H. and S.G.; writing—original draft preparation, J.G., H.H. and S.G.; writing—review and editing, J.G., H.H. and S.G.; visualization, J.G. and S.G.; supervision, J.G. and S.G.; project administration, J.G. and H.H.; funding acquisition, J.G. and S.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study was supported by the Social Science Fund of Tianjin of China grant (TJYY19-013), the Key Technology R&D Program of Henan Province (232102320190), and the Center for Blockchains and Electronic Markets funded by the Carlsberg Foundation under grant no. CF18-1112, and the Special Fund for Young Talents in Henan Agricultural University under grant no.30500930.

## References

1. Dudorova, N.V.; Belan, B.D. The Energy Model of Urban Heat Island. *Atmosphere* **2022**, *13*, 457, doi:10.3390/atmos13030457.
2. Stathopoulou, E.; Mihalakakou, G.; Santamouris, M.; Bagiorgas, H.S. On the impact of temperature on tropospheric ozone concentration levels in urban environments. *J Earth Syst Sci* **2008**, *117*, 227–236, doi:10.1007/s12040-008-0027-9.
3. Santamouris, M.; Kolokotsa, D. On the impact of urban overheating and extreme climatic conditions on housing, energy, comfort and environmental quality of vulnerable population in Europe. *Energy and Buildings* **2015**, *98*, 125–133, doi:10.1016/j.enbuild.2014.08.050.
4. O’Malley, C.; Piroozfar, P.; Farr, E.R.P.; Pomponi, F. Urban Heat Island (UHI) mitigating strategies: A case-based comparative analysis. *Sustainable Cities and Society* **2015**, *19*, 222–235, doi:10.1016/j.scs.2015.05.009.
5. Chun, B.; Guldmann, J.-M. Spatial statistical analysis and simulation of the urban heat island in high-density central cities. *Landscape and Urban Planning* **2014**, *125*, 76–88, doi:10.1016/j.landurbplan.2014.01.016.
6. Qian, Y.; Zhou, W.; Hu, X.; Fu, F. The Heterogeneity of Air Temperature in Urban Residential Neighborhoods and Its Relationship with the Surrounding Greenspace. *Remote Sensing* **2018**, *10*, 965, doi:10.3390/rs10060965.
7. Li, X.; Zhou, W. Optimizing urban greenspace spatial pattern to mitigate urban heat island effects: Extending understanding from local to the city scale. *Urban Forestry & Urban Greening* **2019**, *41*, 255–263, doi:10.1016/j.ufug.2019.04.008.
8. Huang, C.; Ye, X.; Deng, C.; Zhang, Z.; Wan, Z. Mapping Above-Ground Biomass by Integrating Optical and SAR Imagery: A Case Study of Xixi National Wetland Park, China. *Remote Sensing* **2016**, *8*, 647, doi:10.3390/rs8080647.
9. Yao, Z.; Liu, J.; Zhao, X.; Long, D.; Wang, L. Spatial dynamics of aboveground carbon stock in urban green space: a case study of Xi'an, China. *J. Arid Land* **2015**, *7*, 350–360, doi:10.1007/s40333-014-0082-9.

10. Sun, Y.; Xie, S.; Zhao, S. Valuing urban green spaces in mitigating climate change: A city-wide estimate of aboveground carbon stored in urban green spaces of China's Capital. *Glob Change Biol* **2019**, *25*, 1717–1732, doi:10.1111/gcb.14566.
11. Ren, Z.; Zheng, H.; He, X.; Zhang, D.; Shen, G.; Zhai, C. Changes in spatio-temporal patterns of urban forest and its above-ground carbon storage: Implication for urban CO<sub>2</sub> emissions mitigation under China's rapid urban expansion and greening. *Environment International* **2019**, *129*, 438–450, doi:10.1016/j.envint.2019.05.010.
12. Lv, H.; Wang, W.; He, X.; Xiao, L.; Zhou, W.; Zhang, B. Quantifying Tree and Soil Carbon Stocks in a Temperate Urban Forest in Northeast China. *Forests* **2016**, *7*, 200, doi:10.3390/f7090200.
13. Chen, M.; Jia, W.; Du, C.; Shi, M.; Henebry, G.M.; Wang, K. Carbon saving potential of urban parks due to heat mitigation in Yangtze River Economic Belt. *Journal of Cleaner Production* **2023**, *385*, 135713, doi:10.1016/j.jclepro.2022.135713.
14. Lin, W.; Wu, T.; Zhang, C.; Yu, T. Carbon savings resulting from the cooling effect of green areas: A case study in Beijing. *Environmental Pollution* **2011**, *159*, 2148–2154, doi:10.1016/j.envpol.2011.02.035.
15. Jiang, Y.; Lin, W. A Comparative Analysis of Retrieval Algorithms of Land Surface Temperature from Landsat-8 Data: A Case Study of Shanghai, China. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5659, doi:10.3390/ijerph18115659.
16. Jimenez-Munoz, J.C.; Sobrino, J.A.; Skokovic, D.; Mattar, C.; Cristobal, J. Land Surface Temperature Retrieval Methods From Landsat-8 Thermal Infrared Sensor Data. *IEEE Geosci. Remote Sens. Lett.* **2014**, *11*, 1840–1843, doi:10.1109/LGRS.2014.2312032.
17. Sekertekin, A.; Bonafoni, S. Land Surface Temperature Retrieval from Landsat 5, 7, and 8 over Rural Areas: Assessment of Different Retrieval Algorithms and Emissivity Models and Toolbox Implementation. *Remote Sens.* **2020**, *12*, 294, doi:10.3390/rs12020294.
18. Sobrino, J.A.; Jiménez-Muñoz, J.C.; Paolini, L. Land surface temperature retrieval from LANDSAT TM 5. *Remote Sensing of Environment* **2004**, *90*, 434–440, doi:10.1016/j.rse.2004.02.003.
19. Estoque, R.C.; Murayama, Y.; Myint, S.W. Effects of landscape composition and pattern on land surface temperature: An urban heat island study in the megacities of Southeast Asia. *Science of The Total Environment* **2017**, *577*, 349–359, doi:10.1016/j.scitotenv.2016.10.195.
20. Du, C.; Song, P.; Wang, K.; Li, A.; Hu, Y.; Zhang, K.; Jia, X.; Feng, Y.; Wu, M.; Qu, K.; et al. Investigating the Trends and Drivers between Urbanization and the Land Surface Temperature: A Case Study of Zhengzhou, China. *Sustainability* **2022**, *14*, 13845, doi:10.3390/su142113845.
21. Jia, W.; Zhao, S. Trends and drivers of land surface temperature along the urban-rural gradients in the largest urban agglomeration of China. *Science of The Total Environment* **2020**, *711*, 134579, doi:10.1016/j.scitotenv.2019.134579.
22. Zhang, K.; Yun, G.; Song, P.; Wang, K.; Li, A.; Du, C.; Jia, X.; Feng, Y.; Wu, M.; Qu, K.; et al. Discover the Desirable Landscape Structure of Urban Parks for Mitigating Urban Heat: A High Spatial Resolution Study Using a Forest City, Luoyang, China as a Lens. *International Journal of Environmental Research and Public Health* **2023**, *20*, 3155, doi:10.3390/ijerph20043155.
23. Grimm, N.B.; Faeth, S.H.; Golubiewski, N.E.; Redman, C.L.; Wu, J.; Bai, X.; Briggs, J.M. Global Change and the Ecology of Cities. *Science* **2008**, *319*, 756–760, doi:10.1126/science.1150195.
24. Chang, Y.; Xiao, J.; Li, X.; Middel, A.; Zhang, Y.; Gu, Z.; Wu, Y.; He, S. Exploring diurnal thermal variations in urban local climate zones with ECOSTRESS land surface temperature data. *Remote Sensing of Environment* **2021**, *263*, 112544, doi:10.1016/j.rse.2021.112544.
25. Sun, Y.; Zhang, X.; Ren, G.; Zwiers, F.W.; Hu, T. Contribution of urbanization to warming in China. *Nature Clim Change* **2016**, *6*, 706–709, doi:10.1038/nclimate2956.
26. Liu, J.; Tian, Y.; Huang, K.; Yi, T. Spatial-temporal differentiation of the coupling coordinated development of regional energy-economy-ecology system: A case study of the Yangtze River Economic Belt. *Ecological Indicators* **2021**, *124*, 107394, doi:10.1016/j.ecolind.2021.107394.
27. Chen, W.Y. The role of urban green infrastructure in offsetting carbon emissions in 35 major Chinese cities: A nationwide estimate. *Cities* **2015**, *44*, 112–120, doi:10.1016/j.cities.2015.01.005.
28. Maimaitiyiming, M.; Ghulam, A.; Tiyip, T.; Pla, F.; Latorre-Carmona, P.; Halik, Ü.; Sawut, M.; Caetano, M. Effects of green space spatial pattern on land surface temperature: Implications for sustainable urban planning and climate change adaptation. *ISPRS Journal of Photogrammetry and Remote Sensing* **2014**, *89*, 59–66, doi:10.1016/j.isprsjprs.2013.12.010.
29. Hou, H.; Estoque, R.C. Detecting Cooling Effect of Landscape from Composition and Configuration: An Urban Heat Island Study on Hangzhou. *Urban Forestry & Urban Greening* **2020**, *53*, 126719, doi:10.1016/j.ufug.2020.126719.
30. Effati, F.; Karimi, H.; Yavari, A. Investigating effects of land use and land cover patterns on land surface temperature using landscape metrics in the city of Tehran, Iran. *Arab J Geosci* **2021**, *14*, 1240, doi:10.1007/s12517-021-07433-4.

31. Jia, J.; Liu, Y.; Li, H.; Wu, J. Seasonal contrast of the dominant factors for spatial distribution of land surface temperature in urban areas. *Remote Sensing of Environment* **2018**, *215*, 255–267, doi:10.1016/j.rse.2018.06.010.
32. Sun, R.; Chen, L. How can urban water bodies be designed for climate adaptation? *Landscape and Urban Planning* **2012**, *105*, 27–33, doi:10.1016/j.landurbplan.2011.11.018.
33. Chen, A.; Yao, L.; Sun, R.; Chen, L. How many metrics are required to identify the effects of the landscape pattern on land surface temperature? *Ecological Indicators* **2014**, *45*, 424–433, doi:10.1016/j.ecolind.2014.05.002.
34. Yang, L.; Yu, K.; Ai, J.; Liu, Y.; Lin, L.; Lin, L.; Liu, J. The Influence of Green Space Patterns on Land Surface Temperature in Different Seasons: A Case Study of Fuzhou City, China. *Remote Sensing* **2021**, *13*, 5114, doi:10.3390/rs13245114.
35. Zhou, W.; Qian, Y.; Li, X.; Li, W.; Han, L. Relationships between land cover and the surface urban heat island: seasonal variability and effects of spatial and thematic resolution of land cover data on predicting land surface temperatures. *Landscape Ecol* **2014**, *29*, 153–167, doi:10.1007/s10980-013-9950-5.
36. Cai, Q.; Li, E.; Jiang, R. Analysis of the Relationship Between Land Surface Temperature and Land Cover Changes Using Multi-temporal Satellite Data. *Nature Environment and Pollution Technology* **2017**, *16*, 8.
37. Yang, Q.; Huang, X.; Li, J. Assessing the relationship between surface urban heat islands and landscape patterns across climatic zones in China. *Sci Rep* **2017**, *7*, 9337, doi:10.1038/s41598-017-09628-w.
38. Shi, Y.; Liu, S.; Yan, W.; Zhao, S.; Ning, Y.; Peng, X.; Chen, W.; Chen, L.; Hu, X.; Fu, B.; et al. Influence of landscape features on urban land surface temperature: Scale and neighborhood effects. *Science of The Total Environment* **2021**, *771*, 145381, doi:10.1016/j.scitotenv.2021.145381.
39. Chapman, S.; Watson, J.E.M.; Salazar, A.; Thatcher, M.; McAlpine, C.A. The impact of urbanization and climate change on urban temperatures: a systematic review. *Landscape Ecol* **2017**, *32*, 1921–1935, doi:10.1007/s10980-017-0561-4.
40. Du, H.; Wang, D.; Wang, Y.; Zhao, X.; Qin, F.; Jiang, H.; Cai, Y. Influences of land cover types, meteorological conditions, anthropogenic heat and urban area on surface urban heat island in the Yangtze River Delta Urban Agglomeration. *Science of The Total Environment* **2016**, *571*, 461–470, doi:10.1016/j.scitotenv.2016.07.012.
41. Zhao, L.; Lee, X.; Smith, R.B.; Oleson, K. Strong contributions of local background climate to urban heat islands. *Nature* **2014**, *511*, 216–219, doi:10.1038/nature13462.
42. Rakoto, P.Y.; Deilami, K.; Hurley, J.; Amati, M. Revisiting the cooling effects of urban greening: Planning implications of vegetation types and spatial configuration. *Urban For. Urban Green.* **2021**, *64*, 127266, doi:10.1016/j.ufug.2021.127266.
43. Xiang, Y.; Ye, Y.; Peng, C.; Teng, M.; Zhou, Z. Seasonal variations for combined effects of landscape metrics on land surface temperature (LST) and aerosol optical depth (AOD). *Ecological Indicators* **2022**, *138*, 108810, doi:10.1016/j.ecolind.2022.108810.
44. Xiao, R.; Cao, W.; Liu, Y.; Lu, B. The impacts of landscape patterns spatio-temporal changes on land surface temperature from a multi-scale perspective: A case study of the Yangtze River Delta. *Science of The Total Environment* **2022**, *821*, 153381, doi:10.1016/j.scitotenv.2022.153381.
45. Wang, X.; Cheng, H.; Xi, J.; Yang, G.; Zhao, Y. Relationship between Park Composition, Vegetation Characteristics and Cool Island Effect. *Sustainability* **2018**, *10*, 587, doi:10.3390/su10030587.
46. Chen, M.; Jia, W.; Du, C.; Shi, M.; Henebry, G.M.; Wang, K. Carbon saving potential of urban parks due to heat mitigation in Yangtze River Economic Belt. *Journal of Cleaner Production* **2023**, *385*, 135713, doi:10.1016/j.jclepro.2022.135713.
47. Zheng, J.; Tarin, M.W.K.; Chen, G.; Zhang, Q.; Deng, C. The characteristics of plant clusters influence on the cooling effect: A case study in a subtropical Island Park, China. *Global Ecology and Conservation* **2022**, *34*, e02055, doi:10.1016/j.gecco.2022.e02055.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.