

Article

Not peer-reviewed version

Perturbed FLRW Metric Explains the Difference in Measurements of the Hubble Constant

[Wladimir Belayev](#)*

Posted Date: 10 April 2025

doi: 10.20944/preprints202404.0626.v4

Keywords: perturbed FLRW metric; Einstein equations; particle Lagrangian; Hubble tension; gravitational lensing; distance ladder



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Perturbed FLRW Metric Explains the Difference in Measurements of the Hubble Constant

Wladimir Belayev

Affiliation 1 ;bvl2879@gmail.com

Abstract: The expansion of the universe yields two consistent differing values of a Hubble constant depending on the methods of measurement. The aim of this work was to explain the Hubble tension by properties of space corresponding to a radially inhomogeneous metrics. A cosmological model described by this perturbed FLRW metric has zero pressure and exponentially expanding space. But the observed space appears to be compressed in the radial direction. We consider the dependence of the change rate of photon energy on the direction of its motion. It is found applying the principle of the photon's energy integral extremum and the relationship between it and the energy of a material particle obtained using Lagrange mechanics. The change rate of a photon's energy varies depending on whether it moves in a radial direction or has an angular component. The metric coefficients are determined by the difference in measurements of the Hubble constant using gravitational lensing and a distance ladder.

Keywords: Perturbed FLRW metric, Einstein equations, particle Lagrangian, Hubble tension, gravitational lensing; distance ladder

1. Introduction

In the Λ CDM (Lambda Cold Dark Matter) cosmological model, based on the FLRW (Friedmann-Lemaître-Robertson-Walker) metric, matter moves synchronously with the expanding space. Data from the JWST telescope allowed the identification of galaxies with a redshift of $z > 10$ [1]-[3], which led to a departure from the existing Big Bang scenario [4].

The perturbed FLRW metric [5]-[8], describing small deviations from the background space-time, can also serve as the basis for certain cosmological models. Deviations from standard cosmology are investigated by considering flat FLRW space-time in Painlevé–Gullstrand coordinates [9]. With this coordinate choice the space is not expanding, although the distance between galaxies is certainly increasing. In Refs. [10,11] is studied a distorted stereographic projection of hyperconical universes, which are 4D hypersurfaces with linear expansion of space embedded into 5D Minkowski spacetime. This model takes the perspective of explaining Hubble tension [12]. We propose model, which is described by perturbed FLRW metric with spatial inhomogeneity in radial direction.

Determination of the Hubble constant using the cosmic distance ladder [13,14] and gravitational lensing [15,16] yields a difference of about 1/10 of its value. We consider the dependence of the change rate of photon energy on the direction of its motion. It is found applying the principle of the extremum of the photon's energy integral [17]-[21] and the relationship between it and the energy of a material particle obtained using Lagrange mechanics [17].

2. The Solution of the Einstein Equations for the Perturbed Metric

The most common presentation of the spatially flat FLRW cosmology is in terms of the explicit line element

$$ds^2 = c^2 dt^2 - a(t)^2 [dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\varphi^2)], \quad (1)$$

where t is the time measured by an observer associated with the 4-velocity vector $u^i = (1, 0, 0, 0)$. A generalization of this metric is obtained by transforming the time interval

$$cd\hat{t} = cdt - f(t)rdr, \quad (2)$$

where function $f(t)$ has dimension inverse length. The linear element takes the following form:

$$ds^2 = c^2 dt^2 - 2cf(t)rdrdt - (a(t)^2 - f(t)^2 r^2)dr^2 - a(t)^2 r^2 (d\theta^2 + \sin^2 \theta d\varphi^2). \quad (3)$$

For small r , this metric is approximated by the metric of flat expanding spacetime. The components of the Einstein tensor [22] are given by

$$G^{11} = \frac{1}{a^6} \left\{ 3a^4 \dot{a}^2 - 6a^3 f \dot{a} + r^2 \left[2(a^2 f^2 - a^3 f \dot{a}) \dot{f} + 2af^3 \dot{a} - 4a^2 f^2 \dot{a}^2 \right] + r^4 \left[f^4 \dot{a}^2 + 2af^3 \dot{a} \dot{f} \right] \right\}, \quad (4)$$

$$G^{12} = G^{21} = \frac{fr}{a^6} \left[(2af \dot{a} + f \dot{a}^2)r + (2af + 2f \dot{a} - 3a \dot{a}^2)a \right], \quad (5)$$

$$G^{22} = \frac{1}{a^6} \left\{ -2a^3 \ddot{a} - a^2 \dot{a}^2 + 2a^2 \dot{f} + 2af \dot{a} + r^2 (2af \dot{a} \dot{f} + f^2 \dot{a}^2) \right\}, \quad (6)$$

$$G^{33} = G^{44} \sin^2 \theta = \frac{1}{a^6 r^2} \left\{ -2a^3 \ddot{a} - a^2 \dot{a}^2 + 2a^2 \dot{f} + 2af \dot{a} + r^2 (a \ddot{a} f^2 - f^2 \dot{a}^2 + f a^2 \ddot{f} + a^2 \dot{f}^2 + a f \dot{a} \dot{f}) \right\}, \quad (7)$$

where $(\dot{})$ denotes differentiation with respect to $x^1 = ct$.

The components of the contravariant energy-momentum tensor correspond to the energy density, momentum density, energy flux through unit surfaces per unit time and stresses [23]. The energy-momentum tensor of matter in contravariant form is given by

$$T^{ij} = (c^2 \rho + p) u^i u^j - g^{ij} p. \quad (8)$$

The non-zero contravariant coefficients for the metric (3) will be as follows:

$$g^{11} = \frac{a^2 - f^2 r^2}{a^2}, \quad g^{12} = g^{21} = -\frac{fr}{a^2}, \quad g^{22} = -\frac{1}{a^2},$$

$$g^{33} = -\frac{1}{a^2 r^2}, \quad g^{44} = -\frac{1}{a^2 r^2 \sin^2 \theta}. \quad (9)$$

In the absence of angular momentum of matter ($u^3 = u^4 = 0$), the non-zero components of the energy-momentum tensor (8) are given by

$$T^{11} = (c^2 \rho + p) (u^1)^2 - \left(1 - \frac{f^2 r^2}{a^2} \right) p, \quad (10)$$

$$T^{12} = T^{21} = (c^2 \rho + p) u^1 u^2 + \frac{fr}{a^2} p, \quad (11)$$

$$T^{22} = (c^2 \rho + p) (u^2)^2 + \frac{1}{a^2} p, \quad (12)$$

$$T^{33} = T^{44} \sin^2 \theta = \frac{1}{a^2 r^2} p. \quad (13)$$

Let's consider a model with a cosmological constant $\Lambda = 0$. We will seek a solution for a region where the radial velocity u^2 is small, and the term of component T^{22} containing its square is negligible and can be disregarded. Next, we will justify this assumption. Also, we assume that the terms of the

Einstein tensor components (4)-(7) containing r^2 in the numerator are small and can be neglected. Given the satisfaction of this condition and for g^{11} , Einstein equations $G^{ij} = \chi T^{ij} + g^{ij}\Lambda$ with the energy-momentum tensor components (10)-(13), yield

$$\frac{3a\dot{a}^2 - 6f\dot{a}}{a^3} = \chi c^2 \rho, \quad (14)$$

$$\frac{2a\dot{f} + 2f\dot{a} - 3a\dot{a}^2}{a^3} f r = \chi a^2 (c^2 \rho + p) u^1 u^2 + \chi f r p, \quad (15)$$

$$\frac{-2a^2\ddot{a} - a\dot{a}^2 + 2a\dot{f} + 2f\dot{a}}{a^3} = \chi p. \quad (16)$$

The absence of additional energy and momentum influx implies the equality of the corresponding densities T^{12}, T^{21} to zero, i.e., the right-hand side of equation (15). From its left-hand side we obtain

$$2a\dot{f} + 2f\dot{a} - 3a\dot{a}^2 = 0. \quad (17)$$

Solving this equation for f under the condition $a(t_0) = 1$ we find

$$f(t) = \frac{1}{a} \left(f_0 + \frac{3}{2} c \int_{ct_0}^{ct} \dot{a}^2 a dt \right) \quad (18)$$

with $f(t_0) = f_0$. Substituting this into equation (14) under the condition

$$\rho = \frac{\rho_0}{a(t)^3}, \quad (19)$$

with average density of the Universe at the present time $\rho_0 = \rho(t_0)$ yields

$$3a\dot{a}^2 - \frac{6f_0\dot{a}}{a} - \frac{9\dot{a}}{a} c \int_{ct_0}^{ct} \dot{a}^2 a dt = \chi c^2 \rho_0. \quad (20)$$

Expressing the integral from here and differentiating it with respect to x^1 , from the obtained equality we find

$$\chi c^2 \rho_0 \left(1 - \frac{a\ddot{a}}{\dot{a}^2} \right) = -3\dot{a}^2 a + 3\ddot{a} a^2. \quad (21)$$

This equation has a solution

$$a(t) = e^{cA_1(t-t_0)}, \quad (22)$$

where A_1 is a constant.

From equations (16) and (17), we obtain the pressure

$$p = -\frac{2}{c\chi} \frac{d}{dt} \left(\frac{\dot{a}}{a} \right), \quad (23)$$

which, due to (22), turns out to be $p = 0$. In this case, from (15) under the condition of the absence of energy and momentum influx from the outside (17), we find the radial component of the matter velocity in the comoving reference frame.

$$u^2 = 0. \quad (24)$$

Thus, it turns out to be a valid assumption that the radial velocity in the region under consideration is small and may not be taken into account in the component of the energy-momentum tensor (12).

The function (18), with the scale factor of space (22), takes the form

$$f(t) = e^{-cA_1(t-t_0)} f_0 + \frac{1}{2} A_1 \left(e^{2cA_1(t-t_0)} - e^{-cA_1(t-t_0)} \right) \quad (25)$$

and equation (20) yields

$$3A_1^2 - 6f_0A_1 = \chi c^2 \rho_0. \quad (26)$$

3. Change in Photon Energy

Let's determine how the energy of a photon changes in the considered space-time applying the principle of the extremum of the photon's energy integral [17]-[21]. The Euler-Lagrange equations for the covariant components of the energy-momentum vector of a light-like particle of unit energy

$$\frac{dp_\lambda}{d\mu} - F_\lambda = 0 \quad (27)$$

with

$$F_\lambda = \frac{1}{2u_1 u^1} \frac{\partial g_{ij}}{\partial x^\lambda} u^i u^j, \quad (28)$$

corresponding to the energy, take the form

$$\frac{dp_1}{d\mu} = -\frac{1}{u_{1ph} u_{ph}^1} \left[\dot{f} r u_{ph}^1 u_{ph}^2 + (\dot{a}a - \dot{f} f r^2) (u_{ph}^2)^2 + \dot{a}a \Omega^2 \right] \quad (29)$$

under the designation

$$\Omega^2 = r^2 \left[(u_{ph}^3)^2 + \sin^2 \theta (u_{ph}^4)^2 \right]. \quad (30)$$

From the equality of interval (3) to zero, we obtain

$$u_{ph}^2 = \frac{-f r u_{ph}^1 + \sigma \sqrt{a^2 (u_{ph}^1)^2 - (a^2 - f^2 r^2) \Omega^2}}{a^2 - f^2 r^2}, \quad (31)$$

where σ takes on values of ± 1 depending on the direction of the light. For small deviations from radial motion:

$$\Omega^2 \ll (u_{ph}^1)^2 \quad (32)$$

we find

$$u_{ph}^2 = \frac{1}{f r + \sigma a} u_{ph}^1 - \frac{\sigma \Omega^2}{2a u_{ph}^1}. \quad (33)$$

The expression for the time component of the covariant velocity vector is as follows:

$$u_{1ph} = u_{ph}^1 - f r u_{ph}^2 = \frac{a}{a + \sigma f r} u_{ph}^1 + \frac{\sigma f r \Omega^2}{2a u_{ph}^1}. \quad (34)$$

Further we will consider $f r$, \dot{a}/a , \dot{f}/f to be small quantities. After substituting the components u_{ph}^2 and u_{1ph} into equation (29) without accounting for higher order small quantities, it is rewritten in the form

$$\frac{dp_1}{d\mu} = -\frac{\sigma \dot{a} + \dot{f} r}{f r + \sigma a}. \quad (35)$$

Since the covariant and contravariant temporal components of the generalized momenta of a light-like particle

$$p_\lambda = \frac{u_\lambda}{u^1 u_1}, \quad (36)$$

$$p^\lambda = \frac{u^\lambda}{u^1 u_1} \quad (37)$$

are related by

$$p^1 = \frac{u_{ph}^1}{u_{1ph}} p_1 = \left(1 + \sigma \frac{f r}{a} \right) p_1, \quad (38)$$

the rate of energy change can be written in the form

$$\frac{dp^1}{d\mu} = \left(1 + \sigma \frac{fr}{a}\right) \frac{dp_1}{d\mu} + \sigma \frac{1}{a^2 u_{ph}^1} \left[r(fa - af)u_{ph}^1 + fau_{ph}^2 \right]. \quad (39)$$

Substituting u_{ph}^2 from equation (33) and $dp_1/d\mu$ from equation (35) into this expression, we find

$$\frac{dp^1}{d\mu} = -\frac{\dot{f}r + \sigma\dot{a}}{\sigma a} + \frac{1}{\sigma a^2} \left[r(fa - af) + \frac{fa}{fr + \sigma a} - \frac{\sigma f \Omega^2}{2(u_{ph}^1)^2} \right]. \quad (40)$$

Assuming that the terms in this equation containing r are small in the region under consideration, we can write

$$\frac{dp^1}{d\mu} = -\frac{\dot{a}}{a} + \frac{f}{\sigma a^2} - \frac{f \Omega^2}{2a^2 (u_{ph}^1)^2}. \quad (41)$$

In the case of the motion of light emitted towards the observer, we choose $\sigma = -1$. Then, upon substitution of (22) and (25), this expression is transformed to become

$$\begin{aligned} \frac{dp^1}{d\mu} = & -A_1 \left(\frac{3}{2} - \frac{1}{2} e^{-3cA_1(t-t_0)} \right) - f_0 e^{-3cA_1(t-t_0)} \\ & - \frac{\Omega^2}{2(u_{ph}^1)^2} \left[\frac{1}{2} A_1 (1 - e^{-3cA_1(t-t_0)}) + f_0 e^{-3cA_1(t-t_0)} \right]. \end{aligned} \quad (42)$$

In the region under consideration, the energy of a radially moving photon is given by

$$E_{ph} = h\nu_0 \left(1 - r \frac{dp^1}{d\mu} \Big|_{t=t_0} \right), \quad (43)$$

where ν_0 is the value of the photon frequency at the observation point. Since in the cosmological model described by an orthogonal metric, the Hubble constant is expressed as

$$\check{H} = c \frac{\dot{a}(t_0)}{a(t_0)}, \quad (44)$$

from equation (26), when comparing it with the Friedman equation

$$\chi c^2 \rho = \frac{3}{a^2} [k + (\dot{a})^2] - \Lambda, \quad (45)$$

it follows that A_1 can be a quantity of the same order as the Hubble parameter. After substituting the function f (25) without small quantities, equation (42) takes the form

$$\frac{dp^1}{d\mu} \Big|_{t=t_0} = -A_1 - f_0 - \frac{f_0 \Omega^2}{2(u_{ph}^1)^2}. \quad (46)$$

Since in the region under consideration the spacetime metric (3) approximates the Minkowski metric, the affine parameter will be close to $\mu = ct$.

4. Change in the Energy of a Material Particle in the Co-Moving Frame

The Lagrangian of a material particle [17] in space-time (3) is given by

$$L = cm \left\{ (u^1)^2 - 2f(t)ru^1u^2 - \right.$$

$$[a(t)^2 - f(t)^2 r^2](u^2)^2 - a(t)^2 r^2 ((u^3)^2 + \sin^2 \theta (u^4)^2) \}^{1/2}. \quad (47)$$

The motion of a material particle in a gravitational field, like that of a photon, in accordance with equation (27) at $\mu = s$ can be represented [17,21] as the result of the action of generalized forces

$$F_\lambda = \frac{\partial L}{\partial x^\lambda} = \frac{1}{2} cm \frac{\partial g_{ij}}{\partial x^\lambda} u^i u^j, \quad (48)$$

which, however, do not generally form a first-order tensor. The components of the vector of generalized forces associated with F_λ ,

$$F^k = \frac{1}{2} cm g^{k\lambda} \frac{\partial g_{ij}}{\partial x^\lambda} u^i u^j \quad (49)$$

are related to gravitational forces $F^k = cF^k$. The equations of motion (27) for the contravariant momentum are transformed into the form

$$\frac{d\bar{p}^k}{d\mu} = \frac{dp^k}{d\mu} + \frac{d\bar{p}^{\leftrightarrow k}}{d\mu} = F^k, \quad (50)$$

where

$$\frac{d\bar{p}^{\leftrightarrow k}}{d\mu} = g^{k\lambda} \frac{\partial g_{\lambda i}}{\partial x^j} u^j p^i \quad (51)$$

represents the energy imparted to the gravitational field.

Let's consider a particle of matter in a comoving frame, in which it is motionless (24). The component of the force vector (49) corresponding to energy, F_m^1 , becomes zero. The rate of change of energy transferred to the gravitational field (51) will be

$$\frac{dp_m^1}{ds} = cm \frac{f \dot{f} r^2}{a^2}. \quad (52)$$

Due to (50), the energy of the material particle changes at a rate of

$$\frac{dp_m^1}{ds} = -cm \frac{f \dot{f} r^2}{a^2}. \quad (53)$$

The derivative of the function $f(t)$ has the form

$$\dot{f} = -A_1 f_0 e^{-cA_1(t-t_0)} + A_1^2 \left(e^{2cA_1(t-t_0)} + \frac{1}{2} e^{-cA_1(t-t_0)} \right). \quad (54)$$

Now we can express the change in energy of the material particle at the present time as

$$\left. \frac{dp_m^1}{ds} \right|_{t=t_0} = cm A_1 \left(f_0 + \frac{3}{2} A_1 \right) f_0 r^2. \quad (55)$$

In the considered region, the energy of the material particle will be

$$E_m = c \left(cm - r \left. \frac{dp_m^1}{ds} \right|_{t=t_0} \right). \quad (56)$$

5. Cosmological Redshift in the Radial Motion of Photons

The equation (26) is rewritten in the form

$$3(A_1 + f_0)^2 - 12f_0 A_1 - 3f_0^2 = \chi c^2 \rho_0. \quad (57)$$

We assume that f_0 is of the same order of magnitude as A_1 and, consequently, with the Hubble constant. Since in the considered region the quantity fr is small, taking into account the expressions for the rate of change of particle energies (46) and (55), the change in the energy of a material particle will be small in comparison with the change in the equivalent energy of a photon (43) because of

$$\left(\frac{dp_m^1}{ds} \Big|_{t=t_0} \right) / \left(h\nu_0 \frac{dp^1}{d\mu} \Big|_{t=t_0} \right) = o(fr), \quad (58)$$

where the differentiation parameters are

$$s \approx \mu \approx ct. \quad (59)$$

The observed redshift of photon energy is considered as a result of the change in the ratio of their energy (43), (46) to the energy of atoms (56). In a small neighborhood without small higher-order quantities due to (58), this will be

$$\frac{E_{ph}}{E_m} = \frac{h\nu_0}{c^2 m} [1 - r(A_1 + f_0)], \quad (60)$$

from which the value of the Hubble constant at present time

$$H_0 = c(A_1 + f_0), \quad (61)$$

can be deduced, obtained as a result of registering radially moving photons.

In this case, equation (57) takes the form

$$3\left(\frac{H_0}{c}\right)^2 - 12\frac{H_0}{c}f_0 + 9f_0^2 = \chi c^2 \rho_0. \quad (62)$$

6. Observed Length Element

The length element

$$dl^2 = \left(\frac{g_{1p}g_{1q}}{g_{11}} - g_{pq} \right) dx^p dx^q \quad (63)$$

with $p, q = 1, 2, 3$ in the space-time (3) is given by

$$dl = a(t) \sqrt{dr^2 + d\theta^2 + \sin^2 \theta d\varphi^2}. \quad (64)$$

That is, the space is isotropic in proper frame.

The radial elements of the length of the metric (3) at constant time

$$dl' = \sqrt{a(t)^2 - f(t)^2} r^2 dr \quad (65)$$

is less than obtained from (64) with $d\theta = d\varphi = 0$ and approach it with decreasing r . The observed space in the coordinate frame appears to be compressed in the radial direction. This will cause distant objects to appear smaller in size than in isotropic space. A similar effect was discovered when observing galaxies with large z [4]. However, for precise adjustment of the considered model parameters for such distances, a solution is required without the assumption about the insignificance of some terms in the tensor G^{ij} .

7. Difference in Hubble Constant Measurements

Let's consider how the energy of a photon will change in the presence of gravitational lensing. We denote the change in the energy of the photon when moving along the path L_l , which includes gravitational lensing, as

$$\delta p_l^1 = \int_{L_l} \frac{dp^1}{d\mu} d\mu \quad (66)$$

and along the radial path as

$$\delta p_r^1 = \int_{L_r} \frac{dp^1}{d\mu} d\mu. \quad (67)$$

We assume that the coordinate distance Δr from the beginning of the paths to the observer is the same. Under the gravitational lensing, neglecting small quantities, the Hubble constant is determined as follows:

$$H_0 = -\frac{c(\delta p_l^1 - \delta p_r^1)}{\delta\mu_l - \delta\mu_r}, \quad (68)$$

where $\delta\mu_l$ and $\delta\mu_r$ are the change in the affine parameter along each of the paths.

The coordinate system is chosen such that when the photon moves along the path L_l the condition $\theta = \frac{\pi}{2}$ is satisfied. If we take into account only the influence of the deviation from radial motion, then the difference in the time taken for the photon to travel along both paths will be

$$\delta t = \frac{1}{c} \int_0^{\Delta r} (1 - \cos(\alpha r)) dr, \quad (69)$$

where in the first approximation we consider

$$\cos(\alpha(r)) = -\frac{u_{ph}^2}{\sqrt{(u_{ph}^2)^2 + (ru_{ph}^4)^2}}. \quad (70)$$

Due to condition (32), the integral (69) transforms into the form

$$\delta t = \frac{1}{c} \int_0^{\Delta r} \frac{(ru_{ph}^4)^2}{2(u_{ph}^2)^2} dr. \quad (71)$$

In the considered region, we have

$$\delta\mu_l - \delta\mu_r = c\delta t. \quad (72)$$

By substituting the corresponding value from (46) into (66) and (67), we find the additional difference in energy change caused by the discrepancy in the rate of its during radial and angular motion

$$\Delta(\delta p_l^1 - \delta p_r^1) = - \int_0^{\Delta r} \frac{f_0(ru_{ph}^4)^2}{2(u_{ph}^1)^2} dr. \quad (73)$$

The change in the value of the Hubble constant when light moves along a curved path will be

$$\Delta H_0 = -\frac{c\Delta(\delta p_l^1 - \delta p_r^1)}{\delta\mu_l - \delta\mu_r}. \quad (74)$$

Since for the considered motion of the photon, the relations

$$u_{ph}^1 \approx -au_{ph}^2 \approx -u_{ph}^2 \quad (75)$$

hold, by substituting the values (72) and (73) into (74), we obtain

$$\Delta H_0 = cf_0. \quad (76)$$

Let's estimate the values of A_1 and f_0 using the results of obtaining the Hubble constant with and without gravitational lensing. The Hubble constant was found through gravitational lensing of the cosmic microwave background: $68.3 \pm 1.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$, Ref. [15], and of radiation from the supernovae: $66.6_{-3.3}^{+4.1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, Ref. [16]. The values obtained through the extragalactic distance ladder to supernovae are $73.5 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$, Ref. [13], and $75.4_{-3.7}^{+3.8} \text{ km s}^{-1} \text{ Mpc}^{-1}$, Ref. [14]. These results yield the value $\Delta H_0 = cf_0 = -7_{-2.6}^{+2.4} \text{ km s}^{-1} \text{ Mpc}^{-1}$, and due to relation (61) corresponding to the radial motion of the photon we obtain $cA_1 = 81.5 \pm 2.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

When the value of f_0 is negative, the photon moving towards observer from the star additionally loses energy compared to the motion tangential to the circle from the center of observation or in the opposite direction. Also, the negativity of the function f means that space is expanding faster than matter is scattering. The dependence of the spacetime properties on the observer's position is similar to the manifestation of the spacetime curvature in the non-flat FLRW model.

8. Conclusions

Unlike the Λ CDM cosmology, in the presented model the observed space expanded not synchronously with movement of matter. In region corresponding to small perturbations of the FLRW metric the Universe expands with zero pressure while conserving energy and momentum. Solution is obtained that relates exponential proper expansion of space for a model with a zero cosmological constant. The considered metric can also be applied to a cosmological model with a nonzero Λ -term.

In addition to the space expansion, the movement of matter in the radial direction is also established in coordinate frame. The observed space in this frame appears to be compressed in the radial direction. This will cause distant objects to appear smaller in size than in isotropic space.

The redshift in the spectra of galaxies is determined using Lagrange mechanics in its application to the principle of the extremal integral of photon energy. In the considered spacetime, from the observer's standpoint, a photon loses energy faster when moving radially towards them compared to its motion with a tangential component. The light used to obtain redshift through the ladder distance travels along a radial path, unlike gravitational lensing, where its pass has a tangential component. Based on the difference in value of the Hubble constant obtained using gravitational lensing and distance ladder the metric coefficients were found in a local region in which their perturbations are small.

References

1. B. E. Robertson et al., *Nat Astron* **7** (2023) 611.
2. R. P. Naidu et al., *ApJL* **940** (2022) L14.
3. K. N. Hainline et al., *ApJ* **964** (2024) 71.
4. R. P. Gupta, *ApJ* **964** (2024) 55.
5. V. A. Rubakov and P. G. Tinyakov, *Phys. Usp.* **51**, (2008) 759.
6. T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, *Phys. Rep.* **513** (2012) 1.
7. L. A. Glinka, *Journal of Applied Mathematics and Physics* **2**(3) (2014) 112.
8. A. Finke, *MNRAS* **491** (2020) 2330.
9. R. Gaur and M. Visser, *JCAP09 2022* (2022) 030.
10. R. Monjo, *Phys. Rev. D* **96** (2017) 103505.
11. R. Monjo, *Class. Quantum Grav.* **40** (2023) 235002.
12. R. Monjo and R. Campoamor-Stursberg, *Class. Quantum Grav.* **40** (2023) 195006.
13. D Brout. et al., *ApJ* **938** (2022) 110.
14. T. De Jaeger et al., *MNRAS* **514** (2022) 4620.
15. L. Balkenhol et al., *Phys. Rev. D* **108** (2023) 023510.
16. P. L. Kelly et al., *Science* **380** (2023) 6649.

17. W. B. Belayev, *Dinamika v obshhej teorii otноситel'nosti: variacionnye metody* (URSS, Moscow, 2017).
18. D. Yu. Tsipenyuk and W. B. Belayev, *J. Phys.: Conf. Ser.* **251** (2019) 012048.
19. D. Yu. Tsipenyuk and W. B. Belayev, *Rom. Rep. Phys.* **71** (2019) 109.
20. D. Yu. Tsipenyuk and W. B. Belayev, *RENSIT* **11** (2019) 249.
21. D. Yu. Tsipenyuk and W. B. Belayev, *Gravitational waves, fields, and particles in the frame of (1 + 4)D Extended space model*, in *Gravitational Waves - Theory and Observations*, ed. C. Frajuca, art. 2 (IntechOpen, London, 2023).
22. The components of the Einstein tensor were obtained using the Maxima software.
23. L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, 4th edn. (Butterworth-Heinemann, Oxford, 2000).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.