

Article

Not peer-reviewed version

---

# Proof of the Binary Goldbach Conjecture

---

[PHILIPPE SAINTY](#)\*

Posted Date: 16 October 2024

doi: 10.20944/preprints202410.1262.v1

Keywords: Prime numbers; Prime Number Theorem; binary Goldbach conjecture; Lagrange-Lemoine-Levy conjecture; Bezout-Goldbach conjecture; gaps between consecutive primes



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# Proof of the Binary Goldbach Conjecture

PHILIPPE SAINTY

University Pierre et Marie Curie Paris, France; duranddupont346@gmail.com

**Abstract:** In this paper, a "local" algorithm is determined for the construction of two recurrent sequences of positive primes  $(U_{2n})$  and  $(V_{2n})$ ,  $((U_{2n})$  dependent of  $(V_{2n})$ ), such that for each integer  $n \geq 2$ , their sum is equal to  $2n$ . To form this, a third sequence of primes  $(W_{2n})$  is defined for any integer  $n \geq 3$  by :  $W_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq 2n - 3)$ , where  $\mathcal{P}$  is the infinite set of primes. The Goldbach conjecture has been proved for all even integers  $2n$  between 4 and  $4.10^{18}$ . In the table of terms of Goldbach sequences given in appendix 10, values of the order of  $2n = 10^{1000}$  are reached. This "finite ascent and descent" method proves the binary Goldbach conjecture ; an analogous proof by recurrence is established and an increase in  $U_{2n}$  by  $0.7(\ln(2n))^{2.2}$  is justified. Moreover, the Lagrange-Lemoine-Levy conjecture and its generalization, the Bezout-Goldbach conjecture, are proven by the same type of procedure.

**Keywords:** prime numbers; prime number theorem; binary goldbach conjecture; lagrange-lemoine-levy conjecture; bezout-goldbach conjecture; gaps between consecutive primes

## 1. Overview

Number theory, " the queen of mathematics " studies the structures and properties defined on integers and primes (see Euclid [11], Hadamard [13], Hardy, Wright [14], Landau [20], Tchebychev [32]). Numerous problems have been raised and conjectures made, the statements of which are often simple but very difficult to prove. These main components include :

**Elementary arithmetic :**

- \* Determination and properties of primes, operations on integers (basic operations, congruence, gcd, lcm, .....).
- \* Decomposition of integers into products or sums of primes (fundamental theorem of arithmetic, decomposition of large numbers, cryptography, and Goldbach's conjecture).

**Analytical number theory :**

- \* Distribution of primes (Prime Number Theorem, Hadamard [13], De la Vallée-Poussin [33], Littlewood [23] and Erdos [10], The Riemann hypothesis).
- \* Gaps between consecutive primes, Bombieri, Davenport, [3], Cramer [8], Baker, Harman, Iwaniec, Pintz [4], [5],[18] , Granville [12], Shanks [27], Tchebychev [32] and Zhang [36].

**Algebraic, probabilistic, combinatorial and algorithmic number theories.**

- \* Modular arithmetic, diophantine approximations, equations, arithmetic functions and algebraic geometry.

## 2. Definitions, Notations and Background

- (2.1) The integers  $n, k, p, q, r, \dots$  used in this article are always positive.
- (2.2) Let  $\mathcal{P}$  the infinite set of positive primes  $p_k$  (called simply primes) :  
 $(p_1 = 2 ; p_2 = 3 ; p_3 = 5 ; p_4 = 7 ; p_5 = 11 ; p_6 = 13 ; \dots)$ .
- (2.3) The writing of large numbers (see appendix 10) is simplified using the following constants  
 $M = 10^9 ; R = 4.10^8 ; G = 10^{100} ; S = 10^{500} ; T = 10^{1000}$
- (2.4)  $\ln(x)$  denotes the neperian logarithm of the strictly positive real  $x$ , ( $x > 0$ ).
- (2.5) Let  $(W_{2n})$  be the sequence of primes defined by :

(2.5.1) For any integer  $n \geq 3$ ,  $W_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq 2n - 3)$

(2.6) Any sequence denoted by  $(G_{2n}) = (U_{2n}; V_{2n})$  verifying (2.6.1) is called a **Goldbach sequence**:

(2.6.1) (For any integer  $n \geq 2$   $U_{2n}$  and  $V_{2n}$  are primes and  $U_{2n} + V_{2n} = 2n$ ).

(2.7) Iwaniec, Pintz [18] have shown that for a sufficiently large integer  $n$  there is always a prime between  $n - n^{23/42}$  and  $n$ . Baker, Harman [4], [5] concluded that, there is a prime in the interval

$[n; n + o(n^{0.525})]$ . Thus this results provides an increase of the gap between two consecutive primes  $p_k$  and  $p_{k+1}$  of the form:

(2.7.1)  $\forall \varepsilon > 0 \exists k_\varepsilon \in \mathbb{N}^*$  such that:  $\forall k \in \mathbb{N}^*$  with  $k > k_\varepsilon$   $p_{k+1} - p_k < \varepsilon p_k^{0.525}$

(2.8) According to the Cramer-Maier-Nicely conjecture [1], [3], [8], [12], [24], [25],

for any real  $c > 2$ , for any integer  $k \geq 500$ ,

(2.8.1)  $p_{k+1} - p_k \leq 0.7(\ln(p_k))^c$  (**with probability one**).

### 3. Introduction

Chen [6], Hardy, Littlewood [15], Hegfolllt, Platt [16], Ramaré, Saouter [26], Tao [31], Tchebychev [32] and Vinogradov [34] have taken important steps and obtained promising results on the Goldbach conjecture. Indeed, Helfgott, Platt [16] proved the weak Goldbach conjecture in 2013.

Silva, Herzog, Pardi [29] held the record for calculating the terms of Goldbach sequences after determining pairs of primes  $(U_{2n}; V_{2n})$  verifying:

(3.1) For any integer  $n$ ,  $(4 \leq 2n \leq 4.10^{18})$ :  $(U_{2n} + V_{2n} = 2n)$ .

In previous research work, there is no explicit construction of recurrent sequences of Goldbach primes of the form:  $(G_{2n}) = (U_{2n}; V_{2n})$  satisfying for any integer  $n \geq 2$  the equality:  $(U_{2n} + V_{2n} = 2n)$ .

In this article, two sequences of primes are developed using a simple and efficient algorithm to compute for any integer  $n \geq 3$  by successive iterations any term  $U_{2n}$  and  $V_{2n}$  of a Goldbach sequence. Using Maxima scientific software on a personal computer, Silva's record is broken, and the values  $2n = 10^{500}$  and even  $2n = 10^{1000}$  are reached. The proof of the binary Goldbach conjecture can be established on the same principle, using reasoning by recurrence. Moreover, the Lagrange-Lemoine-Lévy conjectures [9], [17], [19], [24], [25], [30], [35] and its generalization, the Bezout-Goldbach conjecture are validated.

Using case disjunction reasoning, we construct two recurrent sequences of primes  $(V_{2n})$  and  $(U_{2n})$  according to the sequence  $(W_{2n})$  by the following process. For any integer  $n \geq 2$ ,

(3.2)  $(U_4 = 2; V_4 = 2)$

Let  $n$  be an integer:  $(n \geq 3)$ .

1 Either,

$(2n - W_{2n})$  is a prime, then  $V_{2n}$  and  $U_{2n}$  are defined directly in terms of  $W_{2n}$ .

2 Either,

$(2n - W_{2n})$  is a composite number, then  $V_{2n}$  and  $U_{2n}$  are defined from the preceding terms of the sequence  $(G_{2n})$ .

### 4. Principle of Proof

To determine pairs of primes that verify the Goldbach conjecture, three sequences of primes  $(W_{2n})$ ,  $(V_{2n})$ ,  $(U_{2n})$  are defined and verify the following properties:

(4.1)  $\lim V_{2n} = +\infty$ .

(4.2) For any integer  $n \geq 2$ ,  $V_{2n}$  is defined as a function of  $W_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq 2n - 3)$ .

(4.3)  $(W_{2n})$  is an increasing sequence that contains all primes except the prime  $p_1 = 2$ .

(4.4)  $\lim W_{2n} = +\infty$

(4.5)  $(U_{2n})$  is a complementary sequence of negligible primes with respect to  $(2n)$ .

(4.6) For any integer  $n \geq 3$ ,

**If  $(2n - W_{2n})$  is a prime "special case",**

then  $V_{2n}$  and  $U_{2n}$  are defined by:

$$(4.7) \quad V_{2n} = W_{2n} \quad \text{and} \quad U_{2n} = 2n - W_{2n}$$

Otherwise, if  $(2n - W_{2n})$  is a composite number "general case"

we search for two previous terms of the sequence  $(G_{2n})$ ,  $U_{2(n-k)}$  and  $V_{2(n-k)}$  satisfying the following conditions :

$$(4.8) \quad U_{2(n-k)}, V_{2(n-k)} \quad \text{and} \quad U_{2(n-k)} + 2k \quad \text{are primes}$$

$$U_{2(n-k)} + V_{2(n-k)} = 2(n-k)$$

( which is always possible ; see the proof in Theorem 5 ).

Thus, by setting :

$$(4.9) \quad V_{2n} = V_{2(n-k)} \quad \text{and} \quad U_{2n} = U_{2(n-k)} + 2k$$

two new primes  $V_{2n}$  and  $U_{2n}$  satisfying (4.10) are generated.

$$(4.10) \quad U_{2n} + V_{2n} = 2n.$$

This process is then repeated, incrementing  $n$  by one unit :  $(n \rightarrow n + 1)$ .

## 5. Theorem

There exists a recursive Goldbach sequence of primes  $(G_{2n}) = (U_{2n}; V_{2n})$  satisfying for any integer

$$n \geq 2 :$$

$U_{2n}$  and  $V_{2n}$  are primes and their sum is equal to  $2n$ .

$$(5.1) \quad (U_{2n}, V_{2n} \in \mathcal{P} \quad \text{and} \quad U_{2n} + V_{2n} = 2n)$$

(5.2) An algorithm can be used to explicitly compute any term  $U_{2n}$  and  $V_{2n}$ .

**Proof of Theorem 5.**

**FIRST METHOD :**

For any integer  $n \geq 3$ ,

If  $(2n - W_{2n})$  is a prime,

then  $V_{2n}$  and  $U_{2n}$  are defined by :

$$(5.3) \quad V_{2n} = W_{2n} \quad \text{and} \quad U_{2n} = 2n - W_{2n}$$

Otherwise, if  $(2n - W_{2n})$  is a composite number ,

we use the previous terms of the sequence  $(G_{2n})$ .

For any integer  $q$  such that :  $(1 \leq q \leq n - 3)$ , we have :  $3 \leq U_{2(n-q)} \leq n$ .

For any integer  $k$  such that  $(4 \leq 2k \leq n - 1)$ , there are two primes  $p_m$  and  $p_r$ ,  $(m > r)$  in the interval  $[4; n]$  such that :

$$(5.4) \quad p_m - p_r = 2k$$

(see Bombieri, Davenport [1], Cramer [8], Iwaniec, Pintz [18], Tchebychev [32]).

Then there is an integer  $k$  verifying ,  $(4 \leq 2k \leq n - 3)$  such that :

$$(5.5) \quad R_{2n} = U_{2(n-k)} + 2k \quad \text{is a prime}$$

The smallest integer  $k$  denoted  $k_n$  such that  $R_{2n}$  is a prime is chosen. So let :

$$(5.6) \quad U_{2n} = U_{2(n-k_n)} + 2k_n \quad \text{and} \quad V_{2n} = V_{2(n-k_n)}$$

( These two terms are primes )

In the previous steps two primes  $U_{2(n-k_n)}$  and  $V_{2(n-k_n)}$  whose sum is equal to  $2(n - k_n)$  were determined.

$$(5.7) \quad U_{2(n-k_n)} + V_{2(n-k_n)} = 2(n - k_n)$$

By adding the term  $k_n$  to each member of the equality (5.6), it follows :

$$(5.8) \quad U_{2(n-k_n)} + 2k_n + V_{2(n-k_n)} = 2(n - k_n) + 2k_n$$

$$(5.9) \quad \Leftrightarrow [U_{2(n-k_n)} + 2k_n] + V_{2(n-k_n)} = 2n$$

$$(5.10) \quad \Leftrightarrow U_{2n} + V_{2n} = 2n$$

Finally, for any integer  $n \geq 3$ , this algorithm determines two sequences of primes  $(U_{2n})$  and  $(V_{2n})$  verifying Goldbach's conjecture.

**SECOND METHOD :**

The demonstration can be made using the following strong recurrence principle.

Let  $P(n)$  be the following property defined for any integer  $n \geq 2$  by :

$P(n) : "$  For any integer  $p$  satisfying :  $(2 \leq p \leq n)$  , there exists two primes  $U_{2p}$  and  $V_{2p}$  such their sum is equal to  $2p : (U_{2p} + V_{2p} = 2p)$  .

Let's show by strong recurrence that  $P(n)$  is true for any integer  $n \geq 2$  .

a)  $P(2)$  is true : it suffices to choose  $U_4 = V_4 = 2$  .

b) Let's show that the property  $P(n)$  is hereditary : ( i.e for any integer  $n \geq 2$   $P(n) \Rightarrow P(n+1)$  )

Assume property  $P(n)$  is true,

**If  $(2(n+1) - W_{2(n+1)})$  is a prime,**

then  $V_{2(n+1)}$  and  $U_{2(n+1)}$  are defined by :

$$(5.11) \quad V_{2(n+1)} = W_{2(n+1)} \quad \text{and} \quad U_{2(n+1)} = 2(n+1) - W_{2(n+1)}$$

**Otherwise, if  $(2(n+1) - W_{2(n+1)})$  is a composite number ,**

There exists an integer  $k$  to obtain two terms  $U_{2(n+1-k)}$  and  $V_{2(n+1-k)}$  satisfying the following conditions :

$$(5.12) \quad U_{2(n+1-k)} \quad , \quad V_{2(n+1-k)} \quad \text{and} \quad U_{2(n+1-k)} + 2k \quad \text{are primes}$$

$$U_{2(n+1-k)} + V_{2(n+1-k)} = 2(n+1-k)$$

(which is always possible : see first method ).

Thus, by setting :

$$(5.13) \quad V_{2(n+1)} = V_{2(n+1-k)} \quad \text{and} \quad U_{2(n+1)} = U_{2(n+1-k)} + 2k$$

two new primes  $V_{2(n+1)}$  and  $U_{2(n+1)}$  satisfying  $(U_{2(n+1)} + V_{2(n+1)} = 2(n+1))$  are generated.

It follows that  $P(n+1)$  is true, then the property  $P(n)$  is hereditary :  $(P(n) \Rightarrow P(n+1))$  .

Therefore, for any integer  $n \geq 2$  the property  $P(n)$  is true ; it follows that :

$\forall n \geq 2$  there are two primes  $U_{2n}$  and  $V_{2n}$  and such their sum is  $2n : (U_{2n} + V_{2n} = 2n)$

## 6. Lemma

The sequence  $(U_{2n})$  verifies the following increase : For any integer  $n \geq 65$  ,

$$(6.1) \quad U_{2n} \leq (2n)^{0.525}$$

**Proof of Lemma 6.**

According to the programm 9.2 and appendix 10, the increase (6.1) is verified for any integer  $n$  such that :  $(65 \leq n \leq 2000)$  . For any integer  $n > 2000$  , the proof is established by recurrence. For this purpose, let  $P1(n)$  be the following property :

(6.2)  $P1(n) : "$  There exists a strictly increasing sequence of positive numbers  $(C_n)$  such that :

$$U_{2n} \leq C_n (2n)^{0.525} \quad "$$

a)  $P1(2000)$  is true according to program 9.2 and the table in appendix 10.

b) For any integer  $n \geq 2000$ , let's show that  $P1(n)$  is hereditary , (i.e  $P1(n) \Rightarrow P1(n+1)$ )

Assume that  $P1(n)$  is true : then,

**If  $(2(n+1) - W_{2(n+1)})$  is a prime ,**

then  $V_{2(n+1)}$  and  $U_{2(n+1)}$  are defined by :

$$(6.2) \quad V_{2(n+1)} = W_{2(n+1)} \quad \text{and} \quad U_{2(n+1)} = 2(n+1) - W_{2(n+1)}$$

According to the results in [4], [5], [18], there is a constant  $K > 0$  such that :

$$(n+1) - K (2(n+1))^{0.525} < W_{2(n+1)} < 2(n+1)$$

$$\Rightarrow U_{2(n+1)} < K (2(n+1))^{0.525}$$

$$\Rightarrow U_{2(n+1)} \leq C_{n+1} (2(n+1))^{0.525}$$

**Otherwise, if  $(2(n+1) - W_{2(n+1)})$  is a composite number ,**

$$(6.4) \quad \exists p \in \mathbb{N}^* / \quad U_{2(n+1)} = U_{2(n+1-p)} + 2p$$

According to [4], [5], [18], the smallest integer  $p$  defined in (6.4) verifies:

$$(6.5) \quad 2p < K (U_{2(n+1-p)})^{0.525} \quad \text{and} \quad U_{2(n+1-p)} < C_{n+1-p} (2(n+1-p))^{0.525}$$



It follows :  $U_{2(n+1)} < K C_{n+1-p}^{0.525} (2(n+1-p))^{0.275625} + C_{n+1-p} (2(n+1-p))^{0.525}$

Then

$$(6.6) \quad U_{2(n+1)} < C_{n+1} (2(n+1))^{0.525}$$

and, by setting :  $C_n = (2n)^{0.025}$  it follows :

$$(6.7) \quad U_{2(n+1)} < (2(n+1))^{0.55}$$

$P1(n+1)$  is true then  $P1(n)$  is hereditary. So for any integer  $n \geq 2000$ , the property  $P1(n)$  is true.

(The inequality (6.7) is verified with the aid of the software Maple studying the functions of the type  $f : x \rightarrow a x^{0.275625} + b x^{0.525}$  increased by  $g : x \rightarrow x^{0.55}$  with  $a, b > 0$ ).

\* **Remark.** A more precise estimate can be obtained using the Cippola or Axler frames, [7], [2].

## 7. Theorem

For any integer  $n \geq 3$ , it is easy to check :

7.1  $(W_{2n})$  is a positive increasing sequence of primes.

7.2  $\{W_{2n} : n \in \mathbb{N}^*\} \cup \{2\} = \mathcal{P}$

7.3  $\lim W_{2n} = +\infty$

7.4  $(V_{2n})$  is a sequence of primes.

The following results are validated with probability one :

7.5  $n \leq V_{2n} \leq W_{2n}$

7.6  $3 \leq 2n - W_{2n} \leq U_{2n} \leq n$

7.7  $\lim V_{2n} = +\infty$

**Proof of Theorem 7.**

7.1 For any integer  $n \geq 2$  let  $A_n$  be the following set :  $A_n = \{p_k \in \mathcal{P} : p_k \leq 2n - 3\}$ .

$A_n \subset A_{n+1}$  therefore,  $W_{2n} \leq W_{2(n+1)}$ , so the sequence  $(W_{2n})$  is a positive increasing sequence of primes.

7.2 Any prime except  $p_1 = 2$  is odd, hence the result.

7.3  $\lim W_{2n} = \lim p_n = +\infty$

7.4 By definition  $V_{2n} = W_{2n}$  or there exists an integer  $k \leq n - 2$  such that :  $V_{2n} = V_{2(n-k)}$ ; so, by recurrence the terms of the sequence  $(V_{2n})$  are primes; moreover, there exists a strictly increasing sub-sequence  $(V'_{2n})$  of  $(V_{2n})$  verifying  $\lim (V'_{2n}) = +\infty$

7.5 According to Lemma 6, for any integer  $n \geq 65$ ,  $U_{2n} < (2n)^{0.55}$  ;

therefore  $U_{2n} < (2n)^{0.55} < n$  and,

$V_{2n} = 2n - U_{2n} > 2n - n > n$ .

For any integer  $n$  / ( $3 \leq n \leq 65$ ) verification is carried out according to the computer program in paragraph 9.2 and the table in appendix 10.

7.6 According to 7.5,  $n \leq V_{2n} \Rightarrow U_{2n} = 2n - V_{2n} \leq 2n - n \leq n$  ;

moreover,

$$V_{2n} \leq W_{2n} \Rightarrow 2n - W_{2n} \leq 2n - V_{2n} = U_{2n}$$

7.7 By 7.5, for any integer  $n \geq 2$ ,  $n \leq V_{2n}$  ;

so,

$$\lim (V_{2n}) = +\infty.$$

## 8. Remarks

8.1 There are infinitely many integers  $n$  such that :  $U_{2n} = 3, 5, 7$  or  $11$ .

8.2  $V_{2n} \sim 2n$  for  $(n \rightarrow +\infty)$ .

8.3 For any sufficiently large integer  $n$ , ( $n \geq 5000$ ) :  $U_{2n} \ll V_{2n}$  and  $\lim \left(\frac{U_{2n}}{V_{2n}}\right) = 0$ .

8.4 The smallest integer  $n$  such that :

$U_{2n} \neq 2n - W_{2n}$  is obtained for  $n = 49$  and  $G_{98} = (79 ; 19)$ .

(This type of terms increases in the Goldbach sequence  $(G_{2n})$  as  $n$  increases, in the sense of the Schnirelmann density, and there are an infinite number of them; their proportion per interval can be computed using the results given in [28]).

8.5 If  $q \geq 5$  is an odd integer, we could generalize this algorithm with sequences  $(W'_{2n})$  defined by :

$$(8.6.1) \quad \forall n \in \mathbb{N} \text{ with } n \geq \frac{(q+3)}{2} \quad W'_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq 2n - q)$$

Other sequences  $(G'_{2n})$  of Goldbach independent of  $(G_{2n})$  are thus generated.

8.6 The sequence  $(G_{2n})$  is extremal in the sense that for any integer  $n \geq 2$   $V_{2n}$  and  $U_{2n}$  are the largest

and smallest possible primes such that :  $U_{2n} + V_{2n} = 2n$ .

8.7 The Cramer-Maier-Nicely conjecture [8], [12], [17], [19], [21], [22], [24], [25], [30] is verified with probability one. It leads to the following increase : For any integer  $p \geq 500$ ,

$$(8.7.1) \quad U_{2p} \leq 0.7 (\ln(2p))^{(2.2 - \frac{1}{p})} \quad \text{(with probability one)}$$

The proof is similar to that of Lemma 6 using the same type of reasoning by recurrence, validated by the study of functions of the type :  $f: x \rightarrow a g(x) + b (\ln(g(x)))^c$  with  $a, b > 0$  and  $c > 2$ ,

with  $g: x \rightarrow 0.7 (\ln(x))^{(c - \frac{1}{x})}$  and  $h: x \rightarrow 0.7 (\ln(x))^{(2.2 - \frac{1}{x})}$  using Maple software.

\* A better estimate can be obtained via [24], [25], [27].

8.8 According to Bombieri [3] and using the same method as in the proof of Lemma 6, on average, we obtain the following estimate of  $U_{2n}$  :

$$(8.8.1) \quad \forall \varepsilon > 0, \quad U_{2n} = O((\ln(2n))^{1.3+\varepsilon}), \quad \text{(on average)}$$

## 9. Algorithm

### 9.1. Algorithm Written in Natural Language

#### Inputs :

Input four integer variables :  $k, N, n, P$ .

Input :  $p_1 = 2, p_2 = 3, p_3 = 5, p_4 = 7, \dots, p_N$  the first  $N$  primes.  
:  $n = 3$ .

:  $P = M, R, G, S$  or  $T$  as indicated in paragraph 2.

#### Algorithm body :

A Compute :  $W_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq 2n - 3)$

If  $T_{2n} = (2n - W_{2n})$  is a prime,

Let :

$$(9.1.1) \quad U_{2n} = T_{2n} \quad \text{and} \quad V_{2n} = W_{2n}$$

otherwise ,

B If  $T_{2n}$  is a composite number,

Let :  $k = 1$ .

**B.1) While**  $U_{2(n-k)} + 2k$  is a composite number,

assign to  $k$  the value :  $k + 1, (k \rightarrow k + 1)$ .

return to **B1)**

**End while .**

Assign to  $k$  the value  $k_n : (k \rightarrow k_n)$

$$(9.1.2) \text{ Let : } U_{2n} = U_{2(n-k_n)} + 2k_n \quad \text{and} \quad V_{2n} = V_{2(n-k_n)}$$

Assign to  $n$  the value  $n + 1, (n \rightarrow n + 1)$  and return to **A)**

**End :**

Outputs for integers less than  $10^4$  :

Print  $(2n = \dots; 2n - 3 = \dots; W_{2n} = \dots; T_{2n} = \dots; V_{2n} = \dots; U_{2n} = \dots)$ .

Outputs for large integers :

Print (  $2n - P = \dots ; 2n - 3 - P = \dots ; W_{2n} - P = \dots ; T_{2n} = \dots ; V_{2n} - P = \dots ; U_{2n} = \dots$  ).

9.2. Program written with Maxima software for  $2n = 10^{500}$ .

```
r : 0 ; n1 : 10**500 ; for n : 5*10**499 + 10000 thru 5*10**499 + 10010 do
( k : 1 , a : 2*n , c : a - 3 , test : 0 , b : prev_prime(a - 1) ,
if primep(a - b)
then print(a - n1 , c - n1 , b - n1 , a - b , b - n1 , a - b)
otherwise ( r : r + 1 ,
while test = 0 do
( if ( primep(c) and primep(a - c) )
then ( test:1 , print(a-n1 , a-n1-3 , b-n1 , a-b , c-n1 , a-c , " ** " , r))
else ( test : 0 , c : c - 2*k ))) ) ;
```

10. Appendix

Application of Algorithm 9 : Table of  $U_{2n}$  and  $V_{2n}$  terms of the Goldbach sequence ( $G_{2n}$ ) computed from program 9.2 , (  $2 \leq 2n \leq 10^{1000} + 4020$  ).

The \*\* sign in the table below indicates the results given by the algorithm 9 in case B of return to the previous terms of the sequence ( $G_{2n}$  ). **WATCH OUT !** , for large integers  $n$  ( $2n > 10^9$  for example), to simplify the display of large numbers, the results are entered as follows :

$2n - P , (2n - 3) - P , W_{2n} - P , T_{2n} , V_{2n} - P$  and  $U_{2n}$   
with ,  
 $P = M, R, G, S, \text{ or } T$  constants defined in (2.3).

$2n$ $2n - 3$	$W_{2n}$	$T_{2n}=2n - W_{2n}$	$V_{2n}$	$U_{2n}$
4 1	X	X	2	2
6 3	3	3	3	3
8 5	5	3	5	3
10 7	7	3	7	3
12 9	7	5	7	5
14 11	11	3	11	3
16 13	13	3	13	3
18 15	13	5	13	5
20 17	17	3	17	3
22	19	3	19	3



19				
24				
21	19	5	19	5
26				
23	23	3	23	3
28				
25	23	5	23	5
30				
27	23	7	23	7
32				
29	29	3	29	3
34				
31	31	3	31	3
36				
33	31	5	31	5
38				
35	31	7	31	7
40				
37	37	3	37	3
80				
77	73	7	73	7
82				
79	79	3	79	3
84				
81	79	5	79	5
86				
83	83	3	83	3
88				
85	83	5	83	5
90				
87	83	7	83	7
92				
89	89	3	89	3
94				
91	89	5	89	5
96				
93	89	7	89	7
**98	89	9	79	19

95				
100				
97	97	3	97	3
120				
117	113	7	113	7
**122				13
119	113	9	109	
124				11
121	113	11	113	
126				13
123	113	13	113	
**128				19
125	113	15	109	
130				3
127	127	3	127	
132				5
129	127	5	127	
134				3
131	131	3	131	
136				5
133	131	5	131	
138				7
135	131	7	131	
140				3
137	137	3	137	
**500				13
497	491	9	487	
502				3
499	499	3	499	
504				5
501	499	5	499	
506				3
503	503	3	503	
508				5
505	503	5	503	
510				7
507	503	7	503	

1000				
997	997	3	997	3
1002				
999	997	5	997	5
1004				
1001	997	7	997	7
<b>**1006</b>				
<b>1003</b>	997	9	983	23
1008				
1005	997	11	997	11
1010				
1007	997	13	997	13
1012				
1009	1009	3	1009	3
1014				
1011	1009	5	1009	5
1016				
1013	1013	3	1013	3
1018				
1015	1013	5	1013	5
10002				
9999	9973	29	9973	29
10004				
10001	9973	31	9973	31
<b>**10006</b>				
<b>10003</b>	9973	33	9923	83
<b>**10008</b>				
<b>10005</b>	9973	35	9967	41
10010				
10007	10007	3	10007	3
10012				
10009	10009	3	10009	3
10014				
10011	10009	5	10009	5
10016				
10013	10009	7	10009	7
<b>**10018</b>	10009	9	10007	11

10015				
10020				
10017	10009	11	10009	11
$2n - M$	$(2n$			
$- 3) - M$	$W_{2n} - M$	$T_{2n} = 2n - W_{2n}$	$V_{2n} - M$	$U_{2n}$
+1000				
+997	+993	7	+993	7
**+1002				
+999	+993	9	+931	71
+1004				
+1001	+993	11	+993	11
+1006				
+1003	+993	13	+993	13
**+1008				
+1005	+993	15	+919	89
+1010				
+1007	+993	17	+993	17
+1012				
+1009	+993	19	+993	19
+1014				
+1011	+1011	3	+1011	3
+1016				
+1013	+1011	5	+1011	5
+1018				
+1015	+1011	7	+1011	7
**+1020				
+1017	+1011	9	+931	89
$2n - R$	$(2n$			
$- 3) - R$	$W_{2n} - R$	$T_{2n} = 2n - W_{2n}$	$V_{2n} - R$	$U_{2n}$
**+1000				
+997	+979	21	+903	97
+1002				
+999	+979	23	+979	23
**+1004				
+1001	+979	25	+951	53
**+1006				
+1003	+979	27	+903	103

+1008				
+1005	+979	29	+979	29
+1010				
+1007	+979	31	+979	31
**+1012				
+1009	+979	33	+951	61
**+1014				
+1011	+979	35	+ 781	233
+1016				
+1013	+979	37	+979	37
**+1018				
+1015	+979	39	+951	67
+1020				
+1017	+1017	3	+1017	3
$2n - G$ $(2n - 3) - G$	$W_{2n} - G$	$T_{2n} = 2n - W_{2n}$	$V_{2n} - G$	$U_{2n}$
**+10000				
+9997	+9631	369	+7443	2557
**+10002				
+9999	+9631	371	+9259	743
+10004				
+10001	+9631	373	+9631	373
**+10006				
+10003	+9631	375	+8583	1423
**+10008				
+ 10005	+9631	377	+6637	3371
+10010				
+10007	+9631	379	+9631	379
**+10012				
+10009	+9631	381	+8583	1429
+10014				
+10011	+9631	383	+9631	383
**+10016				
+10013	+9631	385	+9259	757
**+10018				
+10015	+9631	387	+4491	5527
+10020				
+10017	+9631	389	+9631	389

$2n-S$ $(2n-3)-S$	$W_{2n}-S$	$T_{2n} = 2n - W_{2n}$	$V_{2n}-S$	$U_{2n}$
**+20000 +19997	+18031	1969	+17409	2591
**+20002 +19999	+18031	1971	+ 17409	2593
+20004 +20001	+18031	1973	+18031	1973
**+20006 +20003	+18031	1975	+16663	3343
**+20008 +20005	+18031	1977	+16941	3067
+20010 +20007	+18031	1979	+18031	1979
**+20012 +20009	+18031	1981	+5671	14341
**+20014 +20011	+18031	1983	+4101	15913
**+20016 +20013	+18031	1985	+3229	16787
+20018 +20015	+18031	1987	+18031	1987
**+20020 +20017	+18031	1989	+16941	3079
$2n-T$ $(2n-3)-T$	$W_{2n}-T$	$T_{2n} = 2n - W_{2n}$	$V_{2n}-T$	$U_{2n}$
**+40000 +39997	+29737	10263	+ 21567	18433
**+40002 +39999	+29737	10265	+ 22273	17729
+40004 +40001	+29737	10267	+29737	10267
**+40006 +40003	+29737	10269	+21567	18439
+40008 +40005	+29737	10271	+29737	10271
+40010	+29737	10273	+29737	10273



+ 40007				
**+40012	+29737	10275	+10401	29611
+40009				
**+40014	+29737	10277	-56003	96017
+40011				
**+40016	+29737	10279	+27057	12959
+40013				
**+40018	+29737	10281	+25947	14071
+40015				
**+40020	+29737	10283	+24493	15527
+40017				

## 11. Perspectives and Generalizations

**11.1** Other Goldbach sequences ( $G'_{2n}$ ) and ( $G''_{2n}$ ) independent of ( $G_{2n}$ ) may be studied using the increasing sequences of primes ( $W'_{2n}$ ), (see 8.5) and ( $W''_{2n}$ ) defined by :

For any integer  $n \geq 3$ ,  $W''_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq f(n))$ ,  $f$  being a function defined on the interval

$I = [3; +\infty[$  and satisfying the following conditions:

\*  $f$  is strictly increasing on the interval  $I$ .

\*  $\lim_{x \rightarrow +\infty} f(x) = +\infty$ ;  $f(3) = 3$ .

\*  $\forall x \in I, f(x) \leq 2x - 3$ .

For example, one of the following functions defined on  $I$  can be selected.

- $f : x \rightarrow ax + 3 - 3a$ ; ( $a \in \mathbb{R} : 0 < a \leq 2$ ).
- $g : x \rightarrow [4\sqrt{3x} - 9]$  ( $[x]$  being the integer part of the real number  $x$ ).
- $h : x \rightarrow 6\ln\left(\frac{x}{3}\right) + 3$ .

**11.2** Using this method, it would be interesting to study the Schnirelmann density [28] of certain primes such as 3, 5, 7, 11, ..... in the sequence ( $U_{2n}$ ) for  $n \in [K_N; P_N]$  as a function of  $N$ .

**11.3** It is possible to exceed the values shown in the table ( $2n = 10^{1000}$ ) by optimizing this algorithm, using supercomputers and more efficient software as Maple.

**11.4** Diophantine equations and conjectures of the same nature (Lagrange-Lemoine-Levy conjecture [9], [17], [19],[21],[22], [30]) can be processed using similar reasoning and algorithms.

1) To validate the Lagrange-Lemoine-Levy conjecture, we can study the following sequences of primes ( $WL_{2n}$ ), ( $VL_{2n}$ ) and ( $UL_{2n}$ ) defined by :

For any integer  $n \geq 3$ ,  $WL_{2n} = \text{Sup}(p \in \mathcal{P} : p \leq n - 1)$ ,

a) If  $TL_{2n} = (2n + 1 - 2WL_{2n})$  is a **prime**, then let :  $VL_{2n} = WL_{2n}$  and  $UL_{2n} = TL_{2n}$ .

b) If  $TL_{2n}$  is a **composite number**, then there exists an integer  $k$ , ( $1 \leq k \leq n - 3$ ) such that :  $UL_{2(n-k)} + 2k$  is a **prime**; then let :  $VL_{2n} = VL_{2(n-k)}$  and  $UL_{2n} = UL_{2(n-k)} + 2k$ .

2) Using the same type of reasoning, a generalized Bezout-Goldbach conjecture of the following form can be validated :

a) Let  $K$  and  $Q$  be two odd integers, prime to each other : for any integer  $n$  such that :  $2n \geq 3(K + Q)$ , there exist two primes  $U'''_{2n}$  and  $V'''_{2n}$  verifying :

$$K U'''_{2n} + Q V'''_{2n} = 2n.$$

b) Let  $K$  and  $Q$  be two integers of different parity, prime to each other : for any integer  $n$  such that :

$$2n \geq 3(K + Q), \text{ there are two primes } U'''_{2n} \text{ and } V'''_{2n} \text{ verifying:}$$

$$K U'''_{2n} + Q V'''_{2n} = 2n + 1.$$

## 12. Conclusion

**12.1** An unique recursive and explicit Goldbach sequence  $(G_{2n}) = (U_{2n}; V_{2n})$ , verifying :

$(\forall n \in \mathbb{N} + 2 \quad U_{2n} \text{ and } V_{2n} \text{ are primes} : U_{2n} + V_{2n} = 2n)$ ,

has been developed using an simple and efficient "local" algorithm.

**12.2** Silva's [29] record is broken on a personal computer, and it is possible to reach values of the order of  $2n = 10^{1000}$  with a reasonable computation time ( less than three hours for the evaluation of ten terms  $U_{2n}$  and  $V_{2n}$  ).

**12.3** For a given integer  $n \geq 49$ , the evaluation of the terms  $U_{2n}$  and  $V_{2n}$  does not require the computing of all previous terms  $U_{2k}$  and  $V_{2k}$ ,  $(1 \leq k < n - 1)$ . we just need to know the primes  $p_l$ ,  $V_{2r}$  such that :

$$(12.3.1) \quad p_l \leq 7(\ln(2n))^{1.3} \quad \text{and} \quad 2n - 7(\ln(2n))^{1.3} \leq V_{2r} \leq 2n \quad (\text{on average}).$$

This property allows quick computing of  $U_{2n}$  and  $V_{2n}$  even for values of  $2n$  of the order of  $10^{1000}$ .

**12.4** Therefore, the binary Goldbach and the Lagrange-Lemoine-Levy conjectures are true.

## References

1. L. Adleman, K. Mc Curley, Open Problems in Number Theoretic Complexity, II. Algorithmic number theory (Ithaca, NY,1994), 291–322, Lecture Notes in Comput. Sci., 877, Springer, Berlin, (1994).
2. C. Axler, "New Estimates for the nth Prime" 19.4.2 2 Journal of Integer Sequences, Vol. 22, 30 p., (2019),
3. E. Bombieri, Davenport, "Small differences between prime numbers", Proc. Roy. Soc. Ser. A293, pp. 1-18, (1966).
4. R. C. Baker, and Harman, G. "The difference between consecutive primes". Proc. London Math. Soc. (3) 72, 2 (1996), 261–280.
5. R. C. Baker, Harman, G., and Pintz, J. "The difference between consecutive primes". II. Proc. London Math. Soc. (3) 83, 3 (2001), 532–562.
6. J. R. Chen, "On the representation of a large even integer as the sum of a prime and the product of at most two primes". Kexue Tongbao 17 (1966), pp. 385-386 (Chinese).
7. M. Cipolla, "La determinazione assintotica dell n imo numero primo", Rend. Acad. Sci. Fis. Mat. Napoli 8(3) (1902).
8. H. Cramer, "On the order of magnitude of the difference between consecutive prime numbers", Acta Arithmetica vol. 2, (1986), p.23-46.
9. N. Dawar, "Lemoine's Conjecture: A Limited Solution Using Computers", TechRxiv [ Archive online ] (2023).
10. P. Erdos, "On a new method in elementary number theory which leads to an elementary proof of the prime number theorem", Proc. Natl. Acad. Sci. USA 36, pp. 374-384 (1949).
11. Euclid, (trans. Bernard Vitrac), "Les éléments d'Euclide", Ed. PUF Paris, vol.2, p. 444-446 and p. 339-341, (1994).
12. A. Granville, "Harald Cramér and the distribution of prime numbers", *Scandinavian Actuarial Journal*, 1: 12–28,(1995).
13. J. Hadamard, "On the zeros of the function  $\zeta(s)$  of Riemann". C. R. 122, p.1470-1473 (1896), and "On the distribution of zeros of the function  $\zeta'(s)$  and its arithmetical consequences". S. M. F. Bull. 24, pp. 199-220 (1896).
14. G. H. Hardy & Wright, "An introduction to the Theory of numbers", Oxford : Oxford University Press 621 p. (2008).
15. G. H. Hardy, J. E. Littlewood: Some problems of 'partitio numerorum'; III: On the expression of a number as a sum of primes (Acta Math. Vol. 44: pp. 1 – 70, (1922)
16. H. Helfgott, Platt, "The ternary Goldbach conjecture", Gaz. Math. Soc. Math. Fr. 140, pp. 5-18 (2014). "The weak Goldbach conjecture", Gac. R. Soc. Mat. Esp. 16, no. 4, 709-726 (2013). "Numerical verification of the ternary Goldbach conjecture up to  $8.875 \cdot 10^{30}$ ", Exp. Math. 22, n° 4, 406-409 (2013).
17. L. Hodges, "A lesser-known Goldbach conjecture", *Math. Mag.*, 66 (1993): 45–47.
18. H. Iwaniec, Pintz, "Primes in short intervals". Monatsh. Math. 98, pp. 115-143 (1984).
19. J. O. Kiltinen and P. B. Young, "Goldbach, Lemoine, and a Know/Don't Know Problem", Mathematics Magazine, 58(4) (Sep., 1985), p. 195–203.
20. E. Landau, "Handbuch der Lehre von der Verteilung der Primzahlen", vol. 1 and vol. 2 (1909), published by Chelsea Publishing Company (1953).

21. E. Lemoine, "L'intermédiaire de mathématiciens", vol. 1, 1894, p. 179, vol. 3, 1896, p. 151
22. H. Levy, "On Goldbach's conjecture", Math. Gaz." **47** (1963): 274
23. J. Littlewood, "Sur la distribution des nombres premiers", CRAS Paris, vol. 158, (1914), p. 1869-1875.
24. H. Maier, "Primes in short intervals". Michigan Math. J., 32(2):221–225, 1985.
25. T. R. Nicely, "New maximal prime gaps and first occurrences", Mathematics of Computation, **68** (227): 1311–1315, (1999)
26. O. Ramaré, Saouter, "Short effective intervals containing primes", J. Number theory, 98, No. 1, p.10-33 (2003).
27. D. Shanks, "On Maximal Gaps between Successive Primes", Mathematics of Computation, American Mathematical Society, **18** (88): 646–651, (1964).
28. L. Schnirelmann, "Schnirelmann density", Wikipedia, (on line, internet) and "A proof of the fundamental theorem on the density of sums of sets of positive integers", Annals of Math, 2nd series, vol. 43, no. 3, (1942), pp. 523-527.
29. T. O. e Silva, Herzog, Pardi, "Empirical verification of the even Goldbach conjecture and computation of prime gaps up to  $4.10^{18}$ . Math. Comput. 83, no. 288, pp. 2033-2060 (2014).
30. Z-W. Sun, "On sums of primes and triangular numbers" » [archive], arXiv, 2008 (arXiv 0803.3737).
31. T. Tao, "Every odd number greater than 1 is the sum of at most five primes", Math. Comput. 83, no. 286, pp.997-1038 (2014).
32. P. Tchebychev, "Mémoire sur les nombres premiers" J. math. pures et appliquées, 1ère série, t.17, p. 366-390 et p. 381-382, (1852).
33. C.-J. de La Vallée-Poussin, "Recherches analytiques sur la théorie des nombres premiers", Brux. S. sc. 21 B, pp. 183-256, 281-362, 363-397, vol.21 B, pp. 351-368, (1896).
34. A. Vinogradov, "Representation of an odd number as a sum of three primes". Dokl. Akad.Nauk. SSR, 15:291-294, (1937).
35. E.W. Weisstein, "Levy's Conjecture" » [archive], sur MathWorld, CRC Concise Encyclopédie de mathématiques (CRC Press, ), 733-4, (1999).
36. Y. Zhang, "Bounded gaps between primes", Ann. Math. (2) 179, no. 3, pp.1121-1174 (2014).

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.