
A Combined Thermodynamic and Computational Study of Alkaline Earth Metal Cations Complexation by a Fluorescent calix[4]arene Receptor

Andrea Usenik , [Matija Modrušan](#) , Katarina Leko , [Jakov Borovec](#) , [Sven Marinac](#) , [Lucija Hok](#) , Nikola Cindro , [Robert Vianello](#) , [Gordan Horvat](#) , [Josip Požar](#) , [Tomica Hrenar](#) , [Vladislav Tomišić](#) *

Posted Date: 16 January 2025

doi: 10.20944/preprints202501.1208.v1

Keywords: calixarenes; alkaline earth metal cations; solvent effect; inclusion; fluorescence; thermodynamics



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

A Combined Thermodynamic and Computational Study of Alkaline Earth Metal Cations Complexation by a Fluorescent calix[4]arene Receptor

Andrea Usenik ¹, Matija Modrušan ¹, Katarina Leko ¹, Jakov Borovec ¹, Sven Marinac ¹, Lucija Hok ², Nikola Cindro ¹, Robert Vianello ², Gordan Horvat ¹, Josip Požar ¹, Tomica Hrenar ¹ and Vladislav Tomišić ^{1,*}

¹ Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10 000 Zagreb, Croatia

² Department of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička 54, 10 000 Zagreb, Croatia

* Correspondence: vtomisc@chem.pmf.hr

Abstract: Complexation of alkaline earth metal cations with fluorescent tertiary-amide lower-rim calix[4]arene derivative bearing two phenanthridine moieties was studied experimentally (UV spectrophotometry, fluorimetry, isothermal microcalorimetry, NMR spectroscopy) and computationally (classical molecular dynamics and DFT calculations) at 25 °C. The complexation reactions were studied in acetonitrile, methanol, and ethanol, whereby the solvent effect on cation-binding processes was particularly addressed. The complex stability constants and standard reaction thermodynamic quantities (Gibbs energies, enthalpies, and entropies) were determined. The receptor exhibited particularly high affinity towards alkaline earth metal cations in acetonitrile, with peak affinity for Ca²⁺. The stability of all complexes was significantly lower in ethanol and methanol, where the most stable complex was formed with Sr²⁺. The decrease in cation-binding abilities was a consequence of the differences in solvation of the reactants and products of the complexation reactions (involving inclusion of the solvent molecule in the calixarene *cone*), cation charge density, as well as the cation-ligand binding site compatibility. The reactions were enthalpically controlled in acetonitrile, whereas in methanol and ethanol the binding processes were endothermic, and thus entropy driven. The results of ¹H NMR measurements, MD simulations, and DFT calculations provided an insight into the structure of the complexes and the corresponding adducts with solvent molecules, as well as on the structural aspects behind the differences in complexation thermodynamics. Due to the significant increase of its fluorescence upon cation binding, the studied calixarene derivative was proven to be a promising luminescent sensor for alkaline earth metal cations.

Keywords: calixarenes; alkaline earth metal cations; solvent effect; inclusion; fluorescence; thermodynamics

1. Introduction

A wide possibility of aimed functionalization of calixarenes at their lower and/or upper rim, yielding supramolecular receptors with high affinity and selectivity towards particular species, makes these macrocycles excellent backbones for preparing sensors for various ions and neutral molecules.[1–3] Derivatives of these compounds have found a wide array of chemosensory applications, in molecular recognition,[4–9] as sensors in ion-selective electrodes,[10–12] as extractants,[13–18] surfactants,[19–21] catalysts,[22,23] biomimetics,[24–30] drug delivery systems,[31–33] and nanoparticle components.[33–36]

Among calixarenes capable of binding ionic species, those functionalized at the lower rim with electron-donating moieties, such as the ones containing carbonyl groups, have been the subject of

intensive research due to their ability to form very stable complexes with a number of cations.[13,15,37–41] The reason behind this lies in a well-defined binding site comprised of ether and carbonyl oxygen atoms (amides, esters, ketones). It has been shown that tertiary-amide derivatives have the highest affinity for alkali and alkaline earth metal cations,[13,15,39–42] whereas the stability of the complexes with ester, ketone, and secondary-amide derivatives is significantly lower.[37,43–45] This is caused by the difference in the basicity of carbonyl oxygens and, in the case of secondary amides, by intramolecular hydrogen bonding which significantly lowers their affinity towards cations.[43,44,46–48] The compatibility of cation-binding site and calixarene cavity size also significantly affects the receptors' affinity and selectivity in a series of cations of the same charge but with different ionic radius, such as alkali and alkaline earth metal cations. As an illustration, calix[4]arene derivatives usually form most stable complexes with smaller cations (Li^+ , Na^+ , Mg^{2+} , Ca^{2+}),[13,40] but that also strongly depends on the solvent used.[13,40–42,44,45,48–52] The reaction medium significantly impacts the thermodynamic stability of the complexes through solvation of the cation as well as free and complexed receptor. For instance, calixarene derivatives show much higher cation affinity in acetonitrile compared to protic and polar organic solvents (*e.g.* methanol, formamide, dimethyl sulfoxide) due to the higher cation-solvating abilities of the latter, rendering the complexation thermodynamically less favorable.[13] Additionally, the specific solute-solvent interactions play an important role in the complexation processes, either through hydrogen bonding, or inclusion of solvent molecule in the calixarene hydrophobic *basket*. [41,43,44,50,52–54]

Due to the sensitivity of fluorimetric measurements, calixarenes are frequently functionalized with highly fluorescent moieties, like anthracene, naphthalene, dansyl, pyrene, coumarin, etc.,[5,55–60] allowing detection of various analytes at much lower concentrations compared to other often used spectrometric techniques. Phenanthridinyl group is a good example of such a functionality.[57,61,62] However, calixarenes with the cation binding site comprised of ether oxygen atoms and nitrogen atoms of phenanthridine moieties have moderate affinity towards cations, and their solubility in polar solvents is rather limited.[57] In order to enhance these properties, we have recently prepared the ligand **L** (Figure 1), which, in addition to two phenanthridine subunits, contains two tertiary-amide groups forming an efficient cation-binding site.[61] A comprehensive experimental and computational study of its complexation of alkali metal cations in a wide array of solvents was reported. The results revealed an immense solvent effect on the binding equilibria, with the differences in complex stability constants of up to 8 orders of magnitude depending on the reaction medium.[61] The detailed thermodynamic investigations of solvent influence on the cation-binding reactions involving calixarenes were less frequently conducted with alkaline earth metal cations, since their thermodynamic solvation parameters are less available. As the solvation of multiply charged cations is significantly stronger in polar (organic) solvents, an even more pronounced solvent effect can be expected for divalent cations.

Having in mind the above considerations, herein we have investigated the equilibria of complexation reactions between **L** and alkaline earth metal cations in three solvents of different solvation abilities, namely acetonitrile (MeCN), methanol (MeOH), and ethanol (EtOH). The reactions and structures of the complexes were studied by means of UV, fluorescence, and ^1H NMR spectroscopies, isothermal microcalorimetry, as well as computational methods (classical molecular dynamics simulations and DFT). Such a comprehensive and integrated approach yielded detailed thermodynamic and structural insights into the reactions studied. Furthermore, one of the aims of the study was to address the potential application of compound **L** as a fluorescent sensor for these doubly charged cations, and the fluorescence response of the ligand upon cation binding was shown to be quite suitable with this respect.




Figure 1. Structure of phenanthridine-tertiary-amide calix[4]arene derivative **L**.

2. Results and Discussion

2.1. Complexation in Acetonitrile

An example of spectrophotometric titration of calixarene **L** with barium perchlorate in MeCN is shown in Figure 2a. Significant changes in the UV absorption spectrum of the ligand upon cation binding, *i.e.* bathochromic shift and hyperchromic effect, can be seen. Similar was observed for other alkaline earth metal cations (Figures S1–S5, Supporting Information). Furthermore, these changes were accompanied by the occurrence of several well-defined isosbestic points. From the dependence of absorbance at one of the absorption maxima on cation to **L** molar ratio (Figures 2b and S1b–S5b, it was evident that the absorbance increased linearly up to the approximately equimolar ratio, after which it was almost unchanged. Such finding clearly indicated that very stable complexes of 1:1 stoichiometry were formed in all cases (Table 1), and their stability constants could not be determined by means of direct spectrophotometric titrations.

The results of spectrofluorimetric titration of **L** with Ca^{2+} are presented in Figure 3, whereas those corresponding to titrations with other cations are given in the Supporting Information (Figures S6–S9). As in the case of spectrophotometric titration curves, fluorimetric ones exhibited a linear dependence of relative fluorescence intensity on the cation to ligand molar ratio for all cations with a clear break at $\approx 1:1$ ratio, which suggested that the stability constants of all the corresponding complexes in acetonitrile should be exceptionally high. A large increase in the fluorescence of **L** upon cation complexation was observed, especially in the case of Ca^{2+} and Sr^{2+} (Figures 3 and S8). That could be attributed to a significant suppression of photoinduced electron transfer (PET) upon complexation, consequently leading to a considerable fluorescence enhancement.[55,57,89,90] The differences in the emission properties of **L** and its complexes indicated its potential use as a fluorescent sensor for alkaline earth metal cations in acetonitrile, particularly for calcium and strontium ions.

The results of microcalorimetric titrations of **L** in acetonitrile are presented in Figures 4 and S10–S13. Due to the steep sigmoidal ITC curves in the case of experiments involving Ca^{2+} , Sr^{2+} , and Ba^{2+} , the complex stability constants of their complexes with **L** could not be determined by processing the corresponding data. However, the standard complexation enthalpies were successfully obtained from the titration data before the equivalence point (Table 1; it should be noted that the values determined using perchlorate and triflate salts were quite similar). The exception was Mg^{2+} complexation, since the heats obtained with either of the magnesium salts were irreproducible, and therefore the formation of MgL^{2+} could not be characterized calorimetrically.

As stated above, due to the limited sensitivity of the experimental methods used, the complex stability constants could not be determined by direct titrations. However, sufficiently different characteristic absorption spectra of NaL^+ ($\log K = 10.50$, ref. [61]) and ML^{2+} (M stands for Mg, Ca, Sr, Ba) complexes allowed for determination of the complexation equilibrium constants by means of

competitive spectrophotometric titrations (Figures 5 and S14–S16). Thus determined $K(\text{ML}^{2+})$ values (Table 1) suggest that calixarene **L** is indeed a very potent binder of alkaline earth metal cations in MeCN. The $\Delta_r S^\circ$ values listed in Table 1 were calculated by combining the $\Delta_r H^\circ$ values obtained by direct microcalorimetric titrations and the K (*i.e.* $\Delta_r G^\circ$) values determined by means of competitive spectrophotometric titrations.




Figure 2. a) Spectrophotometric titration of **L** ($c = 1.20 \times 10^{-4} \text{ mol dm}^{-3}$) with $\text{Ba}(\text{ClO}_4)_2$ ($c = 1.31 \times 10^{-3} \text{ mol dm}^{-3}$) in MeCN; $\vartheta = (25.0 \pm 0.1)^\circ\text{C}$; $V_0(\text{L}) = 2.2 \text{ cm}^3$; $l = 1 \text{ cm}$. Spectra are corrected for dilution. b) Dependence of the absorbance of **L** at 357 nm on the cation to ligand molar ratio.




Figure 3. a) Fluorimetric titration of **L** ($c = 4.67 \times 10^{-5} \text{ mol dm}^{-3}$) with $\text{Ca}(\text{ClO}_4)_2$ ($c = 7.27 \times 10^{-4} \text{ mol dm}^{-3}$) in MeCN; $\vartheta = (25.0 \pm 0.1)^\circ\text{C}$; $V_0(\text{L}) = 2.5 \text{ cm}^3$; $\lambda_{\text{ex}} = 330 \text{ nm}$; excitation slit 10 nm, emission slit 5 nm. Spectra are corrected for dilution. b) Relative intensity of fluorescence at 370 nm as a function of cation to ligand molar ratio.




Figure 4. Microcalorimetric titration of **L** ($c = 1.15 \times 10^{-4} \text{ mol dm}^{-3}$) with $\text{Ca}(\text{ClO}_4)_2$ ($c = 1.32 \times 10^{-3} \text{ mol dm}^{-3}$) in MeCN; $\vartheta = (25.0 \pm 0.1) \text{ }^\circ\text{C}$; $V(\text{L}) = 1.42 \text{ cm}^3$. a) Thermogram. b) Dependence of successive enthalpy change on the cation to ligand molar ratio.




Figure 5. a) Competitive spectrophotometric titration of the solution containing **L** ($c = 1.10 \times 10^{-4} \text{ mol dm}^{-3}$) and Na^+ ($c = 1.10 \times 10^{-4} \text{ mol dm}^{-3}$) with $\text{Mg}(\text{ClO}_4)_2$ ($c = 4.23 \times 10^{-3} \text{ mol dm}^{-3}$) in MeCN; $\vartheta = (25.0 \pm 0.1) \text{ }^\circ\text{C}$; $V_0(\text{NaL}^+) = 2.2 \text{ cm}^3$; $l = 1 \text{ cm}$. Spectra are corrected for dilution. b) Dependence of the absorbance of **L** at 342 nm on the cation to ligand molar ratio. ■ experimental; — calculated.

Table 1. Thermodynamic parameters for complexation of **L** with alkaline earth metal cations in acetonitrile at 25 $^\circ\text{C}$. Uncertainties of the last digit are given in parentheses as standard errors of the mean ($N = 3\text{--}5$).

Cation	$\log K (\text{M L}^{2+})^a$	$\frac{\Delta_r G^\circ}{\text{kJ mol}^{-1}}^a$	$\frac{\Delta_r H^\circ}{\text{kJ mol}^{-1}}^b$	$\frac{\Delta_r S^\circ}{\text{J K}^{-1} \text{ mol}^{-1}}^{a,b}$
Mg^{2+}	10.99(3)	-62.7(2)		
Ca^{2+}	11.65(5)	-66.5(3)	-49.4(7)	59.1 ± 2.9
Sr^{2+}	11.17(2)	-63.7(1)	-47.28(4) ^c	91.3 ± 1.7
Ba^{2+}	9.66(2)	-55.2(1)	-37.3(5)	94.1 ± 1.4
			-28.0(4)	
			-28.5(5) ^c	

^a determined by spectrophotometric competitive titrations; ^b determined by ITC; ^c triflate salts used instead of perchlorates.

Formation of all complexes in MeCN is both enthalpically and entropically favorable (Table 1). With the increase of ionic radius, the enthalpic contribution to the standard reaction Gibbs energy

becomes less favorable, whereas the opposite holds for the entropic one, and the two become of almost the same magnitude for Ba^{2+} . Although the entropic contributions in the cases of Sr^{2+} and Ba^{2+} complexation are almost identical, the less negative $\Delta_r H^\circ$ for the reaction of **L** with Ba^{2+} results in the lower stability of the barium complex.

To explain the above findings, the impacts of calixarene binding site-cation interactions, cation and complex (de)solvation, as well as solvent inclusion in the calixarene *cone* need to be taken into account. Of course, ligand solvation also takes a part in the complexation process, but it is unnecessary to consider it when comparing the reactions in a single solvent. The contributions of electrostatic interactions between the cation and calixarene binding site are expected to be stronger for smaller cations of higher charge densities. On the other hand, for these cations the desolvation taking place in the course of complexation is more energetically demanding than in the case of the larger ones. From the cation solvation point of view, the increase of reaction entropy with the cation size (Table 1) may seem unexpected, as for smaller cations the desolvation process is entropically more favorable. Thus, another process related to the solvent should play an important role, namely the exothermic but entropically unfavorable inclusion of acetonitrile molecule in the calixarene *basket*.^[41,43,44,50,52–54,91] Indeed, the occurrence of this process was proven experimentally (see section 3.3), and this finding was supported by molecular dynamics simulations of the ML^{2+} complexes. The results of the latter suggested that all cations were coordinated with 4 ether and both carbonyl oxygen atoms, whereby the average coordination by phenanthridine nitrogen atoms increased steadily with the ionic radius (Table S1). Among the representative structures (Figure 6), those containing included MeCN (denoted as MLMeCN^{2+}) molecules in the calixarene hydrophobic cavity with the nitrile group facing towards the cation and coordinating it were the most abundant in the simulation timeframe. The cation coordination by the included solvent molecule provided additional complex stabilization, whereby the most favorable $E(\text{M}^{2+}-\text{MeCN}_{\text{incl}})$ was with Ca^{2+} , then Mg^{2+} , and the lowest with Sr^{2+} and Ba^{2+} (Table S1; see also section 3.5). This certainly contributes to the differences in the corresponding reaction enthalpies. The complex adducts MgLMeCN^{2+} and CaLMeCN^{2+} were present during the whole simulation time, with no exchange of the included MeCN molecule. In the cases of Sr^{2+} and Ba^{2+} complexes, two other forms (MLMeCN^{2+} with MeCN methyl group facing the cation and ML^{*2+} without included solvent molecule) were also shortly present and two acetonitrile molecules exchanged in the *basket* during simulation. These findings indicated somewhat more entropically disadvantageous inclusion within the complexes with the smaller cations, which is in line with the trend of the observed reaction entropies. Overall, the thermodynamic data regarding complexation in acetonitrile listed in Table 1 suggested partial enthalpy-entropy compensation resulting in similar ML^{2+} stabilities, and that can be accounted for by a combination of all the effects described above.




Figure 6. Representative structures of ML^{2+} complexes and their MeCN adducts obtained by MD simulations at 25 °C. Hydrogen atoms of L are omitted for clarity. Numbers in parentheses represent percentages of total simulation time in which the species existed.

2.2. Complexation in Methanol and Ethanol

Spectrophotometric, fluorimetric, and microcalorimetric titrations of L with alkaline earth metal perchlorates in methanol and ethanol are shown in Figures 7–9 and S21–S36. Despite rather small spectral changes, the MgL^{2+} complex stability constants in MeOH and EtOH were successfully determined by processing the spectrophotometric titration data (Table 2). However, the fluorimetrically obtained spectral data were insufficiently reproducible to allow for the quantitative characterization of the L binding affinity for Mg^{2+} . Stability constants of other ML^{2+} complexes were determined by both spectrometric methods. In addition, the reaction enthalpies and entropies were obtained calorimetrically for Sr^{2+} and Ba^{2+} binding by L in MeOH, as well as Ca^{2+} and Ba^{2+} in EtOH (Table 2). The approximately isoenthalpic complexation of Mg^{2+} in both alcohols and too large dilution heats of Ca^{2+} salts in MeOH prevented reliable microcalorimetric characterization of the corresponding complexation reactions.




Figure 7. a) Spectrophotometric titration of **L** ($c = 1,26 \times 10^{-4} \text{ mol dm}^{-3}$) with $\text{Sr}(\text{ClO}_4)_2$ ($c = 1.96 \times 10^{-3} \text{ mol dm}^{-3}$) in EtOH; $\vartheta = (25.0 \pm 0.1) \text{ }^\circ\text{C}$; $V_0(\text{L}) = 2.2 \text{ cm}^3$; $l = 1 \text{ cm}$. Spectra are corrected for dilution. b) Dependence of the absorbance of **L** at 360 nm on the cation to ligand molar ratio. ■ experimental; — calculated.




Figure 8. a) Fluorimetric titration of **L** ($c = 4.25 \times 10^{-5} \text{ mol dm}^{-3}$) with $\text{Ca}(\text{ClO}_4)_2$ ($c = 3.30 \times 10^{-2} \text{ mol dm}^{-3}$) in MeOH; $\vartheta = (25.0 \pm 0.1) \text{ }^\circ\text{C}$; $V_0(\text{L}) = 2.5 \text{ cm}^3$; $\lambda_{\text{ex}} = 330 \text{ nm}$; excitation slit 10 nm, emission slit 10 nm. Spectra are corrected for dilution. b) Relative intensity of fluorescence at 370 nm on the cation to ligand molar ratio. ■ experimental; — calculated.




Figure 9. Microcalorimetric titration of **L** ($c = 1.53 \times 10^{-4} \text{ mol dm}^{-3}$) with $\text{Ba}(\text{ClO}_4)_2$ ($c = 1.64 \times 10^{-2} \text{ mol dm}^{-3}$) in MeOH; $\vartheta = (25.0 \pm 0.1) \text{ }^\circ\text{C}$; $V(\text{L}) = 1.42 \text{ cm}^3$. a) Thermogram. b) Dependence of successive enthalpy change on the cation to ligand molar ratio. ■ experimental; – calculated.

Table 2. Thermodynamic parameters for complexation of **L** with alkaline earth metal cations in methanol and ethanol at 25 °C. Uncertainties of the last digit are given in parentheses as standard errors of the mean ($N = 3\text{--}5$).

Solvent	Cation	$\log K(\text{ML}^{2+})$			$\frac{\Delta_r G^\circ}{\text{kJ mol}^{-1}}$	$\frac{\Delta_r H^\circ}{\text{kJ mol}^{-1}}$ ^c	$\frac{\Delta_r S^\circ}{\text{J K}^{-1} \text{ mol}^{-1}}$ ^c
		a	b	c			
MeOH	Mg ²⁺	2.74(2)			-15.6(1) ^a	≈0	
	Ca ²⁺	2.82(2)	2.15(4)		-16.1(1) ^a		
	Sr ²⁺	3.29(2)	3.29(2)	3.49(2)	-19.9(1) ^c	12.2(4)	108(1)
	Ba ²⁺	2.96(1)	2.69(1)	2.99(1)	-17.08(1) ^c	13.6(5)	103(2)
EtOH	Mg ²⁺	4.15(2)			-23.7(1) ^a	≈0	
	Ca ²⁺	4.33(2)	3.78(1)	4.30(5)	-24.6(3) ^c	7.2(5)	107(1)
	Sr ²⁺	5.20(2)	5.25(1)	5.09(2)	-29.1(2) ^c	3.5(1)	109(1)
	Ba ²⁺	4.85(3)	4.87(2)	4.95(1)	-28.28(5) ^c	5.7(2)	114(1)

^a spectrophotometry; ^b spectrofluorimetry; ^c microcalorimetry.

Depending on the cation, stability constants of ML^{2+} complexes in EtOH are approximately 1.5 to 2 orders of magnitude higher than in MeOH. This is mostly because of the difference in cation solvation in these solvents (Table S12), [92] as endothermic complexation (Table 2) in both indicates the dominance of desolvation contribution to the reaction enthalpy. On the other hand, the release of rather strongly bound solvent molecules into the bulk is entropically favorable, which is apparently the main complexation driving force.

As in MeCN, the results of molecular dynamics simulations suggest that all cations are coordinated by all oxygen atoms in the calixarene binding site (Table S2). The solvent adducts with both orientations of the included methanol or ethanol molecule (alkyl group towards the cation, MLMeOH^{2+} , MLEtOH^{2+} ; hydroxyl group towards the cation, MLMeOH'^{2+} , MLEtOH'^{2+} ; Figures 10 and S46) are the dominant forms of the complexes, although the species with “empty” basket (ML^{*2+}) are also present in the case of larger cations. No coordination of Mg^{2+} by phenanthridine nitrogen atoms was observed (Tables S2 and S3), whereas unlike in MeCN, for larger cations the average number of coordinating phenanthridine N atoms in MeOH is almost independent of the cation radius, and in EtOH it peaks at Sr^{2+} . This value is generally lower in MeOH than EtOH (Tables S2 and S3), suggesting that stronger coordination of metal cations by oxygen atom of the included methanol molecule “pulls” the cation away from the phenanthridine subunits more strongly. A larger number of exchanged alcohol molecules in the calixarene basket compared to MeCN was observed (Tables S1–S3), indicating lower extent of solvent inclusion in the former cases.




Figure 10. Representative structures of ML^{2+} complexes and their MeOH adducts obtained by MD simulations at 25 °C. Hydrogen atoms of **L** are omitted for clarity. Numbers in parentheses represent percentages of total simulation time in which the species existed.

2.3. NMR Study of Solvent Inclusion in the Hydrophobic Cavity of **L** and ML^{2+} Complexes

In our recent paper,[61] we have examined the inclusion of solvent molecules in the hydrophobic cavity of **L** and its complexes with alkali metal cations by molecular dynamics simulations. To confirm this process experimentally, herein we have recorded and compared the ¹H NMR spectra of the ligand and its complexes with alkaline earth metal cations in deuterated chloroform, acetonitrile,

methanol, and ethanol. The signals of the upper-rim aromatic protons (Ar-H) of free **L** are significantly downfield shifted in CD₃CN compared to CDCl₃ (Figures 11 and S48, Table S4), which can serve as an indication of acetonitrile molecule inclusion into the calixarene *basket*. [13,52,91] The shift is much less pronounced, although observable, in the case of deuterated methanol and ethanol, implying lower extent of adduct formation. These findings are in agreement with the results of previously [61] performed molecular dynamics simulations.

The signals of all **L** protons in its complexes with alkaline earth metal cations are considerably shifted with respect to the free ligand in all solvents examined (Figures 11 and S49–S55, Tables S4–S8). As in the case of uncomplexed **L**, the CD₃CN *vs.* CDCl₃ difference in chemical shifts of Ar-H protons of ML²⁺ complexes is substantial ($\Delta\delta$ up to 0.45 ppm), whereas the corresponding differences for CD₃OD and C₂D₅OD are lower ($\Delta\delta$ up to 0.14 ppm). These downfield shifts of proton signals relative to chloroform solutions again reveal the inclusion of both acetonitrile and alcohol molecules in the calixarene *cone*. In the case of latter solvents, the $\Delta\delta$ values for complexes are larger than those for the free ligand, indicating higher affinity of the complex species for inclusion of solvent molecules and possibly their different orientations in the *basket*, as suggested by MD and quantum chemical calculations.




Figure 11. ¹H NMR spectra of **L** (left) and BaL²⁺ complex (right) in deuterated chloroform, acetonitrile, methanol, and ethanol at 25 °C.

2.4. Comparison of Thermodynamic Complexation Parameters in Acetonitrile and Alcohols

The complex stability constants are by far the highest in acetonitrile, four to eight orders of magnitude lower in ethanol, and six to nine orders of magnitude lower in methanol (Figure 12). Stability of the investigated complexes decreases in the following sequences:

Ca²⁺ > Sr²⁺ ≈ Mg²⁺ > Ba²⁺ (MeCN), Sr²⁺ > Ba²⁺ ≈ Ca²⁺ ≈ Mg²⁺ (MeOH), and

Sr²⁺ > Ba²⁺ > Ca²⁺ ≈ Mg²⁺ (EtOH). Selectivity of **L** is not pronounced in either of the solvents, with the largest differences in binding affinities being for Ca²⁺ and Ba²⁺ in acetonitrile ($K(\text{CaL}^{2+}) / K(\text{BaL}^{2+}) \approx 100$) and Sr²⁺ and Mg²⁺ in ethanol ($K(\text{SrL}^{2+}) / K(\text{MgL}^{2+}) \approx 10$).

As can be seen by inspecting the data in Tables 1 and 2 and Figure 13, the complexation reactions are enthalpically controlled in acetonitrile, whereas in alcohol solvents they are entropy driven and either endothermic (Ca²⁺, Sr²⁺, Ba²⁺) or almost isoenthalpic (Mg²⁺). We were able to determine all thermodynamic reaction parameters for the formation of SrL²⁺ and BaL²⁺ complexes in all reaction media examined, and comparison between the enthalpic and entropic contributions to the corresponding reaction Gibbs energies is presented in Figures S37 and 13. The main reason for significantly higher complex stability in MeCN compared to MeOH and EtOH is the favorable enthalpic contribution in the former case, whereas the entropic one differs much less among the solvents. Similar can be concluded for Mg²⁺ and Ca²⁺.




Figure 12. Comparison of stability constants of L complexes with alkaline earth metal cations in acetonitrile, methanol, and ethanol. Ionic radii (r) corresponding to cations with coordination number 6 are taken from ref. [93].




Figure 13. Standard thermodynamic parameters for complexation of L with Ba²⁺ in MeCN, MeOH, and EtOH at 25 °C.

As already noted, to more deeply understand the differences in $\Delta_r X^\circ$ ($X = G, H, S$) values given in Tables 1 and 2, the effects of all processes taking place upon complexation should be considered. The available enthalpies of transfer of Ca²⁺, Sr²⁺, and Ba²⁺ from MeCN to MeOH (Table S12, ref. [92]) show that this process is very exothermic leading to negative $\Delta_t G^\circ(\text{M}^{2+}, \text{MeCN} \rightarrow \text{MeOH})$ (Table S12, ref.

[92]) and thus considerably favouring the complexation in MeCN compared to MeOH, particularly in the case of smaller cations with higher q/r ratio. Similar, though less significant, can be concluded for ethanol as a solvent. Compared to MeCN, the peak affinity of L in alcohols is shifted from Ca^{2+} to the larger Sr^{2+} cation due to the less unfavorable desolvation of the latter relative to the former (Tables S10–S12).

The energetically beneficial contributions of electrostatic interactions between the cation and calixarene binding site and solvent inclusion into the hydrophobic *basket* of the complex ($\Delta_r H^\circ$ and $\Delta_r S^\circ < 0$ for both) to $\Delta_r G^\circ$ are more pronounced for the smaller cations in all solvents explored. According to the MD and NMR results, the extent of solvent inclusion is higher in acetonitrile than in alcohol solutions. This process enthalpically favors and entropically disfavors complexation reaction. Therefore, from the aspects of cation desolvation and solvent inclusion processes, a huge difference in complexation enthalpies in MeCN vs. MeOH and EtOH (Tables 1 and 2) is easily understandable. Besides, the relatively low difference in (positive) reaction entropies in these solvents (lower than expected based solely on cation desolvation) can also be accounted for by considering the larger extent of entropically disadvantageous acetonitrile inclusion compared to that of methanol and ethanol.

2.5. Quantum Chemical Calculations

To explain various structural phenomena arising in the investigated complexes, extensive quantum chemical calculation of ML^{*2+} , MLMeCN^{2+} , and MLMeCN'^{2+} complexes with alkaline earth metal cations were performed. Geometry optimization for each complex was followed by ground-state geometry validation *via* harmonic frequency calculations. Optimized structures were overlaid according to the least-squares-fit plane through the ether oxygen atoms of L and presented in Figures 14 and S56. In the ML^{*2+} complexes without included solvent molecule the position of alkaline earth metal cation is shifted within the ligand from Mg^{2+} to Ba^{2+} (Fig. S56 and Table 3). Two clusters can be identified, one comprising Mg^{2+} and Ca^{2+} complexes and the other containing those with Sr^{2+} and Ba^{2+} . In the former, Mg^{2+} and Ca^{2+} are closer to the least-squares-fit plane (distances are 0.678 and 0.770 Å from the plane, respectively, Table 3) and the phenanthridine units are “pushed away” from the cations (distances of 4.038 and 3.968 Å, Table S9). In the second cluster Sr^{2+} and Ba^{2+} are further away from the plane (distances are 1.225 and 1.399 Å, respectively, Table 3), whereas the phenanthridine units are much closer to the cations (distances of 3.132 and 3.145 Å, Table Table S9).

Table 3. Interatomic distances in complexes of L and cations calculated at the B3LYP-D3BJ/def2SVP level of the theory.

Cation	ML^{*2+}	MLMeCN^{2+}	MLMeCN'^{2+}
	$r / \text{Å}$ (plane ^a to cation)		
Mg^{2+}	0.678	0.684	0.451
Ca^{2+}	0.770	0.788	0.566
Sr^{2+}	1.225	1.237	0.948
Ba^{2+}	1.399	1.411	1.207
	$r / \text{Å}$ (plane ^a to solvent ^b)		
Mg^{2+}	–	4.757	3.112
Ca^{2+}	–	4.821	3.280
Sr^{2+}	–	4.839	3.082
Ba^{2+}	–	4.808	3.027
	$r / \text{Å}$ (cation to solvent ^b)		
Mg^{2+}	–	5.494	3.564
Ca^{2+}	–	5.655	3.847
Sr^{2+}	–	6.129	4.030
Ba^{2+}	–	6.262	4.234

^a Least-squares-fit plane through the 4 ether oxygen atoms of the calixarene units. ^b Solvent center of mass.

The influence of the included MeCN molecule on the complex structural characteristics was probed using the different orientations of the solvent observed in MD simulations (Figure 6). The distances between the cation and the least-squares-fit plane are almost the same for the solvent-free species and the adduct with acetonitrile methyl group pointing towards the cation (MLMeCN²⁺, Table 3), suggesting that there are no strong electrostatic interactions between the cations and MeCN molecule. On the other hand, in the cases where the acetonitrile molecule is oriented with the nitrile group facing cation (MLMeCN¹²⁺), these distances change significantly (Table 3), *i.e.* cations are shifted in the direction of the plane by more than 0.2 Å which indicates their much stronger interaction (coordination) with the included solvent. This can also be clearly seen from the position of the MeCN molecule. In the former case, the distances of the solvent molecule from the plane are ≈4.8 Å for all cations, whereas in the latter, MeCN is much closer (for more than 1.5 Å) to the plane, and consequently to the metal ion. As in the solvent-free species, in the MLMeCN²⁺ and MLMeCN¹²⁺ adducts the distances between Sr²⁺ and Ba²⁺ and phenanthridine nitrogens are considerably shorter than those corresponding to Mg²⁺ and Ca²⁺ (Table S9). This implies that, contrary to smaller cations, nitrogen atoms participate in the coordination of the larger ones.

All the above-described findings are in accordance with those obtained by classical molecular dynamics simulations and additionally support conclusions drawn from the experimental results.




Figure 14. Optimized geometries of a) MLMeCN²⁺ and b) MLMeCN¹²⁺ complexes with alkaline earth cations: Mg²⁺ (white), Ca²⁺ (orange), Sr²⁺ (green) and Ba²⁺ (blue) calculated by B3LYP-D3BJ/def2SVP method.

3. Materials and Methods

3.1. Physicochemical Measurements

Chemicals. Acetonitrile (J. T. Baker, HPLC Gradient Grade) was used without additional purification, whereas methanol (J. T. Baker, HPLC Gradient Grade) and ethanol (Sigma Aldrich, Spectranal) were distilled prior to use. The salt solutions for complexation studies were prepared by direct weighing and dissolution of the following salts in volumetric flasks: magnesium perchlorate hexahydrate (Sigma Aldrich, 99 %), magnesium triflate (Mg(trf)₂, Sigma Aldrich, 97 %), calcium perchlorate tetrahydrate (Sigma Aldrich, 99 %), calcium triflate (Ca(trf)₂, Sigma Aldrich, 99.9 %), strontium perchlorate trihydrate (Alpha Aesar, 98 %), barium perchlorate (Fluka, ≥ 98 %), barium triflate (Ba(trf)₂, Sigma Aldrich, 98 %), and sodium perchlorate (Sigma Aldrich 98+ %). The perchlorate and triflate salts were chosen based on their inertness regarding ion pairing. The triflate salts were also used in order to investigate the potential influence of water present in perchlorate salts and to avoid large dilution heats of the latter in ITC experiments. For ¹H NMR measurements the following solvents were used: CDCl₃ (Eurisotop, 99.80 % D), CD₃CN (Eurisotop, 99.80 % D), CD₃OD (Eurisotop, 99.80 % D), C₂D₅OD (Cambridge Isotope Laboratories, 99 % D).

UV absorption spectrometry. Spectrophotometric titrations were performed using UV-Vis spectrophotometers Agilent Cary 50 and Cary 60. The experiments were conducted in a way that the solution of **L** ($V_0 = 2.2 \text{ cm}^3$, $l = 1 \text{ cm}$) was titrated with the appropriate salt solution. The spectra were recorded with a step of 1 nm and an integration time of 0.2 s with baseline correction, at a constant temperature of $(25.0 \pm 0.1) \text{ }^\circ\text{C}$. For titration of **L** with Mg^{2+} , Ca^{2+} , Sr^{2+} , and Ba^{2+} , the solutions of following concentrations were used: 1.0×10^{-4} to $1.6 \times 10^{-4} \text{ mol dm}^{-3}$ (**L**), 9.2×10^{-4} to $1.1 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Mg}(\text{ClO}_4)_2$), 1.5×10^{-3} to $1.1 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{ClO}_4)_2$), $2.1 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{trf})_2$), 1.1×10^{-3} to $9.8 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Sr}(\text{ClO}_4)_2$),

1.3×10^{-3} to $1.0 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{ClO}_4)_2$), and $2.1 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{trf})_2$).

For competitive titrations in MeCN, solution of **L** ($c = 1.0 \times 10^{-4}$ to $1.1 \times 10^{-4} \text{ mol dm}^{-3}$) and sodium perchlorate ($c = 1.0 \times 10^{-4}$ to $1.1 \times 10^{-4} \text{ mol dm}^{-3}$) was titrated with the solutions of alkaline earth metal salts of the following concentrations: $4.23 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Mg}(\text{ClO}_4)_2$), $1.6 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{ClO}_4)_2$), $1.5 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Sr}(\text{ClO}_4)_2$), and $4.4 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{ClO}_4)_2$). The experiments were performed at least in triplicate. The obtained spectrophotometric data were processed using the HYPERQUAD software package.[63]

Fluorimetry. Spectrofluorimetric titrations were carried out using the Agilent Cary Eclipse spectrofluorimeter. Experiments were conducted in a way that the solution of **L** ($V_0 = 2.5 \text{ cm}^3$) was titrated with the corresponding salt solution. The spectra were recorded with a step of 2 nm and an integration time of 0.4 s at a constant temperature of $(25.0 \pm 0.1) \text{ }^\circ\text{C}$. For titration of **L** with Mg^{2+} , Ca^{2+} , Sr^{2+} , and Ba^{2+} , the solutions of following concentrations were used: 4.1×10^{-5} to $4.8 \times 10^{-4} \text{ mol dm}^{-3}$ (**L**), $6.3 \times 10^{-4} \text{ mol dm}^{-3}$ ($\text{Mg}(\text{ClO}_4)_2$), $4.7 \times 10^{-4} \text{ mol dm}^{-3}$ ($\text{Mg}(\text{trf})_2$), 7.3×10^{-4} to $3.3 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{ClO}_4)_2$), 4.1×10^{-4} to $9.8 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Sr}(\text{ClO}_4)_2$), and 5.7×10^{-4} to $3.1 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{ClO}_4)_2$). Where necessary, overlap between the excitation and emission spectra of the ligand were taken into account during fitting. The experiments were repeated at least three times. The obtained fluorimetric data were processed using the HYPERQUAD software package.[63]

Isothermal titration calorimetry. Microcalorimetric titrations were carried out using a Malvern MicroCal VP-ITC microcalorimeter with a reaction cell volume of 1.42 cm^3 . The instrument reliability was verified by carrying out the microcalorimetric titrations of 18-crown-6 (Sigma Aldrich, 99%) with BaCl_2 (Sigma Aldrich, 99.9 %) in H_2O at $25 \text{ }^\circ\text{C}$. The obtained thermodynamic complexation parameters ($\Delta_r H^\circ = -32.2 \text{ kJ mol}^{-1}$; $T\Delta_r S^\circ = -10.7 \text{ kJ mol}^{-1}$; $K = 5772 \text{ dm}^3 \text{ mol}^{-1}$) were in excellent agreement with the literature values ($\Delta_r H^\circ = 31.42 \text{ kJ mol}^{-1}$; $T\Delta_r S^\circ = -9.90 \text{ kJ mol}^{-1}$; $K = 5900 \text{ dm}^3 \text{ mol}^{-1}$).[64]

For titration of **L** with Mg^{2+} , Ca^{2+} , Sr^{2+} , and Ba^{2+} , the solutions of following concentrations were used: 1.1×10^{-4} to $1.6 \times 10^{-4} \text{ mol dm}^{-3}$ (**L**), 9.2×10^{-4} to $1.1 \times 10^{-2} \text{ mol dm}^{-3}$ ($\text{Mg}(\text{ClO}_4)_2$), 1.3×10^{-3} to $1.7 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{ClO}_4)_2$), $1.8 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ca}(\text{trf})_2$), 1.2×10^{-3} to $8.1 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Sr}(\text{ClO}_4)_2$), 1.2×10^{-3} to $1.9 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{ClO}_4)_2$), and $2.1 \times 10^{-3} \text{ mol dm}^{-3}$ ($\text{Ba}(\text{trf})_2$).

Experiments were conducted at a constant temperature of $25 \text{ }^\circ\text{C}$ by addition of the alkaline earth metal salt solution to the solution of ligand **L**, or the solution containing NaL^+ in competition experiments, by means of an automated burette ($V = 300 \text{ } \mu\text{L}$). The measured enthalpy changes were corrected for titrant dilution heats. The experiments were carried out at least in triplicate. Thus obtained microcalorimetric data were processed using Microcal OriginPro 7.0 and OriginPro 7.5 or in the case of competitive titrations using the HypDH program.[65]

NMR studies ^1H NMR experiments were conducted on Bruker Avance III HD 400 MHz/54 mm Ascend equipped with 5 mm PA BBI 1H/D-BB probe with z -gradient and automatic tuning. All proton spectra were obtained using 64K data points, spectral width of 20 ppm based on 16 or 128 (in the case of acetonitrile solutions) scans. Solutions of free ligand were prepared by dissolving the solid in the certain solvent, whereas in the case of measurements involving ML^{2+} complexes, large excess of salt was added in the solution to ensure that all present ligand was complexed. In the case of complex solutions in chloroform, ligand solution and excess of salt were mixed and the spectra were recorded daily until no further changes of chemical shifts were observed.

3.2. Computational Investigations

Molecular dynamics simulations. The molecular dynamics simulations were carried out by means of the GROMACS package (version 2018.6).[66–71] Intramolecular and nonbonded intermolecular interactions were modelled by the CHARMM36 (Chemistry at HARvard Macromolecular Mechanics) force field.[72] Partial charges of phenanthridine atoms were calculated for a model compound, namely 6-(phenoxyethyl)phenanthridine with CGENFF web server.[73–75] The initial structures of calixarene complexes were made by placing a cation between ether and carbonyl oxygen atoms of the lower-rim substituents. The ML^{2+} species (M denotes alkaline earth metal) were solvated in a cubical box (edge length 60 Å) of acetonitrile, methanol, or ethanol with periodic boundary conditions. The solvent boxes were equilibrated prior to solvation of the complexes. The solute concentration in such a box was about 0.01 mol dm⁻³. In all simulations an energy minimization procedure was performed followed by 50.5 ns of NpT production simulation, with the time constant of 1 ps. The first 0.5 ns of production simulation were discarded in the data analysis. The integrator used for the propagation and for the temperature control was stochastic dynamics algorithm[76] with a time step of 1 fs. During simulation, the temperature and pressure were kept at 298 K and 1 bar, respectively. The cutoff radius for nonbonded van der Waals and short-range Coulomb interactions was 15 Å. Long-range Coulomb interactions were treated by the Ewald method as implemented in the PME (Particle Mesh Ewald) procedure.[77] The representative molecular structures of ML^{2+} complexes were obtained by principal component analysis on the coordination matrix whose rows contained cation-carbonyl, cation-ether oxygen atoms, and cation-phenanthridine nitrogen atoms distances during simulation. Angles between metal cations and carbonyl groups were added to the coordination matrix as well. The chosen structures were closest to the centroids of the most populous clusters in space defined by the first two principal components. Figures of molecular structures were created using VMD software.[78]

Quantum chemical calculations. Optimizations of geometries for all investigated complexes were performed using the hybrid functional B3LYP[79,80] with the D3 version of Grimme's dispersion[81] and Becke-Johnson damping in combination with the def2-SVP[82,83] basis set. The initial geometries of complexes with and without the solvent molecule for the optimization procedure were assembled from the first optimized structure and re-optimized. To confirm that the obtained geometries were local minima, harmonic frequency calculations were performed and analyzed.[84,85] The standard Gibbs energies of formation were calculated at $T = 298.15$ K and $p = 101\,325$ Pa. All quantum chemical calculations were carried out using the Gaussian 16 program package.[86] The least-squares fit plane through ether oxygen atoms of L was determined by using the advanced regression module implemented in our code *moonee*.[87,88]

4. Conclusions

Thermodynamics of alkaline earth metal cations binding by the fluorescent calix[4]arene derivative L was investigated in acetonitrile, methanol, and ethanol. Complexation reactions in acetonitrile are both enthalpically and entropically favorable, with the overall complex stability being the highest in the case of CaL^{2+} . With increasing cation radius, the binding process becomes less exothermic, while the reaction entropy rises. Contrary to acetonitrile, reactions in methanol and ethanol are enthalpically unfavorable and thus entropy driven. This is mostly due to the energetically quite demanding desolvation of divalent cations in alcohols (more so in MeOH than EtOH). As a consequence, the complex stability constants are far lower in ethanol compared to acetonitrile, and even lower in methanol. In addition, the peak affinity in alcohols is shifted from Ca^{2+} to Sr^{2+} .

Another process which plays a significant role in determining the complexation equilibria, namely the inclusion of solvent molecule in the calixarene *cone*, was observed experimentally and corroborated by means of computational methods. The effect of this enthalpically advantageous and entropically unfavorable process is of considerable importance, particularly for binding of alkaline earth metal ions. This is so because, in contrast to complexes of L with alkali metal cations,[61] the

included MeCN, MeOH, or EtOH molecule provides an additional energetically favorable coordination of doubly charged second-group cations (of higher charge density) by partially negative nitrogen or oxygen solvent atom. The results of DFT computations carried out for acetonitrile adducts showed that the positions of cation and solvent molecule in the complex were quite dependent on the ion size as well as the orientation of the included solvent molecule. The distances between cations and MeCN molecule from the least-squares plane defined by ether oxygen atoms were much shorter when acetonitrile was oriented with nitrile group facing the cation than for its opposite orientation, as a result of the solvent participation in metal coordination in the former case.

The substantial differences between the emission properties of **L** and its complexes suggested that the investigated calixarene derivative could serve as a quite sensitive fluorescent sensor for alkaline earth metal cations, particularly for Ca²⁺ and Sr²⁺ in acetonitrile.

Supplementary Material: The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

Acknowledgements: This research was funded by Croatian Science Foundation (projects MacroSol, grant number IP-2019-04-9560, CalixCORE, grant number IP-2024-05-3012, and TargetCh, grant number IP-2022-10-9525) and European Regional Development Fund (infrastructural project CluK, grant number KK.01.1.1.02.0016).

Conflicts of Interest: There are no conflicts to declare.

References

1. Asfari, Z. *Calixarenes 2001*; Böhmer, V., Harrowfield, J., Vicens, J., Eds.; Kluwer Academic Publishers: Dordrecht, 2001; ISBN 978-0-306-47522-1.
2. Gutsche, C.D. *Calixarenes: An Introduction*; Monographs in supramolecular chemistry; 2nd ed.; RSC publishing: Cambridge, 2008; ISBN 978-0-85404-258-6.
3. *Calixarenes and Beyond*; Neri, P., Sessler, J.L., Wang, M.-X., Eds.; Springer International Publishing: Cham, 2016; ISBN 978-3-319-31865-3.
4. Baldini, L.; Sansone, F.; Casnati, A.; Ungaro, R. Calixarenes in Molecular Recognition. In *Supramolecular Chemistry*; Gale, P.A., Steed, J.W., Eds.; Wiley, 2012 ISBN 978-0-470-74640-0.
5. Kumar, R.; Sharma, A.; Singh, H.; Suating, P.; Kim, H.S.; Sunwoo, K.; Shim, I.; Gibb, B.C.; Kim, J.S. Revisiting Fluorescent Calixarenes: From Molecular Sensors to Smart Materials. *Chem. Rev.* **2019**, *119*, 9657–9721, doi:10.1021/acs.chemrev.8b00605.
6. Macreadie, L.K.; Gilchrist, A.M.; McNaughton, D.A.; Ryder, W.G.; Fares, M.; Gale, P.A. Progress in Anion Receptor Chemistry. *Chem* **2022**, *8*, 46–118, doi:10.1016/j.chempr.2021.10.029.
7. Mohammed, F.A.; Xiao, T.; Wang, L.; Elmes, R.B.P. Macrocyclic Receptors for Anion Recognition. *Chem. Commun.* **2024**, *60*, 11812–11836, doi:10.1039/D4CC04521A.
8. Prata, J.V.; Barata, P.D. Fostering Protein–Calixarene Interactions: From Molecular Recognition to Sensing. *RSC Adv.* **2016**, *6*, 1659–1669, doi:10.1039/C5RA19887A.
9. Crowley, P.B. Protein–Calixarene Complexation: From Recognition to Assembly. *Acc. Chem. Res.* **2022**, *55*, 2019–2032, doi:10.1021/acs.accounts.2c00206.
10. Chung, T.D.; Kim, H. Electrochemistry of Calixarene and Its Analytical Applications. *J. Inclusion Phenom. Mol. Recognit. Chem.* **1998**, *32*, 179–193, doi:10.1023/A:1008059326436.
11. Beer, P.D.; Gale, P.A. Anion Recognition and Sensing: The State of the Art and Future Perspectives. *Angew. Chem. Int. Ed.* **2001**, *40*, 486–516, doi:10.1002/1521-3773(20010202)40:3<486::AID-ANIE486>3.0.CO;2-P.
12. Danil De Namor, A.F.; Alharthi, S.; El Gamouz, A.; Al Hakawati, N.; Cox, B.G. Calix[4] Based Hg(II) Ion Selective Electrodes: A Thermodynamic Protocol to Address the Selectivity versus the Hosting Capacity Paradigm in the Selection of the Carrier. *Electrochim. Acta* **2018**, *290*, 686–694, doi:10.1016/j.electacta.2018.09.101.
13. Danil de Namor, A.F.; Cleverley, R.M.; Zapata-Ormachea, M.L. Thermodynamics of Calixarene Chemistry. *Chem. Rev.* **1998**, *98*, 2495–2526, doi:10.1021/cr970095w.

14. Arnaud-Neu, F.; Barrett, G.; Fanni, S.; Marrs, D.; McGregor, W.; McKervey, M.A.; Schwing-Weill, M.-J.; Vetrogon, V.; Wechsler, S. Extraction and Solution Thermodynamics of Complexation of Alkali and Alkaline-Earth Cations by Calix[4]Arene Amides. *J. Chem. Soc., Perkin Trans. 2* **1995**, 453, doi:10.1039/p29950000453.
15. Creaven, B.S.; Donlon, D.F.; McGinley, J. Coordination Chemistry of Calix[4]Arene Derivatives with Lower Rim Functionalisation and Their Applications. *Coord. Chem. Rev.* **2009**, 253, 893–962, doi:10.1016/j.ccr.2008.06.008.
16. Podyachev, S.N.; Kashapova, N.E.; Syakaev, V.V.; Sudakova, S.N.; Zainullina, R.R.; Gruner, M.; Habicher, W.D.; Barsukova, T.A.; Yang, F.; Konovalov, A.I. Mercury(II) and Silver(I) Receptors Based on Tetrathiacalix[4]Arene Hydrazones. *J. Incl. Phenom. Macrocycl. Chem.* **2014**, 78, 371–380, doi:10.1007/s10847-013-0307-0.
17. Sviben, I.; Galić, N.; Tomišić, V.; Frkanec, L. Extraction and Complexation of Alkali and Alkaline Earth Metal Cations by Lower-Rim Calix[4]Arene Diethylene Glycol Amide Derivatives. *New J. Chem.* **2015**, 39, 6099–6107, doi:10.1039/C5NJ00805K.
18. Aparicio-Aragon, W.B.; Ramos, T.D.; Namor, A.F.D.D. Effect of the Number of Donor Atoms in Calix[4]Arene Derivatives for Selective Complexation and Extraction of Mercury(II) Cations, Thermodynamics and Extraction Studies. *ACES* **2021**, 11, 165–179, doi:10.4236/aces.2021.112011.
19. Francisco, V.; Basílio, N.; Garcia-Rio, L.; R. Leis, J.; F. Maques, E.; Vázquez-Vázquez, C. Novel Catanionic Vesicles from Calixarene and Single-Chain Surfactant. *Chem. Commun.* **2010**, 46, 6551–6553, doi:10.1039/C0CC01806F.
20. Basílio, N.; Garcia-Rio, L.; Martín-Pastor, M. Calixarene-Based Surfactants: Evidence of Structural Reorganization upon Micellization. *Langmuir* **2012**, 28, 2404–2414, doi:10.1021/la204004h.
21. Basílio, N.; Gómez, B.; Garcia-Rio, L.; Francisco, V. Using Calixarenes To Model Polyelectrolyte Surfactant Nucleation Sites. *Chem. Eur. J.* **2013**, 19, 4570–4576, doi:10.1002/chem.201203377.
22. Homden, D.M.; Redshaw, C. The Use of Calixarenes in Metal-Based Catalysis. *Chem. Rev.* **2008**, 108, 5086–5130, doi:10.1021/cr8002196.
23. Sachdeva, G.; Vaya, D.; Srivastava, C.M.; Kumar, A.; Rawat, V.; Singh, M.; Verma, M.; Rawat, P.; Rao, G.K. Calix[n]Arenes and Its Derivatives as Organocatalysts. *Coord. Chem. Rev.* **2022**, 472, 214791, doi:10.1016/j.ccr.2022.214791.
24. Bakirci, H.; Koner, A.L.; Dickman, M.H.; Kortz, U.; Nau, W.M. Dynamically Self-Assembling Metalloenzyme Models Based on Calixarenes. *Angew. Chem. Int. Ed.* **2006**, 45, 7400–7404, doi:10.1002/anie.200602999.
25. Schühle, D.T.; Peters, J.A.; Schatz, J. Metal Binding Calixarenes with Potential Biomimetic and Biomedical Applications. *Coord. Chem. Rev.* **2011**, 255, 2727–2745, doi:10.1016/j.ccr.2011.04.005.
26. Nimse, S.B.; Kim, T. Biological Applications of Functionalized Calixarenes. *Chem. Soc. Rev.* **2013**, 42, 366–386, doi:10.1039/C2CS35233H.
27. Rebilly, J.-N.; Colasson, B.; Bistri, O.; Over, D.; Reinaud, O. Biomimetic Cavity-Based Metal Complexes. *Chem. Soc. Rev.* **2015**, 44, 467–489, doi:10.1039/C4CS00211C.
28. Pan, Y.; Hu, X.; Guo, D. Biomedical Applications of Calixarenes: State of the Art and Perspectives. *Angew. Chem. Int. Ed.* **2021**, 60, 2768–2794, doi:10.1002/anie.201916380.
29. Martins, J.N.; Raimundo, B.; Rioboo, A.; Folgar-Cameán, Y.; Montenegro, J.; Basílio, N. Photoswitchable Calixarene Activators for Controlled Peptide Transport across Lipid Membranes. *J. Am. Chem. Soc.* **2023**, 145, 13126–13133, doi:10.1021/jacs.3c01829.
30. Lazar, A.-N.; Perret, F.; Perez-Lloret, M.; Michaud, M.; Coleman, A.W. Promises of Anionic Calix[n]Arenes in Life Science: State of the Art in 2023. *Eur. J. Med. Chem* **2024**, 264, 115994, doi:10.1016/j.ejmech.2023.115994.
31. Webber, M.J.; Langer, R. Drug Delivery by Supramolecular Design. *Chem. Soc. Rev.* **2017**, 46, 6600–6620, doi:10.1039/C7CS00391A.
32. Gaeta, C.; La Manna, P.; De Rosa, M.; Soriente, A.; Talotta, C.; Neri, P. Supramolecular Catalysis with Self-assembled Capsules and Cages: What Happens in Confined Spaces. *ChemCatChem* **2021**, 13, 1638–1658, doi:10.1002/cctc.202001570.

33. Fan, X.; Guo, X. Development of Calixarene-Based Drug Nanocarriers. *J. Mol. Liq.* **2021**, *325*, 115246, doi:10.1016/j.molliq.2020.115246.
34. Zhang, F.; Sun, Y.; Tian, D.; Shin, W.S.; Kim, J.S.; Li, H. Selective Molecular Recognition on Calixarene-Functionalized 3D Surfaces. *Chem. Commun.* **2016**, *52*, 12685–12693, doi:10.1039/C6CC05876K.
35. Kongor, A.R.; Mehta, V.A.; Modi, K.M.; Panchal, M.K.; Dey, S.A.; Panchal, U.S.; Jain, V.K. Calix-Based Nanoparticles: A Review. *Top. Curr. Chem.* **2016**, *374*, 28, doi:10.1007/s41061-016-0029-z.
36. Wei, A. Calixarene-Encapsulated Nanoparticles: Synthesis, Stabilization, and Self-Assembly. In *Calixarenes and Beyond*; Neri, P., Sessler, J.L., Wang, M.-X., Eds.; Springer International Publishing: Cham, 2016; pp. 921–939 ISBN 978-3-319-31865-3.
37. Danil de Namor, A.F.; Kowalska, D.; Castellano, E.E.; Piro, O.E.; Sueros Velarde, F.J.; Villanueva Salas, J. Lower Rim Calix(4)Arene Ketone Derivatives and Their Interaction with Alkali Metal Cations. Structural and Thermodynamic (Solution and Complexation) Characterisation of the Tetraphenyl Ketone Derivative and Its Sodium Complex. *Phys. Chem. Chem. Phys.* **2001**, *3*, 4010–4021, doi:10.1039/b104355m.
38. Śliwa, W. Calixarene Complexes with Transition Metal Ions. *J. Incl. Phenom. Macrocycl. Chem.* **2005**, *52*, 13–37, doi:10.1007/s10847-005-0083-6.
39. Śliwa, W.; Girek, T. Calixarene Complexes with Metal Ions. *J. Incl. Phenom. Macrocycl. Chem.* **2010**, *66*, 15–41, doi:10.1007/s10847-009-9678-7.
40. Danil de Namor, A.F. Thermodynamics of Calixarene-Ion Interactions. In *Calixarenes 2001*; Asfari, Z., Böhmer, V., Harrowfield, J., Vicens, J., Saadioui, M., Eds.; Springer Netherlands: Dordrecht, 2001; pp. 346–364 ISBN 978-0-306-47522-1.
41. Horvat, G.; Frkanec, L.; Cindro, N.; Tomišić, V. A Comprehensive Study of the Complexation of Alkali Metal Cations by Lower Rim Calix[4]Arene Amide Derivatives. *Phys. Chem. Chem. Phys.* **2017**, *19*, 24316–24329, doi:10.1039/c7cp03920d.
42. Danil de Namor, A.F.; Matsufuji-Yasuda, T.T.; Zegarra-Fernandez, K.; Webb, O.A.; El Gamouz, A. An Enchiridion of Supramolecular Thermodynamics: Calix[N]Arene (N=4,5,6) Tertiary Amide Derivatives and Their Ionic Recognition. *Croat. Chem. Acta* **2013**, *86*, 1–19, doi:10.5562/cca2170.
43. Horvat, G.; Stilinović, V.; Hrenar, T.; Kaitner, B.; Frkanec, L.; Tomišić, V. An Integrated Approach (Thermodynamic, Structural, and Computational) to the Study of Complexation of Alkali-Metal Cations by a Lower-Rim Calix[4]Arene Amide Derivative in Acetonitrile. *Inorg. Chem.* **2012**, *51*, 6264–6278, doi:10.1021/ic300474s.
44. Horvat, G.; Stilinović, V.; Kaitner, B.; Frkanec, L.; Tomišić, V. The Effect of Specific Solvent-Solute Interactions on Complexation of Alkali-Metal Cations by a Lower-Rim Calix[4]Arene Amide Derivative. *Inorg. Chem.* **2013**, *52*, 12702–12712, doi:10.1021/ic4019184.
45. Požar, J.; Nikšić-Franjić, I.; Cvetnić, M.; Leko, K.; Cindro, N.; Pičuljan, K.; Borilović, I.; Frkanec, L.; Tomišić, V. Solvation Effect on Complexation of Alkali Metal Cations by a Calix[4]Arene Ketone Derivative. *J. Phys. Chem. B* **2017**, *121*, 8539–8550, doi:10.1021/acs.jpcc.7b05093.
46. Nomura, E.; Takagaki, M.; Nakaoka, C.; Uchida, M.; Taniguchi, H. Effects on Metal Binding of Circular NH- -OC Intramolecular Hydrogen Bonding in Hydrophilic Pseudocavities of *p*-Tert-Butylcalix[4]Arenes Carrying Amino Acid Moieties. *J. Org. Chem.* **1999**, *64*, 3151–3156, doi:10.1021/jo982251k.
47. Frkanec, L.; Višnjevac, A.; Kojić-Prodić, B.; Žinić, M. Calix[4]Arene Amino Acid Derivatives. Intra- and Intermolecular Hydrogen-Bonded Organisation in Solution and the Solid State. *Chem. Eur. J.* **2000**, *6*, 442–453, doi:10.1002/(SICI)1521-3765(20000204)6:3<442::AID-CHEM442>3.0.CO;2-1.
48. Tomišić, V.; Galić, N.; Bertoša, B.; Frkanec, L.; Simeon, V.; Žinić, M. Hydrogen Bonding and Solvent Effects on Complexation of Alkali Metal Cations by Lower Rim Calix[4]Arene Tetra(O-[N-Acetyl-D-Phenylglycine Methyl Ester]) Derivative. *J. Incl. Phenom. Macrocycl. Chem.* **2005**, *53*, 263–268, doi:10.1007/s10847-005-2024-9.
49. Požar, J.; Cvetnić, M.; Usenik, A.; Cindro, N.; Horvat, G.; Leko, K.; Modrušan, M.; Tomišić, V. The Role of Triazole and Glucose Moieties in Alkali Metal Cation Complexation by Lower-Rim Tertiary-Amide Calix[4]Arene Derivatives. *Molecules* **2022**, *27*, 470, doi:10.3390/molecules27020470.
50. Danil de Namor, A.F.; Chahine, S.; Kowalska, D.; Castellano, E.E.; Piro, O.E. Selective Interaction of Lower Rim Calix[4]Arene Derivatives and Bivalent Cations in Solution. Crystallographic Evidence of the Versatile

- Behavior of Acetonitrile in Lead(II) and Cadmium(II) Complexes. *J. Am. Chem. Soc.* **2002**, *124*, 12824–12836, doi:10.1021/ja020764+.
51. Leko, K.; Bregović, N.; Cvetnić, M.; Cindro, N.; Tranfić Bakić, M.; Požar, J.; Tomišić, V. Complexation of Alkali Metal Cations by a Tertiary Amide Calix[4]Arene Derivative in Strongly Cation Solvating Solvents. *Croat. Chem. Acta* **2017**, *90*, doi:10.5562/cca3200.
 52. Danil de Namor, A.F.; Chahine, S.; Castellano, E.E.; Piro, O.E. Solvent Control on the Selective, Nonselective, and Absent Response of a Partially Substituted Lower Rim Calix(4)Arene Derivative for Soft Metal Cations (Mercury(II) and Silver(I)). Structural and Thermodynamic Studies. *J. Phys. Chem. A* **2005**, *109*, 6743–6751, doi:10.1021/jp0514067.
 53. de Araujo, A.S.; Piro, O.E.; Castellano, E.E.; Danil de Namor, A.F. Combined Crystallographic and Solution Molecular Dynamics Study of Allosteric Effects in Ester and Ketone P-Tert-Butylcalix[4]Arene Derivatives and Their Complexes with Acetonitrile, Cd(II), and Pb(II). *J. Phys. Chem. A* **2008**, *112*, 11885–11894, doi:10.1021/jp804198c.
 54. Danil de Namor, A.F.; De Sueros, N.A.; McKervey, M.A.; Barrett, G.; Neu, F.A.; Schwing-Weill, M.J. The Solution Thermodynamics of Ethyl P-Tert-Butylcalix[4]Arene Tetraacetate and Its Alkali Metal Complexes in Acetonitrile and Methanol. *J. Chem. Soc., Chem. Commun.* **1991**, 1546, doi:10.1039/c39910001546.
 55. Kim, J.S.; Quang, D.T. Calixarene-Derived Fluorescent Probes. *Chem. Rev.* **2007**, *107*, 3780–3799, doi:10.1021/cr068046j.
 56. Mako, T.L.; Racicot, J.M.; Levine, M. Supramolecular Luminescent Sensors. *Chem. Rev.* **2019**, *119*, 322–477, doi:10.1021/acs.chemrev.8b00260.
 57. Tranfić Bakić, M.; Jadreško, D.; Hrenar, T.; Horvat, G.; Požar, J.; Galić, N.; Sokol, V.; Tomaš, R.; Alihodžić, S.; Žinić, M.; et al. Fluorescent Phenanthridine-Based Calix[4]Arene Derivatives: Synthesis and Thermodynamic and Computational Studies of Their Complexation with Alkali-Metal Cations. *RSC Adv.* **2015**, *5*, 23900–23914, doi:10.1039/C5RA01905B.
 58. Galić, N.; Burić, N.; Tomaš, R.; Frkanec, L.; Tomišić, V. Synthesis and Cation Binding Properties of Fluorescent Calix[4]Arene Derivatives Bearing Tryptophan Units at the Lower Rim. *Supramol. Chem.* **2011**, *23*, 389–397, doi:10.1080/10610278.2010.521832.
 59. Bregović, N.; Cindro, N.; Frkanec, L.; Tomišić, V. Complexation of Fluoride Anion and Its Ion Pairs with Alkali Metal Cations by Tetra-Substituted Lower Rim Calix[4]Arene Tryptophan Derivative. *Supramol. Chem.* **2016**, *28*, 608–615, doi:10.1080/10610278.2016.1154147.
 60. Tranfić Bakić, M.; Leko, K.; Cindro, N.; Portada, T.; Hrenar, T.; Frkanec, L.; Horvat, G.; Požar, J.; Tomišić, V. Synthesis of Fluorescent Diphenylanthracene-Based Calix[4]Arene Derivatives and Their Complexation with Alkali Metal Cations. *Croat. Chem. Acta* **2017**, *90*, 711–725, doi:10.5562/cca3308.
 61. Leko, K.; Usenik, A.; Cindro, N.; Modrušan, M.; Požar, J.; Horvat, G.; Stilinović, V.; Hrenar, T.; Tomišić, V. Enhancing the Cation-Binding Ability of Fluorescent Calixarene Derivatives: Structural, Thermodynamic, and Computational Studies. *ACS Omega* **2023**, *8*, 43074–43087, doi:10.1021/acsomega.3c06509.
 62. Modrušan, M.; Cindro, N.; Usenik, A.; Leko, K.; Glazer, L.; Tomaš, R.; Horvat, G.; Požar, J.; Tomišić, V. Complexation of Alkali and Alkaline Earth Metal Cations by Fluorescent Glycoconjugated Calix[4]Arene Derivative: Thermodynamic and Computational Studies. *Croat. Chem. Acta* **2024**, *97*, doi:10.5562/cca4115.
 63. Gans, P.; Sabatini, A.; Vacca, A. Investigation of Equilibria in Solution. Determination of Equilibrium Constants with the HYPERQUAD Suite of Programs. *Talanta* **1996**, *43*, 1739–1753, doi:10.1016/0039-9140(96)01958-3.
 64. Briggner, L.-E.; Wadsö, I. Test and Calibration Processes for Microcalorimeters, with Special Reference to Heat Conduction Instruments Used with Aqueous Systems. *J. Biochem. Biophys. Methods* **1991**, *22*, 101–118, doi:10.1016/0165-022X(91)90023-P.
 65. Gans, P.; Sabatini, A.; Vacca, A. Simultaneous Calculation of Equilibrium Constants and Standard Formation Enthalpies from Calorimetric Data for Systems with Multiple Equilibria in Solution. *J. Solution Chem.* **2008**, *37*, 467–476, doi:10.1007/s10953-008-9246-6.
 66. Berendsen, H.J.C.; Van Der Spoel, D.; Van Drunen, R. GROMACS: A Message-Passing Parallel Molecular Dynamics Implementation. *Comput. Phys. Commun.* **1995**, *91*, 43–56, doi:10.1016/0010-4655(95)00042-E.

67. Lindahl, E.; Hess, B.; Van Der Spoel, D. GROMACS 3.0: A Package for Molecular Simulation and Trajectory Analysis. *J. Mol. Model.* **2001**, *7*, 306–317, doi:10.1007/s008940100045.
68. Van Der Spoel, D.; Lindahl, E.; Hess, B.; Groenhof, G.; Mark, A.E.; Berendsen, H.J.C. GROMACS: Fast, Flexible, and Free. *J. Comput. Chem.* **2005**, *26*, 1701–1718, doi:10.1002/jcc.20291.
69. Hess, B.; Kutzner, C.; Van Der Spoel, D.; Lindahl, E. GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J. Chem. Theory Comput.* **2008**, *4*, 435–447, doi:10.1021/ct700301q.
70. Pronk, S.; Páll, S.; Schulz, R.; Larsson, P.; Bjelkmar, P.; Apostolov, R.; Shirts, M.R.; Smith, J.C.; Kasson, P.M.; Van Der Spoel, D.; et al. GROMACS 4.5: A High-Throughput and Highly Parallel Open Source Molecular Simulation Toolkit. *Bioinformatics* **2013**, *29*, 845–854, doi:10.1093/bioinformatics/btt055.
71. Abraham, M.J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J.C.; Hess, B.; Lindahl, E. GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers. *SoftwareX* **2015**, *1–2*, 19–25, doi:10.1016/j.softx.2015.06.001.
72. Best, R.B.; Zhu, X.; Shim, J.; Lopes, P.E.M.; Mittal, J.; Feig, M.; MacKerell, A.D. Optimization of the Additive CHARMM All-Atom Protein Force Field Targeting Improved Sampling of the Backbone ϕ , ψ and Side-Chain χ_1 and χ_2 Dihedral Angles. *J. Chem. Theory Comput.* **2012**, *8*, 3257–3273, doi:10.1021/ct300400x.
73. Allouche, A.-R. Gabedit—A Graphical User Interface for Computational Chemistry Softwares. *J. Comput. Chem.* **2011**, *32*, 174–182, doi:10.1002/jcc.21600.
74. Vanommeslaeghe, K.; MacKerell, A.D. Automation of the CHARMM General Force Field (CGenFF) I: Bond Perception and Atom Typing. *J. Chem. Inf. Model.* **2012**, *52*, 3144–3154, doi:10.1021/ci300363c.
75. Yu, W.; He, X.; Vanommeslaeghe, K.; MacKerell, A.D. Extension of the CHARMM General Force Field to Sulfonyl-containing Compounds and Its Utility in Biomolecular Simulations. *J. Comput. Chem.* **2012**, *33*, 2451–2468, doi:10.1002/jcc.23067.
76. Goga, N.; Rzepiela, A.J.; De Vries, A.H.; Marrink, S.J.; Berendsen, H.J.C. Efficient Algorithms for Langevin and DPD Dynamics. *J. Chem. Theory Comput.* **2012**, *8*, 3637–3649, doi:10.1021/ct3000876.
77. Darden, T.; York, D.; Pedersen, L. Particle Mesh Ewald: An $N \cdot \log(N)$ Method for Ewald Sums in Large Systems. *J. Chem. Phys.* **1993**, *98*, 10089–10092, doi:10.1063/1.464397.
78. Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual Molecular Dynamics. *J. Mol. Graph. Model.* **1996**, *14*, 33–38, doi:10.1016/0263-7855(96)00018-5.
79. Lee, C.; Yang, W.; Parr, R.G. Development of the Colle-Salvetti Correlation-Energy Formula into a Functional of the Electron Density. *Phys. Rev. B* **1988**, *37*, 785–789, doi:10.1103/PhysRevB.37.785.
80. Peverati, R.; Truhlar, D.G. Screened-Exchange Density Functionals with Broad Accuracy for Chemistry and Solid-State Physics. *Phys. Chem. Chem. Phys.* **2012**, *14*, 16187–16191, doi:10.1039/C2CP42576A.
81. Grimme, S.; Ehrlich, S.; Goerigk, L. Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *J. Comput. Chem.* **2011**, *32*, 1456–1465, doi:10.1002/jcc.21759.
82. Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305, doi:10.1039/B508541A.
83. Weigend, F. Accurate Coulomb-Fitting Basis Sets for H to Rn. *Phys. Chem. Chem. Phys.* **2006**, *8*, 1057–1065, doi:10.1039/B515623H.
84. Hrenar, T.; Primožič, I.; Fijan, D.; Majerić Elenkov, M. Conformational Analysis of Spiro-Epoxides by Principal Component Analysis of Molecular Dynamics Trajectories. *Phys. Chem. Chem. Phys.* **2017**, *19*, 31706–31713, doi:10.1039/c7cp05600a.
85. Primožič, I.; Hrenar, T.; Baumann, K.; Krišto, L.; Križič, I.; Tomić, S. Mechanochemical and Conformational Study of N-Heterocyclic Carbonyl-Oxime Transformations. *Croat. Chem. Acta* **2014**, *87*, 153–160, doi:10.5562/cca2476.
86. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa,

- M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, Gaussian 16, Revision A.03, Gaussian, Inc., Wallingford CT, 2016.
87. Jović, O.; Smolić, T.; Primožič, I.; Hrenar, T. Spectroscopic and Chemometric Analysis of Binary and Ternary Edible Oil Mixtures: Qualitative and Quantitative Study. *Anal. Chem.* **2016**, *88*, 4516–4524, doi:10.1021/acs.analchem.6b00505.
 88. Ramić, A.; Skočibušić, M.; Odžak, R.; Čipak Gašparović, A.; Milković, L.; Mikelić, A.; Sović, K.; Primožič, I.; Hrenar, T. Antimicrobial Activity of Quasi-Enantiomeric Cinchona Alkaloid Derivatives and Prediction Model Developed by Machine Learning. *Antibiotics* **2021**, *10*, 659, doi:10.3390/antibiotics10060659.
 89. Lakowicz, J.R. *Principles of Fluorescence Spectroscopy*; 3rd ed.; Springer: New York, 2006; ISBN 978-0-387-31278-1.
 90. Valeur, B.; Leray, I. Ion-Responsive Supramolecular Fluorescent Systems Based on Multichromophoric Calixarenes: A Review. *Inorganica Chim. Acta* **2007**, *360*, 765–774, doi:10.1016/j.ica.2006.07.027.
 91. Požar, J.; Preočanin, T.; Frkanec, L.; Tomišić, V. Thermodynamics of Complexation of Alkali Metal Cations by a Lower-Rim Calix[4]Arene Amino Acid Derivative. *J. Solution Chem.* **2010**, *39*, 835–848, doi:10.1007/s10953-010-9550-9.
 92. Marcus, Y. *Ion Properties*; Dekker: New York Basel, 1997; ISBN 978-0-8247-0011-9.
 93. Haynes, W.M. *CRC Handbook of Chemistry and Physics*; 95th ed.; CRC Press, Hoboken, 2014;

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.