

Article

Not peer-reviewed version

Characterization of Antibiotic Resistance and Prevalence of Diarrheagenic *E. coli* Strains Isolated from Stool Samples in a Hospital Setting in Mali

Aminata MAIGA , [Lassina DOUMBIA](#) ^{*} , N'Faly Tounkara , [Tidiani Sinayoko](#) , Rabiataou Sanogo , Ibrahim Keita , Ami Diakite , Moussa Diawara , Oumar Agaly Dicko , Famakan Dembele , [Mohamed Ag Baraika](#) , [Yacouba Cissoko](#) , Drissa Kone , [Aldiouma Guindo](#) , Ibrahim Izétiégouma Maïga , [Ousmane Koita](#)

Posted Date: 14 October 2025

doi: 10.20944/preprints202510.1105.v1

Keywords: *E. coli*; DEC; gene; Multidrug resistance; Integron; Mali



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Characterization of Antibiotic Resistance and Prevalence of Diarrheagenic *E. coli* Strains Isolated from Stool Samples in a Hospital Setting in Mali

Aminata Maiga ¹, Lassina Doumbia ^{2,*}, N’faly Touunkara ³, Tidiani Sinayoko ², Rabiathou Sanogo ⁴, Ibrahim Keita ², Ami Diakite ², Moussa Diawara ³, Oumar Agaly Dicko ³, Famakan Dembele ³, Mohamed Ag Baraika ⁵, Yacouba Cissoko ⁵, Drissa Kone ³, Aldiouma Guindo ⁵, Ibrahim Izétiégouma Maïga ¹ and Ousmane Koita ⁵

- ¹ Faculté de Médecine et Odontostomatologie, USTTB, Laboratoire de Biologie Médicale et Hygiène Hospitalière CHU du Point G
- ² Faculté des Sciences et Techniques, USTTB, Laboratoire de Biologie Moléculaire Appliquée
- ³ Laboratoire de Biologie Médicale et Hygiène Hospitalière CHU du Point G
- ⁴ Institut National de Santé Publique
- ⁵ Université des Sciences, des Techniques et des Technologies de Bamako (USTTB)
- * Correspondence: dolslassina@yahoo.fr

Abstract

Introduction. Antibiotic resistance is a major global health challenge that disproportionately affects low-resource countries, particularly those in West Africa. *E. coli*, a major pathogen in childhood diarrhea, is both a prominent infectious agent and a reservoir of resistance genes, including resistance to last-resort antibiotics such as carbapenems. **Methodology.** The study focused on 98 clinical *E. coli* isolates collected from stool samples of patients in a hospital setting in Bamako. The analyses included screening for DEC-specific virulence genes, detection of resistance genes across various classes of antibiotics (e.g., beta-lactams, carbapenems, fluoroquinolones), and identification of class I, II, and III integrons. The *bla_{NDM}* gene was sequenced to identify mutations associated with carbapenem resistance. **Results.** Among the isolates, 85.7% carried at least one virulence gene. Of these, half involved co-infections, commonly combining EPEC, EAEC, and ETEC strains. Regarding antibiotic resistance, 94.9% of isolates harbored at least one resistance gene, and 50% were multidrug-resistant. The most frequently detected genes were *bla_{TEM}*, *qnrS1*, and *aphA3*. Class II integrons were significantly associated with multidrug resistance ($p = 0.01$). Sequencing of the *bla_{NDM}* gene revealed point mutations likely to affect protein function, suggesting an evolution toward increased resistance to carbapenems. **Conclusion.** The high prevalence of multidrug-resistant diarrheagenic *E. coli* strains in this study highlights the local antibiotic pressure and the serious health threat it represents. This study shows that the new β -lactamase *bla_{NDM}* gene has disseminated in the hospital environment of Bamako. It should be noted that this will become a major challenge for clinicians.

Keywords: *E. coli*; DEC; gene; multidrug resistance; integron; Mali

Introduction

Bacterial resistance to antibiotics is one of the major threats to global public health, with several million deaths estimated each year. Sub-Saharan Africa, particularly its western region, appears to be especially affected by this phenomenon [1]. However, even industrialized countries experience significant health and economic consequences [2,3].

Among the primary causes of this resistance are the excessive and inappropriate use of antibiotics in both human and animal health, poor management stewardship, and the intrinsic resistance specific to certain bacterial species [4].

In 2019, *Escherichia coli* (*E. coli*) was one of the six pathogens responsible for the highest number of deaths related to antibiotic resistance, ranking first [1]. As a member of the *Enterobacteriaceae* family, *E. coli* is a commensal bacterium in the mammalian digestive tract. While most strains are harmless, some are pathogenic and can cause severe intestinal or extra-intestinal infections [5]. The World Health Organization (WHO) ranks *E. coli*, alongside other *Enterobacteriaceae*, among the priority pathogens due to the growing threat they pose to human health [6].

E. coli is a common cause of diarrhea in children worldwide. In Mali, this bacterium is implicated in more than 30% of diarrhea cases in children under five years of age. These strains show increased resistance to several classes of antibiotics, notably beta-lactams and quinolones, although they remain relatively sensitive to imipenem, a carbapenem [7]. Moreover, multiple multidrug-resistant strains have been isolated from children suffering from acute diarrhea [8].

Beyond its multidrug-resistant profile, *E. coli* also plays a key role as a reservoir of resistance genes, due to its ability to acquire such genes from other bacteria and transmit them through horizontal gene transfer mechanisms [9,10]. This resistance is more pronounced in clinical strains compared to environmental strains [11].

In this context, this study aims to characterize the molecular profile of virulence and antibiotic resistance genes in clinical *E. coli* strains, with a particular focus on carbapenems, a class that remains understudied in Mali.

Materials and Method

The study focused on 98 clinical *E. coli* isolates collected from stool samples of patients between 2020 and 2021 at the laboratory of the CHU of Point G, Bamako, Mali, and stored at -80 °C. The isolates were subsequently analyzed at the Laboratory of Applied Molecular Biology (LBMA) using standard PCR.

Molecular analyses included screening for DEC-specific virulence genes, detection of resistance genes across various antibiotic classes, and identification of class 1, 2, and 3 integrons (Table 1).

Frozen isolates were thawed and cultured on Mueller-Hinton (MH) agar for DNA extraction. Then, a full loop of fresh pure *E. coli* culture was stored at -80 °C in 1.5% (v/v) glycerol in enumeration broth for further analysis.

DNA Extraction

Reference strains were provided by the National Institute of Public Health (NIPH) of Mali. DNA was extracted from *E. coli* isolates and the reference strain using a full loop of colonies in 100 µL of ultrapure water following the Salting-out method. The purified DNA was eluted in 70 µL of TE buffer and stored at -20 °C for further amplification.

Identification of Virulence Genes

Strains harboring at least one DEC-associated virulence gene were classified as Diarrheagenic *E. coli* (DEC). Several genes characteristic of Diarrheagenic *E. coli* (DEC) were targeted (see Table 1):

- *bfpA* and *eae* for typical enteropathogenic *E. coli* (EPEC),
- *agg* and *aaic* for enteroaggregative *E. coli* (EAEC),
- *lt* and *st* for enterotoxigenic *E. coli* (ETEC).

Three microliters of DNA from the reference strains provided by the NIPH, the negative control (sterile ultrapure water), and the *E. coli* isolates were used for multiplex PCR with specific primers (Table 1), as described by Vilchez S et al., 2009 [8].

Additionally, to confirm the multiplex PCR results, DNA extracted from freshly cultured colonies on MH agar was subjected to single PCR. Each specific primer was tested independently in single PCR to confirm suspected DEC isolates identified by multiplex PCR.

Identification of Resistance and Integron Genes

Specific resistance genes to different antibiotic families were evaluated: ESBL genes (Extended-Spectrum Beta-Lactamases) for beta-lactams; *bla*NDM and *bla*IMP for carbapenems; *tetA* for tetracyclines; *qnrS1* for quinolones; *catA1* for chloramphenicols, and *aphA3* for aminoglycosides.

PCR was used to detect the presence of antibiotic resistance genes, as well as class 1, 2, and 3 integrons genes, and to assess their distribution among the isolates. Integrons play a major role in the acquisition, expression, and dissemination of antibiotic resistance, particularly resistance integrons.

All amplification was carried out using the PTC 200 thermocycler (MJ Research, USA) with specific primer sequences as listed in Table 1. The reaction total volume was 25 µl containing 3 µL of DNA, 1x Buffer (Mg²⁺ free), 3 mM MgCl₂, 0.4 mM deoxynucleotide triphosphates (dNTPs) (Invitrogen, USA), 0.4 µM of each primer, and 0.025 U of Taq polymerase (Invitrogen, USA).

The PCR program for all reactions was as follows: initial denaturation at 94 °C for 5 minutes, followed by 35 cycles of 94 °C for 30 seconds (denaturation), annealing at 44 to 60 °C for 45 seconds (variable according to the specific primers), and 72 °C for 1 minute (extension). The annealing temperatures were: OXA (44 °C), QnrS1 and IMP (50 °C), NDM (52 °C), SHV (60 °C), TEM (55 °C), catA (59 °C), tetA (55 °C), CTX-M (57.2 °C), aphA-3 (51.9 °C), Int1 (59 °C), Int2 (55 °C), and Int3 (57 °C), respectively. A final elongation step was performed at 72 °C for 10 minutes. All PCR products were visualized on a 1.5% agarose gel stained with ethidium bromide.

Table 1. List of primers Used.

DEC type	Target gene	Primer	Primer sequence 5' - 3'	PCR product Size bp	Reference
ETEC	<i>eltB</i>	LT-F	TCTCTATGTGCATACGGAGC	322	[12]
		LT-R	CCATACTGATTGCCGCAAT		
	<i>estA</i>	ST-F	GTCAAACCAGTA(G/A)GGTCTTCAAAA	147	[12]
		ST-R	CCCGGTACA(G/A)GGAGGATTACAACA		
EHEC	<i>vt1</i>	VT1-F	GAAGAGTCCGTGGGATTAC	130	[12]
		VT1-R	AGCGATGCAGCTATTAATAA		
	<i>vt2</i>	VT2-F	ACCGTTTTTCAGATTTT(G/A)CACATA	298	[12]
		VT2-R	TACACAGGAGCAGTTTCAGACAGT		
EPEC	<i>EaeA</i>	Eae-F	CACACGAATAAACTGACTAAAATG	376	[12]
		Eae-R	AAAAACGCTGACCCGCACCTAAAT		
	<i>bfpA</i>	bfpA-F	TTCTTGGTGCTTGC GTGTCTTTT	367	[12]
		bfpA-R	TTTTGTTTGTGTATCTTTGTAA		
EIEC	<i>Ial</i>	SHIG-F	CTGGTAGGTATGGTGAGG	320	[12]
		SHIG-R	CCAGGCCAACAATTATTTC		
EAEC	<i>pCVD432</i>	EA-F	CTGGCGAAAGACTGTATCAT	630	[12]
		EA-R	AAATGTATAGAAATCCGCTGTT		
Resistance genes	<i>bla</i> _{OXA}	OXA 1-F	ATGAAAAACACAATACATATC	890	[13]
		OXA 1-R	AATTTAGTGTGTTTAGAATGG		
	<i>bla</i> _{SHV}	SHV-F	TTATCTCCCTGTTAGCCACC	800	[13,14]
		SHV-R	GATTTGCTGATTTGCTCGG		
	<i>bla</i> _{TEM}	TEM-F	ATAAAATCCTGAAGACGAAA	850	[13]
		TEM-R	GACAGTTACCAATGCTTAATC		
	<i>bla</i> _{CTX-M-3/15/22}	CTX-M-F	GTTACAATGTGTGAGAAGCAG	593	[15]
		CTX-M-R	CCGTTTCCGCTATTACAAAC		
	<i>catA1</i>	catA 1-F	CGCCTGATGAATGCTCATCCG	450	[14]
		catA 1-R	CCTGCCACTCATCGCAGTAC		
	<i>tetA</i>	tetA-F	GTAATTCTGAGCACTGTCTGC	956	[14]
		tetA-R	CTGCCTGGACAACATTGCTT		
	<i>aphA-3</i>	aphA-3-F	GGGACCACCTATGATGTGGAACG	600	[14]
		aphA-3-R	CAGGCTTGATCCCCAGTAAGTC		[16]
	<i>bla</i> _{NDM}	NDM-1 F	GGTTTGGCGATCTGGTTTTC	621	
		NDM-1 R	CGGAATGGCTCATCACGATC		
	<i>bla</i> _{IMP}	IMP F	CACTTGGTTTGTGGAACGTG	192	This study

					GenBank: CP090265.1
		IMP_R	CAATAGTTAACCCCGCCAAA		
	<i>QnrS1</i>	QnrS1_F	ACGCACGGAAGTCTATACCG	154	This study GenBank: CP090265.1
		QnrS1_R	ACGACATTCGTCAACTGCAA		
Integron	<i>intI1</i>	Int1_F	ACATGTGATGGCGACGCACGA	580	[17]
		Int1_R	ATTTCTGTCCTGGCTGGCGA		
	<i>intI2</i>	Int2_F	CACGGATATGCGACAAAAAGGT	806	[17]
		Int2_R	GTAGCAAACGACTGACGAAATG		
	<i>intI3</i>	Int3_F	AACTCTTGCACCGTTCGGAT	542	This study GenBank: CP047278.1
		Int3_R	CAGGAGGTTTCAGACGTTGCT		

Sequencing

Among isolates carrying the *bla*_{NDM} gene, one isolate was sequenced by the Sanger to identify mutations associated with carbapenem resistance.

The amplified product was purified using Exonuclease I and Alkaline Phosphatase enzymes, followed by thermal cycling and ethanol precipitation. Sequencing was performed on the CEQ™ 8000 DNA Analyzer (Beckman Coulter). The resulting sequence was compared to database entries using the NCBI BLAST search tool. Mutations were analyzed using Geneious Prime 2023.0 software.

Phylogenetic analysis

We retrieved the most closely related sequences from GenBank as of June 3rd, 2024. The sequence dataset was aligned using BioEdit, version 7.7.1 (5/10/2021). Based on the resulting alignments, we performed a maximum likelihood (ML) phylogenetic reconstruction using MEGA version 7.0.26, applying the Tamura-Nei model and assessing branch support with 1,000 bootstrap replicates.

Data Analysis

Data were analyzed using STATA software version 14.0. The Chi-square test or Fisher’s exact test was used, as appropriate, to determine the statistical significance of the data. A *p-value* less than 0.05 was considered statistically significant.

Beta-lactam resistance was defined as the presence of at least one beta-lactam resistance gene. Multidrug-resistant (MDR) was defined as the presence of at least one resistance gene from three different classes of antibiotics. Figures were generated using Excel and the online platform Flourish (<https://flourish.studio/>), employing chord diagrams to illustrate associations between variables in cases of co-infections, multi-resistance, and the presence of different integrons.

Results

This study focused on the molecular characterization of virulence and antibiotic resistance genes profiles of *E. coli* strains isolated in a clinical setting. Of the 98 isolate, 84 (approximately 86%) were identified as DEC, based on the detection of at least one specific virulence gene (Figure 1).




Figure 1. Prevalence of *E. coli* strains based on the presence of virulence genes.

Among the 84 DEC strains identified, 50% were mono-infections, predominantly EPEC (21.4%), followed by EAEC (20.2%) and ETEC (8.3%). The remaining 50% involved mixed infections, with the EAEC–EPEC combination being the most common (27.4%), followed by EPEC–ETEC (9.5%), EAEC–ETEC (3.6%), and the triple combination EAEC–EPEC–ETEC (9.5%) (Figure 2).




Figure 2. Distribution of mixed infections among DEC pathotypes based on detected virulence genes.

Detection of resistance genes

A total of 93 out of 98 *E. coli* isolates (94.9%) carried at least one antibiotic resistance gene, and 50% (49/98) were identified as multidrug-resistant (MDR).

Beta-lactam resistance genes were detected in 77.6% (76/98) of isolates, with *bla*_{TEM} being the most frequently identified gene. Regarding the overall prevalence of resistance genes, *qnrS1* was the most common (66.3%), followed by *bla*_{TEM} (49%), *aphA3* (45.9%), *bla*_{CTX-M} (35.7%), *bla*_{IMP} (23.5%), *tetA* (22.5%), *bla*_{NDM} (16.3%), and *catA1* (10.2%). Carbapenem resistance genes were detected in 38.8% of isolates (Table 2).

Multidrug resistance associated with the *bla*_{NDM} resistance gene was observed in 4 isolates, of which 3 also carried an integron gene. Additionally, 11 isolates harbored the *bla*_{TEM}-*bla*_{CTX-M}-*qnrS1* gene combination, and 8 of them also carried an integron gene. Finally, 7 isolates carried the *bla*_{Oxa}-*bla*_{CTX-M}-*qnrS1* gene combination, all of which were positive for integron gene.

Table 2. Distribution of resistance genes according antibiotic families among *E. coli* isolates.

Antibiotic families	Gene	Present n (%)	Absent n (%)
Overall, N = 98		93 (94.9)	5 (5.1)
Beta-lactams	Overall	76 (77.6)	22 (22.4)
	<i>bla</i> _{TEM}	48 (49.0)	50 (51.0)
	<i>bla</i> _{CTX-M}	35 (35.7)	63 (64.3)
	<i>bla</i> _{SHV}	25 (25.5)	73 (74.5)
	<i>bla</i> _{OXA}	16 (16.3)	82 (83.7)
Carbapenems	Overall	38 (38.8)	60 (61.2)
	<i>bla</i> _{IMP}	23 (23.5)	75 (76.5)
	<i>bla</i> _{NDM}	16 (16.3)	82 (83.7)
Quinolone	<i>qnrS1</i>	65 (66.3)	33 (33.7)
Chloramphenicol	<i>catA1</i>	10 (10.2)	88 (89.8)
Tetracycline	<i>tetA</i>	22 (22.5)	76 (77.5)
Aminoglycoside	<i>aphA3</i>	45 (45.9)	53 (54.1)
Integron	Overall	37 (37.8)	61 (62.2)
	<i>Int 1</i>	35 (35.7)	63 (64.3)
	<i>Int 2</i>	25 (25.5)	73 (74.5)
	<i>Int 3</i>	31 (31.6)	67 (68.4)

Co-detection of resistance genes was observed between resistance genes across most classes of antibiotics investigated. However, the most frequent co-detections were observed between beta-lactams and quinolones (53 isolates), followed by beta-lactams and aminoglycosides (37 isolates), beta-lactams and carbapenems (33 isolates), aminoglycosides and quinolones (31 isolates), quinolones and carbapenems (26 isolates), and beta-lactams and tetracyclines (21 isolates) (Figure 3). These co-detections contribute to the high risk of multidrug resistance, significantly reducing therapeutic options.




Figure 3. Distribution of resistance genes across different classes of antibiotics in *E. coli* strains.

Integron Distribution and Resistance Genes

Integron genes were detected among the *E. coli* isolates as follows: *Int1* (35.7%, 35/98), *Int3* (31.6%, 31/98), and *Int2* (25.5%, 25/98) (Table 3). A statistically significant association was observed between class 2 integrons and isolates harboring more than three resistance genes ($p = 0.011$), suggesting a strong link between class 2 integrons and multidrug resistance. Although class 1 and

class 3 integrons were also found among isolates with more than three resistance genes (60% and 61.3%, respectively), their associations were not statistically significant ($p > 0.05$) (Table 3).

Table 3. Distribution of integron genes based on the number of co-detection resistance genes.

Number of resistance genes	Integron 1, (n=34)		Integron 2, (n=25)		Integron 3, (n=30)	
	n (%)	p-value	n (%)	p-value	n (%)	p-value
1	4 (11,4)	0,962	1 (4)	0,185	3 (9,7)	0,791
2	9 (25,7)	0,214	6 (24)	0,236	8 (25,8)	0,262
3 or more	21 (60)	0,14	18 (72)	0,011	19 (61,39)	0,128

Analysis of blaNDM sequencing results

Sequencing of the PCR-amplified blaNDM product showed high similarity to blaNDM-5. The encoded protein was characterized by several amino acid substitutions, including Val→Leu at position 88 and Met→Leu at position 154. It differed from previously described enzymes by additional substitutions at positions 132 (Leu→Met) and 166 (Asp→Hist), which may alter the structure or function of the protein (Table 4). These novel mutations could contribute to reduced susceptibility of *E. coli* strains to expanded-spectrum cephalosporins and carbapenems.

Table 4. Analysis of mutations in blaNDM sequencing results.

Position	Mutation Type	Mutation consequence
88	G → T substitution	Valine - Leucine change
98	C → A substitution	Silent mutation
132	C → A substitution	Leucine - Methionine change
154	A → C substitution	Methionine-Leucine change
166	A → C substitution	Asparagine - Histidine change

➤ **Phylogenetic Diversity of NDM-Producing *Escherichia coli* Strains Including a Malian Isolate**

The single sequence from this study is closely related to the sequence from China.




Figure 4. Phylogeny of blaNDM sequence from this study.

Maximum-likelihood phylogeny of blaNDM sequence from this study reconstructed with MEGA version 7.0.26. Phylogeny was inferred by using the Maximum Likelihood method based on

the Tamura-Nei model, and branch support was evaluated with bootstrap approximation using 1,000 replicates. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. We used six sequences from BLAST in GenBank, along with the single sequence from this study. The sequences from this study is shown in red.

Discussion

Enteropathogenic *E. coli* (EPEC) strains play a central role in the occurrence of diarrhea, especially in children [18,19]. In this study, 85.7% of the isolates were identified as EPEC, representing a higher prevalence than that reported in other African countries: 61.1% in Niger, 55.9% in Ethiopia [20,21], and 45% in Burkina Faso [22]. Conversely, lower rates have been observed in Colombia (17.9%) [18], Kenya (22.46%) [19], and India (17.4%) [23]. Such variability may be attributed to differences in the target population, sanitation conditions, and detection methods.

In this study, EPEC strains were the most frequently detected, followed by EAEC and ETEC. This trend is consistent with findings from Kenya [19], South Korea [21], and India [23], although the latter did not report any ETEC strains. Other studies, however, have reported the opposite pattern, with a predominance of EAEC followed by EPEC [24–26]. Although the overall prevalence of EPEC is generally low, these strains are highly contagious in children and can lead to more severe forms of diarrhea [27]. We did not differentiate between typical and atypical EPEC strains, but previous research suggests that typical EPEC strains are more commonly associated with diarrheal illness in developing countries [28].

Additionally, 50% of DEC strains in this study exhibited mixed infections involving two or three virulence genes. These combinations may enhance the pathogenic potential of the strains and exacerbate the severity of infection [29,30]. Similar associations have been reported in sub-Saharan Africa, although at lower frequencies [20,22,26]. Regarding antibiotic resistance, 50% of the DEC strains in this study were multidrug-resistant (MDR), defined as the presence of at least one resistance gene in three different antibiotic classes. By comparison, MDR rate were 63.2% in Ethiopia [21], 95.3% in Nigeria [31], and 42.07% in Iran [32]. In Mali, a study also reported high levels of beta-lactam resistance, with 13 different variants identified [7].

The most frequently detected resistance genes in this study were *bla*_{TEM} (49%), *bla*_{CTX-M} (35.7%), and *bla*_{SHV} (25.5%). These results align with those obtained by Guindo et al. in Bamako [33] and are consistent with findings from other studies. For instance, in Iran, the prevalence of these genes was 93.2% for *bla*_{TEM}, 20.5% for *bla*_{CTX-M}, and 2.3% for *bla*_{SHV} [34]. Co-detection of resistance genes across multiple antibiotic classes was also observed, particularly between beta-lactams and quinolones, and between beta-lactams and aminoglycosides. A similar, though less frequent, association was reported in Iran (11.4%) [34]. The coexistence of ESBL genes and fluoroquinolone resistance has also been documented in other studies [35]. We also found isolates carrying both the *bla*_{NDM} gene and integron genes.

The high prevalence of multidrug resistant (MDR) *E. coli* observed in this study may be attributed to the strong selection pressure exerted by excessive and inappropriate antibiotic use in Mali. Historical data support this hypothesis; for instance, a 2002 study conducted in community health centers (CSCOM) reported an antibiotic prescription rate of 61.6%, with a substantial proportion classified as inappropriate [36]. Such prescribing practices not only promote resistance but also reduce the effectiveness of commonly used treatment over time. Beta-lactams remain the most frequently prescribed class, followed by aminoglycosides [37]. This pattern of antibiotic use is concerning, as it may contribute to the continued selection of resistant strains, particularly in environments with limited diagnostic capacity and antimicrobial stewardship programs.

We also observed a high prevalence of class 1 integrons, followed by class 3 and class 2 integrons. These genetic elements facilitate the acquisition and dissemination of resistance genes. A similar integron distribution profile was reported by Guindo et al. in Mali [7]. In contrast, a study conducted in Iran found that class 2 integrons (76.8%) were more strongly associated with multidrug resistance [32]. In our study, class 2 integrons were significantly associated with the presence of more than three resistance genes. For class 1 and class 3 integrons, associations were observed but did not reach statistical significance. This contrasts with the findings of Singh et al., who reported a significant association only for class 1 integrons [38].

Finally, the detection of the *bla_{NDM}* gene, which encodes carbapenemase production, underscores the emergence of highly resistant strains. To our knowledge, this is the first report of NDM β -lactamase gene with mutations in Mali, highlighting its spread within hospital settings in Bamako. These findings emphasize the urgent need for enhanced surveillance and molecular studies to mitigate the clinical impact of this emerging resistance threat. This gene was first identified in 2008 [39], and the sequence obtained in our study reveals two characteristic mutations at positions 88 (Val→Leu) and 154 (Met→Leu) [40]. The presence of NDM-5 in clinical isolates constitutes a major public health concern, given the limited therapeutic options available for treating infections caused by such strains.

One limitation of this study is the low number of sequenced samples due to the limited resources and its retrospective design. However, a key strength is the large number of resistance genes investigated using a molecular approach. Finally, additional studies using high-throughput sequencing (NGS) are needed to better characterize the genetic diversity of *E. coli* strains and to deepen our understanding of the molecular mechanisms underlying their resistance.

Conclusions

This study highlights the prevalence of diarrheagenic *E. coli* (DEC) strains, associated with the widespread distribution of antibiotic resistance genes. These findings illustrate the significant threat that multidrug-resistant DEC strains pose to public health.

References

1. Murray, C.J.L., et al. (2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*, 399(10325), 629-655.
2. Fair, R.J., & Tor, Y. (2014). Antibiotics and Bacterial Resistance in the 21st Century. *SAGE Open*.
3. World Health Organization. (2024). Antimicrobial resistance.
4. Marston, H.D., et al. (2016). Antimicrobial Resistance. *JAMA*, 316(11), 1193.
5. Nataro, J.P., & Kaper, J.B. (1998). Diarrheagenic *Escherichia coli*. *Clinical Microbiology Reviews*, 11(1), 142-201.
6. World Health Organization. (2017). WHO names 12 bacteria that pose the greatest threat to human health. *The Guardian*.
7. Guindo, I., et al. (2022). Pathogenicity Factors and Antibiotic Resistance of *Escherichia coli* Strains Isolated from Diarrheic Children Aged 0–59 Months in Community Settings in Bamako.
8. Diarra, B., et al. (2024). High frequency of antimicrobial resistance in *Salmonella* and *Escherichia coli* causing diarrheal diseases at the Yirimadio community health facility, Mali. *BMC Microbiology*, 24(1), 35.
9. Poirel, L., et al. (2018). Antimicrobial Resistance in *Escherichia coli*. *Microbiology Spectrum*, 6(4), 6.4.14.
10. Rodrigo, L. (2020). *E. Coli Infections: Importance of Early Diagnosis and Efficient Treatment*. BoD – Books on Demand, 236.
11. World Health Organization. (2019). Exploring the antimicrobial resistance profiles of WHO critical priority list bacterial strains. *BMC Microbiology*.
12. Vilchez, S., et al., Prevalence of diarrhoeagenic *Escherichia coli* in children from León, Nicaragua. *Journal of Medical Microbiology*, 2009. 58(Pt 5): p. 630-637.

13. Weill, F.X., et al., Multidrug resistance in *Salmonella enterica* serotype Typhimurium from humans in France (1993 to 2003). *Journal of Clinical Microbiology*, 2006. **44**(3): p. 700-708.
14. Letchumanan, V., et al., Prevalence and antimicrobial susceptibility of *Vibrio parahaemolyticus* isolated from retail shrimps in Malaysia. *Frontiers in Microbiology*, 2015. **6**: p. 33-33.
15. Pagani, L., et al., Multiple CTX-M-type extended-spectrum beta-lactamases in nosocomial isolates of Enterobacteriaceae from a hospital in northern Italy. *Journal of Clinical Microbiology*, 2003. **41**(9): p. 4264-4269.
16. Awoke T, T.B., Aseffa A, Sebre S, Seman A, Yeshtela B, Abebe T, Mihret A. , Detection of blaKPC and blaNDM carbapenemase genes among *Klebsiella pneumoniae* isolates in Addis Ababa, Ethiopia: Dominance of blaNDM. *PLoS One*. 2022 Apr 27;17(4):e0267657. doi: 10.1371/journal.pone.0267657.
17. Ploy, M.C., et al., Molecular characterization of integrons in *Acinetobacter baumannii*: description of a hybrid class 2 integron. *Antimicrobial Agents and Chemotherapy*, 2000. **44**(10): p. 2684.
18. Rúgeles, L.C., et al. (2010). Molecular characterization of diarrheagenic *Escherichia coli* strains from stools samples and food products in Colombia. *International Journal of Food Microbiology*, 138(3), 282-286.
19. Okumu, N.O., et al. (2023). Risk factors for diarrheagenic *Escherichia coli* infection in children aged 6–24 months in peri-urban community, Nairobi, Kenya. *PLOS Global Public Health*, 3(11), e0002594.
20. Abdoulaye, O., et al. (2020). Prevalence of Virulence Genes of *Escherichia Coli* During Acute Gastroenteritis in Children in Niamey. *HEALTH SCIENCES AND DISEASE*, 21(7).
21. Wolde, A., et al. (n.d.). Molecular Characterization and Antimicrobial Resistance of Pathogenic *Escherichia coli* Strains in Children from Wolaita Sodo, Southern Ethiopia. *Journal of Tropical Medicine*.
22. *Clinical Microbiology and Infection*. (2012). Diarrhoeagenic *Escherichia coli* detected by 16-plex PCR in children with and without diarrhoea in Burkina Faso.
23. Shetty, V.A., et al. (2012). Prevalence and Characterization of Diarrheagenic *Escherichia coli* Isolated from Adults and Children in Mangalore, India. *Journal of Laboratory Physicians*, 4(1), 24-29.
24. Hegde, A., Ballal, M., & Shenoy, S. (2012). Detection of diarrheagenic *Escherichia coli* by multiplex PCR. *Indian Journal of Medical Microbiology*, 30(3), 279-284.
25. Konaté, A., et al. (2017). Molecular characterization of diarrheagenic *Escherichia coli* in children less than 5 years of age with diarrhea in Ouagadougou, Burkina Faso. *EuJMI*, 7(3), 220-228.
26. Moyo, S.J., et al. (2007). Identification of diarrheagenic *Escherichia coli* isolated from infants and children in Dar es Salaam, Tanzania. *BMC Infectious Diseases*, 7(1), 92.
27. Gambushe, S.M., Zishiri, O.T., & El Zowalaty, M.E. (2022). Review of *Escherichia coli* O157:H7 Prevalence, Pathogenicity, Heavy Metal and Antimicrobial Resistance, African Perspective. *Infection and Drug Resistance*, 15, 4645-4673.
28. Trabulsi, L.R., Keller, R., & Gomes, T.A.T. (2002). Typical and Atypical Enteropathogenic *Escherichia coli*. *Emerging Infectious Diseases*, 8(5).
29. Santos, A.C.d.M., et al. (2020). Diversity of Hybrid- and Hetero-Pathogenic *Escherichia coli* and Their Potential Implication in More Severe Diseases. *Frontiers in Cellular and Infection Microbiology*, 10, 339.
30. *Nature Reviews Microbiology*. (2004). Pathogenic *Escherichia coli*.
31. Medugu, N., et al. (2022). Molecular characterization of multi drug resistant *Escherichia coli* isolates at a tertiary hospital in Abuja, Nigeria. *Scientific Reports*, 12(1), 14822.
32. Kargar, M., et al. (2014). High Prevalence of Class 1 to 3 Integrons Among Multidrug-Resistant Diarrheagenic *Escherichia coli* in Southwest of Iran. *Osong Public Health and Research Perspectives*, 5(4), 193-198.
33. CHU du Point G. (2008). Prevalence of extended-spectrum beta-lactamase-producing enterobacteria from 2006 to 2008.
34. Egwu, E., et al. (2023). Antimicrobial susceptibility and molecular characteristics of beta-lactam and fluoroquinolone resistant *E. coli* from human clinical samples in Nigeria. *Scientific African*, 21, e01863.
35. Rupp, M.E., & Fey, P.D. (2003). Extended spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae: considerations for diagnosis, prevention and drug treatment. *Drugs*, 63(4), 353-365.
36. Diawara, A., et al. (2007). Prescription practices in community health centers (CSCOM) and medicine usage by populations.

37. Coulibaly, Y., et al. (2014). Study of antibiotic prescription in Malian hospitals. *Revue Malienne d'Infectiologie et de Microbiologie*, 2-8.
38. Singh, T., et al. (2021). Integron mediated antimicrobial resistance in diarrheagenic *Escherichia coli* in children: in vitro and in silico analysis. *Microbial Pathogenesis*, 150, 104680.
39. Yong, D., et al. (2009). Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrobial Agents and Chemotherapy*, 53(12), 5046-5054.
40. Hornsey, M., Phee, L., & Wareham, D.W. (2011). A novel variant, NDM-5, of the New Delhi metallo- β -lactamase in a multidrug-resistant *Escherichia coli* ST648 isolate recovered from a patient in the United Kingdom. *Antimicrobial Agents and Chemotherapy*, 55(12), 5952-5954.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.