

Article

Not peer-reviewed version

# Demographic Characteristics and Survival in Young Onset Colorectal Neuroendocrine Neoplasms

Deepak Vadehra , [Sahithi Sonti](#) , Beas Siromoni , Mrinalini Ramesh , [Debduti Mukhopadhyay](#) , [Adrienne Groman](#) , Kristopher Attwood , [Renuka Iyer](#) , [Sarbjit Mukherjee](#) \*

Posted Date: 5 September 2024

doi: 10.20944/preprints202409.0261.v1

Keywords: young-onset neuroendocrine neoplasms; young-onset adenocarcinoma; young-onset colorectal cancer (YOCRC)



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Article*

# Demographic Characteristics and Survival in Young Onset Colorectal Neuroendocrine Neoplasms

Deepak Vadehra <sup>1</sup>, Sahithi Sonti <sup>1</sup>, Beas Siromoni <sup>2</sup>, Mrinalini Ramesh <sup>3</sup>,  
Debduti Mukhopadhyay <sup>3</sup>, Adrienne Groman <sup>4</sup>, Kris Attwood <sup>4</sup>, Renuka Iyer <sup>1</sup>  
and Sarbajit Mukherjee <sup>1,\*</sup>

<sup>1</sup> Department of Medicine, Roswell Park Comprehensive Cancer Center, Buffalo, NY 14263, USA

<sup>2</sup> School of Health Sciences, University of South Dakota, Vermillion, SD 57069, USA

<sup>3</sup> Department of Internal Medicine, University at Buffalo, Buffalo, NY 14204, USA

<sup>4</sup> Department of Biostatistics and Bioinformatics, Roswell Park Comprehensive Cancer Center, Buffalo, NY, USA, 14263

\* Correspondence: sarbajit.mukherjee@roswellpark.org; Tel.: +1-716-845-1300; Fax: +1-716-845-8935

**Abstract:** Recent epidemiologic studies have revealed an upward trend in young-onset colorectal cancer (YOCRC) overall, whereas specific data on young-onset colorectal neuroendocrine neoplasms (YONEN) remain limited. This study investigates demographic characteristics and survival trends in YONEN and compares these with those of young-onset colorectal adenocarcinoma (YOADC), the most common histologic subtype of YOCRC. A retrospective analysis was conducted from 2000 to 2019 using the Surveillance, Epidemiology, and End Results (SEER) database. Survival outcomes were assessed using univariate and multivariable Cox proportional models, with demographic differences evaluated via Wilcoxon Rank Sum and Chi-square tests. Out of 61,705 patients aged 20-49 with colorectal cancer, 8% had NEN, and 92% had adenocarcinoma. The YONEN cohort had a higher proportion of Black patients and a lower proportion of White patients than the YOADC cohort (21% vs. 13% and 44% vs. 57%, respectively). NEN was more commonly found in the rectum (79%), and adenocarcinoma was mostly colonic (57%) in origin. YONEN patients had better survival than YOADC patients. Multivariate analysis in YONEN patients revealed that Hispanic patients had better overall survival compared to White patients (HR 0.67, 95% CI 0.47-0.95,  $p=0.024$ ). Racial disparities should be investigated further to aid in policymaking and targeted interventions.

**Keywords:** young-onset neuroendocrine neoplasms; young-onset adenocarcinoma; young-onset colorectal cancer (YOCRC)

## 1. Introduction

Colorectal cancer (CRC) is the second leading cause of cancer-related death in the United States, with adenocarcinoma comprising the majority. According to the American Cancer Society, an estimated 52,550 individuals will succumb to CRC in 2023 [1]. A recent study by Tan et al. examined the mortality trends in colorectal cancer in the US [2]. The study revealed that the incidence rates in CRC decreased significantly from 26.42 to 15.98 per 100,000 individuals, with an Average Annual Percent Change (AAPC) of -2.41. However, the Age-Adjusted Mortality Rate (AAMR) of rectosigmoid cancer went up from 0.82 to 1.08 per 100,000 individuals, with an AAPC of +1.10. Males and Black patients had the highest AAMRs, with rates of 23.90 and 26.93 per 100,000 individuals, respectively. Moreover, the overall AAMR of CRC decreased for those aged 50 years and older but worsened from 1.02 to 1.58 per 100,000 individuals for YOCRC patients, with an AAPC of +0.75. These results show that disparities in CRC mortality persist across age, sex, race, geographic region, and urbanization level, underscoring the necessity for targeted public health interventions.

A study which investigated the impact of race in receiving guideline-concordant care for Young-Onset Colorectal Cancer (YOCRC) in the United States revealed significant findings [3]. Black patients with YOCRC were more likely to be deprived of surgery (adjusted odds ratio [aOR] 1.15, 95% confidence interval [CI] 1.07 to 1.24), have standard (less than 12) lymph nodes examined (aOR 1.11, 95% CI 1.05 to 1.17), and not receive chemotherapy (aOR 1.22, 95% CI 1.17 to 1.27) compared to Caucasian patients. Black patients with rectal cancer were more likely not to have complete staging (aOR 1.90, 95% CI 1.77 to 2.04), not undergo surgery (aOR 1.38, 95% CI 1.30 to 1.45) or chemotherapy (aOR 1.68, 95% CI 1.56 to 1.82), not start radiotherapy (aOR 1.20, 95% CI 1.14 to 1.27), not finish radiotherapy (aOR 1.20, 95% CI 1.12 to 1.30), and be given treatment in the incorrect order (aOR 1.25, 95% CI 1.16 to 1.34).

The projected figures for Young-Onset Colorectal Cancer (YOCRC) reveal a troubling pattern within the 20 to 49-year-old age bracket. By 2040, colorectal cancer is anticipated to rank as the second most prevalent cancer in this demographic, following breast cancer [4]. Significantly, instances and fatalities related to colorectal cancer in younger adults have shown an upward trajectory over the past decade and are projected to continue increasing over the next two decades. This concerning trend may be attributed to factors such as sedentary lifestyles, poor dietary habits, obesity, and a lack of routine screening in this age group [4]. These forecasts emphasize the critical need for heightened awareness, early detection, and screening programs to tackle the escalating burden of colorectal cancer among younger cohorts.

From a histological perspective, adenocarcinoma is the most common subtype of CRC, followed by neuroendocrine neoplasms (NEN), which are a rare subgroup of young-onset colorectal cancers [5]. According to the WHO system, the grades for neuroendocrine neoplasms (NEN) include grades 1, grade 2, grade 3 which are distinguished by the mitotic rate and Ki-67 indices [6]. Tumors are further divided into neuroendocrine carcinomas (NEC), which are high-grade with further subdivision into multiple categories. One of these categories includes Mixed adenoneuroendocrine carcinomas (MaNEC). Mixed Neuroendocrine Non-neuroendocrine neoplasms (MiNEN), on the other hand, are a rare group of NENs that consist of a neuroendocrine and a non-neuroendocrine component, both exceeding 30%. They can be either well or poorly differentiated and was included as a separate category of NENs in the 2019 WHO classification [6]. A recent study by Abboud et al. showed that the rise in YOCRC may be attributed to the fact that the incidence of colorectal NENs is increasing at a rate even faster than adenocarcinoma in the young population [7]. It was observed that there has been a substantial increase in the incidence of neuroendocrine neoplasms compared to adenocarcinomas (ADC) in this population. Specifically, the incidence of NENs showed a much more significant rise than that of ADC, with an average annual percentage change (AAPC) of 2.65 for NENs compared to 0.91 for ADC. This difference in AAPC between NENs and ADC was statistically significant ( $p = 0.01$ ), indicating a notable disparity in the trends of these histopathological subtypes. These findings underscore the necessity for increased awareness and targeted screening strategies to address the rising incidence of colorectal cancer, particularly neuroendocrine neoplasms, in the younger population, ultimately aiming to enhance early detection and improve patient outcomes.

Similarly, Lumsdaine et al. conducted a population-based study that identified a substantial rise in young-onset colorectal cancer (CRC), specifically in rectal cancer, with a focus on the increasing trend of rectal neuroendocrine neoplasms (NENs) [8]. The incidence of rectal NENs exhibited a significant increase across all age groups, particularly notable in individuals aged 45-54 and those over 55 years. In the younger age brackets of 20-44 and 45-54 years, the annual percent changes (APCs) for rectal NENs were calculated at 2.9 and 6.1, respectively, indicating a notable upward trajectory in incidence rates. Notably, the surge in rectal NENs played a substantial role in the overall increase in rectal cancer cases, with statistics revealing that NENs contributed significantly, accounting for 26.74% and 53.47% of the total increase in the respective age groups. These findings underscore the increasing impact of rectal NENs on the prevalence of young-onset colorectal cancer, underscoring the necessity for further research and clinical focus to address this emerging trend and its implications on patient care and management strategies.

Given the rarity of NENs and their typically slow progression, limited data exists concerning these tumors in younger individuals. Therefore, there is an unmet need to learn about overall survival (OS), disease-specific survival (DSS), and the factors affecting survival in young-onset colorectal NEN. This study aimed to explore patterns and disparities in survival rates among young individuals diagnosed with colorectal NEN and compare them with young-onset colorectal adenocarcinoma patients as well as those with average onset colorectal NEN patients. Additionally, we aimed to identify factors affecting survival in colorectal NEN.

## 2. Materials and Methods

### 2.1. Study Design and Population

We conducted a retrospective study on YOCRC patients in the US between 2000 and 2019 using the Surveillance, Epidemiology, and End Results (SEER) database. The SEER database compiles cancer-specific incidence data from population-based registries, covering approximately 35% of the US population [9]. Our primary endpoint was to estimate overall survival (OS), defined as the time from diagnosis to death from any cause. Our secondary endpoint was Disease-specific survival (DSS), defined as the time from diagnosis to death specifically due to cancer. For the purpose of our analysis, we included adenocarcinomas, neuroendocrine tumors, neuroendocrine carcinomas, MiNEN and MANEC.

### 2.2. Covariates

Key variables of interest included the patient's demographic characteristics such as age, race, sex, and disease characteristics. Clinical variables of interest included disease stage, grade, surgery, primary cancer site, and year of diagnosis. Race was categorized into non-Hispanic whites, non-Hispanic blacks, Hispanics, Asian Americans, and Native Americans. The stage was classified according to the AJCC Classification System [10] as Stage 0, 1, 2, 3, and 4. Grade was classified as 1, 2, 3, and undifferentiated. The primary cancer site was categorized as colon, rectum, or rectosigmoid, and the year of diagnosis was divided into five periods: 2000-2003, 2004-2007, 2008-2011, 2012-2015, and 2016-2019.

### 2.3. Statistical Analysis

Demographic and clinical characteristics of the overall sample were summarized. For categorical variables, frequencies and relative frequencies were provided and compared using the Pearson chi-square test while contrasting YONENs and YOADCs. The Kruskal-Wallis test, a nonparametric test, was used for ordinal or continuous variables. The effects of various factors on OS and DSS were estimated using Cox proportional hazard models adjusted for age at diagnosis (20-39, 40-49,  $\geq 50$ ), sex, race, stage, grade, primary site, tumor size, surgery, and year of diagnosis. Result estimates were expressed as hazard ratios (HR) with 95% confidence intervals (CI). The Kaplan-Meier method calculated OS and DSS. Yearly mortality rates per 100,000 were calculated to examine differences among histology and age groups. Unknown stage, unknown grade and unknown race were included in the model as they are SEER reportable statuses. Variables that could not be estimated, were removed from the model. All model assumptions, proportional hazards and goodness of fit were evaluated visually using standard residual plots (Schoenfeld residuals versus time and standardized residuals versus predicted values). The significance level was denoted by  $p < 0.05$ . SAS, version 9.4 statistical software (SAS Institute Inc., Cary, NC) was used for all statistical analyses.

## 3. Results

There were 61,705 patients in the young-onset colorectal cancer (YOCRC) group, of which 5,128 belonged to the young-onset neuroendocrine neoplasms (YONEN) cohort, and the remaining 56,577 cases were part of the young-onset adenocarcinoma (YOADC) group. Fifty-two percent of the YONEN group were female, compared with 45.8% in the YOADC group ( $p < 0.001$ ). In the YONEN

group, 43.6% were White, 20.6% were Black, 18.2% were Hispanic, 11.8% were Asian, and 1.2% were Native American. In the YOADC group, 58% were White, 17.6% Hispanic, 13.15% Black, 10.3% Asian, and 1% Native American ( $p < 0.001$ ). The primary disease site for most YONENs was the rectum (78.7%), followed by the colon (17.3%) and the rectosigmoid region (4%). Conversely, the majority of YOADCs had the colon as their primary disease site (61.2%), with 28.3% in the rectum and 10.5% in the rectosigmoid ( $p < 0.001$ ). The most common stage at diagnosis for YONENs was Stage I (21%), followed by 6.3% with Stage IV disease. However, a significant majority (67.2%) did not have a reported stage in the SEER database. For the YOADC group, 29% had Stage III disease at diagnosis, another 23.6% had Stage IV disease, with only 14.8% not having a reported stage ( $p < 0.001$ ). The demographic characteristics are described in Table 1.

We also compared young and average onset NEN patients. We found that of patients from the younger group ( $< 50$  years), 52% of the group were female patients, compared with 48.9% in the average onset group (patients aged 50 and above). Hispanic patients were overrepresented in the younger age group (18.2%), compared to the average onset group (13.5%). YONEN patients had a higher proportion of rectal tumors compared to the average onset NEN patients (79% vs 69%). Patients in the average onset group were also found to have a significantly higher proportion of grade III tumors (8%) compared to YONEN patients (4.3%),  $p < 0.001$ . This comparison is detailed in Table 2.

We analyzed overall survival and disease-specific survival for YONEN and YOADC patients. YONENs had a 1-year survival rate of 0.94 (95% CI: 0.93, 0.94) and a 5-year survival rate of 0.88 (95% CI: 0.87, 0.89), whereas YOADCs had a 1-year survival rate of 0.90 (95% CI: 0.90, 0.90) and a 5-year survival rate of 0.63 (95% CI: 0.63, 0.64) with median follow up time of 105 months. The 1-year disease-specific survival (DSS) rate for YONENs was 0.95 (95% CI: 0.94, 0.95) compared to 0.91 (95% CI: 0.91, 0.91) for YOADCs. The 5-year DSS rate for YONENs was 0.91 (95% CI: 0.90, 0.92) versus 0.66 (95% CI: 0.65, 0.66) for YOADCs with median follow up time of 97 months. Survival rates by age group are detailed in Tables 3 and 4 as well as the Kaplan-Meier curves in Figures A1 and A2.

Next, we examined the mortality trends. The rate of deaths per 100,000 decreased from 49,554 in 2000-2003 to 15,527 in the period between 2016-2019. Despite the rising incidence, there was a numerical improvement in mortality rates in recent years: the YONEN mortality rate decreased from 20,403 per 100,000 in 2000-2003 to 6,705.3 per 100,000 in 2016-2019 (Figure 1). A similar magnitude of decrease was observed in the YOADC group, from 51,928.6 per 100,000 in 2000-2003 to 16,331.9 in 2016-2019 ( $p < 0.001$ ) (Figure 1).

We also examined survival differences between the young-onset ( $< 50$ ) and average onset ( $\geq 50$ ) colorectal neuroendocrine neoplasms cohorts and found that the overall survival in the young onset cohort was better than the average onset NEN cohort. The 5-year overall survival rate between the young-onset and average onset groups was 0.89 (95% CI: 0.88, 0.89) versus 0.65 (95% CI: 0.64, 0.66) after a median follow up time of 96 months. Similarly, the 5-year disease-specific survival rate for the young versus average onset groups was 0.92 (95% CI: 0.92, 0.92) versus 0.76 (95% CI: 0.75, 0.77) after a median follow up time of 82 months. This is likely related to the fact that younger patients have fewer comorbidities and better functional status, and hence can tolerate more aggressive treatments, leading to better survival rates. Survival rates of the young onset versus average onset colorectal NEN are detailed in Tables 5 and 6.

Finally, we assessed factors affecting overall survival and disease-specific survival in the YONEN population. Higher grade was associated with worse overall survival (OS) (Grade III vs. Grade I; HR = 11.88,  $p < .001$  and Undifferentiated vs. Grade I; HR = 11.23,  $p < .001$ ) and worse DSS (Grade III vs. Grade I; HR = 29.05 and Undifferentiated vs. Grade I; HR = 31.49,  $p < .001$ ). Rectal NENs were associated with improved overall survival (HR = 0.63,  $p = 0.008$ ), as was the Hispanic race (HR = 0.67,  $p = 0.024$ ). (Tables A1 and A2).




Figure 1. Mortality rates in YOADC and YONEN in the last two decades (per 100,000).

Table 1. Demographic and Clinical Characteristics of Young-Onset Colorectal Neuroendocrine neoplasms and Adenocarcinomas.

		Neuroendocrine neoplasms	Adenocarcinoma	p-value
Age	20-39	1714 (33.4%)	14164 (25%)	p<0.001
	40-49	3414 (66.6%)	42413 (75%)	
Sex	Male	2456 (47.9%)	33108 (53.7%)	p<0.001
	Female	2672 (52.1%)	28597 (46.3%)	
Race	White	2235 (43.6%)	32525 (57.5%)	p<0.001
	Black	1054 (20.6%)	7429 (13.1%)	
	Asian	603 (11.8%)	5830 (10.3%)	
	Native American	61 (1.2%)	573 (1.0%)	
	Hispanic	935 (18.2%)	9937 (17.6%)	
	Not reported	240 (4.7%)	283 (0.5%)	
Year of diagnosis	2000-2003	794 (15.5%)	9748 (17.2%)	p=0.035

	2004-2007	950 (18.5%)	10911 (19.3%)	
	2008-2011	1054 (20.6%)	11129 (19.7%)	
	2012-2015	1122 (21.9%)	11545 (20.4%)	
	2016-2019	1208 (23.6%)	13244 (23.4%)	
Disease Site	Colon	887 (17.3%)	34614 (61.2%)	p<0.001
	Rectosigmoid	204 (4.0%)	5954 (10.5%)	
	Rectum	4037 (78.7%)	16009 (28.3%)	
Disease Grade	I	1760 (34.3%)	3848 (6.8%)	p<0.001
	II	302 (5.9%)	36234 (64%)	
	III	220 (4.3%)	9405 (16.6%)	
	Undifferentiated	112 (2.2%)	971 (1.7%)	
	Not reported	2734 (53.3%)	6119 (10.8%)	
Overall Stage	I	1076 (21.0%)	6873 (12.1%)	p<0.001
	II	102 (2.0%)	10695 (18.9%)	
	III	129 (2.5%)	16526 (29.2%)	
	IV	323 (6.3%)	13349 (23.6%)	
	Not reported	3448 (67.2%)	8372 (14.8%)	

**Table 2.** Colorectal Neuroendocrine Neoplasms– Descriptive Statistics by Age (< 50 vs ≥ 50).

		< 50	≥ 50	p-value
Sex	Male	2456 (47.9%)	10648 (51.1%)	p<0.001
	Female	2672 (52.1%)	10209 (48.9%)	
Race	White	2235 (43.6%)	10420 (50.0%)	p<0.001
	Black	1054 (20.6%)	4214 (20.2%)	
	Asian	603 (11.8%)	2490 (11.9%)	
	Native American	61 (1.2%)	158 (0.8%)	
	Hispanic	935 (18.2%)	2825 (13.5%)	
	Not reported	240 (4.7%)	750 (3.6%)	
Year of diagnosis	2000-2003	794 (15.5%)	2773 (13.3%)	p<0.001
	2004-2007	950 (18.5%)	3579 (17.2%)	
	2008-2011	1054 (20.6%)	4428 (21.2%)	
	2012-2015	1122 (21.9%)	4931 (23.6%)	
	2016-2019	1208 (23.6%)	5146 (24.7%)	
Disease Site	Colon	887 (17.3%)	5668 (27.2%)	p<0.001
	Rectosigmoid	204 (4.0%)	864 (4.1%)	
	Rectum	4037 (78.7%)	14325 (68.7%)	
Disease Grade	I	1760 (34.3%)	6973 (33.4%)	p<0.001
	II	302 (5.9%)	1272 (6.1%)	
	III	220 (4.3%)	1678 (8.0%)	

	Undifferentiated	112 (2.2%)	684 (3.3%)	
	Not reported	2734 (53.3%)	10250 (49.1%)	
Overall Stage	In situ	50 (1.0%)	192 (0.9%)	p<0.001
	I	1076 (21.0%)	3945 (18.9%)	
	II	102 (2.0%)	473 (2.3%)	
	III	129 (2.5%)	1080 (5.2%)	
	IV	323 (6.3%)	1760 (8.4%)	
	Not reported	3448 (67.2%)	13407 (64.3%)	
Tumor size in mm	Mean/StdErr	15.1/0. 5	20.3/0. 4	p<0.001

Table 3. Age-specific OS Rates at 1 Year and 5 Years in YONEN and YOADC patients.

Age group (years)	1-yr survival rate (NEN) (95% CI) (N=5128)	5-yr survival rate (NEN) (95% CI) (N=5128)	1-yr survival rate (ADC) (95% CI) (N=56,577)	5-yr survival rate (ADC) (95% CI) (N=56,577)
20-39	0.96 (0.94, 0.96)	0.91(0.89, 0.92)	0.90 (0.89, 0.90)	0.62 (0.61, 0.63)
40-49	0.93 (0.92, 0.94)	0.87 (0.86, 0.88)	0.90 (0.90, 0.90)	0.64 (0.63, 0.64)

Table 4. Age-specific DSS rates at 1 year and 5 years in YONEN and YOADC patients.

Age group (years)	1-yr disease-specific survival rate (NEN) (95% CI) (N=5128)	5-yr disease-specific-survival rate (NEN) (95% CI) (N=5128)	1-yr disease-specific-survival rate (ADC) (95% CI) (N=56,577)	5-yr disease-specific-survival rate (ADC) (95% CI) (N=56,577)
20-39	0.96 (0.95, 0.97)	0.93 (0.91, 0.94)	0.90 (0.90, 0.91)	0.64 (0.63, 0.65)
40-49	0.94 (0.93, 0.95)	0.90 (0.89, 0.91)	0.91(0.91, 0.91)	0.66 (0.66, 0.67)

**Table 5.** Age-specific OS Rates between young-onset (< 50) vs average onset (>= 50) colorectal NENs at 1 Year and 5 Years.

Age group (years)	1-yr disease-specific survival rate (NEN)  (95% CI)  (N=25,985)	5-yr disease specific-survival rate (NEN)  (95% CI)  (N=25,985)
< 50	0.94 (0.94, 0.95)	0.89 (0.88, 0.89)
>= 50	0.80 (0.80, 0.81)	0.65 (0.64, 0.66)

**Table 6.** Age-specific DSS rates between young-onset (< 50) vs average onset (>= 50) colorectal NENs at 1 Year and 5 Years.

Age group (years)	1-yr disease-specific survival rate (NEN)  (95% CI)  (N=25,985)	5-yr disease specific-survival rate (NEN)  (95% CI)  (N=25,985)
< 50	0.95 (0.95, 0.96)	0.92 (0.92, 0.92)
>= 50	0.85 (0.84, 0.85)	0.76 (0.75, 0.77)

4. Discussion

The increase in colorectal cancer in young people is attributable to the rise in both neuroendocrine neoplasms and adenocarcinomas. However, Abboud et al. have shown that the rate of increase in neuroendocrine neoplasms in the population under 50 years is significantly higher than that of adenocarcinoma. Adolescents and young adults (AYA) represent a unique population Up to age 39 years of age. These patients are distinctive, as rare cancers are overrepresented in this group [11,12]. Given that this population does not routinely undergo screening colonoscopies, we wanted to see the distribution pattern of adenocarcinoma versus neuroendocrine neoplasms in these patients compared to those in the 40-49 age group. Similar to Abboud et al., we noted that the AYA population is overrepresented with NEN compared to adenocarcinomas (Table 1).

We also noted several demographic differences between the YONEN and average onset neuroendocrine neoplasm populations. When we compared these two groups, we found that female, and Hispanic patients were overrepresented in the younger population and that the primary tumor site was more likely to be in the rectum. This calls for targeted interventions in younger female and Hispanic patients. Besides, since these tumors are commonly found in the rectum, screening via flexible sigmoidoscopy should be considered in yearly testing.

A considerably larger fraction of YONEN patients were Black when compared to those with adenocarcinoma (20.6% vs 13.1%) (P<0.001). This is consistent with what has been seen in other studies. In a SEER analysis [13] of all NENs, it was seen that Black patients had a higher incidence and worse survival when compared to other races. However, our multivariate analysis did not identify Black race as an individual prognostic factor. This could be due to relatively small sample size in our study. In another study, Herring et al. showed significant differences in gene expression between Black and White pancreatic NEN (pNEN) patients, indicating potential disparities in tumor microenvironment that could affect outcomes [14]. RNA sequencing of pNENs from Black and White patients identified 372 markedly differentially expressed genes and 179 enriched gene sets, with key pathways associated with angiogenesis, blood vessel formation, cell migration, and immune

response. Black patients showed enrichment in gene sets associated with blood vessel formation and cellular migration, while immune response pathways were downregulated in this group. These findings suggest distinct tumor biology in NENs from Black patients that may contribute to the disparate outcomes observed in this population, highlighting the importance of further validation and consideration of genetic ancestry in future studies.

Another critical difference we saw was the primary site of the tumor: 78.7% of YONENs were in the rectum, whereas 61.2% of YOADCs were primarily in the colon. Interestingly, we noted essential survival differences in the YONEN vs YOADC population. Despite the rapid increase in the incidence of NENs, the median overall survival is high when compared to YOADCs. We looked at the overall survival rate in YONENs in the adolescent and young adult (AYA, 20-39 years) and 40-49-year-old subgroups. The age-specific OS rate slightly worsened with increasing age (0.96 vs 0.93 1-year survival rate and 0.91 vs 0.87 5-year survival rate, respectively). Nevertheless, this trend was not seen in the YOADCs (0.90 vs. 0.90 for a 1-year survival rate and 0.62 vs 0.64 for a 5-year survival rate). It may be hypothesized that while the rise in incidence in YOCRC is attributable to the exponential increase in YONENs, the poor OS in this population is primarily still due to YOADCs, suggesting that adenocarcinoma in the young-onset cohort, specifically in the AYA cohort is an aggressive subtype [15,16].

Even with the differences in overall survival, mortality rates for both YOADC and YONENs consistently decreased over the last two decades in our study. This is in line with the decrease in mortality in all NENs, seen in a SEER analysis from 2017 [17] and an improved OS in the YOADC population seen in an NCDB analysis conducted in 2021 [18]. This may be attributable to multiple factors: earlier screening and more treatment options in our therapeutic armamentarium. Research into available therapies is moving at a fast pace, with developments such as the tremendous success of PRRT [19,20].

Finally, we looked at the potential factors affecting mortality in the YONEN population and, predictably, found that higher grade was associated with worse overall survival. We also found that Hispanic patients had better outcomes than White patients, corroborating existing literature. A SEER analysis studying racial/ethnic disparities in non-pancreatic NENs found that Hispanic patients had the best overall survival when compared with non-Hispanic White and Black patients [21]. The study by Gosku et al. uncovered notable disparities in survival outcomes across racial and ethnic groups, shedding light on the nuanced impact of race and ethnicity on disease prognosis. Hispanic patients emerged as a cohort with distinct survival advantages, showcasing better overall survival rates than non-Hispanic White patients, with a Hazard Ratio of 0.89 (0.81–0.97). This lower risk of mortality among Hispanic individuals underscores a significant disparity in outcomes that warrants further investigation. Moreover, when examining specific primary tumor sites, Hispanic patients demonstrated superior overall survival in locations such as the small intestine and rectum, with Hazard Ratios of 0.81 (0.69–0.96) and 0.79 (0.63–0.99), respectively [20]. These findings suggest a potential biological or treatment-related advantage for Hispanic patients in these particular anatomical sites, highlighting the complexity of factors influencing survival disparities in neuroendocrine neoplasms.

The improved survival rates of Hispanics/Latinos compared to Non-Hispanic Whites can be attributed to a complex interplay of genetic, behavioral, cultural, and environmental factors [22]. Genetic variances across racial/ethnic groups may influence survival outcomes. Behavioral differences, such as smoking patterns, also contribute, with Hispanics/Latinos potentially engaging in behaviors that confer a survival advantage. Despite often having lower socioeconomic status, this group may experience similar or better health outcomes, suggesting the influence of other factors. Cultural aspects like *familismo*, a strong family-oriented philosophy prevalent in Hispanic culture, could play a crucial role in promoting better health outcomes [22]. Additionally, survival advantages may arise not just from differences in healthcare access, but also from disparities in environmental exposures, cultural influences, and treatment approaches across racial/ethnic groups [22]. These elements together underscore the multifaceted reasons behind the superior survival outcomes observed in this demographic. Nevertheless, our findings highlight the critical role of race and

ethnicity as independent prognostic factors in neuroendocrine neoplasms, emphasizing the importance of tailored interventions and personalized treatment strategies to address disparities and improve overall survival rates for diverse patient populations.

In moving forward from the findings of this study on young onset colorectal neuroendocrine neoplasms, several key areas warrant further investigation to advance our understanding and improve patient outcomes. Better documentation at diagnosis of stage would aid identifying differences in stage at presentation if any. Investigating genetic markers could lead to personalized treatment approaches, while examining the impact of lifestyle factors such as diet, exercise, and environmental exposures could provide valuable insights into disease development and progression. In addition, there are data showing that higher pain and stress scores using validated scales can correlate with poor outcomes in other diseases and the impact of these markers warrants further study in this group where anxiety and stress are expected to be high, underscoring the importance of databases that capture these measures [23]. Evaluating novel treatment modalities, including immunotherapy and targeted therapies, through randomized clinical trials, will help identify optimal strategies for YONEN patients.

Long-term follow-up studies are essential to assess survival outcomes and quality of life, helping to optimize patient care over time. Additionally, research on the impact of health insurance and care disparities is pivotal for addressing inequities, especially among minority groups [24,25]. Prior studies have documented disparities in relative survival that often impacts minority groups [26]. Investigations have found that minority patients often have inadequate insurance that in turn results in increased risk of locally advanced disease on diagnosis [27]. Investigating barriers to healthcare access and designing interventions to reduce these disparities can improve outcomes [28]. Healthcare policy analysis should focus on evaluating existing policies and advocating for targeted interventions to reduce treatment disparities [29]. Implementing strategies like mobile screening programs and outreach oncology clinics can serve medically underserved communities, advancing health equity [30,31]. These research directions aim to tailor interventions to individual needs and enhance the quality of care and survival rates for YONEN patients.

Our study is not without limitations. Firstly, the study's retrospective design may introduce biases and limitations in data collection, analysis, and interpretation. Secondly, we utilized data from the Surveillance, Epidemiology, and End Results (SEER) database, which may have limitations in terms of data accuracy, completeness, and consistency. The quality of the data in the SEER database, especially the significant amount of unreported data on disease stage and grade, might have impacted some of the study's findings like the multivariate analysis. Besides, the grades reported in the SEER database are not concordant with the WHO grading classification of NENs [32]. We also removed the variables that could not be estimated from the multivariate model. Finally, the study may not have accounted for all potential confounding variables that could influence the outcomes of interest, such as comorbidities, or treatment variations: like time to treatment initiation, quality of surgery, or type of chemotherapy, which were not available in the SEER database. Despite these limitations, our study was one of the largest studies to look at the demographic characteristics and survival of YONEN patients at a population level.

## 5. Conclusions

In conclusion, our study identified several socio-demographic disparities in the YONEN and YOADC patients. We also found that despite increasing incidence rates, mortality rates have been steadily decreasing in this population. Further research is needed to understand the disparities further for resource allocation as well as implementation of appropriate healthcare policies.

**Author Contributions:** D.V. and S.M. were involved in the conception of the study; D.V., S.S. S.M and B.S. were involved in data curation and wrote the manuscript; K.A. and A.G. were involved in data analysis and methodology; R.I., M.R. and D.M. reviewed and edited the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported through funding by the National Cancer Institute (Grant number: P30CA016056). The study’s design and decision to publish were independent of any involvement from the funding sources.

**Ethics Approval:** Not Applicable.

**Consent for Publication:** Since de-identified retrospectively collected publicly available data was utilized, informed consent requirement was not required.

**Data Availability:** All the data used for this study is from the publicly available SEER database.

**Conflicts of Interest:** Sarbajit Mukherjee serves as a volunteer guidelines panel member at the National Comprehensive Cancer Network and American Society of Clinical Oncology. He received research funding from the National Comprehensive Cancer Network and Ipsen Biopharmaceuticals/North American Neuroendocrine Tumor Society which were paid to the institute. Dr. Mukherjee received consulting fee from Merck, Eisai, and Beigene. Sarbajit Mukherjee and all other authors declare no other conflict of interest.

Abbreviations

Young-onset adenocarcinoma	(YOADC)
Young-onset neuroendocrine neoplasms	(YONEN)
Surveillance, Epidemiology, and End Results	(SEER)
Overall survival	(OS)
Disease-specific survival	(DSS)
Colorectal cancer	(CRC)
Young-Onset Colorectal Cancer	(YOCRC)
Gastroenteropancreatic neuroendocrine neoplasms	(GEP-NENs)
National Cancer Institute	(NCI)
Rural-Urban Continuum Code	(RUCC)
American Joint Committee on Cancer	(AJCC)
Hazard Ratio	(HR)
Confidence Intervals	(CI)
Adolescent and young adult	(AYA)
Mixed neuroendocrine non-neuroendocrine neoplasms	(MiNEN)
Mixed adenoneuroendocrine carcinoma	(MANEC)
Adolescents and young adults	(AYA)

Appendix A

**Table A1.** Factors affecting overall survival in YONEN (N=2636).


Variable	Reference Group	Estimate (95% CL)	p-value
Age			
40-49	20-39	1.14 (0.89, 1.46)	0.297
Gender			
Female	Male	0.87 (0.70, 1.08)	0.204
Grade			
II	I	1.41 (0.82, 2.43)	0.211
III	I	11.88 (8.16, 17.29)	<0.001
Undifferentiated	I	11.23 (7.37, 17.12)	<0.001
Not Reported	I	2.13 (1.48, 3.08)	<0.001
Race			
Asian	White	0.96 (0.65, 1.41)	0.842
Black	White	1.20 (0.91, 1.58)	0.203
Hispanic	White	0.67 (0.47, 0.95)	0.024
Native American	White	1.45 (0.53, 3.94)	0.471
Site			
Rectosigmoid	Colon	0.82 (0.47, 1.41)	0.468
Rectum	Colon	0.63 (0.45, 0.89)	0.008
Stage			
I	0	1.21 (0.17, 8.90)	0.849
II	0	1.32 (0.16, 10.58)	0.794
III	0	3.00 (0.40, 22.35)	0.283
IV	0	7.86 (1.07, 57.64)	0.043

Not Reported	0	1.75 (0.24, 12.70)	0.582
<b>Surgical Procedure</b>			
Colectomy	None	0.58 (0.40, 0.84)	0.003
Local Procedure/Wedge Resection	None	0.33 (0.24, 0.45)	<0.001
Total Proctectomy	None	0.31 (0.13, 0.71)	0.006
Total Proctocolectomy	None	0.42 (0.13, 1.36)	0.150
<b>Year of Diagnosis</b>			
2008-2011	2004-2007	0.92 (0.67, 1.26)	0.605
2012-2015	2004-2007	1.17 (0.83, 1.64)	0.371
2016-2019	2004-2007	1.33 (0.91, 1.95)	0.140


**Table A2.** Factors affecting disease specific survival in YONEN (N=2636).

Variable	Reference Group	Estimate (95% CL)	p-value
<b>Age</b>			
40-49	20-39	1.16 (0.87, 1.54)	0.317
<b>Gender</b>			
Female	Male	0.78 (0.61, 1.00)	0.052
<b>Grade</b>			
II	I	2.57 (1.33, 4.99)	0.005
III	I	29.05 (18.12, 46.56)	<0.001
Undifferentiated	I	31.49 (18.96, 52.32)	<0.001
Not Reported	I	2.17 (1.33, 3.53)	0.002
<b>Race</b>			
Asian	White	1.15 (0.74, 1.78)	0.546

Variable	Reference Group	Estimate (95% CL)	p-value
Black	White	1.02 (0.73, 1.43)	0.905
Hispanic	White	0.76 (0.52, 1.11)	0.155
Native American	White	0.49 (0.07, 3.57)	0.484
<b>Site</b>			
Rectosigmoid	Colon	1.00 (0.56, 1.77)	0.995
Rectum	Colon	0.45 (0.31, 0.65)	<0.001
<b>Surgical Procedure</b>			
Colectomy	None	0.40 (0.27, 0.58)	<0.001
Local Procedure/Wedge Resection	None	0.15 (0.10, 0.23)	<0.001
Procedure Not Reported	None	0.35 (0.05, 2.50)	0.294
Total Proctectomy	None	0.44 (0.19, 1.03)	0.059
Total Proctocolectomy	None	0.57 (0.18, 1.83)	0.342
<b>Year of Diagnosis</b>			
2008-2011	2004-2007	0.99 (0.69, 1.41)	0.954
2012-2015	2004-2007	1.04 (0.72, 1.51)	0.816
2016-2019	2004-2007	0.77 (0.51, 1.17)	0.220



**Figure A1.** OS Rates in YOADC and YONEN patients (N=45,827).



**Figure A2.** DSS rates in YOADC and YONEN patients (N=45,827).

## References

1. Siegel, R. L.; Wagle, N. S.; Cercek, A.; Smith, R. A.; Jemal, A. Colorectal cancer statistics, 2023. *CA Cancer J Clin* 2023, 73 (3), 233-254. <https://doi.org/10.3322/caac.21772> From NLM.
2. Tan, J. Y.; Yeo, Y. H.; Ng, W. L.; Fong, Z. V.; Brady, J. T. How have US colorectal cancer mortality trends changed in the past 20 years? *Int J Cancer* 2024. <https://doi.org/10.1002/ijc.34926> From NLM.
3. Nogueira, L. M.; May, F. P.; Yabroff, K. R.; Siegel, R. L. Racial Disparities in Receipt of Guideline-Concordant Care for Early-Onset Colorectal Cancer in the United States. *J Clin Oncol* 2024, 42 (12), 1368-1377. <https://doi.org/10.1200/jco.23.00539> From NLM.
4. Rahib, L.; Wehner, M. R.; Matrisian, L. M.; Nead, K. T. Estimated Projection of US Cancer Incidence and Death to 2040. *JAMA Netw Open* 2021, 4 (4), e214708-e214708. <https://doi.org/10.1001/jamanetworkopen.2021.4708> (accessed 5/30/2024).
5. Stewart, S. L.; Wike, J. M.; Kato, I.; Lewis, D. R.; Michaud, F. A population-based study of colorectal cancer histology in the United States, 1998-2001. *Cancer* 2006, 107 (5 Suppl), 1128-1141. <https://doi.org/10.1002/cncr.22010> From NLM.
6. Helderma, N. C.; Suerink, M.; Kiliç, G.; van den Berg, J. G.; Nielsen, M.; Tesselaar, M. E. T. Relation between WHO Classification and Location- and Functionality-Based Classifications of Neuroendocrine Neoplasms of the Digestive Tract. *Neuroendocrinology* 2023, 114 (2), 120-133. <https://doi.org/10.1159/000534035> (accessed 7/13/2024).

7. Abboud, Y.; Fraser, M.; Qureshi, I.; Hajifathalian, K. Early-Onset Colorectal Cancer: Are Neuroendocrine Tumors or Adenocarcinomas the Culprit? Analysis of the Largest U.S. Cancer Incidence Database, 2001-2020. *J Clin Med* 2024, 13 (4). <https://doi.org/10.3390/jcm13041098> From NLM.
8. Lumsdaine, C. T.; Liu-Smith, F.; Li, X.; Zell, J. A.; Lu, Y. Increased incidence of early onset colorectal adenocarcinoma is accompanied by an increased incidence of rectal neuroendocrine tumors. *Am J Cancer Res* 2020, 10 (6), 1888-1899. From NLM.
9. Seer incidence data—Seer data & software. (n.d.). SEER. Retrieved April 29, 2024, from <https://seer.cancer.gov/data/index.html>
10. Weiser, M. R. AJCC 8th Edition: Colorectal Cancer. *Ann Surg Oncol* 2018, 25 (6), 1454-1455. <https://doi.org/10.1245/s10434-018-6462-1> From NLM.
11. Barr, R. D.; Ferrari, A.; Ries, L.; Whelan, J.; Bleyer, W. A. Cancer in Adolescents and Young Adults: A Narrative Review of the Current Status and a View of the Future. *JAMA Pediatr* 2016, 170 (5), 495-501. <https://doi.org/10.1001/jamapediatrics.2015.4689> From NLM.
12. Biller, L. H.; Ng, K. Pioneering a new care model for young-onset colorectal cancer: innovations in clinical care and scientific discovery. *Colorectal Cancer* 2020, 9 (3), CRC18. <https://doi.org/10.2217/crc-2020-0011>.
13. Shen, C.; Gu, D.; Zhou, S.; Xu, Y.; Sarshekeh, A. M.; Halperin, D.; Shih, Y. T.; Yao, J. C.; Dasari, A. Racial Differences in the Incidence and Survival of Patients With Neuroendocrine Tumors. *Pancreas* 2019, 48 (10), 1373-1379. <https://doi.org/10.1097/mpa.0000000000001431> From NLM.
14. Herring, B.; Guenter, R.; Dhall, D.; Chen, H.; Yates, C.; Bart, R. J. Transcriptomic influences of racial disparities in black patients with pancreatic neuroendocrine tumors. In Basic Science, Proceedings of the 15th Annual Multidisciplinary NET Medical Symposium NANETS 2022; Endocrine Abstracts; abstr 21456
15. Georgiou, A.; Khakoo, S.; Edwards, P.; Minchom, A.; Kouvelakis, K.; Kalaitzaki, E.; Nobar, N.; Calamai, V.; Ifijen, M.; Husson, O.; et al. Outcomes of Patients with Early Onset Colorectal Cancer Treated in a UK Specialist Cancer Center. *Cancers (Basel)* 2019, 11 (10). <https://doi.org/10.3390/cancers11101558> From NLM.
16. Sheth Bhutada, J.; Hwang, A.; Liu, L.; Deapen, D.; Freyer, D. R. Poor-Prognosis Metastatic Cancers in Adolescents and Young Adults: Incidence Patterns, Trends, and Disparities. *JNCI Cancer Spectr* 2021, 5 (3). <https://doi.org/10.1093/jncics/pkab039> From NLM.
17. Dasari, A.; Shen, C.; Halperin, D.; Zhao, B.; Zhou, S.; Xu, Y.; Shih, T.; Yao, J. C. Trends in the Incidence, Prevalence, and Survival Outcomes in Patients With Neuroendocrine Tumors in the United States. *JAMA Oncol* 2017, 3 (10), 1335-1342. <https://doi.org/10.1001/jamaoncol.2017.0589> From NLM.
18. Cheng, E.; Blackburn, H. N.; Ng, K.; Spiegelman, D.; Irwin, M. L.; Ma, X.; Gross, C. P.; Tabung, F. K.; Giovannucci, E. L.; Kunz, P. L.; et al. Analysis of Survival Among Adults With Early-Onset Colorectal Cancer in the National Cancer Database. In *JAMA Netw Open*, 2021; Vol. 4, p e2112539.
19. Singh, S.; Halperin, D. M.; Myrehaug, S.; Herrmann, K.; Pavel, M.; Kunz, P. L.; Chasen, B.; Capdevila, J.; Tafuto, S.; Oh, D.-Y.; et al. [177Lu]Lu-DOTA-TATE in newly diagnosed patients with advanced grade 2 and grade 3, well-differentiated gastroenteropancreatic neuroendocrine tumors: Primary analysis of the phase 3 randomized NETTER-2 study. *Journal of Clinical Oncology* 2024, 42 (3\_suppl), LBA588-LBA588. [https://doi.org/10.1200/JCO.2024.42.3\\_suppl.LBA588](https://doi.org/10.1200/JCO.2024.42.3_suppl.LBA588).
20. Strosberg, J. R.; Caplin, M. E.; Kunz, P. L.; Ruszniewski, P. B.; Bodei, L.; Hendifar, A.; Mittra, E.; Wolin, E. M.; Yao, J. C.; Pavel, M. E.; et al. (177)Lu-Dotatate plus long-acting octreotide versus high-dose long-acting octreotide in patients with midgut neuroendocrine tumours (NETTER-1): final overall survival and long-term safety results from an open-label, randomised, controlled, phase 3 trial. *Lancet Oncol* 2021, 22 (12), 1752-1763. [https://doi.org/10.1016/s1470-2045\(21\)00572-6](https://doi.org/10.1016/s1470-2045(21)00572-6) From NLM.
21. Goksu, S. Y.; Ozer, M.; Beg, M. S.; Sanford, N. N.; Ahn, C.; Fangman, B. D.; Goksu, B. B.; Verma, U.; Sanjeevaiah, A.; Hsiehchen, D.; et al. Racial/Ethnic Disparities and Survival Characteristics in Non-Pancreatic Gastrointestinal Tract Neuroendocrine Tumors. *Cancers* 2020, 12 (10), 2990.
22. Klugman, M.; Xue, X.; Ginsberg, M.; Cheng, H.; Rohan, T.; Hosgood, H. D., 3rd. Hispanics/Latinos in the Bronx Have Improved Survival in Non-Small Cell Lung Cancer Compared with Non-Hispanic Whites. *J Racial Ethn Health Disparities* 2020, 7 (2), 316-326. <https://doi.org/10.1007/s40615-019-00660-2> From NLM.
23. Gandhi, S.; Ballman, K.V.; Holmes, M.D.; Visvanathan, K.; Symington, B.; Carvan, M.; Matyka, C.; Weiss, A.; Winer, E.P.; Razaq, W.; Carey, L.A.; Partridge, A.H.; Chen, W.Y. Association of higher baseline stress and pain with clinical outcomes: Secondary analysis from Alliance A011502. In *Quality Care/Health Services Research, Proceedings of the ASCO Annual Meeting, Chicago, U.S.A, 2024; J Clin Oncol* 42, 2024; suppl 16; abstr 11016

24. Carethers, J. M. Racial and ethnic disparities in colorectal cancer incidence and mortality. *Adv Cancer Res* 2021, 151, 197-229. <https://doi.org/10.1016/bs.acr.2021.02.007> From NLM.
25. Hao, S.; Snyder, R. A.; Irish, W.; Parikh, A. A. Association of race and health insurance in treatment disparities of colon cancer: A retrospective analysis utilizing a national population database in the United States. *PLoS Med* 2021, 18 (10), e1003842. <https://doi.org/10.1371/journal.pmed.1003842> From NLM.
26. Zaki, T. A.; Liang, P. S.; May, F. P.; Murphy, C. C. Racial and Ethnic Disparities in Early-Onset Colorectal Cancer Survival. *Clin Gastroenterol Hepatol* 2023, 21 (2), 497-506.e493. <https://doi.org/10.1016/j.cgh.2022.05.035> From NLM.
27. Ko, N. Y.; Hong, S.; Winn, R. A.; Calip, G. S. Association of Insurance Status and Racial Disparities With the Detection of Early-Stage Breast Cancer. *JAMA Oncology* 2020, 6 (3), 385-392. <https://doi.org/10.1001/jamaoncol.2019.5672> (accessed 7/13/2024).
28. Dickerson, J. C.; Ragavan, M. V.; Parikh, D. A.; Patel, M. I. Healthcare delivery interventions to reduce cancer disparities worldwide. *World J Clin Oncol* 2020, 11 (9), 705-722. <https://doi.org/10.5306/wjco.v11.i9.705> From NLM.
29. Hollis, R. H.; Chu, D. I. Healthcare Disparities and Colorectal Cancer. *Surg Oncol Clin N Am* 2022, 31 (2), 157-169. <https://doi.org/10.1016/j.soc.2021.11.002> From NLM.
30. Kale, S.; Hirani, S.; Vardhan, S.; Mishra, A.; Ghode, D. B.; Prasad, R.; Wanjari, M. Addressing Cancer Disparities Through Community Engagement: Lessons and Best Practices. *Cureus* 2023, 15 (8), e43445. <https://doi.org/10.7759/cureus.43445> From NLM.
31. Richardson-Parry, A.; Baas, C.; Donde, S.; Ferraiolo, B.; Karmo, M.; Maravic, Z.; Münter, L.; Ricci-Cabello, I.; Silva, M.; Tinianov, S.; et al. Interventions to reduce cancer screening inequities: the perspective and role of patients, advocacy groups, and empowerment organizations. *Int J Equity Health* 2023, 22 (1), 19. <https://doi.org/10.1186/s12939-023-01841-6> From NLM.
32. Puneekar, S. R.; Kaakour, D.; Masri-Lavine, L.; Hajdu, C.; Newman, E.; Becker, D. J. Characterization of a Novel Entity of G3 (High-grade Well-differentiated) Colorectal Neuroendocrine Tumors (NET) in the SEER Database. *Am J Clin Oncol* 2020, 43 (12), 846-849. <https://doi.org/10.1097/coc.0000000000000761> From NLM.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.