

Negative Speed of Light, and the Imaginary Set of Base Planck Units

Szymon Łukaszyk*

Łukaszyk Patent Attorneys, ul. Głowackiego 8, 40-052 Katowice, Poland

The reflectance R of monolayer graphene for the normal incidence of electromagnetic radiation is known to be remarkably defined only by π and the fine-structure constant α . It is shown in this paper that the reflectance (or the sum of transmittance and absorptance) of monolayer graphene, expressed as a quadratic equation with respect to the fine-structure constant α must unsurprisingly introduce the 2nd fine-structure constant α_2 , as the 2nd root of this equation, as well as two π -like constants for a convex and a saddle surface. It turns out that this 2nd fine-structure constant is negative, and the sum of its reciprocal with the reciprocal of the fine-structure constant α is independent of the reflectance value R and remarkably equals $-\pi$. Particular algebraic definition of the fine-structure constant $\alpha^{-1} = 4\pi^3 + \pi^2 + \pi \approx 137.036$, containing the free π term and agreeing with the physical definition of this dimensionless constant to the 5th significant digit, when introduced to this sum, yields $\alpha_2^{-1} = -4\pi^3 - \pi^2 - 2\pi \approx -140.178$. Assuming universal validity of the physical definition of α , α_2 defines the negative speed of light in vacuum c_n . The average of this speed c_n and the speed of light in vacuum c is in the range of the Fermi velocity (10^6 m/s). Furthermore, the 2nd negative fine-structure constant α_2 alone introduces the imaginary set of base Planck units.

Keywords: Planck units; the fine-structure constant; speed of light in vacuum

I. INTRODUCTION

Numerous publications provide Fresnel coefficients for the normal incidence of electromagnetic radiation (EMR) on monolayer graphene (MLG), which are remarkably defined only by π and the fine-structure constant α having the reciprocal

$$\alpha^{-1} = \left(\frac{q_P}{e}\right)^2 = \frac{4\pi\epsilon_0\hbar c}{e^2} \approx 137.036, \quad (1)$$

where e is the elementary charge, q_P is the Planck charge, ϵ_0 is vacuum permittivity, \hbar is the reduced Planck constant, and c is the speed of light in vacuum.

Transmittance (T) of MLG

$$T = \frac{1}{\left(1 + \frac{\pi\alpha}{2}\right)^2} \approx 97.746\% \quad (2)$$

for normal EMR incidence was derived from the Fresnel equation in the thin-film limit [1] (Eq. 3), whereas spectrally flat absorptance (A) $A \approx \pi\alpha \approx 2.3\%$ was reported [2, 3] for photon energies between about 0.5 and 2.5 eV. T was related to reflectance (R) [4] (Eq. 53) as $R = \pi^2\alpha^2 T/4$, i.e.,

$$R = \frac{\frac{1}{4}\pi^2\alpha^2}{\left(1 + \frac{\pi\alpha}{2}\right)^2} \approx 0.013\%, \quad (3)$$

The above formulas for T and R , as well as the formula for the absorptance

$$A = \frac{\pi\alpha}{\left(1 + \frac{\pi\alpha}{2}\right)^2} \approx 2.241\%, \quad (4)$$

were also derived [5] (Eqs. 29-31) based on the thin film model (setting $n_s = 1$ for substrate).

The sum of transmittance (2) and the reflectance (3) at normal EMR incidence on MLG was also derived [6] (Eq. 4a) as

$$\begin{aligned} T + R &= 1 - \frac{4\sigma\eta}{4 + 4\sigma\eta + \sigma^2\eta^2 + k^2\chi^2} \\ &= \frac{1 + \frac{1}{4}\pi^2\alpha^2}{\left(1 + \frac{\pi\alpha}{2}\right)^2} \approx 97.759\%, \end{aligned} \quad (5)$$

where

$$\eta = \frac{4\pi\alpha\hbar}{e^2} = \frac{1}{\epsilon_0 c} \quad (6)$$

is the impedance of vacuum, $\sigma = e^2/(4\hbar) = \pi\alpha/\eta$ is the MLG conductivity [7], and $\chi = 0$ is the electric susceptibility of vacuum.

These coefficients are thus well-established theoretically and experimentally confirmed [1–3, 6, 8, 9].

As a consequence of the conservation of energy

$$(T + A) + R = 100\%. \quad (7)$$

In other words, the transmittance in the Fresnel equation describing the reflection and transmission of EMR at normal incidence on a boundary between different optical media is, in the case of the 2-dimensional (boundary) of MLG, modified to include its absorption.

The paper is structured as follows. Section II shows that Fresnel coefficients for the normal incidence of EMR on MLG introduce the second, negative fine-structure constant α_2 . Section III shows that this second fine-structure constant introduces the negative speed of light in vacuum c_n , which in turn introduces the imaginary set of base Planck units. Section IV shows that the negativity of the α_2 alone is sufficient to introduce the α_2 -set of Planck units. Section V presents Fresnel

* szymon@patent.pl

coefficients for the normal incidence of EMR on MLG expressed just by π . Section VI shows that Fresnel coefficients for the normal incidence of EMR on MLG introduce two π -like constants for a convex and a saddle surface having respectively positive and negative Gaussian curvatures. Section VII concludes and discusses the findings of this study.

II. THE SECOND FINE-STRUCTURE CONSTANT

The reflectance $R = 0.013\%$ (3) of MLG can be expressed as a quadratic equation with respect to α

$$\frac{1}{4}(R-1)\pi^2\alpha^2 + R\pi\alpha + R = 0, \quad (8)$$

having two roots with reciprocals

$$\alpha^{-1} = \frac{\pi - \pi\sqrt{R}}{2\sqrt{R}} \approx 137.036, \quad \text{and} \quad (9)$$

$$\alpha_2^{-1} = \frac{-\pi - \pi\sqrt{R}}{2\sqrt{R}} \approx -140.178. \quad (10)$$

Therefore, the quadratic equation (8) introduces the second, negative fine-structure constant α_2 .

The sum of the reciprocals of these fine-structure constants (9) and (10)

$$\alpha^{-1} + \alpha_2^{-1} = \frac{\pi - \pi\sqrt{R} - \pi - \pi\sqrt{R}}{2\sqrt{R}} = -\pi, \quad (11)$$

is remarkably independent of the reflectance R . The same result can be obtained for the sum of T and A , as shown in Appendix .

Furthermore, this result is intriguing in the context of a peculiar algebraic definition of the fine-structure constant [10]

$$\alpha^{-1} = 4\pi^3 + \pi^2 + \pi \approx 137.036 \quad (12)$$

that contains a *free* π term and agrees with the physical definition (1) of α to the 5th significant digit. Therefore, using Equations (11) and (12), we can express the negative reciprocal of the 2nd fine-structure constant α_2^{-1} that emerged in the quadratic equation (8) also as a function of π only

$$\alpha_2^{-1} = -\pi - \alpha_1^{-1} = -4\pi^3 - \pi^2 - 2\pi \approx -140.178. \quad (13)$$

This result supports the validity of the algebraic definition (12).

But how can this negative value be interpreted physically?

III. NEGATIVE SPEED OF LIGHT AND THE α_2 -SET OF PLANCK UNITS BASED ON IT

Almost all physical constants of $\alpha = e^2/(4\pi\epsilon_0\hbar c)$ in the physical definition of the fine-structure constant (1) are posi-

tive¹, whereas the charge e is squared. Only the velocity can be negative, as it is a *directional* quantity. Therefore, if

$$\alpha^{-1} = \frac{\pi - \pi\sqrt{R}}{2\sqrt{R}} = \frac{4\pi\epsilon_0\hbar c}{e^2} \approx 137.036, \quad (14)$$

then

$$\alpha_2^{-1} = \frac{-\pi - \pi\sqrt{R}}{2\sqrt{R}} = \frac{4\pi\epsilon_0\hbar c_n}{e^2} \approx -140.178, \quad (15)$$

where c_n is the negative speed of light in vacuum that, using Equations (11) with (1) and (15), amounts

$$\begin{aligned} \frac{4\pi\epsilon_0\hbar c}{e^2} + \frac{4\pi\epsilon_0\hbar c_n}{e^2} &= -\pi \\ c_n &= -\frac{e^2}{4\epsilon_0\hbar} - c \approx -3.066653 \times 10^8 \text{ [m/s]}, \end{aligned} \quad (16)$$

which is greater than the speed of light in vacuum c in modulus, whereas their average

$$\frac{c + c_n}{2} \approx -3.436417 \times 10^6 \text{ [m/s]} \quad (17)$$

is in the range of the Fermi velocity.

Furthermore, we can express the $e^2/(4\epsilon_0\hbar)$ term in (16) using the impedance of vacuum (6) as $c_n = -c(\alpha\pi + 1)$ which, using the algebraic definitions of α (12) and α_2 (13), yields the following relation between the speed of light in vacuum c , negative speed of light c_n , the fine-structure constant α and the negative fine-structure constant α_2

$$c\alpha = c_n\alpha_2 \quad (=v_e), \quad (18)$$

where v_e is the electron's velocity at the first circular orbit in the Bohr model of the atom.

The concept of the negative speed of light in vacuum c_n might seem to be a *nonsense* or unsettling for the reader. If this is the case, the reader is advised to skip the subsequent Section IV.

The negative speed of light in vacuum c_n (16) introduces the imaginary set of base Planck units $\{q_{Pi}, \ell_{Pi}, m_{Pi}, t_{Pi}, T_{Pi}\}$ that redefined by square roots containing $c_n < 0$ raised to an odd (1, 3, 5) power become imaginary and bivalued²:

$$\begin{aligned} q_{Pi} &= \pm \sqrt{4\pi\epsilon_0\hbar c_n} = \pm q_P \sqrt{\frac{\alpha}{\alpha_2}} \\ &\approx \pm i1.8969 \times 10^{-18} \text{ [C]} \quad (> q_P), \end{aligned} \quad (19)$$

$$\begin{aligned} \ell_{Pi} &= \pm \sqrt{\frac{\hbar G}{c_n^3}} = \pm \ell_P \sqrt{\frac{\alpha_2^3}{\alpha^3}} \\ &\approx \pm i1.5622 \times 10^{-35} \text{ [m]} \quad (< \ell_P), \end{aligned} \quad (20)$$

¹ Vacuum permittivity ϵ_0 is the value of the absolute dielectric permittivity of classical vacuum. Thus, ϵ_0 cannot be negative. The Planck constant h is the uncertainty principle parameter. Thus, it cannot be negative; negative probabilities do not seem to withstand Occam's razor.

² Base Planck units themselves admit negative values as negative square roots.

$$m_{Pi} = \pm \sqrt{\frac{\hbar c_n}{G}} = \pm m_P \sqrt{\frac{\alpha}{\alpha_2}} \quad (21)$$

$$\approx \pm i 2.2012 \times 10^{-8} \text{ [kg]} \quad (> m_P),$$

$$t_{Pi} = \pm \sqrt{\frac{\hbar G}{c_n^5}} = \pm t_P \sqrt{\frac{\alpha_2^5}{\alpha^5}} \quad (22)$$

$$\approx \pm i 5.0942 \times 10^{-44} \text{ [s]} \quad (< t_P),$$

$$T_{Pi} = \pm \sqrt{\frac{\hbar c_n^5}{G k_B^2}} = \pm T_P \sqrt{\frac{\alpha^5}{\alpha_2^5}} \quad (23)$$

$$\approx \pm i 1.4994 \times 10^{32} \text{ [K]} \quad (> T_P),$$

and can be expressed, using (18), in terms of base Planck units ℓ_P , m_P , t_P , and T_P ³.

Both charge $\alpha_2 q_{Pi}^2 = \alpha q_P^2$ (19) and mass $\alpha_2 m_{Pi}^2 = \alpha m_P^2$ (21) are related with base Planck units in the same way. It is not surprising: both Coulomb's law and Newton's law of universal gravitation are inverse-square laws. We also note that the relations between time (22) and temperature α_2 Planck units (23) are inverted: $\alpha^5 t_{Pi}^2 = \alpha_2^5 t_P^2$, whereas $\alpha_2^5 T_{Pi}^2 = \alpha^5 T_P^2$. Furthermore, eliminating α and α_2 from (19), (21)-(23) yields the following relations

$$\frac{q_P^2}{m_P^2} = \frac{q_{Pi}^2}{m_{Pi}^2} = 4\pi\epsilon_0 G \quad (24)$$

and

$$T_P^2 t_P^2 = T_{Pi}^2 t_{Pi}^2 = \frac{\hbar^2}{k_B^2}, \quad (25)$$

independent on the velocity of light.

Planck units derived from imaginary base units (20)-(23) are not imaginary in general.

The α_2 Planck volume

$$\ell_{Pi}^3 = \pm \left(\frac{\hbar G}{c_n^3} \right)^{3/2} = \pm \ell_P^3 \sqrt{\frac{\alpha_2^9}{\alpha^9}} \quad (26)$$

$$\approx \pm i 3.8127 \times 10^{-105} \text{ [m}^3\text{]} \quad (< \ell_P^3),$$

the α_2 Planck momentum

$$p_{Pi} = \pm m_{Pi} c_n = \pm \sqrt{\frac{\hbar c_n^3}{G}} = \pm m_P c \sqrt{\frac{\alpha^3}{\alpha_2^3}} \quad (27)$$

$$\approx \pm i 6.7504 \text{ [kg m/s]} \quad (> m_P c),$$

the α_2 Planck energy

$$E_{Pi} = \pm m_{Pi} c_n^2 = \pm \sqrt{\frac{\hbar c_n^5}{G}} = \pm E_P \sqrt{\frac{\alpha^5}{\alpha_2^5}} \quad (28)$$

$$\approx \pm i 2.0701 \times 10^9 \text{ [J]} \quad (> E_P),$$

and the α_2 Planck acceleration

$$a_{Pi} = \pm \frac{c_n}{t_{Pi}} = \pm \sqrt{\frac{c_n^7}{\hbar G}} = \pm a_P \sqrt{\frac{\alpha^7}{\alpha_2^7}} \quad (29)$$

$$\approx \pm i 6.0198 \times 10^{51} \text{ [m/s}^2\text{]} \quad (> a_P),$$

are imaginary and bivalued.

However, the α_2 Planck force

$$F_{Pi} = \pm \frac{E_{Pi}}{\ell_{Pi}} = \pm \frac{c_n^4}{G} = \pm F_P \frac{\alpha^4}{\alpha_2^4} \quad (30)$$

$$\approx \pm 1.3251 \times 10^{44} \text{ [N]} \quad (> F_P),$$

and the α_2 Planck density

$$\rho_{Pi} = \pm \frac{m_{Pi}}{\ell_{Pi}^3} = \pm \frac{c_n^5}{\hbar G^2} = \pm \rho_P \frac{\alpha^5}{\alpha_2^5} \quad (31)$$

$$\approx \pm 5.7735 \times 10^{96} \text{ [kg/m}^3\text{]} \quad (> \rho_P),$$

are real and bivalued.

On the other hand, the α_2 Planck area

$$\ell_{Pi}^2 = \frac{\hbar G}{c_n^3} = \ell_P^2 \frac{\alpha_2^3}{\alpha^3} \approx -2.4406 \times 10^{-70} \text{ [m}^2\text{]} \quad (< \ell_P^2), \quad (32)$$

is strictly negative. According to the holographic principle, each bit of information is physically represented on the holographic boundary by the Planck area [11–13] (having a positive value), whereas these areas are triangular [14]⁴. Notably, out of all omnidimensional (i.e. present in all complex dimensions n) convex n -polytopes and n -balls, only n -simplices (i.e., triangles for $n = 2$) are bivalued, admitting both positive and negative volumes and surfaces [15].

IV. ALTERNATIVE DERIVATION OF THE α_2 -SET OF PLANCK UNITS WITHOUT THE NEGATIVE SPEED OF LIGHT

The α_2 -set of Planck units, discussed in the preceding section, can also be derived without the negative speed of light c_n (16).

Using the definition of the fine-structure constant $\alpha = e^2/q_P^2$ (1) for the negative α_2 (10) or (13), we see that it requires an

³ The notation $\ell_{Pi} (< \ell_P)$, for example, means that the absolute value of the imaginary Planck length (20) is lower than the absolute value of the Planck length $\ell_P = \sqrt{\hbar G/c^3}$.

⁴ This is also compatible with the Causal Dynamical Triangulation (CDT) approach, which does not assume a pre-existence of a dimensional space, but focuses on the evolution of the spacetime as such.

introduction of some imaginary Planck charge q_{Pi} , so that its square would yield a negative α_2

$$\frac{q_{Pi}^2}{e^2} = \alpha_2^{-1} \approx -140.178 < 0. \quad (33)$$

Thus, using the value of the elementary charge e in (33) and the value of the α_2 (10) or (13), the imaginary Planck charge (33) can be derived without the negative speed of light c_n as

$$q_{Pi} = \pm \sqrt{\frac{e^2}{\alpha_2}}. \quad (34)$$

Furthermore, the concept of the negative speed of light c_n is not required to derive the imaginary Planck mass, which, using (34), (24) and the values of the Planck mass and charge is

$$m_{Pi} = \pm \frac{q_{Pi} m_P}{q_P}. \quad (35)$$

In the derivation of (35), we have used the relation (24) that was derived on the basis of the negative speed of light, but this relation, as such, is independent of any particular value of the speed of light, be it positive or negative.

Knowing that the Planck charge is $q_P = \sqrt{4\pi\epsilon_0\hbar c}$ we can solve it for the speed of light to find $c = q_P^2/(4\pi\epsilon_0\hbar)$ and introduce it to the Planck force $F_P = \pm c^4/G$ making it is independent on the speed of light.

This relation must be valid also for the α_2 Planck force that therefore can be derived without the negative speed of light c_n and using the imaginary Planck charge (34) as

$$F_{P2} = \pm \frac{1}{G} \left(\frac{q_{Pi}^2}{4\pi\epsilon_0\hbar} \right)^4. \quad (36)$$

Knowing the value of the α_2 Planck force (36) and mass (35) we can derive the α_2 Planck acceleration

$$a_{Pi} = \pm \frac{F_{P2}}{m_{Pi}}, \quad (37)$$

and the remaining α_2 Planck units, using the imaginary Planck charge (34) instead of the negative speed of light c_n .

V. ALGEBRAIC FRESNEL COEFFICIENTS FOR THE NORMAL INCIDENCE OF EMR ON MLG

With algebraic definitions of α (12) and α_2 (13) transmittance T (2), reflectance R (3) and absorptance A (4) of MLG for normal EMR incidence can be expressed just by π .

For $\alpha^{-1} = 4\pi^3 + \pi^2 + \pi$ (12) they become

$$T(\alpha) = \frac{4(4\pi^2 + \pi + 1)^2}{(8\pi^2 + 2\pi + 3)^2} \approx 97.746\%, \quad (38)$$

$$A(\alpha) = \frac{4(4\pi^2 + \pi + 1)}{(8\pi^2 + 2\pi + 3)^2} \approx 2.241\%, \quad (39)$$

while for $\alpha_2^{-1} = -4\pi^3 - \pi^2 - 2\pi$ (13) they become

$$T(\alpha_2) = \frac{4(4\pi^2 + \pi + 2)^2}{(8\pi^2 + 2\pi + 3)^2} \approx 102.279\%, \quad (40)$$

$$A(\alpha_2) = -\frac{4(4\pi^2 + \pi + 2)}{(8\pi^2 + 2\pi + 3)^2} \approx -2.292\%, \quad (41)$$

with

$$R(\alpha) = R(\alpha_2) = \frac{1}{(8\pi^2 + 2\pi + 3)^2} \approx 0.013\%. \quad (42)$$

Obviously $(T(\alpha) + A(\alpha)) + R(\alpha) = (T(\alpha_2) + A(\alpha_2)) + R(\alpha_2) = 1$ as required by the law of conservation of energy (7), whereas each conservation law is associated with a certain symmetry, as asserted by Noether's theorem. Nonetheless, physical interpretation of $T(\alpha_2) > 1$ and $A(\alpha_2) < 0$ requires further research. $A(\alpha) > 0$ implies a *sink*, whereas $A(\alpha_2) < 0$ implies a *source*, whereas the opposite holds true for the transmittance T , as illustrated schematically in Fig 1. Perhaps, the negative absorptance and transmittance exceeding 100% for α_2 (10) or (13) could be explained in terms of graphene spontaneous emission.




Figure 1. Illustration of the concepts of negative absorptance and excessive transmittance of EMR under normal incidence on MLG.

VI. FRESNEL COEFFICIENTS FOR THE NORMAL INCIDENCE OF EMR ON MLG INTRODUCE TWO CIRCUMFERENCE TO DIAMETER RATIOS FOR CONVEX AND FOR SADDLE SURFACE

The quadratic equation (8) describing the reflectance R of MLG under normal incidence of EMR (or alternatively (A.1)) can also be solved for π yielding two roots

$$\pi_{R\alpha_1} = \frac{2\sqrt{R}}{\alpha_*(1 - \sqrt{R})}, \quad \text{and} \quad (43)$$

$$\pi_{R\alpha_2} = \frac{-2\sqrt{R}}{\alpha_*(1 + \sqrt{R})}, \quad (44)$$

dependent on R and α_* , where α_* indicates α or α_2 . This can be further evaluated, using the algebraic definition of R (42) (which is the same for both α and α_2), yielding four, yet only three distinct, possibilities

$$\pi_{R\alpha_1} = -\pi \frac{4\pi^2 + \pi + 1}{4\pi^2 + \pi + 2} = -\pi(\alpha_2\pi + 1) \approx -3.0712, \quad (45)$$

$$\pi_{R\alpha_2} = \pi_{R\alpha_1} = \pi \approx 3.1416, \quad \text{and} \quad (46)$$

$$\pi_{R\alpha_2} = -\pi \frac{4\pi^2 + \pi + 2}{4\pi^2 + \pi + 1} = -\pi(\alpha\pi + 1) \approx -3.2136. \quad (47)$$

The absolute value of $\pi_{R\alpha_1}$ (45) corresponds to a convex surface having a positive Gaussian curvature, whereas the absolute value of $\pi_{R\alpha_2}$ (47) - to a negative Gaussian curvature. Their product $\pi_{R\alpha_1}\pi_{R\alpha_2} = \pi^2$. It remains to be found, whether each of them describes the ratio of circumference of a circle drawn on the respective surface to its diameter (π_c) or the ratio of the area of this circle to the square of its radius (π_a). These definitions produce different results on curved surfaces, whereas $\pi_a > \pi_c$ on convex surfaces, while $\pi_a < \pi_c$ on saddle surfaces [16].

VII. CONCLUSIONS

We have shown that the reflectance of graphene under the normal incidence of electromagnetic radiation (EMR), expressed as the quadratic equation with respect to the fine-structure constant α must introduce the 2nd negative fine-structure constant α_2 .

The sum of the reciprocal of this 2nd fine-structure constant α_2 with the reciprocal of the fine-structure constant α (1) is independent of the reflectance value R and remarkably equals simply $-\pi$.

Particular algebraic definition of the fine-structure constant $\alpha^{-1} = 4\pi^3 + \pi^2 + \pi$ (12), containing the free π term, when introduced to this sum, yields $\alpha_2^{-1} = -4\pi^3 - \pi^2 - 2\pi < 0$.

Assuming universal validity of the physical definition of the fine-structure constant α (1), the 2nd fine-structure constant α_2

(13) defines the negative speed of light c_n (16), the average of which and the speed of light in vacuum is in the range of the Fermi velocity (10^6 m/s).

Furthermore, the negative fine-structure constant α_2 alone introduces the imaginary set of five base Planck units (19)-(23), whereas mass and time units (24), as well as temperature and time units (25) are directly related to base Planck units.

The Planck volume (26), momentum (27), energy (28), and acceleration (29) derived from imaginary base units (20)-(23) are imaginary and bivalued, whereas the Planck force (30) and density (31) are real and bivalued.

The square of the imaginary Planck length (20) introduces the negative Planck area (32). If Planck areas correspond to bits of information [11-13] and if they are triangular [14], then the fact that among n -balls, n -cubes, n -orthoplices, and n -simplices only the latter admit negative volumes and surfaces [15] for any complex dimension n , suggests a physical interpretation for this negative Planck area (32).

The findings of this study, in particular the meaning of the relation (25), and the negative ratios $\pi_{R\alpha_1}$ (45) and $\pi_{R\alpha_2}$ (47), require further research in the context of emergent dimensionality [14, 15, 17, 18].

In the context of the results of this study, monolayer graphene, a truly 2-dimensional material with no thickness⁵, is a *keyhole* to other, unperceived [14], dimensionalities. Graphene history is also instructive. Discovered in 1947 [20], graphene was long considered *academic material* until it was eventually pulled in 2004 [21] from graphite by means of ordinary Scotch tape. These fifty-seven years, along with twenty-nine years (1935-1964) between the condemnation of quantum theory as *incomplete* [22] and Bell's mathematical theorem [23] asserting that it is not true, and the fifty-eight years (1964-2022) between the formulation of this theorem and this year's Nobel prize in physics for its experimental *loophole-free* confirmation, should remind us that Max Planck, the genius who discovered Planck units, has also discovered Planck's principle.

ACKNOWLEDGMENTS

I truly thank my wife for her support when this research [24, 25] was conducted.

Appendix: Other Quadratic Equations

The quadratic equation for the sum of transmittance (2) and absorptance (4), putting $C_{TA} \doteq T + A$, is

$$\frac{1}{4}C_{TA}\pi^2\alpha^2 + (C_{TA} - 1)\pi\alpha + (C_{TA} - 1) = 0, \quad (\text{A.1})$$

⁵ Thickness of MLG is reported [19] as 0.37 nm with other reported values up to 1.7 nm. However, taking into account that 0.335 nm is the established inter-layer distance and thus the thickness of bilayer graphene, these results do not seem credible.

and has two roots with reciprocals

$$\alpha^{-1} = \frac{C_{TA}\pi}{2(1 - C_{TA} + \sqrt{1 - C_{TA}})} \approx 137.036, \quad (\text{A.2})$$

and

$$\alpha_2^{-1} = \frac{C_{TA}\pi}{2(1 - C_{TA} - \sqrt{1 - C_{TA}})} \approx -140.178, \quad (\text{A.3})$$

whereas their sum $\alpha^{-1} + \alpha_2^{-1} = -\pi$ is also independent of T and A .

Other quadratic equations do not feature this property. For example, the sum of $T+R$ (5) expressed as the quadratic equa-

tion and putting $C_{TR} \doteq T + R$, is

$$\frac{1}{4}(C_{TR} - 1)\pi^2\alpha^2 + C_{TR}\pi\alpha + (C_{TR} - 1) = 0, \quad (\text{A.4})$$

and has two roots with reciprocals

$$\alpha^{-1} = \frac{\pi(C_{TR} - 1)}{-2C_{TR} + 2\sqrt{2C_{TR} - 1}} \approx 137.036, \quad (\text{A.5})$$

and

$$\alpha_{TR}^{-1} = \frac{\pi(C_{TR} - 1)}{-2C_{TR} - 2\sqrt{2C_{TR} - 1}} \approx 0.0180, \quad (\text{A.6})$$

whereas their sum

$$\alpha_{TR_1}^{-1} + \alpha_{TR_2}^{-1} = \frac{-\pi C_{TR}}{C_{TR} - 1} \approx 137.054 \quad (\text{A.7})$$

is dependent on T and R .

-
- [1] A. B. Kuzmenko, E. van Heumen, F. Carbone, and D. van der Marel, "Universal dynamical conductance in graphite," *Physical Review Letters*, vol. 100, p. 117401, Mar. 2008. arXiv:0712.0835 [cond-mat].
- [2] K. F. Mak, M. Y. Sfeir, Y. Wu, C. H. Lui, J. A. Misewich, and T. F. Heinz, "Measurement of the Optical Conductivity of Graphene," *Physical Review Letters*, vol. 101, p. 196405, Nov. 2008.
- [3] R. R. Nair, P. Blake, A. N. Grigorenko, K. S. Novoselov, T. J. Booth, T. Stauber, N. M. R. Peres, and A. K. Geim, "Universal Dynamic Conductivity and Quantized Visible Opacity of Suspended Graphene," *Science*, vol. 320, pp. 1308–1308, June 2008. arXiv:0803.3718 [cond-mat].
- [4] T. Stauber, N. M. R. Peres, and A. K. Geim, "Optical conductivity of graphene in the visible region of the spectrum," *Physical Review B*, vol. 78, p. 085432, Aug. 2008.
- [5] X. Wang and B. Chen, "Origin of Fresnel problem of two dimensional materials," *Scientific Reports*, vol. 9, p. 17825, Dec. 2019.
- [6] M. Merano, "Fresnel coefficients of a two-dimensional atomic crystal," *Physical Review A*, vol. 93, p. 013832, Jan. 2016.
- [7] T. Ando, Y. Zheng, and H. Suzuura, "Dynamical Conductivity and Zero-Mode Anomaly in Honeycomb Lattices," *Journal of the Physical Society of Japan*, vol. 71, pp. 1318–1324, May 2002.
- [8] S.-E. Zhu, S. Yuan, and G. C. A. M. Janssen, "Optical transmittance of multilayer graphene," *EPL (Europhysics Letters)*, vol. 108, p. 17007, Oct. 2014.
- [9] I. G. Ivanov, J. U. Hassan, T. Iakimov, A. A. Zakharov, R. Yakimova, and E. Jánzén, "Layer-number determination in graphene on SiC by reflectance mapping," *Carbon*, vol. 77, pp. 492–500, Oct. 2014.
- [10] P. Varlaki, L. Nadai, and J. Bokor, "Number Archetypes in System Realization Theory Concerning the Fine Structure Constant," in *2008 International Conference on Intelligent Engineering Systems*, (Miami, FL), pp. 83–92, IEEE, Feb. 2008.
- [11] J. D. Bekenstein, "Black Holes and Entropy," *Phys. Rev. D*, vol. 7, pp. 2333–2346, Apr 1973.
- [12] G. t. Hooft, "Dimensional Reduction in Quantum Gravity," 1993.
- [13] E. Verlinde, "On the origin of gravity and the laws of Newton," *Journal of High Energy Physics*, vol. 2011, p. 29, Apr. 2011.
- [14] S. Łukaszyk, *Black Hole Horizons as Patternless Binary Messages and Markers of Dimensionality*. Nova Science Publishers, 2023.
- [15] S. Łukaszyk, "On the Omnidimensional Convex Polytopes and n-Balls in Negative, Fractional and Complex Dimensions," preprint, MATHEMATICS & COMPUTER SCIENCE, Oct. 2022.
- [16] S. Mahajan, "Calculation of the pi-like circular constants in curved geometry," ResearchGate, Nov. 2013.
- [17] S. Łukaszyk, "Four Cubes," Feb. 2021. arXiv:2007.03782 [math].
- [18] S. Łukaszyk, "Novel Recurrence Relations for Volumes and Surfaces of n-Balls, Regular n-Simplices, and n-Orthoplices in Real Dimensions," *Mathematics*, vol. 10, p. 2212, June 2022.
- [19] H. Jussila, H. Yang, N. Granqvist, and Z. Sun, "Surface plasmon resonance for characterization of large-area atomic-layer graphene film," *Optica*, vol. 3, p. 151, Feb. 2016.
- [20] P. R. Wallace, "Erratum: The Band Theory of Graphite [Phys. Rev. 71, 622 (1947)]," *Physical Review*, vol. 72, pp. 258–258, Aug. 1947.
- [21] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, "Electric Field Effect in Atomically Thin Carbon Films," *Science*, vol. 306, pp. 666–669, Oct. 2004.
- [22] A. Einstein, B. Podolsky, and N. Rosen, "Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?," *Physical Review*, vol. 47, pp. 777–780, May 1935.
- [23] J. S. Bell, "On the Einstein Podolsky Rosen paradox," *Physica Physique Fizika*, vol. 1, pp. 195–200, Nov. 1964.
- [24] S. Łukaszyk, "A short note about graphene and the fine structure constant," 2020.
- [25] S. Łukaszyk, "A short note about the geometry of graphene," 2020.
- [26] C. J. Shearer, A. D. Slattery, A. J. Stapleton, J. G. Shapter, and C. T. Gibson, "Accurate thickness measurement of graphene," *Nanotechnology*, vol. 27, p. 125704, Mar. 2016.