

Article

Not peer-reviewed version

Antiproliferative Potential of *Eugenia uniflora* L. Leaf Essential Oil in Normal and Tumoral Human Colon Cells

[Ana G. Sánchez](#)^{*}, Macarena Menoni, Pamela Lombardo, [Eduardo Dellacassa](#), María Angélica Severi, Gabriela Ferragut, Beatriz Vignale, [Juan Cedano](#), María José Zuluaga, [Deborah J. Keszenman](#)^{*}

Posted Date: 29 May 2025

doi: 10.20944/preprints202505.2299.v1

Keywords: *Eugenia uniflora*; essential oil; colorectal cancer; CCD 841 CoN cells; Caco-2 cells; chemoprevention; chemotherapy; antiproliferative effect; cytotoxicity



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Antiproliferative Potential of *Eugenia uniflora* L. Leaf Essential Oil in Normal and Tumoral Human Colon Cells

Ana G. Sánchez ^{1,*}, Macarena Menoni ¹, Pamela Lombardo ², Eduardo Dellacassa ³,
María Angélica Severi ⁴, Gabriela Ferragut ⁴, Beatriz Vignale ⁵, Juan Cedano ⁶, María José Zuluaga ¹
and Deborah J. Keszenman ^{1,7,*}

¹ Grupo de Biofísicoquímica, Departamento de Ciencias Biológicas, CENUR Litoral Norte, Universidad de la República, Salto 50000, Uruguay

² Fitopatología, Departamento de Ciencias Biológicas, CENUR Litoral Norte, Universidad de la República, Salto 50000, Uruguay

³ Laboratorio de Biotecnología de Aromas, Departamento de Química Orgánica, Facultad de Química, Universidad de la República, Montevideo 11800, Uruguay

⁴ Laboratorio de Inmunología "Dr. Alberto Nieto", Departamento de Ciencias Biológicas, CENUR Litoral Norte, Universidad de la República, Salto 50000, Uruguay

⁵ Departamento de Producción Vegetal, Estación Experimental de Facultad de Agronomía, Salto 50000, Uruguay

⁶ Departament de Bioquímica i Biologia Molecular, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

⁷ Citizen Science Program, Biology Department, Bard College, Annandale-on-Hudson, NY 12504, USA

* Correspondence: agsanchez@unorte.edu.uy (A.G.S.); dkeszen@gmail.com (D.J.K.)

Abstract: Background/Objectives: Natural products are important in healthcare because of their accessibility and linkage to a healthy lifestyle. However, their effectiveness is uncertain due to insufficient scientific data. Cancer patients are frequent users of natural products to relieve symptoms or for chemoprevention. *Eugenia uniflora* leaf essential oil (EO), traditionally used for digestive disorders, emerges as a potential antineoplastic agent. We investigated the cytotoxic and antiproliferative effects of *E. uniflora* EO in human normal CCD 841 CoN and tumoral Caco-2 colonic cell lines. **Methods:** CCD 841 CoN and Caco-2 cells were exposed to different concentrations of *E. uniflora* EO, and the cytotoxicity was determined by MTT and Trypan Blue assays. Cell proliferation kinetics were analyzed at a low EO concentration, and the induction of DNA damage and oxidative stress was assessed by Comet and Cellular ROS assays. **Results:** Both cell lines exhibited cytotoxicity produced by the EO and decreased cell viability of the exposed cells and their progeny. CCD 841 CoN proliferation was impaired by low EO concentration, while the proliferation kinetics of the Caco-2 cells was modified. EO treatment induced variable DNA damage and oxidative stress depending on the cell line. **Conclusions:** Our results suggest that *E. uniflora* EO may prevent the proliferation of normal cells, inducing loss of viability. The EO produced cytotoxic and antiproliferative effects in tumoral cells by inducing DNA damage and increased oxidative stress. These effects support the consideration of *E. uniflora* EO (or its bioactive compounds) as a potential agent for the chemoprevention and treatment of colorectal cancer.

Keywords: *Eugenia uniflora*; essential oil; colorectal cancer; CCD 841 CoN cells; Caco-2 cells; chemoprevention; chemotherapy; antiproliferative effect; cytotoxicity

1. Introduction

Natural products have played a vital role in healthcare across different cultures throughout history. Today, nearly 80% of the global population depends on traditional medicine and natural products to meet their health needs. These resources are often preferred in developing countries because of their accessibility, affordability, and connection to local cultural practices. In contrast, in developed countries, their use is linked to a healthy lifestyle and a greater concern for environmental sustainability [1–3].

Nowadays, a wide variety of natural products and herbal medicines are available over the counter in many countries, giving rise to a growing global market for these products. Some are used as potential therapeutic agents for diverse diseases, and others are consumed in preventive medicine. However, many commercialized products are unregulated, and their effectiveness is uncertain due to insufficient scientific data. Patients with certain pathological conditions, such as cancer, are more frequent users of herbal medicines to relieve symptoms, ameliorate adverse effects from radiotherapy or chemotherapy, improve quality of life, and as chemopreventive agents to prevent further recurrence or metastasis. Nonetheless, observational studies have indicated that the concomitant use of natural products with cancer therapies may negatively interact, producing adverse side effects by lowering or neutralizing the benefits of conventional treatments [1,2,4–7]. Although the amount of scientific data related to the components and biological effects of many natural products and the clinical uses and outcomes of herbal medicine has been increasing notoriously in the last 20 years, more research is still needed to identify and characterize the bioactive compounds and their benefits and risks of their use as disease prevention and treatment.

Cancer is one of the ten leading causes of morbidity and mortality worldwide. It constitutes a complex pathological and often multisystemic condition affecting any part of the human body. In 2022, almost 20 million new cases were diagnosed (including non-melanoma skin cancer), and 9.7 million deaths due to cancer were recorded globally [8]. Colorectal cancer (CRC), which includes colon, rectum, and anus, is the third most common cancer and ranks second in terms of mortality. Although stabilizing or declining trends of the incidence rates for all ages combined have been observed in high-income countries, a rise in CRC among younger adults has been reported [9,10]. Also, there is a striking, steady rise in the incidence rate in some regions of the world, such as Eastern Europe and South America [11,12]. Behavioral and dietary changes in lifestyles, especially occurring during early life or young adulthood, lead to an increase in the influence of risk factors such as increased consumption of red or processed meat, less fiber in the diet, and sedentarism [8,13].

In South America, numerous native plant species have been traditionally used for medicinal purposes [14–17]. Among them, those belonging to the Myrtaceae family are particularly noteworthy due to their use in traditional medicine by their antidiarrheal, antimicrobial, antioxidant, antirheumatic, and anti-inflammatory properties [14,18]. Within this family, *Eugenia uniflora* L., commonly known as Pitanga, is widely distributed in tropical and subtropical regions of northeastern Argentina, southern Brazil, and Uruguay [19,20]. Its fruits are globular berries, varying in color, with a bright appearance and a distinctive, intense flavor that ranges from sweet to slightly acidic. Pitanga fruits are rich in vitamins A and C, phosphorus, calcium, and iron [21,22]. Infusions, decoctions, and alcoholic extracts of the leaves are used in traditional medicine because of their diuretic, anti-inflammatory, carminative, and antidiarrheal effects [14]. The leaf essential oil (EO) has been reported as possessing antifungal, antibacterial, cytotoxic, and antiparasitic activities [21,23–31].

E. uniflora EO has been reported to contain bioactive compounds with potential antitumor properties, including antioxidant capacity, apoptotic activity, and anti-inflammatory effects [29,32,33]. SiHa (HPV 16-positive) human cervical cancer cells treated with *Eugenia uniflora* leaf aqueous crude extract exhibit decreased viability, apoptotic cell death, and reduced migration. However, the same treatment does not modify the viability of normal human lymphocytes [34]. In addition, other studies have reported cytotoxic effects of the EO in various cancer cell lines [24]. Since *E. uniflora* EO is frequently used as herbal medicine for digestive disorders and could emerge as a natural product candidate to provide bioactive compounds for colon cancer therapy and prevention, we studied antineoplastic properties of *E. uniflora* leaf EO in human normal CCD 841 CoN and

tumoral Caco-2 colonic cells lines by determining the effect on cell proliferation kinetics, cell viability and induction of DNA damage and oxidative stress. Our main results present evidence of the cytotoxic and antiproliferative effects of *E. uniflora* EO in both human normal CCD 841 CoN and tumoral Caco-2 cell lines, as shown by the increased lethality, loss of viability, and altered proliferation kinetics. In addition, variable levels of DNA damage and oxidative stress were observed, depending on the cell line. These findings support the recognition of *Eugenia uniflora* as a promising plant species with potential for chemoprevention and treatment of CRC. This contribution aims to be particularly relevant in the ongoing research of bioactive compounds with antineoplastic properties, offering new possibilities for biotechnological applications in cancer therapy and preventive medicine.

2. Materials and Methods

2.1. Herbal Material

Leaves of *Eugenia uniflora* were collected from the plant specimen VIII-7, located in the Introduction Garden of the Experimental Station of the Faculty of Agronomy (Universidad de la República) in Salto, Uruguay (GPS coordinates: 31°22'S 57°56'W). This specimen was selected to represent the plant population to extract the essential oil (EO). Botanical identification was carried out by the Botany Department of the Faculty of Agronomy, CENUR Litoral Norte, Salto. A voucher specimen was deposited in the herbarium of the “Museo Jardín Botánico Prof. Atilio Lombardo” in Montevideo, Uruguay, and identified with the catalog number *E. uniflora* MVJB 29158.

Four independent collections of herbal material were carried out during Spring 2018 (October-November) and Summer 2019 (February-March). It is worth mentioning that, in each collection, just enough leaves to yield the appropriate amount of EO were harvested to preserve the plant. This environmental aspect should be considered when using native plants as a source of potential nutraceutical or pharmacological agents since the uncontrolled extraction of any plant species' components can result in their disappearance.

After collection, the leaves were dispersed on a flat surface and allowed to dry at room temperature under shaded conditions for one week. After grinding, the plant material was weighed, and 5600 g was used.

2.2. Essential Oil Extraction and Solutions

The EO was extracted by hydrodistillation using a Clevenger-type apparatus. The obtained EO was stored in amber glass bottles at 4°C for later use and analysis. For the biological assays, the EO was diluted as follows. An initial stock solution was prepared by diluting the original extracted EO 1:10 in a 10% dimethyl sulfoxide (DMSO, Sigma, USA) solution in phosphate-buffered saline (PBS, Sigma-Aldrich, USA). The calculated concentration was based on the EO's mass and density (0.96 g/mL), and appropriate dilutions were made to obtain different concentrations, as indicated in Table 1.

Table 1. *E. uniflora* essential oil (EO) dilutions from the stock solution.

	Dilutions				
	1/2000	1/1000	1/500	1/200	1/100
[EO] (mg/mL)	0.66	1.32	2.64	6.60	13.2

2.3. Essential Oil Analysis: Gas Chromatography Coupled with Mass Spectrometry (GC-MS)

GC-MS analyses were conducted using a Shimadzu QP 2020 mass spectrometer (Shimadzu Corporation, Kyoto, Japan) equipped with reference libraries [35–37] and a BP 20 bonded fused-silica capillary column (25 m × 0.25 mm i.d.), coated with a 0.25-µm-thick layer of poly(ethylene glycol) (SGE, Ringwood, Australia). The experimental conditions were the same as those reported by

Dellacassa et al. [38,39]. The temperature program was as follows: 40°C (6 min), 40-180°C at 3°C/min, then to 220°C at 10°C/min, 220°C (20 min); injector temperature, 250°C; carrier gas, helium at 122.2 kPa (51.6 cm/s); injection mode, split; split ratio, 1:40; the sample volume injected was 1 µL of a 1:10 solution in dichloromethane; the interface temperature, 250°C; energy, 70 eV; acquisition mass range, 40-400 a.m.u.

The EO components were tentatively identified by comparing the mass spectra of the samples with the data system [35–37] and based on arithmetic retention indices (LRI) of either reference substances or literature values. Compounds were considered tentatively identified when experimental and reported LRI did not differ by more than 10 units and when the similarity between the mass spectrum of each chromatographic peak and spectrum of the mass spectral libraries was at least 80%. Whenever possible, MS identification was confirmed with authentic references. The relative amounts of components present in the EO were expressed as percentages of the peak area of total chromatograms. Values less than 0.01 were considered as statistically insignificant.

2.4. Cell Culture

The human normal colon cell line CCD 841 CoN was purchased directly from ATCC USA (CRL-1790), and cells were cultured in α -Minimum Essential Medium (α -MEM, Gibco, USA) supplemented with 10% fetal bovine serum (FBS, CAPRICORN Scientific, USA) in T25 flasks. Culture passages of CCD 841 CoN between +19 to +25 were used in the experiments. The human colon carcinoma cell line Caco-2 was kindly provided by Andrés Lisozaín (Laboratory of Virology, CENUR Litoral Norte, Salto, Universidad de la República, Uruguay). Cells were cultured in T25 flasks (Cellstar®) containing nutrient medium Dulbecco's Modified Eagle's Medium (DMEM, Gibco, USA) supplemented with 10% FBS. No antibiotic or antifungal supplements were added to the culture media. Both cell lines were maintained in a humidified incubator at 37°C, 5% CO₂, and a stable pH of 7.40. Subconfluent cultures (60 - 80 % confluency) were used in all the experiments.

Caco-2 cells were subcultured twice a week, and CCD 841 CoN cells every 6–8 days. Culture medium was replaced with fresh medium every 48 h for the Caco-2 and every 96 h for the CCD 841 CoN cells. Adherent cells were washed twice with 1X PBS at pH 7.40. Cell detachment was performed using a 0.25% Trypsin-EDTA solution (CAPRICORN Scientific, Germany) at 37°C for 3-5 min. Microscopic assessment of viable and nonviable cells was performed using a 0.4% Trypan Blue solution (TB, Sigma-Aldrich, USA) diluted in 1X PBS, and cell counts were recorded by a hemocytometer.

2.5. MTT Assay

CCD 841 CoN and Caco-2 cell cultures that showed greater than 95% cell viability were used. In 96-well plates, 3.0×10^4 cells/well of CCD 841 CoN and 1.6×10^4 cells/well of Caco-2 were seeded in a final volume of 100 µL of culture medium in triplicate, and plates were incubated at 37°C, 5% CO₂. After 24 h, *E. uniflora* EO was added at different concentrations, and cells were incubated for 14 h at 37°C and 5% CO₂. Cells in culture media without EO were used as untreated negative controls. Cells treated with 1% DMSO as solvent control, and blanks of culture medium, DMSO, and Pitanga extract were included in the 96-well plates. Three repeats of controls, blanks, and treatments were done.

After treatment, cells were washed with 1X PBS, and 100 µL of MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) stock solution (5 mg/mL in 1X PBS and 1,1 M glucose) was added to allow the formation of formazan crystals by viable cells, as described by Mosmann et al. [40]. After 4 h of incubation at 37°C and 5% CO₂ in the dark, the optical density (O.D.) was measured using an ELISA plate reader at a 540 nm wavelength. The survival fraction was calculated using the equation:

$$\text{Survival fraction (\%)} = \frac{\text{O.D. treated samples}}{\text{O.D. control samples}} \times 100$$

2.6. Trypan Blue Viability Assay

In 6-well plates, 5.0×10^4 cells/well were seeded in α -MEM + 10% FBS (CCD 841 CoN) or DMEM + 10% FBS (Caco-2), and incubated in a humidified atmosphere of 5 % CO₂ at 37°C. One 6-well plate was prepared for each time point. After 24 h, *E. uniflora* EO was added to each well at the following concentrations: 0, 0.66, 1.32, 2.64, 6.60, and 13.2 mg/mL. After 0, 4, 8, 12, 16, 20, 24, 28, and 32 h of incubation, cells were harvested by trypsinization and centrifuged. Cell pellets were resuspended in 1X PBS, and 100 μ L of the cell suspension was mixed with 10 μ L of 0.4% TB solution for counting in a hemocytometer [41]. At least three independent experiments were performed. The unstained cells were counted as viable cells, while the blue-stained cells were counted as dead. The percentage of cell viability was calculated according to the following equation:

$$\text{Viability (\%)} = \frac{\text{Number of viable cells}}{\text{Total cell number}} \times 100$$

2.7. Assessment of Cell Proliferation Kinetics

To determine the proliferation kinetics of CCD 841 CoN cell line, 2.0×10^5 cells/plate in α -MEM + 10% FBS were seeded in 60 mm Petri dishes (2 dishes per time point). Due to the slow growth of this cell line, the culture medium was replaced every 96 hours. To study Caco-2 cells, 60 mm Petri dishes (2 dishes per time point) were seeded with 6.4×10^4 cells/plate in DMEM + FBS and incubated in a humidified environment at 5% CO₂ and 37°C for the duration of the study. Culture medium was replaced every 48 h with fresh medium.

CCD 841 CoN cell proliferation was monitored for 17 days, and Caco-2 proliferation was observed for up to 14 days by tracking the increase in cell number as a function of time. At each time point, cells were harvested by washing and trypsinization. Cell pellets were resuspended in 3 mL of 1X PBS, and 100 μ L of this cell suspension was mixed with 10 μ L of 0.4% TB solution for counting viable cells.

To assess the effect of *E. uniflora*, the proliferation kinetics of both cell lines -CCD 841 CoN and Caco-2- treated with the EO were studied at a concentration of 0.66 mg/mL. The selection of the EO concentration was based on the results obtained from the MTT assay, which indicated the concentration that yields the lowest lethality.

The analysis of cell proliferation kinetics was performed by applying the following logistic model of cell population growth:

$$N(t) = \frac{N_{\max}}{1 + e^{-\ln 2 / DT (t - t_{1/2})}}$$

where $N(t)$ corresponds to the number of cells/mL quantified at time t ; N_{\max} corresponds to the number of cells/mL observed in the stationary phase of growth; DT corresponds to the cell doubling time; $t_{1/2}$ corresponds to the time it takes for the cell population to reach half of N_{\max} . The lag phase time (t_{lag}) represents the time required for the cell population to initiate proliferation [42].

2.8. DNA Damage Assay

DNA damage was evaluated in cultured cells of both cell lines using the single-cell gel electrophoresis, or Comet assay [43]. Briefly, control and treated samples containing 1.5×10^5 cells were embedded in 0.75% low-melting point agarose (Invitrogen, USA) and then plated onto a slide coated with 1% running agarose (SeaKem Agarose, Lonza, USA). Cell samples treated with Bleomycin 40 μ g/mL for 2 h were used as a positive control. Once solidified, the slides were treated with detergents and high ionic strength solutions to remove lipids, carbohydrates, and proteins, leaving only DNA molecules whose strands were separated due to the alkaline lysis buffer treatment (lysis buffer: 2.5 M NaCl, 100 mM EDTA, 10 mM Tris-HCl, $\text{NH}_2\text{C}(\text{CH}_2\text{OH})_3 \cdot \text{HCl}$ and NaOH to adjust the pH to 10, then 1% Triton-X-100 and 10% DMSO were added; Sigma, USA). The slides were placed in a horizontal electrophoresis tank with alkaline running buffer (0.3 M NaOH, 1 mM EDTA) at 4°C. The running parameters were 15 V and 280-300 mA (voltage: 1 V/cm) for 20 min. Once the electrophoresis was completed, the slides were placed in the neutralization buffer (0.4 M Tris, pH

7.50) and stained for 3 min with 6 mg/mL DAPI. Images were obtained using a Nikon Eclipse Ts2R inverted epifluorescence microscope with a 361-389/435 nm UV filter at 400X magnification.

2.9. Reactive Oxygen Species Detection Assay

CCD 841 CoN and Caco-2 cells were cultured in 96-well culture plates by adding 200 μ L/well of a 1.5×10^4 cell suspension in α -MEM + 10% FBS or DMEM + 10% FBS, pH 7.40, respectively. The cells were incubated at 37°C, 5% CO₂ for 24 h to allow adhesion. To detect Reactive Oxygen Species (ROS) in both cell lines, the Cellular ROS Assay Kit (ab186027, Abcam, USA) was used according to the manufacturer's recommended instructions. Treated cells were exposed to different concentrations of *E. uniflora* EO ($0 \leq [\text{EO}] \leq 13.2$ mg/mL) for 2 h. Hydrogen peroxide (200 μ M) was used as a positive control for ROS production. The presence of ROS in control and treated samples was quantified by measuring the increase in fluorescence at 540 nm using a microplate reader.

2.10. Statistical Analysis

Data from at least three independent experiments are presented as mean \pm Standard Error of the Mean (S.E.M.). Graphs were obtained using the statistical program SigmaPlot v.15.0, and where error bars were not shown, they were smaller than the size of the symbol. Significance calculations using the t-test were made using SigmaPlot. In all cases, statistically significant differences were considered when $p < 0.05$.

3. Results

3.1. Chemical Characterization of *E. uniflora* EO

The *E. uniflora* EO extracted by hydrodistillation was chemically analyzed by GC-MS. The average EO yield of the hydrodistillation process was 0.98 % (v/w). Forty-nine compounds were identified, representing 95.51% of the total, of which 12 were monoterpenes (4.21 %) and 33 sesquiterpenes (91.18 %) (Table 2). The main sesquiterpenes were hydroxycalamenene (41.04 %), δ -cadinene (18.73 %), and τ -muurolol (12.72 %).

Table 2. *E. uniflora* EO chemical compounds identified by GC-MS.

LRI ¹	Compound ²	%	LRI	compound	%
1082	Hexanal	0.01	1666	α -Humulene	0.07
1110	β -Pinene	0.01	1668	γ -Gurjunene	0.06
1122	Sabinene	0.01	1699	Bicyclogermacrene	0.02
1162	α -Terpinene	0.01	1682	β -Humulene	0.13
1167	α -Phellandrene	0.01	1684	trans- β -Guaiene	0.01
1169	Myrcene	1.10	1686	β -Chamigrene	3.71
1198	Limonene	0.25	1688	Zonarene	0.48
1209	β -Phellandrene	0.26	1689	γ -Muurolene	5.22
1239	2-Pentylfuran	0.01	1708	Germacrene D	0.22
1250	trans- β -Ocimene	0.71	1720	α -Cadinene	0.41
1234	cis- β -Ocimene	1.82	1722	Viridiflorene	0.71
1270	p-Cymene	0.03	1723	α -Muurolene	0.55
1282	Terpinolene	0.03	1746	cis-Muurola-3,5-diene	0.41
1351	1-Hexanol	0.01	1755	d-Cadinene	18.73
1351	allo-Ocimene	0.01	1806	Germacrene B	0.55
1380	(Z)-3-Hexen-1-ol	0.07	1808	trans-Calamenene	0.02
1460	α -Cubebene	0.06	1818	cis-Calamenene	0.27
1523	β -Bourbonene	0.03	1918	β -Calacorene	0.06

1600	Guaia-6,9-diene	0.03	1901	α -Colacorene	0.05
1626	β -Copaene	0.59	1930	Palustrol	0.01
1636	trans-Caryophyllene	0.23	2082	Globulol	0.54
1637	Aromadendrene	0.17	2089	Viridiflorol	3.85
1642	trans-Muurola-3,5-diene	0.17	2185	τ -Muurolol	12.72
1652	α -Guaiene	0.01	2325	Hydroxy calamenene	41.04
1658	γ -Elemene	0.05			
				Total	95.51

¹Linear retention index based on a series of n-hydrocarbons reported according to their elution order on Carbowax 20M; ²Identity tentatively assigned by comparing mass spectra with those obtained from the literature [35–37].

3.2. Microscopic Morphology of Colonic Cell Lines

Microscopic observation of the normal CCD 841 CoN cell culture under optimal conditions showed elongated cells that grew disorganized, resulting in a monolayer 7 days after seeding (168 h) (Figure 1A). Tumoral Caco-2 cells, on the other hand, exhibited a polygonal appearance, and the culture formed an organized monolayer 144 h after seeding (Figure 1B).




Figure 1. Microphotographs of normal CCD 841 CoN (A) and tumoral Caco-2 (B) cell cultures, grown and maintained under optimal conditions, captured at 20X magnification.

3.3. Evaluation of the Cytotoxic Effect of *E. uniflora* EO on Cell Lines

To study the cytotoxic effect of *E. uniflora* EO on the normal CCD 841 CoN and tumor Caco-2 colonic cell lines, the survival fraction to different concentrations of EO (0, 0.66, 1.32, 2.64, 6.60, and 13.2 mg/mL) was determined by the MTT assay (Figure 2).




Figure 2. Cell survival of colonic cell lines as a function of the concentration of *E. uniflora* EO. Dose-response survival curve of normal CCD 841 CoN (○) and tumor Caco-2 (▽) cells exposed for 14 h to different concentrations of EO in nutrient medium. The S.E.M. intervals are included in the figure.

The dose-response curve for the normal colon cell line CCD 841 CoN, presented in Figure 2 as an open circle, showed that the lowest concentration of EO tested ([EO] = 0.66 mg/mL) did not significantly modify the survival compared to the control. Cell lethality increased exponentially at higher concentrations, reaching about 1% survival fraction at [EO] = 2.64 mg/mL and remained unchanged thereafter. It should be noted that cell survival decreased 40-fold when the EO concentration doubled from 1.32 mg/mL to 2.64 mg/mL.

The survival curve for the tumoral cell line Caco-2 (Figure 2, triangle down) exhibited an exponential decrease with two components. The initial exponential component, characterized by the steepest slope, was observed up to [EO] = 2.64 mg/mL, where cell survival was 4.5%. At higher EO concentrations, the slope decreased, with cell survival dropping to 1% at the highest tested dose, [EO] = 13.2 mg/mL.

The MTT assay provided a snapshot of the cell lethality induced by a continuous 14-hour exposure to EO at the end of the treatment. However, it did not determine whether surviving cells might exhibit delayed lethality after treatment. To investigate the time course of the cytotoxic effects of the EO, cell viability was assessed over the exposure period to [EO] = 0.66 mg/mL, the lowest tested concentration. Briefly, cells were seeded in 6-well culture plates and exposed to this concentration for 32 hours. The numbers of viable and nonviable cells were recorded at regular intervals, and cell viability was calculated.

Figure 3A shows that the total cell count of the EO-treated Caco-2 culture decreased up to 14 h, followed by a transient 2-fold increase at 24 h, and then decreased again until the end of the assay. The cell viability, calculated at the same time points (Figure 3B), decreased as a function of time, reaching a minimum at 20 h, followed by an increase at 24 h, which coincided with the peak in cell number. After this time point, the cell viability, as the cell count, decreased.




Figure 3. Cell viability time course of the Caco-2 tumor cell line treated with 0.66 mg/mL of EO. (A) Total number of cells (viable and nonviable) as a function of time; (B) Percentage of cell viability calculated using the data presented in (A) at the same time points.

To test whether the decrease in cell viability observed in Caco-2 tumor cell cultures continuously exposed to [EO] = 0.66 mg/mL was due to effects on the cell progeny, the proliferation capability was evaluated after treatment removal by scoring the total number of cells. Cells were counted immediately after 24 h or 48 h EO treatments. Also, cell counts were scored after removing EO treatments and incubating cells under optimal culture conditions for 72 h. Figure 4 shows that the number of cells in the culture decreased 25% when the duration of the treatment increased from 24 to 48 h ($p = 0.03$). Cell counts after the post-treatment incubation without EO for 72 h showed a 2.6-fold increase for the 24 h EO treatment ($p = 0.006$) and 1.8-fold for the 48 h one ($p = 0.006$). The difference in proliferation observed with increasing exposure time to the EO was significant ($p = 0.008$). Therefore, the EO treatment duration affected the progeny of treated cells, decreasing their proliferation capacity.




Figure 4. Effect of the *E. uniflora* EO on the progeny of Caco-2 cells treated with [EO] = 0.66 mg/mL during 24 or 48 h. Cells were scored immediately after EO treatment (black bars) or after incubation for 72 h under optimal conditions without EO (white bars). Data from at least 3 independent experiments are included. Asterisks indicate significant differences between samples and treatments, $p < 0.05$.

After 48 h of EO treatment (0.66 mg/mL), the microscopic observation of both cell lines showed striking differences (Figure 5). While the CCD 841 CoN image shows cells with clearly altered morphology (compare Figures 1A–5A), the Caco-2 showed both long adherent and rounded cells in suspension (compare Figures 1B–5B).




Figure 5. Microphotographs of CCD 841 CoN (A) and Caco-2 (B) cells treated with 0.66 mg/mL of *E. uniflora* EO for 48 h (40X magnification).

3.4. Cell Proliferation Kinetics of CCD 841 CoN and Caco-2 Cell Lines

The potential antiproliferative effect of *E. uniflora* EO was investigated by assessing the proliferation kinetics of CCD 841 CoN normal colon cells and Caco-2 tumor cells without and with EO treatment.

As shown in Figure 6A,B, the proliferation kinetics of CCD 841 CoN and Caco-2 were studied under optimal conditions for 17 and 14 days, respectively, and the calculated parameters are presented in Table 3. The observed differences in the proliferation kinetics between the CCD 841 CoN and Caco-2 cell lines are consistent with the normal and tumoral characteristics, respectively. The kinetics of the Caco-2 tumor cell line, as compared to the normal CCD 841 CoN, presented a steeper slope of the exponential phase, indicating faster proliferation. The analysis of the proliferation kinetics parameters showed that the Caco-2 tumor cell line reached the stationary phase with a cell number 8.2 times higher than the normal cell line CCD 841 CoN. The half-time ($t_{1/2}$) values for Caco-2 and CCD 841 CoN were 168 h and 309.5 h, respectively. Regarding the t_{lag} , it was 5.5 times higher in the normal cell line than in the tumoral one.




Figure 6. Proliferation kinetics of the CCD 841 CoN (A) and the Caco-2 (B) under optimal growth conditions and the corresponding cell viability determined at each time point (C). The number of cells per mL (N) as a function of time for CCD 841 CoN was recorded up to 400 h due to slow growth. The grey lines in (A) and (B) correspond to the curve fitting based on the logistic growth model indicated in Materials and Methods. The dotted lines in (C) indicate 95 % confidence intervals for the normal colon cell line CCD 841 CoN (○) and the tumor cell line Caco-2 (▽). Mean values ± S.E.M. are included in the figures.

Table 3. Proliferation kinetics parameters of the CCD 841 CoN and the Caco-2 cell lines without and with treatment with *E. uniflora* EO ([EO] = 0.66 mg/mL). Means of at least 3 experiments ± S.E.M. are presented.

	N _{max} (cells/mL)	DT (h)	t _{1/2} (h)	t _{lag} (h)
Control CCD 841 CoN	1.0x10 ⁶ ± 1.5x10 ⁵	57.9 ± 8.9	309.5 ± 30.8	66.0 ± 3.0
Control Caco-2	8.2x10 ⁶ ± 2.8x10 ⁵	17.1 ± 3.0	168.0 ± 5.1	12.0 ± 0.6
Treated Cado-2	5.2x10 ⁶ ± 2.6x10 ⁵	60.0 ± 3.0	225.0 ± 11.3	0

Cell viability was monitored throughout the proliferation study using the TB staining method. Under optimal culture conditions, cell viability remained above 85% throughout the observation period in both cell lines. However, the Caco-2 tumor cell line exhibited higher cell viability (greater than 95%) than the normal cell line CCD 841 CoN (Figure 6C).

To study the effect of *E. uniflora* EO on cell proliferation, and based on the previous survival results, the lowest concentration of EO (0.66 mg/mL) that allowed the highest cell survival was selected to assess the proliferation kinetics. Treated Caco-2 cells showed proliferation up to 11 days, and afterward, the number of cells declined progressively (Figure 7A). A slower kinetics was evidenced by a lower maximum number of cells (N_{max}) reached before the decline phase, a 3.5-fold increase in DT, and an increase in t_{1/2} from 168.0 h to 225.0 h. It was striking that, despite a decrease in proliferation, the lag phase (t_{lag}) decreased to 0 (Table 3). Simultaneously with the quantification of the proliferation kinetics, Caco-2 cell viability was evaluated as a function of time (Figure 7B). Cell viability was not notably affected until 200 h of the proliferation study, after which it began to decline with a steep slope.




Figure 7. Proliferation kinetics (A) and cell viability (B) of the Caco-2 tumor colon cell line treated with 0.66 mg/mL of EO added to the culture medium. Mean values ± S.E.M. are included in the figures.

In the case of CCD 841 CoN, it was not possible to obtain a proliferation curve since immediately after the exposure to the EO, the cells presented nonviable characteristics, with notorious cell death observed after 48 h of incubation, making it not possible to perform subsequent counts (Figure 5B).

3.5. DNA Damage Produced by the *E. uniflora* EO

DNA damage induced by *E. uniflora* EO in CCD 841 CoN and Caco-2 cells was assessed using the Comet assay. Figure 8 shows DAPI-stained images of control and EO-treated cells of CCD 841 CoN (A and B, respectively) and Caco-2 (C and D, respectively). Cells treated with bleomycin, a radiomimetic antitumor antibiotic that produces single and double DNA strand breaks and base loss, thus inducing DNA fragmentation [44], were used as a positive control (Figure 8E). The image of treated CCD 841 CoN (Figure 8B) shows amorphous cell structures and inhomogeneous staining, characteristic of extensive cell destruction. In comparison, Figure 8D shows the images of comets formed by the migration of DNA fragments from the nucleus due to strand breaks produced by 0.66 mg/mL of EO in Caco-2 cells.




Figure 8. DNA damage produced by *E. uniflora* EO in the CCD 841 CoN and Caco-2 cell lines determined using the Comet assay. Microphotographs of CCD 841 CoN cells untreated (A) and treated with 0.66 mg/mL of *E. uniflora* EO for 14 h (B); Caco-2 cells untreated (C) and treated with 0.66 mg/mL of EO for 14 h (D). (E) Positive control: Caco-2 cells treated with 40 µg/mL of bleomycin.

3.6. Oxidative Stress Produced by *E. uniflora* EO

To investigate whether *E. uniflora* EO produces modifications in the level of oxidative stress that may be one possible molecular mechanism underlying the lethal and antiproliferative effects, the levels of ROS induced by treatment with different concentrations of EO were determined in CCD 841 CoN and Caco-2 cells at 2 h after treatment using the Cellular ROS Assay Kit. In the normal colon cell line CCD 841 CoN (Figure 9, white bars), the ROS levels increased significantly at the highest concentration of EO ($p = 0.0002$). At lower EO concentrations, no significant ROS level variation was observed. In the Caco-2 cell line, the ROS levels increased with the EO concentration (Figure 9, striped bars). Significantly higher fluorescence values were observed in the tumoral cell line compared to the CCD 841 CoN at all EO concentrations tested. Interestingly, the spontaneous levels of ROS ($[EO] = 0$ mg/mL) were similar in both normal and tumor cell lines ($p = 0.574$).




Figure 9. ROS levels induced in normal colonic cells CCD 841 CoN (white bars) and tumoral Caco-2 cells (striped bars) by treatment with different concentrations of *E. uniflora* EO for 2 h. Measurements were made 2 h post-treatment with EO. Single asterisks in each cell line indicate significant differences between untreated and treated samples. Double asterisks indicate significant differences in ROS levels between CCD 841 CoN and Caco-2 cell lines ($p < 0.05$).

4. Discussion

Cancer—in particular colorectal cancer—represents a significant global public health concern, contributing to high mortality rates, substantial morbidity, and considerable socioeconomic burdens [8,10]. This worldwide health challenge demands innovative research to develop new tools for prevention, rapid diagnosis, and effective treatment of the disease. Natural products have been used to treat diseases for centuries in various cultures. Nowadays, societal cultural changes are leading to an increased interest and appreciation for natural products, particularly in health and wellness. Also, a shift towards more natural and sustainable practices, including traditional healing systems and herbal remedies, has occurred, especially in developed countries. From the scientific point of view, natural products are well-recognized sources of bioactive compounds with various biomedical applications [45–47].

The native plant *Eugenia uniflora* (Pitanga), traditionally used in South America to treat various health-related issues, including gastrointestinal diseases, emerges as a good candidate to identify biological compounds with potential chemopreventive and chemotherapeutic properties. Several studies on *E. uniflora* leaf extracts have shown antimicrobial activity [48,49], antioxidant capacity [33], and a few lines of evidence have specifically reported possible anticarcinogenic effects [24,34]. To gain more insight into the use of *E. uniflora* as a natural herbal remedy for digestive disorders and to investigate the antineoplastic potential of *E. uniflora* leaf essential oil in colorectal cancer, the immediate and delayed cytotoxicity and the effect on cell proliferation were studied in normal CCD 841 CoN and tumoral Caco-2 colonic cell lines.

To evaluate the cytotoxic effects of the *E. uniflora* EO, the cell survival and viability were assessed by MTT assay and Trypan Blue staining in both CCD 841 CoN and Caco-2 cell lines upon treatment with different concentrations of the EO. The dose-response survival curve (Figure 2) shows that the *E. uniflora* EO produced lethal effects on both normal and tumoral cells in a concentration-dependent manner. In the case of the normal CCD 841 CoN cells, no significant lethality was observed at the lowest concentration ([EO] = 0.66 mg/mL). However, the cell viability of the cell culture was severely compromised, as observed upon microscopic inspection. The comparison of the microscopic image of CCD 841 CoN cells incubated under optimal conditions to that of EO-treated cells clearly showed

drastic cell morphological changes, including variable cell shape modifications, cell disruption, and nuclear condensation, which are typical indicators of apoptosis or necrosis and therefore of nonviable cells (compare Figures 1A–5A). This high cytotoxic effect of the EO in the normal human colonic cell line was unexpected since the MTT results showed no lethality. It should be noted that MTT assessment of cell health constitutes a snapshot of functioning mitochondria after 14 h of EO exposure. However, the microscopic observation was performed after incubating the cells with EO for 48 h. In vitro and in vivo studies have shown that *E. uniflora* EO does not affect normal cells, as demonstrated by the lack of viability loss in normal human lymphocytes [34] and also the absence of animal lethality after oral administration to rodents [24,29,33,50]. A possible explanation for the observed cytotoxicity in the present study could be that EO treatment was performed in proliferating non-confluent CCD 841 CoN cultures, where cells were actively cycling. Considering this interpretation, this result suggests that the possible underlying mechanism involved in Pitanga EO effects could be related to cell proliferation.

To confirm the lethality produced by low EO concentrations in the Caco-2 tumor cell line and to assess the time course of the cytotoxic effect, the cell viability was determined as a function of exposure time to [EO] = 0.66 mg/mL, using the Trypan Blue assay (Figure 3). An initial decrease in the cell count (Figure 3A) and viability (Figure 3B) was observed. It should be noted that the viability fell below the value at $t = 0$ after a brief increase. Based on this result, the effect of the EO treatment on the cell progeny was evaluated, assessing the proliferation capability of cells after 72 h in optimal conditions post-treatment removal. Figure 4 shows a significant decrease in the number of cells after a 48 h EO treatment compared to the 24 h one, evidencing that longer exposures decrease the proliferation capacity of the progeny. The microscopic observation of Caco-2 cells incubated under optimal conditions showed a polygonal appearance (Figure 1B). However, after the 48 h low EO concentration treatment, the culture appearance changed, exhibiting both long adherent and suspended rounded cells (Figure 5B). Our results are in agreement with other studies, where *E. uniflora* EO cytotoxicity was observed in various tumoral cell lines, including MCF-7 (breast), HCT-116 (colon), AGP-01 (malignant gastric ascites), and SKMEL-19 (melanoma), even at low concentrations [24,51,52].

Dysregulated cell proliferation is one of the hallmarks of carcinogenesis [53]. As a step toward determining the potential chemopreventive and/or therapeutic properties of *E. uniflora* EO in CRC, we studied the effect of this EO on cell proliferation. The analysis of the proliferation kinetics under optimal culture conditions of normal CCD 841 CoN and Caco-2 cell lines (Figure 6A,B, Table 3) showed that, as expected, the tumoral Caco-2 cell proliferation was faster and reached a higher number of cells in the stationary phase compared to normal CCD 841 CoN cells. Also, the cell viability determined during cell proliferation (Figure 6C) evidenced the tumoral characteristic of the Caco-2 cell line, showing higher viability than the normal CCD 841 CoN throughout the proliferation study.

To study the potential antiproliferative effect of *E. uniflora* EO, the lowest concentration of EO (0.66 mg/mL) that ensures maximum cell survival was selected. Interestingly, although the CCD 841 CoN cell line exhibited 100% survival upon treatment with [EO] = 0.66 mg/mL, the proliferation was impaired, and no viable cells were observed (Figure 5B). In the case of the tumoral Caco-2 cells, and unlike the normal CCD 841 CoN, the analysis of the proliferation kinetics and cell viability upon *E. uniflora* EO treatment (Figure 7A,B, Table 3) evidenced that the prolonged exposure to a low concentration of EO produced the progressive loss of cell viability, contributing to a slower proliferation kinetics and the decline phase observed after 11 days. This finding confirms the modification of the proliferation capability of the progeny shown in Figure 4.

Several studies have reported antiproliferative effects of the *Eugenia* sp extracts or their single chemical components in cancer cells, assessed by cytotoxicity assays, such as MTT and apoptosis determinations [24,54–56]. To our knowledge, cell proliferation kinetics studies using normal and tumoral human colon cell lines and their comparison have not been reported. Our findings demonstrate that the *E. uniflora* leaf EO affects cell proliferation, producing lethal events and probably altering the cell cycle. In the chemical characterization of the Pitanga EO, 49 compounds were

identified (Table 2), of which 12 were monoterpenes and 33 sesquiterpenes, mainly hydroxycalamenene, δ -cadinene, and τ -muurolol. Various monoterpenes and sesquiterpenes with anti-inflammatory, antioxidant, and antiproliferative properties were confirmed. Myrcene combined with other monoterpenes, such as β -pinene, has been shown to inhibit colon carcinoma tumorigenesis in in vivo and in vitro assays [57]. Other monoterpenes detected, such as terpinolene and limonene, have been proposed as potential anticarcinogenic agents, both as chemopreventives and chemotherapeutics [58–61]. Among the sesquiterpenes, calamenene inhibits the cell proliferation of various human tumor lines, including lung (A549), breast (MCF7), liver (HepG2), cervical (HeLa), and prostate (PC-3) cancer cells [62]. Also, δ -cadinene acts by inhibiting ovarian cancer cell proliferation through cell cycle arrest and induction of apoptosis [63]. Little information is available on the biological effects of τ -muurolol, although anti-inflammatory and antioxidant properties have been reported in EOs containing it [64].

To explore further the possible mechanisms underlying the cytotoxic and antiproliferative effects of *E. uniflora* EO, the induction of DNA damage and oxidative stress was assessed by Comet assay and Cellular ROS assay, respectively. Our results showed genomic damage in the normal CCD 841 CoN and Caco-2 cell lines, as revealed by the observation of DNA fragmentation, even at low doses (Figure 8). In agreement with the CCD 841 CoN cell viability results, extensive cell destruction was observed in the EO-treated samples (compare Figure 8A,B). In treated Caco-2 cells, DNA damage was also observed, as shown by the presence of comet tails images (Figures 8C,D); however, the overall DNA fragmentation was less dramatic than that of normal cells.

It is known that in normal cells, the concentration of ROS remains low and is regulated by antioxidant defenses. In tumor cells, an imbalance occurs that leads to an increased demand for ROS during proliferation. However, increased oxidative stress can also generate genomic alterations, leading to apoptosis and cell death. Therefore, increasing oxidative stress may constitute an antineoplastic mechanism [65]. The determination of the ROS levels as a function of the concentration of *E. uniflora* EO in CCD 841 CoN cells showed a significant increase only at the highest concentration tested ([EO] = 13.2 mg/mL) (Figure 9, white bars). This result indicates that the detrimental effect of the Pitanga EO on normal CCD 841 CoN cells' cytotoxicity and viability is unlikely to be related to an increase in oxidative stress. In contrast, Caco-2 cells exhibited a significant increase in oxidative stress as a function of the EO concentration (Figure 9, striped bars), thus contributing to the detrimental effect on cell viability. Research on *Drosophila melanogaster* indicates that exposure to *E. uniflora* EO can increase ROS production, lipid peroxidation, and alterations in antioxidant enzyme activity [66,67]. Ascari et al. [51] reported pro-oxidative effects and cytotoxicity of two compounds isolated from *E. uniflora* leaf extract in A549 lung adenocarcinoma epithelial cells and IMR90 human lung fibroblasts. In the same line of evidence about the role of the EO in ROS induction, several studies have demonstrated that *E. uniflora* EO increases the antioxidant cellular defenses such as glutathione and superoxide dismutase [24,33], suggesting that the EO triggers ROS production, leading to the induction of an antioxidant cellular response.

5. Conclusions

In conclusion, the present study shows evidence of the cytotoxic and antiproliferative effects of *Eugenia uniflora* essential oil in human normal CCD 841 CoN and tumoral Caco-2 cell lines. Even exposure to low concentrations of EO produced lethality and loss of viability of the treated cells and their progeny. Correspondingly, the detrimental effect on cell proliferation, DNA damage, and oxidative stress production in both cell lines led to the consideration of this essential oil as a potential antineoplastic agent. Our results suggest that the *E. uniflora* EO may prevent the proliferation of normal cells, as shown by the loss of viability when proliferating and the lack of proliferation of the EO-treated human CCD 841 CoN colonic cells. Moreover, the cytotoxic and antiproliferative effects of the EO on the treated tumoral Caco-2 cells support the need for further investigation to consider the *E. uniflora* EO as a whole or their isolated bioactive compounds as antineoplastic therapeutic agents.

Author Contributions: Conceptualization: A.G.S. and D.J.K.; methodology: M.M., P.L., E.D., M.J.Z., D.J.K.; formal analysis: A.G.S., P.L., E.D., D.J.K.; investigation: A.G.S., M.M., P.L., E.D., M.A.S., G.F., B.V., J.C., D.J.K.; writing—original draft preparation: A.G.S., M.M., D.J.K.; writing—review and editing: A.G.S. and D.J.K.; supervision: D.J.K.; project administration: A.G.S., M.A.S., D.J.K.; funding acquisition: D.J.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by “United Nations Development Programme”, grant number 5381 – Proyecto Global ABS UND/GEF "Fortalecimiento de recursos humanos, marco legal y capacidades institucionales para la implementación del Protocolo de Nagoya".

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available upon reasonable request from the corresponding authors.

Acknowledgments: The authors (A.G.S., M.M., and D.J.K.) acknowledge Prof. R. Daniel Peluffo (Biophysical Chemistry Group, Biological Sciences Department, CENUR Litoral Norte, Universidad de la República) for access to resources and support. The authors thank Natalia Ibarгойen for technical support.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

EO Essential Oil
TB Trypan Blue
DT Cell Doubling Time
FBS Fetal Bovine Serum
PBS Phosphate buffer saline
CRC Colorectal Cancer

References

1. Dzobo, K. 2.20 - The Role of Natural Products as Sources of Therapeutic Agents for Innovative Drug Discovery. In *Comprehensive Pharmacology*, Kenakin, T., Ed.; Elsevier: Oxford, 2022; pp. 408-422.
2. Ekor, M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Pharmacology* **2014**, Volume 4 - 2013, doi:10.3389/fphar.2013.00177.
3. Keszenman, D.J., German, P. Traditional medicine usage of pitanga's leaves for medicinally valuable infusions is directing the search for potential chemotherapeutic agents against colon cancer. In *ABS is Genetic Resources for Sustainable Development*; UNDP – Global Environmental Finance Sustainable Development Cluster Bureau for Policy and Programme Support United Nations Development Programme 304: 2018; pp. 212-219
4. Allegra, S.; De Francia, S.; Turco, F.; Bertaggia, I.; Chiara, F.; Armando, T.; Storto, S.; Mussa, M.V. Phytotherapy and Drugs: Can Their Interactions Increase Side Effects in Cancer Patients? *Journal of Xenobiotics* **2023**, *13*, 75-89.
5. Asimwe, J.B.; Nagendrappa, P.B.; Atukunda, E.C.; Kamatenesi, M.M.; Nambozi, G.; Tolo, C.U.; Ogwang, P.E.; Sarki, A.M. Prevalence of the Use of Herbal Medicines among Patients with Cancer: A Systematic Review and Meta-Analysis. *Evidence-Based Complementary and Alternative Medicine* **2021**, *2021*, 9963038, doi:https://doi.org/10.1155/2021/9963038.
6. Tayeb, B.A.; Kusuma, I.Y.; Osman, A.A.M.; Minorics, R. Herbal compounds as promising therapeutic agents in precision medicine strategies for cancer: A systematic review. *Journal of Integrative Medicine* **2024**, *22*, 137-162, doi:https://doi.org/10.1016/j.joim.2024.02.001.

7. Youn, B.-Y.; Kim, J.-H.; Jo, Y.-K.; Yoon, S.; Im, J.-Y.; Kim, H.-J.; Lee, J.-D.; Ko, S.-G. Current Characteristics of Herbal Medicine Interventions for Cancer on Clinical Databases: A Cross-Sectional Study. *Integrative Cancer Therapies* **2023**, *22*, 15347354231218255, doi:10.1177/15347354231218255.
8. Bray, F.; Laversanne, M.; Sung, H.; Ferlay, J.; Siegel, R.L.; Soerjomataram, I.; Jemal, A. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians* **2024**, *74*, 229-263, doi:https://doi.org/10.3322/caac.21834.
9. Chambers, A.C.; Dixon, S.W.; White, P.; Williams, A.C.; Thomas, M.G.; Messenger, D.E. Demographic trends in the incidence of young-onset colorectal cancer: a population-based study. *British Journal of Surgery* **2020**, *107*, 595-605, doi:10.1002/bjs.11486.
10. Howren, A.; Sayre, E.C.; Loree, J.M.; Gill, S.; Brown, C.J.; Raval, M.J.; Farooq, A.; De Vera, M.A. Trends in the Incidence of Young-Onset Colorectal Cancer With a Focus on Years Approaching Screening Age: A Population-Based Longitudinal Study. *JNCI: Journal of the National Cancer Institute* **2021**, *113*, 863-868, doi:10.1093/jnci/djaa220.
11. Musetti, C.; Garau, M.; Alonso, R.; Piñeros, M.; Soerjomataram, I.; Barrios, E. Colorectal Cancer in Young and Older Adults in Uruguay: Changes in Recent Incidence and Mortality Trends. *International Journal of Environmental Research and Public Health* **2021**, *18*, 8232.
12. Piñeros, M.; Laversanne, M.; Barrios, E.; Cancela, M.d.C.; de Vries, E.; Pardo, C.; Bray, F. An updated profile of the cancer burden, patterns and trends in Latin America and the Caribbean. *The Lancet Regional Health – Americas* **2022**, *13*, doi:10.1016/j.lana.2022.100294.
13. Clinton, S.K.; Giovannucci, E.L.; Hursting, S.D. The World Cancer Research Fund/American Institute for Cancer Research Third Expert Report on Diet, Nutrition, Physical Activity, and Cancer: Impact and Future Directions. *The Journal of Nutrition* **2020**, *150*, 663-671, doi:https://doi.org/10.1093/jn/nxz268.
14. Arrillaga, B. *Plantas usadas en medicina natural.*; 1997.
15. Davies, P. Fichas técnicas de cultivo. In *Estudios de domesticación y cultivo de especies medicinales y aromáticas nativas.*, FPTA-INIA, Ed.; INIA: Montevideo, Uruguay., 2004; pp. 37 - 120.
16. Paz, E.A.; Bassagoda, M.J.; Ferreira, F. *Yuyos: uso racional de las plantas medicinales*; Ed. Fin de Siglo: 1992.
17. Tabakian, G. Estudio comparativo de plantas medicinales vinculadas a tradiciones indígenas y europeas en Uruguay. *Bonplandia* **2019**, *28*, 135-158, doi:10.30972/bon.2823855.
18. Stefanello, M.É.A.; Pascoal, A.C.R.F.; Salvador, M.J. Essential Oils from Neotropical Myrtaceae: Chemical Diversity and Biological Properties. *Chemistry & Biodiversity* **2011**, *8*, 73-94, doi:https://doi.org/10.1002/cbdv.201000098.
19. Brussa, C.A., Grela González, A. *Flora arbórea del Uruguay : con énfasis en las especies de Rivera y Tacuarembó*; 2007; p. 544.
20. Jolochín Manorani, G. Estudios biogeográficos en poblaciones uruguayas de *Eugenia uniflora* L. Universidad de la República, Facultad de Ciencias, 2016.
21. Bagetti, M.; Facco, E.M.P.; Piccolo, J.; Hirsch, G.E.; Rodriguez-Amaya, D.; Kobori, C.N.; Vizzotto, M.; Emanuelli, T. Physicochemical characterization and antioxidant capacity of pitanga fruits (*Eugenia uniflora* L.). *Food Science and Technology* **2011**, *31*.
22. Vizzotto, M. Fitoquímicos em pitanga (*Eugenia uniflora* L.): seu potencial na prevencao e comabte a doencas. In Proceedings of the III Simpósio nacional do morango II Encontro sobre pequenas frutas e frutas nativas do Mercosul. , Pelotas, Brasil, 2006; pp. 29-34.
23. Ferragut, G., Lombardo, P., Severi, M.A., Vignale, B., Cedano, J., Dellacassa, E., Pérez, E. Estudios de bioactividad de extractos de plantas nativas uruguayas y su papel inmunológico. In Proceedings of the XV Jornadas de la Sociedad Uruguaya de Biociencias. , Piriápolis, Uruguay, 2014.

24. Fidelis, E.M.; Savall, A.S.P.; de Oliveira Pereira, F.; Quines, C.B.; Ávila, D.S.; Pinton, S. Pitanga (*Eugenia uniflora* L.) as a source of bioactive compounds for health benefits: A review. *Arabian Journal of Chemistry* **2022**, *15*, 103691, doi:<https://doi.org/10.1016/j.arabjc.2022.103691>.
25. Lombardo, P. Caracterización química y bioactividad de aceites esenciales contra patógenos de los cítricos. Universidad de la República, Montevideo, Uruguay, 2015.
26. Lombardo, P.; Guimaraens, A.; Franco, J.; Dellacassa, E.; Pérez Faggiani, E. Effectiveness of essential oils for postharvest control of *Phyllosticta citricarpa* (citrus black spot) on citrus fruit. *Postharvest Biology and Technology* **2016**, *121*, 1-8, doi:<https://doi.org/10.1016/j.postharvbio.2016.07.002>.
27. Rodrigues, K.A.d.F.; Amorim, L.V.; Oliveira, J.M.G.d.; Dias, C.N.; Moraes, D.F.C.; Andrade, E.H.d.A.; Maia, J.G.S.; Carneiro, S.M.P.; Carvalho, F.A.d.A. *Eugenia uniflora* L. Essential Oil as a Potential Anti-Leishmania Agent: Effects on *Leishmania amazonensis* and Possible Mechanisms of Action. *Evidence-Based Complementary and Alternative Medicine* **2013**, *2013*, 279726, doi:<https://doi.org/10.1155/2013/279726>.
28. Sánchez, A.G.; Ferragut, G.; Lombardo, P.; Severi, M.A.; Cedano, J.; Vignale, B.; Vázquez, A.; Dellacassa, E.; Keszenman, D.J. . Efectos celulares de extractos de Pitanga. In Proceedings of the 8vo Encuentro Nacional de Frutos Nativos, Rocha, Uruguay, 2017.
29. Victoria, F.N.; Lenardão, E.J.; Savegnago, L.; Perin, G.; Jacob, R.G.; Alves, D.; Silva, W.P.d.; Motta, A.d.S.d.; Nascente, P.d.S. Essential oil of the leaves of *Eugenia uniflora* L.: Antioxidant and antimicrobial properties. *Food and Chemical Toxicology* **2012**, *50*, 2668-2674, doi:<https://doi.org/10.1016/j.fct.2012.05.002>.
30. Vignale, B.; Cabrera, D.; Rodríguez, P.; Machado, G. Selección de frutales nativos en Uruguay. *Horticultura Argentina* **2016**, *35*, 19-29.
31. Weyerstahl, P.; Marschall-Weyerstahl, H.; Christiansen, C.; Oguntimein, B.O.; Adeoye, A.O. Volatile Constituents of *Eugenia uniflora* Leaf Oil1. *Planta Med* **1988**, *54*, 546-549, doi:10.1055/s-2006-962544.
32. Borsoi, F.T.; Rosa, B.B.d.S.; Filomena, M.; Oliveira, F.D.L.d.; Dulce, B.M.; and Kempka, A.P. *Eugenia uniflora* L. seed and pulp extracts: phytochemical profile, cytotoxic potential, antitumoral activity, and α -amylase and α -glucosidase inhibition capacity. *Natural Product Research* **2023**, *37*, 3862-3867, doi:10.1080/14786419.2022.2153128.
33. Falcão, T.R.; de Araújo, A.A.; Soares, L.A.L.; de Moraes Ramos, R.T.; Bezerra, I.C.F.; Ferreira, M.R.A.; de Souza Neto, M.A.; Melo, M.C.N.; de Araújo, R.F.; de Aguiar Guerra, A.C.V.; et al. Crude extract and fractions from *Eugenia uniflora* Linn leaves showed anti-inflammatory, antioxidant, and antibacterial activities. *BMC Complementary and Alternative Medicine* **2018**, *18*, 84, doi:10.1186/s12906-018-2144-6.
34. Núñez, J.G.; Pinheiro, J.d.S.; Silveira, G.F.; Beckenkamp, A.; Buffon, A.; Bruno, A.N. Antineoplastic potential of the aqueous crude extract of *Eugenia uniflora* L. in human cervical cancer. *Brazilian Journal of Pharmaceutical Sciences* **2018**, *54*.
35. Adams, R. Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy. *Carol Stream* **2005**, *16*, 65-120.
36. McLafferty, F.W.; Stauffer, D.B.; Loh, S.Y. Comparative evaluations of mass spectral data bases. *J Am Soc Mass Spectrom* **1991**, *2*, 438-440, doi:10.1016/1044-0305(91)85011-t.
37. Mondello, L. *Mass spectra of flavors and fragrances of natural and synthetic compounds*; Wiley Blackwell Hoboken, NJ, USA: 2015; Volume 10.
38. Dellacassa, E.; Lorenzo, D.; Paz, D. Procesos de extracción aplicados a la obtención de productos aromáticos de origen vegetal. In *Estudios en domesticación y cultivos de especies medicinales y aromáticas nativas.*, Davies, P., Ed.; FPTA-INIA; INIA: Montevideo, Uruguay, 2004a; pp. 155-160.

39. Dellacassa, E., Lorenzo, D., Paz, D. . Caracterización fisicoquímica de los aceites esenciales. . In *Estudios en domesticación y cultivos de especies medicinales y aromáticas nativas.*, Davies, P., Ed.; FPTA-INIA; INIA: Montevideo, Uruguay., 2004b; pp. 161 - 169.
40. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* **1983**, *65*, 55-63, doi:10.1016/0022-1759(83)90303-4.
41. Avelar-Freitas, B.A.; Almeida, V.G.; Pinto, M.C.; Mourao, F.A.; Massensini, A.R.; Martins-Filho, O.A.; Rocha-Vieira, E.; Brito-Melo, G.E. Trypan blue exclusion assay by flow cytometry. *Braz J Med Biol Res* **2014**, *47*, 307-315, doi:10.1590/1414-431X20143437.
42. Keszenman, D.J.; Salvo, V.A.; Nunes, E. Effects of bleomycin on growth kinetics and survival of *Saccharomyces cerevisiae*: a model of repair pathways. *J Bacteriol* **1992**, *174*, 3125-3132, doi:10.1128/jb.174.10.3125-3132.1992.
43. Benitez-Bribiesca, L.; Sanchez-Suarez, P. Oxidative damage, bleomycin, and gamma radiation induce different types of DNA strand breaks in normal lymphocytes and thymocytes. A comet assay study. *Ann N Y Acad Sci* **1999**, *887*, 133-149, doi:10.1111/j.1749-6632.1999.tb07928.x.
44. Keszenman, D.J.; Carmen Candreva, E.; Nunes, E. Cellular and molecular effects of bleomycin are modulated by heat shock in *Saccharomyces cerevisiae*. *Mutat Res* **2000**, *459*, 29-41, doi:10.1016/s0921-8777(99)00056-7.
45. Atanasov, A.G.; Zotchev, S.B.; Dirsch, V.M.; Orhan, I.E.; Banach, M.; Rollinger, J.M.; Barreca, D.; Weckwerth, W.; Bauer, R.; Bayer, E.A.; et al. Natural products in drug discovery: advances and opportunities. *Nature Reviews Drug Discovery* **2021**, *20*, 200-216, doi:10.1038/s41573-020-00114-z.
46. Chaachouay, N.; Zidane, L. Plant-Derived Natural Products: A Source for Drug Discovery and Development. *Drugs and Drug Candidates* **2024**, *3*, 184-207.
47. Costa-Font, J.; Sato, A.; Saenz-de-Miera, B. Cultural persistence of self-assessed health: A study of first- and second-generation migrants. *J Migr Health* **2025**, *11*, 100280, doi:10.1016/j.jmh.2024.100280.
48. Benfatti, C.S.; Cordova, S.M.d.; Guedes, A.; Magina, M.D.A.; Cordova, C.M.M.d. Atividade antibacteriana in vitro de extratos brutos de espécies de *Eugenia* sp frente a cepas de mollicutes. *Revista Pan-Amazônica de Saúde* **2010**, *1*, 33-39.
49. de Souza Prestes, L.; Damé Schuch, L.F.; Hörnke Alves, G.; Ziemann dos Santos, M.A.; Alves Rodrigues, M.R.; Araújo Meireles, M.C. Evaluación de la actividad bactericida de aceites esenciales de hojas de guayabo, pitango y arazá. *Revista Cubana de Plantas Medicinales* **2011**, *16*, 324-330.
50. Schmeda-Hirschmann, G.; Theoduloz, C.; Franco, L.; Ferro, E.; de Arias, A.R. Preliminary pharmacological studies on *Eugenia uniflora* leaves: xanthine oxidase inhibitory activity. *J Ethnopharmacol* **1987**, *21*, 183-186, doi:10.1016/0378-8741(87)90128-0.
51. Ascari, J.; Felipe Maciel Pereira, M.; Schaffka, V.M.; Nunes, D.S.; Magalhães, C.G.; Santos, J.S.; Granato, D.; Carmo, M.; Azevedo, L.; Archilha, M.; et al. Selina-1,3,7(11)-trien-8-one and Oxidoselina-1,3,7(11)-trien-8-one from *Eugenia uniflora* Leaf Essential Oil and Their Cytotoxic Effects on Human Cell Lines. *Molecules* **2021**, *26*, doi:10.3390/molecules26030740.
52. Figueiredo, P.L.B.; Pinto, L.C.; da Costa, J.S.; da Silva, A.R.C.; Mourão, R.H.V.; Montenegro, R.C.; da Silva, J.K.R.; Maia, J.G.S. Composition, antioxidant capacity and cytotoxic activity of *Eugenia uniflora* L. chemotype-oils from the Amazon. *Journal of Ethnopharmacology* **2019**, *232*, 30-38, doi:https://doi.org/10.1016/j.jep.2018.12.011.
53. Hanahan, D.; Weinberg, R.A. Hallmarks of cancer: the next generation. *Cell* **2011**, *144*, 646-674, doi:10.1016/j.cell.2011.02.013.

54. Aranha, E.S.P.; de Azevedo, S.G.; dos Reis, G.G.; Silva Lima, E.; Machado, M.B.; de Vasconcellos, M.C. Essential oils from *Eugenia* spp.: In vitro antiproliferative potential with inhibitory action of metalloproteinases. *Industrial Crops and Products* **2019**, *141*, 111736, doi:https://doi.org/10.1016/j.indcrop.2019.111736.
55. de Oliveira, G.G.F.V.; Longue, M.F.; Pescinelli, L.M.R.; Charret, T.S.; Nogueira, T.S.R.; Pereira, M.T.M.; Vieira, I.J.C.; Abreu, L.S.; Pascoal, V.D.B.; Pascoal, A.C.R.F. *Eugenia brasiliensis*: Analysis of the Chemical Profile and Evaluation of Cytotoxic Potential. *Chem Biodivers* **2025**, e202500429, doi:10.1002/cbdv.202500429.
56. Ismiyati, N.; Putri, D.; Kusumastuti, S.A.; Febriansyah, R. Antiproliferative Effect of Ethanol Extract *Eugenia uniflora* Lam. Leaves on T47D Cells. *Indonesian Journal of Cancer Chemoprevention* **2012**, *3*, 370, doi:10.14499/indonesianjancchemoprev3iss2pp370-375.
57. Spyridopoulou, K.; Tiptiri-Kourpeti, A.; Lampri, E.; Fitsiou, E.; Vasileiadis, S.; Vamvakias, M.; Bardouki, H.; Goussia, A.; Malamou-Mitsi, V.; Panayiotidis, M.I.; et al. Dietary mastic oil extracted from *Pistacia lentiscus* var. *chia* suppresses tumor growth in experimental colon cancer models. *Sci Rep* **2017**, *7*, 3782, doi:10.1038/s41598-017-03971-8.
58. Agus, H.H.; Sarp, C.; Cemiloglu, M. Oxidative stress and mitochondrial impairment mediated apoptotic cell death induced by terpinolene in *Schizosaccharomyces pombe*. *Toxicol Res (Camb)* **2018**, *7*, 848-858, doi:10.1039/c8tx00100f.
59. Jia, S.S.; Xi, G.P.; Zhang, M.; Chen, Y.B.; Lei, B.; Dong, X.S.; Yang, Y.M. Induction of apoptosis by D-limonene is mediated by inactivation of Akt in LS174T human colon cancer cells. *Oncol Rep* **2013**, *29*, 349-354, doi:10.3892/or.2012.2093.
60. Okumura, N.; Yoshida, H.; Nishimura, Y.; Kitagishi, Y.; Matsuda, S. Terpinolene, a component of herbal sage, downregulates AKT1 expression in K562 cells. *Oncol Lett* **2012**, *3*, 321-324, doi:10.3892/ol.2011.491.
61. Menezes, I.O.; Scherf, J.R.; Martins, A.O.B.P.B.; Ramos, A.G.B.; Quintans, J.d.S.S.; Coutinho, H.D.M.; Ribeiro-Filho, J.; de Menezes, I.R.A. Biological properties of terpinolene evidenced by in silico, in vitro and in vivo studies: A systematic review. *Phytomedicine* **2021**, *93*, 153768, doi:https://doi.org/10.1016/j.phymed.2021.153768.
62. Zhong, Z.-C.; Zhao, D.-D.; Liu, Z.-D.; Jiang, S.; Zhang, Y.-L. A New Human Cancer Cell Proliferation Inhibition Sesquiterpene, Dryofraterpene A, from Medicinal Plant *Dryopteris fragrans* (L.) Schott. *Molecules* **2017**, *22*, 180.
63. Hui, L.M.; Zhao, G.D.; Zhao, J.J. δ -Cadinene inhibits the growth of ovarian cancer cells via caspase-dependent apoptosis and cell cycle arrest. *Int J Clin Exp Pathol* **2015**, *8*, 6046-6056.
64. Chien, T.C.; Lo, S.F.; Ho, C.L. Chemical composition and anti-inflammatory activity of *Chamaecyparis obtusa* f. *formosana* wood essential oil from Taiwan. *Nat Prod Commun* **2014**, *9*, 723-726.
65. Sosa, V.; Moliné, T.; Somoza, R.; Paciucci, R.; Kondoh, H.; ME, L.L. Oxidative stress and cancer: an overview. *Ageing Res Rev* **2013**, *12*, 376-390, doi:10.1016/j.arr.2012.10.004.
66. da Cunha, F.A.B.; Wallau, G.L.; Pinho, A.I.; Nunes, M.E.M.; Leite, N.F.; Tintino, S.R.; da Costa, G.M.; Athayde, M.L.; Boligon, A.A.; Coutinho, H.D.M.; et al. *Eugenia uniflora* leaves essential oil induces toxicity in *Drosophila melanogaster*: involvement of oxidative stress mechanisms. *Toxicology Research* **2015**, *4*, 634-644, doi:10.1039/c4tx00162a.
67. de Carvalho, N.R.; Rodrigues, N.R.; Macedo, G.E.; Bristot, I.J.; Boligon, A.A.; de Campos, M.M.; Cunha, F.A.B.; Coutinho, H.D.; Klamt, F.; Merritt, T.J.S.; et al. *Eugenia uniflora* leaf essential oil promotes mitochondrial dysfunction in *Drosophila melanogaster* through the inhibition of oxidative phosphorylation. *Toxicol Res (Camb)* **2017**, *6*, 526-534, doi:10.1039/c7tx00072c.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.