

Brief Report

Not peer-reviewed version

On Melanocytic Nevi Analysis

[Antonella Chesca](#)*

Posted Date: 3 April 2026

doi: 10.20944/preprints202401.0665.v10

Keywords: epiderm; analysis; optical microscope; electronic microscope; future trens



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Brief Report

On Melanocytic Nevi Analysis

Antonella Chesca

Transylvania University of Brasov, Romania; anto.chesca@gmail.com

Abstract

Epiderm is composed by specific layers, functions and implications in the life The purpose of the study is to analyse and to identify structural characteristics reffering to melanocytic nevi, in youth patients. Congenital melanocytic nevi are pigmented lesions that are usually present a birth. They are generally benign, but a small percentage (especially the larger ones) can potentially transform into malignant melanoma. Using both optical and electronic microscope, could be possible a better description related specificity in melanocytic nevi characteristics. Future trends, are important key points in management, including preventive and prophylactic methods.

Keywords: epiderm; analysis; optical microscope; electronic microscope; future trens

Introduction

In order to define a disease, must have in attention a lot of differents factors such as historical, or social and cultural key points. Results of research studies, show us that some connective cells such as fibroblasts, lose their identity, in pathological conditions. [1] Another specific cells, namely melanocytes are known that having a specific structural point that is consider important in structural pathological description. [2] Referring to melanocytic nevi, in medical specific field of study and of research, various pigmented lesions of the epiderm, known as nevi, could be observe in different pars of the body. [3] For a proper diagnostic, an atypical nevus,can be biopsied.[4] In this direction, is important to practice a biopsy beside the extended clinical evaluation in melanocytic nevi A great point in this field of research, could be possible the genetic susceptibility for morphological and functional alterations, in nevi with that surrounding nevi changes. [5] A complete medical examination, play a great point for establishing the medical conduct, for a healthy status improving.[6,7] Structural analysis describe specific cells namely melanocytes as aggregated in ‘nests’, which conduct forming the nevus cells.[8] To the youth patients researchers found specific cells knowing as melanocytes. This specific cells could be found in areas of the epiderm of the parts of the body. [9,10] Theoretical and practical studies, show that melanocytic nevi developing in utero present genetical differences from those that appear later.[11,12] In the present field, we can mention about various informations from scientific literature, referring to specific nevi. [13] Also from literature and from practicum actually are known different scientific informatiuons about extending melanocytic *nevi*, having specific scientific names.[14] Because are many cases in all of the world, diagnosed as melanocitic nevi, we can mention that currently, the proper treatment of epidermal nevi is challenging. [15,18]Congenital melanocytic nevi it is known as a subject of research that offer controversy. [19] Clinical monitoring in congenital melanocytic nevi is important for diagnosis and for possible medical treatment strategies applications. [20] A complete examination of the human body, during a medical examination, is important. [21,22] Best to mention that the nowadays higher incidence in melanoma is in accompaniment of the nevi existence of the body and of the increase exposure to the ultraviolet light. [23,24] Practical biopsy is important for diagnosis.[25]One of an important point in the diagnosis of melanocytic nevi is to differentiate melanocitic nevi from a possible melanoma. [26,27]. An earlier diagnosis of the melanoma play a great role in idea that neoplastic lessions could be develop from pre-existing nevi in many cases. [28] Unfortunately, the epidermal melanoma is growing faster., depending of various conditions. [29,30]

Melanocytic nevi considerations

From many types of nevi, in the next short written text, we will describe a little bit on Ito's nevus and Ota's nevus. These two types of nevi could be observed in pregnancy, at birth and also at puberty. Their presence is in concordance with hormonal changes. Research studies described possible malignant status in Ota's nevus, rarely in Ito's nevus. [31,32] These two previously mentioned types of nevi, namely Ito and Ota, do not differ from a histological point of view. Ito's nevus and Ota's nevus are distinguished by specific location on the body. So, typically, Ito's nevus occurs in the arm region and Ota's nevus could be found on the face. [33,34] Ota's nevus could also be found in the supply areas of the first two branches of the trigeminal nerve. [34–37] Structurally, Ito's nevus presents as a slate-blue/gray-blue macula in the shoulder/breast and lateral arm region in the supply area of the brachial nerve, in infants or prior to puberty. [38] It is known that a specific sign of melanoma within the existing Ito's nevus is a typical nodule. [39] In rarely malignant cases in patients diagnosed with Ito's nevi have been reported in addition, typical nodules. [40,41]

From birth age, congenital melanocytic nevi (CMN) are known as one of the frequent skin lesions. [42] From research results and conclusions, could be found rarely medical namely, neuroend differentiation. To a specific analysis, is possible to observe specific areas of cells with myxoid stroma in addition. Possible resemble later than, as neurofibromas. [42]

From a current research perspective we can mention that in utero, specific stem cells from the neuroectoderm play a significant role such as migration to the skin as melanoblasts. Mechanism refers to a differentiation process into melanocytes. In addition, mutations arising in specific cells can occur to well known mosaicism. Good to know that in the early embryogenesis, multipotent progenitor cells can be affected, leading to the presence of multiple congenital melanocytic nevi and also to extracutaneous alterations. [43,44]

In addition to previously above mentioned idea, congenital melanocytic nevi occur as a result of in-utero somatic mutations. In this idea, genes play a great role. So there are known the mitogen-activated protein kinase (MAPK) pathway (mainly NRAS and BRAF). More than, their specific mutations refers to damages in the development of cutaneous and/or extracutaneous previously mentioned mosaicism. [45] Additionally to congenital melanocytic nevi, proliferative nodules (PN) constitute nodular lesions. [46] All described epidermal alterations, are factors incriminate in differentiating proliferative nodule from melanoma.

Good to mention that neurocutaneous melanosis is a disease where congenital melanocytic nevi are associated with melanocyte proliferation. Besides, satellite lesions are especially at risk. Clinically are signs and specific symptoms So could be describe neurological symptoms, with possible intracranial pressure. [47] From literature data there are known a lot of types of melanocytic nevi. [48] Then possible surgical intervention and pathological diagnosis we can take into consideration. [49] Management directions play also a significant role as future directions. [50] From medical point of view we can mention that in case of an atypical nevus could be practice a proper biopsy. [51] Using this previously mentioned procedure is important to extend tissue excision in the unaffected structure. [51] We can mention a lot about possible complications that include in pre-existing immune disorders, new exacerbations. [52]

Normal epiderm and pathological key points used for diagnosis

Normal epiderm is composed by specific layers, functions and implications in the human life. There are known a number of layers, namely basale stratum, spinosum stratum, granulosum stratum, lucidum stratum and corneous stratum. In abnormal conditions, depending on diverse factors, epidermal layers could possibly be affected structurally. Morphologically, specific cellularity, cell organelles, junctions, and not only suffer damages. Knowing this and referring to epidermal nevi, previously mentioned possible damages play a role in pathological status appearance, including nevi in youth patients. From another perspective, in a normal epiderm, there are known four cell populations. Namely of specific cells are keratinocytes, melanocytes, Langerhans and Merkel cells. Referring to keratinocytes we can mention that comprise the bulk of the epiderm. [53,54] Melanocytes are interesting cells by their function. So these cells synthesize and store pigment namely melanin in

specific organelles known as melanosomes. The role of melanocytes is the transfer of melanosomes via dendrites, primarily to keratinocytes.[55] Langerhans cells are antigen-presenting cells found in the epiderm. [56] Merkel cells are neuroendocrine origin. This cells are known by the implication as adapting mechanoreceptors.[57,58] [Also important to found and to know a little bit about cause of developing the nevi in young peoples. [59] Genetical causes is important in that cases..[60]

One of the important aspects in nevi research studies is the problem reffering to dysplastic nevi category. This category on nevi is considered being a special one atypical in this dermatologically subject.[61–63] Etiology in dysplastic nevi category unfortunately remain nowadays unclear. Genetically aspect are important in diagnosis. So phenotypic expression is important to detemin and to know more exactly on dysplastic nevi. Clark nevi category is an example in this dermatologically direction. In dysplastic nevi appearance, important are also environmental factors in addition with skin sensibility. Studies in youth patients area actually relatively concludent. [64–66] Objectives in dysplastic nevi category management, reffers to some medical directions. In this way, we can include history in appearance, etiology key points, patophysiology aspects, dermatological analyse, diagnosis. Genetic analyse is a modern one. Using this previously study domain, important is to identify BRAF, p16, or p53 modifications in genetically expression. Medical treatment possibilities in dysplastic nevi is established by an interprofessional specialized team. [67–69] (Table I).

Table I. Dysplastic nevi management key points.

Ethiology	including historically aspects
Pathophysiology	including genetical analyse
Analyse	structural, ultrastructural, genetical
Diagnostic	established by an interprofessional team

Practical methods for diagnostic and their results

For the purpose of the study we can mention about classic laboratory technique used and about the materials needed. In the specific laboratory, were followed the steps of the classic method, using Hematoxylin &Eosin staining. The samples used were from male and female youth patients, before mature age, from urban and rural residence. This are examined using the optical microscope. In addition can be used also specialized microscopes and electron microscopes. TEM (transmission electron microscope) is usually used for structural analyse. The operative pieces are intended to bring in the pathological anatomy service for macroscopic examination for diagnosis.

Epiderm protect us during the life, from different factors. For a morphological analyse, structural and ultrastructural characteristics could be describes, using optical and electron microscope. Structural analyse of the epiderm, using colour laboratory techniques, is able to describe the specific layers with their characteristics. More than, using electron microscope, specific compounds as filaggrin which is knowing as an important epidermal protein and/or tight junction located in the granular layer of the epiderm, could be observed. For this purpose, transmission electron microscope examination, is consider one proper method for analyse. Scanning electron microscopy is also a modern method for analyse, which offer results that demonstrate abnormalities in the epiderm ultrastructure. The human body is covers by skin and the epiderm contein differents types of glands, as sebaceous glands and sudoripar glands. In this study direction, it is known a typical physiopathologic mechanism for the functionality of the body, including epidermal compounds and their body sorroundings. Histopathological analyse describe various modifications to the melanocytic nevi aspect, located on various regions of the body. So we can mention asymmetry, irregular form, cytologic atypia, and mitotic activity. Medical specialists, describe and conclude that to benign melanocytic nevi, could be possible a describtion for atypical pathological characteristics of nevi and more important to mention characteristics when benign nevi are traumatized. Epiderm is a barrier, but is able for conducing to an illness status if include modifications in structural compounds. Histopathological analyse describe the melanocytic nevi located on various regions of the body,with asymmetry, irregular form, cytologic atypia and mitotic activity. More than, medical

specialists, describe and conclude related to the structural aspects in benign traumatized melanocytic nevi. In this field, dermoscopy play a role for a proper diagnostic. Dermoscopy play a role for a proper diagnostic important in practice to all ages, including, youth age and children. Immunohistochemistry (IHC) method has limitations from variability in sensitivity and specificity. [Figure 1]. (Table II.) (Table III.)




Figure 1. Layers of the normal epiderm.

Table II. IHC method in diagnostic.

Immunohistochemistry (IHC)	a process that involves using of selective antibodies to target and stain antigens
Immunohistochemistry (IHC)	known to be specific to a particular condition.
Immunohistochemistry (IHC)	a process that creates a color signal that can be visualized on light microscopy.

Table III. Light microscopy implications in epidermal nevi.

Light microscopy	The 100x magnification usually requires a medium between the lens and the slide, such as oil immersion
Light microscopy	Most microscopes used in dermatopathology are compound microscopes.
Light microscopy	Is the primary means for the magnified examination of tissue blocks.
Light microscopy	Generally shows cells as colorless, necessitating the need for various stains. Most tissue is initially stained with hematoxylin and eosin (H&E).

Management strategies

Key point in dermatological diagnosis are important to establish and to know, for management directions. (Table IV) Disease diagnostic and future trends, are important key points in management directions. As future directions good to mention idea including preventive and prophylactic methods. For each patient is important to improve a skin evaluation. (Table V) In this context a simple

clinical skin evaluation could be accompanied by biopsy, excision, surgical practice and other accompanying activities. (Table VI) (Table VII)

Table IV. Nevi characteristics for management directions

Review the management options available for melanocytic nevi.	Describe the presentation of a patient with melanocytic nevi.
Identify the risk factors for melanocytic nevi.	Outline interprofessional team strategies for improving care coordination and communication to advance the management of melanocytic nevi and improve outcomes.

Table V. Nevi management directions depending on patient age

children	Screening methods, dermatologist visit with diagnosis and medical indications
youth	Etiology factors, dermatologist diagnosis, future trends
mature	dermatologist diagnosis, planning preventive methods for avoiding a presumptive recidive
olders	dermatologist diagnosis, planning preventive methods for avoiding a presumptive malignancy recidive

Table VI. Nevi characteristics for differential diagnostic

Atypical mole	Basal cell carcinoma	Café au lait spots
Cutaneous melanoma	Nevi of Ota and Ito	Nevus spilus
Cockade nevus	Nodular lesions	Pyogenic granuloma

An interest in medicine refers to congenital melanocytic nevi (CMN). [70–72] In this direction must be carefully to strategies for management with their applicability in routine practice. [73] A first diagnostic point in attention are screening methods in CMN. Imaginal methods are important and for this, magnetic resonance imaging (MRI) and their results play a great role for diagnostic. [74,75] A next point in management strategies, refers to a medical visit to dermatologist. [76] Practical biopsy also is important in idea to find a possible malignancy degeneration, and then a faster medical decision application in practice. [77,78]

Table VII. Management strategies and considerations in congenital melanocytic nevi (CMN) analysis for diagnostic

Screening	Magnetic resonance imaging (MRI)	malignant melanoma (MM)
Visit to Dermatology	medical specialists with competence in concordance with patients age	Congenital melanocytic nevi (CMN)
Biopsy	pain, ulceration, age, and change in color, thickness, and size	Congenital melanocytic nevi (CMN) for diagnosis, taking in consideration malignancy potential

Discussions

Great interest in knowing epidermal compounds. So, the epiderm, is composed of a number of specific layers. Specific cells are known. One of the role of the epiderm is implication in different injuries. Alterations in the compounds of the epiderm layers, contribute to the visual signs of pathologic conditions. One research direction, refer to the role of benign melanocytic lesions with alterations, which conduct to malignant cutaneous melanoma. Related to melanocytic nevi, in some circumstances, could be possible that the prognosis be poor having in attention the healthy of the patients having comorbidities. Pathological analyse and diagnosis referring to melanocytic nevi located on different regions from the body can find asymmetry, irregularity, cytologic atypia, and mitotic activity. Medical team including dermatologists, pathologists and dermatopathologists play a great role, in idea referring to differentiate benign melanocytic nevi from malignant melanoma. This is important in order to avoid unnecessary surgical intervention or a treatment. Management and a better clinical evaluation, is a key point for a proper next time abordation in epidermal pathologically compounds as nevi. Structural analyze to the epidermal alterations is important for diagnosis.

Conclusions

Techniques for the laboratory diagnosis, as a key point in monitoring pathological status to patients diagnosed with melanocytic nevi, conduct to a proper quality of life. Implication of an interprofessional team is a condition that play a great medical role. Research studies described possible malignant status in nevi, in time. Future trends, new laboratory methods and techniques for diagnosis. are in attention, for the next coming period of time.

References

1. Salzer M.C., Lafzi A., Berenguer-Llargo A., Youssif C., Castellanos A., Solanas G., Peixoto F.O., Stephan-Otto Attolini C., Prats N., Aguilera M., Martín-Caballero J., Heyn H., Benitah S.A. Identity noise and adipogenic traits characterize dermal fibroblast aging. *Cell*. 2018;175:1575–1590.e22.
2. Palicka GA, Rhodes AR. Acral melanocytic nevi: prevalence and distribution of gross morphologic features in white and black adults. *Arch Dermatol*. 2010 Oct;146(10):1085-94.
3. Wang M, Xu Y, Wang J, Cui L, Wang J, Hu XB, Jiang HQ, Hong ZJ, Yuan SM. Surgical Management of Plantar Melanoma: A Retrospective Study in One Center. *J Foot Ankle Surg*. 2018 Jul-Aug;57(4):689-693.
4. Richtig E. ASCO Congress 2018: melanoma treatment. *Memo*. 2018;11(4):261-265.
5. Wang B, Qu XL, Chen Y. Identification of the potential prognostic genes of human melanoma. *J Cell Physiol*. 2019 Jun;234(6):9810-9815.
6. Bristow I, Bower C. Melanoma of the Foot. *Clin Podiatr Med Surg*. 2016 Jul;33(3):409-22.
7. Lallas A, Kyrgidis A, Koga H, Moscarella E, Tschandl P, Apalla Z, Di Stefani A, Ioannides D, Kittler H, Kobayashi K, Lazaridou E, Longo C, Phan A, Saida T, Tanaka M, Thomas L, Zalaudek I, Argenziano G. The BRAAFF checklist: a new dermoscopic algorithm for diagnosing acral melanoma. *Br J Dermatol*. 2015 Oct;173(4):1041-9.
8. Tronnier M. Melanotische Flecke und melanozytäre Nävi. In: Plewig G, Ruzicka T, Kaufmann R, Hertl M. *Braun-Falcos Dermatol. Venerol. Allergol*. Springer Berlin Heidelberg, 7. Auflage, 2016.
9. Tolleson WH. Human Melanocyte Biology, Toxicology, and Pathology. *J Environ Sci Health Part C* 2005; 23: 105–61.
10. Thomas AJ, Erickson CA. The making of a melanocyte: the specification of melanoblasts from the neural crest. *Pigment Cell Melanoma Res* 2008; 21: 598–610.
11. Colebatch AJ, Ferguson P, Newell F et al. Molecular genomic profiling of melanocytic nevi. *J Invest Dermatol* 2019; 139: 1762–8.
12. Kinsler VA, Thomas AC, Ishida M et al. Multiple congenital melanocytic nevi and neurocutaneous melanosis are caused by postzygotic mutations in codon 61 of NRAS. *J Invest Dermatol* 2013; 133: 2229–36.
13. Bandyopadhyay D. Halo nevus. *Indian Pediatr* 2014; 51: 850.

14. Fernandez-Flores A. Eponyms, Morphology, and Pathogenesis of some less mentioned types of melanocytic nevi. *Am J Dermatopathol* 2012; 34: 607–18.
15. Kim JJ (Department of Dermatology, Henry Ford Health System, Detroit, MI 48202, USA), Chang MW, Shwayder T. Topical tre-tinoin and 5-fluorouracil in the treatment of linear verrucous epidermal nevus. *J Am Acad Dermatol*. 2000 Jul;43(1 Pt 1):129–132.
16. Brown HM, Gorlin RJ. Oral mucosal involvement in nevus unius lateris (Ichthyosis Hysterix). *Arch Dermatol*. 1960 Apr;81:509–515.
17. Zvulunov A (Soroka Medical Center, Ben-Gurion University, Beer-Sheva Israel), Grunwald MH, Halvy S. Topical calcipotriol for treatment of inflammatory linear verrucous epidermal nevus. *Arch Dermatol*. 1997 May;133(5):567–568.
18. Boyce S (Washington Institute of Dermatologic Laser Surgery, Washington, DC 20037, USA), Alster T. CO2 laser treatment of epidermal nevi: Long-term success. *Dermatol Surg*. 2002 Jul;28(7):611–614.
19. Arad Ehud, Zuker M. Ronald, The shifting paradigm in the management of giant congenital melanocytic nevi: review and clinical applications, *Plast Reconstr Surg*, 2014 Feb;133(2):367–376.
20. Mologousis MA, Tsai SY, Tissera KA, Levin YS, Hawryluk EB., Updates in the Management of Congenital Melanocytic Nevi., *Children (Basel)*. 2024 Jan 2;11(1):62.
21. Bristow I, Bower C. Melanoma of the Foot. *Clin Podiatr Med Surg*. 2016 Jul;33(3):409–22.
22. Lallas A, Kyrgidis A, Koga H, Moscarella E, Tschandl P, Apalla Z, Di Stefani A, Ioannides D, Kittler H, Kobayashi K, Lazaridou E, Longo C, Phan A, Saida T, Tanaka M, Thomas L, Zalaudek I, Argenziano G. The BRAAFF checklist: a new dermoscopic algorithm for diagnosing acral melanoma. *Br J Dermatol*. 2015 Oct;173(4):1041–9.
23. Bartoš V, Kullová M. Malignant Melanomas of the Skin Arising on the Feet. *Klin Onkol*. 2018 Summer;31(4):289–292.
24. Palicka GA, Rhodes AR. Acral melanocytic nevi: prevalence and distribution of gross morphologic features in white and black adults. *Arch Dermatol*. 2010 Oct;146(10):1085–94.
25. Bristow IR, de Berker DA, Acland KM, Turner RJ, Bowling J. Clinical guidelines for the recognition of melanoma of the foot and nail unit. *J Foot Ankle Res*. 2010 Nov 01;3:25.
26. Gershenwald JE, Ross MI. Sentinel-lymph-node biopsy for cutaneous melanoma. *N Engl J Med*. (2011) 364:1738–45. 10.1056/NEJMct1002967.
27. Eggermont AM, Spatz A, Robert C. Cutaneous melanoma. *Lancet*. (2014) 383:816–27. 10.1016/S0140-6736(13)60802-8.
28. Elder DE. Precursors to melanoma and their mimics: nevi of special sites. *Mod Pathol*. (2006) 19(Suppl. 2):S4–20. 10.1038/modpathol.3800515.
29. Gershenwald JE, Ross MI. Sentinel-lymph-node biopsy for cutaneous melanoma. *N Engl J Med*. (2011) 364:1738–45. 10.1056/NEJMct1002967.
30. Eggermont AM, Spatz A, Robert C. Cutaneous melanoma. *Lancet*. (2014) 383:816–27. 10.1016/S0140-6736(13)60802-8.
31. van Krieken JH, Boom BW, Scheffer E. Malignant transformation in a naevus of Ito. A case report. *Histopathology* 1988; 12: 100–2.
32. Wise SR, Capra G, Martin P et al. Malignant melanoma transformation within a nevus of Ito. *J Am Acad Dermatol* 2010; 62: 869–74.
33. Pérez ME, Bley C, Cárdenas C. Nevus of Ota, a classic presentation. *Med Clin (Barc)* 2019; 153: 92.
34. Agarwal P, Patel BC. Nevus of Ota and Ito. *StatPearls* 2020.
35. Kang HY, Kang WH. Bilateral type of nevus of Ota presenting as agminated lentigines. *Eur J Dermatol EJD* 2003; 13: 205–6.
36. Turnbull JR, Assaf C, Zouboulis C, Tebbe B. Bilateral naevus of Ota: a rare manifestation in a Caucasian. *J Eur Acad Dermatol Venereol JEADV* 2004; 18: 353–5.
37. Hori Y, Kawashima M, Oohara K, Kukita A. Acquired, bilateral nevus of Ota-like macules. *J Am Acad Dermatol* 1984; 10: 961–4.
38. Que SKT, Weston G, Suchecki J, Ricketts J. Pigmentary disorders of the eyes and skin. *Clin Dermatol* 2015; 33: 147–58.

39. Martínez-Peñuela A, Iglesias ME, Mercado MR et al. Malignant transformation of a nevus of Ito: description of a rare case. *Actas Dermosifiliogr* 2011; 102: 817–20.
40. Wise SR, Capra G, Martin P et al. Malignant melanoma transformation within a nevus of Ito. *J Am Acad Dermatol* 2010; 62: 869–74.
41. Martínez-Peñuela A, Iglesias ME, Mercado MR et al. Malignant transformation of a nevus of Ito: description of a rare case. *Actas Dermosifiliogr* 2011; 102: 817–20.
42. Zayour M, Lazova R. Congenital melanocytic nevi. *Clin Lab Med*. 2011 Jun;31(2):267-80.
43. Schaffer JV. Update on melanocytic nevi in children. *Clin Dermatol*. 2015 May-Jun;33(3):368-86.
44. Kinsler VA, O'Hare P, Bulstrode N, Calonje JE, Chong WK, Hargrave D, Jacques T, Lomas D, Sebire NJ, Slater O. Melanoma in congenital melanocytic naevi. *Br J Dermatol*. 2017 May;176(5):1131-1143.
45. Price HN. Congenital melanocytic nevi: update in genetics and management. *Curr Opin Pediatr*. 2016 Aug;28(4):476-82.
46. Vergier B, Laharanne E, Prochazkova-Carlotti M, de la Fouchardière A, Merlio JP, Kadlub N, Avril MF, Bodemer C, Lacoste C, Boralevi F, Taieb A, Ezzedine K, Fraitag S. Proliferative Nodules vs Melanoma Arising in Giant Congenital Melanocytic Nevi During Childhood. *JAMA Dermatol*. 2016 Oct 01;152(10):1147-1151.
47. Flores-Sarnat L. Neurocutaneous melanocytosis. *Handb Clin Neurol*. 2013;111:369-88.
48. Bandyopadhyay D. Halo nevus. *Indian Pediatr* 2014; 51: 850
49. Bristow IR, de Berker DA, Acland KM, Turner RJ, Bowling J. Clinical guidelines for the recognition of melanoma of the foot and nail unit. *J Foot Ankle Res*. 2010 Nov 01;3:25.
50. Richtig E. ASCO Congress 2018: melanoma treatment. *Memo*. 2018;11(4):261-265.
51. Crompton JG, Gilbert E, Brady MS. Clinical implications of the eighth edition of the American Joint Committee on Cancer melanoma staging. *J Surg Oncol*. 2019 Jan;119(2):168-174.
52. Richtig E. ASCO Congress 2018: melanoma treatment. *Memo*. 2018;11(4):261-265.
53. Bergstresser PR, Taylor JR. Epidermal 'turnover time'—a new examination. *Br J Dermatol*. 1977 May;96(5):503-9.
54. Hoath SB, Leahy DG. The organization of human epidermis: functional epidermal units and phi proportionality. *J Invest Dermatol*. 2003 Dec;121(6):1440-6.
55. Delevoeye C. Melanin transfer: the keratinocytes are more than gluttons. *J Invest Dermatol*. 2014 Apr;134(4):877-879.
56. Pasparakis M, Haase I, Nestle FO. Mechanisms regulating skin immunity and inflammation. *Nat Rev Immunol*. 2014 May;14(5):289-301.
57. Van Keymeulen A, Mascre G, Youseff KK, Harel I, Michaux C, De Geest N, Szpalski C, Achouri Y, Bloch W, Hassan BA, Blanpain C. Epidermal progenitors give rise to Merkel cells during embryonic development and adult homeostasis. *J Cell Biol*. 2009 Oct 05;187(1):91-100.
58. Patel P, Hussain K. Merkel cell carcinoma. *Clin Exp Dermatol*. 2021 Jul;46(5):814-819.
59. Colebatch AJ, Ferguson P, Newell F et al. Molecular genomic profiling of melanocytic nevi. *J Invest Dermatol* 2019; 139: 1762–8
60. Kinsler VA, Thomas AC, Ishida M et al. Multiple congenital melanocytic nevi and neurocutaneous melanosis are caused by postzygotic mutations in codon 61 of NRAS. *J Invest Dermatol* 2013; 133: 2229–36.
61. Strazzulla LC, Li X, Zhu K, Okhovat JP, Lee SJ, Kim CC. Clinicopathologic, misdiagnosis, and survival differences between clinically amelanotic melanomas and pigmented melanomas. *J Am Acad Dermatol*. 2019 May;80(5):1292-1298.
62. de Unamuno Bustos B, Sahuquillo Torralba A, Moles Poveda P, Pérez Simó G, Simarro Farinos J, Llavador Ros M, Palanca Suela S, Botella Estrada R. Telomerase Expression in a Series of Melanocytic Neoplasms. *Actas Dermosifiliogr (Engl Ed)*. 2019 Apr;110(3):212-219.
63. Salmi S, Siiskonen H, Sironen R, Tyynelä-Korhonen K, Hirschovits-Gerz B, Valkonen M, Auvinen P, Pasonen-Seppänen S. The number and localization of CD68+ and CD163+ macrophages in different stages of cutaneous melanoma. *Melanoma Res*. 2019 Jun;29(3):237-247.
64. Bsirini C, Smoller BR. Histologic mimics of malignant melanoma. *Singapore Med J*. 2018 Nov;59(11):602-607.

65. Colebatch AJ, Scolyer RA. Trajectories of premalignancy during the journey from melanocyte to melanoma. *Pathology*. 2018 Jan;50(1):16-23.
66. Tan SY, Strazzulla LC, Li X, Park JJ, Lee SJ, Kim CC. Association of clinicopathological features of melanoma with total naevus count and a history of dysplastic naevi: a cross-sectional retrospective study within an academic centre. *Clin Exp Dermatol*. 2018 Jul;43(5):566-572.
67. Winkelmann RR, Farberg AS, Glazer AM, Cockerell CJ, Sober AJ, Siegel DM, Leachman SA, High WA, Markowitz O, Berman B, Pariser DM, Goldenberg G, Rosen T, Rigel DS. Integrating Skin Cancer-Related Technologies into Clinical Practice. *Dermatol Clin*. 2017 Oct;35(4):565-576.
68. Madigan LM, Treyger G, Kohen LL. Compliance with serial dermoscopic monitoring: An academic perspective. *J Am Acad Dermatol*. 2016 Dec;75(6):1171-1175.
69. Raghavan D. Cutaneous manifestations of genitourinary malignancy. *Semin Oncol*. 2016 Jun;43(3):347-52.
70. Macneal P., Patel B.C. *Congenital Melanocytic Nevi*. StatPearls Publishing LLC; Treasure Island, FL, USA: 2022.
71. Kanada K.N., Merin M.R., Munden A., Friedlander S.F. A prospective study of cutaneous findings in newborns in the United States: Correlation with race, ethnicity, and gestational status using updated classification and nomenclature. *J. Pediatr*. 2012;161:240–245.
72. Alikhan A., Ibrahimi O.A., Eisen D.B. Congenital melanocytic nevi: Where are we now? Part I. Clinical presentation, epidemiology, pathogenesis, histology, malignant transformation, and neurocutaneous melanosis. *J. Am. Acad. Dermatol*. 2012;67:495.e1–495.e17.
73. Kadonaga J.N., Frieden I.J. Neurocutaneous melanosis: Definition and review of the literature. *J. Am. Acad. Dermatol*. 1991;24:747–755.
74. Agero A.L., Benvenuto-Andrade C., Dusza S.W., Halpern A.C., Marghoob A.A. Asymptomatic neurocutaneous melanocytosis in patients with large congenital melanocytic nevi: A study of cases from an Internet-based registry. *J. Am. Acad. Dermatol*. 2005;53:959–965.
75. Mosa A., Ho E.S., Heinelt M., Wong K., Neuhaus K. Management of congenital melanocytic nevi in the plastic surgery clinic: Families' expectations and their persistent concern about malignancy. *Pediatr. Dermatol*. 2019;36:876–881.
76. Jahnke M.N., O'Haver J., Gupta D., Hawryluk E.B., Finelt N., Kruse L., Jen M., Horii K.A., Frieden I.J., Price H., et al. Care of Congenital Melanocytic Nevi in Newborns and Infants: Review and Management Recommendations. *Pediatrics*. 2021;148:e2021051536.
77. Yélamos O., Arva N.C., Obregon R., Yazdan P., Wagner A., Guitart J., Gerami P. A comparative study of proliferative nodules and lethal melanomas in congenital nevi from children. *Am. J. Surg. Pathol*. 2015;39:405–415.
78. Kim J.Y., Kim S.M., Park K.D., Jang Y.H., Lee W.J., Lee S.J. A retrospective case series of 10 patients with malignant melanomas arising from small- and medium-sized congenital melanocytic nevi in South Koreans. *Indian J. Dermatol. Venereol. Leprol*. 2020;87:293–297.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.