Preprints.org

Article

Not peer-reviewed version

Proof of the Riemann Hypothesis

Yoshinori Shimizu *

Posted Date: 20 October 2025

doi: 10.20944/preprints202505.2110.v3

Keywords: Riemann hypothesis; Fredholm determinant; operator theory; analytic number theory

Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Proof of the Riemann Hypothesis

Yoshinori Shimizu

Independent Researcher; usagin.work@gmail.com

Abstract

This paper constructs two parallel approaches—Weil-type positivity and the Herglotz-type m-function—and connects them via a common core consisting of narrow-band equivalence ($\eta < \log 2$) and the uniqueness principle, thereby reaching the Riemann Hypothesis (RH) for the completed Riemann function ξ, and further establishing the Generalized Riemann Hypothesis GRH(π) for self-dual GL(d)-type L-functions. In the Weil route, we show that the measures μ_L and μ_ξ appearing on the operator side and the number-theoretic side of the distributionally normalized explicit formula agree in the narrow band (and, by densification, extend to $\mathcal{F}_{\log}$). Combining this with the known Weil equivalence theorem $Q_\xi \geq 0 \;\Leftrightarrow\;$ RH yields the RH Main Theorem (Theorem 8.23). In the Herglotz route, we construct, via a band-limited window Φ, the operator-side $m_L^{(\Phi)}$ and the number-theoretic side $M_\pi^{(\Phi)}$, and prove their equality over the entire complex plane by Poisson smoothing and the uniqueness of the Herglotz representation. From self-adjointness and the positivity of the Nevanlinna measure, we deduce that all nontrivial zeros lie on the critical line, arriving at the GRH(π) Main Theorem (Theorem 10.35 / Theorem 10.39). In the wide band, finite prime sums and endpoint contributions are absorbed into the regularized determinant $\det_2$ and its generating function. By precisely calibrating constants arising from the conductor, Archimedean terms, and the order of vanishing at the endpoints, we ensure robustness in error control. As applications, we show that the L-functions of Dirichlet characters, Hecke characters, holomorphic GL(2) cusp forms, and Maaß newforms satisfy axioms (AL1)–(AL5), and that GRH(π) follows immediately from the arguments in this chapter alone (Proposition 10.43, Corollary 10.44). Global conventions on the Fourier transform, boundary values, the Cayley transform, $\det_2$, and others are compiled in the appendix to ensure reproducibility and transparency in constant management.

Keywords: Riemann hypothesis; Fredholm determinant; operator theory; analytic number theory

---

1. Introduction

1.1. Background and Problem Setting [1–5]

Formulation of the Riemann Hypothesis (RH).

For the complex variable $s = \sigma + it$,

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} \qquad (\operatorname{Re} s > 1)$$

is extended by the usual analytic continuation to the whole complex plane (except for the pole at $s = 1$), and we use the completed form

$$\xi(s) \;:=\; \tfrac{1}{2}\, s(s-1)\, \pi^{-s/2}\, \Gamma\!\left(\tfrac{s}{2}\right) \zeta(s).$$

The function $\xi(s)$ is entire (of order ≤ 1) and satisfies the functional equation $\xi(s) = \xi(1-s)$. Let ρ be a nontrivial zero lying in the critical strip $0 < \operatorname{Re} s < 1$; then

$$\textbf{(RH)} \qquad \operatorname{Re} \rho = \tfrac{1}{2}.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

The purpose of this paper is to present the above statement concerning the distribution of the zeros of ξ as an equivalent framework via two distinct routes: the *Weil-type positivity* and the *Herglotz (m-function)* routes. For the latter, the scope is extended to the *Generalized Riemann Hypothesis* (GRH(π)) for the completed L-function $\Lambda(s,\pi)$ of a self-dual $GL(d)$-type general L-function (see Section 8 and Section 10 for details).

Framework for General L-functions (Outline).

For a self-dual $GL(d)$-type L-function $L(s,\pi)$, the associated completed form $\Lambda(s,\pi)$ contains the analytic conductor Q_π and Archimedean factors, and satisfies the functional equation $\Lambda(s,\pi) = \varepsilon_\pi\,\Lambda(1-s,\pi)$ with $|\varepsilon_\pi| = 1$. For the set of nontrivial zeros ρ_π,

$$\mathbf{GRH}(\pi) \qquad \operatorname{Re}\rho_\pi = \tfrac{1}{2}$$

is the object of study (for the axiomatic framework and normalization, see Section 10.1).

Test Space and Bilinear Form Used in This Paper (Preview of Their Roles).

We adopt the Fourier conventions

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\,e^{-it\lambda}\,dt, \qquad (f*g)(t) = \int_{\mathbb{R}} f(u)\,g(t-u)\,du, \qquad \widetilde{f}(t) = f(-t)$$

(the consistency with the Poisson/Hilbert formulas will be confirmed in the relevant sections of the main text). In v1.1, for even, real Schwartz functions, we adopt

$$\mathcal{F}_{\log} := \left\{ f \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} : \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2+|\lambda|)\,d\lambda < \infty \right\}$$

as the basic test space (rigorously defined in Section 8). With the "height measure" in the critical strip

$$\mu_\xi := \sum_\rho \delta_{\operatorname{Im}\rho} \quad \text{(counting multiplicity)},$$

we define

$$Q_\xi(f) := \langle \mu_\xi,\, f*\widetilde{f}\rangle = \left\langle \mu_\xi,\, |\widehat{f}|^2 \right\rangle, \qquad f \in \mathcal{F}_{\log}.$$

Then, by the *Weil equivalence*,

$$\boxed{Q_\xi(f) \ge 0 \ \ (\forall f \in \mathcal{F}_{\log}) \iff \text{RH}}$$

holds (see Section 8.4). In the first half of this paper (Section 6–Section 8), starting from the *small-bandwidth test functions*

$$A_\eta := \{\, \phi \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} : \operatorname{supp}\widehat{\phi} \subset [-\eta,\eta] \,\}, \quad \eta < \log 2,$$

we initiate from the *small-bandwidth equivalence* (agreement between the operator side and the arithmetic side) to extend $Q_\xi \ge 0$ to $\mathcal{F}_{\log}$ (by densification), and combine this with the equivalence theorem to conclude RH (Section 8.3–Section 8.4).

Two Routes (Placement in the Main Text).

- *Weil Positivity Route (Section 6–Section 8):* Starting from the *small-bandwidth agreement* ($\eta < \log 2$) between the operator side and the arithmetic side in the explicit formula arranged as a distribution, we densify $Q_\xi(f) \ge 0$ ($f \in A_\eta$) as $\eta \uparrow \log 2$ and $A_\eta \nearrow \mathcal{F}_{\log}$. By restating the known *Weil equivalence theorem* ($Q_\xi \ge 0 \iff$ RH) and combining, we arrive at the main theorem RH.
- *Herglotz Route (Section 10):* Constructing the operator-side $m_L^{(\Phi)}$ and the arithmetic-side $M_\pi^{(\Phi)}$ via a finite-bandwidth window Φ, we prove *agreement on the whole plane* from Poisson smoothing and

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

the uniqueness of the Herglotz representation. From self-adjointness and the positivity of the Nevanlinna measure, we deduce the *real-axis nature of poles* and establish GRH(π) as the main theorem. RH is recovered as the special case of ζ (ξ).

Appendices, Conventions, and Reproducibility.

The global conventions for the Fourier transform, boundary values, Cayley transform, and the regularized determinant $\det_2$ used in this paper are consolidated in the appendices, so that the assumptions, normalizations, and coefficient correspondences on which each chapter's statements depend can be referenced in an auditable form. In particular, the choice of small bandwidth $\eta_0 < \log 2$, the finite part (normalization of distributions), and constants in the Riemann–von Mangoldt type main term are unified via correspondence tables in the appendices and cross-references with the main text.

1.2. Main Results of This Paper (Summary) [1–3]

This paper reaches the Riemann Hypothesis (RH) and the Generalized Riemann Hypothesis (GRH(π)) for self-dual $GL(d)$-type L-functions via two routes: *Weil-type positivity* and *Herglotz (m-function)*. Under the notation fixed in Section 1.1 (in particular, the test space $\mathcal{F}_{\log}$ and the bilinear form Q_ξ), the main conclusions are summarized in the following three points.

(A) Main Theorem on RH (Weil Route; Conclusion of Section 8).

The *Weil-type bilinear form* $Q_\xi(f)$ on $\mathcal{F}_{\log}$ is always nonnegative, and by combining with the known "Weil equivalence theorem," RH is concluded:

$$\boxed{\forall f \in \mathcal{F}_{\log} \;\Rightarrow\; Q_\xi(f) \geq 0 \iff \mathrm{RH}}$$

(Theorem 8.19 in Section 8.3 establishes $Q_\xi \geq 0$, and by combining with Theorem 8.21 (equivalence theorem) in Section 8.4, we obtain the *Main Theorem on RH* = Theorem 8.23.) For the construction of Q_ξ and the definition of $\mathcal{F}_{\log}$, see Section 8.2–Section 8.3.

(B) Main Theorem on RH (Herglotz Route; Conclusion of Section 8).

From the construction of the m-function using a finite-bandwidth "window" Φ and from Poisson smoothing / uniqueness of the Herglotz representation, we prove the *equality of windowed m-functions* $m_L^{(\Phi)} \equiv -\xi'/\xi$. Through the analysis of its zeros and poles (Stieltjes inversion and pole distribution), we exclude *non-real poles* and reach RH (Section 8.6–Section 8.7; the final conclusion coincides with the Main Theorem on RH in Section 8). This route shares the same *small-bandwidth equivalence* $\to$ *uniqueness principle* as (A) and is mutually reinforcing.

(C) Generalization (Self-dual $GL(d)$-type L-functions; Conclusion of Section 10).

Under the *general axiomatic framework* ((AL1)–(AL5) in Section 10.1), the generalized bilinear form Q_π on $\mathcal{F}_{\log}$ is always nonnegative (Theorem 10.38), and by the *Weil equivalence theorem (generalized form)*,

$$\boxed{\forall f \in \mathcal{F}_{\log} \;\Rightarrow\; Q_\pi(f) \geq 0 \iff \mathrm{GRH}(\pi)}$$

holds (Theorem 10.39). At the same time, in the Herglotz route (Section 10.6–Section 10.7), GRH(π) is obtained from the equality of *windowed m-functions* $M_\pi^{(\Phi)}$ and the reality of poles (Theorem 10.35). Thus, the two systems, Weil / Herglotz, close consistently for both the ξ case and the general $\Lambda(s, \pi)$ case.

(D) *Immediate Application* to Specific Classes.

The four classes — Dirichlet ($d = 1$), Dedekind of number fields, Hecke characters, and self-dual $GL(2)$ newforms (elliptic modular / Maaß) — satisfy the axioms (AL1)–(AL5) of Section 10.1

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Proposition 10.43). Therefore, the above main theorems (Theorem 10.35 / 10.39) apply immediately, and $GRH(\pi)$ *holds within the discussion of this chapter alone* (Corollary 10.44).

(E) Robustness (Error Budget and Tolerances).

We organize the reduction of bandwidth ($\eta \uparrow \log 2$), evaluation of endpoint contributions, order control of $\det_2$, and minimization of assumptions for the uniqueness principle, and visualize the *allowable error* from three aspects: small bandwidth / large bandwidth / generating functions (Section 8.5 and Section 10.10). Dependencies of constants in implementation (conductor Q_π, Archimedean factors, vanishing orders at endpoints, etc.) are consolidated in correspondence tables (Appendix A).

(F) Summary of This Section (Correspondence with Chapter Structure).

(A)–(B) provide RH in the conclusion of Section 8 (Theorem 8.23), and (C)–(D) provide GRH(π) in the conclusion of Section 10 (Theorem 10.35 / 10.39 and Proposition 10.43 / Corollary 10.44). (E) is the *integration of robustness* in Section 8.5 and Section 10.10, ensuring the portability of the overall strategy (small-bandwidth equivalence $\to$ densification / uniqueness $\to$ Weil / Herglotz).

1.3. Proof Strategy (Overview of Two Routes) [1,2,6]

This paper constructs in parallel two distinct routes — *Weil-type positivity* and *Herglotz (m-function)* — each leading to RH (the ξ case), and further to GRH(π) (general case) for general self-dual $GL(d)$-type $\Lambda(s,\pi)$. These are connected by a common core of *small-bandwidth equivalence* and the *uniqueness principle*, while prime finite sums and endpoint contributions that appear in the large bandwidth regime are absorbed and controlled within the framework of the *regularized determinant* $\det_2$ and generating functions. Below, the inputs, outputs, and key points are stated explicitly.

(I) Weil Positivity Route: Composition of $Q \geq 0$ and the Equivalence Theorem (Section 6–Section 8).

First, for the finite-bandwidth test functions

$$A_\eta \;:=\; \{\, \phi \in \mathcal{S}(\mathbb{R}) \text{ even, real} :\ \operatorname{supp} \widehat{\phi} \subset [-\eta,\eta] \,\}, \qquad 0 < \eta < \log 2,$$

we establish *small-bandwidth equivalence* between the operator side and the arithmetic side (Section 6). Then, through the limiting process $\eta \uparrow \log 2$ and dominated convergence (including evaluation of the finite part), we *densify* to $\mathcal{F}_{\log}$ (defined in Section 8.2) to obtain

$$Q_\xi(f) \;:=\; \langle \mu_\xi,\ f * \widetilde{f} \rangle \;=\; \langle \mu_\xi,\ |\widehat{f}|^2 \rangle \;\geq 0 \qquad (\forall f \in \mathcal{F}_{\log})$$

(Section 8.3). Finally, we restate the *Weil equivalence theorem* (within the framework of this paper) and combine:

$$Q_\xi(f) \geq 0 \ (\forall f \in \mathcal{F}_{\log}) \iff \text{RH}$$

to conclude RH (Section 8.4).
Input/Output:

$$\boxed{\text{Small-bandwidth equivalence (Section 6)}} \xrightarrow[\eta\uparrow\log 2]{\text{Densification (Section 8.3)}} \boxed{Q_\xi \geq 0 \text{ on } \mathcal{F}_{\log}} \xrightarrow[]{\text{Weil equivalence (Section 8.4)}} \boxed{\text{RH}}.$$

(II) Herglotz Route: Equality of Windowed m-functions and Reality of Poles (Section 8H, Section 10.6–Section 10.7).

Fix a finite-bandwidth "window" Φ (time-side convolution kernel $W_\Phi = \check{\Phi} * \check{\Phi}^{\sim}$), and construct the *operator-side* Weyl–Titchmarsh type m-function $m_L^{(\Phi)}$ and the *arithmetic-side* Herglotz function $M^{(\Phi)}$ via the zero measure μ. By combining the *small-bandwidth equivalence* of Section 6 with the *uniqueness principle* of Section 8.1–Section 8.2, we show that the two agree *on the whole plane up to a polynomial difference*, and remove this difference from the asymptotics at infinity. Through Herglotz property (positivity of the Nevanlinna measure), we deduce the *reality of poles* and obtain RH (the ξ case) / GRH(π) (general case).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Input/Output:

$$\boxed{\text{Small-bandwidth equivalence (Section 6) + Uniqueness (Section 8.1–Section 8.2)}} \Longrightarrow \boxed{m_{\mathrm{op}}^{(\Phi)} \equiv M_{\mathrm{arith}}^{(\Phi)}}$$
$$\Longrightarrow \boxed{\text{Poles are on the real axis}} \Longrightarrow \boxed{\text{RH / GRH}(\pi)}.$$

(III) Extension to General (Self-dual) $GL(d)$ and Consistency of the Two Routes (Section 10).

For self-dual $GL(d)$ types satisfying the general axioms (AL1)–(AL5) (Section 10.1), the *Weil route* shows the equivalence of $Q_\pi \geq 0$ and GRH(π) (Section 10.8), and the *Herglotz route* shows the reality of poles from equality of windowed m-functions (Section 10.6–Section 10.7). Both routes are connected by a single skeleton: *small-bandwidth equivalence* $\Rightarrow$ *large-bandwidth difference (prime finite sum + endpoint term)* $\Rightarrow$ $\det_2$ *coefficient identification* $\Rightarrow$ *Herglotz/Weil* (including generating functions and functional calculus in Section 10.3–Section 10.5). Dirichlet / Hecke / Dedekind / self-dual $GL(2)$ satisfy the axioms (Proposition 10.43), and by Theorem 10.35 (Herglotz) or Theorem 10.39 (Weil), *GRH*(π) *holds from the discussion within this chapter alone* (Corollary 10.44).

(IV) Common Core and Error Management (Section 6.4, Section 8.1–Section 8.2, Section 10.10, Appendix A).

- **Small bandwidth** $\eta < \log 2$ **and endpoint** $\eta = \log 2$**:** In the small bandwidth regime, prime finite sums vanish (except for the $p = 2$ endpoint), and endpoint contributions are explicitly controlled by the half-rule (incorporated into the remainder if necessary).
- **Uniqueness principle:** Serves as the key to lift from *family-uniform vanishing* of the shrunken bandwidth family and *small disk agreement* to agreement on the whole domain.
- $\det_2$ **and generating functions:** Constrain large-bandwidth differences in both *order* and *coefficient*, and provide the uniform bounds needed for m-function identification and transport of Q.
- **Robustness:** Integrates error budgets from the three aspects — small bandwidth / large bandwidth / generating functions — and makes constant dependencies explicit (conductor $Q(\pi)$, vanishing order at endpoints, Weyl error, order of $\det_2$).

(V) Summary (Relation to the Whole Paper).

(I) combines positivity in Section 8.3 (Theorem 8.19) and the equivalence theorem in Section 8.4 (Theorem 8.21) to reach RH (Theorem 8.23); (II) provides m-function equality $\Rightarrow$ reality of poles $\Rightarrow$ RH/GRH(π) in Section 8H and Section 10.6–Section 10.7; (III) connects Section 10.8 (Weil route) and Section 10.6–Section 10.7 (Herglotz route) via the *same skeleton*, and (IV) guarantees its *portability and error management*.

1.4. Technical Essentials and Notation [2,6,7]

In this section, we fix in advance the conventions, symbols, and basic objects used throughout the paper. We collectively define the Fourier conventions, test spaces (finite-bandwidth family A_η and basic space $\mathcal{F}_{\log}$), the bilinear form Q, the Herglotz (m-function) / Cayley phase, and the regularized determinant $\det_2$. Each item will be restated and refined in later rigorous sections (Section 6–Section 10), but here we present them in a minimally self-contained form to facilitate back-and-forth referencing.

Fourier Conventions and Basic Operations.

We unify the Fourier transform, convolution, and reflection as

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\, e^{-it\lambda}\, dt, \qquad (f * g)(t) = \int_{\mathbb{R}} f(u)\, g(t-u)\, du, \qquad \widetilde{f}(t) = f(-t) \tag{1}$$

(the inverse transform is denoted by $\mathbb{F}^{-1}$, adopting the normalization consistent with (1)). Unless otherwise stated, test functions are assumed to be "even and real." Global auxiliary conventions

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Poisson/Hilbert kernels, boundary values, continuous connection of phases, etc.) are consolidated in the appendix.

Finite-Bandwidth Family A_η and Basic Test Space $\mathcal{F}_{\log}$.

For bandwidth $0 < \eta < \log 2$, we define

$$A_\eta := \left\{ \phi \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} : \operatorname{supp} \widehat{\phi} \subset [-\eta, \eta] \right\} \tag{2}$$

(*small bandwidth*). Also, we define the basic test space $\mathcal{F}_{\log}$ by

$$\mathcal{F}_{\log} := \left\{ f \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} : \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty \right\} \tag{3}$$

The bandwidth cutoff is taken using a smooth even cutoff $m_\eta \in C_c^\infty([-\eta, \eta])$, setting $\widehat{f_\eta} := m_\eta \widehat{f}$ and $f_\eta := \mathbb{F}^{-1}\widehat{f_\eta}$ (so $f_\eta \in A_\eta$). A_η becomes dense in $\mathcal{F}_{\log}$ as $\eta \uparrow \log 2$ (see Section 8).

Zero Measure and Bilinear Form Q.

From the imaginary parts of the nontrivial zeros $\rho = \frac{1}{2} + i\gamma$ of ξ, define $\mu_\xi := \sum_\rho \delta_\gamma$, and from the imaginary parts of the zeros of the completed form $\Lambda(s, \pi)$ associated with a self-dual $GL(d)$-type π, define μ_{Ξ_π} (counting multiplicity; the generalized finite part is taken according to the convention in the main text). For any $f \in \mathcal{F}_{\log}$, define

$$Q_\xi(f) := \langle \mu_\xi,\, f * \widetilde{f} \rangle = \langle \mu_\xi,\, |\widehat{f}|^2 \rangle, \qquad Q_\pi(f) := \langle \mu_{\Xi_\pi},\, |\widehat{f}|^2 \rangle \tag{4}$$

(where $\langle \cdot, \cdot \rangle$ denotes the duality between distributions and test functions). As a property of closure under small bandwidth, if $f \in A_\eta$ then $f * \widetilde{f} \in A_\eta$. In this paper, through *small-bandwidth equivalence* in Section 6 and *densification* in Section 8, we establish $Q \geq 0$ and combine it with the Weil equivalence to obtain RH/GRH (Section 8, Section 10).

Windowed m-function and Cayley Phase.

For a finite-bandwidth "window" Φ, define the time-side convolution kernel by

$$W_\Phi := \check{\Phi} * \widetilde{\check{\Phi}} \qquad (\check{\Phi} := \mathbb{F}^{-1}\Phi) \tag{5}$$

On the operator side, we construct the Weyl–Titchmarsh type m-function $m_L^{(\Phi)}$, and on the arithmetic side, the Herglotz function $M_\pi^{(\Phi)}$ based on the zero measure (Section 8, Section 10), and prove their equality from *small-bandwidth equivalence* + *uniqueness principle*. Boundary values are written as $M(t + i0)$ for the nontangential limit from the upper half-plane, and the Cayley transform

$$S(t) = \frac{1 - i\, M(t + i0)}{1 + i\, M(t + i0)} = e^{-i\phi(t)} \tag{6}$$

defines the phase ϕ (jump by π at poles; the phase is extended monotonically by continuously connecting the principal value).

Regularized Determinant $\det_2$.

For a self-adjoint operator K of Hilbert–Schmidt class, define the regularized determinant by

$$\det_2(I - zK) := \prod_j (1 - z\lambda_j)\, e^{z\lambda_j} \tag{7}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

where $\{\lambda_j\}$ are the eigenvalues (Weierstrass factor of genus 1). Its derivative is

$$\partial_z \log \det_2(I - zK) \;=\; -\sum_j \frac{\lambda_j}{1 - z\lambda_j} \;+\; \operatorname{Tr} K \tag{8}$$

(the normalization includes $-\operatorname{Tr} K$), from which order ≤ 2 follows. In §7, we present the consistency between the zero distribution of $\det_2$ and the Weyl main term / HS condition, and the design of upper bounds (shrinking bandwidth family).

Small-Bandwidth Endpoints and Management of Endpoint Terms.

The difference at the bandwidth endpoint η is decomposed into "prime finite sum + endpoint term" (Section 10.1). For smooth windows ($\Phi \in C_c^\infty$), endpoint terms vanish by vanishing conditions of arbitrary order, and even for piecewise smooth windows they are boundedly controlled by vanishing of endpoint values and higher derivatives (constants depend on the conductor and the endpoint order). This endpoint management is essential for transfer from small bandwidth $\eta < \log 2$ to large bandwidth (densification / uniqueness).

Remarks (Summary of Symbols).

$\widehat{\cdot}$: Fourier, $\widetilde{f}$: reflection, $\check{\Phi}$: inverse transform, $\mathcal{F}_{\log}$: (14), A_η: (2), Q_ξ, Q_π: (4), W_Φ: (5), $S = e^{-i\phi}$: (6), $\det_2$: (29)–(8). Hereafter, these symbols are used consistently with *the same normalization*.

1.5. Scope of Application [3,8–13]

The main theorems of the two routes in this paper (Weil positivity / Herglotz) apply to *self-dual* $GL(d)$-type L-functions satisfying the axioms (AL1)–(AL5) in Section 10.1 (analytic continuation, functional equation, Euler product, normalization of Archimedean factors, and good behavior of the conductor). Here we list the representative classes actually treated in this paper and the key points to be confirmed for application (see Section 10.8–Section 10.10, Proposition 10.43, Corollary 10.44 for details). Notation and conventions follow Section 1.4.

(i) Riemann ζ and its completion ξ ($d = 1$, basic example).

The completed form

$$\xi(s) = \tfrac{1}{2}\, s(s-1)\, \pi^{-s/2} \Gamma\!\left(\tfrac{s}{2}\right) \zeta(s)$$

satisfies (AL1)–(AL5) and is self-dual. In Section 8, from small-bandwidth equivalence $\Rightarrow$ densification we obtain $Q_\xi \ge 0$, and by combining with the Weil equivalence (Section 8.4) we conclude RH (Theorem 8.23). In the Herglotz route as well, we arrive at the same conclusion from equality of windowed m-functions and reality of poles (Section 8.6–Section 8.7).

(ii) Dirichlet L ($d = 1$, real (self-conjugate) characters).

For a primitive real Dirichlet character χ,

$$\Lambda(s,\chi) = \left(\frac{q}{\pi}\right)^{(s+\kappa)/2} \Gamma\!\left(\frac{s+\kappa}{2}\right) L(s,\chi) \quad (\kappa \in \{0,1\})$$

is self-dual and satisfies (AL1)–(AL5). Therefore, by the main theorem in Section 10,

$$Q_\pi \ge 0 \iff \mathrm{GRH}(\pi) \quad (\pi \leftrightarrow \chi)$$

holds (Theorem 10.39), and $\mathrm{GRH}(\pi)$ also follows from the Herglotz route (Theorem 10.35). For non-self-conjugate characters (complex χ), since in this paper the statements are made under the self-duality assumption, we refer, when necessary, to standard reductions such as self-dualization by $\pi \oplus \tilde{\pi}$ or symmetric square, etc. (see Section 10.9).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(iii) Dedekind ζ_K of a number field K ($d = 1$, special case of Hecke characters).

The completed form

$$\Lambda(s, \zeta_K) \;=\; Q_K^{s/2} \prod_{j=1}^{d} \Gamma_{\mathbb{R}}(s + \mu_j)\, \zeta_K(s)$$

is self-dual and fits (AL1)–(AL5) (here Q_K is the conductor from the discriminant, and $\Gamma_{\mathbb{R}}$ is the Archimedean factor determined by the combination of real and complex embeddings). Hence by the main theorem in Section 10, GRH(ζ_K) follows (Proposition 10.43, Corollary 10.44).

(iv) Hecke characters ($d = 1$, self-conjugate unitary idele class characters over number fields).

For a unitary Hecke character ω (self-conjugate), the associated $L(s, \omega)$ has a completed form $\Lambda(s, \omega)$ that is self-dual and satisfies (AL1)–(AL5). Thus both Theorem 10.35 and Theorem 10.39 apply, and GRH(π) ($\pi \leftrightarrow \omega$) is established within this chapter.

(v) Self-dual $GL(2)$ newforms (automorphic forms: holomorphic / Maaß).

For a self-dual automorphic representation π of level N with trivial central character (holomorphic newform of weight k or Maaß newform), the standard L-function

$$\Lambda(s, \pi) \;=\; Q_\pi^{s/2} \prod_{j=1}^{2} \Gamma_{\mathbb{R}}(s + \mu_j)\, L(s, \pi)$$

satisfies (AL1)–(AL5). In the Weil route, $Q_\pi \geq 0 \iff$ GRH(π) (Theorem 10.39); in the Herglotz route, from equality of windowed $M_\pi^{(\Phi)}$ and the reality of poles, we obtain GRH(π) (Theorem 10.35).

(vi) Error / Constant Dependence and Appendix References.

In any class, contributions from bandwidth endpoints, the degree of the conductor Q_π and Archimedean factors, the order estimates of $\det_2$, and the vanishing order in Poisson smoothing appear in the error constants. This paper summarizes the integration of errors in Section 10.10, and correspondence tables of constants and normalizations are consolidated in the Appendix A.

Summary.

Each of the classes (i)–(v) satisfies the assumptions of Proposition 10.43, and by Corollary 10.44, *GRH(π) follows immediately from the discussion within this chapter alone*. The ζ case (i) coincides with the main theorem on RH in Section 8, and for general self-dual $GL(d)$ ($d \geq 2$) the two routes in Section 10 complete the proof.

1.6. Organization of This Paper

This paper is structured around two main pillars: the *Weil positivity route* ($Q \geq 0 \Rightarrow$ RH/GRH) and the *Herglotz (m-function) route* (equality of windowed m-functions $\Rightarrow$ reality of poles $\Rightarrow$ RH/GRH). The roles of each chapter and their contributions to the two routes are as follows (the notation and conventions fixed in Section 1 are common to all chapters, and the appendices centrally manage reference tables and normalizations).

Section 2 (Foundations: Conventions, Distributions, Kernels).

We organize the minimal tools of distribution theory and the conventions for Fourier/Poisson/Hilbert, and fix the framework for treating the explicit formula as a distribution. We standardize the window Φ and convolution kernel, finite part (principal value / finite part), and the method of taking boundary values, providing a *common base language* for subsequent small-bandwidth equivalence and m-function construction.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Section 3 (Distributional Form of the Explicit Formula and Preparation for Small Bandwidth).

We rewrite prime sums and zero measures in the same distributional framework, and for families of finite-bandwidth test functions prepare the decomposition "difference from large bandwidth = prime finite sum + endpoint term." This decomposition serves as the input for *small-bandwidth equivalence* in Section 6 and as groundwork for *absorption via* $\det_2$ in §7.

Section 4 (Operator Side I: Convolution Operators and Regularization).

We bring convolution operators obtained from finite-bandwidth kernels into the Hilbert–Schmidt class, and establish the framework of the *regularized determinant* $\det_2$. We present Weierstrass factorization, order estimates, consistency between the Weyl main term and $\det_2$, and introduce a *mechanism on the generating function side to constrain difference terms* appearing in the large-bandwidth regime.

Section 5 (Operator Side II: m-functions and the Uniqueness Principle).

We construct the Weyl–Titchmarsh type *windowed m-function* $m^{(\Phi)}$, and via the Herglotz representation and the Cayley transform (phase ϕ), prepare the analytic structure compatible with *reality of poles*. At the same time, we establish the *uniqueness principle* based on small-disk agreement and family-uniform vanishing, providing the logical core for m-function identification in Section 8 and Section 10.

Section 6 (Small-Bandwidth Equivalence).

We prove the *small-bandwidth agreement* ($\eta < \log 2$) between the operator side and the arithmetic side. We make explicit the half-rule for endpoint contributions and the disappearance of prime finite sums (except $p = 2$), positioning this as the starting point of $Q \geq 0$ in the Weil route and as the *origin* of m-function equality in the Herglotz route.

§7 ($\det_2$, Weyl, and Upper Bound Design).

We give the consistency between the zero distribution of $\det_2$ and the Weyl main term, and design *uniform upper bounds* for shrinking-bandwidth families. This shows that differences (prime finite sum + endpoint term) arising in large bandwidth can be absorbed on the generating function side, and prepares the *framework for the error budget* needed for densification / uniqueness in Section 8 and Section 10.

Section 8 (The ξ Case: Main Theorem on RH).

We define the basic space $\mathcal{F}_{\log}$ and, combining the small-bandwidth equivalence of Section 6 with the upper bounds of §7, establish $Q_\xi \geq 0$ (Theorem 8.19). By combining with the known Weil equivalence (Theorem 8.21), we obtain the *Main Theorem on RH* (Theorem 8.23). In parallel, we complete the Herglotz route in the ξ case (equality of windowed $m \Rightarrow$ reality of poles), showing that the two routes *converge to the same point*.

Section 9 (Conclusions and Guidelines: Overview of Robustness and Applications).

We organize the *integration of errors* from the three aspects of small bandwidth / large bandwidth / generating functions, and visualize constant dependencies (conductor, endpoint vanishing order, order of $\det_2$).
We also provide an overview of the application policy toward Section 10 for Dirichlet / Hecke / Dedekind / self-dual $GL(2)$, serving as a bridge from the *ζ case (Section 8) to the general case (Section 10)*.

Section 10 (Generalization: Self-dual $GL(d)$-type L-functions).

Under axioms (AL1)–(AL5), in the Weil route we have $Q_\pi \geq 0 \iff \mathrm{GRH}(\pi)$ (Theorem 10.39), and in the Herglotz route we deduce the reality of poles from equality of windowed $M_\pi^{(\Phi)}$ (Theorem 10.35). Proposition 10.43 confirms that Dirichlet / Hecke / Dedekind / self-dual $GL(2)$ satisfy the

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

axioms, and together with Corollary 10.44 concludes that $GRH(\pi)$ *holds from the discussion within this chapter alone*. Finally, Section 10.10 completes the *integration of errors* for the general case.

Appendices.

We present Fourier conventions, boundary values, Cayley phase, normalization of $\det_2$, vanishing conditions for endpoint terms, and *correspondence tables* of conductors, Archimedean terms, and main term constants. Symbols and constants in the main text can be traced in a unified manner through the tables in the appendices.

2. Space and Generator

2.1. Introduction of Space and Generator [7,14]

Important Note (Regarding the Double Definition of R_{PW})

In this chapter (Section 2), the symbol R_{PW}, representing the *Paley–Wiener type region*, is defined from two viewpoints:

(Op) **Operator-side definition** $R_{\mathrm{PW}}^{\mathrm{op}}$: Based on the Paley–Wiener type image obtained from the convolution operator arising from a finite-bandwidth kernel (coming from the window Φ) (Fourier conventions, boundary values, and finite part handling follow the conventions of Section 2).

(Ar) **Arithmetic-side definition** $R_{\mathrm{PW}}^{\mathrm{arith}}$: Based on the Paley–Wiener type image induced by arranging the explicit formula as a distribution and by the *small-bandwidth* test function family A_η ($\eta < \log 2$) and its limit yielding the basic space (see Section 1.4, Section 8).

This double definition is *intentional*, in order to naturally guarantee both the extension of the scope of application and the proof techniques (Weil positivity / Herglotz). The following points are made explicit:

(1) **Absence of contradiction**: On the core generated by the small-bandwidth family A_η fixed in Section 1.4,

$$R_{\mathrm{PW}}^{\mathrm{core}} := \overline{\bigcup_{0<\eta<\log 2} A_\eta}^{\text{natural topology}}$$

we have *identical agreement* between $R_{\mathrm{PW}}^{\mathrm{op}}$ and $R_{\mathrm{PW}}^{\mathrm{arith}}$. Therefore, the claims, constants, and estimates used in this paper *do not depend on the choice of definition*.

(2) **Notation policy**: Hereafter, unless otherwise stated, R_{PW} denotes the object obtained by *naturally identifying* the two. Only when it is necessary to emphasize a specific construction will the superscripts $R_{\mathrm{PW}}^{\mathrm{op}}$ / $R_{\mathrm{PW}}^{\mathrm{arith}}$ be used.

(3) **Extension and uniqueness**: By the small-bandwidth equivalence (Section 6) and the uniqueness principle (Section 5), the identical identification on $R_{\mathrm{PW}}^{\mathrm{core}}$ is *continuously extended* to both constructions. Differences in normalization concerning endpoint contributions and regularization ($\det_2$) follow the correspondence tables in the Appendix A and are consistent in either route (Weil / Herglotz).

From the above, we emphasize that the definitions and propositions in this chapter have a *single meaning* regardless of the choice of R_{PW}.

Positioning of this subsection (Relation to the overall strategy).

This section corresponds to the strategy of the paper and provides the *foundation* that connects to the subsequent self-adjointization, Hilbert–Schmidt inclusion, compactness of the resolvent (Section 2.3–Section 2.4), and further to the Weyl-type main term in Section 3 and the explicit formula in Section 6. Notation follows Section 1.4 (Fourier conventions, etc.).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Basic setting and fixed notation

Hereafter, we take as fixed parameters $\alpha > \frac{1}{2}$ and $\Lambda > 0$. With the weight

$$w(\tau) := \langle \tau \rangle^{2\alpha} = (1+\tau^2)^{\alpha}$$

we define the weighted Hilbert space

$$H_\alpha := \Big\{ f \in L^2(\mathbb{R}) : \ \|f\|_{H_\alpha}^2 := \int_{\mathbb{R}} w(\tau)\, |f(\tau)|^2 \, d\tau < \infty \Big\}.$$

The Fourier transform conventions follow Section 1.4:

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(\tau)\, e^{-i\tau\xi}\, d\tau, \qquad f(\tau) = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{f}(\xi)\, e^{i\tau\xi}\, d\xi.$$

The bandwidth space

$$PW_\Lambda := \{ f \in L^2(\mathbb{R}) : \ \operatorname{supp} \widehat{f} \subset [-\Lambda, \Lambda] \}$$

is defined as a closed subspace of $L^2(\mathbb{R})$ (hereafter, definitions and reasoning for operators are performed on $PW_\Lambda \cap H_\alpha$ as needed). The frequency cut-off projection

$$P_\Lambda := \mathbb{F}^{-1}\, \mathbf{1}_{[-\Lambda,\Lambda]}\, \mathbb{F}$$

is the (self-adjoint) orthogonal projection on $L^2(\mathbb{R})$. The weight multiplication operator is

$$(Uf)(\tau) := w(\tau)^{1/2} f(\tau)$$

and we standardize to denote it by U (consistent with N2 in Section 1.3). As a *core*,

$$\mathcal{C} := \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$$

is used. The differential operators are

$$R := -\partial_\tau, \qquad \mathrm{Re} := R - \frac{w'}{2w}\, I$$

(where Re denotes the notation for formal adjointization with respect to the H_α inner product). As a candidate for the generator,

$$L := -\, i\, R_{PW}$$

is introduced, where R_{PW} denotes the appropriate self-adjoint restriction (closure) of R to PW_Λ (definition clarified in the lemma below).

Strong commutativity on the Fourier side and restricted operator

Lemma 1 (Strong commutativity and essential self-adjointness of the restriction)**.** *Let $\alpha > \frac{1}{2}$, $\Lambda > 0$. $R = -\partial_\tau$ is self-adjoint on $L^2(\mathbb{R})$ and $\mathbb{F} R \mathbb{F}^{-1} = i\, M_\xi$ (multiplication operator). Therefore, R strongly commutes with P_Λ, and the restriction*

$$R|_{PW_\Lambda} : \ D\big(R|_{PW_\Lambda}\big) \subset PW_\Lambda \longrightarrow PW_\Lambda, \qquad D\big(R|_{PW_\Lambda}\big) = \{ f \in PW_\Lambda : \ \xi\, \widehat{f}(\xi) \in L^2 \}$$

is essentially self-adjoint. The closure is denoted by R_{PW}.

Proof. $\mathbb{F} R \mathbb{F}^{-1} = i M_\xi$ is standard. P_Λ coincides with $M_{\mathbf{1}_{[-\Lambda,\Lambda]}}$ and therefore strongly commutes with M_ξ. From strong commutativity, the self-adjointness of the multiplication operator iM_ξ on the closed set $[-\Lambda, \Lambda]$ is restricted to PW_Λ (spectral theorem). Therefore $R|_{PW_\Lambda}$ is essentially self-adjoint, and its closure R_{PW} is self-adjoint. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Equivalence of the two representations (in the sense of unitary equivalence)

Lemma 2 (Equivalence of the two representations of R_{PW}). *Let $\alpha > \frac{1}{2}$, $\Lambda > 0$, and on the common core $\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ consider*

$$S_0 f := -\partial_\tau f, \qquad S_1 f := -\partial_\tau f - \frac{w'}{2w} f.$$

The map $U : H_\alpha \to L^2(\mathbb{R})$ is an isometry (unitary), and

$$U \operatorname{Re} U^{-1} = R \quad \textit{(operator equality; in the distributional sense)}$$

holds. In particular,

(i) R_{PW} is uniquely determined by Lemma 1, and the generator $L := -iR_{PW}$ is self-adjoint;

(ii) $U^{-1} R_{PW} U$ is a self-adjoint operator on H_α, and its differential representation coincides with that given by Re *(equal to S_1 on $\mathcal{C}$).*

Therefore, the restriction on the Fourier side *($R|_{PW_\Lambda}$) and the* adjointized representation on the real side *(*Re*) give, in the sense of unitary equivalence, the same self-adjoint generator.*

Proof. U is a unitary between H_α and L^2, and by distributional calculation

$$U \left(-\partial_\tau - \frac{w'}{2w} \right) U^{-1} \;=\; -\partial_\tau$$

holds (boundary terms vanish in integration by parts on $\mathcal{C}$). Therefore, Re and R are unitarily equivalent. By Lemma 1, $R|_{PW_\Lambda}$ is essentially self-adjoint and its closure R_{PW} is self-adjoint. By unitary equivalence, $U^{-1} R_{PW} U$ is also self-adjoint, and its differential representation on $\mathcal{C}$ coincides with S_1. The claim follows. □

Remark 1 (Note on bandwidth preservation). *Multiplication on the time side, $Uf = w^{1/2} f$, does not in general preserve bandwidth, so it is not necessary to assume $U(PW_\Lambda) \subset PW_\Lambda$. In this paper, as in Lemma 2, we represent R_{PW} from two perspectives in terms of* unitary equivalence*, and all properties used in the subsequent self-adjointization (Section 2.3), functional calculus (Section 4), and explicit formula (Section 6) — such as spectral properties, counting, and Schatten class properties — are treated as unitary invariants.*

Standard notation and references used hereafter

Hereafter, we fix the notation

$$\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda, \qquad R_{PW} = \overline{R|_{PW_\Lambda}}, \qquad L := -iR_{PW}.$$

References to $\varphi(L)$ via Borel functional calculus (Section 4), $\mathcal{S}_p$ and $\det_2(I \pm zK)$ (Section 4.3), and to the small-bandwidth explicit formula and difference distributions (Section 6.1–Section 6.3) are all made according to the conventions in Section 1.4.
With this, the preparations for the remainder of Section 2 (domain, graph norm and self-adjointization, Hilbert–Schmidt inclusion of embeddings) are complete.

2.2. Domain and Graph Norm

Position of this subsection (relation to overall strategy).

This subsection deals with the *refinement of the domain* and the *foundation of the graph norm* for the generator candidate $L = -iR_{PW}$ introduced in Section 2.1 (Lemmas 1, 2), in order to establish the subsequent self-adjointness (Section 2.3) and the Hilbert–Schmidt property of the embedding (Section 2.4). In particular, we explicitly show that the graph norm defined using Re $:= R - \frac{w'}{2w} I$ (Section 2.1) is equivalent to H^1_α, and that $\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ is a *core*.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Definition (minimal domain and maximal domain)

Hereafter, fix $\alpha > \frac{1}{2}$, $\Lambda > 0$. With the weight $w(\tau) = (1+\tau^2)^\alpha$,

$$R := -\partial_\tau, \qquad \mathrm{Re} := R - \frac{w'}{2w} I$$

and $\mathcal{C} := \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ as the common dense subspace.

Definition 1 (Minimal domain and maximal domain). *The* minimal domain *of* Re *is defined by*

$$D_{\min}(\mathrm{Re}) := \overline{\mathcal{C}}^{\|\cdot\|_{\mathrm{gr}}}, \qquad \|f\|_{\mathrm{gr}}^2 := \|f\|_{H_\alpha}^2 + \|\mathrm{Re} f\|_{H_\alpha}^2,$$

that is, the closure of $\mathcal{C}$ *with respect to the graph norm. Furthermore, the* maximal domain *is*

$$D_{\max}(\mathrm{Re}) := \{ f \in H_\alpha : \mathrm{Re} f \in H_\alpha \}.$$

The domain of the restricted operator Re_{PW} *is written as* $D(\mathrm{Re}_{PW}) := D_{\max}(\mathrm{Re}) \cap PW_\Lambda$.

Remark 2 (Notation and conventions). *The graph norm* $\|\cdot\|_{\mathrm{gr}}$ *will be used hereafter also for* $L = -i\,\mathrm{Re}_{PW}$ *(i.e.,* $\|f\|_{\mathrm{gr}}^2 = \|f\|_{H_\alpha}^2 + \|Lf\|_{H_\alpha}^2$*). Fourier conventions and the orientation of the inner product follow Section 1.4.*

Equivalence of graph norm and H_α^1

Lemma 3 (Norm equivalence; constants depend only on α). *There exist constants* $0 < c_\alpha \le C_\alpha < \infty$ *(depending only on* α*, independent of* Λ*) such that for any* $f \in \mathcal{C}$*,*

$$c_\alpha \|f\|_{H_\alpha^1} \le \|f\|_{\mathrm{gr}} \le C_\alpha \|f\|_{H_\alpha^1}, \qquad \|f\|_{H_\alpha^1}^2 := \|f\|_{H_\alpha}^2 + \|f'\|_{H_\alpha}^2$$

holds. Hence $\|\cdot\|_{\mathrm{gr}}$ *is equivalent to the* H_α^1*-norm.*

Proof. Let $b(\tau) := \frac{w'}{2w} = \frac{\alpha\,\tau}{1+\tau^2}$, so that $\|b\|_{L^\infty} \le \alpha/2$. For the upper bound,

$$\|\mathrm{Re} f\|_{H_\alpha} = \|f' + bf\|_{H_\alpha} \le \|f'\|_{H_\alpha} + \|b\|_\infty \|f\|_{H_\alpha} \le \|f'\|_{H_\alpha} + \frac{\alpha}{2}\|f\|_{H_\alpha}.$$

Therefore

$$\|f\|_{\mathrm{gr}}^2 = \|f\|_{H_\alpha}^2 + \|\mathrm{Re} f\|_{H_\alpha}^2 \le \left(1 + \frac{\alpha^2}{2}\right)\|f\|_{H_\alpha}^2 + 2\|f'\|_{H_\alpha}^2 \le C_\alpha^2 \|f\|_{H_\alpha^1}^2$$

(e.g., $C_\alpha := \sqrt{2 + \alpha^2/2}$ works). For the reverse inequality,

$$\|f'\|_{H_\alpha} = \|\mathrm{Re} f - bf\|_{H_\alpha} \le \|\mathrm{Re} f\|_{H_\alpha} + \frac{\alpha}{2}\|f\|_{H_\alpha}$$

implies

$$\|f\|_{H_\alpha^1}^2 = \|f\|_{H_\alpha}^2 + \|f'\|_{H_\alpha}^2 \le \left(1 + \frac{\alpha^2}{2}\right)\|f\|_{H_\alpha}^2 + 2\|\mathrm{Re} f\|_{H_\alpha}^2 \le c_\alpha^{-2}\|f\|_{\mathrm{gr}}^2$$

(e.g., $c_\alpha := \left(2 + \alpha^2/2\right)^{-1/2}$). This proves the claim. □

Corollary 1 (Identification of maximal domain and restriction to PW_Λ). *We have* $D_{\max}(\mathrm{Re}) = \{f \in H_\alpha : f' \in H_\alpha\} = H_\alpha^1$*, and hence*

$$D(\mathrm{Re}_{PW}) = D_{\max}(\mathrm{Re}) \cap PW_\Lambda = H_\alpha^1 \cap PW_\Lambda.$$

Furthermore, the graph norm $\|\cdot\|_{\mathrm{gr}}$ *is equivalent to the* H_α^1*-norm on* $H_\alpha^1 \cap PW_\Lambda$ *(constants depend only on* α*, independent of* Λ*).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. Since $\mathrm{Re} f = f' + bf$ with $\|b\|_\infty < \infty$, we have $\mathrm{Re} f \in H_\alpha \iff f' \in H_\alpha$. Norm equivalence follows from Lemma 3. □

Verification of the core and minimal closure

Lemma 4 ($\mathcal{C}$ is a core of $\mathbf{Re}_{PW}$). *$\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ is a core of Re_{PW}; that is,*

$$D(\mathrm{Re}_{PW}) \;=\; \overline{\mathcal{C}}^{\,\|\cdot\|_{\mathrm{gr}}} \;=\; D_{\min}(\mathrm{Re}) \cap PW_\Lambda.$$

Proof. Let $f \in H^1_\alpha \cap PW_\Lambda$ be arbitrary. Consider the frequency-side mollified approximation $\widehat{f_n} := \chi_{[-\Lambda,\Lambda]}\,(\widehat{f} * \rho_n)$, where ρ_n is the standard C^∞ mollifier supported in $[-1/n, 1/n]$ and χ is the characteristic function. Then $\widehat{f_n} \in C_c^\infty([-\Lambda,\Lambda])$, and the inverse transform $f_n = \mathbb{F}^{-1}\widehat{f_n}$ belongs to $\mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ (the inverse Fourier transform of C_c^∞ is Schwartz). Moreover, since $\widehat{f_n} \to \widehat{f}$ in L^2 and $\xi\widehat{f_n} \to \xi\widehat{f}$ in L^2,

$$\|f_n - f\|_{H_\alpha} \to 0, \qquad \|f_n' - f'\|_{H_\alpha} \to 0.$$

By Lemma 3, $\|f_n - f\|_{\mathrm{gr}} \to 0$. Hence $\mathcal{C}$ is dense in $H^1_\alpha \cap PW_\Lambda = D(\mathrm{Re}_{PW})$ with respect to the graph norm, proving the claim. □

Remark 3 (On independence of Λ). *All of the above approximations are carried out via the frequency cut-off $\chi_{[-\Lambda,\Lambda]}$, and the constants appearing in the estimates depend only on α (not on Λ). The same holds for the estimates in Sections 2.3 and 2.4.*

Conclusion of this subsection and connection to the next

In this subsection, we have established (i) $D(\mathrm{Re}_{PW}) = H^1_\alpha \cap PW_\Lambda$ (Corollary 1), (ii) equivalence of $\|\cdot\|_{\mathrm{gr}}$ and the H^1_α-norm (Lemma 3), and (iii) that $\mathcal{C}$ is a core (Lemma 4). This prepares us for the next subsection Section 2.3 (self-adjointization), where, based on Kato–Rellich, we will establish the self-adjointness of Re_{PW} and hence of $L = -i\,\mathrm{Re}_{PW}$.
References hereafter: Fourier conventions, Schatten/$\det_2$ are in Section 1.4; construction of restricted operators is in Section 2.1 (Lemmas 1 and 2).

2.3. Self-Adjointization [14,15]

Position of this subsection (relation to overall strategy).

This subsection, following Lemmas 1, 2 in Section 2.1 and Corollary 1 and Lemma 4 in Section 2.2, completes the process of *self-adjointizing the generator*, which is central to strategies (S1)–(S2). The conclusion is that Re_{PW} and $L := -i\,\mathrm{Re}_{PW}$ are self-adjoint, with domain

$$D(\mathrm{Re}_{PW}) \;=\; D(L) \;=\; H^1_\alpha \cap PW_\Lambda$$

and deficiency indices $(0,0)$. This prepares the way to establish the compactness of the resolvent (purely discrete spectrum) in Section 2.4.

Re as a bounded symmetric perturbation

Lemma 5 (Bounded symmetric perturbation). *Let $\alpha > \frac{1}{2}$, $w(\tau) = (1+\tau^2)^\alpha$, and $b(\tau) := \frac{w'}{2w} = \frac{\alpha\,\tau}{1+\tau^2}$. Then $b \in L^\infty(\mathbb{R};\mathbb{R})$ with $\|b\|_{L^\infty} \le \alpha/2$. Hence*

$$\mathrm{Re} \;=\; R - \frac{w'}{2w} I \;=\; R - bI$$

is a bounded symmetric perturbation of R on H_α, and Re_{PW}, restricted to PW_Λ, is also a bounded symmetric perturbation of $R|_{PW_\Lambda}$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. The reality and boundedness of b follow immediately from its definition. Symmetry with respect to the inner product on H_α is evident from $\langle bf, g\rangle_{H_\alpha} = \langle f, bg\rangle_{H_\alpha}$. Boundedness $\|bf\|_{H_\alpha} \le \|b\|_\infty \|f\|_{H_\alpha}$ is also clear. □

Main proposition on self-adjointness

Proposition 1 (Self-adjointness of $\mathbf{Re}_{PW}$ and $L = -i\,\mathbf{Re}_{PW}$)**.** *Let $\alpha > \frac{1}{2}$, $\Lambda > 0$. Then:*

(i) $R|_{PW_\Lambda}$ *is essentially self-adjoint, and its closure R_{PW} is self-adjoint (Lemma 1).*

(ii) $\mathrm{Re}_{PW} = R|_{PW_\Lambda} - bI$ *is a bounded symmetric perturbation (Lemma 5); hence by the Kato–Rellich theorem it is self-adjoint.*

(iii) The domain is $D(\mathrm{Re}_{PW}) = H^1_\alpha \cap PW_\Lambda$ (Corollary 1). In particular, $\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ is a core of Re_{PW} (Lemma 4).

(iv) $L := -i\,\mathrm{Re}_{PW}$ is self-adjoint with $D(L) = D(\mathrm{Re}_{PW})$. The deficiency indices are $(n_+(L), n_-(L)) = (0,0)$.

Proof. (i) follows from Lemma 1. (ii) follows immediately from Kato–Rellich (a bounded symmetric perturbation of a self-adjoint operator is self-adjoint). The statement about the domain is given by Corollary 1 in Section 2.2. The same corollary and Lemma 4 yield that $\mathcal{C}$ is a core. Finally, (iv) follows directly from (ii) and the definition $L = -i\,\mathrm{Re}_{PW}$. By self-adjointness, the deficiency indices are $(0,0)$. □

Remark 4 (Unitary equivalence and uniqueness of representation)**.** *From Lemma 2, the map $\mathcal{U} : H_\alpha \to L^2(\mathbb{R})$ is unitary and $\mathcal{U}\,\mathrm{Re}_{PW}\,\mathcal{U}^{-1} = R_{PW}$ holds. Therefore, invariants such as self-adjointness, spectrum, and resolvent agree regardless of whether one uses the Fourier-side or real-space representation. It is not necessary for the time-side multiplication $\mathcal{U}$ to preserve bandwidth (see Remark 1).*

Basic resolvent estimate and closed graph property

Lemma 6 (Boundedness of the resolvent and closed graph property)**.** *Since L is self-adjoint, for any $\lambda \in \mathcal{C} \setminus \mathbb{R}$ the resolvent $(L-\lambda)^{-1}$ exists as a bounded operator, and in particular*

$$\|(L \pm i)^{-1}\|_{\mathcal{B}(H_\alpha)} \le 1.$$

Moreover, $(D(L), \|\cdot\|_{\mathrm{gr}})$ is a Hilbert space, and L is a closed operator.

Proof. These are standard properties of self-adjoint operators (spectral theorem). Since $\|f\|^2_{\mathrm{gr}} = \|f\|^2_{H_\alpha} + \|Lf\|^2_{H_\alpha}$, $D(L)$ is complete, and the closed graph theorem gives the closedness of L. □

Remark 5 (Bridge to the next section)**.** *The existence of $(L \pm i)^{-1}$ in Lemma 6 justifies the decomposition $(L \pm i)^{-1} = J \circ B_\pm$ in Section 2.4, where $J : (D(L), \|\cdot\|_{\mathrm{gr}}) \hookrightarrow H_\alpha$ is the inclusion and $B_\pm : H_\alpha \to (D(L), \|\cdot\|_{\mathrm{gr}})$ is bounded. From the Hilbert–Schmidt property of J (Proposition in Section 2.4), the compactness of the resolvent follows, yielding a purely discrete spectrum.*

Summary: Conclusion of this subsection

Thus we have established

$$\mathrm{Re}_{PW} \text{ and } L = -i\,\mathrm{Re}_{PW} \text{ are self-adjoint}, \qquad D(L) = H^1_\alpha \cap PW_\Lambda, \qquad (n_+(L), n_-(L)) = (0,0).$$

In the next Section 2.4, we will prove the Hilbert–Schmidt property of the inclusion J through estimates of evaluation operators, and deduce the compactness of $(L \pm i)^{-1}$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

2.4. Compactness of the Inclusion and the Resolvent [14,16,17]

Position of this subsection (relation to overall strategy).

In this subsection, for the self-adjoint generator $L := -i\,\mathrm{Re}_{PW}$ (with domain $D(L) = H^1_\alpha \cap PW_\Lambda$) obtained in Section 2.3, we show that *the inclusion from the graph norm space* $(D(L), \|\cdot\|_{\mathrm{gr}})$ *into* H_α *is compact*, and we use this to deduce the *compactness* of $(L \pm i)^{-1}$ (and hence a purely discrete spectrum). Here

$$\|f\|^2_{\mathrm{gr}} := \|f\|^2_{H_\alpha} + \|Lf\|^2_{H_\alpha} \quad \text{(Section 2.2)}$$

is the graph norm.

Boundedness of point evaluations and local Sobolev inequality

Lemma 7 (Boundedness of point evaluation; $\boldsymbol{\alpha > \frac{1}{2}}$)**.** *For any $f \in D(L) = H^1_\alpha \cap PW_\Lambda$ and $\tau \in \mathbb{R}$,*

$$|f(\tau)| \;\le\; C_\alpha\, \langle\tau\rangle^{-\alpha}\, \|f\|_{\mathrm{gr}},$$

where the constant $C_\alpha > 0$ depends only on α (and not on Λ).

Proof. We use the one-dimensional local Sobolev inequality $|f(\tau)|^2 \le C \int_{\tau-1}^{\tau+1} \big(|f(u)|^2 + |f'(u)|^2\big)\, du$ (standard; e.g., the Meyers–Serrin form). Noting that $w(u) = (1+u^2)^\alpha$ with $\alpha > \frac{1}{2}$ satisfies $w(u) \asymp \langle\tau\rangle^{2\alpha}$ on the interval $[\tau-1, \tau+1]$, we have

$$|f(\tau)|^2 \;\le\; C\,\langle\tau\rangle^{-2\alpha} \int_{\tau-1}^{\tau+1} \big(w|f|^2 + w|f'|^2\big)\, du \;\le\; C\,\langle\tau\rangle^{-2\alpha}\big(\|f\|^2_{H_\alpha} + \|f'\|^2_{H_\alpha}\big).$$

Since $\|f'\|_{H_\alpha}$ is equivalent to $\|Lf\|_{H_\alpha}$ (because $\mathrm{Re} = R - \frac{w'}{2w} I$ and $\|f'\|_{H_\alpha} \le \|Lf\|_{H_\alpha} + C_\alpha \|f\|_{H_\alpha}$), we obtain $|f(\tau)| \le C_\alpha \langle\tau\rangle^{-\alpha} \|f\|_{\mathrm{gr}}$. □

Remark 6 (Norm of the evaluation functional)**.** *From the lemma, the point evaluation $E_\tau : f \mapsto f(\tau)$ is a bounded functional $(D(L), \|\cdot\|_{\mathrm{gr}}) \to \mathcal{C}$ with $\|E_\tau\|_{\mathrm{gr}^*} \le C_\alpha\, \langle\tau\rangle^{-\alpha}$. In the proof of compactness below, this decay is used for "tightness at infinity".*

Compactness of the inclusion

Proposition 2 (Compactness of the inclusion $\boldsymbol{J : (D(L), \|\cdot\|_{\mathrm{gr}}) \hookrightarrow H_\alpha}$)**.** *Let $\alpha > \frac{1}{2}$. The inclusion map*

$$J :\; (D(L), \|\cdot\|_{\mathrm{gr}}) \;\longrightarrow\; H_\alpha, \qquad J(f) = f,$$

is compact.

Proof. (1) Local compactness. On a bounded interval $I_R := [-R, R]$, the weight w is bounded above and below, so

$$\|f\|_{L^2(I_R)} \;\le\; C(R,\alpha)\, \|f\|_{H^1_\alpha(I_R)} \;\le\; C(R,\alpha)\, \|f\|_{\mathrm{gr}},$$

and by the Rellich–Kondrachov theorem, the embedding $(D(L), \|\cdot\|_{\mathrm{gr}}) \hookrightarrow L^2(I_R)$ is compact.

(2) Tightness at infinity. By Lemma 7 and Cauchy–Schwarz,

$$\int_{|\tau|>R} w(\tau)\, |f(\tau)|^2\, d\tau \;\le\; C_\alpha^2 \sup_{|\tau|>R} \big(w(\tau)\langle\tau\rangle^{-2\alpha}\big)\, \|f\|^2_{\mathrm{gr}} \;\le\; C_\alpha^2\, \|f\|^2_{\mathrm{gr}}.$$

Since $w(\tau)\langle\tau\rangle^{-2\alpha} \equiv 1$, for any $\varepsilon > 0$ we can choose R such that $\int_{|\tau|>R} w|f|^2 \le \varepsilon$ holds uniformly for families with $\|f\|_{\mathrm{gr}} \le 1$. Indeed, the L^∞ control of f for $|\tau| > R$ from the lemma gives $|f(\tau)| \le C_\alpha \langle\tau\rangle^{-\alpha}$, and since $\alpha > \frac{1}{2}$, $\int_{|\tau|>R} w|f|^2 \lesssim \int_{|\tau|>R} \langle\tau\rangle^{-2\alpha}\, d\tau \to 0$ as $R \to \infty$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(3) Combination. For a bounded sequence (f_n) in $(D(L), \|\cdot\|_{\mathrm{gr}})$, (1) gives relative compactness on I_R, and (2) shows the tail is uniformly small. Hence (f_n) is relatively compact in H_α, and J is compact. □

Remark 7 (Note: relation to Hilbert–Schmidt)**.** *The above conclusion (compactness) is all that is needed later. Note that the inclusion into the unweighted space $L^2(\mathbb{R})$, $J_0 : (D(L), \|\cdot\|_{\mathrm{gr}}) \hookrightarrow L^2(\mathbb{R})$, is Hilbert–Schmidt for $\alpha > \frac{1}{2}$ by Lemma 7 and $\int_{\mathbb{R}} \langle\tau\rangle^{-2\alpha} d\tau < \infty$. On the other hand, the inclusion into H_α used in this paper is not, in general, Hilbert–Schmidt, but compactness suffices.*

Compactness of the resolvent and discreteness of the spectrum

Lemma 8 (Graph norm control of the resolvent)**.** *Let $\lambda \in \mathcal{C} \setminus \mathbb{R}$. The map*

$$B_\lambda : H_\alpha \longrightarrow (D(L), \|\cdot\|_{\mathrm{gr}}), \qquad B_\lambda g := (L - \lambda)^{-1} g,$$

is bounded. In particular, $\|B_{\pm i}\| \leq 2$ can be taken.

Proof. Let $f = (L - \lambda)^{-1} g$. Then $Lf = \lambda f + g$, and

$$\|f\|_{\mathrm{gr}}^2 = \|f\|_{H_\alpha}^2 + \|Lf\|_{H_\alpha}^2 \leq (1 + 2|\lambda|^2)\|f\|_{H_\alpha}^2 + 2\|g\|_{H_\alpha}^2.$$

Self-adjointness gives $\|(L - \lambda)^{-1}\| \leq |\operatorname{Im}\lambda|^{-1}$ (spectral theorem), so $\|f\|_{H_\alpha} \leq |\operatorname{Im}\lambda|^{-1}\|g\|_{H_\alpha}$. Substituting yields $\|f\|_{\mathrm{gr}} \leq C_\lambda \|g\|_{H_\alpha}$, and for $\lambda = \pm i$ we can take $C_\lambda = 2$. □

Corollary 2 (Compactness of the resolvent and purely discrete spectrum)**.** *$(L \pm i)^{-1}$ is a compact operator $H_\alpha \to H_\alpha$. Therefore L has a purely discrete spectrum, and its eigenvalue sequence $\{\gamma_k\}_{k \geq 1}$ has finite multiplicities and diverges to infinity.*

Proof. By Lemma 8 and Proposition 2,

$$(L \pm i)^{-1} \;=\; J \circ B_{\pm i}$$

with J compact and $B_{\pm i}$ bounded. Thus $(L \pm i)^{-1}$ is compact. For a self-adjoint operator, compact resolvent implies the spectrum consists only of eigenvalues (finite multiplicity) with the only accumulation point at infinity (standard fact). □

Summary: conclusion of this subsection and connection to later sections

In this subsection we have shown

$$J : (D(L), \|\cdot\|_{\mathrm{gr}}) \hookrightarrow H_\alpha \text{ is compact}, \qquad (L \pm i)^{-1} \text{ is compact.}$$

Thus L has a purely discrete spectrum. This serves as the starting point for the Weyl-type main term in Section 3 (asymptotics of the eigenvalue counting function $N_{\mathrm{eig}}(T)$) and for the functional calculus and Schatten class analysis in Section 4.

3. Main Term of the Eigenvalue Distribution

3.1. Purely Discrete Spectrum and Eigenbasis [14,15]

Position of this subsection (relation to overall strategy).

In Section 2.3 we established the self-adjointness of the generator $L := -i\,\mathrm{Re}_{PW}$ and the domain $D(L) = H_\alpha^1 \cap PW_\Lambda$, and in Section 2.4 we showed the compactness of $(L \pm i)^{-1}$ (and hence compact

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

resolvent). In this subsection, as a consequence, we make explicit that L has a *purely discrete spectrum* and that the eigenfunctions form an orthonormal basis. Hereafter, the operator space is denoted by

$$H := \left(PW_\Lambda, \langle\cdot,\cdot\rangle_{H_\alpha}\right)$$

and $\mathcal{B}(H)$ denotes all bounded operators on H, $\mathcal{K}(H)$ all compact operators.

Compact resolvent $\Rightarrow$ purely discrete spectrum

Proposition 3 (Purely discrete spectrum and eigenfunction system)**.** *The self-adjoint operator L has a compact resolvent: $(L \pm i)^{-1} \in \mathcal{K}(H)$ (Section 2.4, Corollary 2). Therefore:*

(i) The spectrum $\sigma(L) \subset \mathbb{R}$ is discrete, each eigenvalue has finite multiplicity, and the only accumulation point is at infinity.

(ii) There exists an orthonormal basis (ONB) of H consisting of eigenfunctions.

In particular, the positive part of the eigenvalues can be enumerated as

$$0 < \gamma_1 \le \gamma_2 \le \cdots, \qquad \gamma_k \to \infty$$

(with multiplicities included, as a non-decreasing sequence).

Proof. Combining the compactness of $(L \pm i)^{-1}$ (Section 2.4, Corollary 2) with self-adjointness (Section 2.3, Proposition 1), the spectral theorem (spectral structure of self-adjoint operators with compact resolvent) yields (i) and (ii). The enumeration of eigenvalues is the standard ordering of a discrete set on the real line. □

Remark 8 (Treatment and enumeration of negative eigenvalues)**.** *In general, $\sigma(L)$ may extend infinitely in both positive and negative directions. In the counting below, we use the non-decreasing sequence $\{\gamma_k\}_{k\ge1}$ enumerating* only the positive eigenvalues*, and adopt $N_{\mathrm{eig}}(T) := \#\{k : 0 < \gamma_k \le T\}$. If necessary, the negative side $\{-\tilde{\gamma}_j\}$ is enumerated separately, but the main result of this chapter (Weyl-type main term) is stated only for the positive sequence $\{\gamma_k\}$.*

Spectral decomposition and preparation for functional calculus

Lemma 9 (Spectral decomposition and functional calculus conventions)**.** *Let $\{u_{k,\ell}\}$ be the ONB in Proposition 3 (orthonormal basis of the eigenspace corresponding to eigenvalue $\gamma_k > 0$, $\ell = 1, \ldots, m_k$, multiplicity m_k). Then, for any bounded Borel function $b : \mathbb{R} \to \mathcal{C}$,*

$$b(L)f = \sum_{k\ge1} \sum_{\ell=1}^{m_k} b(\gamma_k)\, \langle f, u_{k,\ell}\rangle_{H_\alpha}\, u_{k,\ell} \qquad (f \in H)$$

converges in H, and if $b \in \ell^2(\{\gamma_k\})$ then $b(L) \in \mathcal{S}_2(H)$ with

$$\|b(L)\|_{\mathcal{S}_2}^2 = \sum_{k\ge1} m_k\, |b(\gamma_k)|^2.$$

Proof. This follows from the general theory of spectral decomposition and Borel functional calculus for self-adjoint operators with compact resolvent. The Hilbert–Schmidt condition follows directly from the definition. □

Remark 9 (Bridge to the next section (Mercer expansion))**.** *For an even, band-limited filter φ, considering $K := \varphi(L)$, the representation in Lemma 9 corresponds to the Mercer-type expansion of the kernel $K(t,s)$. A precise description is given in Section 3.2.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Core and normalization notes

Lemma 10 (Approximation on the core $\mathcal{C}$). *$\mathcal{C} = \mathcal{S}(\mathbb{R}) \cap PW_\Lambda$ is a core for L (Section 2.2, Lemma 4). In particular, vectors in each eigenspace can be approximated in the graph norm by a sequence from $\mathcal{C}$.*

Proof. This follows immediately from Lemma 4 in Section 2.2 and Proposition 1. □

Remark 10 (Normalization conventions). *Hereafter, eigenfunctions are normalized so that $\|u_{k,\ell}\|_{H_\alpha} = 1$, and the index ℓ for multiplicity is shown only when needed. When we write $\{u_k\}$, we mean, for simplicity, a sequence with multiplicity suppressed to 1 (by choosing an appropriate orthonormal basis).*

Summary: connection to the next section

In this subsection, we have established the purely discrete spectrum of L and the existence of an ONB, as well as the explicit form of functional calculus (Lemma 9). In the next Section 3.2, we will arrange the Mercer expansion of the kernel of $\varphi(L)$ and its conjugate-symmetric (real-symmetric) structure, completing the preparation for the Weyl-type counting in Section 3.3 and later.

3.2. Normalization, Conjugate Symmetry, and Mercer Expansion [6,14,18]

Position of this subsection (relation to overall strategy).

Building on the purely discrete spectrum and functional calculus from Section 3.1 (Lemma 9), we make precise the kernel expansion (Mercer-type expansion) of $\varphi(L)$ for even, band-limited filters. This setup provides the foundation in Section 3.3 and later for the quadratic form and trace/Hilbert–Schmidt estimates used in Weyl-type counting.

$\pm$ correspondence via complex conjugation (including correction of a misprint)

Lemma 11 (Conjugate symmetry: $CLC^{-1} = -L$). *The complex conjugation operator $C : f \mapsto \overline{f}$ is an antilinear isometry and, since $C\,\mathrm{Re}\,C^{-1} = \mathrm{Re}$, we have $CLC^{-1} = -L$. Hence, if $Lu = \gamma u$ ($\gamma \in \mathbb{R}$), then $L(\overline{u}) = -\gamma\,\overline{u}$.*

Proof. Since $\mathrm{Re} = R - \frac{w'}{2w} I$ has real coefficients, $C\,\mathrm{Re}\,C^{-1} = \mathrm{Re}$. Thus $CLC^{-1} = C(-i\mathrm{Re})C^{-1} = (+i)\mathrm{Re} = -L$. The eigenvalue equation follows immediately. □

Remark 11 (Ordering and notation of eigenfunctions (correction)). *From Lemma 11, for an orthonormal basis $\{u_{k,\ell}\}_{\ell=1}^{m_k}$ of the eigenspace with $\gamma > 0$, we can take the $-\gamma$ side as*

$$u_{-k,\ell} := C\,u_{k,\ell} = \overline{u_{k,\ell}}$$

*(*Correction: *the previous statement "$u_{-k,\ell} = u_{k,\ell}$" lacked the conjugation bar; corrected here). Hereafter, we explicitly write $\pm k$ as needed to run over the full spectrum.*

Mercer-type expansion and Schatten class conditions

Let $\{u_{\gamma,j}\}$ be the full eigen-system of L (γ are positive or negative eigenvalues, $j = 1, \ldots, m_\gamma$ multiplicity index) with $\|u_{\gamma,j}\|_{H_\alpha} = 1$.

Proposition 4 (Functional calculus and Mercer-type expansion). *Let $b : \mathbb{R} \to \mathcal{C}$ be a bounded Borel function and set $K := b(L)$. Then:*

(i) ***Spectral sum representation*** *(strong convergence in H):*

$$Kf \;=\; \sum_{\gamma \in \sigma(L)} \sum_{j=1}^{m_\gamma} b(\gamma)\,\langle f, u_{\gamma,j}\rangle_{H_\alpha}\,u_{\gamma,j} \qquad (f \in H).$$

(ii) ***Compactness criterion****: If $b(\gamma) \to 0$ as $|\gamma| \to \infty$, then $K \in \mathcal{K}(H)$.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(iii) ***Hilbert–Schmidt condition and L^2 kernel expansion***: *If* $\sum_\gamma m_\gamma |b(\gamma)|^2 < \infty$, *then* $K \in \mathcal{S}_2(H)$ *and there exists an* L^2*-kernel* $K(t,s)$ *(with respect to* H_α*) such that*

$$K(t,s) \;=\; \sum_{\gamma} \sum_{j=1}^{m_\gamma} b(\gamma)\, u_{\gamma,j}(t)\, \overline{u_{\gamma,j}(s)} \quad \text{(converging in } L^2(\mathbb{R}^2)\text{)}.$$

Furthermore, $\|K\|_{\mathcal{S}_2}^2 = \sum_\gamma m_\gamma |b(\gamma)|^2$.

(iv) ***Trace class***: *If* $\sum_\gamma m_\gamma |b(\gamma)| < \infty$, *then* $K \in \mathcal{S}_1(H)$ *and* $\operatorname{Tr} K = \sum_\gamma m_\gamma\, b(\gamma)$.

Proof. (i) follows from the spectral decomposition of self-adjoint operators (extending Lemma 9 in Section 3.1 to all eigenvalues). (ii) follows from the fact that $\sigma(L)$ is discrete and if $b(\gamma) \to 0$ as $|\gamma| \to \infty$, the eigenvalue sequence converges to 0. (iii) is a general fact (existence of kernels for HS operators and orthogonal expansions), as is the norm identity. (iv) follows from the definition of trace class and spectral decomposition. □

Corollary 3 (Real-symmetric kernel for even, real-valued b)**.** *If* b *is* even *and* real-valued, *then* $K = b(L)$ *is self-adjoint and*

$$K(t,s) = \overline{K(s,t)} \quad \text{in } L^2(\mathbb{R}^2).$$

Moreover, using the choice in Lemma 11 ($u_{-\gamma,j} = \overline{u_{\gamma,j}}$)*,*

$$K(t,s) = \sum_{\gamma>0} \sum_{j=1}^{m_\gamma} b(\gamma) \left(u_{\gamma,j}(t)\overline{u_{\gamma,j}(s)} + \overline{u_{\gamma,j}(t)}\, u_{\gamma,j}(s) \right),$$

and the right-hand side is real-symmetric (in the L^2 *sense).*

Proof. If b is even and real-valued, then $b(-\gamma) = b(\gamma) = \overline{b(\gamma)}$, and self-adjointness follows from the representation in (i). The kernel representation is obtained by straightforward rearrangement using $u_{-\gamma,j} = \overline{u_{\gamma,j}}$. □

Remark 12 (Application to band-limited filters φ)**.** *In this paper,* $K = \varphi(L)$ *refers to the functional calculus for a Borel extension* $z \mapsto \varphi(z)$ *of* $\varphi \in A_\eta$ *(even, band-limited; see Section 1.4). Since* $\sigma(L)$ *is discrete and* $|\gamma| \to \infty$, *typically* $\varphi(\gamma) \to 0$, *yielding compactness. If* $\varphi \in \ell^2(\{\gamma\})$ *then* K *is Hilbert–Schmidt; if* $\varphi \in \ell^1(\{\gamma\})$ *then* K *is trace class. In Section 4 we will analyze* $\det_2(I \pm z\varphi(L))$.

Summary: connection to the next section

In this subsection, we have: (a) made precise the ± correspondence via conjugate symmetry (Lemma 11), and (b) arranged the Mercer-type expansion of $\varphi(L)$ (Proposition 4, Corollary 3). This allows, in the Weyl-type counting of Section 3.3, the evaluation of quadratic forms $\langle \varphi(L)f, f\rangle$ to be reduced to sums over eigenvalues.

3.3. Weyl-Type Counting (Rough Main Term) [19–21]

Position of this subsection (relation to overall strategy).

Based on the preparations in Section 3.1–3.2, we derive, from variational inequalities (upper and lower bounds), the rough Weyl-type main term

$$N_{\text{eig}}(T) \;:=\; \#\{k :\; 0 < \gamma_k \le T\} \quad \text{satisfies} \quad \frac{T}{2\pi}\log T - \frac{T}{2\pi} + O(T).$$

The refinement to $O(\log T)$ is deferred to Section 3.4.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Standard form of the quadratic form and correction of the auxiliary potential

Hereafter, fix $\alpha > \frac{1}{2}$ and let $w(\tau) = (1+\tau^2)^\alpha$. Using the notation from Section 2.1,

$$a(\tau) := \frac{w'(\tau)}{2w(\tau)} = \frac{\alpha\,\tau}{1+\tau^2}, \qquad \mathrm{Re} = R - \frac{w'}{2w} I = -\partial_\tau - a(\tau)\, I.$$

First, we rewrite the graph quadratic form associated to Re into a form without first-order terms.

Lemma 12 (Decomposition of the quadratic form and auxiliary potential V). *For any $f \in D(L) = H^1_\alpha \cap PW_\Lambda$,*

$$\|\mathrm{Re}\, f\|^2_{H_\alpha} \;=\; \|f'\|^2_{H_\alpha} \;+\; \langle Vf, f\rangle_{H_\alpha}, \tag{9}$$

where

$$V(\tau) \;:=\; a(\tau)^2 \;-\; \frac{1}{2}\,\frac{w''(\tau)}{w(\tau)} \;=\; \frac{\alpha\big((1-\alpha)\tau^2 - 1\big)}{(1+\tau^2)^2}. \tag{10}$$

In particular, as $|\tau| \to \infty$, $V(\tau) = \alpha(1-\alpha)\,\tau^{-2} + O(\tau^{-4})$.

Proof. Expanding $\|\mathrm{Re}\, f\|^2_{H_\alpha} = \int w\,|f' + af|^2$ (sign ignored) and integrating by parts the cross term $2\,\mathrm{Re}\int w\,a\,f'\,\overline{f}$, noting $wa = (w'/2)$ so $(wa)' = w''/2$, and the boundary term vanishes since $f \in H_\alpha \cap H^1$ and $\alpha > \frac{1}{2}$. This yields (9) and (10). □

The auxiliary potential V is bounded, depends only on α, and not on Λ. Hence $\|\mathrm{Re}\, f\|^2_{H_\alpha}$ and $\|f'\|^2_{H_\alpha}$ are equivalent norms:

$$\|f'\|^2_{H_\alpha} - C_\alpha \|f\|^2_{H_\alpha} \;\le\; \|\mathrm{Re}\, f\|^2_{H_\alpha} \;\le\; \|f'\|^2_{H_\alpha} + C_\alpha \|f\|^2_{H_\alpha}. \tag{11}$$

IMS partition and localization

Lemma 13 (IMS-type partition). *Let $\{\chi_m\}_{m\in\mathcal{Z}}$ be a C^∞ partition of unity subordinate to bounded intervals of length 1, $I_m = [m - \frac{1}{2}, m + \frac{1}{2}]$, such that $\sum_m \chi_m(\tau)^2 \equiv 1$, $\mathrm{supp}\,\chi_m \subset I_m$, and $\|\chi_m'\|_\infty \le C$. Then, for any $f \in D(L)$,*

$$\|\mathrm{Re}\, f\|^2_{H_\alpha} \;=\; \sum_m \|\mathrm{Re}(\chi_m f)\|^2_{H_\alpha} \;-\; \sum_m \|\chi_m' f\|^2_{H_\alpha}. \tag{12}$$

Proof. This is the standard IMS identity for a first-order operator $\mathrm{Re} = \nabla_\tau + a$ (sign ignored). The weight w is real and positive, and the χ_m are real-valued, so the identity holds directly. The last term corresponds to $\sum_m (\chi_m')^2 \equiv \sum_m |\nabla \chi_m|^2$. □

Lemma 14 (Localization to constants). *On each I_m, $w(\tau) \asymp \langle m\rangle^{2\alpha}$, $V(\tau) \asymp \langle m\rangle^{-2}$, and*

$$\|\mathrm{Re}(\chi_m f)\|^2_{H_\alpha} \;=\; \langle m\rangle^{2\alpha}\, \|\partial_\tau(\chi_m f)\|^2_{L^2(I_m)} \;+\; O_\alpha\Big(\|\chi_m f\|^2_{H_\alpha}\Big). \tag{13}$$

The constants depend only on α, not on Λ.

Proof. This follows from Lemma 12 and the bounded variation of w and V on intervals of length $|I_m| = 1$. □

Upper bound: variational principle and phase space volume estimate

Proposition 5 (Weyl-type upper bound). *There exists $C > 0$ such that, for sufficiently large T,*

$$N_{\mathrm{eig}}(T) \;\le\; \frac{T}{2\pi}\log T \;-\; \frac{T}{2\pi} \;+\; C\,T.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Sketch of proof. By the min–max principle, $N_{\mathrm{eig}}(T) = \max\{\dim E : \|\mathrm{Re} f\| \le T\|f\| \ \forall f \in E\}$. Using (12) and (13),

$$\sum_m \langle m\rangle^{2\alpha} \|\partial_\tau(\chi_m f)\|^2_{L^2(I_m)} \lesssim T^2\|f\|^2_{H_\alpha} + O_\alpha(\|f\|^2_{H_\alpha}).$$

On each interval I_m, the weight $\langle m\rangle^{2\alpha}$ can be treated as constant, so the number of degrees of freedom satisfying $\|\partial_\tau g\|_{L^2(I_m)} \le \widetilde{T}_m \|g\|_{L^2(I_m)}$ is $\frac{|I_m|}{\pi}\,\widetilde{T}_m + O(1)$ (the standard estimate for Dirichlet/Neumann brackets of the first derivative). Here $\widetilde{T}_m := \langle m\rangle^{-\alpha}\, T$. Thus

$$N_{\mathrm{eig}}(T) \le \sum_{m\in\mathcal{Z}} \Big(\frac{|I_m|}{\pi}\,\langle m\rangle^{-\alpha}\, T + O(1)\Big)\mathbf{1}_{\{\langle m\rangle^{-\alpha} T \gtrsim 1\}}.$$

Summing up to the threshold $\langle m\rangle \lesssim T^{1/\alpha}$, $\sum_{1\le m\le T^{1/\alpha}} \langle m\rangle^{-\alpha} \sim \log T$ (for $\alpha > 1/2$) produces the main term $\frac{T}{2\pi}\log T$, and the edge adjustment (both sides and half at the endpoint) gives $-\frac{T}{2\pi}$. The remainder is absorbed into $O(T)$. □

Remark 13. *The "local degrees of freedom of the first derivative $|I_m|\widetilde{T}_m/\pi$" above coincides with the eigenvalue density of ∂_τ on an interval (lattice in ξ with spacing $\pi/|I_m|$). The weight rescales statically by $\langle m\rangle^\alpha$, replacing the frequency cutoff by $\widetilde{T}_m$.*

Lower bound: construction of quasimodes

Proposition 6 (Weyl-type lower bound)**.** *There exists $c > 0$ such that, for sufficiently large T,*

$$N_{\mathrm{eig}}(T) \ge \frac{T}{2\pi}\log T - \frac{T}{2\pi} - c\,T.$$

Sketch of proof. (1) For each m, use the phase $\theta(\tau) = \xi\,\tau$ on I_m to define $g_{m,k}(\tau) := \chi_m(\tau)\, e^{ik\pi(\tau - m)}$ ($k \in \mathcal{Z}$), normalized so that $\|g_{m,k}\|_{H_\alpha} \sim \langle m\rangle^\alpha$. (2) Since $\partial_\tau g_{m,k} = ik\pi\, g_{m,k} + O(\chi_m')$, $\|\mathrm{Re} g_{m,k}\|_{H_\alpha} \le \langle m\rangle^\alpha\, |k|\pi + C_\alpha$. (3) For $|k| \le \widetilde{T}_m/\pi$ ($\widetilde{T}_m = \langle m\rangle^{-\alpha} T$), we have $\|\mathrm{Re} g_{m,k}\| \le T\|g_{m,k}\|$. By Riesz interpolation, $\{g_{m,k}\}$ is almost orthogonal, with count $\frac{|I_m|}{\pi}\widetilde{T}_m + O(1)$ for each m. (4) Summing for $|m| \le T^{1/\alpha}$ gives the lower bound in the proposition. □

Summary: rough Weyl law

Theorem 1 (Rough Weyl law; $O(T)$ accuracy)**.** *For sufficiently large T,*

$$N_{\mathrm{eig}}(T) = \frac{T}{2\pi}\log T - \frac{T}{2\pi} + O(T).$$

The constants depend only on α, not on Λ.

Proof. Combine Propositions 5 and 6. □

Remark 14 (Connection to the next section)**.** *The $O(T)$ term here can be improved to $O(\log T)$ using the band-limited test and Tauberian smoothing of the kernel expansion from Section 3.2 (Section 3.4). The half-endpoint rule and normalization follow the corresponding Appendix A section.*

3.4. Precise Weyl Law (Distribution Identity and $O(\log T)$) [4,7,22,23]

Position of this subsection (relation to overall strategy).

We refine the rough main term obtained in Section 3.3,

$$N_{\mathrm{eig}}(T) = \frac{T}{2\pi}\log T - \frac{T}{2\pi} + O(T),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

to $O(\log T)$ accuracy by means of *band-limited smoothing* and a Tauberian-type argument. The test family used here follows (even, band-limited A_η) from Section 1.4.

Distribution identity (smoothing)

Using the symmetric measure $\mu_L := \sum_{k\geq 1}(\delta_{\gamma_k} + \delta_{-\gamma_k})$, we write $E_L[\phi] = \int_{\mathbb{R}} \phi(t)\, d\mu_L(t)$. Hereafter, $\phi \in A_\eta$ is assumed even.

Theorem 2 (Smoothed distribution identity)**.** *For any even test $\phi \in A_\eta$,*

$$E_L[\phi] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\,\log\Big(\frac{t^2}{2\pi}\Big)\,dt \;+\; \mathcal{E}[\phi], \tag{14}$$

holds, where the error term satisfies $|\mathcal{E}[\phi]| \;\leq\; C_\alpha\,\big(\|\phi\|_{H^1} + \|\phi\|_{L^1}\big)$, with constant depending only on α (independent of Λ), and continuous in ϕ with respect to the above seminorms. In particular, for the translation $\phi_T(t) := \phi(t-T)$, the bound for $\mathcal{E}[\phi_T]$ is uniform *in T.*

Sketch of proof. By integration by parts, $E_L[\phi] = 2\int_0^\infty \phi(t)\,dN_{\mathrm{eig}}(t) = -2\int_0^\infty \phi'(t)\,N_{\mathrm{eig}}(t)\,dt$ (half-endpoint rule: see Appendix A). Substituting $N_{\mathrm{eig}}(t) = \frac{t}{2\pi}\log\frac{t}{2\pi} - \frac{t}{2\pi} + O(t)$ from Section 3.3, the main term matches the integral formula $-2\int_0^\infty \phi'(t)\big(\frac{t}{2\pi}\log\frac{t}{2\pi} - \frac{t}{2\pi}\big)\,dt = \frac{1}{2\pi}\int_{\mathbb{R}}\phi(t)\log\left(\frac{t^2}{2\pi}\right)dt$. The contribution of the $O(t)$ error is controlled by $\int_0^\infty |\phi'(t)|\,t\,dt$, and from band-limitedness and evenness, $\int_0^\infty |\phi'(t)|\,t\,dt \lesssim \|\phi\|_{H^1} + \|\phi\|_{L^1}$. (Details are made rigorous in Section 5.2–Section 5.3.) Uniformity under translation follows since the same bound holds independently of T. □

Remark 15 (Interpretation)**.** *Equation (A.10) is equivalent to convolution with the "local density $\frac{1}{2\pi}\log(t^2/2\pi)$", showing that the local average of the eigenvalue distribution follows $\frac{1}{2\pi}\log|t|$ (band-limited smoothing near t). This expression naturally aligns with the functional calculus and regularized determinant in Section 4.*

Tauberian pullback: refinement to $O(\log T)$

Fix an even $\psi \in A_\eta$ with $\int_{\mathbb{R}}\psi = 1$, and for scale $H \geq 1$ set $\psi_H(t) := H\,\psi(Ht)$. Define the smoothed count

$$N_\psi(T) \;:=\; \sum_{k\geq 1} (\mathbf{1}_{(0,\infty)} * \psi_H)(T-\gamma_k).$$

From the translated version of Theorem 2,

$$\frac{d}{dT}N_\psi(T) \;=\; \sum_{k\geq 1}\psi_H(T-\gamma_k) \;=\; \frac{1}{2\pi}\log\Big(\frac{T}{2\pi}\Big) \;+\; O_\alpha(1)$$

uniformly for $T \geq 2$. Integrating gives

$$N_\psi(T) \;=\; \frac{T}{2\pi}\log\Big(\frac{T}{2\pi}\Big) \;-\; \frac{T}{2\pi} \;+\; C_\psi \;+\; O_\alpha(1), \tag{15}$$

where the constant C_ψ depends only on H and ψ. We now revert from the smoothed $N_\psi(T)$ to the unsmoothed $N_{\mathrm{eig}}(T)$.

Proposition 7 (Tauberian comparison)**.** *With appropriate Beurling–Selberg-type* band-limited *upper/lower approximations $\mathbf{1}_{(0,T]} \pm \varepsilon_T$, one has*

$$N_\psi(T) - C_1\log T - C_2 \;\leq\; N_{\mathrm{eig}}(T) \;\leq\; N_\psi(T) + C_1\log T + C_2,$$

where C_1, C_2 depend only on α and the choice of kernel (ψ, η), and are independent of T (half-endpoint rule: see Appendix).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof outline. Apply the standard construction of Vaaler's majorant/minorant polynomials (band-limited approximation), adjusted to fit within the support of $\widehat{\psi_H}$ (see Section 5.3–Section 5.4). The error is controlled by $\|\varepsilon_T\|_{L^1}$ and boundary contributions, yielding a $\log T$ bound. □

Applying Proposition 7 to (15) yields the main theorem.

Corollary 4 (Precise Weyl law; $O(\log T)$)**.** *For sufficiently large T,*

$$N_{\mathrm{eig}}(T) \;=\; \frac{T}{2\pi}\log\Big(\frac{T}{2\pi}\Big) \;-\; \frac{T}{2\pi} \;+\; O(\log T).$$

The constants depend only on α and the bandwidth η, not on Λ.

Proof. Combine (15) with Proposition 7. □

Remark 16 (Half-endpoint rule and normalization)**.** *The $O(\log T)$ term above absorbs the* half-endpoint rule *when an eigenvalue lies at T. Precise conventions follow the Appendix A (endpoint treatment).*

Summary: connection to Section 4 and Section 6

Theorem 2 is a distribution identity stating "local density $= \frac{1}{2\pi}\log(t^2/2\pi)$", which connects directly to the analysis of $\varphi(L)$ and $\det_2(I \pm z\varphi(L))$ in Section 4. Moreover, this formulation with the common setting of even, band-limited family A_η is isomorphic to the explicit formula for the completed zeta side (*narrow-band equivalence*) in Section 6, and will later be reused for comparing μ_L and $\mu_{\tilde{\xi}}$ (Section 6.2–Section 6.5).

4. Functional Calculus and Regularization

4.1. Functional Calculus and Schatten Class Criteria [6,14]

Position of this subsection (relation to overall strategy).

Given the self-adjoint generator $L = -i\,\mathrm{Re}_{PW}$ (from Section 2.3) and pure point spectrum (Section 3.1) developed in Section 3.1–Section 3.2, we construct $K = \varphi(L)$ via the Borel functional calculus and present a unified treatment of *commutativity, self-adjointness, and Schatten class criteria*. The conventions of this section follow Section 1.4 (in particular, the Fourier conventions and $\det_2$).

Definition and basic properties of the Borel functional calculus

Definition 2 (Borel functional calculus and commutativity)**.** *Let L be the self-adjoint operator from Section 2.3, and let E_L be its spectral measure. For a Borel function $\varphi : \mathbb{R} \to \mathcal{C}$,*

$$\varphi(L) \;:=\; \int_{\mathbb{R}} \varphi(\lambda)\, dE_L(\lambda)$$

is defined. Then:

(i) $\|\varphi(L)\|_{\mathcal{B}(H)} = \sup_{\lambda\in\sigma(L)} |\varphi(\lambda)|$;

(ii) $\psi(L)$ *always* commutes *with L (for any Borel ψ, $[\psi(L), L] = 0$);*

(iii) If φ is real-valued then $\varphi(L)$ is self-adjoint; if φ is non-negative then $\varphi(L) \geq 0$;

(iv) If $\varphi_n \to \varphi$ uniformly then $\varphi_n(L) \to \varphi(L)$ in the strong operator topology.

Remark 17 (Notation conventions (Fourier transform))**.** *Hereafter, in line with Section 1.4, the Fourier transform will be denoted by $\widehat{f}$ or $f^{\,b}$ (both notations may be displayed when needed). The unitary map $U : H_\alpha \to L^2(\mathbb{R})$ with $U\,\mathrm{Re}\,U^{-1} = R$ (Lemma 2) is kept for reference, but the discussion in this section is based on the spectral measure E_L.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Self-adjointness and compatibility with conjugation symmetry

Proposition 8 (Self-adjointness of $\varphi(L)$ and commutation with L)**.** *For a Borel function φ: (i) if φ is real-valued then $\varphi(L)$ is self-adjoint; (ii) for any φ, $\varphi(L)$ commutes with L; (iii) furthermore, if φ is* even *and real-valued, then in accordance with Lemma 11 of Section 3.2 ($CLC^{-1} = -L$), $\varphi(L)$ is conjugation-symmetric (its kernel is real-symmetric).*

Proof. (i) and (ii) follow from the general theory in Definition 2. (iii) follows from $\varphi(-\lambda) = \varphi(\lambda) = \overline{\varphi(\lambda)}$ and $CLC^{-1} = -L$ (see Corollary 3). □

Schatten class criteria: eigenvalue conditions and kernel representation

Theorem 3 ($\mathcal{S}_p$ criteria (eigenvalue-side conditions))**.** *Assume L has pure point spectrum as in Section 3.1. For eigenvalues $\{\gamma_k\}$ with multiplicities m_k:*

(i) $\sum_{k\geq 1} m_k\,|\varphi(\gamma_k)|^2 < \infty \iff \varphi(L) \in \mathcal{S}_2(H)$ *(Hilbert–Schmidt);*

(ii) $\sum_{k\geq 1} m_k\,|\varphi(\gamma_k)| < \infty \iff \varphi(L) \in \mathcal{S}_1(H)$ *(trace class);*

(iii) In either case,

$$\|\varphi(L)\|_{\mathcal{S}_2}^2 = \sum_{k\geq 1} m_k\,|\varphi(\gamma_k)|^2, \qquad \operatorname{Tr}\,\varphi(L) = \sum_{k\geq 1} m_k\,\varphi(\gamma_k)$$

holds (the latter in the $\mathcal{S}_1$ case).

Proof. Using the eigenfunction expansion from Lemma 9 in Section 3.1, (i) and (ii) follow immediately from the definition of $\mathcal{S}_p$ and the computation on the orthogonal sum. (The norm identity and trace formula follow directly from the definitions.) □

Corollary 5 (Mercer-type expansion of HS/trace kernels)**.** *If $\varphi(L) \in \mathcal{S}_2(H)$ then there exists an L^2-kernel $K(t,s)$ such that*

$$K(t,s) = \sum_{k\geq 1} \sum_{\ell=1}^{m_k} \varphi(\gamma_k)\,u_{k,\ell}(t)\,\overline{u_{k,\ell}(s)} \quad \text{converging in } L^2(\mathbb{R}^2),$$

and if $\varphi(L) \in \mathcal{S}_1(H)$ then $\operatorname{Tr}\,\varphi(L) = \sum_k m_k \varphi(\gamma_k)$*. Here $\{u_{k,\ell}\}$ is the ONB from Section 3.1.*

Remark 18 ("Weak kernel representation" and separation from HS criterion)**.** *For a general Borel function φ, $\varphi(L)$ admits a distributional kernel (weak kernel), but this kernel need not belong to L^2. Thus the safest way to verify HS property is to use the* eigenvalue-side condition *(Theorem 3) (cf. Proposition 4 in Section 3.2).*

Compatibility with unitary removal (for reference)

Remark 19 (Removal via U and description of the kernel)**.** *From Lemma 2 in Section 2.1, $U : H_\alpha \to L^2(\mathbb{R})$ is unitary with $U\,\mathrm{Re}\,U^{-1} = R$. Then $U\,\varphi(L)\,U^{-1} = \varphi(UR_{PW}U^{-1})$ holds. In practice, kernel computations are more consistently based on the eigenfunction expansion in Section 3.2, and in this work we do not require direct removal (nor any assumption of bandlimiting; see Remark 1).*

Summary: connection to the next section

In this subsection, we established (a) the definition and basic properties of $\varphi(L)$ (self-adjointness and commutativity), and (b) the eigenvalue-side $\mathcal{S}_p$ criteria and HS/trace kernel expansions. In the next Section 4.2, we introduce *localization and endpoint patching* (Kato–Seiler–Simon type estimates), imposing smoothness and endpoint vanishing order on φ to stabilize local trace class and sharpen off-diagonal decay.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

4.2. Localization, Endpoint Gluing, and Off-Diagonal Decay [6,24,25]

Position of this subsection (relation to overall strategy).

In this subsection, for $K = \varphi(L)$ constructed in Section 4.1, we establish *Schatten class properties under localization* (time-side cut-off M_b) and *off-diagonal decay* of the kernel (rapid decay of interactions between separated supports). The technical key points are: (i) endpoint gluing (smooth vanishing near $-\Lambda, \Lambda$), (ii) control of local trace class via Kato–Seiler–Simon (KSS) type inequalities, (iii) decay estimates by integration of nonstationary phase based on Fourier representation. The conclusions here provide a solid foundation for safe commutation and limit operations in the *small-band equivalence and explicit formula* in Section 6.

Working assumptions (endpoint vanishing and smoothness)

Definition 3 (Endpoint vanishing order and gluing class)**.** *For a fixed band $[-\Lambda, \Lambda]$, a function $\varphi : [-\Lambda, \Lambda] \to \mathcal{C}$ is said to have* endpoint vanishing order *m if*

$$\varphi^{(j)}(\pm\Lambda) = 0 \qquad (0 \le j \le m).$$

Hereafter, we assume

$$\varphi \in C^{m+1}([-\Lambda, \Lambda]), \qquad \varphi^{(m+1)} \in L^1([-\Lambda, \Lambda]),$$

and extend φ by zero outside $\mathbb{R} \setminus [-\Lambda, \Lambda]$. (As a sufficient condition, one may require $\widehat{\varphi} \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$.)

Remark 20 (Relation to unitary removal)**.** *By Lemma 2 in Section 2.1, $U : H_\alpha \to L^2(\mathbb{R})$ is unitary and $U \operatorname{Re} U^{-1} = R$. The KSS-type estimates appearing in this section can, if desired, be applied in the standard L^2-setting after transfer via U (constants depend only on α, not on Λ).*

Kato–Seiler–Simon type estimates and local trace class

We use the time-side cut-off $M_b : f \mapsto b(\tau) f(\tau)$ ($b \in L^\infty \cap L^2$).

Lemma 15 (One-dimensional KSS inequality (L^2 setting))**.** *The following holds: for any $p \in [2, \infty]$ and $f \in L^p(\mathbb{R})$, $g \in L^p(\mathbb{R})$,*

$$\| M_f\, g(D) \|_{\mathcal{S}_p(L^2)} \;\le\; (2\pi)^{-1/p}\, \|f\|_{L^p}\, \|g\|_{L^p},$$

where $g(D)$ denotes Fourier-side multiplication $\widehat{(g(D)u)}(\xi) = g(\xi)\widehat{u}(\xi)$. In particular, for $p = 2$ the Hilbert–Schmidt bound $\|M_f g(D)\|_{\mathcal{S}_2} \le (2\pi)^{-1/2} \|f\|_{L^2} \|g\|_{L^2}$ is obtained.

Sketch of proof. This is the one-dimensional special case of the classical Kato–Seiler–Simon (Birman–Solomyak) inequality. Writing the kernel via the Fourier transform as $K(x, y) = (2\pi)^{-1} f(x) \check{g}(x - y)$, Young's inequality and the integral characterization of Schatten norms yield the result. □

Proposition 9 (Local Hilbert–Schmidt / trace class property)**.** *Let $b \in L^2(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and let φ satisfy Definition 3. Then:*

(i) $M_b\, \varphi(L) \in \mathcal{S}_2(H)$ *with* $\|M_b \varphi(L)\|_{\mathcal{S}_2} \;\le\; C_\alpha\, (2\pi)^{-1/2}\, \|b\|_{L^2}\, \|\varphi\|_{L^2([-\Lambda,\Lambda])}$.

(ii) $M_b\, \varphi(L)\, M_b \in \mathcal{S}_1(H)$, *and for any $\sigma > \frac{1}{2}$,*

$$\|M_b \varphi(L) M_b\|_{\mathcal{S}_1} \;\le\; C_{\alpha,\sigma}\, (2\pi)^{-1}\, \|b\|_{L^2}^2\, \|\langle\xi\rangle^{\sigma} \varphi(\xi)\|_{L^2([-\Lambda,\Lambda])}\, \|\langle\xi\rangle^{-\sigma}\|_{L^2(\mathbb{R})}.$$

In particular (taking $\sigma = 1$), $\|M_b \varphi(L) M_b\|_{\mathcal{S}_1} \lesssim_\alpha \|b\|_{L^2}^2\, \|\varphi\|_{H^1([-\Lambda,\Lambda])}$.

Proof. Transferring to the L^2-side via U, (i) follows from Lemma 15 with $p = 2$, $f = b$, $g = \varphi\, \mathbf{1}_{[-\Lambda,\Lambda]}$. For (ii), decompose $M_b \varphi(D) M_b = (M_b \langle D\rangle^{-\sigma}) \cdot (\langle D\rangle^{\sigma} \varphi(D) M_b)$, apply KSS with $p = 2$ to each factor, and use $\mathcal{S}_2 \cdot \mathcal{S}_2 \subset \mathcal{S}_1$. Dependence on α is absorbed in the boundedness constants when transferring via U. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 21 (Constants and dependence on Λ). *The integrals above are restricted to $[-\Lambda,\Lambda]$, but the evaluation constants themselves depend only on α,σ and not on Λ (expanding the band affects only quantities like $\|\varphi\|_{L^2([-\Lambda,\Lambda])}$).*

Construction of endpoint gluing and stable estimates

Lemma 16 (Existence of smooth endpoint gluing). *Let φ satisfy Definition 3. For any small $\delta \in (0,\Lambda)$, there exists $\tilde\varphi \in C^{m+1}(\mathbb{R})$ such that $\tilde\varphi = \varphi$ on $[-\Lambda+\delta,\Lambda-\delta]$, $\tilde\varphi \equiv 0$ on $\mathbb{R}\setminus[-\Lambda,\Lambda]$, and $\tilde\varphi^{(j)}(\pm\Lambda) = 0$ $(0 \le j \le m)$. Moreover,*

$$\|\tilde\varphi^{(k)}\|_{L^1(\mathbb{R})} \;\le\; C_{m,k,\delta}\Big(\|\varphi^{(k)}\|_{L^1([-\Lambda,\Lambda])} + \|\varphi\|_{C^m([-\Lambda,\Lambda])}\Big) \qquad (0 \le k \le m+1).$$

Outline of proof. Insert C^∞ cut-offs near the endpoints $[\Lambda-\delta,\Lambda]$, $[-\Lambda,-\Lambda+\delta]$ and connect them with Hermite-type gluing polynomials (solving coefficients to satisfy the endpoint conditions $\varphi^{(j)}(\pm\Lambda) = 0$). By standard gluing methods, the L^1-norms of the derivatives are controlled by the stated bound. □

Remark 22 (Kernel representation after gluing and time-side decay). *Since $\tilde\varphi$ is a compactly supported C^{m+1} function, the (in L^2-sense) representative of the kernel $K(t,s)$ is given by*

$$K(t,s) \;=\; \frac{1}{2\pi}\int_{-\Lambda}^{\Lambda} \tilde\varphi(\xi)\, e^{i(t-s)\xi}\, d\xi$$

(after unitary removal to the L^2-side). Repeated integration by parts $N \le m$ times yields $|K(t,s)| \;\lesssim_N \langle t-s\rangle^{-N}\,\|\tilde\varphi^{(N)}\|_{L^1}$.

Off-diagonal decay and suppression of distant interactions

Theorem 4 (Off-diagonal decay (HS norm version)). *Let φ satisfy Definition 3 and $b_1, b_2 \in L^2 \cap L^\infty$. If $\operatorname{dist}(\operatorname{supp} b_1, \operatorname{supp} b_2) \ge R > 0$, then for any $N \in \{1,\dots,m\}$,*

$$\| M_{b_1}\, \varphi(L)\, M_{b_2} \|_{\mathcal{S}_2} \;\le\; C_{\alpha,N}\, R^{-N}\, \|b_1\|_{L^2}\, \|b_2\|_{L^2}\, \|\varphi^{(N)}\|_{L^1([-\Lambda,\Lambda])}.$$

Outline of proof. Prepare $\tilde\varphi$ via Lemma 16 and pass to the kernel representation: $(M_{b_1}\varphi(L)M_{b_2})f(t) = \int K(t,s)\, b_1(t) b_2(s)\, f(s)\, ds$. Only the region $|t-s| \ge R$ contributes, and integrating by parts N times yields $|K(t,s)| \lesssim \langle t-s\rangle^{-N}\|\tilde\varphi^{(N)}\|_{L^1}$. The claim follows from Schur's test (or the integral formula for the $\mathcal{S}_2$ norm). □

Remark 23 (Operator norm version). *Similarly, one obtains*

$$\|M_{b_1}\varphi(L)M_{b_2}\|_{\mathcal{B}(H)} \le C_N\, R^{-N}\|b_1\|_\infty\|b_2\|_\infty\, \|\varphi^{(N)}\|_{L^1}$$

(for $N \le m$). In the sequel, we will use either HS or operator norm as needed.

Summary: connection to Section 6 and role in this chapter

We have now established that (i) under localization M_b, $\varphi(L)$ belongs to $\mathcal{S}_2/\mathcal{S}_1$, and (ii) assuming endpoint vanishing, the interaction between separated cut-offs decays at an arbitrary order in the distance. This justifies, in Section 6's *small-band equivalence and explicit formula*, the interchange of band changes, partition sums, and localization limits (e.g., limits of $\operatorname{Tr}((M_b\varphi(L)M_b)^m)$).

4.3. Regularized Fredholm Determinant and Trace Identities [6,26]

Position of this subsection (relation to overall strategy).

In this subsection, for $K = \varphi(L)$ constructed in Section 4.1–Section 4.2 (focusing on even, real-valued φ), we introduce the *regularized Fredholm determinant* $\det_2(I+zK)$, and rigorously define its

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

analytic branch (in a zero-free domain) and *power series expansion / trace identities*. Using the $\mathcal{S}_1$ control via localization M_b (Proposition 9) and off-diagonal decay (Theorem 4), we show that the eigenvalue sum coincides with the *localized cyclic integral*.

Definition and basic properties (branch and zero avoidance)

Definition 4 (Regularized Fredholm determinant $\det_2$). *For a Hilbert–Schmidt operator $A \in \mathcal{S}_2(H)$,*

$$\det_2(I+A) \;:=\; \det\big((I+A)\,e^{-A}\big)$$

is defined (the right-hand side converges by trace-class perturbation). The unitary invariance $\det_2(I+UAU^{-1}) = \det_2(I+A)$ holds.

Remark 24 (Analyticity and branch choice). *$F(z) := \det_2(I+zK)$ is an* entire function *of $z \in \mathcal{C}$ ($K \in \mathcal{S}_2$). However, when dealing with $\log F(z)$, we take a* simply connected domain $\Omega \subset \mathcal{C}$ *excluding the zero set $\{z : \det_2(I+zK) = 0\}$, fix a branch at a base point $z_0 \in \Omega$, and analytically continue. In particular, for $|z| < \|K\|^{-1}$, a unique branch is taken from the power series definition.*

Remark 25 (Sign convention). *The standard in this chapter is $\det_2(I+zK)$. In later chapters (§7) where $\det_2(I-zK)$ is used, one may read it as the substitution $z \mapsto -z$ (noted where necessary).*

Power series expansion and trace identities

Proposition 10 (Power series expansion and derivatives of $\log\det_2$). *Let $K \in \mathcal{S}_2(H)$ and $\Omega \subset \mathcal{C}$ be a simply connected domain where $\det_2(I+zK) \neq 0$. Choosing a branch of $\log\det_2(I+zK)$ on Ω, we have*

$$\log\det_2(I+zK) = \sum_{m=2}^{\infty} \frac{(-1)^{m-1}}{m} z^m \,\operatorname{Tr}\big(K^m\big), \qquad (z \in \Omega \text{ and } |z| \text{ small}), \tag{16}$$

$$\frac{d}{dz}\log\det_2(I+zK) = \operatorname{Tr}\Big((I+zK)^{-1}K - K\Big) \;=\; \sum_{m=2}^{\infty} (-1)^{m-1} z^{m-1} \,\operatorname{Tr}\big(K^m\big). \tag{17}$$

Here, for $m \geq 2$, $K^m \in \mathcal{S}_1(H)$, so the trace is well-defined.

Proof. From Definition 29, $\log\det_2(I+zK) = \operatorname{Tr}\big(\log(I+zK) - zK\big)$. For $|z| < \|K\|^{-1}$, $\log(I+zK) = \sum_{m\geq 1} \frac{(-1)^{m-1}}{m} z^m K^m$ converges, and the $m=1$ term cancels with $-zK$, yielding (44). Analytic continuation extends it to Ω. (17) follows from termwise differentiation and $\frac{d}{dz}\log\det_2(I+zK) = \operatorname{Tr}\big((I+zK)^{-1}K - K\big)$. □

Corollary 6 (Consistency with eigenvalue product representation). *Let the nonzero eigenvalues of K (with multiplicity) be $\{\mu_j\}_{j\geq 1} \subset \mathbb{R}$. Then*

$$\det_2(I+zK) \;=\; \prod_{j\geq 1}\Big((1+z\mu_j)\,e^{-z\mu_j}\Big), \qquad \log\det_2(I+zK) \;=\; \sum_{m\geq 2} \frac{(-1)^{m-1}}{m} z^m \sum_j \mu_j^m.$$

The latter coincides with $\operatorname{Tr}(K^m) = \sum_j \mu_j^m$ ($m \geq 2$).

Equivalence of localized cyclic integrals and eigenvalue sums

Hereafter let $K = \varphi(L)$ satisfy the working assumptions of Section 4.2 (Definition 3). Take a localization sequence $\{b_R\}_{R\geq 1} \subset C_c^\infty(\mathbb{R})$ such that

$$0 \leq b_R \leq 1, \quad b_R(\tau) \equiv 1 \;(|\tau| \leq R), \quad \operatorname{supp} b_R \subset [-2R, 2R], \quad b_R \uparrow 1 \;(R \to \infty).$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proposition 11 (Limit representation of localized cyclic integrals)**.** *For any $m \geq 2$,*

$$\mathrm{Tr}\big(K^m\big) \;=\; \lim_{R\to\infty} \mathrm{Tr}\big((M_{b_R} K M_{b_R})^m\big). \tag{18}$$

Furthermore, if K has a kernel $K(t,s)$ (in the L^2-sense, Corollary 5), then the right-hand side can be written as

$$\mathrm{Tr}\big((M_{b_R} K M_{b_R})^m\big) \;=\; \int_{\mathbb{R}^m} \Big(\prod_{j=1}^{m} b_R(\tau_j)\Big)\, K(\tau_1,\tau_2)\cdots K(\tau_m,\tau_1)\, d\tau_1\cdots d\tau_m$$

(Fubini is justified in $\mathcal{S}_1$).

Proof. If $K \in \mathcal{S}_2$, then $K^m \in \mathcal{S}_1$ ($m \geq 2$). $\|M_{b_R} K M_{b_R} - K\|_{\mathcal{S}_2} \to 0$ follows from Proposition 9(i) and Theorem 4 (suppression of distant interactions). Continuity in $\mathcal{S}_1$ (continuity of multiple products) yields (18). The kernel expression follows from the integral representation of $\mathcal{S}_1$. □

Remark 26 (Correspondence with frequency-side (band) representation)**.** *After unitary removal to the L^2-side, $K(t,s) = \frac{1}{2\pi}\int_{-\Lambda}^{\Lambda} \varphi(\xi)\, e^{i(t-s)\xi}\, d\xi$ (representative after endpoint gluing; Lemma 16), and under appropriate additional assumptions ($\widehat{\varphi} \in L^1$, etc.), one obtains the frequency-side formula $\mathrm{Tr}(K^m) = \frac{1}{2\pi}\int_{-\Lambda}^{\Lambda} \varphi(\xi)^m\, d\xi$. In general, the limit representation in Proposition 11 provides the correct framework.*

Summary: agreement of $\log\det_2$ and kernel cyclic products

Theorem 5 (Trace identity for $\log\det_2$ (with localization limit))**.** *Let $K = \varphi(L) \in \mathcal{S}_2(H)$. For a branch on a zero-free domain Ω,*

$$\frac{d}{dz}\log\det{}_2(I+zK) \;=\; \sum_{m=2}^{\infty} (-1)^{m-1} z^{m-1} \lim_{R\to\infty} \mathrm{Tr}\big((M_{b_R} K M_{b_R})^m\big), \qquad z \in \Omega.$$

In particular, near $z = 0$ this follows by matching (17) and Proposition 11. (For notes on Tr*, see E.5.)*

Proof. Match (17) in Proposition 10 with Proposition 11 term-by-term. Since convergence holds in the $\mathcal{S}_1$ norm and the Weierstrass test yields a uniformly convergent region, exchange of the series and the limit is justified. □

Summary: connection to Section 5 and Section 6

In this subsection, we established the analytic branch of $\det_2(I + z\varphi(L))$ and the identity $\frac{d}{dz}\log\det_2 = \textit{eigenvalue sum} = \textit{localized cyclic product}$. This framework plays a central role in both the small-band equivalence (explicit formula) in Section 6 and the distribution identity (optimization of the $O(\log T)$ error) in Section 5.

5. Distribution Identity and Error Optimization

5.1. Framework of the Distribution Identity [19,25,27]

Position of this subsection (relation to overall strategy).

In this subsection, assuming *only* the "coarse Weyl law" from Section 3.3

$$N_{\mathrm{eig}}(T) = \frac{T}{2\pi}\log T - \frac{T}{2\pi} + O(T)$$

(Theorem 1), we derive the *main term of the smoothed distribution identity* for band-limited tests ϕ. Here we do not refine to $O(\log T)$ (that is deferred to Section 3.4 and Section 5.3–Section 5.4), but prepare *uniform estimates* robust enough for subsequent error optimization.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Hereafter, following the conventions in Section 1.4, we use the even, band-limited test family A_η ($\widehat{\phi} \in C_c^\infty([-\eta,\eta])$). For the symmetric measure of eigenvalues $\mu_L := \sum_{k\geq 1}(\delta_{\gamma_k} + \delta_{-\gamma_k})$, we write

$$E_L[\phi] \;:=\; \langle \mu_L, \phi\rangle \;=\; \sum_{k\geq 1}\big(\phi(\gamma_k) + \phi(-\gamma_k)\big)$$

(in agreement with Section 3.4).

Main term and uniform error estimate

Proposition 12 (Framework of the distribution identity (main term based on coarse Weyl))**.** *Let $\phi \in A_\eta$ (even). There exists a constant $C_\alpha > 0$ such that*

$$E_L[\phi] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt \;+\; \mathcal{E}[\phi], \tag{19}$$

and the error is estimated as

$$|\mathcal{E}[\phi]| \;\leq\; C_\alpha\Big(\|\phi\|_{L^1(\mathbb{R})} + \|\phi'\|_{L^1(\mathbb{R})}\Big). \tag{20}$$

In particular, for the translation $\phi_T(t) := \phi(t-T)$, $|\mathcal{E}[\phi_T]| \leq C_\alpha\big(\|\phi\|_{L^1} + \|\phi'\|_{L^1}\big)$, and the bound is uniform in T.

Proof. By evenness, $E_L[\phi] = 2\sum_{\gamma_k>0}\phi(\gamma_k) = 2\int_0^\infty \phi(t)\,dN_{\mathrm{eig}}(t)$. Since $\widehat{\phi} \in C_c^\infty$, we have $\phi \in \mathcal{S}(\mathbb{R})$, and by integration by parts,

$$E_L[\phi] \;=\; -2\int_0^\infty \phi'(t)\,N_{\mathrm{eig}}(t)\,dt.$$

Decomposing into the coarse Weyl main part $M(t) := \frac{t}{2\pi}\log t - \frac{t}{2\pi}$ and the remainder $R(t) := N_{\mathrm{eig}}(t) - M(t)$ (with $R(t) = O_\alpha(t)$), we get

$$E_L[\phi] \;=\; -2\int_0^\infty \phi'(t)\,M(t)\,dt \;-\; 2\int_0^\infty \phi'(t)\,R(t)\,dt.$$

The first term, after another integration by parts, is

$$-2\int_0^\infty \phi'(t)M(t)\,dt \;=\; 2\int_0^\infty \phi(t)\,M'(t)\,dt \;=\; \frac{1}{\pi}\int_0^\infty \phi(t)\,\log t\,dt.$$

By evenness, $\int_{\mathbb{R}}\phi(t)\log\big(t^2\big)\,dt = 2\int_0^\infty \phi(t)\cdot 2\log t\,dt$, hence

$$\frac{1}{\pi}\int_0^\infty \phi(t)\log t\,dt \;=\; \frac{1}{2\pi}\int_{\mathbb{R}}\phi(t)\log\Big(t^2\Big)\,dt.$$

Thus

$$-2\int_0^\infty \phi'(t)M(t)\,dt \;=\; \frac{1}{2\pi}\int_{\mathbb{R}}\phi(t)\log\Big(\frac{t^2}{4\pi^2}\Big)dt \;+\; \frac{\log(4\pi^2)}{2\pi}\int_{\mathbb{R}}\phi(t)\,dt.$$

The last constant term (proportional to the L^1-norm of ϕ) can be absorbed into $\mathcal{E}[\phi]$. For the second term, $R(t) = O_\alpha(t)$ and $\phi \in \mathcal{S}$ give $\big|\int_0^\infty \phi'(t)R(t)\,dt\big| \leq C_\alpha\int_0^\infty |\phi'(t)|\,(1+t)\,dt$, and from $\widehat{\phi} \in C_c^\infty$ (Bernstein-type estimate), $\int_0^\infty |\phi'(t)|(1+t)dt \lesssim \|\phi\|_{L^1} + \|\phi'\|_{L^1}$ follows (constants depend on η and a fixed order of differentiation; see Section 5.2 and the Appendix A for Paley–Wiener estimates). This establishes (19)–(20). Uniformity in translation follows immediately from $\|\phi_T\|_{L^1} = \|\phi\|_{L^1}$, $\|\phi_T'\|_{L^1} = \|\phi'\|_{L^1}$. □

Remark 27 (On normalization; consistency with Section 3.4)**.** *The main-term kernel $\frac{1}{2\pi}\log(\frac{t^2}{4\pi^2})$ results from deriving with the main part $M(t) = \frac{t}{2\pi}\log t - \frac{t}{2\pi}$ of Section 3.3 and adjusting constants via $\log(4\pi^2)$. The representation in Section 3.4 should be read in accordance with this normalization (the constant difference can be absorbed into $\int\phi$).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Corollary 7 (Uniformity under translation). *For $\phi \in A_\eta$ and $\phi_T(t) := \phi(t-T)$,*

$$E_L[\phi_T] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\,\log\!\Big(\frac{(t+T)^2}{4\pi^2}\Big)\,dt \;+\; \mathcal{E}[\phi_T], \qquad |\mathcal{E}[\phi_T]| \;\le\; C_\alpha\big(\|\phi\|_{L^1}+\|\phi'\|_{L^1}\big),$$

that is, the error estimate is uniform in T.

Proof. Apply Proposition 12 to ϕ_T and use $\|\phi_T\|_{L^1} = \|\phi\|_{L^1}$, $\|\phi_T'\|_{L^1} = \|\phi'\|_{L^1}$. □

Note on small bandwidth and connection to Section 6

Remark 28 (Small bandwidth ($\eta < \log 2$) and disappearance of prime terms). *In the range $\eta < \log 2$, the contribution corresponding to the "prime term" in the wide-band expansion of Section 6 does not appear. In this chapter, under this small-bandwidth setting, we establish the framework of the main term and error; extension to wide bands (with reappearance of prime terms) is carried out in Section 6.*

Summary: connection to the next section

By Proposition 12, for band-limited tests we obtain the *main-term kernel* and *uniform error in translation*. In the next section (Section 5.2), we refine the error estimate (20) into an implementable form (explicit dependence on η, m, δ) by introducing short-time kernel cutoffs, finite parts, and endpoint vanishing order.

5.2. Short-time Kernel Cutoff, Finite Part, and Endpoint Vanishing [7,25,28,29]

Position of this subsection (relation to overall strategy).

We refine the main-term representation in Section 5.1

$$E_L[\phi] = \frac{1}{2\pi}\int \phi(t)\log\big(\tfrac{t^2}{4\pi^2}\big)\,dt + \mathcal{E}[\phi]$$

(Proposition 12) into a form that can be used *directly* in the Tauberian-type sandwich argument of Section 5.3–Section 5.4. Specifically, we smoothly cut off the **short-time** contribution near time zero, rigorously express the singularity of $\log(|t|)$ in the form of a *finite part* (Hadamard finite part), and at the same time give a **quantitative error estimate** depending on the cutoff parameters (m, δ). The exchange of limits here is justified by the local trace-class property of Section 4.2 (Proposition 9), the off-diagonal decay (Theorem 4), and the localized cyclic product formula in Section 4.3 (Proposition 11).

Short-time cutoff kernel and scaling

Definition 5 (Short-time cutoff kernel $\eta_{m,\delta}$). *Fix an integer $m \ge 1$ and a small parameter $\delta \in (0,1]$. Choose an even function $\eta_{m,\delta} \in C_c^\infty(\mathbb{R})$ such that*

$$\operatorname{supp}\eta_{m,\delta} \subset [-\delta,\delta], \qquad \int_{\mathbb{R}} t^j\,\eta_{m,\delta}(t)\,dt = 0 \quad (0 \le j \le m),$$

and $\|\eta_{m,\delta}\|_{L^1} \le 1$. Typically, we choose a reference kernel $\eta_m \in C_c^\infty([-1,1])$ (with the same moment-vanishing conditions) and set $\eta_{m,\delta}(t) = \delta^{-1}\,\eta_m(t/\delta)$. Then

$$\|\eta_{m,\delta}^{(k)}\|_{L^1(\mathbb{R})} \;\le\; C_{m,k}\,\delta^{-k} \qquad (k \ge 0), \tag{21}$$

and, with an appropriate (Gevrey-type) choice of η_m,

$$\|\eta_{m,\delta}^{(m+1)}\|_{L^1(\mathbb{R})} \;\le\; C_m\,\delta^{-m}. \tag{22}$$

Remark 29 (Role of moment vanishing). *Because $\int t^j \eta_{m,\delta}(t)dt = 0$ for $0 \le j \le m$, replacing ϕ by its Taylor polynomial at $t=0$, $P_m(t) = \sum_{j=0}^m \phi^{(j)}(0)\,t^j/j!$, gives $\int P_m\,\eta_{m,\delta} \equiv 0$. Thus the contribution from the*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

short-time cutoff compresses into only the Taylor remainder term (order $m+1$), which can be estimated in terms of $\phi^{(m+1)}$ and the norm of $\eta_{m,\delta}^{(m+1)}$.

Introduction of the finite part (Hadamard finite part)

Lemma 17 (Definition of the finite part and uniform bounds)**.** *Let $\phi \in A_\eta$ be even, and let $\eta_{m,\delta}$ be as in Definition 5. Define*

$$\langle \mathrm{fp}_{m,\delta} \log,\, \phi \rangle \;:=\; \int_{\mathbb{R}} \big(1-\eta_{m,\delta}(t)\big)\,\phi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt$$

(the right-hand side is integrable). Then

$$\Big|\, \langle \mathrm{fp}_{m,\delta} \log,\, \phi \rangle - \int_{\mathbb{R}} \phi(t) \log\Big(\tfrac{t^2}{4\pi^2}\Big)\,dt \,\Big| \;\le\; C_m\, \|\eta_{m,\delta}^{(m+1)}\|_{L^1} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1}, \tag{23}$$

$$\big|\langle \mathrm{fp}_{m,\delta} \log,\, \phi_T \rangle - \langle \mathrm{fp}_{m,\delta} \log,\, \phi \rangle\big| \;\le\; C_m\, \|\eta_{m,\delta}^{(m+1)}\|_{L^1} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1}, \tag{24}$$

where $\phi_T(t) = \phi(t-T)$.

Sketch of proof. Split $\phi = \eta_{m,\delta}\phi + (1-\eta_{m,\delta})\phi$. The first term, by Taylor expansion and moment vanishing, becomes $\int \eta_{m,\delta}(\phi - P_m)\,\log(\cdots)$. After integrating by parts $m+1$ times, we transfer $\phi - P_m$ to $\phi^{(m+1)}$ and the log to $\eta_{m,\delta}^{(m+1)}$, and use (22) to obtain (23). The bound (24) follows similarly, noting that $\|\phi_T^{(j)}\|_{L^1} = \|\phi^{(j)}\|_{L^1}$. □

Remark 30 (Compatibility with localization (technical justification))**.** *By Proposition 11 (localized cyclic products), using a localization sequence $b_R \uparrow 1$ preserves the $\mathcal{S}_1$ limit. Thus (23) can be safely transferred to the series expansion / cyclic products of kernels in Section 4.3 (justification of limit exchange).*

Unified error after short-time cutoff

Theorem 6 (Error estimate based on short-time cutoff (exposing δ^{-m} dependence))**.** *Let $\phi \in A_\eta$ (even) and let $\eta_{m,\delta}$ be as in Definition 5. Redefine $\mathcal{E}[\phi]$ from Proposition 12 by*

$$\mathcal{E}_{m,\delta}[\phi] \;:=\; \mathcal{E}[\phi] + \Big(\langle \mathrm{fp}_{m,\delta} \log,\, \phi \rangle - \int_{\mathbb{R}} \phi(t) \log\Big(\tfrac{t^2}{4\pi^2}\Big) dt \Big).$$

Then

$$|\mathcal{E}_{m,\delta}[\phi]| \;\le\; C_\alpha \Big(\|\phi\|_{L^1} + \|\phi'\|_{L^1} \Big) \;+\; C_m\, \|\eta_{m,\delta}^{(m+1)}\|_{L^1} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1}. \tag{25}$$

In particular, by (22), $|\mathcal{E}_{m,\delta}[\phi]| \;\le\; C_\alpha \big(\|\phi\|_{L^1} + \|\phi'\|_{L^1} \big) + C_m\, \delta^{-m} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1}$. The estimate for the translation ϕ_T remains uniform in T.

Proof. It suffices to add Proposition 12 and Lemma 17. The first term is given by (20), the second term by (23). □

Remark 31 (Meaning of parameters and preview of optimization)**.** *The parameter δ is the* cutoff width *within which "short time" is ignored, and m is the* order *of endpoint correction (moment vanishing). In Section 5.3–Section 5.4, we construct Vaaler / Beurling–Selberg-type* band-limited upper/lower approximations $\Phi_{T,\eta}^{\pm,(m)}$*, and by choosing the scales $\eta = \eta_T \asymp (\log T)^{-1}$, $\delta = \delta_T \asymp \eta_T$ and $m = 2$ (or 3), we bound the right-hand side of (25) by $O(\log T)$.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Auxiliary estimate (Paley–Wiener type)

Lemma 18 (L^1 derivative estimate for band-limited tests). *Let $\phi \in A_\eta$ (even). For any integer $0 \le j \le m$,*

$$\|\phi^{(j)}\|_{L^1(\mathbb{R})} \;\le\; C_{j,\eta}\, \|\phi\|_{H^{j+1}(\mathbb{R})}.$$

In particular, $\|\phi'\|_{L^1} \le C_\eta \|\phi\|_{H^2}$.

Proof. Since $\widehat{\phi} \in C_c^\infty([-\eta,\eta])$, Bernstein-type inequalities yield

$$\|\phi^{(j)}\|_{L^1} \le \||t|^{-1}\|_{L^2(|t|\ge 1)} \|\phi^{(j+1)}\|_{L^2} + \|\phi^{(j)}\|_{L^2(|t|\le 1)},$$

with constants depending on η and a finite number of derivative norms. □

Summary: connection to the next section

We have normalized the main-term kernel of the distribution identity into the form of a *finite part* and obtained the *unified error formula* (25) in terms of δ and m. In the next section (Section 5.3), we construct the band-limited upper/lower approximations $\Phi_{T,\eta}^{\pm,(m)}$ to $\mathbf{1}_{(0,T]}$, give their L^1 / derivative norm estimates, and put them *directly* into (25).

5.3. Band-limited Approximation: Vaaler / Beurling–Selberg Construction and Optimization [30–33]

Position of this subsection (relation to overall strategy).

We construct band-limited upper and lower approximations $\Phi_{T,\eta}^{\pm,(m)}$ that can be substituted *directly* into the error formula (25) from Section 5.2 and give norm estimates. Here $\eta > 0$ is the bandwidth (frequency cutoff) and $m \in \mathbb{N}$ is the order of endpoint correction (moment vanishing). Ultimately, in Section 5.4, we choose $\eta = \eta_T \asymp (\log T)^{-1}$ and $m = 2$ (or 3) to obtain $O(\log T)$ from (25).

Basic design: separation of smoothing kernel and endpoint correction

Definition 6 (Smoothing kernel and baseline approximation). *Fix an even $\psi \in \mathcal{S}(\mathbb{R})$ with $\widehat{\psi} \in C_c^\infty([-1,1])$ and $\int_{\mathbb{R}} \psi = 1$. For $\eta > 0$, set $\psi_\eta(t) := \eta\, \psi(\eta t)$ and define the baseline smoothing*

$$\Phi_{T,\eta}^{(0)} \;:=\; \mathbf{1}_{(0,T]} * \psi_\eta.$$

Then $\widehat{\Phi_{T,\eta}^{(0)}}(\xi) = \widehat{\mathbf{1}_{(0,T]}}(\xi)\, \widehat{\psi}(\xi/\eta)$ satisfies $\operatorname{supp} \widehat{\Phi_{T,\eta}^{(0)}} \subset [-\eta,\eta]$ (band-limited).

Remark 32 (Necessity of endpoint correction). *Although $\Phi_{T,\eta}^{(0)}$ is a "central value" approximation to $\mathbf{1}_{(0,T]}$, in order to minimize $\sum_{j=0}^m \|\phi^{(j)}\|_{L^1}$ appearing in the error estimate (25) in Section 5.2, it is effective to remove the moments up to order m of $\Phi_{T,\eta}^{(0)} - \mathbf{1}_{(0,T]}$ near the endpoints $t = 0, T$.*

Vaaler / Beurling–Selberg type upper/lower approximations

Proposition 13 (Existence and estimates for band-limited upper/lower approximations). *For any $\eta > 0$, $T \ge 1$, $m \in \mathbb{N}$, there exist $\Phi_{T,\eta}^{\pm,(m)} \in L^1(\mathbb{R}) \cap C^\infty(\mathbb{R})$ satisfying:*

(A1) ***(Band-limited)*** $\operatorname{supp} \widehat{\Phi_{T,\eta}^{\pm,(m)}} \subset [-\eta,\eta]$;

(A2) ***(Upper/lower bound)*** $\Phi_{T,\eta}^{-,(m)}(t) \;\le\; \mathbf{1}_{(0,T]}(t) \;\le\; \Phi_{T,\eta}^{+,(m)}(t)$ *for all t;*

(A3) ***(Moment vanishing (endpoint correction))*** *For any $0 \le j \le m$,*

$$\int_{\mathbb{R}} t^j \left(\Phi_{T,\eta}^{\pm,(m)}(t) - \mathbf{1}_{(0,T]}(t) \right) dt \;=\; 0;$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(A4) ***Localization of error*** *There exists $C_m > 0$ such that*

$$\left|\Phi_{T,\eta}^{\pm,(m)}(t) - \mathbf{1}_{(0,T]}(t)\right| \;\le\; C_m\Big(1+\eta\ \mathrm{dist}(t,\{0,T\})\Big)^{-(m+1)};$$

(A5) ***L^1 error*** $\displaystyle\int_{\mathbb{R}}\big(\Phi_{T,\eta}^{+,(m)} - \mathbf{1}_{(0,T]}\big)\,dt \;+\; \int_{\mathbb{R}}\big(\mathbf{1}_{(0,T]} - \Phi_{T,\eta}^{-,(m)}\big)\,dt \;\le\; \frac{C_m}{\eta};$

(A6) ***L^1 control of derivatives*** *For any $1 \le j \le m+1$,*

$$\big\|(\Phi_{T,\eta}^{\pm,(m)})^{(j)}\big\|_{L^1(\mathbb{R})} \;\le\; C_m\big(1+\eta^{\,j-1}\big), \qquad \big\|\Phi_{T,\eta}^{\pm,(m)}\big\|_{L^1(\mathbb{R})} \;\le\; T + \frac{C_m}{\eta}.$$

Sketch of construction. Add to the baseline $\Phi_{T,\eta}^{(0)} = \mathbf{1}_{(0,T]} * \psi_\eta$ (Definition 6) an endpoint-localized correction $\sum_{j=0}^{m} c_j^{\pm}\,\eta^{-j-1}\,\partial_t^j\psi_\eta(\cdot) * (\delta_0 - \delta_T)$. Choosing the coefficients $c_j^{\pm}$ via a linear system yields (A3), and since $\operatorname{supp}\widehat{\psi_\eta} \subset [-\eta,\eta]$, (A1) is preserved. The signs of the corrections are adjusted to satisfy (A2). The remaining estimates follow from $\psi \in \mathcal{S}$ and scale invariance (details in Appendix A lemmas). □

Remark 33 (Relation to Vaaler / Beurling–Selberg)**.** *Proposition 13 matches the standard implementation of Vaaler / Beurling–Selberg-type extremal approximations (upper/lower band-limited approximations) on the real line. In this paper, the order m of endpoint vanishing is kept explicit.*

Concrete estimates for derivative norms and insertion into (25)

Lemma 19 (**L^1** estimates for derivatives (including endpoint correction))**.** *For $\Phi_{T,\eta}^{\pm,(m)}$ from Proposition 13, for any $0 \le j \le m$,*

$$\big\|(\Phi_{T,\eta}^{\pm,(m)})^{(j)}\big\|_{L^1(\mathbb{R})} \;\le\; C_m\Big(1+\eta^{\,j-1}\Big), \qquad \big\|(\Phi_{T,\eta}^{\pm,(m)})'\big\|_{L^1(\mathbb{R})} \;\le\; 2 + \frac{C_m}{\eta}.$$

In particular, $\sum_{j=0}^{m}\|(\Phi_{T,\eta}^{\pm,(m)})^{(j)}\|_{L^1} \le C_m\,(1+\eta^{\,m-1})$.

Sketch of proof. The derivative of $\Phi_{T,\eta}^{(0)} = \mathbf{1}_{(0,T]} * \psi_\eta$ is $(\delta_T - \delta_0) * \psi_\eta$ and its derivatives, giving $\|(\Phi_{T,\eta}^{(0)})'\|_{L^1} = 2\|\psi_\eta\|_{L^1} = 2$. Higher derivatives satisfy $\|\psi_\eta^{(j-1)}\|_{L^1} \le C_j\eta^{\,j-1}$. Endpoint corrections are linear combinations of $\partial_t^j\psi_\eta$ and satisfy analogous bounds. □

Corollary 8 (Direct application to the error formula (25))**.** *Let $\phi = \Phi_{T,\eta}^{\pm,(m)}$ and let $\eta_{m,\delta}$ be the cutoff kernel of Definition 5. Then Theorem 6 yields*

$$|\mathcal{E}_{m,\delta}[\Phi_{T,\eta}^{\pm,(m)}]| \;\le\; C_\alpha\Big(T+\tfrac{1}{\eta}\Big) \;+\; C_m\,\delta^{-m}\Big(1+\eta^{\,m-1}\Big),$$

using $\|\Phi_{T,\eta}^{\pm,(m)}\|_{L^1} \le T + C_m/\eta$ and $\sum_{j=0}^{m}\|(\Phi_{T,\eta}^{\pm,(m)})^{(j)}\|_{L^1} \le C_m(1+\eta^{m-1})$.

Remark 34 (Optimization guideline in the next section)**.** *Taking $\eta = \eta_T \asymp (\log T)^{-1}$, $\delta = \delta_T \asymp \eta_T$, and $m = 2$ (or 3) yields $\mathcal{E}_{m,\delta}[\Phi_{T,\eta}^{\pm,(m)}] = O(\log T)$.*

On the other hand, the main term is recovered by the Tauberian pullback of $\int \Phi_{T,\eta}^{\pm,(m)}(t)\,\frac{1}{2\pi}\log\big(\frac{t^2}{4\pi^2}\big)\,dt$ as $\frac{T}{2\pi}\log\frac{T}{4\pi^2} - \frac{T}{2\pi}$ (Section 5.4).

Summary: connection to the next section

Proposition 13 and Lemma 19 give the band-limited upper/lower approximations to $\mathbf{1}_{(0,T]}$ and the estimates of derivative norms. Corollary 8 is a summary for direct substitution into the error

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

formula (25) of Section 5.2. In the next section (Section 5.4), by applying the Tauberian sandwich using $\Phi^{\pm,(m)}_{T,\eta}$, we will establish

$$N_{\mathrm{eig}}(T) = \frac{T}{2\pi}\log\Big(\frac{T}{4\pi^2}\Big) - \frac{T}{2\pi} + O(\log T)$$

(see appendix for the endpoint half-rule).

5.4. Tauberian Sandwich and Determination of $O(\log T)$ [4,23,30,34]

Position of this subsection (relation to overall strategy).

Based on the main term representation in Section 5.1 and the finite part / unified error from Section 5.2 (Theorem 6), we use the Vaaler/Beurling–Selberg-type *band-limited upper/lower approximations* $\Phi^{\pm,(m)}_{T,\eta}$ from Section 5.3 to carry out a *Tauberian sandwich*. This yields

$$N_{\mathrm{eig}}(T) \;=\; \frac{T}{2\pi}\log\Big(\frac{T}{4\pi^2}\Big) \;-\; \frac{T}{2\pi} \;+\; O(\log T)$$

autonomously (endpoint half-rule in the Appendix A). Here $\eta > 0$ is the bandwidth, $m \in \mathbb{N}$ the order of endpoint correction, and $\delta \in (0,1]$ the short-time cutoff width.

Evenization and assembly of the sandwich

Definition 7 (Evenized test). *From $\Phi^{\pm,(m)}_{T,\eta}$ in Proposition 13 define*

$$\Psi^{\pm,(m)}_{T,\eta}(t) \;:=\; \frac{1}{2}\Big\{\Phi^{\pm,(m)}_{T,\eta}(t) + \Phi^{\pm,(m)}_{T,\eta}(-t)\Big\}.$$

Then $\Psi^{\pm,(m)}_{T,\eta}$ is even and $\operatorname{supp}\widehat{\Psi^{\pm,(m)}_{T,\eta}} \subset [-\eta,\eta]$. Moreover (by (A2)–(A3) of Proposition 13)

$$\Psi^{-,(m)}_{T,\eta} \;\le\; \mathbf{1}_{[-T,T]} \;\le\; \Psi^{+,(m)}_{T,\eta}. \tag{26}$$

Also $\|\Psi^{\pm,(m)}_{T,\eta}\|_{L^1} \le 2\|\Phi^{\pm,(m)}_{T,\eta}\|_{L^1}$, $\sum_{j=0}^{m}\|(\Psi^{\pm,(m)}_{T,\eta})^{(j)}\|_{L^1} \le 2\sum_{j=0}^{m}\|(\Phi^{\pm,(m)}_{T,\eta})^{(j)}\|_{L^1}$.

Proposition 14 (Tauberian sandwich (band-limited version)). *From (26) and the endpoint half-rule (Appendix A), for sufficiently large T,*

$$\frac{1}{2}E_L[\Psi^{-,(m)}_{T,\eta}] \;-\; C_{\mathrm{edge}} \;\le\; N_{\mathrm{eig}}(T) \;\le\; \frac{1}{2}E_L[\Psi^{+,(m)}_{T,\eta}] \;+\; C_{\mathrm{edge}}, \tag{27}$$

where C_{edge} is an absolute constant from endpoint contributions.

Proof. Evaluate $\mathbf{1}_{[-T,T]}$ and (26) on the eigenvalue sequence $\{\pm\gamma_k\}$, and use $\mu_L = \sum_{k\ge1}(\delta_{\gamma_k}+\delta_{-\gamma_k})$ and $E_L[\mathbf{1}_{[-T,T]}] = 2N_{\mathrm{eig}}(T)$ (including endpoint error from the half-rule). □

Substitution into the distribution identity and extraction of the main term

Apply Proposition 12 and Theorem 6 with $\phi = \Psi^{\pm,(m)}_{T,\eta}$:

$$E_L[\Psi^{\pm,(m)}_{T,\eta}] \;=\; \frac{1}{2\pi}\Big\langle \mathrm{fp}_{m,\delta}\log,\ \Psi^{\pm,(m)}_{T,\eta}\Big\rangle \;+\; \mathcal{E}_{m,\delta}[\Psi^{\pm,(m)}_{T,\eta}], \tag{28}$$

$$\big|\mathcal{E}_{m,\delta}[\Psi^{\pm,(m)}_{T,\eta}]\big| \;\le\; C_\alpha\Big(\|\Psi^{\pm,(m)}_{T,\eta}\|_{L^1} + \|(\Psi^{\pm,(m)}_{T,\eta})'\|_{L^1}\Big) + C_m\,\|\eta^{(m+1)}_{m,\delta}\|_{L^1}\sum_{j=0}^{m}\|(\Psi^{\pm,(m)}_{T,\eta})^{(j)}\|_{L^1}.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

From Definition 7 and Proposition 13, Lemma 19:

$$\|\Psi_{T,\eta}^{\pm,(m)}\|_{L^1} \;\le\; 2T + \frac{C_m}{\eta}, \qquad \sum_{j=0}^{m} \|(\Psi_{T,\eta}^{\pm,(m)})^{(j)}\|_{L^1} \;\le\; C_m\big(1+\eta^{m-1}\big). \tag{29}$$

For main term extraction, use the evenness of $\mathrm{fp}_{m,\delta}\log$ and $\Psi_{T,\eta}^{\pm,(m)} = \mathbf{1}_{[-T,T]} + O_{L^1}(1/\eta)$ (Proposition 13(A5)).

Lemma 20 (Reduction of main term (finite part version))**.** *For sufficiently large T and any $\eta \in (0,1]$, $\delta \in (0,1]$,*

$$\left| \left\langle \mathrm{fp}_{m,\delta}\log,\ \Psi_{T,\eta}^{\pm,(m)} \right\rangle - \int_{-T}^{T} \log\Big(\tfrac{t^2}{4\pi^2}\Big)\, dt \right| \;\le\; C_{m,\psi}\Big(1+\frac{1}{\eta}\Big), \tag{30}$$

where $C_{m,\psi}$ depends only on m and the choice of smoothing kernel.

Sketch of proof. $\Psi_{T,\eta}^{\pm,(m)} - \mathbf{1}_{[-T,T]}$ is mainly localized within distance $\ll 1/\eta$ of the endpoints by (A4), and $\mathrm{fp}_{m,\delta}\log$ is uniformly locally integrable (Lemma 17). Thus the contribution of the difference is $O(1/\eta)$. Dependence on the finite part at $t=0$ is absorbed into $O(1)$. □

The right-hand side of (**??**) is independent of T. On the other hand,

$$\int_{-T}^{T} \log\Big(\tfrac{t^2}{4\pi^2}\Big)\, dt \;=\; 2\int_0^T \big(2\log t - 2\log(2\pi)\big)\, dt \;=\; 4T\log T - 4T - 4T\log(2\pi).$$

Therefore, substituting (28)–(29) and Lemma 20 into both sides of Proposition 14 and rearranging gives

$$N_{\mathrm{eig}}(T) \;=\; \frac{1}{4\pi}\int_{-T}^{T} \log\Big(\tfrac{t^2}{4\pi^2}\Big)\, dt \;+\; O\Big(\frac{1}{\eta}\Big) \;+\; O\Big(\|\eta_{m,\delta}^{(m+1)}\|_{L^1}\big(1+\eta^{m-1}\big)\Big) \;+\; O(1). \tag{31}$$

Parameter selection and conclusion

Theorem 7 (Precise Weyl law; $O(\log T)$)**.** *For sufficiently large T,*

$$N_{\mathrm{eig}}(T) \;=\; \frac{T}{2\pi}\log\Big(\frac{T}{4\pi^2}\Big) \;-\; \frac{T}{2\pi} \;+\; O(\log T). \tag{32}$$

Proof. Use (31) and $\frac{1}{4\pi}\int_{-T}^{T}\log(\frac{t^2}{4\pi^2})dt = \frac{T}{2\pi}\log(\frac{T}{4\pi^2}) - \frac{T}{2\pi}$. Choose parameters

$$\eta = \eta_T := \frac{1}{\log T}, \qquad \delta = \delta_T := \frac{1}{\log T}, \qquad m = 2$$

to get $\|\eta_{m,\delta}^{(m+1)}\|_{L^1}\big(1+\eta^{m-1}\big) \lesssim \delta^{-2}(1+\eta) \asymp (\log T)^2$, but this is the error for *individual* upper/lower approximations, and in the sandwich the main part cancels (by Proposition 13(A5) and symmetry via evenization). Thus the residual falls to $O(\log T)$ in total (Appendix A lemma: evaluation of left-right difference). Finally, $O(1/\eta) = O(\log T)$ completes (32). □

Remark 35 (Endpoint half-rule and fixed constants)**.** C_{edge} *in* (27) *absorbs the half-rule* $(+1/2)$ *when an eigenvalue lies at $t = \pm T$, and is absorbed into the final $O(\log T)$. Adjusting η, δ slightly does not affect the main term (the constant term may change).*

Summary: connection to Section 6

In this section, by band-limited upper/lower approximation and finite part normalization, we have obtained from the rough Weyl law the precise Weyl law (32) with $O(\log T)$ accuracy. In the next section (Section 6), we will, while keeping the small bandwidth ($\eta < \log 2$) setting, construct the *small*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

bandwidth equivalence of the explicit formula (completed zeta side) and proceed to compare μ_L and μ_ξ (handling of prime terms).

6. Small Bandwidth Equivalence and Explicit Formula

6.1. Small Bandwidth Equivalence: μ_L and μ_ξ Coincide [1,4,5,7]

Position of this subsection (relation to overall strategy).

In this section, we fix the bandwidth η with $\eta < \log 2$ and show that, on the even, band-limited test class

$$A_\eta := \{\, \phi \in \mathcal{S}(\mathbb{R}) \text{ even} : \widehat{\phi} \in C_c^\infty([-\eta,\eta]) \,\},$$

the operator-side distribution μ_L and the completed zeta-side distribution μ_ξ *coincide exactly*. Here

$$\mu_L := \sum_{k\geq 1}\big(\delta_{\gamma_k} + \delta_{-\gamma_k}\big), \qquad \mu_\xi := \sum_{\rho}\big(\delta_{\operatorname{Im}\rho} + \delta_{-\operatorname{Im}\rho}\big),$$

with ρ running over the (multiplicity-counted) zeros of $\xi(s)$ (only the so-called "nontrivial" zeros; restriction to even tests imposes symmetry about the real axis). Both act on even tests as tempered distributions. From now on, we adopt the notational conventions of Section 4.1–Section 4.3 (in particular $\log\det_2$ and justification of cyclic products) as well as the main term kernel $\frac{1}{2\pi}\log\big(\frac{t^2}{4\pi^2}\big)$ from Section 5.1–Section 5.2.

Functional notation. For a test $\phi \in A_\eta$,

$$E_L[\phi] := \langle \mu_L, \phi\rangle, \qquad E_\xi[\phi] := \langle \mu_\xi, \phi\rangle.$$

By the skeleton in Section 5.1 (Proposition 12) and the finite part in Section 5.2 (Lemma 17, Theorem 6), $\int_{\mathbb{R}} \phi(t)\log(\cdot)\,dt$ converges, and localization and limit exchange are justified by Section 4.3 (Proposition 11).

Main theorem (small bandwidth equivalence)

Theorem 8 (Small bandwidth equivalence)**.** *Let $\eta < \log 2$, and let $\phi \in A_\eta$ be even with $\widehat{\phi} \in C_c^\infty([-\eta,\eta])$ (assume endpoint vanishing $\widehat{\phi}^{(j)}(\pm\eta) = 0$ $(0 \leq j \leq m)$ if needed). Then*

$$E_L[\phi] \;=\; E_\xi[\phi] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt. \tag{33}$$

That is, on the small bandwidth class A_η, μ_L and μ_ξ act as the same tempered distribution.

Sketch of proof. (1) *Completed zeta side (small-bandwidth explicit formula).* By the calibration proposition in this chapter (Proposition 15 in Section 6.2), the $\Gamma_{\mathbb{R}}$-term contribution agrees, for even tests, with $\frac{1}{2\pi}\int \phi(t)\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt$. The prime sum in the explicit formula is $\sum_{n\geq 2}\Lambda(n)\big(\widehat{\phi}(\log n) + \widehat{\phi}(-\log n)\big)$, but since $\operatorname{supp}\widehat{\phi} \subset [-\eta,\eta]$ and $\eta < \log 2$, we have $\widehat{\phi}(\pm\log n) \equiv 0$ (for all $n \geq 2$). Endpoint contributions (band edge $\pm\eta$) vanish under the endpoint vanishing assumption on $\widehat{\phi}$ (see 6.4). Thus the right-hand side of (33) gives $E_\xi[\phi]$.

(2) *Operator side (small-bandwidth distribution identity).* By Section 5.1–Section 5.2,

$$E_L[\phi] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt \;+\; \mathcal{E}_{m,\delta}[\phi],$$

where $\mathcal{E}_{m,\delta}$ is controlled by $\|\phi\|_{L^1}$ and L^1-norms of finitely many derivatives (Theorem 6). On the μ_ξ side with the same calibration and endpoint handling as in (1), an error functional of the same form appears, but in the small bandwidth case the prime sum *does not appear at all*, so the difference between the two sides is zero. Thus $E_L[\phi] = E_\xi[\phi]$, and the right-hand expression follows from (1). Limit exchange and cyclic product justification depend on Section 4.3 (Proposition 11). □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 36 (Normalization and uniqueness). *The main term kernel* $\frac{1}{2\pi}\log\left(\frac{t^2}{4\pi^2}\right)$ *matches the normalization throughout Section 5 and is identified with the Archimedean term by the calibration in Section 6.2 (Proposition 15). Therefore (33) expresses content independent of normalization.*

Corollary 9 (Distributional equality in small bandwidth). *Under* $\eta < \log 2$, μ_L *and* μ_{ξ} *agree as tempered distributions on* A_η*:*

$$\forall\, \phi \in A_\eta, \qquad \langle \mu_L - \mu_{\xi}, \phi \rangle = 0.$$

In particular, the same holds for the band-limited upper/lower approximations $\Phi^{\pm,(m)}_{T,\eta}$ *of Section 5.3 (evenized as* $\Psi^{\pm,(m)}_{T,\eta}$*), which are used in the sandwich of Section 5.4.*

Remark 37 (Boundary $\eta = \log 2$ and connection to 6.4). *In the boundary case* $\eta = \log 2$*, the values* $\widehat{\phi}(\pm\eta)$ *may give* endpoint contributions *from the prime sum. In this paper, assuming* $\widehat{\phi}^{(j)}(\pm\eta) = 0$ $(0 \le j \le m)$ *removes the boundary term (Lemma 22), and (33) holds in the limiting sense. See Section 6.4 for details.*

Summary: connection to next section

Thus, for small bandwidth $\eta < \log 2$, μ_L and μ_{ξ} have been shown to *coincide exactly*. In the next Section 6.2, we independently prove the *calibration proposition* for the Archimedean term ($\Gamma_{\mathbb{R}}$-term Fourier image = main term kernel), and then in Section 6.3 we present the return of the prime term (finite sum) and a difference representation in the large bandwidth case ($\eta \ge \log 2$).

6.2. Archimedean Calibration: Agreement between $\Gamma_{\mathbb{R}}$ *Term and Main Term Kernel [4,5,29,35]*

Position of this subsection (relation to overall strategy).

The key to the small bandwidth equivalence in Section 6.1 is to show that the *Archimedean term* (arising from the Γ-factor) appearing in the explicit formula *coincides exactly* (in the sense of the finite part) with the main term kernel used in Section 5.1–Section 5.2:

$$K_0(t) \;:=\; \frac{1}{2\pi}\log\Big(\frac{t^2}{4\pi^2}\Big).$$

In this section we establish this calibration as an independent proposition. From now on, the Fourier conventions and definition of the finite part follow Section 4.1 and Section 5.2 (Lemma 17).

Notation. $\Gamma_{\mathbb{R}}(s) := \pi^{-s/2}\Gamma(s/2)$ (Archimedean factor of the completed ζ), $\psi(z) := \frac{\Gamma'(z)}{\Gamma(z)}$ (digamma). For an even test $\phi \in A_\eta$, define the action of the Archimedean distribution by

$$\mathcal{A}[\phi] \;:=\; \frac{1}{2\pi}\int_{\mathbb{R}} \phi(t)\,\mathrm{Re}\Big(\psi\Big(\frac{1}{4}+\frac{it}{2}\Big)\Big)\,dt \;-\; \frac{\log\pi}{2\pi}\int_{\mathbb{R}}\phi(t)\,dt \tag{34}$$

(this matches the $\Gamma_{\mathbb{R}}$-term in the explicit formula; by evenness it suffices to take the real part).

Main proposition (calibration identity)

Proposition 15 (Archimedean calibration). *Let* $\phi \in A_\eta$ *be even. For any endpoint vanishing order* $m \ge 1$ *and cutoff width* $\delta \in (0,1]$*,*

$$\mathcal{A}[\phi] \;=\; \frac{1}{2\pi}\Big\langle \mathrm{fp}_{m,\delta}\log,\ \phi\Big\rangle \;=\; \frac{1}{2\pi}\int_{\mathbb{R}}\big(1-\eta_{m,\delta}(t)\big)\,\phi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt. \tag{35}$$

In particular, the finite part is unique (independent of δ *and* m*) within the scope of Lemma 17, and in the limit* $\delta \downarrow 0$ *we have* $\mathcal{A}[\phi] = \int \phi(t)K_0(t)\,dt$ *(in the distributional sense).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Outline of proof. *(i) Integral representation of the digamma and evenization.* For the real part $\operatorname{Re}\,\psi(\sigma+it)$ there is the classical integral representation

$$\operatorname{Re}\,\psi(\sigma+it) \;=\; \int_0^\infty \Big(\frac{e^{-x}}{x} - \frac{e^{-\sigma x}\cos(tx)}{1-e^{-x}}\Big)\,dx, \qquad \sigma>0.$$

Taking $\sigma=\frac14$, substituting into (34), convolving with the even test ϕ, and using Fubini (absolute integrability from $\widehat{\phi}\in C_c^\infty$) to interchange integrals, we obtain

$$\mathcal{A}[\phi] \;=\; \frac{1}{2\pi}\int_0^\infty \Big(\frac{e^{-x}}{x} - \frac{e^{-x/4}}{1-e^{-x}}\Big)\Big(\int_{\mathbb{R}}\phi(t)\,dt\Big)\,dx \;+\; \frac{1}{2\pi}\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\Big(\int_{\mathbb{R}}\phi(t)\,(1-\cos(tx))\,dt\Big)\,dx.$$

The first term cancels with $-\frac{\log\pi}{2\pi}\int\phi$ (since $\int_0^\infty(\frac{e^{-x}}{x}-\frac{e^{-x/4}}{1-e^{-x}})\,dx=\log\pi$), hence

$$\mathcal{A}[\phi] \;=\; \frac{1}{2\pi}\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\Big(\int_{\mathbb{R}}\phi(t)\,(1-\cos(tx))\,dt\Big)\,dx. \tag{36}$$

(ii) Application of Fourier–Paley–Wiener. Since ϕ is even with $\widehat{\phi}\in C_c^\infty([-\eta,\eta])$, $\int_{\mathbb{R}}\phi(t)\cos(tx)\,dt=\widehat{\phi}(x)$ (normalized by our convention). Thus (36) becomes

$$\mathcal{A}[\phi] \;=\; \frac{1}{2\pi}\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\,(\widehat{\phi}(0)-\widehat{\phi}(x))\,dx.$$

Near $x=0$, the singularity is $\frac{e^{-x/4}}{1-e^{-x}}=\frac1x+\frac12+O(x)$, while $\widehat{\phi}(0)-\widehat{\phi}(x)=\frac{x^2}{2}\widehat{\phi}''(0)+O(x^4)$ (by evenness), so the integral is absolutely convergent when interpreted as a *finite part* (see Lemma 17).

(iii) Identification of the finite part: recovery of log *kernel.* Using integration by parts and $1-\cos(tx)=2\sin^2(tx/2)$,

$$\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\,(1-\cos(tx))\,dx \;=\; \int_0^\infty\Big(\frac1x+r(x)\Big)\,(1-\cos(tx))\,dx,$$

where $r(x)\in L^1(0,\infty)$. The first part $\int_0^\infty x^{-1}(1-\cos(tx))dx=\frac{\pi}{2}|t|$ is standard, while the second is the Fourier transform of an L^1-kernel and is smooth and even. Thus for even tests ϕ,

$$\mathcal{A}[\phi] \;=\; \frac{1}{2\pi}\int_{\mathbb{R}}\phi(t)\,\Big(\pi|t| \;+\; \widetilde{R}(t)\Big)\,dt,$$

with $\widetilde{R}\in C^\infty$ even. Since $\frac{d}{dt}\big(\frac{1}{2\pi}\log(t^2)\big)=\frac{1}{\pi}\,\mathrm{pv}\,\frac1t$ and $\mathcal{F}\big(\mathrm{pv}\,\frac1t\big)=-i\pi\,\mathrm{sgn}$, the primitive of $\pi|t|$ is $\frac{1}{2\pi}\log(t^2)$ (as a distribution). The remainder $\widetilde{R}$ is C^∞ and even, so $\int\phi(t)\,\widetilde{R}(t)\,dt$ is a constant multiple of $\int\phi(t)\,dt$, which is exactly canceled by the $-\frac{\log\pi}{2\pi}\int\phi$ term in (34). Altogether, we obtain $\mathcal{A}[\phi]=\frac{1}{2\pi}\langle \mathrm{fp}_{m,\delta}\log,\,\phi\rangle$ (with independence of δ,m from Lemma 17, (23)). □

Remark 38 (Absorption of constants and uniqueness)**.** *The smooth remainder $\widetilde{R}$ in the above argument is limited to an even constant term (by finite Fourier support and evenness). This is canceled by $-\frac{\log\pi}{2\pi}\int\phi$ in* (34)*, and this calibration makes it* exactly *match the main term kernel K_0. Therefore* (35) *holds regardless of the choice of finite part (Lemma 17) and in the limit $\delta\downarrow 0$.*

Corollary 10 (Application to Section 6.1)**.** *For any $\phi\in A_\eta$ (even), $\mathcal{A}[\phi]=\int\phi(t)\,K_0(t)\,dt$ holds. Therefore, the main term expression in* (33) *of Section 6.1 is in complete agreement with the $\Gamma_{\mathbb{R}}$-term of the explicit formula.*

Summary: connection to next section

In this section we have shown that the Archimedean term agrees exactly, in the sense of the finite part, with the main term kernel $K_0(t)=\frac{1}{2\pi}\log(\frac{t^2}{4\pi^2})$. In the next Section 6.3, we present as a theorem that in the *large bandwidth* case $\eta\geq\log 2$, the prime term returns as a finite sum (difference representation).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

6.3. Wide-Band Version of the Explicit Formula: Finite Sum Representation of Prime Terms [1,3–5]

Position of this subsection (relation to overall strategy).

In Section 6.1 we showed that in the *small bandwidth* case $\eta < \log 2$, the equality $\mu_L = \mu_\xi$ holds. In this section we move to the *wide bandwidth* case $\eta \geq \log 2$, and formulate, using the Archimedean calibration in Section 6.2 (Proposition 15) and the finite part from Section 5.2 (Lemma 17), that the difference between them appears as a *finite sum of prime terms*. The *boundary terms* arising from the endpoints (bandwidth boundaries $\pm\eta$) will be estimated in Section 6.4 (Lemma 22).

Recollection of assumptions and notation

From now on we work with the even band-limited test class

$$A_\eta \;:=\; \{\, \phi \in \mathcal{S}(\mathbb{R}) \text{ even} :\; \widehat{\phi} \in C_c^\infty([-\eta,\eta]) \,\},$$

assuming if necessary the vanishing at the band edge $\widehat{\phi}^{(j)}(\pm\eta) = 0$ $(0 \leq j \leq m)$ (endpoint vanishing order m). The main term kernel is $K_0(t) = \frac{1}{2\pi}\log\!\left(\frac{t^2}{4\pi^2}\right)$ (cf. Proposition 15 in Section 6.2).

Main theorem (wide-band difference = finite sum of prime terms)

Theorem 9 (Wide-band explicit formula)**.** *Let $\eta \geq \log 2$ and $\phi \in A_\eta$ be even, with endpoint vanishing order $m \geq 1$. Then*

$$E_L[\phi] \;-\; E_\xi[\phi] \;=\; -\sum_{\substack{n\geq 2\\ \log n\leq \eta}} \Lambda(n)\left(\widehat{\phi}(\log n) + \widehat{\phi}(-\log n)\right) \;+\; \mathcal{B}_{\eta,m}[\phi], \tag{37}$$

where $\mathcal{B}_{\eta,m}$ is a boundary term *localized at the band edges $|\xi| = \eta$, and provided $\widehat{\phi}$ vanishes to order m at $\pm\eta$ (Lemma 22 in Section 6.4),*

$$|\mathcal{B}_{\eta,m}[\phi]| \;\leq\; C_{m,\eta} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1(\mathbb{R})} \tag{38}$$

with constants depending only on η, m and the constants in Section 4.2.

Sketch of proof. Applying Section 6.2 to the standard form of the explicit formula, we have for even tests ϕ,

$$E_\xi[\phi] \;=\; \int_{\mathbb{R}} \phi(t)\, K_0(t)\, dt \;-\; \sum_{n\geq 2} \Lambda(n)\left(\widehat{\phi}(\log n) + \widehat{\phi}(-\log n)\right) \;+\; \mathcal{E}_{\text{edge}}[\phi].$$

Here $\mathcal{E}_{\text{edge}}$ is the (finite part) boundary contribution arising from the band edges and low-frequency regularization. On the other hand, from Section 5.1–Section 5.2 we have $E_L[\phi] = \int \phi K_0 +$ (finite-part adjustment of the same form), and on the μ_L side there is no prime term. Subtracting the two expressions and using $\operatorname{supp}\widehat{\phi} \subset [-\eta,\eta]$ to eliminate n with $\widehat{\phi}(\pm\log n) = 0$ yields (37). The bound (38) for the boundary term is given in Lemma 22 of Section 6.4. □

Remark 39 (Significance of the finiteness of the sum)**.** *Since* $\operatorname{supp}\widehat{\phi} \subset [-\eta,\eta]$*, only* $\{n \geq 2 : \log n \leq \eta\}$ *contribute, so the prime term on the right-hand side is a* finite sum*. Thus analytically, the handling of the prime term is controlled solely by η.*

Upper bound for the prime term (basic form)

Lemma 21 (Upper bound for the finite sum of prime terms)**.** *Let $\eta \geq \log 2$ and $\phi \in A_\eta$ be even. Then*

$$\begin{aligned}\left|\sum_{\substack{n\geq 2\\ \log n\leq \eta}} \Lambda(n)\left(\widehat{\phi}(\log n) + \widehat{\phi}(-\log n)\right)\right| \;&\leq\; 2\,\|\widehat{\phi}\|_{L^\infty([-\eta,\eta])} \sum_{n\leq e^\eta} \Lambda(n) \\ &\leq\; 2\,\|\phi\|_{L^1(\mathbb{R})} \cdot \eta\, e^\eta. \end{aligned} \tag{39}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. The first inequality follows from the triangle inequality and evenness. The basic Fourier bound $\|\widehat{\phi}\|_\infty \le \|\phi\|_{L^1}$ and the trivial bound $\sum_{n\le x}\Lambda(n) \le \sum_{n\le x}\log n \le x\log x$ (with $x = e^\eta$) yield (39). □

Remark 40 (Room for improvement of the bound)**.** *Sharper bounds (e.g. of type $\ll e^\eta$ or $\ll e^\eta/\eta$) can be obtained using known estimates from analytic number theory, but for our purposes the rough form (39) suffices. Also, assuming higher order vanishing at the band edge, $\|\widehat{\phi}\|_\infty$ can be controlled by $\sum_{j\le m}\|\phi^{(j)}\|_{L^1}$ via a Bernstein-type bound (see Lemma 18 in Section 5.2).*

Summary: connection to next section (boundary handling)

By Theorem 9, in the wide-band case we have

$$\mu_L - \mu_\xi \;=\; -\sum_{\log n\le\eta}\Lambda(n)\,(\delta_{\log n}+\delta_{-\log n}) \;\text{ (inverse Fourier image)} \;\; + \text{ boundary term } \mathcal{B}_{\eta,m}.$$

The prime term is a finite "point spectrum" contribution whose size is bounded in terms of η and $\|\widehat{\phi}\|_\infty$ (or $\sum_{j\le m}\|\phi^{(j)}\|_{L^1}$) by Lemma 21. The remaining term $\mathcal{B}_{\eta,m}$ is estimated in Section 6.4 (Lemma 22) using endpoint vanishing and the finite part.

6.4. Treatment of the Band Edge: $\eta = \log 2$ and Endpoint Vanishing [7,25,29]

Position of this subsection (relation to overall strategy).

In Theorem 9 of Section 6.3 we presented, for $\eta \ge \log 2$,

$$E_L[\phi] - E_\xi[\phi] \;=\; -\sum_{\log n\le\eta}\Lambda(n)\,(\widehat{\phi}(\log n)+\widehat{\phi}(-\log n)) \;+\; \mathcal{B}_{\eta,m}[\phi].$$

The aim of this section is to give a quantitative estimate of the boundary term $\mathcal{B}_{\eta,m}$, localized at the *band edges* $|\xi| = \eta$, in terms of the order m of endpoint vanishing ($\widehat{\phi}^{(j)}(\pm\eta) = 0$), and to connect *continuously* to the small-band equivalence (Theorem 8) in the limit $\eta \downarrow \log 2$. Justification of finite parts, cyclic products, and exchange of limits relies on Section 5.2 (Lemma 17, Theorem 6) and Section 4.3 (Proposition 11).

Edge localization decomposition and technical preparation

Definition 8 (Edge localization decomposition)**.** *Take a small $\kappa \in (0,\eta)$, and choose $\chi_{\text{in}}, \chi_\pm \in C_c^\infty(\mathbb{R})$ such that*

$$\chi_{\text{in}} \equiv 1 \text{ on } [-\eta+\kappa, \eta-\kappa], \qquad \operatorname{supp}\chi_\pm \subset [\pm\eta-\kappa, \pm\eta+\kappa], \qquad \chi_{\text{in}}+\chi_+ +\chi_- \equiv 1.$$

For $\phi \in A_\eta$ decompose in the Fourier side as $\widehat{\phi} = \widehat{\phi}_{\text{in}} + \widehat{\phi}_{\text{edge}}$, where $\widehat{\phi}_{\text{in}} := \chi_{\text{in}}\widehat{\phi}$, $\widehat{\phi}_{\text{edge}} := \chi_+\widehat{\phi} + \chi_-\widehat{\phi}$. The corresponding time-side functions are denoted $\phi_{\text{in}}, \phi_{\text{edge}}$.

Remark 41 (Effect of endpoint vanishing (Taylor remainder form))**.** *If $\widehat{\phi}^{(j)}(\pm\eta) = 0$ for $0 \le j \le m$, then near each endpoint*

$$\chi_\pm(\xi)\,\widehat{\phi}(\xi) \;=\; (\xi\mp\eta)^{m+1}\,h_\pm(\xi), \qquad h_\pm \in C_c^\infty([\pm\eta-\kappa, \pm\eta+\kappa]),$$

(Taylor remainder). In this case $\phi_{\text{edge}}(t) = i^{m+1}\,\partial_t^{m+1}\big(e^{\pm i\eta t}h_\pm^\vee(t)\big)$. Hence the L^1 and derivative norms of ϕ_{edge} are controlled by $\|h_\pm^\vee\|_{W^{m+1,1}}$, which by Bernstein/Paley–Wiener type estimates (Lemma 18) can be bounded in terms of $\sum_{j\le m}\|\phi^{(j)}\|_{L^1}$ with constants depending only on m, η, κ.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Main result: bound on the boundary term

Lemma 22 (Estimate for the band edge term). *Let $\eta \geq \log 2$ and $\phi \in A_\eta$ be even, with $\widehat{\phi}^{(j)}(\pm\eta) = 0$ for $0 \leq j \leq m$. Then the boundary term $\mathcal{B}_{\eta,m}[\phi]$ in Theorem 9 satisfies*

$$|\mathcal{B}_{\eta,m}[\phi]| \;\leq\; C_{m,\eta,\kappa} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1(\mathbb{R})}, \tag{40}$$

where the constant depends only on m, η and the partition width κ, and can be absorbed into the constants from Section 4.2.

Sketch of proof. By Definition 8, decompose $E_L[\phi] - E_{\tilde{\xi}}[\phi]$ into interior and edge components. The interior component has $\operatorname{supp} \widehat{\phi}_{\text{in}} \subset [-\eta+\kappa, \eta-\kappa]$, so by the same computation as in Section 6.1 (no prime term) it vanishes. Thus the difference comes only from the edge component, which coincides with $\mathcal{B}_{\eta,m}[\phi]$.

From the preceding remark, ϕ_{edge} can be written as $i^{m+1}\partial_t^{m+1}(e^{\pm i\eta t} h_\pm^\vee)$. Under the justification for $\det_2$ and cyclic products (see Section 4.3, Proposition 11), the boundary term is given by the difference of traces with ϕ_{edge} inserted into the kernel cyclic product. Using the Kato–Seiler–Simon type inequality (Section 4.2, Proposition 9) and off-diagonal decay (Theorem 4) we obtain $|\mathcal{B}_{\eta,m}[\phi]| \lesssim \sum_{j\leq m} \|\phi_{\text{edge}}^{(j)}\|_{L^1}$. Finally, applying the expression for ϕ_{edge} and Lemma 18 gives $\|\phi_{\text{edge}}^{(j)}\|_{L^1} \leq C_{m,\eta,\kappa} \sum_{\ell\leq m} \|\phi^{(\ell)}\|_{L^1}$, yielding (40). □

Remark 42 (Dependence of constants and practical choice). *The constant $C_{m,\eta,\kappa}$ decreases monotonically with the partition width κ, and may worsen as $\kappa \downarrow 0$. In practice it is sufficient to fix $\kappa = \eta/4$, allowing uniform choice of constants even in the limit $\eta \to \log 2$.*

Limit $\eta \downarrow \log 2$ and connection to the small band

Proposition 16 (Continuous connection from band edge to small band). *Let $\{\eta_\nu\}_{\nu\geq 1}$ satisfy $\eta_\nu \downarrow \log 2$. For each ν, let $\phi_\nu \in A_{\eta_\nu}$ be even with $\widehat{\phi_\nu}^{(j)}(\pm\eta_\nu) = 0$ for $0 \leq j \leq m$, and assume $\sum_{j=0}^m \|\phi_\nu^{(j)}\|_{L^1} \leq M$ holds uniformly in ν. Then*

$$\lim_{\nu\to\infty} \mathcal{B}_{\eta_\nu,m}[\phi_\nu] \;=\; 0. \tag{41}$$

Consequently, Theorem 9 connects continuously in the limit $\eta = \log 2$ to Theorem 8 (small-band equivalence).

Proof. From Lemma 22 and the assumption $\sum_{j\leq m} \|\phi_\nu^{(j)}\|_{L^1} \leq M$ we have $|\mathcal{B}_{\eta_\nu,m}[\phi_\nu]| \leq C_{m,\eta_\nu,\kappa} M$. Fixing κ as a constant fraction of η_ν ensures $C_{m,\eta_\nu,\kappa}$ is uniformly bounded in ν. On the other hand, due to endpoint vanishing, the endpoint-neighborhood contribution of $\widehat{\phi_\nu}$ has a Taylor remainder factor $(\xi \mp \eta_\nu)^{m+1}$ and support length $\sim \kappa$ fixed, so the $W^{m+1,1}$ norm of $\phi_{\nu,\text{edge}}$ can be made uniformly small in ν (continuity under *shift* of the endpoint, not shrinking of the partition width). Propagating this as in the previous proof yields (41). □

Convention. In the small-band case we always use the strict form $\eta < \log 2$, and treat the boundary $\eta = \log 2$ via endpoint vanishing (Appendix A.16).

Remark 43 (Case without endpoint vanishing). *If $\widehat{\phi}(\pm\eta) \neq 0$, then the boundary term will in general not vanish, and an endpoint contribution from the prime 2 may remain in the limit $\eta = \log 2$. In the framework of this paper, we adopt the assumption of endpoint vanishing (at least $m \geq 1$) in order to have continuous connection to the small band.*

Summary: connection to next section

Lemma 22 shows that the *boundary term* in the wide-band difference can be controlled solely in terms of the order m of endpoint vanishing and the bandwidth η. Proposition 16 ensures that as

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

$\eta \downarrow \log 2$, there is a continuous connection to the small-band equivalence (Theorem 8). In the next Section 6.5, we summarize the conclusions of this chapter and prepare the bridge to §7 (next chapter; $\det_2(I \pm z\varphi(L))$ and the Cayley transform).

6.5. Summary and Bridge to §7

Conclusions of this chapter.

In this chapter, we established the *equivalence* and *structure of the difference* between the operator-side distribution μ_L and the completed zeta-side distribution μ_{ξ} on the space of even, band-limited test functions $A_\eta = \{\phi \in \mathcal{S}(\mathbb{R}) \text{ even} : \widehat{\phi} \in C_c^\infty([-\eta,\eta])\}$. Throughout, we consistently used the normalization

$$K_0(t) := \frac{1}{2\pi}\log\Big(\frac{t^2}{4\pi^2}\Big)$$

(see Section 6.2, Proposition 15).

(S1) **Small-band equivalence ($\eta < \log 2$).** By Theorem 8,

$$\forall\, \phi \in A_\eta, \qquad E_L[\phi] \;=\; E_{\xi}[\phi] \;=\; \int_{\mathbb{R}} \phi(t)\, K_0(t)\, dt.$$

That is, μ_L and μ_{ξ} act as the *same* tempered distribution on A_η.

(S2) **Archimedean calibration.** By Proposition 15, the $\Gamma_{\mathbb{R}}$-term in the explicit formula coincides exactly with K_0 in the finite-part sense. Thus the handling of the main term is in complete agreement with Section 5.

(S3) **Wide band ($\eta \geq \log 2$) difference = prime term + boundary term.** From Theorem 9 and Lemma 21,

$$E_L[\phi] - E_{\xi}[\phi] \;=\; -\sum_{\log n \leq \eta} \Lambda(n)\,\big(\widehat{\phi}(\log n) + \widehat{\phi}(-\log n)\big) \;+\; \mathcal{B}_{\eta,m}[\phi],$$

where $\mathcal{B}_{\eta,m}$ is the boundary term localized at the band edge, and by Lemma 22 $|\mathcal{B}_{\eta,m}[\phi]| \leq C_{m,\eta} \sum_{j=0}^{m} \|\phi^{(j)}\|_{L^1}$. The prime term is a *finite sum* due to $\operatorname{supp}\widehat{\phi} \subset [-\eta,\eta]$ (with $\#\{n : \log n \leq \eta\} \leq e^{\eta}$).

(S4) **Continuity at the boundary ($\eta \downarrow \log 2$).** By Proposition 16, under endpoint vanishing $\widehat{\phi}^{(j)}(\pm\eta) = 0$ for $0 \leq j \leq m$, we have $\mathcal{B}_{\eta,m}[\phi] \to 0$. Thus the wide-band difference formula connects *continuously* to the small-band equivalence (Theorem 8).

Operator-side vs. zeta-side correspondence table (summary).

- Main term: $E_L[\phi]$ and $E_{\xi}[\phi]$ are both calibrated by $\int \phi K_0$ (Proposition 15).
- Prime term: Contributes to the wide-band difference *only* through $\widehat{\phi}(\pm\log n)$ (Theorem 9).
- Boundary term: Localized contribution from the band edge $|\xi| = \eta$ can be controlled by the order m of endpoint vanishing, and vanishes as $\eta \downarrow \log 2$ (Lemma 22, Proposition 16).

Technical support from Section 4–Section 5.

The localized trace-class property (Proposition 9), off-diagonal decay (Theorem 4), and localized cyclic products (Proposition 11), together with the finite part (Lemma 17), underpin the justification of limit exchanges and equalities in this chapter. The Tauberian-type sandwich of Section 5.4 is consistent with the small-band equivalence in this chapter, and matches the normalization of the main term (K_0).

Bridge to §7: $\det_2$ and the Cayley transform

The distribution-level equalities and differences obtained in this chapter will be lifted, in the next §7, via the *regularized Fredholm determinant* and the *Cayley transform*, to analytic-function-theoretic $\det_2$-type generating functions:

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

- Operator side: For $K = \varphi(L)$, the logarithmic derivative of $\det_2(I + zK)$ is $\sum_{m\geq 2}(-1)^{m-1} z^{m-1} \operatorname{Tr}(K^m)$ (Proposition 10).
- Distribution side: $\operatorname{Tr}(K^m)$ is a localized cyclic product = kernel cyclic product (Proposition 11).
- Zeta side: In the small band, $E_L = E_{\xi}$; in the wide band, the prime term (finite sum) carries the difference (Theorem 9).

This allows, through coefficient identification in $\log\det_2$, a comparison of the operator-side analytic data and the completed zeta-side explicit formula within the same framework (see §7 for details).

Summary: work plan for the next chapter

In the next chapter §7:

(N1) Use the Cayley transform to move from the self-adjoint L to a unitary on the unit circle, fixing the choice of $\varphi(L)$;

(N2) Establish the analytic properties of $\det_2(I \pm z\varphi(L))$ (branches, zero set, Jensen-type formulas);

(N3) Reflect the equivalence/difference from Section 6 (finite-sum prime term) in the coefficients of $\log\det_2$, matching the generating functions on both sides.

This will lift the distribution-level equalities to the analytic function level, connecting to the core claims in the subsequent sections.

7. Regularized Fredholm Determinant and Cayley Transform

7.1. Setup and Conventions: Unification via $\Phi(L)$ [6,14,24]

Position of this subsection (relation to the overall strategy).

In this chapter, for the self-adjoint operator L (as set up in Section 4.1), we relegate the Cayley transform to a *supplementary* role and adopt the functional calculus $\Phi(L)$ as the main route. From now on we write

$$\mathbf{K} := \Phi(L)$$

and first establish sufficient conditions for $\mathbf{K}$ to be Hilbert–Schmidt class ($\mathcal{S}_2$). Under this assumption, we discuss the entire-function property and coefficient expansion of $\det_2(I + z\mathbf{K})$ in Section 7.2, and in Section 7.3 connect $\operatorname{Tr}(\mathbf{K}^r)$ to the distributional equivalence of Section 6. Justification for exchanging limits, localization, and cyclic products can be found in Section 4.2 (Proposition 9, Theorem 4) and Section 4.3 (Proposition 11).

Notation and conventions.

- *Band-limited tests* from Section 5–Section 6 are denoted by ϕ, while in this chapter the *functional calculus kernel* is denoted by Φ.
- $\Phi : \mathbb{R} \to \mathbb{R}$ is assumed to be *even* and *real*-valued and smooth; when necessary, we use $\Phi \in \mathcal{S}(\mathbb{R})$ or convolution with a band-limited approximation.
- The symmetric measure corresponding to the eigenvalue sequence $\{\pm\gamma_k\}_{k\geq 1}$ is $\mu_L := \sum_{k\geq 1}(\delta_{\gamma_k} + \delta_{-\gamma_k})$ (see Section 5.1).

Sufficient condition for Hilbert–Schmidt and its derivation

First, we identify the Hilbert–Schmidt norm of $\mathbf{K} = \Phi(L)$ via the spectral measure as $\|\mathbf{K}\|_{\mathcal{S}_2}^2 = \sum_{k\geq 1} |\Phi(\gamma_k)|^2$ (Lemma below). Using the Weyl law (Section 3.4, Theorem 7) stating that $dN_{\mathrm{eig}}(t) \sim \frac{1}{2\pi}\log t\, dt$ (main term), we obtain an estimate $\sum |\Phi(\gamma_k)|^2 \approx \int_1^{\infty} |\Phi(t)|^2 \log t\, dt$.

Lemma 23 (Spectral representation and $\mathcal{S}_2$ norm)**.** *For the functional calculus of a self-adjoint L, $\mathbf{K} = \Phi(L)$ satisfies*

$$\|\mathbf{K}\|_{\mathcal{S}_2}^2 = \sum_{\gamma_k > 0} \left(|\Phi(\gamma_k)|^2 + |\Phi(-\gamma_k)|^2 \right) = \int_{\mathbb{R}} |\Phi(t)|^2 \, d\mu_L(t).$$

In particular, if Φ is even then $\|\mathbf{K}\|_{\mathcal{S}_2}^2 = 2\sum_{\gamma_k > 0} |\Phi(\gamma_k)|^2$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. By the spectral theorem, $L = \int t\, dE_L(t)$ and $\Phi(L) = \int \Phi(t)\, dE_L(t)$. The $\mathcal{S}_2$ norm is the square root of $\mathrm{Tr}(\mathbf{K}^*\mathbf{K})$, and orthogonality in the eigen-decomposition yields the claim. □

Proposition 17 (Sufficient conditions for $\Phi(L) \in \mathcal{S}_2$)**.** *Let $\Phi : \mathbb{R} \to \mathbb{R}$ be even, real, and measurable. If any of the following holds, then $\Phi(L) \in \mathcal{S}_2$:*

(i) $\int_0^1 |\Phi(t)|^2\, dt < \infty$ *and* $\int_1^\infty |\Phi(t)|^2 \log(2+t)\, dt < \infty$;

(ii) More strongly, $\Phi \in \mathcal{S}(\mathbb{R})$;

(iii) Or Φ is band-limited with $\widehat{\Phi} \in C_c^\infty([-\eta, \eta])$.

In this case,

$$\|\Phi(L)\|_{\mathcal{S}_2}^2 \;\le\; C\Big(\int_0^1 |\Phi(t)|^2\, dt \;+\; \int_1^\infty |\Phi(t)|^2 \log(2+t)\, dt\Big), \tag{42}$$

where $C > 0$ depends only on the constants in the Weyl law of Section 3.4 (Theorem 7).

Sketch of proof. From Lemma 23, $\|\Phi(L)\|_{\mathcal{S}_2}^2 = \int |\Phi(t)|^2\, d\mu_L(t) = \int_0^\infty |\Phi(t)|^2\, dN_{\mathrm{eig}}(t)$. By integration by parts and $N_{\mathrm{eig}}(t) = \frac{t}{2\pi}\log\frac{t}{4\pi^2} - \frac{t}{2\pi} + O(\log t)$ (Theorem 7),

$$\int_1^\infty |\Phi(t)|^2\, dN_{\mathrm{eig}}(t) \;\ll\; \int_1^\infty |\Phi(t)|^2 \log t\, dt \;+\; \int_1^\infty t\,\big|(|\Phi|^2)'(t)\big|\,\frac{dt}{t},$$

and the latter can be absorbed into $\int |\Phi|^2 + \int |\Phi'|^2$ using $|(|\Phi|^2)'| \le 2|\Phi||\Phi'|$ and Young/Hardy-type estimates (trivial if $\Phi \in \mathcal{S}$). In the band-limited case, the Paley–Wiener-type inequality (Lemma 18) yields $\||\Phi|'\|_{L^2} \lesssim \eta\, \|\Phi\|_{L^2}$, giving (42). □

Remark 44 (On evenness and the Cayley transform)**.** *The evenness assumption aligns with the symmetric spectrum $\{\pm\gamma_k\}$ and is effective in simplifying multiplicity counting in trace-class arguments (Section 4.3). Although one could equivalently work with $\Psi(U)$ via the Cayley transform $U = (L - iI)(L + iI)^{-1}$, we here unify the discussion in terms of $\Phi(L)$ (technical justification in Section 4.2).*

Summary: connection to the next section

In this subsection, we fixed the convention of taking $\mathbf{K} = \Phi(L)$ as the main object, and obtained sufficient conditions (Proposition 17) to ensure $\mathbf{K} \in \mathcal{S}_2$. This sets the stage for Section 7.2, where we establish the *entire-function property*, *coefficient expansion*, and *zero structure* (Theorem 10) of the *regularized Fredholm determinant* $\det_2(I + z\mathbf{K})$, and in Section 7.3 transport $\mathrm{Tr}(\mathbf{K}^r)$ to the distributional equivalence/difference of Section 6.

7.2. Regularized Fredholm Determinant: Entire Functionality and Expansion [6,26,36]

Position of this subsection (relation to the overall strategy).

For $\mathbf{K} = \Phi(L) \in \mathcal{S}_2$ fixed in Section 7.1 (Proposition 17), we establish the *entire-function property*, *zero structure*, *coefficient expansion* (starting index $r \ge 2$), and *growth estimates* for the regularized Fredholm determinant

$$\det_2(I + z\mathbf{K}).$$

This will form the basis for the coefficient identification in Section 7.3.

Definition and basic properties

Definition 9 (Regularized determinant (Hilbert–Schmidt class))**.** *Let $\{\lambda_j\}_{j\ge1}$ (counted with algebraic multiplicity) be the nonzero eigenvalues of $\mathbf{K} \in \mathcal{S}_2$. The* regularized Fredholm determinant $\det_2(I + z\mathbf{K})$ *is defined by*

$$\det_2(I + z\mathbf{K}) \;:=\; \prod_{j\ge1}\Big\{(1 + z\lambda_j)\, e^{-z\lambda_j}\Big\}, \tag{43}$$

a Weierstrass-type regularization; the Hilbert–Schmidt property $\sum_j |\lambda_j|^2 < \infty$ guarantees convergence.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 45 (Relation to trace class). *If* $\mathbf{K} \in \mathcal{S}_1$ *then* $\det_2(I + z\mathbf{K}) = \det(I + z\mathbf{K})\, e^{-z \operatorname{Tr}\mathbf{K}}$, *so the linear term is removed by the normalization. Here we focus on* $\mathbf{K} \in \mathcal{S}_2$ *and do not assume* $\operatorname{Tr}\mathbf{K}$ *exists.*

Entire function property, zero structure, and coefficient expansion

Theorem 10 (Entire function property and expansion of $\det_2(I + z\mathbf{K})$). *Let* $\mathbf{K} \in \mathcal{S}_2$. *Then:*

(a) $\det_2(I + z\mathbf{K})$ *is an* entire function *on* $z \in \mathcal{C}$.

(b) Its zeros occur precisely at $z = -\lambda_j^{-1}$ *for* $\lambda_j \neq 0$, *with multiplicity equal to the algebraic multiplicity of the eigenvalue.*

(c) The power series expansion near the origin is

$$\log \det_2(I + z\mathbf{K}) \;=\; \sum_{r\geq 2} \frac{(-1)^{r-1}}{r}\, z^r \,\operatorname{Tr}(\mathbf{K}^r), \tag{44}$$

with derivative form

$$\frac{d}{dz} \log \det_2(I + z\mathbf{K}) \;=\; \sum_{r\geq 2} (-1)^{r-1}\, z^{r-1} \,\operatorname{Tr}(\mathbf{K}^r). \tag{45}$$

All coefficients are finite ($\mathbf{K}^2 \in \mathcal{S}_1$ *and* $\mathbf{K}$ *is bounded), and (44) converges as a holomorphic power series around the origin, coinciding with the* entire function *given by analytic continuation.*

Sketch of proof. By eigenvalue expansion, (43) is a Weierstrass regularized product; for Hilbert–Schmidt class it converges over the whole plane, defining an entire function. (b) follows immediately from the zero locations of the factors. For (c), apply $\log\big((1+w)e^{-w}\big) = \sum_{r\geq 2}(-1)^{r-1} w^r / r$ to each eigenvalue $w = z\lambda_j$, taking $|z|$ sufficiently small to permit termwise summation. The coefficients $\operatorname{Tr}(\mathbf{K}^r)$ are finite for $r \geq 2$ since $\mathbf{K}^2 \in \mathcal{S}_1$ and $\mathbf{K}$ is bounded. Analyticity then extends the expansion to all z. □

Remark 46 (Significance of starting index $r \geq 2$). *Because (44) starts at* $r = 2$, *there is no need for* $\operatorname{Tr}\mathbf{K}$. *This is the core of the normalization in* $\det_2$. *Thus, in coefficient identification (Section 7.3) we only deal with* $\operatorname{Tr}(\mathbf{K}^r)$ *for* $r \geq 2$.

Growth estimate and preparation for Jensen/Carleman application

Proposition 18 (Quadratic bound on growth; order ≤ 2). *Let* $\mathbf{K} \in \mathcal{S}_2$. *There exist absolute constants* $C_1, C_2 > 0$ *such that for any* $z \in \mathcal{C}$,

$$\log\big|\det_2(I + z\mathbf{K})\big| \;\leq\; C_1\, |z|^2\, \|\mathbf{K}\|_{\mathcal{S}_2}^2 \;+\; C_2\, \log\big(1 + |z|\, \|\mathbf{K}\|\big). \tag{46}$$

In particular, $\det_2(I + z\mathbf{K})$ *is an* entire function of order ≤ 2, *and Jensen (or Carleman) type zero-counting estimates apply.*

Sketch of proof. Use the product representation (43) and a splitting argument. For any z, let $J := \{j : |z\lambda_j| \geq \frac{1}{2}\}$ (finite). The finite "head" $\prod_{j\in J}$ can be crudely bounded by $\ll \exp\big(C\,|z|^2 \sum_{j\in J} |\lambda_j|^2\big)$. For the "tail" $\prod_{j\notin J}$, apply $|\log(1+w) - w| \leq |w|^2$ for $|w| \leq \frac{1}{2}$ to each $w = z\lambda_j$, giving

$$\log\Big|\prod_{j\notin J} (1 + z\lambda_j) e^{-z\lambda_j}\Big| \;\leq\; \sum_{j\notin J} |z\lambda_j|^2 \;\leq\; |z|^2\, \|\mathbf{K}\|_{\mathcal{S}_2}^2.$$

Finally, absorb the finite head contribution using the crude bound with $\|\mathbf{K}\|$ to obtain (46). □

Corollary 11 (Rough upper bound on zero count). *Let* $n(r)$ *be the total number (counted with multiplicity) of zeros in* $|z| \leq r$. *From Jensen's formula and (46),*

$$n(r) \;\ll\; r^2\, \|\mathbf{K}\|_{\mathcal{S}_2}^2 \;+\; \log\big(1 + r\, \|\mathbf{K}\|\big),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

with an absolute implied constant.

Remark 47 (Self-adjoint case)**.** *In our setting,* $\mathbf{K} = \Phi(L)$ *with* Φ *even and real, so* $\mathbf{K}$ *is self-adjoint. Thus* $\lambda_j \in \mathbb{R}$*, and the zeros* $-\lambda_j^{-1}$ *lie only on the real axis (for* $\lambda_j \neq 0$*).*

Summary: connection to the next section

In this subsection we established the entire-function property of $\det_2(I + z\mathbf{K})$ (Theorem 10), the coefficient expansion (44) starting at $r \geq 2$, and the growth estimate of order ≤ 2 (Proposition 18). This enables us, in the next Section 7.3, to directly transfer $\mathrm{Tr}(\mathbf{K}^r)$ to the distributional equivalence / wide-band difference of Section 6, identifying the coefficients in (44) on both the operator and zeta sides.

7.3. Coefficient Identification: $\mathrm{Tr}(\Phi(L)^r)$ *and Transfer of Distributional Equivalence [6,14,29]*

Position of this subsection (relation to the overall strategy).

For the expansion in Section 7.2

$$\log \det_2(I + z\Phi(L)) \;=\; \sum_{r\geq 2} \frac{(-1)^{r-1}}{r}\, z^r \;\mathrm{Tr}\big(\Phi(L)^r\big) \tag{47}$$

(eq. (44) in Theorem 10), we directly *transfer* the coefficients $\mathrm{Tr}(\Phi(L)^r)$ to the *distributional equivalence* between μ_L and μ_{ξ} established in Section 6. Hereafter, the Fourier convention follows Section 6, $\widehat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-it\xi}\, dt$.

Expressing the trace as a distributional moment

Definition 10 (Inverse Fourier transform of $\boldsymbol{\Phi}$ and $\boldsymbol{r}$-fold convolution)**.** *Let* $\check{\Phi}(t) := \mathbb{F}^{-1}_{\lambda\to t}\Phi(\lambda)$ *(with this convention* $\widehat{\check{\Phi}} = \Phi$*). For an integer* $r \geq 2$*, denote by* $\check{\Phi}^{(*r)}$ *the r-fold time convolution.*

Lemma 24 (Trace as a distributional moment)**.** *For a self-adjoint L and even, real* Φ*, for any integer* $r \geq 2$*,*

$$\mathrm{Tr}\big(\Phi(L)^r\big) \;=\; \sum_{k\geq 1}\Big(\Phi(\gamma_k)^r + \Phi(-\gamma_k)^r\Big) \;=\; \big\langle \mu_L,\, \Phi(\cdot)^r \big\rangle. \tag{48}$$

Furthermore, since $\Phi(\lambda)^r = \widehat{\check{\Phi}^{(*r)}}(\lambda)$*,*

$$\mathrm{Tr}\big(\Phi(L)^r\big) \;=\; \big\langle \mu_L,\, \widehat{\check{\Phi}^{(*r)}} \big\rangle. \tag{49}$$

Proof. By the spectral theorem, the eigenvalues of $\Phi(L)$ are $\{\Phi(\pm\gamma_k)\}$ (with multiplicity). This gives the first equality by definition. Next, $\Phi(\lambda) = \widehat{\check{\Phi}}(\lambda)$ implies $\Phi(\lambda)^r = \widehat{\check{\Phi}^{(*r)}}(\lambda)$ (convolution theorem for the Fourier transform). Substituting $\lambda = \pm\gamma_k$ and summing yields (49). □

Remark 48 (Alternative proof via localized cyclic product)**.** *Write* $\Phi(L) = \int \check{\Phi}(t)\, e^{-itL}\, dt$*, so that* $\Phi(L)^r = \int \check{\Phi}^{(*r)}(t)\, e^{-itL}\, dt$*. By Proposition 11 (localized cyclic product),* $\mathrm{Tr}(\Phi(L)^r) = \int \check{\Phi}^{(*r)}(t)\; \mathrm{Tr}\big(e^{-itL}\big)\, dt$*. Since* $\mathrm{Tr}\big(e^{-itL}\big) = \langle \mu_L, e^{-it(\cdot)}\rangle$*, this agrees with (49).*

Small band: complete coefficient equality

Proposition 19 (Coefficient identification in the small band)**.** *Let* $\check{\Phi} \in C_c^\infty([-\eta_0, \eta_0])$ *be even and real, and fix an integer* $r \geq 2$*. Set* $\eta_r := r\,\eta_0$*. If* $\eta_r < \log 2$ *and (if necessary) endpoint vanishing* $\big(\widehat{\Phi^r}\big)^{(j)}(\pm\eta_r) = 0$ *for* $0 \leq j \leq m$ *holds, then*

$$\mathrm{Tr}\big(\Phi(L)^r\big) \;=\; \big\langle \mu_L,\, \Phi^r \big\rangle \;=\; \big\langle \mu_{\xi},\, \Phi^r \big\rangle \;=\; \int_{\mathbb{R}} \Phi(t)^r\, K_0(t)\, dt. \tag{50}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. By Lemma 24, $\mathrm{Tr}(\Phi(L)^r) = \langle \mu_L, \Phi^r \rangle$. The Fourier transform of Φ^r is $\check{\Phi}^{(*r)}$, whose support lies in $[-\eta_r, \eta_r]$. Thus $\Phi^r \in A_{\eta_r}$. By Theorem 8 (small-band equivalence), $\langle \mu_L, \Phi^r \rangle = \langle \mu_\xi, \Phi^r \rangle = \int \Phi^r K_0$. □

Wide band: coefficient difference = finite prime sum + boundary term

Theorem 11 (Coefficient difference formula in the wide band)**.** *Under the assumptions of Proposition 19, let $\eta_r \geq \log 2$ (assume endpoint vanishing order $m \geq 1$). Then*

$$\mathrm{Tr}\big(\Phi(L)^r\big) \;-\; \big\langle \mu_\xi,\, \Phi^r \big\rangle \;=\; -\sum_{\substack{n\geq 2\\ \log n \leq \eta_r}} \Lambda(n)\left(\check{\Phi}^{(*r)}(\log n) + \check{\Phi}^{(*r)}(-\log n)\right) \;+\; \mathcal{B}_{\eta_r,m}\big[\Phi^r\big], \tag{51}$$

where $\mathcal{B}_{\eta_r,m}$ is the boundary term localized at the band edge, and by Lemma 22,

$$\big|\mathcal{B}_{\eta_r,m}[\Phi^r]\big| \;\leq\; C_{m,\eta_r} \sum_{j=0}^{m} \big\|\partial_t^j(\Phi^r)\big\|_{L^1(\mathbb{R})}. \tag{52}$$

Proof. From Lemma 24, $\mathrm{Tr}(\Phi(L)^r) = \langle \mu_L, \Phi^r \rangle$. Apply Theorem 9 to $\phi = \Phi^r \in A_{\eta_r}$ and use $\widehat{\phi} = \check{\Phi}^{(*r)}$ to obtain (51). Boundary estimation is by Lemma 22. □

Lemma 25 (Rough bound for the finite prime sum)**.** *In the situation of Theorem 11,*

$$\left|\sum_{\log n \leq \eta_r} \Lambda(n)\left(\check{\Phi}^{(*r)}(\log n) + \check{\Phi}^{(*r)}(-\log n)\right)\right| \;\leq\; 2\,\|\Phi^r\|_{L^1(\mathbb{R})}\,\eta_r\, e^{\eta_r}. \tag{53}$$

Proof. Apply Lemma 21 to $\phi = \Phi^r$, using $\|\widehat{\phi}\|_\infty = \|\check{\Phi}^{(*r)}\|_\infty \leq \|\phi\|_{L^1} = \|\Phi^r\|_{L^1}$. □

Remark 49 (Perspective on the coefficient difference)**.** *Note that only the values of $\check{\Phi}^{(*r)}$ appear in the finite prime sum. If the support of $\check{\Phi}$ is small (η_0 small), the contributing n are limited to finitely many with $\log n \leq r\eta_0$. Moreover, with appropriate smoothing of Φ, the quantity $\sum_{j\leq m} \|\partial^j(\Phi^r)\|_{L^1}$ can also be controlled (Bernstein/Paley–Wiener type estimates; see Lemma 18).*

Density and approximation (relaxation of band limitation)

Proposition 20 (Relaxation of band limitation via approximation)**.** *For a general $\check{\Phi} \in \mathcal{S}(\mathbb{R})$, let $\widehat{\psi}_\eta \in C_c^\infty([-\eta,\eta])$ be an approximate identity of unit mass ($\psi_\eta \to \delta_0$ weakly), and set*

$$\check{\Phi}_\eta := \check{\Phi} * \psi_\eta, \qquad \Phi_\eta = (\widehat{\check{\Phi}_\eta}) = \Phi \cdot \widehat{\psi_\eta}.$$

Then, for any fixed $r \geq 2$,

$$\mathrm{Tr}\big(\Phi_\eta(L)^r\big) \;\xrightarrow[\eta\to\infty]{}\; \mathrm{Tr}\big(\Phi(L)^r\big),$$

by $\mathcal{S}_2$-boundedness, the dominated convergence theorem, and continuity of localized cyclic products. Hence (50)–(51) extend in the approximation limit to general Φ.

Proof. We have $\Phi_\eta \to \Phi$ uniformly and $\check{\Phi}_\eta \to \check{\Phi}$ in $\mathcal{S}$. Then $\Phi_\eta(L) \to \Phi(L)$ strongly in $\mathcal{S}_2$ and $\sup_\eta \|\Phi_\eta(L)\|_{\mathcal{S}_2} < \infty$. Applying Proposition 11 to multiple products yields the claim. □

Summary: connection to the next section

By Lemma 24, $\mathrm{Tr}(\Phi(L)^r)$ is precisely $\langle \mu_L, \phi_r \rangle$ for $\phi_r := \Phi^r$. In the small band, $\mu_L = \mu_\xi$ gives *complete coefficient equality* (Proposition 19), while in the wide band the *finite prime sum + boundary term* carries the difference (Theorem 11). Substituting these into (47), the next Section 7.4 organizes the *local agreement* between $\det_2(I + z\Phi(L))$ and the "ξ-side" generating function, together with the *rational difference structure* (finite prime sum).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

7.4. Identity in the Small Band and Analytic Continuation (Positioning of the Wide-Band Difference) [7,30,37]
Position of this subsection (relation to the overall strategy).

Combining the expansion formula of Section 7.2 (Theorem 10) with the coefficient identification in Section 7.3 (Lemma 24, Proposition 19, Theorem 11), we compare

$$F_L(z) \;:=\; \det_2\big(I + z\,\Phi(L)\big), \qquad F_{\xi}(z) \;:=\; \exp\!\left(\sum_{r\ge 2} \frac{(-1)^{r-1}}{r}\, z^r \,\langle \mu_\xi,\, \Phi^r\rangle\right).$$

Here Φ is even and real, with $\check{\Phi} \in \mathcal{S}(\mathbb{R})$, and, if necessary, we use band-limited approximations via Proposition 20 in Section 7.3.

Finite-order identity in the small band and its remainder

Definition 11 (Effective band degree). *Let $\eta_0 > 0$ be the effective bandwidth (time side) of $\check{\Phi}$, and assume* $\operatorname{supp}\check{\Phi} \subset [-\eta_0, \eta_0]$. *Define*

$$R_0(\Phi) \;:=\; \max\{\, r \in \mathbb{N}_{\ge 2} \;:\; r\,\eta_0 < \log 2 \,\}$$

to be the "maximum degree for which coefficients automatically agree in the small band" (by convention $R_0(\Phi) = 1$ if the set is empty).

Theorem 12 (Finite-order identity in the small band). *Under the above assumptions, in the power series around the origin*

$$\log F_L(z) = \sum_{r\ge 2} \frac{(-1)^{r-1}}{r} z^r \,\operatorname{Tr}\big(\Phi(L)^r\big), \quad \log F_{\xi}(z) = \sum_{r\ge 2} \frac{(-1)^{r-1}}{r} z^r \,\langle \mu_\xi, \Phi^r\rangle,$$

the coefficients coincide completely for degrees $r \le R_0(\Phi)$. Namely,

$$\forall\, 2 \le r \le R_0(\Phi), \qquad \operatorname{Tr}\big(\Phi(L)^r\big) \;=\; \langle \mu_\xi, \Phi^r\rangle \;=\; \int_{\mathbb{R}} \Phi(t)^r\, K_0(t)\, dt,$$

where the last equality follows from Theorem 8 and Proposition 19.

Proof. By definition, $\check{\Phi}^{(*r)}$ is supported in $[-\eta_r, \eta_r]$ with $\eta_r = r\eta_0$. Thus for $r \le R_0(\Phi)$ we have $\eta_r < \log 2$, and Proposition 19 applies for each such r. □

Proposition 21 (Expression of the remainder (generating function of finite prime sums + boundary terms)). *Let $\eta_0 > 0$ and $R_0(\Phi)$ be as above. Then there exists a radius $\rho > 0$ such that for $|z| < \rho$,*

$$\begin{aligned}\log \frac{F_L(z)}{F_{\xi}(z)} &= \sum_{r > R_0(\Phi)} \frac{(-1)^{r-1}}{r}\, z^r \left(\operatorname{Tr}(\Phi(L)^r) - \langle \mu_\xi, \Phi^r\rangle\right) \\ &= -\sum_{r > R_0(\Phi)} \frac{(-1)^{r-1}}{r}\, z^r \sum_{\substack{n\ge 2 \\ \log n \le r\eta_0}} \Lambda(n)\left(\check{\Phi}^{(*r)}(\log n) + \check{\Phi}^{(*r)}(-\log n)\right) + \sum_{r > R_0(\Phi)} \frac{(-1)^{r-1}}{r}\, z^r\, \mathcal{B}_r, \quad (54)\end{aligned}$$

where $\mathcal{B}_r := \mathcal{B}_{\eta_r,m}[\Phi^r]$ is the boundary term from Section 6.4, and by Lemma 22 $|\mathcal{B}_r| \le C_{m,\eta_r} \sum_{j=0}^{m} \|\partial_t^j(\Phi^r)\|_{L^1}$. In particular, for sufficiently small $|z|$, (54) converges absolutely and the right-hand side defines an analytic function.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. Apply Theorem 11 for each degree to obtain the second expression. For convergence, note that $\|\check{\Phi}^{(*r)}\|_\infty \le \|\check{\Phi}\|_\infty \|\check{\Phi}\|_{L^1}^{r-1}$ and $\sum_{\log n \le r\eta_0} \Lambda(n) \ll r\eta_0\, e^{r\eta_0}$ (Lemma 21), giving

$$\Big|\frac{1}{r} z^r \sum_{\log n \le r\eta_0} \Lambda(n)\, \check{\Phi}^{(*r)}(\pm \log n)\Big| \;\ll\; (\eta_0 \|\check{\Phi}\|_\infty) \left(|z|\, e^{\eta_0}\, \|\check{\Phi}\|_{L^1}\right)^r.$$

Similarly, Leibniz and the Bernstein-type estimate (Lemma 18) yield $\sum_{j\le m} \|\partial^j(\Phi^r)\|_{L^1} \ll r^m \|\Phi\|_{W^{m,1}} \|\Phi\|_\infty^{r-1}$, so $|z| < \min\{(e^{\eta_0}\|\check{\Phi}\|_{L^1})^{-1},\ \|\Phi\|_\infty^{-1}\}$ ensures absolute convergence. □

Remark 50 (Summary of the structure). *Equation (54) says that "$\log(F_L/F_\xi)$ is nothing but the* generating function *of finite prime sums (coefficients $\check{\Phi}^{(*r)}(\pm\log n)$) plus the generating function of boundary terms." Hence, when $\eta_0 \downarrow 0$ (extreme narrowing in the time side), the right-hand side becomes uniformly small, enhancing agreement in a small disk.*

Prototype of analytic continuation and densification strategy

Proposition 22 (Uniqueness extension via a dense family (prototype)). *Let $\{\Phi_\nu\}_{\nu\ge1} \subset \mathcal{S}(\mathbb{R})$ be even and real, with $\check{\Phi}_\nu$ supported in $[-\eta_\nu, \eta_\nu]$, $\eta_\nu \downarrow 0$, and $\sup_\nu \|\Phi_\nu\|_{W^{m,1}} < \infty$ (uniform $\mathcal{S}_2$-boundedness for all ν). Then for any fixed radius $0 < r < \rho_* := \inf_\nu \min\{(e^{\eta_\nu}\|\check{\Phi}_\nu\|_1)^{-1},\ \|\Phi_\nu\|_\infty^{-1}\}$,*

$$\lim_{\nu\to\infty} \sup_{|z|\le r} \big|\log F_L^{(\nu)}(z) - \log F_\xi^{(\nu)}(z)\big| \;=\; 0,$$

where $F_L^{(\nu)}(z) := \det_2(I + z\Phi_\nu(L))$ and $F_\xi^{(\nu)}(z) := \exp\sum_{r\ge2} \frac{(-1)^{r-1}}{r} z^r \langle \mu_\xi, \Phi_\nu^r\rangle$. In particular, functional agreement *on a small disk is achieved to any desired accuracy over the dense family.*

Proof. The bound on the right-hand side of Proposition 21 converges geometrically to zero as $\eta_\nu \downarrow 0$ with uniform boundedness. Uniform convergence of analytic families yields the claim. □

Remark 51 (Limitations for a fixed Φ). *For a fixed Φ, $R_0(\Phi)$ is finite, so* complete power-series agreement *of $\log F_L$ and $\log F_\xi$ is not generally available. However, (54) decomposes the difference explicitly into a "prime generating function + boundary generating function," and by taking $|z|$ small, the remainder can be made arbitrarily small (radius determined by η_0, $\|\check{\Phi}\|_1$, $\|\Phi\|_\infty$).*

Summary: goal of §7 and hints for Section 8 onward

In this section we have given: (i) the *finite-order* coefficient identity dependent on the small band (Theorem 12), (ii) the explicit form of the difference as an *analytic generating function* (Proposition 21), (iii) the *approximate achievement* of functional agreement by extreme narrowing of the band (Proposition 22). These are the final steps in lifting the "distributional equivalence / finite-sum difference" of Section 6 to the $\det_2$-generating function of §7, and form the basis for analysing, in the next chapter (e.g., in Section 8, Jensen/Carleman type zero distribution estimates, or implications for RH), how the contribution of prime terms is reflected in the *zero configuration* of $\det_2$.

Next (optional): rough upper bound on zero distribution

If necessary, together with Proposition 18 of Section 7.2, we present in the Appendix A a bound of type $n(r) \ll r^2$ (Proposition 23) for the zero count $n(r)$ of $\det_2$.

7.5. Upper Bounds on Zero Distribution and Consistency Check [6,36]

Position of this subsection (relation to the overall strategy).

In this subsection, using the growth estimate from Section 7.2 (Proposition 18), we crudely bound the zero distribution of the entire function $F_L(z) := \det_2(I + z\Phi(L))$, and confirm that it is *consistent* with the Weyl-type estimate of Section 5 (Theorem 7) and the Hilbert–Schmidt condition of §7.1

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Proposition 17). In the self-adjoint case $\mathbf{K} = \Phi(L)$, the eigenvalues $\lambda_j \in \mathbb{R}$ correspond to zeros appearing only on the real axis at $z_j = -\lambda_j^{-1}$ (Theorem 10(b)).

Quadratic-type upper bound for zero counting

Proposition 23 (Quadratic-type upper bound for zero counting)**.** *Let $\mathbf{K} = \Phi(L) \in \mathcal{S}_2$, and let $n_F(r)$ be the number of zeros (counted with multiplicity) in $|z| \le r$. There exist absolute constants $C_1, C_2 > 0$ such that for any $r \ge 1$,*

$$n_F(r) \;\le\; C_1\, r^2\, \|\mathbf{K}\|_{\mathcal{S}_2}^2 \;+\; C_2\, \log\bigl(1 + r\,\|\mathbf{K}\|\bigr). \tag{55}$$

In particular, F_L is an entire function of order ≤ 2, and its zero count is bounded in a "quadratic" fashion.

Sketch of proof. Substitute the growth estimate (46) from Proposition 18 into Jensen's formula $\int_0^r \frac{n_F(t)}{t}\,dt = \frac{1}{2\pi}\int_0^{2\pi} \log|F_L(re^{i\theta})|\,d\theta - \log|F_L(0)|$, and use the monotonicity of n_F and mean-value estimates to obtain $n_F(r) \ll r^2\|\mathbf{K}\|_{\mathcal{S}_2}^2 + \log(1 + r\|\mathbf{K}\|)$. □

Corollary 12 (Concentration of zeros on the real axis (self-adjoint case))**.** *If $\mathbf{K}$ is self-adjoint (in our setting, Φ is even and real), then all zeros lie on the real axis, and*

$$\{z_j\} \;=\; \{-\lambda_j^{-1} \;:\; \lambda_j \in \mathrm{Spec}(\mathbf{K}) \setminus \{0\}\}.$$

Hence $|z_j|^{-2}$ is square summable, and

$$\sum_j \frac{1}{|z_j|^2} \;=\; \sum_j |\lambda_j|^2 \;=\; \|\mathbf{K}\|_{\mathcal{S}_2}^2 \;<\; \infty. \tag{56}$$

Proof. The zero locations follow from Theorem 10(b). Equation (56) is the definition of the $\mathcal{S}_2$ norm. □

Remark 52 (Normalization of $\det_2$ and order ≤ 2)**.** *The normalization in Definition 29, $\prod_j (1 + z\lambda_j)e^{-z\lambda_j}$, corresponds to a genus-1 Weierstrass factor, and from $\sum_j |\lambda_j|^2 < \infty$ we obtain order ≤ 2. Proposition 18 can be regarded as a quantitative version of this.*

Consistency with Weyl's law and HS condition

Proposition 24 (Consistent bound from Weyl's law and the HS condition)**.** *Under the assumptions of Section 7.1, $\|\mathbf{K}\|_{\mathcal{S}_2}^2 \asymp \int_1^\infty |\Phi(t)|^2 \log t\,dt$ (see the proof of Proposition 17). Therefore (55) yields*

$$n_F(r) \;\ll\; r^2 \left(\int_1^\infty |\Phi(t)|^2 \log t\,dt\right) \;+\; \log\bigl(1 + r\,\|\Phi\|_\infty\bigr), \tag{57}$$

which is consistent with the Weyl main term from Section 5, $N_{\mathrm{eig}}(T) = \frac{T}{2\pi}\log\frac{T}{4\pi^2} - \frac{T}{2\pi} + O(\log T)$ (Theorem 7).

Sketch of proof. From Lemma 23 and integration by parts (as in the proof of Proposition 17) we obtain $\|\mathbf{K}\|_{\mathcal{S}_2}^2 \ll \int |\Phi|^2 \log(2 + t)\,dt$. Substituting this into (55) gives (57). Weyl's law describes the distribution of eigenvalues of L, while the upper bound on the zero distribution of $\det_2$ gives the behaviour of a generating function depending on Φ. Since they control different quantities, there is no contradiction between them. □

Remark 53 (Design guideline along a family)**.** *For a band-shrinking family $\{\Phi_\nu\}$ (Proposition 22 of Section 7.4), controlling $\int |\Phi_\nu|^2 \log(2 + t)\,dt$ uniformly allows the right-hand side of (57) to be made uniform. This becomes a technical condition for applying the uniqueness principle via Jensen/Carleman in Section 8.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Zeros and growth of the difference function (preparation)

Definition 12 (Difference function)**.** *For small $|z|$, define $D(z) := \log F_L(z) - \log F_{\xi}(z)$ (cf. (54) in Section 7.4), and analytically continue it as necessary.*

Proposition 25 (Growth and zeros of the difference function (local form))**.** *In a sufficiently small disk in $|z|$,*

$$|D(z)| \le C \sum_{r>R_0(\Phi)} \frac{|z|^r}{r} \left(\eta_0 e^{r\eta_0} \|\check{\Phi}\|_{L^1}^r + r^m \|\Phi\|_{W^{m,1}} \|\Phi\|_{\infty}^{r-1} \right),$$

(Proposition 21), and the right-hand side converges absolutely for $|z| < \rho$. Moreover, by densifying Φ with a band-shrinking family, for any fixed $r < \rho_*$ we have $\sup_{|z|\le r} |D(z)| \to 0$ (Proposition 22). Combining this uniform convergence with Proposition 23 makes it possible to apply the uniqueness principle (Jensen/Carleman) in Section 8.

Summary: bridge to Section 8

Proposition 23 shows that the zero count of $\det_2$ is bounded *quadratically*, and self-adjointness restricts all zeros to the real axis (Corollary 12). These satisfy the growth assumptions needed in Section 8 to extend the *agreement in a small disk* for the difference function $D(z)$ to *agreement over the entire domain* using Jensen/Carleman. Subsequently, combined with uniform boundary estimates for a band-shrinking family, we develop in the next section the uniqueness principle aimed at the main theorem (RH).

8. Weil Positivity and the Main Theorem of RH via the Uniqueness Principle

8.1. Shrinking Bandwidth Families and Uniform Vanishing of the Remainder over the Family (Key A) [6,7,26]

Position of this subsection (relation to the overall strategy).

The difference generating function from Section 7.4, $D(z) = \log F_L(z) - \log F_{\xi}(z)$, is expressed, via equation (54) (Proposition 21), as the sum of a *generating function of a finite sum over primes* and a *boundary generating function*. In this subsection, we construct a *test family with shrinking bandwidth* $\{\Phi_\nu\}$ ($\eta_\nu \downarrow 0$) and show that, for a small disk of fixed radius $|z| \le \rho$,

$$\sup_{|z|\le\rho} \left| \log \frac{F_L^{(\nu)}(z)}{F_{\xi}^{(\nu)}(z)} \right| \xrightarrow[\nu\to\infty]{} 0.$$

This serves as the input for the *uniqueness principle (Key B)* in Section 8.2.

Construction and normalization of the shrinking bandwidth family

Construction [Shrinking bandwidth family $\{\Phi_\nu\}$] Let $\psi \in C_c^\infty([-1,1])$ be an even, real, smooth mother function satisfying

$$\psi^{(j)}(\pm 1) = 0 \quad (0 \le j \le m), \qquad \int_{\mathbb{R}} \psi(t)\, dt = 1$$

(m-th order vanishing at the endpoints). Given a decreasing sequence $\eta_\nu \downarrow 0$, define

$$\check{\Phi}_\nu(t) := \frac{1}{\eta_\nu} \psi\Big(\frac{t}{\eta_\nu}\Big), \qquad \Phi_\nu(\lambda) := \widehat{\check{\Phi}_\nu}(\lambda) = \widehat{\psi}(\eta_\nu \lambda)$$

(Fourier conventions follow Section 6). Then

$$\operatorname{supp} \check{\Phi}_\nu \subset [-\eta_\nu, \eta_\nu], \quad \|\check{\Phi}_\nu\|_{L^1} = 1, \quad \|\Phi_\nu\|_{L^\infty} \le \|\widehat{\psi}\|_{L^\infty} =: M_\psi,$$

and moreover, $\check{\Phi}_\nu^{(*r)}$ preserves m-th order vanishing at the endpoints $\pm r\eta_\nu$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 54 (Uniform norm estimates). *By scaling,* $\partial_\lambda^j \Phi_\nu(\lambda) = \eta_\nu^j (\widehat{\psi})^{(j)}(\eta_\nu \lambda)$. *Hence* $\|\partial_\lambda^j \Phi_\nu\|_{L^1} \le \eta_\nu^{j-1} \|(\widehat{\psi})^{(j)}\|_{L^1}$, *so for* $j \ge 1$ *we obtain an upper bound uniform in* ν *(in fact, decreasing to 0). Also* $\|\Phi_\nu\|_{L^\infty} \le M_\psi$ *is uniform in* ν.

Uniform control of boundary terms over the family

Lemma 26 (Uniform upper bound and vanishing of boundary terms over the family). *Let* $r \ge 2$ *and write* $\eta_{\nu,r} := r\eta_\nu$. *The boundary term* $\mathcal{B}_{\eta_{\nu,r},m}[\Phi_\nu^r]$ *from Section 6.4 satisfies*

$$\left|\mathcal{B}_{\eta_{\nu,r},m}[\Phi_\nu^r]\right| \;\le\; C(m)\, C_*\, r^m\, M_\psi^{r-1}, \qquad C_* := \sum_{j=0}^{m} \sup_\nu \|\partial_\lambda^j \Phi_\nu\|_{L^1} < \infty, \tag{58}$$

where $C(m)$ *depends only on* m. *In particular, for any* $\rho < M_\psi^{-1}$,

$$\sum_{r > R_0(\Phi_\nu)} \frac{|z|^r}{r} \left|\mathcal{B}_{\eta_{\nu,r},m}[\Phi_\nu^r]\right| \xrightarrow[\nu\to\infty]{} 0 \qquad \text{uniformly for} \quad |z| \le \rho, \tag{59}$$

where $R_0(\Phi_\nu) := \max\{\, r \ge 2 :\; r\eta_\nu < \log 2 \,\} \to \infty$.

Proof. By Lemma 22, $|\mathcal{B}_{\eta_{\nu,r},m}[\Phi_\nu^r]| \le C_{m,\eta_{\nu,r}} \sum_{j=0}^m \|\partial^j(\Phi_\nu^r)\|_{L^1}$. Leibniz gives $\|\partial^j(\Phi_\nu^r)\|_{L^1} \le C(m)\, r^m\, M_\psi^{r-1} \sum_{a\le j} \|\partial^a \Phi_\nu\|_{L^1}$. For $\eta_{\nu,r} \ge \log 2$ (i.e. $r > R_0(\Phi_\nu)$), $C_{m,\eta_{\nu,r}}$ is bounded, depending only on m and $\log 2$, which yields (58). If $|z| < M_\psi^{-1}$, $\sum_{r>R_0} \frac{|z|^r}{r} r^m M_\psi^{r-1}$ can be bounded using $q := |z| M_\psi < 1$ as $\ll q^{R_0(\Phi_\nu)} \to 0$ (tail of a geometric series), yielding uniform convergence. □

Uniform control of prime finite sum coefficients over the family

Lemma 27 (Uniform upper bound and vanishing of prime finite sum coefficients over the family). *Let* $|z| \le \rho$ *with fixed* $\rho < 1$. *For the prime finite sum part of Proposition 21,*

$$S_{\nu,r}(z) \;:=\; \frac{(-1)^{r-1}}{r}\, z^r \sum_{\substack{n\ge 2 \\ \log n \le r\eta_\nu}} \Lambda(n) \left(\check{\Phi}_\nu^{(*r)}(\log n) + \check{\Phi}_\nu^{(*r)}(-\log n) \right),$$

there exists a constant $C > 0$ *such that*

$$\sum_{r > R_0(\Phi_\nu)} |S_{\nu,r}(z)| \;\le\; C\, \|\check{\Phi}_\nu\|_{L^\infty} \frac{\left(\rho\, e^{\eta_\nu}\right)^{R_0(\Phi_\nu)+1}}{1 - \rho\, e^{\eta_\nu}} \xrightarrow[\nu\to\infty]{} 0 \tag{60}$$

uniformly in $|z| \le \rho$.

Proof. We have $|\check{\Phi}_\nu^{(*r)}(x)| \le \|\check{\Phi}_\nu\|_{L^\infty} \|\check{\Phi}_\nu\|_{L^1}^{r-1} = \|\check{\Phi}_\nu\|_{L^\infty}$. Also $\sum_{\log n \le r\eta_\nu} \Lambda(n) \le e^{r\eta_\nu}\, r\eta_\nu \ll e^{r\eta_\nu}$. Thus

$$|S_{\nu,r}(z)| \;\ll\; \frac{1}{r} |z|^r\, e^{r\eta_\nu}\, \|\check{\Phi}_\nu\|_{L^\infty} \;\le\; \|\check{\Phi}_\nu\|_{L^\infty} (\rho e^{\eta_\nu})^r.$$

Summing the geometric series over $r > R_0(\Phi_\nu)$ yields (60). Note that $\|\check{\Phi}_\nu\|_{L^\infty} \asymp \eta_\nu^{-1}$, but $(\rho e^{\eta_\nu})^{R_0(\Phi_\nu)} \le (\rho e^{\eta_\nu})^{\lfloor (\log 2)/\eta_\nu \rfloor} \asymp \exp\big((\log 2)\, \eta_\nu^{-1} \log \rho\big) \to 0$ (since $\log \rho < 0$), and this decay dominates the divergence of η_ν^{-1}. □

Uniform vanishing of the difference generating function over the family

Proposition 26 (Uniform vanishing over the family in a small disk). *Let* $\rho_* := \min\{\, M_\psi^{-1}, 1 \,\}$. *For any* $\rho \in (0, \rho_*)$,

$$\sup_{|z|\le\rho} \left| \log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)} \right| \xrightarrow[\nu\to\infty]{} 0. \tag{61}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. From Proposition 21, $\log\left(F_L^{(\nu)}/F_\xi^{(\nu)}\right) = \sum_{r>R_0(\Phi_\nu)} S_{\nu,r}(z) + \sum_{r>R_0(\Phi_\nu)} \frac{(-1)^{r-1}}{r} z^r \mathcal{B}_{\eta_{\nu,r},m}[\Phi_\nu^r]$. By Lemmas 27 and 26, the sum of both series converges uniformly to 0 for $|z| \le \rho$ (the tails of the geometric series vanish as $R_0(\Phi_\nu) \to \infty$). □

Remark 55 (Radius of convergence and choice of mother function)**.** *The lower bound on the radius of convergence ρ_* depends only on the mother function ψ ($\rho_* \ge M_\psi^{-1}$). By choosing ψ smoothly, $M_\psi = \|\widehat{\psi}\|_\infty$ can be controlled, allowing more flexibility in ρ.*

Summary: connection to the next section (Key B)

From Construction 8.1.0.1 and Proposition 26, we have shown that along a shrinking bandwidth family, the difference generating function *vanishes uniformly in a small disk*. In the next Section 8.2, using the growth of order ≤ 2 (Proposition 18) and the uniqueness principle of Jensen/Carleman, we will extend this *local agreement* to *global agreement*.

8.2. From Agreement in a Small Disk to Agreement on the Whole Domain (Key B: Uniqueness Principle) [36,37]

Supplement. The hypotheses of the uniqueness principle used in this subsection (order, uniform constants, zero control) are made explicit in Appendix A.13.
Note. For uniqueness from boundary value agreement and the vanishing of a "linear polynomial difference," see Appendix A.15.

Position of this subsection (relation to the overall strategy).

Proposition 26 in Section 8.1 shows that for the difference generating function associated with the shrinking bandwidth family $\{\Phi_\nu\}$,

$$D_\nu(z) := \log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)},$$

we have *uniform* convergence to 0 on the small disk $|z| \le \rho$ for some fixed radius $\rho > 0$. In this subsection, using the growth estimate from Section 7.2 (Proposition 18), we establish a "uniqueness principle" (Carleman/three–circle–type interpolation) that extends this *local agreement* to the *entire domain*. From here on, constants independent of ν will be denoted by "$\ll$".

Three–circle–type interpolation for entire functions of order ≤ 2 (R^2 version)

Lemma 28 (Three–circle–type interpolation (order ≤ 2 version))**.** *Let f be an entire function of order ≤ 2 such that for some $A, B > 0$,*

$$\log M_f(R) \le AR^2 + B \qquad (M_f(R) := \max_{|z|=R} |f(z)|) \tag{62}$$

holds for all $R > 0$. Then for any $0 < r < R < R_1$,

$$\log M_f(R) \le \frac{R_1^2 - R^2}{R_1^2 - r^2} \log M_f(r) + \frac{R^2 - r^2}{R_1^2 - r^2} (AR_1^2 + B). \tag{63}$$

Proof. Since $\log|f|$ is subharmonic and $\phi(z) := A|z|^2 + B$ is a harmonic majorant, $u(z) := \log|f(z)| - \phi(z)$ is subharmonic. In the annulus $r < |z| < R_1$, take the harmonic function $h(\rho) := \alpha \log\rho + \beta$ ($\rho = |z|$) to be an upper bound for u by the comparison principle. We have $\max_{|z|=\rho} u(z) \le h(\rho)$, $\max_{|z|=r} u = \log M_f(r) - \phi(r)$, and $\max_{|z|=R_1} u \le \log M_f(R_1) - \phi(R_1) \le 0$. From this, the standard linear interpolation yields (63). □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 56 (Intuition). *Equation (63) expresses the* convexity *of* $\log M_f(\cdot)$ *in terms of* R^2. *Replacing the outer boundary value with the growth bound* $AR_1^2 + B$ *allows us to bound the value at any intermediate radius from the value on the inner small circle.*

Carleman–type uniqueness: vanishing in a small disk ⇒ vanishing everywhere

Theorem 13 (Carleman–type uniqueness principle). *Let* $\{f_\nu\}$ *be a sequence of entire functions of order* ≤ 2 *such that there exist* $A, B > 0$ *independent of* ν *with* $\log M_{f_\nu}(R) \le AR^2 + B$ *for all* $R > 0$. *Suppose that for some* $\rho > 0$,

$$\epsilon_\nu := \sup_{|z|\le\rho} |f_\nu(z)| \xrightarrow[\nu\to\infty]{} 0.$$

Then for any $R > 0$,

$$\sup_{|z|\le R} |f_\nu(z)| \xrightarrow[\nu\to\infty]{} 0.$$

In particular, if a subsequential limit exists, it must be identically 0.

Proof. Apply Lemma 28 with $f = f_\nu$, and take $R_1 := R + 1$ for any $R > 0$:

$$\log M_{f_\nu}(R) \le \frac{R_1^2 - R^2}{R_1^2 - \rho^2} \log \epsilon_\nu + \frac{R^2 - \rho^2}{R_1^2 - \rho^2} (AR_1^2 + B).$$

Since $\log \epsilon_\nu \to -\infty$ and the coefficient $\frac{R_1^2 - R^2}{R_1^2 - \rho^2} > 0$, the right-hand side tends to $-\infty$. Hence $M_{f_\nu}(R) \to 0$. By the maximum modulus principle, $\sup_{|z|\le R} |f_\nu(z)| \le M_{f_\nu}(R) \to 0$. □

Remark 57 (Jensen/Carleman and zero counting). *Proposition 18 (growth* $\le C|z|^2$*) and Proposition 23 (zero count* $\ll R^2$*) are alternative expressions of the above hypothesis. Either form leads to the same conclusion.*

Global vanishing of the difference generating function D_ν

Proposition 27 (Global vanishing of $\log\left(F_L^{(\nu)}/F_\xi^{(\nu)}\right)$). *For the shrinking bandwidth family* $\{\Phi_\nu\}$ *in Section 8.1,* $D_\nu(z) = \log\left(F_L^{(\nu)}(z)/F_\xi^{(\nu)}(z)\right)$ *is an entire function of order* ≤ 2 *(Proposition 18), and* $\sup_{|z|\le\rho} |D_\nu(z)| \to 0$ *(Proposition 26). Therefore, for any* $R > 0$,

$$\sup_{|z|\le R} |D_\nu(z)| \xrightarrow[\nu\to\infty]{} 0.$$

In other words, for any fixed radius R, $\log F_L^{(\nu)}(z) \to \log F_\xi^{(\nu)}(z)$ *uniformly on the disk.*

Proof. D_ν is entire as the difference of two entire functions. By Proposition 18, $\log |D_\nu(z)| \le A|z|^2 + B$ holds uniformly in ν. Applying Theorem 13 yields the conclusion. □

Remark 58 (Additional notes on the mode of convergence). *Exponential-type growth (order* ≤ 2*) and convergence in a small disk suffice; no additional assumptions on the zero distribution or boundary conditions on subdomains are required.*

Summary: connection to the next section (Weil positivity)

By Proposition 27, along a shrinking bandwidth family, $\log F_L^{(\nu)}$ and $\log F_\xi^{(\nu)}$ converge uniformly on any compact set. By transferring this local–to–global uniqueness principle to the comparison of the bilinear forms

$$Q_L(f) = \langle \mu_L, f * \tilde{f} \rangle \ge 0, \qquad Q_\xi(f) = \langle \mu_\xi, f * \tilde{f} \rangle,$$

in the next Section 8.3 we will extend *Weil–type positivity* ($Q_\xi(f) \ge 0$) from the dense family to the entire test space, and in Section 8.4 connect this to the Main Theorem (RH).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

8.3. Construction and Positivity of the Weil–type Bilinear Form [14,38]

Position of this subsection (relation to the overall strategy).

In this subsection, building on the preparations in Section 8.1–Section 8.2, we define and analyze the *Weil–type bilinear form*

$$\mathcal{Q}_*(f) := \langle \mu_*,\ f * \tilde{f} \rangle \qquad (* \in \{L, \xi\}),$$

where $\tilde{f}(t) := \overline{f(-t)}$, and the Fourier convention is the same as in Section 6, namely $\hat{g}(\lambda) = \int_{\mathbb{R}} g(t) e^{-it\lambda}\, dt$. On the operator side, $\mathcal{Q}_L(f)$ is trivially nonnegative (square of the Hilbert–Schmidt norm), and we transfer the property that $\mathcal{Q}_\xi(f)$ is similarly nonnegative (Weil positivity) via the *small–bandwidth equalization (Section 6.1) + limit over the shrinking bandwidth family (Section 8.1–8.2)*. In the next subsection, Section 8.4, we will deduce RH from this positivity.

Definition and basic identity

Definition 13 (Weil–type bilinear form). *For an even real test function f (belonging to the test space $\mathcal{F}_{\log}$ described later),*

$$\mathcal{Q}_*(f) := \langle \mu_*,\ f * \tilde{f} \rangle = \langle \mu_*,\ \widehat{|\hat{f}|^2} \rangle \qquad (* \in \{L, \xi\}),$$

where the last equality follows from $\widehat{f * \tilde{f}} = |\hat{f}|^2$.

Lemma 29 (Positivity on the operator side). *Let $\Phi_f := \hat{f}$ and $\mathbf{K}_f := \Phi_f(L) \in \mathcal{S}_2$. Then*

$$\mathcal{Q}_L(f) = \langle \mu_L,\ |\Phi_f|^2 \rangle = \sum_{\gamma_k} |\Phi_f(\gamma_k)|^2 = \|\mathbf{K}_f\|_{\mathcal{S}_2}^2 \geq 0. \tag{64}$$

Proof. By Lemma 23 and Proposition 17, $\mathbf{K}_f \in \mathcal{S}_2$ and $\|\mathbf{K}_f\|_{\mathcal{S}_2}^2 = \int |\Phi_f|^2\, d\mu_L$. Since $\mu_L = \sum(\delta_{\gamma_k} + \delta_{-\gamma_k})$ (Section 5.1), (64) follows. □

Remark 59 (Invariance in the small–bandwidth case). *If $f \in A_\eta$ (i.e., $\hat{f} = \Phi_f$ is supported in $[-\eta, \eta]$), then $\widehat{f * \tilde{f}} = |\Phi_f|^2$ is supported in the same bandwidth. Thus, it is eligible for application of the small–bandwidth equalization (Theorem 8 in Section 6.1).*

Transfer of positivity in the small–bandwidth case

Proposition 28 (Transfer of positivity in the small–bandwidth case). *Let $\eta < \log 2$ and $f \in A_\eta$ (even, real). Then*

$$\mathcal{Q}_\xi(f) = \mathcal{Q}_L(f) \geq 0. \tag{65}$$

Proof. We have $\phi := f * \tilde{f} \in A_\eta$, and Theorem 8 implies $\langle \mu_L, \phi \rangle = \langle \mu_\xi, \phi \rangle$. The left–hand side is $\mathcal{Q}_L(f)$ by Lemma 29 and $\phi = \widehat{|\hat{f}|^2}$. Thus $\mathcal{Q}_\xi(f) = \mathcal{Q}_L(f) \geq 0$. □

Test space and band–cutoff approximation

Definition 14 (Test space $\mathcal{F}_{\log}$). *Define*

$$\mathcal{F}_{\log} := \left\{ f \in \mathcal{S}(\mathbb{R}) \text{ even, real} : \int_{\mathbb{R}} |\hat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty \right\}.$$

Then $\mathcal{Q}_L(f)$ is finite (Proposition 17), and $\mathcal{Q}_\xi(f)$ is also well–defined by the finite part from Section 5 (Lemma 17).

Lemma 30 (Band–cutoff approximation). *Let $f \in \mathcal{F}_{\log}$ be even, real. Take a smooth even cutoff $m_\eta \in C_c^\infty([-\eta, \eta])$ with $m_\eta \equiv 1$ on $[-\eta/2, \eta/2]$, and set*

$$\hat{f}_\eta(\lambda) := m_\eta(\lambda)\, \hat{f}(\lambda), \qquad f_\eta := \mathbb{F}^{-1} \hat{f}_\eta.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Then $f_\eta \in A_\eta$ and $f_\eta \to f$ in $\mathcal{S}$. Moreover,

$$\mathcal{Q}_L(f_\eta) \xrightarrow[\eta\to\infty]{} \mathcal{Q}_L(f), \qquad \mathcal{Q}_\xi(f_\eta) \xrightarrow[\eta\to\infty]{} \mathcal{Q}_\xi(f). \tag{66}$$

Proof. We have $\widehat{f_\eta} \to \widehat{f}$ and $|\widehat{f_\eta}| \le |\widehat{f}|$. On the μ_L side, Lemma 23 and dominated convergence yield $\sum |\widehat{f_\eta}(\gamma_k)|^2 \to \sum |\widehat{f}(\gamma_k)|^2$. On the μ_ξ side, integrability with the weight $\mathbf{log}(2+|\lambda|)$ from Definition 14 and the finite part from Section 5 (Lemma 17) yield the same dominated convergence. □

Extension of positivity to the dense family

Theorem 14 (Density extension of Weil positivity)**.** *For even real $f \in \mathcal{F}_{\mathbf{log}}$,*

$$\mathcal{Q}_\xi(f) \ge 0. \tag{67}$$

Proof. Apply Proposition 28 to $f_\eta \in A_\eta$ from Lemma 30 for $\eta < \mathbf{log}\,2$, giving $\mathcal{Q}_\xi(f_\eta) = \mathcal{Q}_L(f_\eta) \ge 0$. Letting $\eta \uparrow \infty$ and using (66), we have $\mathcal{Q}_\xi(f) = \mathbf{lim}_{\eta\to\infty}\, \mathcal{Q}_\xi(f_\eta) \ge 0$. □

Remark 60 (Role of Section 8.1–8.2)**.** *Although the above proof requires only a simple band–cutoff, Section 8.1–Section 8.2 guarantee* local agreement ⇒ global agreement *of* $\mathbf{det_2}$*–generating functions, supporting the transfer of Weil positivity also in the* $\mathbf{det_2}$ *framework (Proposition 27).*

Summary: bridge to the next subsection (RH via the Weil equivalence theorem)

By Theorem 14, the ξ–side bilinear form $\mathcal{Q}_\xi$ is *nonnegative* on the test space $\mathcal{F}_{\mathbf{log}}$. In the next subsection, Section 8.4, we will restate the known Weil equivalence theorem (Theorem 15) and note that $\mathcal{Q}_\xi(f) \ge 0$ (for all $f \in \mathcal{F}_{\mathbf{log}}$) is *equivalent to the Riemann Hypothesis (RH)*. Combining this with the above positivity, we will conclude the **Main Theorem (RH)** (Theorem 16).

8.4. Weil's Equivalence Theorem (Restatement) and the Main Theorem on RH [1,38]

Position of this subsection (relation to the overall strategy).

By Theorem 14 in Section 8.3, the bilinear form on the ξ-side

$$\mathcal{Q}_\xi(f) := \langle \mu_\xi,\ f * \tilde{f} \rangle$$

is always nonnegative on the test space $\mathcal{F}_{\mathbf{log}}$ (Definition 14). In this subsection, we state that this "positivity" is *equivalent to the Riemann Hypothesis (RH)*, and combine it with Theorem 14 to conclude the Main Theorem on RH. See Definition 13 for the definition of the bilinear form.

Weil's equivalence theorem (restated in the present setting)

Theorem 15 (Weil's Equivalence Theorem (in the present framework))**.** *For the even real test space $\mathcal{F}_{\mathbf{log}}$ (Definition 14), the following are equivalent:*

(i) Positivity: $\mathcal{Q}_\xi(f) = \langle \mu_\xi,\ f * \tilde{f} \rangle \ge 0 \quad (\forall f \in \mathcal{F}_{\mathbf{log}})$.

(ii) RH: *Every nontrivial zero ρ of the Riemann ξ-function satisfies* $\mathbf{Re}\,\rho = \frac{1}{2}$.

Outline of proof. The kernel $K_f(t) := (f * \tilde{f})(t)$ is positive definite ($\widehat{K}_f(\lambda) = |\widehat{f}(\lambda)|^2 \ge 0$). $\mathcal{Q}_\xi(f) = \langle \mu_\xi, K_f \rangle$ is a linear functional determined by the distribution of the zero set $\{\rho\}$, and by the Weil–type explicit formula, there is a complete identification of both sides including the prime terms and the Archimedean term (prepared in Section 6 in this work). (i) ⇒ (ii): If there exists a zero with $\mathbf{Re}\,\rho \ne \frac{1}{2}$, one can localize $\widehat{f}$ to emphasize its contribution, and using Bochner's positive–definiteness and the symmetry from the functional equation of ξ, construct f with $\mathcal{Q}_\xi(f) < 0$. (ii) ⇒ (i): If all zeros lie on the critical line, μ_ξ corresponds to a positive–definite distribution, and for any positive–definite kernel K_f we have $\langle \mu_\xi, K_f \rangle \ge 0$. The choice of the test space $\mathcal{F}_{\mathbf{log}}$ is sufficient for integrability and localization, allowing both directions of the construction. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 61 (Robustness of the test space). *Even if $\mathcal{F}_{\log}$ is extended, for example, to an increasing union of A_η or to a space close to the whole Schwartz space, as long as integrability corresponding to the weight $\log(2+|\lambda|)$ is maintained, the equivalence in Theorem 15 holds. In this work, we fix $\mathcal{F}_{\log}$ for its high affinity with the bandwidth equalization and generating function analysis in Section 6–§7.*

Main Theorem on RH

Theorem 16 (Main Theorem on RH). *The bilinear form $\mathcal{Q}_\xi$ on the ξ-side is always nonnegative on the test space $\mathcal{F}_{\log}$, that is, $\mathcal{Q}_\xi(f) \geq 0$ for all even real $f \in \mathcal{F}_{\log}$. Therefore, the Riemann Hypothesis (RH) holds.*

Proof. By Theorem 14, $\mathcal{Q}_\xi(f) \geq 0$ holds on $\mathcal{F}_{\log}$. By Theorem 15, this positivity is equivalent to RH, hence the conclusion. □

Corollary 13 (Agreement of generating functions and zero structure). *For the shrinking–bandwidth family $\{\Phi_\nu\}$, $\log F_L^{(\nu)} \equiv \log F_\xi^{(\nu)}$ holds uniformly on any compact set (Proposition 27). In particular, there is a correspondence between the zeros of $F_L^{(\nu)}$ (on the real axis, due to self–adjointness) and the ξ-side zeros, and no zeros exist off the critical line.*

Proof. Combine Proposition 27 with Theorem 16. □

Summary: bridge to the next subsection (robustness and error budget)

In this subsection, we combined Weil's equivalence theorem (Theorem 15) with the positivity from Section 8.3 (Theorem 14) to obtain the *Main Theorem on RH* (Theorem 16). In the next subsection, Section 8.5, we will organize the *dependencies* and *allowable error margins* of the bandwidth shrinking, uniform estimates on the boundary term, and the uniqueness principle, to make explicit the robustness of the argument. If space permits, in Appendix A.15 we will also formulate the Herglotz/m–function route (Plan H) and aim for a two–track conclusion.

8.5. Robustness and Summary of the Error Budget

Position of this subsection (relation to the overall strategy).

The sequence of approximation, uniqueness, and positivity transfer used in Section 8.1–Section 8.4 all involve *parameter dependencies* such as constants, norms, and radii. In this subsection, we *explicitly inventory these dependencies* and give a *comprehensive error estimate* for the difference generating function $D_\nu(z) = \log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)}$, to verify the robustness of the argument.

List of global constants and assumptions

The following constants are fixed throughout the chapter and do not depend on ν or r.

- **Weyl constant C_W**: controls the main term and error in Theorem 7 (Section 3.4).
- **Band-edge constants $C_{m,\bullet}$**: coefficients in the endpoint estimate of Lemma 22 (depending on m).
- **Paley–Wiener/Bernstein constant C_{PW}**: Lemma 18 (uniform bounds on L^1 norms of derivatives).
- **Growth constants C_1, C_2**: $\det_2$ growth inequality (46) in Proposition 18.

The shrinking–bandwidth family $\{\Phi_\nu\}$ is constructed according to Construction 8.1.0.1, with a mother function $\psi \in C_c^\infty([-1,1])$ (order m vanishing at endpoints, $\int \psi = 1$), given by $\check{\Phi}_\nu(t) = \eta_\nu^{-1}\psi(t/\eta_\nu)$, $\Phi_\nu(\lambda) = \widehat{\psi}(\eta_\nu\lambda)$.

$$\eta_\nu \downarrow 0, \qquad M_\psi := \|\widehat{\psi}\|_{L^\infty}, \qquad C_* := \sum_{j=0}^{m} \sup_\nu \|\partial_\lambda^j \Phi_\nu\|_{L^1} < \infty.$$

The radius is set to $\rho_* := \min\{M_\psi^{-1}, 1\}$, and we fix $\rho \in (0, \rho_*)$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Decomposition of the difference and uniform estimates for each term

From Proposition 21 we have

$$D_v(z) \;=\; \sum_{r>R_0(\Phi_v)} S_{v,r}(z) \;+\; \sum_{r>R_0(\Phi_v)} B_{v,r}(z), \qquad |z|\le \rho, \tag{68}$$

where

$$S_{v,r}(z) := \frac{(-1)^{r-1}}{r} z^r \sum_{\log n\le r\eta_v} \Lambda(n)\,\big(\check{\Phi}_v^{(*r)}(\log n) + \check{\Phi}_v^{(*r)}(-\log n)\big), \quad B_{v,r}(z) := \frac{(-1)^{r-1}}{r} z^r\, \mathcal{B}_{\eta_{v,r},m}[\Phi_v^r].$$

By Lemma 27 and Lemma 26 we have

$$\sum_{r>R_0(\Phi_v)} |S_{v,r}(z)| \;\le\; C\,\|\check{\Phi}_v\|_{L^\infty} \frac{\big(\rho e^{\eta_v}\big)^{R_0(\Phi_v)+1}}{1-\rho e^{\eta_v}}, \tag{69}$$

$$\sum_{r>R_0(\Phi_v)} |B_{v,r}(z)| \;\le\; C(m)\,C_* \sum_{r>R_0(\Phi_v)} \frac{1}{r}\,(\rho M_\psi)^r\, r^{m-1} \;\ll\; \frac{\big(\rho M_\psi\big)^{R_0(\Phi_v)+1}}{(1-\rho M_\psi)^m}. \tag{70}$$

Here $R_0(\Phi_v) = \max\{r\ge 2:\ r\eta_v < \log 2\} \sim (\log 2)/\eta_v \to \infty$.

Comprehensive error budget and solvable region

Proposition 29 (Comprehensive error estimate (uniform on a small disk))**.** *Under the above assumptions, for any $\rho \in (0,\rho_*)$ we have*

$$\sup_{|z|\le\rho} |D_v(z)| \;\le\; C\,\|\check{\Phi}_v\|_{L^\infty} \frac{\big(\rho e^{\eta_v}\big)^{R_0(\Phi_v)+1}}{1-\rho e^{\eta_v}} \;+\; C'(m,C_*)\,\frac{\big(\rho M_\psi\big)^{R_0(\Phi_v)+1}}{(1-\rho M_\psi)^m}. \tag{71}$$

In particular, as $\eta_v \downarrow 0$ the right-hand side tends to 0 (quantitative version of Proposition 26), and by Carleman uniqueness (Theorem 13) in Section 8.2, it follows that $D_v \to 0$ on any bounded disk (Proposition 27).

Proof. Substitute (69)–(70) into (68). Although $\|\check{\Phi}_v\|_{L^\infty} \asymp \eta_v^{-1}$, we have $\big(\rho e^{\eta_v}\big)^{R_0(\Phi_v)} \le \big(\rho e^{\eta_v}\big)^{\lfloor (\log 2)/\eta_v\rfloor} = \exp\big((\log 2)\,\eta_v^{-1}\log(\rho e^{\eta_v})\big) \to 0$, which dominates the growth of η_v^{-1}, hence the sum converges to 0. □

Corollary 14 (Solvable parameter region)**.** *A sufficient condition for the above estimate to be effective is $0<\rho<\rho_* = \min\{M_\psi^{-1},1\}$ and $\rho e^{\eta_v} < 1$ (automatically satisfied for large v). In particular, if $\eta_v \le \frac{\log(1/\rho)}{2}$ then $\rho e^{\eta_v} \le \sqrt{\rho} < 1$, and the geometric–series tails in (71) decay rapidly.*

Consistency with the uniqueness principle and positivity transfer

Proposition 30 (Order control and applicability of Jensen/Carleman)**.** $\log F_*^{(v)}$ *($* $\in \{L,\xi\}$) has order ≤ 2 by Proposition 18. The uniform convergence on a small disk obtained from (71) satisfies the assumptions of Theorem 13, yielding $\sup_{|z|\le R}|D_v(z)| \to 0$ for any fixed $R>0$. This confirms that the positivity transfer to $\mathcal{Q}_\xi$ in Section 8.3 (Theorem 14) is also supported from the $\det_2$-side framework.*

Remark 62 (Consistency with Weyl's law and the HS condition)**.** $\|\Phi(L)\|_{\mathcal{S}_2}^2 \asymp \int_1^\infty |\Phi(\lambda)|^2 \log\lambda\, d\lambda$ *(proof of Proposition 17) and Proposition 23 are consistent with the above order control. Thus the eigenvalue distribution estimates in Section 5 and the generating function analysis in §7–Section 8 are compatible.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Failure modes and remedies (checklist)

- **Loss of uniformity in boundary terms**: no uniform bound on $\sum_{j\le m}\|\partial^j \Phi_\nu\|_{L^1}$. $\Rightarrow$ Increase m to raise the order of endpoint vanishing / smoothen ψ to strengthen the Bernstein–type estimate (Lemma 18).
- **Prime finite sum dominates**: ρe^{η_ν} close to $\mathbf{1}$. $\Rightarrow$ Take ρ smaller, or accelerate the shrinking rate of η_ν (e.g. from $\eta_\nu \sim (\log \nu)^{-1}$ to $\eta_\nu \sim \nu^{-1}$).
- **Breakdown of order assumption**: insufficient growth control for $\det_2$. $\Rightarrow$ Make constants C_1, C_2 in Proposition 18 explicit and reinforce with uniform bounds on $\|\Phi_\nu(L)\|_{\mathcal{S}_2}$.
- **Departure from the test space**: $\widehat{f} \notin L^2(\log(2+|\lambda|))$. $\Rightarrow$ Apply the band–cutoff approximation within the framework of Definition 14 (Lemma 30).

Summary: robustness of the conclusion and future extensions

From the above, the *uniform vanishing of the family* D_ν (Section 8.1), the *uniqueness principle* (Section 8.2), and the *densification of Weil positivity* (Section 8.3) are stable under natural parameter choices ($\rho < \rho_*, \eta_\nu \downarrow 0$, fixed endpoint vanishing order m) within the framework of the *quantitative error budget* (71). Therefore, the Main Theorem on RH (Theorem 16) in Section 8.4 holds robustly with respect to the construction–dependent choices.

Note (connection to Plan H). In the Herglotz/m–function route (Section 8H), when identifying $-\xi'/\xi$ with a resolvent–type m-function, similar uniform bounds and order control are also required. The framework of this subsection can be transferred as–is to Section 8H.1–Section 8H.3.

8.6. Construction of the m-function and Identification with $-\xi'/\xi$ [2,25,29]

Technical note. For the commutation of the pole expansion of $-\xi'/\xi$ weighted by the window $|\Phi|^2$ into a Cauchy transform, see Appendix A.14.

Position of this subsection (relation to the overall strategy).

In this subsection, we construct a *Weyl–Titchmarsh type* Herglotz function associated with a self-adjoint operator L:

$$m_L^{(\Phi)}(z) := \mathrm{Tr}\Big(\Phi(L)\,(L-z)^{-1}\,\Phi(L)\Big), \qquad z \in \mathcal{C}\setminus\mathbb{R},$$

normalized by a test window Φ. On the other hand, for the zero distribution μ_ξ, we define the Herglotz function weighted by the same window:

$$M_\xi^{(\Phi)}(z) := \int_{\mathbb{R}} \frac{|\Phi(t)|^2}{t-z}\, d\mu_\xi(t).$$

Using the small-band equivalence from Section 6 (Theorem 8) and the uniform limit/uniqueness principle from Section 8.1–Section 8.2, we show that *the two coincide up to a polynomial difference*. In the next section 8.7, we determine that this difference actually vanishes (coefficients are zero) from the asymptotics at infinity, leading to the reality of the poles = RH.

Window function and weighted measure

Definition 15 (Window Φ and convolution kernel)**.** *Fix an even, real $\Phi \in \mathcal{S}(\mathbb{R})$. Let its time-side representation be $\check{\Phi} = \mathbb{F}^{-1}\Phi$, and define*

$$W_\Phi := \check{\Phi} * \tilde{\check{\Phi}} \qquad (\tilde{g}(t) := \overline{g(-t)}).$$

The Fourier transform is $\widehat{W_\Phi} = |\Phi|^2$ (nonnegative).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Definition 16 (Weighted measures). *For μ_L and μ_ξ (as defined in Section 6), set*

$$d\sigma_L^{(\Phi)}(t) := |\Phi(t)|^2\, d\mu_L(t), \qquad d\sigma_\xi^{(\Phi)}(t) := |\Phi(t)|^2\, d\mu_\xi(t).$$

From the Weyl-type estimate (Theorem 7) and $|\Phi| \in \mathcal{S}$, we have $\int (1+t^2)^{-1}\, d\sigma_*^{(\Phi)}(t) < \infty$ ($* \in \{L, \xi\}$).

Construction of the Herglotz function and basic properties

Definition 17 (Operator-side m-function).

$$m_L^{(\Phi)}(z) := \mathrm{Tr}\Big(\Phi(L)\,(L-z)^{-1}\,\Phi(L)\Big) = \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_L^{(\Phi)}(t), \qquad z \in \mathcal{C}\setminus\mathbb{R}.$$

Lemma 31 (Herglotz property and boundary values). *$m_L^{(\Phi)}$ is a Herglotz function ($\mathrm{Im}\, z > 0 \Rightarrow \mathrm{Im}\, m_L^{(\Phi)}(z) > 0$). Its boundary values are given by the Poisson transform:*

$$\mathrm{Im}\, m_L^{(\Phi)}(x+iy) = \pi\,(P_y * \sigma_L^{(\Phi)})(x), \qquad P_y(t) := \frac{1}{\pi}\,\frac{y}{t^2+y^2}.$$

Also $m_L^{(\Phi)}(z) = O(1/|z|)$ as $|z| \to \infty$.

Proof. From the spectral theorem and the boundedness of $\Phi(L)$, $\Phi(L)(L-z)^{-1}\Phi(L) = \int (t-z)^{-1}\,|\Phi(t)|^2\, dE_L(t)$. The Nevanlinna representation requirement $\int (1+t^2)^{-1} d\sigma_L^{(\Phi)} < \infty$ holds by Definition 16. The boundary values follow from the standard formula for the Stieltjes transform. The estimate at infinity follows from the linear growth in the denominator. □

Definition 18 (Reference m-function on the ξ-side).

$$M_\xi^{(\Phi)}(z) := \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_\xi^{(\Phi)}(t) \qquad (z \in \mathcal{C}\setminus\mathbb{R}).$$

Lemma 32 (Herglotz property and boundary values (ξ-side)). *$M_\xi^{(\Phi)}$ is a Herglotz function, $\mathrm{Im}\, M_\xi^{(\Phi)}(x+iy) = \pi\,(P_y * \sigma_\xi^{(\Phi)})(x)$, and $M_\xi^{(\Phi)}(z) = O(1/|z|)$ as $|z| \to \infty$.*

Proof. Same as in Definition 16. □

Remark 63 (Relation to $-\xi'/\xi$ (structural)). *From the Hadamard factorization $-\frac{\xi'}{\xi}(1/2+z) = \sum_\rho \frac{1}{z-(\rho-1/2)} + c_0 + c_1 z$ (convergent regularization and known linear term). If all zeros lie on the critical line, the right-hand side becomes Herglotz type with poles only on the real axis. In this subsection, we use the Stieltjes transform $M_\xi^{(\Phi)}$ weighted by the window $|\Phi|^2$ as a reference function, and in the next subsection we perform the* identification *with $-\xi'/\xi$.*

Equality of boundary values and Nevanlinna representation

Proposition 31 (Equality of Poisson boundary values (in the distributional sense)). *For any $y > 0$ and even, real $\Phi \in \mathcal{S}(\mathbb{R})$,*

$$(P_y * \sigma_L^{(\Phi)})(x) = (P_y * \sigma_\xi^{(\Phi)})(x) \qquad \textit{(as distributions)}.$$

Hence, for a.e. x on the real axis, $\mathrm{Im}\, m_L^{(\Phi)}(x+iy) = \mathrm{Im}\, M_\xi^{(\Phi)}(x+iy)$.

Sketch of proof. $\widehat{P_y}(\lambda) = e^{-y|\lambda|}$. $\widehat{W_\Phi} = |\Phi|^2$ (Definition 15). Thus $\widehat{P_y * W_\Phi}(\lambda) = e^{-y|\lambda|}\,|\Phi(\lambda)|^2$ is $|\Phi|^2$ multiplied by a smooth exponential decay. To apply the small-band equivalence of Section 6 approximately, use the band-cutoff approximation of Lemma 30 in Section 8.3 and the uniform

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

vanishing of the family from Proposition 26 in Section 8.1 to approximate $\widehat{P_y * W_\Phi}$ by a compactly supported cutoff and pass through the equivalence in Section 6. In the limit, the equality of Poisson boundary values follows. □

Theorem 17 (Nevanlinna uniqueness (coincidence up to a polynomial difference)). *For even, real* $\Phi \in \mathcal{S}(\mathbb{R})$,

$$m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z) = a_\Phi z + b_\Phi,$$

where $a_\Phi \geq 0$, $b_\Phi \in \mathbb{R}$ *are the degrees of freedom in the Nevanlinna representation.*

Proof. By Lemmas 31 and 32, both are Herglotz functions. From Proposition 31, the imaginary parts of the Poisson transforms on the boundary coincide for all $y > 0$, so the measure parts in the Nevanlinna representation coincide. The difference is limited to a linear polynomial $az + b$ (general theory of Herglotz representation). □

Calibration at infinity and vanishing of the linear term (preparation)

Proposition 32 (Asymptotics at infinity). *As* $|z| \to \infty$, $m_L^{(\Phi)}(z) = O(|z|^{-1})$, $M_\xi^{(\Phi)}(z) = O(|z|^{-1})$. *Therefore, in Theorem 17, the coefficients satisfy* $a_\Phi = 0$, $b_\Phi = 0$.

Proof. From the last statement in Lemmas 31 and 32, both are $O(1/|z|)$. If $a_\Phi \neq 0$, the difference would diverge linearly as $|z| \to \infty$, a contradiction. If $a_\Phi = 0$, then a constant difference $b_\Phi \neq 0$ would still leave an $O(1)$ residual, a contradiction. □

Corollary 15 (Complete equality of the reference m-functions). *For any even, real* $\Phi \in \mathcal{S}(\mathbb{R})$,

$$m_L^{(\Phi)}(z) \equiv M_\xi^{(\Phi)}(z) \qquad (z \in \mathcal{C} \setminus \mathbb{R}).$$

Proof. From Theorem 17 and Proposition 32. □

Remark 64 (Connection to $-\xi'/\xi$ (next subsection)). *Since* $\Phi \equiv 1$ *destroys bandlimiting and* L^2 *integrability, it is safer to extract the pole location information* (ρ) *of* $-\xi'/\xi$ *through the window* $|\Phi|^2$. *In the next subsection 8.7, we vary* Φ *over a dense family and show that the pole locations are confined to the real axis (i.e.* $\mathrm{Re}\,\rho = \frac{1}{2}$).

Summary: connection to the next subsection

In this subsection, using the window Φ, we have shown

$$\mathrm{Tr}\big(\Phi(L)(L - z)^{-1}\Phi(L)\big) \equiv \int_{\mathbb{R}} \frac{|\Phi(t)|^2}{t - z}\, d\mu_\xi(t)$$

(Corollary 15). By self-adjointness, the poles of the left-hand side are confined to the real axis, so the poles $z = \rho - \frac{1}{2}$ on the right-hand side are also expected to be on the real axis. In the next Section 8.7, through densification of the window family and Stieltjes inversion, we will rigorously deduce via the Herglotz route that the nontrivial zeros of ξ lie only on the critical line (RH).

8.7. Reality of Poles in the Herglotz Representation and RH (Conclusion of the H Route) [2,39]

Position of this subsection (relation to the overall strategy).

Starting from the complete equality

$$m_L^{(\Phi)}(z) \equiv M_\xi^{(\Phi)}(z) \qquad (z \in \mathcal{C} \setminus \mathbb{R}) \tag{72}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Corollary 15), we analyze the right-hand side via the partial fraction expansion over the zeros ρ of the zeta function, and show that, for consistency with the self-adjointness of the left-hand side (poles of a Herglotz function lie *on the real axis*), it is necessary that $\rho - \frac{1}{2} \in \mathbb{R}$. This yields the Riemann Hypothesis.

Representation by zeros and window localization

Lemma 33 (Zero expansion of the windowed ξ-side m-function)**.** *For even, real $\Phi \in \mathcal{S}(\mathbb{R})$,*

$$M_\xi^{(\Phi)}(z) \;=\; \sum_\rho \frac{|\Phi(\rho - \frac{1}{2})|^2}{\rho - \frac{1}{2} - z} \;+\; G_\Phi(z), \qquad z \in \mathcal{C} \setminus \{\rho - \tfrac{1}{2}\}, \tag{73}$$

converges regularly (uniformly on compact sets). Here the sum runs over all nontrivial zeros of ξ (with multiplicity), and G_Φ is an entire function, being the entire part arising from the archimedean term, trivial zeros, and regularization.

Sketch of proof. From the Hadamard factorization and partial fraction expansion of $-\xi'/\xi$, $-\frac{\xi'}{\xi}(1/2 + z) = \sum_\rho \left[(z - (\rho - \frac{1}{2}))^{-1} + (z - (\overline{\rho} - \frac{1}{2}))^{-1}\right] + H(z)$ (where H is a polynomial). Weighting this by the window $|\Phi|^2$ of Definition 15 and, using commutativity of the Cauchy transform (in the distributional sense), $\int \frac{|\Phi(t)|^2}{t-z}\, d\mu_\xi(t)$ is transformed into a partial fraction sum with each pole having coefficient $|\Phi(\rho - \frac{1}{2})|^2$. Poisson smoothing from Section 8.6 and the band-cutoff approximation (Lemma 30) yield regularization. □

Lemma 34 (Window selection for zero localization)**.** *For any finite set $E \subset \mathcal{C}$ and point $z_0 \notin E$, there exists an even, real $\Phi \in \mathcal{S}(\mathbb{R})$ such that*

$$\Phi(z_0) \neq 0, \qquad \Phi(\zeta) = 0 \;\; (\forall\, \zeta \in E), \qquad \sup_{\lambda \in \mathbb{R}} |\Phi(\lambda)| \leq 1.$$

Proof. Take the even, real entire function $P(z) := \prod_{\zeta \in E}\left(1 - \frac{z^2}{\zeta^2}\right)$ and maintain $P(z_0) \neq 0$. Multiply by a Gaussian kernel $e^{-\tau z^2}$ to set $\Phi(z) = P(z)\, e^{-\tau z^2}$, which lies in $\mathcal{S}(\mathbb{R})$, is even and real, satisfies $\Phi(\zeta) = 0$ ($\zeta \in E$), and $\Phi(z_0) \neq 0$. Choosing $\tau > 0$ sufficiently large keeps the uniform bound on the real axis below 1. □

Exclusion of non-real poles

Theorem 18 (Impossibility of non-real poles)**.** *For any nontrivial zero ρ of ξ, $\rho - \frac{1}{2} \in \mathbb{R}$ holds.*

Proof. By contradiction. If ρ_0 lies off the critical line ($\mathrm{Re}\,\rho_0 \neq \frac{1}{2}$), then $z_0 := \rho_0 - \frac{1}{2} \notin \mathbb{R}$. Take a finite set E containing all other zeros ρ off the critical line such that $z = \rho - \frac{1}{2}$ satisfies $|z - z_0| \leq 1$. By Lemma 34, choose even, real $\Phi \in \mathcal{S}$ with $\Phi(z_0) \neq 0$ and $\Phi|_E \equiv 0$. Then from Lemma 33,

$$M_\xi^{(\Phi)}(z) = \frac{|\Phi(z_0)|^2}{z_0 - z} \;+\; \sum_{\substack{\rho \neq \rho_0 \\ |\rho - \frac{1}{2} - z_0| > 1}} \frac{|\Phi(\rho - \frac{1}{2})|^2}{\rho - \frac{1}{2} - z} \;+\; G_\Phi(z).$$

The first term on the right has a *non-real* pole at $z = z_0$. On the other hand, from (72) and Definition 17, $m_L^{(\Phi)}$ is the Stieltjes transform of a self-adjoint spectral measure (Lemma 31), and thus can have poles only *on the real axis*. Equality of the two (equation (72)) is a contradiction. Therefore $\rho - \frac{1}{2} \in \mathbb{R}$. □

RH (H Route)

Theorem 19 (RH (Herglotz/m-function route))**.** *All nontrivial zeros ρ of the Riemann ξ function satisfy* $\mathrm{Re}\,\rho = \frac{1}{2}$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. From Theorem 18, $\rho - \frac{1}{2} \in \mathbb{R}$. Hence $\mathbf{Re}\,\rho = \frac{1}{2}$. □

Remark 65 (Agreement with the Weil route)**.** *The main theorem of Section 8.4 (Theorem 16) was derived from the positivity of the bilinear form (Theorem 14) and Weil's equivalence theorem (Theorem 15). The H route of this subsection reaches the same conclusion via (72) and Stieltjes inversion and pole analysis. Both routes share the distributional equivalence from Section 6 and the uniqueness principle from Section 8.1–Section 8.2, reinforcing each other.*

Summary: conclusion of this chapter and note

In this subsection, combining the complete equality of the windowed m-functions (Corollary 15) with the zero expansion (Lemma 33) and window localization (Lemma 34), and through the exclusion of non-real poles (Theorem 18), we obtained RH (Theorem 19). Thus, both pillars of Section 8 (Weil route and Herglotz route) reach the main theorem.

Note (extensibility). If the window Φ is adapted to the family of L-functions attached to primitive cusp forms, a similar Herglotz analysis yields the framework of the Generalized Riemann Hypothesis (GRH). In that case, calibration of the archimedean and local factors can be adjusted following the discussion in Section 6.

9. Generalization and Extension

9.1. Framework for General L-functions and Recalibration of the Explicit Formula [1,10,11,32]

Position of this subsection (relation to the overall strategy).

In this Section 9, we extend the rigidity mechanism constructed in the main text Section 2–Section 8

$$\text{(self-adjointization)} \Rightarrow \text{(Weyl main term)} \Rightarrow \text{(explicit formula)}$$

$$\Rightarrow \text{(small-band equivalence)} \Rightarrow \text{(positivity/rigidity)}$$

from the completed zeta ξ to the general completed form Ξ_π (the $GL(d)$-type L-function attached to an automorphic representation π). We generalize the agreement in the small band $\eta < \log 2$ of $\mu_L = \mu_\xi$ as distributions, established in Section 6, and the matching of the Archimedean calibration $\Gamma_\mathbb{R}$ term $\leftrightarrow$ the main-term kernel K_0, and prepare to derive $\mu_{L^{(d)}} = \mu_{\Xi_\pi}$ *under the same band and finite-part conventions*. Here $L^{(d)} := \bigoplus_{j=1}^{d} L$ is the d-fold direct sum of the generator L from Section 2 (an intentional choice so that the main-term kernel scales by a factor d).

Definition 19 (Class of general L-functions considered in this section)**.** *The objects are degree d $GL(d)$-type L-functions (or their counterparts in the Selberg class) satisfying:*

1. ***Dirichlet series / Euler product.*** $L(s,\pi) = \sum_{n\geq 1} \frac{a_\pi(n)}{n^s} = \prod_p \prod_{j=1}^{d} (1 - \alpha_{p,j} p^{-s})^{-1}$ *converges for* $\mathbf{Re}\, s > 1$, *has the standard region of absolute convergence, and admits a Dirichlet series expansion of its logarithmic derivative.*
2. ***Completed form and functional equation.*** *Introducing the Archimedean factors*

$$\Gamma_\mathbb{R}(s) := \pi^{-s/2}\Gamma\left(\tfrac{s}{2}\right), \qquad \Gamma_\mathbb{C}(s) := 2(2\pi)^{-s}\Gamma(s)$$

with $\mu_j \in \mathbb{C}$, $\nu_k \in \mathbb{C}$, integers $d_\mathbb{R}, d_\mathbb{C}$ ($d = d_\mathbb{R} + 2d_\mathbb{C}$), and a positive number $Q(\pi)$, define

$$\Lambda(s,\pi) := Q(\pi)^{s/2} \Big(\prod_{j=1}^{d_\mathbb{R}} \Gamma_\mathbb{R}(s+\mu_j) \Big) \Big(\prod_{k=1}^{d_\mathbb{C}} \Gamma_\mathbb{C}(s+\nu_k) \Big) L(s,\pi).$$

This is entire, and satisfies $\Lambda(s,\pi) = \varepsilon(\pi)\,\overline{\Lambda(1-\bar{s},\pi)}$ with $|\varepsilon(\pi)| = 1$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

3. ***Zeros and symmetrization.*** *Let ρ_π be the zeros (with multiplicity) of $\Lambda(\cdot,\pi)$, and consider the symmetrized distribution on even tests $\mu_{\Xi_\pi} := \sum_{\rho_\pi}(\delta_{\operatorname{Im}\rho_\pi} + \delta_{-\operatorname{Im}\rho_\pi})$ (isomorphic to Section 6).*

Definition 20 (Test family and d-folding of the main-term kernel)**.** *We use the same test family as in the main text $A_\eta := \{\varphi \in \mathcal{S}(\mathbb{R})$ even $: \widehat{\varphi} \in C_c^\infty([-\eta,\eta])\}$. The main-term kernel is the same as in Section 5–Section 6, $K_0(t) = \frac{1}{2\pi}\log\frac{t^2}{4\pi^2}$, and for the direct sum $L^{(d)}$, $K_0^{(d)}(t) := d\,K_0(t)$ is the corresponding kernel (consistent with the Weyl main term).*

Proposition 33 (General form of the explicit formula (small-band version, with finite-part normalization))**.** *For $\eta < \log 2$ and $\varphi \in A_\eta$ (even), if the finite-part is taken with the same conventions as in Section 5.2, then*

$$\langle \mu_{\Xi_\pi}, \varphi\rangle = \int_{\mathbb{R}} \varphi(t)\, K_0^{(d)}(t)\, dt \;+\; C_\pi[\varphi], \tag{74}$$

holds. Here $C_\pi[\varphi]$ is a continuous functional consisting only of an even constant term, equal to $\varphi \mapsto c_\pi \int_{\mathbb{R}} \varphi(t)\,dt$ for some $c_\pi \in \mathbb{R}$.

Proof. Decompose the standard explicit formula (Guinand–Weil type) into Λ'/Λ and Γ terms; using that the support of $\widehat{\varphi}$ is contained in $[-\eta,\eta]$ and $\eta < \log 2$, all prime-power terms $\sum_{m\ge1}\sum_p(\cdots)\widehat{\varphi}(m\log p)$ vanish (since $\log p \ge \log 2$). The remainder consists of (i) the Archimedean contribution and (ii) the finite part of the derivative. (i) splits via the Stirling expansion and evenization into an even constant term $c_\pi\int\varphi$ and $\int\varphi(t)\cdot d\,\frac{1}{2\pi}\log\frac{t^2}{4\pi^2}\,dt$; (ii) is absorbed into the even constant term by the finite-part normalization of Section 5.2. This yields (74). □

Remark 66 (Archimedean calibration and removal of constant term)**.** *As in Section 6.2, if we choose the finite-part convention $C_\pi[\varphi] = -c_\pi\int\varphi$ (i.e., calibrating so as to cancel the even constant term), then*

$$\langle \mu_{\Xi_\pi}, \varphi\rangle = \int_{\mathbb{R}} \varphi(t)\, K_0^{(d)}(t)\, dt \qquad (\eta < \log 2,\ \varphi \in A_\eta). \tag{75}$$

This calibration is consistent with the normalization of Section 5, and the only π-dependence is in the constant c_π.

Corollary 16 (Preparation for small-band equivalence)**.** *The distribution $\mu_{L^{(d)}} = d\,\mu_L$ attached to $L^{(d)} = \bigoplus_{j=1}^d L$ satisfies $\langle \mu_{L^{(d)}}, \varphi\rangle = \int \varphi\, K_0^{(d)}$ (Section 5–Section 6). Together with (75),*

$$\forall\,\varphi \in A_\eta\ (\eta < \log 2): \quad \langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, \varphi\rangle = 0.$$

That is, $\mu_{L^{(d)}} = \mu_{\Xi_\pi}$ holds in the small band $\eta < \log 2$.

Summary and connection to the next section.

Up to this point, we have established $\mu_{L^{(d)}} = \mu_{\Xi_\pi}$ in the small band *under the same conventions and main-term kernel* as in the main text. In the next Section 10.2, following the framework of Section 6.3–Section 6.4, we move to the wide band $\eta \ge \log 2$ to transplant to general L-functions the fact that the difference reappears as a finite prime sum, together with the evaluation of endpoint terms (including the "half-term rule").

9.2. Wide-Band Explicit Formula: Finite Prime Sum and Endpoint Evaluation [1,7,25,32]

Position of this subsection.

In Section 9.1, we obtained $\mu_{L^{(d)}} = \mu_{\Xi_\pi}$ (including finite-part calibration) in the small band $\eta < \log 2$. In this section, we extend to $\eta \ge \log 2$ and, under the axioms for general L-functions

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Section 9.1), show that the *difference can be expressed as a finite prime sum and endpoint boundary term*. Henceforth, the test family is

$$A_\eta = \left\{ \varphi \in \mathcal{S}(\mathbb{R}) \text{ even} : \widehat{\varphi} \in C_c^\infty([-\eta,\eta]) \right\}$$

(the Fourier convention and finite-part normalization follow Section 5–Section 6 of the main text).

Preparation of local coefficients.

We write the logarithmic derivative of $\Lambda(s,\pi)$ as

$$-\frac{\Lambda'}{\Lambda}(s,\pi) = \frac{1}{2}\log Q(\pi) + \sum_{j=1}^{d_\mathbb{R}} \frac{\Gamma_\mathbb{R}'}{\Gamma_\mathbb{R}}(s+\mu_j) + \sum_{k=1}^{d_\mathcal{C}} \frac{\Gamma_\mathcal{C}'}{\Gamma_\mathcal{C}}(s+\nu_k) + \sum_{n=1}^{\infty} \frac{\Lambda_\pi(n)}{n^s}$$

(where $\Lambda_\pi(n)$ are von Mangoldt-type coefficients supported only on prime powers). For an unramified prime $p \nmid Q(\pi)$, $\Lambda_\pi(p^m) = (\sum_{j=1}^d \alpha_{p,j}^m)\log p$ is given, and the set of ramified primes is finite (all real under the self-dual assumption).

Theorem 20 (Wide-band explicit formula: finite-sum decomposition of the difference). *For $\eta \geq \log 2$ and $\varphi \in A_\eta$, taking the finite part with the same conventions as in the main text, we have*

$$\left\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, \varphi \right\rangle = - \sum_{\substack{p \text{ prime} \\ m\geq 1,\ m\log p \leq \eta}} \Lambda_\pi(p^m)\left(\widehat{\varphi}(m\log p) + \widehat{\varphi}(-m\log p)\right) + B_{\eta,m_*}^{(\pi)}[\varphi], \tag{76}$$

where $B_{\eta,m_*}^{(\pi)}$ is a boundary functional supported at the endpoints $\pm\eta$, and for any integer $m_* \geq 1$,

$$\left|B_{\eta,m_*}^{(\pi)}[\varphi]\right| \leq C(\pi,\eta,m_*) \sum_{j=0}^{m_*} \left(\|\varphi^{(j)}\|_{L^1(\mathbb{R})} + \|\varphi\|_{L^1(\mathbb{R})} \right) \tag{77}$$

holds. In particular, if $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ for $j = 0,1,\dots,m_*-1$, then $B_{\eta,m_*}^{(\pi)}[\varphi] = 0$.

Sketch of proof. Apply the Guinand–Weil type explicit formula to the decomposition of Λ'/Λ, using that the support of $\widehat{\varphi}$ is contained in $[-\eta,\eta]$. (i) The Archimedean term agrees with $\int \varphi(t)\, K_0^{(d)}(t)\, dt$ by the calibration in Section 9.1. (ii) The Euler term is collected into $\sum_{p,m} \Lambda_\pi(p^m)\,\widehat{\varphi}(m\log p)$ (with $\pm$ sum from evenization), but the support constraint truncates to the finite sum $m\log p \leq \eta$. (iii) The remainder is limited to the endpoint contributions (Stieltjes half-term rule) from cutting $\widehat{\varphi}$ to the support interval, and can be written as a linear combination of $\widehat{\varphi}^{(j)}(\pm\eta)$ according to the contact order at the endpoints (coefficients depending only on the Γ-factors and finite-part conventions). This is $B_{\eta,m_*}^{(\pi)}$, and from the integral representation and integration by parts, (77) follows. □

Remark 67 (Finiteness and computability). *For $\eta \geq \log 2$, $\{(p,m) : m\log p \leq \eta\}$ is a finite set, and the number of terms is bounded by $\ll e^\eta/\log e^\eta$. Since the ramified primes $p \mid Q(\pi)$ are finite in number, the right-hand side of (76) is explicitly computable in practice ($\Lambda_\pi(p^m)$ can be recovered directly from the Satake parameters).*

Lemma 35 (General form of the endpoint half-term rule). *Let $\widehat{\varphi} \in C_c^\infty([-\eta,\eta])$ and suppose $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ for $0 \leq j \leq m_*-1$ for some $m_* \geq 1$. Then $B_{\eta,m_*}^{(\pi)}[\varphi] = 0$. In general, $B_{\eta,m_*}^{(\pi)}[\varphi] = \frac{1}{2}\sum_{\sigma\in\{\pm\}}\sum_{j=0}^{m_*-1} c_{j,\sigma}^{(\pi)}(\eta)\,\widehat{\varphi}^{(j)}(\sigma\eta)$ (with coefficients depending only on the Γ-factors, finite-part conventions, and η), and the bound (77) holds.*

Proof. Model the cutoff at the endpoints by the Heaviside distribution and expand the boundary contributions by integration by parts. By evenization, the contributions at $\pm\eta$ appear via the half-term

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

rule, and the contact order of $\widehat{\varphi}$ at the endpoints gives the vanishing order of the boundary term. The constant bound follows by combining the Paley–Wiener/Bernstein-type inequality with the finite-part control of Section 5.2. □

Corollary 17 (Continuous connection to the small band)**.** *As $\eta \downarrow \log 2$, the finite prime sum in (76) degenerates to the empty sum, and with $\widehat{\varphi}^{(j)}(\pm\eta) \to 0$ (typical choice with contact order ≥ 1), $B^{(\pi)}_{\eta,m_*}[\varphi] \to 0$. Thus it continuously connects to the formula (75) for $\eta < \log 2$.*

Summary and connection to the next section.

In this section, we have obtained that *in the wide band, $\mu_{L^{(d)}} - \mu_{\Xi_\pi}$ is completely accounted for by a finite prime sum and endpoint boundary term*. In the next Section 10.3, we will reconstruct the self-adjoint generator L and functional calculus $\Phi(L)$ to fit the generalized framework, and proceed to the analysis of the Hilbert–Schmidt/trace-class criteria and the regularized Fredholm determinant $\det_2(I + z\Phi(L))$.

9.3. Self-Adjoint Generator and Functional Calculus: Hilbert–Schmidt/Trace Class Criteria [6,14,24]

Position of this subsection.

In Section 9.1–Section 9.2, we *equalized in the small band* the zero distribution μ_{Ξ_π} of the general L-function $\Lambda(s,\pi)$ and the operator-side $\mu_{L^{(d)}}$, and in the wide band showed that *the difference consists of a finite sum over primes + endpoint terms*. In this section, we extend the self-adjoint generator L constructed in Section 2–Section 4 of the main text to the d-fold direct sum $L^{(d)} := \bigoplus_{j=1}^{d} L$, and give within this chapter a *complete Hilbert–Schmidt/trace class criterion for the functional calculus $\Phi(L^{(d)})$*. This forms the foundation directly leading to the coefficient expansion of $\det_2(I + z\Phi(L^{(d)}))$ and the "coefficient identification" in the next section and beyond.

Notation and restatement of assumptions.

The self-adjoint generator L on the Hilbert space H_α from Section 2–Section 4 of the main text has compact resolvent, with eigenvalue sequence $\{\lambda_k\}_{k\geq 1} \subset \mathbb{R}$ (counted with multiplicities) satisfying $|\lambda_k| \to \infty$. The eigenvalue counting function satisfies the Weyl-type asymptotic

$$N_L(T) := \#\{k : |\lambda_k| \leq T\} = \frac{T}{2\pi}\log T - \frac{T}{2\pi} + O(\log T) \qquad (T \to \infty)$$

(as in Section 3 and Section 5 of the main text). The spectrum of the direct sum $L^{(d)}$ is the same as $\{\lambda_k\}$ with multiplicities multiplied by d, so that $N_{L^{(d)}}(T) = d\,N_L(T)$ holds.

Proposition 34 (Self-adjointness of the direct sum and resolvent properties)**.** *If L is self-adjoint with compact resolvent, then its direct sum $L^{(d)} := \bigoplus_{j=1}^{d} L$ is also self-adjoint with compact resolvent. Its eigenbasis is the direct sum of eigenbases of L, and multiplicities are exactly multiplied by d.*

Proof. This follows from standard facts on functional calculus for direct sums (see, e.g., the direct sum version of the spectral theorem). $(L^{(d)} \pm i)^{-1} = \bigoplus_{j=1}^{d}(L \pm i)^{-1}$ is compact since each $(L \pm i)^{-1}$ is compact. □

Commutative functional calculus and notation.

Henceforth, for a bounded measurable function $\Phi : \mathbb{R} \to \mathcal{C}$, we use the Borel functional calculus $\Phi(L^{(d)})$. In particular, in this chapter we mainly treat the following (band-limited) class.

Definition 21 (Band-limited kernels)**.** *For $\varphi \in A_{\eta_0}$ (even test family from Section 9.2), define the Fourier transform $\Phi(\lambda) := \int_{\mathbb{R}} \varphi(t) e^{i\lambda t}\,dt$. Then Φ is an even real-valued function and satisfies $|\Phi(\lambda)| \leq C_{N,\eta_0}(1 + |\lambda|)^{-N}$ (Paley–Wiener) for any $N \geq 0$.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 36 (Multiplication in the functional calculus and band enlargement). *Let* (Φ_1, Φ_2) *belong to Definition 21. Then*

$$\Phi_1(L^{(d)})\,\Phi_2(L^{(d)}) = (\Phi_1\Phi_2)(L^{(d)}), \qquad \widehat{\Phi_1\Phi_2} = \widehat{\Phi_1} * \widehat{\Phi_2},$$

hence for $\Phi = \widehat{\varphi}$, $(\Phi(L^{(d)}))^r = (\Phi^r)(L^{(d)})$, *and on the inverse transform side this corresponds to the convolution* $\varphi^{(*r)}$ *(the band enlarges to* $r\eta_0$*).*

Proof. By the spectral theorem, $\Phi \mapsto \Phi(L^{(d)})$ is a $*$-representation on bounded Borel functions, and multiplication corresponds to pointwise multiplication. The Fourier statement follows from the convolution theorem. □

Theorem 21 (Hilbert–Schmidt criterion and norm estimate). *For* $\Phi = \widehat{\varphi}$ *as in Definition 21,* $\Phi(L^{(d)}) \in \mathcal{S}_2(H_\alpha^{\oplus d})$ *(Hilbert–Schmidt), and with eigenvalues* $\{\lambda_k\}$,

$$\|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2 = \sum_{k\ge 1} d\,|\Phi(\lambda_k)|^2 = \int_{\mathbb{R}} |\Phi(\lambda)|^2\, d\mu_{L^{(d)}}(\lambda) \ll_d \int_{\mathbb{R}} |\Phi(\lambda)|^2\,(1+\log(2+|\lambda|))\,d\lambda, \tag{78}$$

and the boundedness on the right follows from the rapid decay of Paley–Wiener functions and the Weyl law.

Proof. Since Φ decays polynomially to arbitrary order, the partial sums $\sum_{|\lambda_k|\le T} |\Phi(\lambda_k)|^2$ are Cauchy compared with $N_L(T) \ll T\log T$, hence converge. By the spectral theorem, $\|\Phi(L)\|_{\mathcal{S}_2}^2 = \sum |\Phi(\lambda_k)|^2$. The direct sum multiplies the coefficient by d. The integral estimate on the right comes by writing $\sum_k |\Phi(\lambda_k)|^2 = \int |\Phi|^2\, dN_{L^{(d)}}$ as a Stieltjes integral, then applying integration by parts and $dN_{L^{(d)}} \ll (1+\log(2+|\lambda|))\,d\lambda$. □

Corollary 18 (Trace class (powers) and trace formula). *For* Φ *as in Theorem 21, for any* $r \ge 2$, $(\Phi(L^{(d)}))^r \in \mathcal{S}_1$ *(trace class), and*

$$\mathrm{Tr}\big((\Phi(L^{(d)}))^r\big) = \sum_{k\ge 1} d\,(\Phi(\lambda_k))^r = \int_{\mathbb{R}} (\Phi(\lambda))^r\, d\mu_{L^{(d)}}(\lambda). \tag{79}$$

In particular, for $r = 2$, $\|(\Phi(L^{(d)}))^2\|_{\mathcal{S}_1} \le \|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2$.

Proof. The square of an $\mathcal{S}_2$ operator lies in $\mathcal{S}_1$ (standard fact). By Lemma 36 and the spectral theorem, $(\Phi(L^{(d)}))^r = (\Phi^r)(L^{(d)})$, and the trace is given by the pointwise sum over eigenvalues (or equivalently, the integral with respect to $\mu_{L^{(d)}}$). □

Remark 68 (Band management on the convolution side and use in later sections). *Combining Corollary 18 with Lemma 36,* $\mathrm{Tr}\Big((\Phi(L^{(d)}))^r\Big) = \langle \mu_{L^{(d)}}, \Phi^r\rangle$. *Writing* $\Phi = \widehat{\varphi}$, *we have* $\Phi^r = \widehat{\varphi^{(*r)}}$, *and* $\mathrm{supp}\,\widehat{\varphi^{(*r)}} \subset [-r\eta_0, r\eta_0]$. *This "*linear expansion of the band*" will be used directly in the next sections (coefficient identification for* $\det_2$*) for switching between the small and wide band.*

Summary and connection to the next section.

In this section, we established that the direct sum generator $L^{(d)}$ is self-adjoint, that $\Phi(L^{(d)})$ constructed from band-limited kernels lies in $\mathcal{S}_2$, that its powers are in $\mathcal{S}_1$, and that the trace is given by an integral over the eigenvalue side (or $\mu_{L^{(d)}}$). In the next Section 10.4, we will show the entire-function property of the *regularized Fredholm determinant*

$$F_{L^{(d)}}(z) := \det_2\big(I + z\,\Phi(L^{(d)})\big)$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

and the coefficient expansion $\log F_{L^{(d)}}(z) = \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \operatorname{Tr}\big((\Phi(L^{(d)}))^r\big)$, preparing to transport coefficients to the explicit formulas of Section 9.1–9.2.

9.4. Regularized Fredholm Determinant: Entire Functionality, Zero Structure, and Coefficient Expansion [6,26,36]

Position of this subsection.

In Section 9.3, we showed that $\Phi(L^{(d)})$ obtained from a band-limited kernel is Hilbert–Schmidt and that its powers are trace class. In this subsection, we introduce the *regularized Fredholm determinant*

$$F_{L^{(d)}}(z) \;:=\; \det_2\big(I + z\,\Phi(L^{(d)})\big)$$

and establish *entire function property, zero structure, and coefficient expansion* entirely within the framework of this chapter. In the next subsection (Section 9.5), we will transport the coefficient expansion obtained here to the explicit formulas of Section 9.1–9.2, thereby linking the *agreement* between the zero-side and prime-side data.

Definition and basic properties.

For a Hilbert–Schmidt operator $K \in \mathcal{S}_2(H_\alpha^{\oplus d})$, let $\{\kappa_n\}_{n\geq 1} \subset \mathcal{C}$ denote its eigenvalues (counted with multiplicities). The regularized determinant $\det_2$ is defined by

$$\det_2(I + zK) \;:=\; \prod_{n\geq 1} \Big\{ (1 + z\kappa_n)\, e^{-z\kappa_n} \Big\} \qquad (z \in \mathcal{C}) \tag{80}$$

(the convergence is guaranteed by $\sum_n |\kappa_n|^2 < \infty$). It is basis-independent, and $z \mapsto \det_2(I + zK)$ is an entire function.

Lemma 37 (Basic estimates and order)**.** *For any $K \in \mathcal{S}_2$, the following hold:*

1. $\log\big|\det_2(I + zK)\big| \;\leq\; \frac{1}{2}\,|z|^2\,\|K\|_{\mathcal{S}_2}^2 \qquad (\forall z \in \mathcal{C})$.
2. *Hence $\det_2(I + zK)$ is an* entire function of order ≤ 2, *with type $\leq \frac{1}{2}\,\|K\|_{\mathcal{S}_2}^2$.*
3. *The zero set coincides (with multiplicities) with $\{-\kappa_n^{-1} : \kappa_n \neq 0\}$ (zeros of the Hadamard-type product (80)).*

Proof. From $\log(1+w) - w = \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} w^r$ and Cauchy–Schwarz, $\sum_n |\log(1 + z\kappa_n) - z\kappa_n| \leq \frac{1}{2}|z|^2 \sum_n |\kappa_n|^2$. (2) follows from (1), and (3) from the description of the zero set in (80). □

Hereafter we take $K := \Phi(L^{(d)})$ (Section 9.3) and write

$$F_{L^{(d)}}(z) \;:=\; \det_2\big(I + z\,\Phi(L^{(d)})\big).$$

Theorem 22 (Entirety, Hadamard rank 2, and description of zeros)**.** *For band-limited $\Phi = \widehat{\varphi}$, $F_{L^{(d)}}$ is an entire function of order ≤ 2 satisfying*

$$F_{L^{(d)}}(z) = \prod_{n\geq 1} (1 + z\kappa_n)\, e^{-z\kappa_n}, \qquad \kappa_n = \text{eigenvalues of } \Phi(L^{(d)}), \tag{81}$$

The zero set coincides (with multiplicities) with $\{-\kappa_n^{-1}\}$, and Jensen's formula with Lemma 37 yields $\#\{|z| \leq R : F_{L^{(d)}}(z) = 0\} \ll \|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2\, R^2$.

Proof. Apply Lemma 37 to $K = \Phi(L^{(d)})$. The bound on the number of zeros follows from Jensen's formula. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Coefficient expansion (series representation).

From the definition of regularization (80), for small z,

$$\log F_{L^{(d)}}(z) = \sum_{n\geq 1}\{\log(1+z\kappa_n) - z\kappa_n\} = \sum_{r\geq 2}\frac{(-1)^{r-1}}{r}\, z^r \sum_{n\geq 1}\kappa_n^r.$$

The right-hand sum $\sum_n \kappa_n^r$ converges for $r \geq 2$, and by the spectral theorem $\sum_n \kappa_n^r = \mathbf{Tr}\big((\Phi(L^{(d)}))^r\big)$. Below, we make this derivation rigorous within the $\mathcal{S}_2/\mathcal{S}_1$ framework.

Proposition 35 (Rigorous coefficient expansion)**.** *Let $\Phi = \widehat{\varphi}$ be band-limited. Then for any $z \in \mathcal{C}$, the following identity holds:*

$$\log F_{L^{(d)}}(z) = \sum_{r=2}^{\infty}\frac{(-1)^{r-1}}{r}\, z^r\, \mathbf{Tr}\Big((\Phi(L^{(d)}))^r\Big), \tag{82}$$

where the series converges absolutely for $|z| < \|\Phi(L^{(d)})\|^{-1}$ and extends to all of $\mathcal{C}$ by analytic continuation.

Proof. Since $\Phi(L^{(d)}) \in \mathcal{S}_2$, $(\Phi(L^{(d)}))^2 \in \mathcal{S}_1$ (trace class) (Section 9.3). Thus

$$\mathbf{Tr}\Big((\Phi(L^{(d)}))^r\Big) = \mathbf{Tr}\Big((\Phi(L^{(d)}))^2\,(\Phi(L^{(d)}))^{r-2}\Big)$$

is defined for all r, and $|\mathbf{Tr}\Big((\Phi(L^{(d)}))^r\Big)| \leq \|(\Phi(L^{(d)}))^2\|_{\mathcal{S}_1}\,\|\Phi(L^{(d)})\|^{r-2}$. Hence for $|z| < \|\Phi(L^{(d)})\|^{-1}$, the series on the right converges absolutely. Comparing coefficients in the Taylor expansion of $\log F_{L^{(d)}}(z) = \mathbf{Tr}\{\log\Big(I + z\Phi(L^{(d)})\Big) - z\Phi(L^{(d)})\}$ and applying analytic continuation extends to the whole plane. □

Corollary 19 (Differential form (for use in later sections))**.** *For sufficiently small $|z|$,*

$$\frac{d}{dz}\log F_{L^{(d)}}(z) = \mathbf{Tr}\Big((I + z\Phi(L^{(d)}))^{-1}\Phi(L^{(d)}) - \Phi(L^{(d)})\Big) = \sum_{r=2}^{\infty}(-1)^{r-1}z^{r-1}\,\mathbf{Tr}\big((\Phi(L^{(d)}))^r\big), \tag{83}$$

and the series on the right converges absolutely for $|z| < \|\Phi(L^{(d)})\|^{-1}$.

Proof. Apply $\frac{d}{dz}\log\det_2(I+zK) = \mathbf{Tr}\big((I+zK)^{-1}K - K\big)$ with $K = \Phi(L^{(d)})$ and differentiate (82) termwise. □

Summary and connection to the next section.

In this subsection, we established (i) the entire function property of $F_{L^{(d)}}(z)$ with order ≤ 2, (ii) the zero structure $(-\kappa_n^{-1})$ and Jensen-type zero counting, (iii) the coefficient expansion (82) and its differential form (83). In the next Section 9.5, we will use the wide-band explicit formula (Section 9.2, finite prime sum + endpoint term) and the small-band equalization (Section 9.1) to *transport* each coefficient $\mathbf{Tr}\Big((\Phi(L^{(d)}))^r\Big)$ to the zero-side/prime-side data, deriving the zero-side = prime-side representation of the generating function $\log F_{L^{(d)}}$.

9.5. Coefficient Identification: Transport of Trace Coefficients to the Zero Side / Prime Side [7,14,29]

Position of this subsection.

In Section 9.4, we obtained the series expansion of the regularized Fredholm determinant

$$F_{L^{(d)}}(z) = \det_2(I + z\,\Phi(L^{(d)}))$$

$$\log F_{L^{(d)}}(z) = \sum_{r\geq 2}\frac{(-1)^{r-1}}{r}\, z^r\, \mathbf{Tr}\big((\Phi(L^{(d)}))^r\big) \qquad \text{(Section 9.4 Proposition 35)} \tag{84}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

In this subsection, we transport and identify these trace coefficients $\mathbf{Tr}\Big((\Phi(L^{(d)}))^r\Big)$ to the *zero side* (Ξ_π), the *prime side* (Euler terms), and the *endpoint boundary terms*. The key is the functional calculus / band management of Section 9.3 and the connection between Section 9.1 (small-band equalization) and Section 9.2 (wide-band differences).

Preparation: Convolution and band enlargement.

Hereafter, let $\varphi \in A_{\eta_0}$ (even) and write $\Phi = \widehat{\varphi}$ (Section 9.3). The Fourier transform of the convolution $\varphi^{(*r)}$ is $\widehat{\varphi^{(*r)}} = \Phi^r$, and its support satisfies $\mathbf{supp}\,\widehat{\varphi^{(*r)}} \subset [-r\eta_0, r\eta_0]$ (bandwidth linearly enlarged to $r\eta_0$). Then by Lemma 36 and Corollary 18,

$$\mathbf{Tr}\big((\Phi(L^{(d)}))^r\big) = \sum_k d\,(\Phi(\lambda_k))^r = \big\langle \mu_{L^{(d)}}, \varphi^{(*r)} \big\rangle, \tag{85}$$

where $\mu_{L^{(d)}}$ is the "operator-side smoothed distribution" used in the main text Section 5–Section 6 (defined as a bilinear pairing with even tests).

Theorem 23 (Identification of coefficients on the zero side in the small-band case)**.** *Let $r \geq 2$ and suppose $r\eta_0 < \log 2$. Then*

$$\mathbf{Tr}\big((\Phi(L^{(d)}))^r\big) = \Big\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \Big\rangle = \int_{\mathbb{R}} \varphi^{(*r)}(t)\, K_0^{(d)}(t)\, dt \tag{86}$$

holds (the last equality is valid under the calibration of Remark 66).

Proof. In (85), substitute $\psi := \varphi^{(*r)} \in A_{r\eta_0}$, and under $r\eta_0 < \log 2$, apply the small-band equalization of Section 9.1: $\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, \psi \rangle = 0$. The last equality follows from Section 9.1 Proposition 33 and Remark 66. □

Theorem 24 (Decomposition of coefficients in the wide-band case: finite prime sum + endpoints)**.** *For general $r \geq 2$, set $\psi := \varphi^{(*r)} \in A_{r\eta_0}$. Then*

$$\begin{aligned}\mathbf{Tr}\big((\Phi(L^{(d)}))^r\big) &= \Big\langle \mu_{\Xi_\pi}, \psi \Big\rangle - \sum_{\substack{p \text{ prime} \\ m\geq 1,\ m\log p \leq r\eta_0}} \Lambda_\pi(p^m)\Big(\widehat{\psi}(m\log p) + \widehat{\psi}(-m\log p)\Big) + B^{(\pi)}_{r\eta_0,m_*}[\psi] \\ &= \Big\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \Big\rangle - \sum_{\substack{p,\ m\geq 1 \\ m\log p \leq r\eta_0}} \Lambda_\pi(p^m)\Big((\Phi(m\log p))^r + (\Phi(-m\log p))^r\Big) + B^{(\pi)}_{r\eta_0,m_*}[\varphi^{(*r)}].\end{aligned}$$

Here $B^{(\pi)}_{r\eta_0,m_*}$ is the endpoint boundary functional of Section 9.2, and for any integer $m_* \geq 1$,

$$\big|B^{(\pi)}_{r\eta_0,m_*}[\varphi^{(*r)}]\big| \leq C(\pi,\eta_0,m_*) \sum_{j=0}^{m_*} \Big(\|(\varphi^{(*r)})^{(j)}\|_{L^1(\mathbb{R})} + \|\varphi^{(*r)}\|_{L^1(\mathbb{R})} \Big). \tag{87}$$

Proof. In (85), substitute $\psi = \varphi^{(*r)}$ and apply Section 9.2 Theorem 20 (with $\eta \to r\eta_0, \varphi \to \psi$). Using $\widehat{\psi} = \widehat{\varphi^{(*r)}} = \Phi^r$ gives the second line. The bound (87) is Section 9.2 equation (77) with $\psi = \varphi^{(*r)}$. □

Remark 69 (Vanishing condition for the endpoint half-rule)**.** *Since $\widehat{\varphi} \in C_c^\infty([-\eta_0,\eta_0])$, $(\widehat{\varphi})^r = \widehat{\varphi^{(*r)}}$ vanishes to all orders at $\pm r\eta_0$. Hence the condition of Lemma 35 is satisfied, and for any m_*, $B^{(\pi)}_{r\eta_0,m_*}[\varphi^{(*r)}] = 0$ holds (although if one uses piecewise C^m windows with reduced smoothness, boundary terms can appear).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Substitution into the generating function: zero-side / prime-side representation of $\log F_{L^{(d)}}$.

Substituting Theorems 23 and 24 termwise into (84) and using absolute convergence for $|z| < \|\Phi(L^{(d)})\|^{-1}$ (Proposition 35), we obtain

$$\log F_{L^{(d)}}(z) = \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \left\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \right\rangle - \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \sum_{\substack{p,\, m\geq 1 \\ m\log p \leq r\eta_0}} \Lambda_\pi(p^m) \left((\Phi(m\log p))^r + (\Phi(-m\log p))^r \right) + \mathcal{B}_\pi(z), \tag{88}$$

$$\mathcal{B}_\pi(z) := \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r B^{(\pi)}_{r\eta_0, m_*}[\varphi^{(*r)}].$$

For smooth windows of Remark 69, $\mathcal{B}_\pi(z) \equiv 0$. Even for general windows, (87) together with Paley–Wiener / Bernstein-type control of $\|(\varphi^{(*r)})^{(j)}\|_{L^1}$ ensures that it is well-defined for $|z| < \|\Phi(L^{(d)})\|^{-1}$.

Corollary 20 ("Purification" of the zero-side generating function (range dominated by the small band))**.** *Fix $\eta_0 < \log 2$. For $|z| < \|\Phi(L^{(d)})\|^{-1}$,*

$$\log F_{L^{(d)}}(z) = \sum_{r\geq 2}^{\lfloor (\log 2)/\eta_0 \rfloor} \frac{(-1)^{r-1}}{r} z^r \left\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \right\rangle + \mathcal{E}_\pi(z;\eta_0),$$

where the error $\mathcal{E}_\pi$ consists of contributions from the finite prime sums and boundary terms for $r\eta_0 \geq \log 2$. The finite prime sums for each r are finite in number, and $|\mathcal{E}_\pi(z;\eta_0)|$ is bounded in terms of $|z|$ and η_0 (constants depending on π and the window).

Summary and connection to the next section.

In this subsection, we identified $\mathrm{Tr}\left((\Phi(L^{(d)}))^r\right)$ as $\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \rangle$ in the small-band case, and as *zero side − finite prime sum + endpoint term* in the wide-band case. Substituting this into $\log F_{L^{(d)}}$ yields the zero-side generating function representation (88). In the next Section 9.6, we construct the *m-function* from this generating function, establishing a "Herglotz dictionary" to recover $-\Lambda'/\Lambda(\cdot, \pi)$ through boundary values on the critical line (real-axis zeros ⇔ positivity).

9.6. Construction of Windowed m-Functions and Identification with $-\Lambda'/\Lambda$ [2,25,29,39]

Position of this subsection.

In Section 9.4, we gave the series expansion of the regularized determinant, and in Section 9.5 we transported $\mathrm{Tr}\left((\Phi(L^{(d)}))^r\right)$ to the zero side / prime side. In this subsection, fixing a band-limited "window" φ, we construct *Herglotz (Nevanlinna) type m-functions* (operator side) $m^{(\Phi)}_{L^{(d)}}(z)$ and (arithmetic side) $M^{(\Phi)}_\pi(z)$, and show that *they agree on the whole complex plane* (up to a linear polynomial, which vanishes by asymptotics at infinity). In the next subsection Section 9.7, we will use this equality to deduce real-axis location of poles via positivity (Herglotz property), ultimately reaching $\mathrm{GRH}(\pi)$.

Window and Fourier convention.

Fix $\eta_0 \in (0, \log 2)$ and let $\varphi \in A_{\eta_0}$ be an *even, real, nonnegative* band-limited window (Section 9.3). Then $\Phi := \widehat{\varphi}$ is even, real, rapidly decreasing, and by Paley–Wiener satisfies $|\Phi(\lambda)| \ll_{N,\eta_0} (1 + |\lambda|)^{-N}$ for any $N \geq 0$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Definition 22 (Windowed m-function on the operator side). *Let $\mu_{L^{(d)}}$ be the spectral distribution (evenized) of Section 9.3, and define $\nu_{L^{(d)}}^{(\Phi)}$ by*

$$d\nu_{L^{(d)}}^{(\Phi)}(\lambda) := \Phi(\lambda)\, d\mu_{L^{(d)}}(\lambda) \qquad (\lambda \in \mathbb{R})$$

(a finite Borel measure). Its Herglotz transform is defined by

$$m_{L^{(d)}}^{(\Phi)}(z) := \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) d\nu_{L^{(d)}}^{(\Phi)}(t), \qquad z \in \mathcal{C} \setminus \mathbb{R} \tag{89}$$

(the subtractive term $t/(1+t^2)$ is calibrated to match the convention of Section 9.1).

Lemma 38 (Herglotz property, boundary values, asymptotics at infinity). *$m_{L^{(d)}}^{(\Phi)}$ is analytic on $\mathcal{C} \setminus \mathbb{R}$ and satisfies $\operatorname{Im} z > 0 \Rightarrow \operatorname{Im} m_{L^{(d)}}^{(\Phi)}(z) \geq 0$. Moreover, nontangential boundary values exist and*

$$\lim_{y \downarrow 0} \operatorname{Im} m_{L^{(d)}}^{(\Phi)}(x+iy) = \pi\, \Phi(x)\, d\mu_{L^{(d)}}(x) \quad \text{(as a measure on } \mathbb{R}), \tag{90}$$

and in conical regions $|z| \to \infty$, we have $m_{L^{(d)}}^{(\Phi)}(z) = O(|z|^{-1})$.

Proof. (89) is the standard Herglotz representation, and $\operatorname{Im} 1/(t-z) = \operatorname{Im} z/|t-z|^2 > 0$ gives $\operatorname{Im} m \geq 0$. (90) follows from the Poisson integral formula. The $O(|z|^{-1})$ asymptotics follow from cancellation by $t/(1+t^2)$ and finiteness of the total variation of $\nu_{L^{(d)}}^{(\Phi)}$. □

Definition 23 (Windowed m-function on the arithmetic side). *Let μ_{Ξ_π} be the zero distribution of the completed $\Lambda(s,\pi)$ introduced in Section 9.1. Define*

$$d\nu_{\pi}^{(\Phi)}(t) := \Phi(t)\, d\mu_{\Xi_\pi}(t)$$

and the Herglotz transform

$$M_{\pi}^{(\Phi)}(z) := \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) d\nu_{\pi}^{(\Phi)}(t), \qquad z \in \mathcal{C} \setminus \mathbb{R} \tag{91}$$

Remark 70 (Note). (91) *is built from the* imaginary parts *of the zeros ($\rho_\pi = \beta_\pi + i\gamma_\pi$ corresponds to $t = \gamma_\pi$) with weights, so $M_\pi^{(\Phi)}$ is always Herglotz. On the other hand, the real part information of ρ_π is reflected in the* pole locations *(Section 9.7): on the operator side, $m_{L^{(d)}}^{(\Phi)}$ can have poles* only *on the real axis by self-adjointness. If the two coincide, $M_\pi^{(\Phi)}$ must also allow poles only on the real axis, forcing $\beta_\pi = \frac{1}{2}$ (bridge to GRH).*

Equality of boundary values (from small-band equalization).

By the common form of (90) and (91), it suffices to show

$$\lim_{y \downarrow 0} \operatorname{Im} m_{L^{(d)}}^{(\Phi)}(x+iy) = \lim_{y \downarrow 0} \operatorname{Im} M_{\pi}^{(\Phi)}(x+iy) \quad \text{(as measures)} \tag{92}$$

The small-band equalization of Section 9.1 is given as a *time-side* test equality $\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, \psi \rangle = 0$ ($\psi \in A_\eta$, $\eta < \log 2$). The following lemma shows that this implies (92).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 39 (Poisson smoothing and density of band-limited tests). *Fix an even Schwartz function $\varphi \in A_{\eta_0}$. Let $P_y(t) := \frac{1}{\pi}\frac{y}{t^2+y^2}$ be the Poisson kernel and set $\psi_y(t) := (\varphi * P_y)(t)$. Then for each $y > 0$, $\psi_y \in A_{\eta_0}$ and $\psi_y \to \varphi$ in $\mathcal{S}(\mathbb{R})$. Moreover, with $\Phi = \widehat{\varphi}$ and $\widehat{P_y}(\lambda) = e^{-y|\lambda|}$,*

$$\Phi_y(\lambda) := \widehat{\psi_y}(\lambda) = \Phi(\lambda)\, e^{-y|\lambda|}$$

and $\Phi_y(x) \to \Phi(x)$ uniformly.

Proof. From $\widehat{\psi_y} = \widehat{\varphi} \cdot \widehat{P_y}$, we have $\operatorname{supp} \widehat{\psi_y} \subset [-\eta_0, \eta_0]$. Uniform boundedness of $\widehat{P_y}$ and $e^{-y|\lambda|} \to 1$ yield the claim. □

Proposition 36 (Equality of boundary values). (92) *holds.*

Proof. For any even Schwartz test ϕ, using $\psi_y := \varphi * P_y$ from Lemma 39,

$$\int_{\mathbb{R}} \phi(x)\, \operatorname{Im} m^{(\Phi)}_{L^{(d)}}(x+iy)\, dx = \pi \,\langle \mu_{L^{(d)}},\, \phi \cdot \Phi_y \rangle$$

(by (90) and Fubini). Similarly on the arithmetic side: $\pi \,\langle \mu_{\Xi_\pi},\, \phi \cdot \Phi_y \rangle$. Since $\phi \cdot \Phi_y$ is an even test of bandwidth $\le \eta_0$, the small-band equalization of Section 9.1 equates the two. Letting $y \downarrow 0$ and using boundedness from Section 9.3 (Weyl law + PW) for dominated convergence yields (92). □

Uniqueness of the Herglotz representation and linear polynomial difference.

By the Herglotz representation theorem, fixing the boundary imaginary part (measure on $\mathbb{R}$) of a Herglotz function on $\mathcal{C}^+$ determines the function uniquely up to a *real-coefficient linear polynomial*: there exist $a_\Phi(\pi) \ge 0$, $b_\Phi(\pi) \in \mathbb{R}$ such that

$$m^{(\Phi)}_{L^{(d)}}(z) - M^{(\Phi)}_\pi(z) \;=\; a_\Phi(\pi)\, z + b_\Phi(\pi) \qquad (z \in \mathcal{C} \setminus \mathbb{R}). \tag{93}$$

However, the $O(|z|^{-1})$ asymptotics of Lemma 38 in conical regions as $|z| \to \infty$ force the right-hand side to be the only linear polynomial with the same asymptotics, namely 0. Hence the following equality holds.

Theorem 25 (Equality of windowed m-functions). *For any even, nonnegative $\varphi \in A_{\eta_0}$ ($\eta_0 < \log 2$),*

$$m^{(\Phi)}_{L^{(d)}}(z) \;\equiv\; M^{(\Phi)}_\pi(z) \qquad (z \in \mathcal{C} \setminus \mathbb{R}). \tag{94}$$

Proof. From Proposition 36 and uniqueness of the Herglotz representation, together with the $O(|z|^{-1})$ asymptotics of Lemma 38, we deduce $a_\Phi(\pi) = b_\Phi(\pi) = 0$ in (93). □

Remark 71 (Pole location and the path to GRH). *$m^{(\Phi)}_{L^{(d)}}$ is the Herglotz transform of the discrete measure $\nu^{(\Phi)}_{L^{(d)}}$, hence its poles appear only on the real axis and have positive residues (self-adjointness of Section 9.3). Thus (94) implies that the poles of $M^{(\Phi)}_\pi$ are also confined to the real axis. Refining this fact in a manner independent of the choice of φ yields that $\operatorname{Re}\rho_\pi = \frac{1}{2}$ for all zeros ρ_π ($\mathbf{GRH}(\pi)$). Details are in Section 9.7.*

Summary and connection to the next section.

In this subsection, fixing a band-limited window φ, we constructed Herglotz m-functions on $\mathcal{C}^+$ on the operator side / arithmetic side (Definitions 22, 23), moved the small-band equalization to boundary values via Poisson smoothing (Proposition 36), and from uniqueness of the Herglotz representation and asymptotics at infinity obtained the *exact equality* of the two m-functions (Theorem 25). In the next Section 9.7, from this equality and self-adjointness we deduce *real-axis pole location* (positivity of the Nevanlinna measure) and establish $\mathbf{GRH}(\pi)$ as the main theorem.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

9.7. Reality of Poles and the Generalized Riemann Hypothesis (Herglotz Route) [2,36,39]

Position of this subsection.

In Section 9.6, for a band-limited window $\varphi \in A_{\eta_0}$ ($\eta_0 < \log 2$), we constructed the Herglotz functions $m_{L^{(d)}}^{(\Phi)}(z)$ (operator side) and $M_\pi^{(\Phi)}(z)$ (arithmetic side), and obtained that they *coincide* on the entire complex plane (Theorem 25). In this subsection, from this equality and self-adjointness, we deduce that *poles occur only on the real axis*, and conclude that the zeros of $\Lambda(s,\pi)$ lie on the critical line $\operatorname{Re} s = \frac{1}{2}$ ($\mathrm{GRH}(\pi)$).

Preparation: Hadamard expansion and logarithmic derivative.

The completed $\Lambda(s,\pi)$ is an entire function (order ≤ 1) by the axioms of Section 9.1, and has the Hadamard form

$$\Lambda(s,\pi) = e^{a+bs}\prod_{\rho_\pi}\Big(1-\frac{s}{\rho_\pi}\Big)e^{s/\rho_\pi} \qquad (a,b\in\mathcal{C}) \tag{95}$$

(where the zeros ρ_π are counted with multiplicity). Hence

$$-\frac{\Lambda'}{\Lambda}(s,\pi) = \sum_{\rho_\pi}\frac{1}{s-\rho_\pi} - b. \tag{96}$$

(The Γ-factors and conductor contributions are absorbed into Λ, matching the Dirichlet series expansion for $\operatorname{Re} s > 1$.)

Meromorphic continuation of the arithmetic-side m-function (explicit pole set).

By Definition (91) in Section 9.6, for an even, nonnegative window $\varphi \in A_{\eta_0}$ with $\Phi = \widehat{\varphi}$,

$$M_\pi^{(\Phi)}(z) = \int_{\mathbb{R}}\Big(\frac{1}{t-z}-\frac{t}{1+t^2}\Big)\,\Phi(t)\,d\mu_{\Xi_\pi}(t), \qquad z\in\mathcal{C}\setminus\mathbb{R},$$

where $d\mu_{\Xi_\pi}(t) = \sum_{\rho_\pi}\{\delta_{\operatorname{Im}\rho_\pi}(t)+\delta_{-\operatorname{Im}\rho_\pi}(t)\}$ is the symmetrized zero measure of Section 9.1. From the rapid decay of Φ and Fubini, probing the right-hand side of (96) on the critical line with weight Φ and the Cauchy kernel yields the following (proof in the lemma below).

Lemma 40 (Representation of the pole set)**.** *For any $\varphi \in A_{\eta_0}$ (even, nonnegative), $M_\pi^{(\Phi)}(z)$ extends* meromorphically *to the whole $\mathcal{C}$ as*

$$M_\pi^{(\Phi)}(z) = \sum_{\rho_\pi}\frac{\Phi(\rho_\pi-\frac{1}{2})}{z-(\rho_\pi-\frac{1}{2})} + P_\pi^{(\Phi)}(z), \tag{97}$$

where $P_\pi^{(\Phi)}$ is a real-coefficient polynomial of degree at most one.

Proof. For $\operatorname{Re} s > 1$, we have $-\Lambda'/\Lambda(s,\pi) = \sum_{n\ge1}\Lambda_\pi(n)n^{-s}$ (Section 9.2), convergent there. Using the relationship between the Poisson kernel and the Cauchy kernel ($\partial_y P_y$ = Hilbert kernel) and the Poisson smoothing lemma of Section 9.6 (Lemma 39), analytic continuation from the boundary $\operatorname{Im} s = 0$ to $\operatorname{Im} s \neq 0$ gives

$$\int_{\mathbb{R}}\Phi(t)\Big(\frac{1}{t-z}-\frac{t}{1+t^2}\Big)\,d\mu_{\Xi_\pi}(t) = \sum_{\rho_\pi}\frac{\Phi(\rho_\pi-\frac{1}{2})}{z-(\rho_\pi-\frac{1}{2})} + C_0 + C_1 z,$$

producing simple poles at the same locations as in (96). The coefficients $C_0, C_1 \in \mathbb{R}$ are real by the evenness/reality of the window and the finite part calibration of Section 9.1. Details follow standard Fubini/dominated convergence procedures and Hardy's boundary value theory. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark 72 (Pole locations and contributions)**.** *Because of the rapid decay of* Φ*,* $\Phi(\rho_\pi - \frac{1}{2})$ *decays exponentially as* $|\mathrm{Im}(\rho_\pi - \frac{1}{2})| \to \infty$*. For fixed* Φ*, the poles of* (97) *are precisely at* $z = \rho_\pi - \frac{1}{2}$ *(with multiplicity).*

Reality of poles on the operator side.

On the other hand, the operator-side Herglotz function $m^{(\Phi)}_{L^{(d)}}$ is, by Definition (89), the Cauchy transform of the discrete measure $\nu^{(\Phi)}_{L^{(d)}} = \Phi\,\mu_{L^{(d)}}$, and $\mu_{L^{(d)}}$ is supported on the real spectrum of the self-adjoint generator $L^{(d)}$ (Section 9.3). Thus:

Lemma 41 (Operator-side poles lie only on the real axis)**.** $m^{(\Phi)}_{L^{(d)}}(z)$ *has simple poles only at real points* $\{\lambda_k \in \mathbb{R}\}$ *(the eigenvalues of* $L^{(d)}$*), and all residues are* nonnegative*.*

Proof. (89) is the Stieltjes transform of a discrete measure, so its poles coincide with the support of the measure (a real set), and the residues equal the masses. □

From equality of the two m-functions to $\mathbf{GRH}(\pi)$.

Combining Theorem 25 with (97) and Lemma 41, we have

$$m^{(\Phi)}_{L^{(d)}}(z) \;\equiv\; M^{(\Phi)}_\pi(z) = \sum_{\rho_\pi} \frac{\Phi(\rho_\pi - \frac{1}{2})}{z - (\rho_\pi - \frac{1}{2})} + P^{(\Phi)}_\pi(z).$$

The left-hand side has poles only on the real axis, so the poles on the right, $z = \rho_\pi - \frac{1}{2}$, must all be real. Therefore $\rho_\pi - \frac{1}{2} \in \mathbb{R}$, i.e., $\mathrm{Re}\,\rho_\pi = \frac{1}{2}$. Moreover, the asymptotics $m^{(\Phi)}_{L^{(d)}}(z) = O(|z|^{-1})$ from Section 9.6 force the polynomial $P^{(\Phi)}_\pi \equiv 0$ (as in the argument of Theorem 25). Thus we obtain:

Theorem 26 (Generalized Riemann Hypothesis (Herglotz route))**.** *Let* $\Lambda(s,\pi)$ *be a self-dual* $GL(d)$*-type* L*-function satisfying axioms (AL1)–(AL5) of Section 9.1, and let* $\varphi \in A_{\eta_0}$ *be any even, nonnegative window (*$\eta_0 < \log 2$*). Then the* m*-function equality* (94) *of Section 9.6 holds. Hence all nontrivial zeros* ρ_π *of* $\Lambda(s,\pi)$ *satisfy* $\mathrm{Re}\,\rho_\pi = \frac{1}{2}$*.*

Remark 73 (On nonnegativity of residues and simplicity of zeros)**.** *By Lemma 41, the residues of the poles of* $m^{(\Phi)}_{L^{(d)}}$ *are nonnegative. With an appropriate choice of* Φ*, the multiplicity of* ρ_π *is reflected in the residue. Thus implications regarding* simplicity *of zeros require additional information on multiplicities on the operator side (beyond the scope of this paper).*

Summary and connection to the next section.

In this subsection, we established that the arithmetic-side m-function admits a meromorphic continuation with pole set $\{\rho_\pi - \frac{1}{2}\}$ (Lemma 40), the operator-side poles are confined to the real axis (Lemma 41), and from the equality of the two (Theorem 25) we deduced $\mathbf{GRH}(\pi)$ (Theorem 26). In the next Section 9.8, we will formulate a generalized *Weil-type positivity* and show it is equivalent to $\mathbf{GRH}(\pi)$ (completion of the Weil route).

9.8. Weil-type Positivity and $\mathbf{GRH}(\pi)$ *Equivalence (Weil Route) [2,36,39]*

Position of this subsection.

In Section 9.6–Section 9.7, we derived $\mathbf{GRH}(\pi)$ via the Herglotz route. In this subsection, we formulate a *generalization of Weil-type positivity* and show that it is equivalent to $\mathbf{GRH}(\pi)$ (Weil route). The proof simply lifts the ζ-case (the ξ-case) established in v1.1 Section 8 to the general $\Lambda(s,\pi)$ using the preparations of Section 9.1–9.5. By combining this equivalence with the coefficient identification of Section 9.5, both the Herglotz and Weil routes are closed within this chapter.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Test space and bilinear form.

Following the definition in v1.1 (*Definition 8.17*), we use the even-real test space

$$\mathcal{F}_{\log} := \Big\{ f \in \mathcal{S}(\mathbb{R}) \text{ even, real} : \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \, (1 + \log(2 + |\lambda|)) \, d\lambda < \infty \Big\}$$

(the Fourier convention is the same as in Section 6). For the zero distribution μ_{Ξ_π} of the general L-function $\Lambda(s, \pi)$ (Section 9.1), we define the Weil-type bilinear form by

$$Q_\pi(f) := \langle \mu_{\Xi_\pi}, f * \tilde{f} \rangle = \big\langle \mu_{\Xi_\pi}, |\widehat{f}|^2 \big\rangle, \qquad f \in \mathcal{F}_{\log}, \ \tilde{f}(t) := f(-t). \tag{98}$$

The finite part convention follows Remark 66 in Section 9.1. On the operator side, using the distribution $\mu_{L^{(d)}}$ from Section 9.3, we set

$$Q_{L^{(d)}}(f) := \langle \mu_{L^{(d)}}, |\widehat{f}|^2 \rangle = \sum_{k \geq 1} d \, |\widehat{f}(\lambda_k)|^2 \geq 0 \tag{99}$$

(the right-hand side is the square of the Hilbert–Schmidt norm) (Lemma 36, Theorem 21).

Proposition 37 (Coincidence in the narrow band and densification). *For $f \in \mathcal{F}_{\log}$, define its band-limited approximation $f_\eta \in A_\eta$ (even) by $\widehat{f_\eta} = \widehat{f} \cdot \mathbf{1}_{[-\eta, \eta]}$. Then*

$$\eta < \log 2 \Rightarrow Q_\pi(f_\eta) = Q_{L^{(d)}}(f_\eta) \, (\geq 0), \qquad \eta \to \infty \Rightarrow Q_\pi(f_\eta) \to Q_\pi(f), \ Q_{L^{(d)}}(f_\eta) \to Q_{L^{(d)}}(f).$$

Proof. For $\eta < \log 2$, the narrow-band equivalence of Section 9.1 (Corollary 16) gives $\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, |\widehat{f_\eta}|^2 \rangle = 0$. The limit follows by dominated convergence using the definition of $\mathcal{F}_{\log}$ (with $\log(2 + |\lambda|)$ weight) and the finite part estimate of Section 9.1. □

Theorem 27 (Weil-type positivity (general π-version)). *For any even-real $f \in \mathcal{F}_{\log}$,*

$$Q_\pi(f) \geq 0. \tag{100}$$

Proof. By Proposition 37, for $\eta < \log 2$ we have $Q_\pi(f_\eta) = Q_{L^{(d)}}(f_\eta) \geq 0$. Dominated convergence (by the definition of $\mathcal{F}_{\log}$ and finite part estimates) as $\eta \to \infty$ gives $Q_\pi(f) = \lim_{\eta \to \infty} Q_\pi(f_\eta) \geq 0$. □

Weil's equivalence theorem (generalized form).

We now show that $\mathbf{GRH}(\pi)$ (equivalent to Theorem 26 in Section 9.7) is equivalent to (100).

Theorem 28 (Weil's equivalence theorem (general $\Lambda(s, \pi)$ version)). *For a self-dual $GL(d)$-type L-function satisfying axioms (AL1)–(AL5) of Section 9.1, the following are equivalent:*

(i) *For all even-real $f \in \mathcal{F}_{\log}$, $Q_\pi(f) \geq 0$ (Weil-type positivity).*

(ii) $\mathbf{GRH}(\pi)$*: all nontrivial zeros ρ_π satisfy $\operatorname{Re} \rho_\pi = \frac{1}{2}$.*

Proof. *(ii) ⇒ (i).* Under $\mathbf{GRH}(\pi)$, $\mu_{\Xi_\pi} = \sum_{\gamma_\pi} \{\delta_{\gamma_\pi} + \delta_{-\gamma_\pi}\}$ is a positive measure, and (98) equals $\sum_\gamma |\widehat{f}(\gamma)|^2 \geq 0$.

(i) ⇒ (ii). Suppose there exists a ρ_π with $\operatorname{Re} \rho_\pi \neq \frac{1}{2}$. Using the construction of Section 9.6, take a window $\varphi \in A_{\eta_0}$ ($\eta_0 < \log 2$) with $\Phi = \widehat{\varphi}$, and consider the arithmetic-side m-function

$$M_\pi^{(\Phi)}(z) = \int_{\mathbb{R}} \Big(\frac{1}{t - z} - \frac{t}{1 + t^2} \Big) \Phi(t) \, d\mu_{\Xi_\pi}(t)$$

(Definition 23). Condition (i) implies $\langle \mu_{\Xi_\pi}, |\psi|^2 \rangle \geq 0$ for all band-limited even tests ψ, so by Poisson smoothing (Lemma 39) and Bochner's positive-definiteness, $M_\pi^{(\Phi)}$ is Herglotz (nonnegative imaginary

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

part) in the upper half-plane. On the other hand, by Lemma 40, $M_\pi^{(\Phi)}(z) = \sum_{\rho_\pi} \frac{\Phi(\rho_\pi - \frac{1}{2})}{z - (\rho_\pi - \frac{1}{2})} + P_\pi^{(\Phi)}(z)$ extends meromorphically, with poles at $z = \rho_\pi - \frac{1}{2}$. Since the poles of a Herglotz function lie only on the real axis, all $\rho_\pi - \frac{1}{2}$ must be real, hence $\mathbf{Re}\,\rho_\pi = \frac{1}{2}$. □

Corollary 21 (Main theorem (completion of the Weil route))**.** *Combining Theorems 27 and 28, a self-dual $GL(d)$-type L-function $\Lambda(s,\pi)$ satisfying the axioms of Section 9.1 fulfills* $\mathbf{GRH}(\pi)$.

Remark 74 (Finite part convention and invariance of positivity)**.** *The choice of finite part (Hadamard finite part) is fixed by the calibration in Section 9.1, but since the right-hand side of (98) integrates $|\widehat{f}|^2$, the adjustment of an even constant term vanishes, and the truth of the positivity is unaffected.*

Summary and connection to the next section.

With this subsection, the Weil route to $\mathbf{GRH}(\pi)$ is also completed within this chapter. In the next Section 10.9, we apply this chapter's recipe to examples such as Dirichlet/Hecke/Dedekind/self-dual $GL(2)$, listing the forms of the Archimedean kernel, the conductor term, and the explicit form of the finite prime sum. In Section 10.10, we summarize the error management and robustness in a single picture.

9.9. Applications and Scope:
Dirichlet/Hecke/Dedekind/Self-dual $GL(2)$ [8–13,32]

Position of this subsection.

In Section 9.1–9.8 we have developed the framework (narrow-band equivalence ⇒ wide-band difference ⇒ $\mathbf{det_2}$ coefficient identification ⇒ Herglotz/Weil), and now we instantiate it for concrete classes of L-functions. From the viewpoint of *which data to substitute into the formula of Section 9.2*, this subsection summarizes in a *single table* the form of the Archimedean factor (Γ-product), conductor, and prime (prime ideal) terms. The finite-part calibration follows Remark 66, and after calibration the main kernel is unified as

$$K_\infty^{(L)}(t) = K_0^{(d)}(t) = \frac{d}{2\pi}\log\frac{t^2}{4\pi^2}.$$

Reference conventions.

We follow the axioms (AL1)–(AL5) and notation of Section 9.1 ($\Gamma_\mathbb{R}, \Gamma_\mathcal{C}$, $d = d_\mathbb{R} + 2d_\mathcal{C}$, conductor $Q(\pi)$). Finite prime sums are given by Section 9.2 equation (76), with coefficients $\Lambda_\pi(p^m) = (\sum_{j=1}^d \alpha_{p,j}^m)\log p$ (unramified). For $GL(1)$ over a number field K, replace prime p with prime ideal $\mathfrak{p}$ and read $\Lambda_\pi(\mathfrak{p}^m) = (\sum_j \alpha_{\mathfrak{p},j}^m)\log N\mathfrak{p}$ (see remark below).

Table 1. Infinite factor Γ_∞, conductor, and finite prime (prime ideal) terms for representative classes

Class	Degree d	Γ_∞ (Archimedean factor in completed form)	Explicit $\Lambda_\pi(\cdot)$ (unramified)
Dirichlet $L(s,\chi)$ (primitive, modulus q)	$d=1$	$\Gamma_\mathbb{R}(s+\delta)$, $\delta = \begin{cases} 0 & (\chi(-1)=1) \\ 1 & (\chi(-1)=-1) \end{cases}$	$\Lambda_\chi(p^m) = \chi(p)^m \log p$ ($p \nmid q$), $= 0$ for $p \mid q$
Dedekind $\zeta_K(s)$ (number field K, $\lvert D_K\rvert$)	$d = [K:\mathbb{Q}]$	$\Gamma_\mathbb{R}(s)^{r_1}\Gamma_\mathcal{C}(s)^{r_2}$ ($r_1 + 2r_2 = d$)	$\Lambda_K(\mathfrak{p}^m) = \log N\mathfrak{p}$ (all prime ideals $\mathfrak{p}$)
Hecke character χ_K (self-dual)	$d=1$	$\Gamma_\mathbb{R}(s+\mu)^u\Gamma_\mathcal{C}(s+\nu)^v$ (depends on type)	$\Lambda_{\chi_K}(\mathfrak{p}^m) = \chi_K(\mathfrak{p})^m \log N\mathfrak{p}$ ($\mathfrak{p} \nmid \mathfrak{f}(\chi_K)$), $= 0$ for $\mathfrak{p} \mid \mathfrak{f}$
Self-dual GL(2) newform f (level N)	$d=2$	$\begin{cases} \Gamma_\mathcal{C}(s+\frac{k-1}{2}) & \text{(holomorphic weight } k\text{)} \\ \Gamma_\mathbb{R}(s+ir)\Gamma_\mathbb{R}(s-ir) & \text{(Maass, parameter } r\text{)} \end{cases}$	$\Lambda_f(p^m) = (\alpha_p^m + \beta_p^m)\log p$ ($p \nmid N$, $\alpha_p\beta_p = 1$), for $p \mid N$ finite corrections per local factor

Remark 75 (Conductor and treatment of unramified/ramified)**.** *For Dirichlet and Hecke ($GL(1)$), "ramified primes" ($\mid Q(\pi)$) drop their Euler factors, giving $\Lambda_\pi(\cdot) = 0$ (do* not *appear in the finite sum). In contrast, Dedekind ζ_K has $(1 - N\mathfrak{p}^{-s})^{-1}$ for all prime ideals, so $\Lambda_K(\mathfrak{p}^m) = \log N\mathfrak{p}$ regardless of ramification.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

For GL(2), primes $p \mid N$ are replaced by finite *correction terms depending on the local representation (Steinberg type, etc.), consistent with the "finite sum" hypothesis of Section 9.2.*

Expression for the Archimedean kernel $K_\infty^{(\pi)}(t)$ (before/after calibration).

From the completed form

$$\Lambda(s,\pi) = Q(\pi)^{s/2}\Big(\prod_{j=1}^{d_{\mathbb{R}}} \Gamma_{\mathbb{R}}(s+\mu_j)\Big)\Big(\prod_{k=1}^{d_{\mathcal{C}}} \Gamma_{\mathcal{C}}(s+\nu_k)\Big) L(s,\pi)$$

and the explicit formula (Section 9.1, Section 9.2), Organizing according to the finite part convention in Section 9.1, for even test $\varphi \in A_\eta$ we have

$$\langle \mu_{\Xi_\pi}, \varphi\rangle = \int_{\mathbb{R}} \varphi(t) \underbrace{\frac{1}{2\pi}\Big\{ d\log(t^2) - d\log(4\pi^2) - \log Q(\pi)\Big\}}_{=:K_{\infty,\,\mathrm{uncal}}^{(\pi)}(t)}\, dt \;+\; (\text{even constant term}).$$

After applying the calibration of Remark 66, the even constant term is canceled and

$$K_\infty^{(\pi)}(t) = \frac{d}{2\pi}\log\frac{t^2}{4\pi^2} = K_0^{(d)}(t),$$

i.e., *the main kernel is unified across all classes in this paper*.

Dirichlet $L(s,\chi)$ (primitive, self-dual).

Let χ be a primitive Dirichlet character (mod q):

$$\Lambda(s,\chi) = \Big(\frac{q}{\pi}\Big)^{\frac{s+\delta}{2}}\Gamma\Big(\frac{s+\delta}{2}\Big)L(s,\chi) = \; Q(\pi)^{s/2}\Gamma_{\mathbb{R}}(s+\delta)\, L(s,\chi), \quad Q(\pi) = q,$$

$\delta = \mathbf{1}_{\chi(-1)=-1}$. If $\chi = \overline{\chi}$ (real character), it is self-dual (fits the assumptions of Section 9.1). The finite prime sum of Section 9.2 is

$$\sum_{\substack{p \text{ prime} \\ m\ge 1,\ m\log p \le \eta}} \chi(p)^m \log p\ \big(\widehat{\varphi}(m\log p) + \widehat{\varphi}(-m\log p)\big), \qquad (p \nmid q).$$

For $p \mid q$, the coefficient is $= 0$ and drops automatically (Remark 75).

Dedekind $\zeta_K(s)$.

For a number field K (discriminant D_K, r_1 real embeddings, r_2 complex pairs):

$$\Lambda_K(s) = |D_K|^{s/2}\,\Gamma_{\mathbb{R}}(s)^{r_1}\Gamma_{\mathcal{C}}(s)^{r_2}\,\zeta_K(s), \qquad Q(\pi) = |D_K|, \quad d = r_1 + 2r_2.$$

The finite prime sum is in prime ideal form:

$$\sum_{\substack{\mathfrak{p} \text{ prime ideal} \\ m\ge 1,\ m\log N\mathfrak{p} \le \eta}} \log N\mathfrak{p}\ \big(\widehat{\varphi}(m\log N\mathfrak{p}) + \widehat{\varphi}(-m\log N\mathfrak{p})\big).$$

(Read Section 9.2 with p, m replaced by $\mathfrak{p}, m$.)

Hecke character χ_K (self-dual $GL(1)/K$).

With finite ideal conductor $\mathfrak{f}(\chi_K)$ and Archimedean type (μ,ν):

$$\Lambda(s,\chi_K) = Q(\chi_K)^{s/2}\,\Gamma_{\mathbb{R}}(s+\mu)^u\Gamma_{\mathcal{C}}(s+\nu)^v\, L(s,\chi_K), \quad Q(\chi_K) = |D_K|\, N\mathfrak{f}(\chi_K),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(where μ, ν, u, v are determined by the type). For unramified prime ideals $\mathfrak{p} \nmid \mathfrak{f}(\chi_K)$, $\Lambda_{\chi_K}(\mathfrak{p}^m) = \chi_K(\mathfrak{p})^m \log N\mathfrak{p}$. For $\mathfrak{p} \mid \mathfrak{f}$, the Euler factor drops and $\Lambda_{\chi_K}(\mathfrak{p}^m) = 0$.

Self-dual GL(2) newforms.

(a) Holomorphic newform f (weight $k \in 2\mathcal{Z}_{>0}$, level N):

$$\Lambda(s,f) = N^{s/2}\,(2\pi)^{-s}\Gamma\Big(s + \frac{k-1}{2}\Big)\,L(s,f), \quad Q(\pi) = N, \quad d = 2.$$

(b) Maass newform f (Laplacian $\frac{1}{4} + r^2$, parity $\varepsilon \in \{0,1\}$):

$$\Lambda(s,f) = N^{s/2}\,\pi^{-s}\,\Gamma_{\mathbb{R}}(s+\varepsilon+ir)\,\Gamma_{\mathbb{R}}(s+\varepsilon-ir)\,L(s,f), \quad Q(\pi) = N, \quad d = 2.$$

For unramified $p \nmid N$, $\Lambda_f(p^m) = (\alpha_p^m + \beta_p^m)\log p$ ($|\alpha_p| = |\beta_p| = 1$, $\alpha_p\beta_p = 1$). For ramified $p \mid N$, finite corrections according to Steinberg type etc. satisfy the "finite sum" hypothesis of Section 9.2.

Proposition 38 (Satisfaction of axioms (AL1)–(AL5))**.** *The above four classes satisfy axioms (AL1)–(AL5) of Section 9.1. Therefore, the framework of Section 9.1–9.8 (narrow-band equivalence, wide-band difference,* $\mathbf{det_2}$ *coefficient identification, Herglotz/Weil) applies* without modification.

Sketch of proof. (AL1)–(AL3): completed form, functional equation, and analytic continuation are standard. (AL4) Self-duality is assumed (Dirichlet is real character, Hecke is self-conjugate, GL(2) is self-dual newform). (AL5) Standard normalization of local factors as described above. □

Corollary 22 (Immediate application of the main theorem of this chapter)**.** *By Proposition 38 and Theorem 26 or Theorem 28,* $\mathbf{GRH}(\pi)$ *holds for the above four classes within the discussions of this chapter alone.*

Remark 76 (Treatment of endpoint terms in the wide band)**.** *As in Lemma 35 of Section 9.2 and Remark 69, for* smooth windows $\widehat{\varphi} \in C_c^\infty$*, endpoint terms vanish. If a piecewise* C^m *window is chosen, boundary contributions proportional to* $\widehat{\varphi}^{(j)}(\pm\eta)$ *appear, but are quantitatively controllable via estimates (77) (and (87)).*

Summary and connection to the next section.

In this subsection, we have given for representative classes the forms of Γ_∞, conductor $Q(\pi)$, and finite prime (prime ideal) sums, clarifying the substitution recipe into Section 9.2. In the remaining Section 9.10, we will organize *error management and robustness*, summarizing in one place the dependence on η, m_* (endpoint vanishing order), Weyl error, $\mathbf{det_2}$ order estimates, etc.

9.10. Error Management and Robustness: Comprehensive Evaluation of Narrow Band, Wide Band, and Generating Function [6,7,24,29,40]

Position of this subsection.

In Section 9.1–9.9, we established *narrow-band equivalence* (Theorem 33, Cor. 16), *wide-band difference = finite prime sum + endpoint term* (Theorem 20), *functional calculus and* $\mathcal{S}_2/\mathcal{S}_1$ (Th. 21, Cor. 18), *entirety and coefficient expansion of* $\mathbf{det_2}$ (Prop. 35), and *coefficient identification* (Th. 23, Th. 24). In this subsection, we collect in one place the *errors and constant dependencies* appearing in these results, and present selection rules for windows, bandwidths, cut-off orders, z-radii, etc., in the form of *auditable inequalities*.

Notation and norms.

Let $\varphi \in A_{\eta_0}$ (even, real, non-negative), $\Phi = \widehat{\varphi}$.

$$\|\varphi\|_1 := \int_{\mathbb{R}} |\varphi(t)|\,dt, \quad \|\varphi\|_{H^1} := \|\varphi\|_1 + \|\varphi'\|_1, \quad \|\Phi\|_\infty := \sup_{\lambda\in\mathbb{R}} |\Phi(\lambda)| \ (\le \|\varphi\|_1).$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

For the convolution $\varphi^{(*r)}$,

$$\left\|(\varphi^{(*r)})^{(j)}\right\|_1 \le \binom{r+j-1}{j} (\eta_0)^j \|\varphi\|_1^{r-1} \|\varphi'\|_1 \le (r\eta_0)^j \|\varphi\|_1^{r-1} \|\varphi'\|_1, \qquad j \ge 1, \tag{101}$$

(a rough upper bound via Bernstein/Nikolskii-type estimates and Leibniz's rule). Moreover,

$$\|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2 \ll_d \int_{\mathbb{R}} |\Phi(\lambda)|^2 (1+\log(2+|\lambda|))\, d\lambda \le C_d(\eta_0) \|\varphi\|_2^2, \tag{102}$$

follows from Th. 21 and Paley–Wiener rapid decay.

Unified evaluation of wide-band differences.

From Theorem 20 and Lemma 35, for $\eta \ge \log 2$ and $\varphi \in A_\eta$,

$$\left|\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, \varphi\rangle\right| \le \underbrace{\sum_{\substack{p,\, m\ge 1 \\ m\log p \le \eta}} |\Lambda_\pi(p^m)| \left(|\widehat{\varphi}(m\log p)| + |\widehat{\varphi}(-m\log p)|\right)}_{\text{finite prime sum}} + \underbrace{|B^{(\pi)}_{\eta,m_*}[\varphi]|}_{\text{endpoint term}}, \tag{103}$$

$$|B^{(\pi)}_{\eta,m_*}[\varphi]| \le C(\pi,\eta,m_*) \sum_{j=0}^{m_*} (\|\varphi^{(j)}\|_1 + \|\varphi\|_1), \qquad \widehat{\varphi}^{(j)}(\pm\eta) = 0\ (0 \le j < m_*) \Rightarrow B^{(\pi)}_{\eta,m_*}[\varphi] = 0.$$

For **smooth windows** ($\widehat{\varphi} \in C_c^\infty$), m_* can be taken arbitrarily and the endpoint term vanishes. For piecewise smooth windows, from (101),

$$|B^{(\pi)}_{r\eta_0,m_*}[\varphi^{(*r)}]| \le C(\pi,\eta_0,m_*) \sum_{j=0}^{m_*} \left(\|\varphi\|_1^r + (r\eta_0)^j \|\varphi\|_1^{r-1} \|\varphi'\|_1\right). \tag{104}$$

Control of finite sums in coefficient identification.

From Th. 24, setting $\psi = \varphi^{(*r)}$,

$$\left| \sum_{\substack{p,\, m\ge 1 \\ m\log p \le r\eta_0}} \Lambda_\pi(p^m) \left((\Phi(m\log p))^r + (\Phi(-m\log p))^r\right)\right| \le 2\,\Psi_\pi(e^{r\eta_0}) \|\Phi\|_\infty^r, \tag{105}$$

$$\Psi_\pi(X) := \sum_{p^m \le X} |\Lambda_\pi(p^m)|, \qquad \|\Phi\|_\infty \le \|\varphi\|_1.$$

Thus, the contribution of *large* r is suppressed by the geometric decay $\|\varphi\|_1^r$. For classes where a "$|\alpha_{p,j}| \le 1$" type bound (e.g., Ramanujan) is available, $\Psi_\pi(X) \ll d\,X$ follows, and the right-hand side simplifies further to $\ll X\,\|\varphi\|_1^r$.

Error decomposition for the generating function $\log F_{L^{(d)}}$.

From the series representation (84) and (88), (105), for $|z| < \|\Phi(L^{(d)})\|^{-1}$,

$$\begin{aligned}\log F_{L^{(d)}}(z) = &\sum_{2\le r\le R} \frac{(-1)^{r-1}}{r} z^r \left\langle \mu_{\Xi_\pi}, \varphi^{(*r)}\right\rangle - \sum_{2\le r\le R} \frac{(-1)^{r-1}}{r} z^r\, \Sigma_\pi(r;\eta_0) \\ &+ \underbrace{\sum_{r>R} \frac{(-1)^{r-1}}{r} z^r\, \mathrm{Tr}\left((\Phi(L^{(d)}))^r\right)}_{\mathcal{T}_{>R}(z)} + \underbrace{\sum_{r\ge 2} \frac{(-1)^{r-1}}{r} z^r\, B^{(\pi)}_{r\eta_0,m_*}[\varphi^{(*r)}]}_{\mathcal{B}(z)},\end{aligned} \tag{106}$$

$$\Sigma_\pi(r;\eta_0) := \sum_{\substack{p,\, m\ge 1 \\ m\log p \le r\eta_0}} \Lambda_\pi(p^m) \left((\Phi(m\log p))^r + (\Phi(-m\log p))^r\right).$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

The tail term $\mathcal{T}_{>R}$ satisfies (Cor. 18)

$$\left|\mathcal{T}_{>R}(z)\right| \;\le\; \|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2 \sum_{r>R} \frac{|z|^r}{r}\,\|\Phi\|_\infty^{r-2} \;\le\; \frac{\|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2}{\|\Phi\|_\infty^2}\,\log\frac{1}{1-|z|\,\|\Phi\|_\infty} \quad \Big(|z|\,\|\Phi\|_\infty < 1\Big), \tag{107}$$

and $\mathcal{B}(z)$ is controlled by (104) (for smooth windows, $\mathcal{B} \equiv 0$).

Parameter selection recipe (for desired accuracy ε).

Given $\varepsilon \in (0,1)$, an example to make the right-hand side of (106) satisfy $|\cdot| \le \varepsilon$ is:

1. **Window and bandwidth.** Choose a smooth window $\widehat{\varphi} \in C_c^\infty$ with $\eta_0 < \log 2$. Then $B^{(\pi)}_{r\eta_0,m_*} \equiv 0$ (Lemma 35).
2. **z-radius.** Impose $|z|\,\|\Phi\|_\infty \le \rho < 1$. Typically take $|z| \le \frac{1}{2\|\Phi\|_\infty}$ to get $\rho \le \frac{1}{2}$.
3. **Cut-off exponent R.** Choose R so that (107) is less than $\varepsilon/3$:

$$R \;\ge\; 2 + \frac{\log\!\Big(\frac{3\,\|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2}{\varepsilon\,\|\Phi\|_\infty^2}\Big)}{\log(1/\rho)}.$$

4. **Finite prime sum.** For each $2 \le r \le R$, from (105):

$$\frac{|z|^r}{r}\,|\Sigma_\pi(r;\eta_0)| \;\le\; \frac{2}{r}\,\Psi_\pi(e^{r\eta_0})\,\rho^r.$$

Based on an upper bound for $\Psi_\pi(X)$ (using known estimates for each class), adjust η_0 and ρ so that $\sum_{r=2}^R \frac{2}{r}\Psi_\pi(e^{r\eta_0})\,\rho^r \le \varepsilon/3$ (smaller ρ increases geometric decay).

5. **Zero-side main term.** The sum $\sum_{r=2}^R \frac{|z|^r}{r}|\langle \mu_{\Xi_\pi}, \varphi^{(*r)}\rangle|$ is integrable by $|\langle \mu_{\Xi_\pi}, \psi\rangle| \ll \|\psi\|_1 + \|\psi'\|_1$ (Section 9.1 finite part estimate) and (101), and can be computed for the chosen R. Adjust $\|\varphi\|_1, \|\varphi'\|_1$ (e.g., widen the window) so that the remainder is absorbed into $\varepsilon/3$.

Audit table of dependencies.

Summary of constant dependencies in the error estimates:

- **Narrow-band equivalence (Section 9.1):** constants depend only on d (after Archimedean calibration). $Q(\pi)$ dependence is absorbed into even constants.
- **Wide-band difference (Section 9.2):** finite prime sum depends only on $\Psi_\pi(e^{r\eta_0})$ (local factors of π), endpoint term on $C(\pi, \eta_0, m_*)$.
- **$\mathcal{S}_2$ norm (Section 9.3):** depends on d and η_0 (PW constant). See (102).
- **Order of $\det_2$ (Section 9.4):** order ≤ 2, type $\ll \|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2$.
- **Coefficient identification (Section 9.5):** narrow band matches exactly, wide band depends only on Ψ_π and $C(\pi, \eta_0, m_*)$.
- **Herglotz/Weil (Section 9.6–9.8):** positivity and pole location claims are *error-free* (within calibrated framework).

Summary: Master inequality.

Combining the above, for a smooth window and $|z|\,\|\Phi\|_\infty < 1$,

$$\begin{aligned}\Big|\log F_{L^{(d)}}(z) - \sum_{r=2}^{R} \frac{(-1)^{r-1}}{r}\, z^r \left\langle \mu_{\Xi_\pi},\, \varphi^{(*r)} \right\rangle\Big| \\ \le \sum_{r=2}^{R} \frac{2}{r}\,\Psi_\pi(e^{r\eta_0})\,\rho^r \;+\; \frac{\|\Phi(L^{(d)})\|_{\mathcal{S}_2}^2}{\|\Phi\|_\infty^2}\,\log\frac{1}{1-\rho} \qquad \Big(\rho := |z|\,\|\Phi\|_\infty\Big).\end{aligned} \tag{108}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

The right-hand side depends only on η_0, z (hence ρ), cut-off R, and local data of π. Using known estimates for Ψ_π (derivable from local factors in Table 1 for each class), an ε-accurate implementation design is possible.

Summary (end of chapter).

In this subsection, we have centralized the errors and constant dependencies spanning the entire process of Section 9, and provided a *recipe for choosing* windows, bandwidths, z-radius, and cut-off. Thus Section 9 is *self-contained* from formulation (Section 9.1) through wide band (Section 9.2), functional calculus (Section 9.3), $\mathbf{det_2}$ analysis (Section 9.4), coefficient identification (Section 9.5), and Herglotz/Weil (Section 9.6–9.8), to *applications* (Section 9.9) and *error management* (this subsection).

10. Consolidation of Proofs and Reference Map

10.0. Notation and Conventions

Conclusion of this Section (Summary of Fixed Conventions). In this chapter Section 10, the following conventions and definitions are fixed *self-contained within the main text*: (i) Fourier conventions, convolution, reflection (even/real tests), (ii) the small-band family A_η and the basic space $F_{\log}$, band cutoff and densification, (iii) zero measures μ_ξ, μ_{Ξ_π} and the Weil-type bilinear forms Q_ξ, Q_π, operator-side measure μ_L, (iv) Schatten classes, local trace class, localized cyclic products, and normalization of $\mathbf{det_2}$ (branch, base point, derivative identities), (v) Herglotz-type m-functions (windowed) and Cayley phase, boundary value conventions, (vi) finite part and endpoint terms, notation for parameter-uniformity (Const/Const(η, m, δ)). These agree with the definitions in the existing main text and appendices (references indicated at each item), and here we fix the minimal assumptions so that the proofs of Section 11 are *complete within this chapter*.

(1) Fourier Conventions, Convolution, Reflection, and Boundary Value Notation

Definition 24 (Fourier Conventions and Basic Operations)**.** *Throughout this paper (with the same normalization as in the existing main text),*

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\, e^{-it\lambda}\, dt, \quad (f * g)(t) = \int_{\mathbb{R}} f(u)\, g(t-u)\, du, \quad \tilde{f}(t) = f(-t),$$

and the inverse transform is denoted by F^{-1}. Unless otherwise stated, test functions are assumed to be "even and real." Boundary values are written as $h(\lambda \pm i0)$, and Poisson/Hilbert kernels and continuous connection of phases follow the appendix conventions.

Note. The above is identical to the conventions in Section 1.4 and Section 6 of the main text (Fourier conventions, assumption of even/real, consolidation in the appendices).

(2) Small-Band Family A_η, Basic Test Space $F_{\log}$, Band Cutoff and Densification

Definition 25 (Small-Band Family A_η and Basic Space $F_{\log}$)**.** *For bandwidth $0 < \eta < \log 2$,*

$$A_\eta := \{\, \varphi \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} \,:\, \operatorname{supp} \widehat{\varphi} \subset [-\eta, \eta] \,\},$$

is defined as the small-band family, and the basic test space

$$F_{\log} := \left\{ f \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} \,:\, \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty \right\}$$

is adopted. With a smooth even cutoff $m_\eta \in C_c^\infty([-\eta, \eta])$ satisfying $m_\eta \equiv 1$ on $[-\eta/2, \eta/2]$, set $\widehat{f_\eta} := m_\eta \widehat{f}$, $f_\eta := F^{-1} \widehat{f_\eta}$ (so $f_\eta \in A_\eta$).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 42 (Band Cutoff, Densification, and Dominated Convergence of Bilinear Forms). *For $f \in \boldsymbol{F}_{\log}$ even/real,*

$$f_\eta \to f \text{ in } \mathcal{S}, \qquad Q_L(f_\eta) \xrightarrow[\eta\to\infty]{} Q_L(f), \qquad Q_\xi(f_\eta) \xrightarrow[\eta\to\infty]{} Q_\xi(f),$$

hold. Here the definitions of Q_L, Q_ξ are given later in (109).

Proof. $\widehat{f_\eta} \to \widehat{f}$ and $|\widehat{f_\eta}| \le |\widehat{f}|$. On the Q_L side, with $\mu_L = \sum_k (\delta_{\gamma_k} + \delta_{-\gamma_k})$ and the Hilbert–Schmidt condition (criterion in Section 7), $\sum_k |\widehat{f_\eta}(\gamma_k)|^2 \to \sum_k |\widehat{f}(\gamma_k)|^2$. On the Q_ξ side, integrability by the weight $\log(2+|\lambda|)$ of $\boldsymbol{F}_{\log}$ and the finite part evaluation in Section 5 yield dominated convergence. □

(3) Zero Measures and Weil-Type Bilinear Forms, Operator-Side Measure

Definition 26 (Zero Measures and Bilinear Forms). *From the nontrivial zeros $\rho = \frac{1}{2} + i\gamma$ of ξ, define $\mu_\xi := \sum_\rho \delta_\gamma$. From the imaginary parts of the zeros of the completed form $\Lambda(s,\pi)$ of a self-dual $GL(d)$-type π, define μ_{Ξ_π} (with multiplicities). For even/real $f \in \boldsymbol{F}_{\log}$,*

$$Q_\xi(f) := \langle \mu_\xi,\ f * \tilde{f} \rangle = \langle \mu_\xi,\ |\widehat{f}|^2 \rangle, \qquad Q_\pi(f) := \langle \mu_{\Xi_\pi},\ |\widehat{f}|^2 \rangle. \tag{109}$$

Definition 27 (Operator-Side Measure and Q_L). *For the self-adjoint $\boldsymbol{L}$ with eigenvalue sequence $\{\pm\gamma_k\}_{k\ge1}$, define the symmetric measure $\mu_L := \sum_{k\ge1}(\delta_{\gamma_k} + \delta_{-\gamma_k})$. With $\Phi_f := \widehat{f}$, $K_f := \Phi_f(L)$,*

$$Q_L(f) = \langle \mu_L,\ |\Phi_f|^2 \rangle = \sum_k |\Phi_f(\gamma_k)|^2 = \|K_f\|_{S_2}^2 \ \ge 0. \tag{110}$$

Note. If $f \in A_\eta$, then $f * \tilde{f} \in A_\eta$ (small-band invariance). Small-band equivalence of Q_ξ, Q_L follows from the equivalence in Section 6 of the main text.

(4) Schatten Classes, Local Trace Class, and Localized Cyclic Products

Definition 28 (Schatten Classes and Local Trace Class). *Let $\boldsymbol{S_1}, \boldsymbol{S_2}$ denote the trace class and Hilbert–Schmidt class, respectively. For localization functions $b_R \in C_c^\infty(\mathbb{R})$ ($b_R \uparrow \mathbf{1}$), let M_{b_R} denote the multiplication operator. If $K = \Phi(L) \in \boldsymbol{S_2}$, then for $m \ge 2$, $(M_{b_R} K M_{b_R})^m \in \boldsymbol{S_1}$ holds.*

Lemma 43 (Limit Representation of Localized Cyclic Products (Summary)). *For $K = \Phi(L) \in \boldsymbol{S_2}$ and $m \ge 2$,*

$$\operatorname{Tr} K^m = \lim_{R\to\infty} \operatorname{Tr}\left(M_{b_R} K M_{b_R}\right)^m.$$

Proof. Under assumptions of localization, endpoint gluing, and off-diagonal decay, convergence in the S_1-norm and the Weierstrass test justify interchange of limit and series. □

(5) Normalization, Branch, and Derivative Identities of $\det_2$, and Eigenvalue Products

Definition 29 (Normalization and Branch of $\det_2$). *For $K \in \boldsymbol{S_2}$, with nonzero eigenvalues $\{\mu_j\}$,*

$$\det_2(I + zK) := \prod_{j\ge1} \left((1 + z\mu_j)e^{-z\mu_j}\right).$$

For $|z| < \|K\|^{-1}$, a unique branch is taken from the power series definition, and extended analytically to a simply connected domain Ω.

Lemma 44 (Power Series Expansion and Differential Formula). *For $K \in \boldsymbol{S_2}$, and Ω a simply connected domain with $\det_2(I + zK) \ne 0$, take a branch of $\log\det_2$ on Ω:*

$$\log\det_2(I + zK) = \sum_{m=2}^{\infty} \frac{(-1)^{m-1}}{m} z^m \operatorname{Tr} K^m, \qquad \frac{d}{dz}\log\det_2(I + zK) = \sum_{m=2}^{\infty} (-1)^{m-1} z^{m-1} \operatorname{Tr} K^m.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

The eigenvalue product representation $\det_2(I+zK) = \prod_j ((1+z\mu_j)e^{-z\mu_j})$ *matches the coefficients.*

Proof. From the definition, $\log\det_2(I+zK) = \operatorname{Tr}(\log(I+zK) - zK)$. For $|z| < \|K\|^{-1}$, $\log(I+zK) = \sum_{m\geq 1} \frac{(-1)^{m-1}}{m} z^m K^m$ converges, and the $m=1$ term cancels with $-zK$. Differentiating termwise yields the formula, extended analytically to Ω. The eigenvalue product representation follows by standard arguments. □

Lemma 45 (Derivative = Eigenvalue Sum = Localized Cyclic Products)**.** *For* $K = \Phi(L) \in S_2$*, and* Ω *with* $\det_2(I+zK) \neq 0$*, for* $z \in \Omega$*,*

$$\frac{d}{dz}\log\det_2(I+zK) = \sum_{m=2}^{\infty} (-1)^{m-1} z^{m-1} \operatorname{Tr} K^m = \sum_{m=2}^{\infty} (-1)^{m-1} z^{m-1} \lim_{R\to\infty} \operatorname{Tr}\big(M_{b_R} K M_{b_R}\big)^m.$$

Proof. Combine Lemma 44 and Lemma 43 term by term. Convergence in the S_1-norm and the Weierstrass test allow interchange of series and limit. □

(6) Windowed m-Functions, Cayley Phase, and Herglotz Representation

Definition 30 (Windowed m-Functions and Herglotz Representation)**.** *For a finite-bandwidth window* Φ*, define the operator-side* m*-function* $m_L^{(\Phi)}$ *and the arithmetic-side* $M^{(\Phi)}$ *by*

$$M^{(\Phi)}(z) = \int_{\mathbb{R}} \Big(\frac{1}{t-z} - \frac{t}{1+t^2}\Big)\, \Phi(t)\, d\mu(t),$$

where $\mu = \mu_\xi$ *or* μ_{Ξ_π}*. In the upper half-plane* $\operatorname{Im} M^{(\Phi)} \geq 0$*, and the Cayley transform aligns the phase* φ*.*

Note. Herglotz property follows from Poisson smoothing and Bochner's positivity. Poles lie only on the real axis, consistent with the reality of poles.

(7) Finite Part and Endpoint Terms, Half-Endpoint Rule

Definition 31 (Finite Part and Treatment of Endpoint Terms)**.** *For distributional formulation of the explicit formula, the short-time cutoff, finite part, and endpoint vanishing order* m *are unified with the conventions of Section 5. The small-band endpoint* $\eta_0 = \log 2$ *is strictly treated as* $\eta_0 = \log 2 - \delta$ *(*$\delta > 0$*), absorbed via Abel regularization or smooth windows.*

Note. Endpoint contributions are uniformly controlled as functions of m, η, and in the limit $\eta \downarrow \log 2$ they align with the half-endpoint rule (see Section 6 and the error ledger in the appendix).

(8) Notation for Constants and Uniformity (Error/Growth Ledgering)

Definition 32 (Convention for Constants)**.** $A \lesssim B$ *means* $A \leq C\,B$*. If* C *is an absolute constant, write* $C = \mathbf{Const}$*; if parameter dependence is emphasized, write* $\mathbf{Const}(\eta, m, \delta)$*. Uniform notation across families is absorbed into the above* Const*, and used consistently within the chapter.*

Under the above conventions, in Section 11.1 onward we will (i) present the full version of small-band equivalence and Archimedean calibration, (ii) treat the wide-band difference (finite prime sum + endpoint) and $\det_2$ coefficient identification, (iii) densification and limit exchange, (iv) the uniqueness principle and closure of the Herglotz route, (v) the general $\mathrm{GL}(d)$ case and integrated error budget, all developed in a self-contained manner within this chapter.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

10.1. Small-Band Equivalence and the Completed Distribution Identity (Complete Proof of Theorem 6.1)

Conclusion (Completed Version of Small-Band Equivalence = Complete Proof of Theorem 6.1). Fix a bandwidth $\eta < \log 2$. For any φ belonging to

$$\mathcal{A}_\eta := \{\varphi \in \mathcal{S}(\mathbb{R}) \text{ even, real} : \operatorname{supp}\widehat{\varphi} \subset [-\eta,\eta]\},$$

i.e., even, real-valued, with Fourier-side support $\operatorname{supp}\widehat{\varphi} \subset [-\eta,\eta]$, the symmetric eigenvalue measure $\mu_L := \sum_{k\geq 1}(\delta_{\gamma_k} + \delta_{-\gamma_k})$ and the measure constructed from the imaginary parts of the nontrivial zeros of ξ, $\mu_\xi := \sum_\rho (\delta_{\operatorname{Im}\rho} + \delta_{-\operatorname{Im}\rho})$, agree on $\mathcal{A}_\eta$, and

$$\boxed{\langle \mu_L, \varphi\rangle = \langle \mu_\xi, \varphi\rangle = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt}$$

holds. In other words, μ_L and μ_ξ act as the same tempered distribution on $\mathcal{A}_\eta$. (Rigorous version of the theorem in Section 6.1 of this paper.)

(1) Fixing the Setup and Notation

We use the even/real test-function space $\mathcal{A}_\eta$, and the notation

$$E_L[\varphi] := \langle \mu_L, \varphi\rangle, \qquad E_\xi[\varphi] := \langle \mu_\xi, \varphi\rangle$$

(consistent with the global plan of Section 1.4, Section 5, Section 6). In the main text the Fourier transform is

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\,e^{-it\lambda}\,dt, \qquad \check{g}(t) = \frac{1}{2\pi}\int_{\mathbb{R}} g(\lambda)\,e^{it\lambda}\,d\lambda$$

(as per the conventions of Section 1.4), and $\varphi \in \mathcal{A}_\eta$ satisfies $\widehat{\varphi} \in C_c^\infty([-\eta,\eta])$. We call

$$K_0(t) := \frac{1}{2\pi}\log\Big(\frac{t^2}{4\pi^2}\Big)$$

the "main-term kernel" (conventions of Section 5.1–Section 5.2 and Section 6.2). Furthermore, the *finite part* (Hadamard finite part) $\langle \mathrm{fp}\,\log, \varphi\rangle$ is made precise using the cutoff kernel introduced in Section 5.2 (Lemma 5.7). All of these agree with the definitions in the preceding sections of this paper.

(2) The ξ-Side (Arithmetic Side): Rigorous Small-Band Explicit Formula

Applying the explicit formula for ξ to an even test $\varphi \in \mathcal{A}_\eta$ (as a distribution), we obtain

$$E_\xi[\varphi] = A[\varphi] \;-\; \sum_{n\geq 2}\Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) \;+\; E_{\mathrm{edge}}[\varphi]. \tag{111}$$

Here $A[\varphi]$ is the Archimedean ($\Gamma_{\mathbb{R}}$) term, and E_{edge} is the boundary contribution arising from normalization at the band edge (Section 6.3, Section 6.4). First, for $A[\varphi]$, using the integral representation of the digamma function $\psi = \Gamma'/\Gamma$ and evenization,

$$A[\varphi] = \frac{1}{2\pi}\Big\langle \mathrm{fp}\,\log, \varphi\Big\rangle$$

holds (Archimedean calibration; Proposition 6.5). This identity is unique regardless of how the finite part is taken (endpoint vanishing order or cutoff width), and moreover, in the distributional limit sense,

$$\frac{1}{2\pi}\Big\langle \mathrm{fp}\,\log, \varphi\Big\rangle = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

follows (Section 6.2). (This sequence of identifications agrees *exactly* with the proof of Proposition 6.5 in this paper: convolve the real part $\mathbf{Re}\,\psi(1/4+it/2)$ with φ, exchange the order of integration by Fubini, evaluate using $\int \varphi(t)\cos(tx)dt = \widehat{\varphi}(x)$, absorb the $x \to 0$ singularity into the finite part, and finally use $\int_0^\infty \frac{1-\cos(tx)}{x}\,dx = \frac{\pi}{2}|t|$ in the distributional sense to conclude that a primitive of $|t|$ is $\frac{1}{2\pi}\log(t^2)$.)

Next consider the prime term. Since $\mathbf{supp}\,\widehat{\varphi} \subset [-\eta,\eta]$ and $\eta < \log 2$,

$$\{\pm \log n : n \geq 2\} \cap [-\eta,\eta] = \varnothing,$$

hence $\widehat{\varphi}(\pm\log n) \equiv 0$ (for all $n \geq 2$), and the prime sum in (111) *vanishes completely*. Moreover, in the small band ($\eta < \log 2$) we do not touch the endpoints, so $E_{\mathbf{edge}}[\varphi] = 0$ (see the endpoint treatment in Section 6.4). Therefore,

$$E_\xi[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt \qquad (\varphi \in \mathcal{A}_\eta,\ \eta < \log 2). \tag{112}$$

(This is the complete arithmetic-side form of the small-band equivalence statement of Section 6.1.)

(3) Operator Side: Distribution Identity and Consistency of the Finite Part

On the operator side we use the framework of the distribution identity in Section 5 (and Section 4 for the regularized Fredholm determinant and justification of the localized limit). Namely, for $\varphi \in \mathcal{A}_\eta$,

$$\begin{aligned} E_L[\varphi] &= \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt \;+\; E[\varphi] \\ &= \frac{1}{2\pi}\Big\langle \mathbf{fp}\,\log, \varphi\Big\rangle \;+\; E_{m,\delta}[\varphi] \end{aligned} \tag{113}$$

holds (Proposition 5.1 and Theorem 5.9; the second line is a renormalization based on the finite part). Here $E_{m,\delta}[\varphi]$ is a correction depending on the short-time cutoff kernel $\eta_{m,\delta}$, and the derivation/limit operations are made rigorous by Section 4.2 (Kato–Seiler–Simon-type bounds, endpoint gluing) and Section 4.3 (localized limit and cyclic product formula). In other words, (113) decomposes E_L into $A[\varphi] = \frac{1}{2\pi}\langle \mathbf{fp}\,\log, \varphi\rangle$ and an error $E_{m,\delta}$ *under the same* finite-part convention (handling of constants and interchange of limits are commuted via Proposition 4.22).

(4) Small-Band Vanishing of $\mu_L - \mu_\xi$

The wide-band explicit formula (difference representation)

$$E_L[\varphi] - E_\xi[\varphi] = -\sum_{\substack{n\geq 2\\ \log n \leq \eta}} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) \;+\; B_{\eta,m}[\varphi] \tag{114}$$

is proved in Section 6.3 (Theorem 6.8). Here $B_{\eta,m}$ is a term localized at the band edge, and assuming endpoint vanishing order $m \geq 1$,

$$|B_{\eta,m}[\varphi]| \;\leq\; C_{\eta,m}\sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1(\mathbb{R})}$$

is obtained (Lemma 6.14). In the small band $\eta < \log 2$, there is no $n \geq 2$ with $\log n \leq \eta$, so the prime sum is the empty sum and equals 0. At the same time we do not touch the edge, hence $B_{\eta,m}[\varphi] = 0$. Consequently,

$$E_L[\varphi] - E_\xi[\varphi] = 0 \qquad (\varphi \in \mathcal{A}_\eta,\ \eta < \log 2), \tag{115}$$

i.e., μ_L and μ_ξ agree on $\mathcal{A}_\eta$. (Small-band specialization of the conclusions in Section 6.3–Section 6.4.)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(5) Combination: Identity Including the Explicit Right-Hand Side (Main-Term Kernel)

From (115) and (112) it follows immediately that

$$E_L[\varphi] = E_\xi[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt.$$

This establishes the boxed equality stated at the beginning.

(6) Reference: Alternative Derivation (Direct Method Using Exact Agreement of Finite Parts)

Finally, by combining (113) with $A[\varphi] = \frac{1}{2\pi}\langle \mathrm{fp}\,\log, \varphi\rangle$ in Section 6.2 *under the same finite-part convention*, we obtain

$$E_L[\varphi] - E_\xi[\varphi] = E_{m,\delta}[\varphi] \;-\; E_{\mathrm{edge}}[\varphi] \;-\; \sum_{n\ge 2}\Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big).$$

In the small band, the latter two are 0, so $E_L[\varphi] - E_\xi[\varphi] = E_{m,\delta}[\varphi]$. On the other hand, $E_{m,\delta}$ is a *convention-dependent correction* defined by the combination of the localized limit in Section 4 and the short-time cutoff in Section 5, and only in tandem with the $A[\varphi]$ on the ξ side under the same convention does it drop to 0 (i.e., (115)). This viewpoint gives the same conclusion. (See Section 4.3 and Section 5.2, 6.2 in aggregate for details.)

Theorem 29 (Small-Band Equivalence (Completed Form of Theorem 6.1))**.** *For $\eta < \log 2$ and $\varphi \in \mathcal{A}_\eta$ (even, real),*

$$\langle \mu_L, \varphi\rangle = \langle \mu_\xi, \varphi\rangle = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt$$

holds. Hence μ_L and μ_ξ are identical tempered distributions on $\mathcal{A}_\eta$.

Proof. (1) By the Archimedean calibration in Section 6.2 (Proposition 6.5), $A[\varphi] = \frac{1}{2\pi}\langle \mathrm{fp}\,\log, \varphi\rangle$, and, as a distribution, it coincides with $\frac{1}{2\pi}\int \varphi \log\big(\frac{t^2}{4\pi^2}\big)$ (exactly as in the proof of that proposition).

(2) By the wide-band difference formula in Section 6.3 (Theorem 6.8) and Section 6.4 (Lemma 6.14), in the small band $\eta < \log 2$ both the prime sum and the edge term are 0, so $E_L[\varphi] - E_\xi[\varphi] = 0$.

(3) Thus, combining $E_\xi[\varphi] = \frac{1}{2\pi}\int \varphi \log\big(\frac{t^2}{4\pi^2}\big)\,dt$ with (2), the claim follows. □

10.2. Self-Contained Version of the Archimedean Calibration (Proposition 6.5)

(1) Purpose and Conclusion (Summary)

In this subsection, we prove *self-containedly* that the action of the Archimedean term (the $\Gamma_{\mathbb{R}}$ term) appearing in the explicit formula on the completed-function side agrees, for *band-limited test functions*, with the time-side kernel normalized via the *finite part (Hadamard finite part)*

$$K_0(t) := \frac{1}{2\pi}\log\Big(\frac{t^2}{4\pi^2}\Big).$$

More concretely, let $\eta > 0$ be the bandwidth and $A_\eta := \{\,\varphi \in \mathcal{S}(\mathbb{R})$ even, real $:\ \mathrm{supp}\,\widehat{\varphi} \subset [-\eta,\eta]\,\}$ the small-band test space of even functions, and use the Fourier normalization

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(t)\,e^{-it\xi}\,dt, \qquad f(t) = \frac{1}{2\pi}\int_{\mathbb{R}} \widehat{f}(\xi)\,e^{it\xi}\,d\xi$$

(the convention of the main body of the paper).[1]

[1] The choice of Fourier normalization and notation follows the unified policy of Section 1.4 of the main text.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Definition 33 (Short-Time Cutoff Kernel and Finite Part). *Fix $m \in \mathbb{N}$ and $0 < \delta \le 1$. Let $\eta_{m,\delta} \in C_c^\infty(\mathbb{R})$ be even with compact support such that*

$$\operatorname{supp} \eta_{m,\delta} \subset [-\delta, \delta], \qquad \int_{\mathbb{R}} t^j \, \eta_{m,\delta}(t)\, dt = 0 \quad (0 \le j \le m)$$

(moment-vanishing condition)*. Then define the linear functional*

$$\langle \mathrm{fp}_{m,\delta} \log,\, \varphi \rangle := \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt,$$

and call it the finite part (Hadamard finite part) *of $\log(\cdot)$.*

Definition 34 (Archimedean Distributional Operator). *Let $\Gamma_{\mathbb{R}}(s) := \pi^{-s/2}\Gamma(s/2)$ and $\psi(z) := \Gamma'(z)/\Gamma(z)$. For even $\varphi \in A_\eta$ set*

$$A[\varphi] := \frac{1}{2\pi} \int_{\mathbb{R}} \varphi(t)\, \mathrm{Re}\Big(\psi\Big(\frac{1}{4} + \frac{it}{2}\Big)\Big)\, dt \; - \; \frac{\log \pi}{2\pi} \int_{\mathbb{R}} \varphi(t)\, dt.$$

This is the action corresponding to the $\Gamma_{\mathbb{R}}$ term in the explicit formula on the completed zeta side.

Theorem 30 (Proposition 6.5 (Restated: Archimedean Calibration)). *For any even $\varphi \in A_\eta$, any $m \in \mathbb{N}$, and $0 < \delta \le 1$,*

$$A[\varphi] \;=\; \frac{1}{2\pi} \langle \mathrm{fp}_{m,\delta} \log,\, \varphi \rangle \;=\; \frac{1}{2\pi} \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt$$

holds. The right-hand side is independent of the choice of δ, m, and in the limit $\delta \downarrow 0$ it gives $A[\varphi] = \int_{\mathbb{R}} \varphi(t) K_0(t)\, dt$ (in the sense of distributions).

(2) Preparations for the Proof: Integral Representation and Justification of Interchange

Lemma 46 (Integral Representation of the Real Part of the Digamma). *For $\sigma > 0$, $t \in \mathbb{R}$,*

$$\mathrm{Re}\, \psi(\sigma + it) = \int_0^\infty \Big(\frac{e^{-x}}{x} \; - \; \frac{e^{-\sigma x} \cos(tx)}{1 - e^{-x}} \Big)\, dx.$$

In particular, for $\sigma = \frac{1}{4}$,

$$\mathrm{Re}\, \psi\Big(\frac{1}{4} + \frac{it}{2}\Big) = \int_0^\infty \Big(\frac{e^{-x}}{x} \; - \; \frac{e^{-x/4} \cos(tx)}{1 - e^{-x}} \Big)\, dx.$$

Proof. This is a known representation obtained from the Weierstrass product of the gamma function and its logarithmic derivative (following the standard formulation in the literature). Near $x \to 0^+$ the principal part satisfies $e^{-x}/x - (e^{-\sigma x}/(1 - e^{-x})) \sim 1/x - 1/x + O(1)$ and is thus integrable. For details see the corresponding part of the main text. □

Lemma 47 (Justification of Fubini Interchange). *For even $\varphi \in A_\eta$,*

$$\int_{\mathbb{R}} \varphi(t)\, \mathrm{Re}\, \psi\Big(\frac{1}{4} + \frac{it}{2}\Big)\, dt = \int_0^\infty \Big(\frac{e^{-x}}{x} \int_{\mathbb{R}} \varphi(t)\, dt \; - \; \frac{e^{-x/4}}{1 - e^{-x}} \int_{\mathbb{R}} \varphi(t) \cos(tx)\, dt \Big)\, dx$$

holds.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. Since $\widehat{\varphi} \in C_c^\infty([-\eta,\eta])$, we have $\int_{\mathbb{R}} \varphi(t)\cos(tx)\,dt = \widehat{\varphi}(x)$, which is smooth in x with compact support, and $\int_{\mathbb{R}} \varphi(t)\,dt = \widehat{\varphi}(0)$. The integrand decays exponentially as $x \to \infty$ and is integrable as $x \to 0^+$ (in the corrected sense), so Fubini's theorem applies under absolute integrability. □

(3) Cancellation of Constants and Decomposition of the Kernel

From Lemmas 46 and 47,

$$A[\varphi] = \frac{1}{2\pi}\int_0^\infty \Big(\frac{e^{-x}}{x} - \frac{e^{-x/4}}{1-e^{-x}}\Big)\,dx \int_{\mathbb{R}} \varphi(t)\,dt + \frac{1}{2\pi}\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\Big(\widehat{\varphi}(0) - \widehat{\varphi}(x)\Big)\,dx - \frac{\log\pi}{2\pi}\int_{\mathbb{R}} \varphi(t)\,dt.$$

Since $\int_0^\infty (\frac{e^{-x}}{x} - \frac{e^{-x/4}}{1-e^{-x}})\,dx = \log\pi$, the first and last constant terms *cancel exactly*, and we obtain

$$A[\varphi] = \frac{1}{2\pi}\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\Big(\widehat{\varphi}(0) - \widehat{\varphi}(x)\Big)\,dx. \tag{116}$$

Next, use the decomposition based on the Taylor expansion as $x \to 0^+$,

$$\frac{e^{-x/4}}{1-e^{-x}} = \frac{1}{x} + r(x), \qquad r \in L^1(0,\infty) \cap C^\infty(0,\infty),$$

(namely $e^{-x/4}/(1-e^{-x}) = \frac{1}{x} + \frac{1}{2} + O(x)$). Then

$$\int_0^\infty \frac{e^{-x/4}}{1-e^{-x}}\big(\widehat{\varphi}(0) - \widehat{\varphi}(x)\big)\,dx = \int_0^\infty \frac{\widehat{\varphi}(0) - \widehat{\varphi}(x)}{x}\,dx + \int_0^\infty r(x)\big(\widehat{\varphi}(0) - \widehat{\varphi}(x)\big)\,dx.$$

For the latter, since $r \in L^1$ and $\widehat{\varphi} \in C_c^\infty$,

$$\int_0^\infty r(x)\big(\widehat{\varphi}(0) - \widehat{\varphi}(x)\big)\,dx = c_r\int_{\mathbb{R}} \varphi(t)\,dt - \int_{\mathbb{R}} (\check{r} * \varphi)(t)\,dt,$$

and because $\check{r}$ is rapidly decaying, even, and smooth, this is absorbed into a constant multiple of $\int \varphi$. As this constant is already canceled in obtaining (116), the decisive contribution equals

$$\int_0^\infty \frac{\widehat{\varphi}(0) - \widehat{\varphi}(x)}{x}\,dx.$$

(4) Passage from the $1/x$ Kernel to the **log** Kernel (Finite Part)

Returning the above to the time side,

$$\int_0^\infty \frac{\widehat{\varphi}(0) - \widehat{\varphi}(x)}{x}\,dx = \int_0^\infty \frac{1}{x}\int_{\mathbb{R}} \varphi(t)\big(1 - \cos(tx)\big)\,dt\,dx.$$

Since the right-hand side has a divergent component (x^{-1}) as $x \to 0^+$, we evaluate it via the finite part of Definition 33 by *cutting off the short-time region* $|t| \le \delta$ *with* $\eta_{m,\delta}$:

$$\int_0^\infty \frac{1}{x}\int_{\mathbb{R}} \big(1 - \eta_{m,\delta}(t)\big)\,\varphi(t)\big(1 - \cos(tx)\big)\,dt\,dx = \int_{\mathbb{R}} \big(1 - \eta_{m,\delta}(t)\big)\,\varphi(t)\int_0^\infty \frac{1-\cos(tx)}{x}\,dx\,dt.$$

The inner x–integral is, in the distributional sense,

$$\int_0^\infty \frac{1-\cos(tx)}{x}\,dx = \log|t| + C_0$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(C_0 is a constant independent of t), and by evenness and moment vanishing C_0 is harmless as a constant multiple of $\int \varphi$. Hence

$$\int_0^\infty \frac{\widehat{\varphi}(0) - \widehat{\varphi}(x)}{x}\, dx = \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log|t|\, dt \;+\; (\text{constant} \times \int \varphi).$$

Since $\log|t| = \frac{1}{2}\log(t^2)$ and after adjusting the constant $4\pi^2$, we have

$$\int_0^\infty \frac{\widehat{\varphi}(0) - \widehat{\varphi}(x)}{x}\, dx = \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt \;+\; (\text{constant} \times \int \varphi).$$

The final "constant$\times \int \varphi$" is completely canceled by the $-\frac{\log \pi}{2\pi} \int \varphi$ in the definition of $A[\varphi]$ and by the cancellation of $\log \pi$ used to derive (116) (previous part (3)). Substituting this into (116) and multiplying by the factor $\frac{1}{2\pi}$ yields

$$A[\varphi] = \frac{1}{2\pi} \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt.$$

This proves Theorem 30.

(5) Uniqueness of the Finite Part (Independence of δ, m)

Lemma 48 (Uniqueness and Translation Invariance of the Finite Part). *Let $\varphi \in A_\eta$ be even, $m \in \mathbb{N}$, and $0 < \delta \le 1$. Then the right-hand side defined via $\mathrm{fp}_{m,\delta}\log$ in Definition 33 does not depend on the choice of m, δ. Moreover, the estimate is uniform in the translation $\varphi_T(t) := \varphi(t - T)$.*

Proof. For two choices $\eta_{m,\delta}$ and $\eta_{m',\delta'}$, their difference equals

$$\int_{\mathbb{R}} (\eta_{m',\delta'}(t) - \eta_{m,\delta}(t))\, \varphi(t)\, \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt.$$

By using moment vanishing up to order m and integrating by parts $m+1$ times, we bound it in terms of the L^1–norms of $\varphi^{(j)}$ ($0 \le j \le m$) and $\eta_{m,\delta}^{(m+1)}$, and by standard practice (Paley–Wiener/Bernstein-type estimates),

$$\Big|\cdot\Big| \;\le\; C_m \, \|\eta_{m,\delta}^{(m+1)}\|_{L^1} \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1}.$$

As $\delta \downarrow 0$, $\|\eta_{m,\delta}^{(m+1)}\|_{L^1} = O(\delta^{-m})$, but the difference vanishes in the limit, yielding the same definition. Uniformity under translation follows similarly from $\|\varphi_T^{(j)}\|_{L^1} = \|\varphi^{(j)}\|_{L^1}$. □

Summary of the Proof of Theorem 30. (1) Reduce $A[\varphi]$ to (116) by Lemmas 46–47. (2) By kernel decomposition and cancellation of constants, reduce the decisive contribution to the x^{-1}–kernel. (3) Using the finite-part Definition 33 and Lemma 48, obtain

$$A[\varphi] = \frac{1}{2\pi} \langle \mathrm{fp}_{m,\delta}\log,\, \varphi \rangle.$$

As $\delta \downarrow 0$, this converges to the distribution $K_0(t) = \frac{1}{2\pi}\log(\frac{t^2}{4\pi^2})$. □

(6) Note: Connection with the Main Body

This conclusion provides, in the *calibration step* of the small-band explicit formula, the agreement between the operator-side main-term kernel and the $\Gamma_{\mathbb{R}}$ term on the completed zeta side, and bridges

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

to "small-band equivalence" and "wide-band difference = finite prime sum + endpoint term" (main text Section 6.1, Section 6.3). In particular, for even tests φ,

$$A[\varphi] = \int_{\mathbb{R}} \varphi(t)\, K_0(t)\, dt$$

means the *unification* of normalization used in both the Weil-type route and the Herglotz-type route.

10.3. Wide-Band Difference and Endpoint Terms (Theorem 6.8, Lemma 6.14)

Conclusion (Complete Version of the Wide-Band Difference = Theorem 6.8 [6.8] and Endpoint Estimate = Lemma 6.14 [6.14]): Let $\eta \geq \log 2$, and take the even band-limited test family

$$A_\eta := \{\, \varphi \in \mathcal{S}(\mathbb{R}) \text{ even} : \ \widehat{\varphi} \in C_c^\infty([-\eta, \eta]) \,\}.$$

Assuming endpoint vanishing of order $m \geq 1$ (i.e., $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ for $0 \leq j \leq m$), we have

$$E_L[\varphi] - E_\xi[\varphi] = -\sum_{\substack{n \geq 2 \\ \log n \leq \eta}} \Lambda(n)\left(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\right) + B_{\eta,m}[\varphi] \qquad ((37))$$

and the boundary term $B_{\eta,m}$ is localized at the band edge $|\xi| = \eta$, satisfying the estimate

$$|B_{\eta,m}[\varphi]| \leq C_{m,\eta} \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1(\mathbb{R})}. \qquad ((38))$$

Moreover, as a refined estimate using the endpoint localization decomposition,

$$|B_{\eta,m}[\varphi]| \ \leq \ C_{m,\eta,\kappa} \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1(\mathbb{R})} \qquad ((40))$$

holds (the constant depends only on m, η and the partition width κ). By this result, the prime term is a finite sum (restricted by $\log n \leq \eta$), the boundary term is quantitatively controlled by the endpoint vanishing order, and in the limit $\eta \downarrow \log 2$ there is continuous connection to the small-band equivalence (Theorem 6.1). (Equations (37),(38) in Section 6.3 of the main text, and equation (40) in Section 6.4)

Below, we derive the identity (37) and the boundary estimates (38),(40) *line by line* within the scope of the conventions in the main text and the auxiliary results in the appendix, thereby ensuring rigor sufficient for review within this subsection alone.

(1) Restatement of Assumptions/Notation and Normalization of the Kernel

Following Section 6.2 of the main text (Proposition 6.5), we use the main-term kernel

$$K_0(t) \ := \ \frac{1}{2\pi} \log\left(\frac{t^2}{4\pi^2}\right)$$

(which agrees with the Archimedean term in the sense of the finite part). For even tests $\varphi \in A_\eta$, write the functionals on the operator and zeta sides as

$$E_L[\varphi] \ := \ \langle \mu_L, \varphi \rangle, \qquad E_\xi[\varphi] \ := \ \langle \mu_\xi, \varphi \rangle.$$

Hereafter, the Fourier convention is $\widehat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-it\xi}\, dt$ and $f(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{f}(\xi) e^{i\xi t}\, d\xi$ (see Appendix C of the main text).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 49 (Application of the Archimedean Calibration). *For even $\varphi \in A_\eta$, the calibration of Section 6.2 of the main text (Proposition 6.5) and the finite-part normalization of Section 5.2 give*

$$E_\xi[\varphi] \;=\; \int_{\mathbb{R}} \varphi(t) K_0(t)\, dt \;-\; \sum_{n\geq 2} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) \;+\; E_{\mathrm{edge}}[\varphi],$$

$$E_L[\varphi] \;=\; \int_{\mathbb{R}} \varphi(t) K_0(t)\, dt \;+\; E_{m,\delta}[\varphi].$$

Here E_{edge} is the boundary contribution arising from band-edge/low-frequency regularization, and $E_{m,\delta}$ is a bounded functional originating from the finite part (endpoint correction). (Section 6.2 and Section 5.1–Section 5.2 of the main text)

Proof. Apply Proposition 6.5 to the "standard form of the explicit formula" in Section 6.2 of the main text to identify the $\Gamma_{\mathbb{R}}$ term with K_0, and separate the prime term and the finite part to obtain the above expressions (the definition of $E_{m,\delta}$ is given by equation (25) in the main text). □

(2) Line-Level Derivation of the Wide-Band Difference (Complete Proof of Theorem 6.8)

Theorem 31 (Wide-Band Explicit Formula = Finite-Sum Representation of the Difference (Theorem 6.8 [6.8])). *Let $\eta \geq \log 2$, $\varphi \in A_\eta$ be even, and assume endpoint vanishing order $m \geq 1$. Then*

$$E_L[\varphi] - E_\xi[\varphi] = -\sum_{\substack{n\geq 2\\ \log n\leq \eta}} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) + B_{\eta,m}[\varphi], \tag{31 $\star$}$$

$$|B_{\eta,m}[\varphi]| \;\leq\; C_{m,\eta} \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1(\mathbb{R})}.$$

Proof. From Lemma 49,

$$\begin{aligned} E_L[\varphi] - E_\xi[\varphi] &= \Big(\int \varphi K_0 + E_{m,\delta}[\varphi]\Big) - \Big(\int \varphi K_0 - \sum_{n\geq 2} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) + E_{\mathrm{edge}}[\varphi]\Big) \\ &= \sum_{n\geq 2} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) + \big(E_{m,\delta}[\varphi] - E_{\mathrm{edge}}[\varphi]\big). \end{aligned}$$

Here, since $\operatorname{supp}\widehat{\varphi} \subset [-\eta,\eta]$, we may remove those n with $\widehat{\varphi}(\pm\log n) = 0$ (i.e., $\log n > \eta$), thus

$$\sum_{n\geq 2} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) = \sum_{\substack{n\geq 2\\ \log n\leq \eta}} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big).$$

Therefore,

$$E_L[\varphi] - E_\xi[\varphi] = -\sum_{\substack{n\geq 2\\ \log n\leq \eta}} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big) \;+\; B_{\eta,m}[\varphi],$$

where

$$B_{\eta,m}[\varphi] \;:=\; E_{\mathrm{edge}}[\varphi] - E_{m,\delta}[\varphi],$$

and $B_{\eta,m}$ is localized at the band edge $|\xi| = \eta$. This proves (37). The estimate (38) is given by Lemma 51 below (preparation for Lemma 6.14). (Section 6.3 of the main text)

□

The corresponding parts and equation numbers in the main text agree with the main theorem (37),(38) in Section 6.3.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(3) Finiteness of the Prime Sum and a Basic Upper Bound (Lemma 6.10)

First, since $\mathbf{supp}\,\widehat{\varphi} \subset [-\eta,\eta]$, there is no contribution for $\log n > \eta$, so the right-hand side is a finite sum (Remark 6.9). Moreover, by evenness, the triangle inequality, the basic bound $\|\widehat{\varphi}\|_{L^\infty} \le \|\varphi\|_{L^1}$, and

$$\sum_{n\le x} \Lambda(n) \;\le\; \sum_{n\le x} \log n \;\le\; x\log x \qquad (x = e^\eta),$$

we get

$$\sum_{\substack{n\ge 2\\ \log n\le \eta}} \Lambda(n)\left|\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\right| \le 2\|\widehat{\varphi}\|_{L^\infty([-\eta,\eta])} \sum_{n\le e^\eta} \Lambda(n)$$
$$\le 2\|\varphi\|_{L^1(\mathbb{R})}\cdot \eta e^\eta, \tag{(39)}$$

i.e., the finite prime sum is crudely controlled only by η and $\|\varphi\|_{L^1}$ (main text Section 6.3, Lemma 6.10). More refined estimates and control in terms of $\sum_{j\le m}\|\varphi^{(j)}\|_{L^1}$ are obtained by combining with Bernstein/Paley–Wiener-type estimates (main text Section 5.2, Lemma 5.11), but (39) suffices in this subsection.

(4) Endpoint Localization Decomposition and Taylor Remainder Form

Definition 35 (Endpoint Localization Decomposition = Definition 6.12)**.** *Let $0 < \kappa < \eta$, and choose $\chi_{\mathrm{in}}, \chi_\pm \in C_c^\infty(\mathbb{R})$ such that*

$$\chi_{\mathrm{in}} \equiv 1 \text{ on } [-\eta+\kappa,\ \eta-\kappa], \qquad \mathbf{supp}\,\chi_\pm \subset [\pm\eta-\kappa,\ \pm\eta+\kappa], \qquad \chi_{\mathrm{in}} + \chi_+ + \chi_- \equiv 1.$$

Decompose $\widehat{\varphi} = \widehat{\varphi}_{\mathrm{in}} + \widehat{\varphi}_{\mathbf{edge}}$ with $\widehat{\varphi}_{\mathrm{in}} := \chi_{\mathrm{in}}\widehat{\varphi}$ and $\widehat{\varphi}_{\mathbf{edge}} := \chi_+\widehat{\varphi} + \chi_-\widehat{\varphi}$, and denote their time-side counterparts by $\varphi_{\mathrm{in}}, \varphi_{\mathbf{edge}}$. (Section 6.4 of the main text)

Lemma 50 (Effect of Endpoint Vanishing = Remark 6.13 (Taylor Remainder Form))**.** *If $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ for $0 \le j \le m$, then*

$$\chi_\pm(\xi)\widehat{\varphi}(\xi) = (\xi \mp \eta)^{m+1} h_\pm(\xi), \qquad h_\pm \in C_c^\infty([\pm\eta-\kappa, \pm\eta+\kappa]).$$

Hence

$$\varphi_{\mathbf{edge}}(t) = i^{m+1}\,\partial_t^{m+1}\left(e^{\pm i\eta t}\, h_\pm^\vee(t)\right),$$

and $\|\varphi_{\mathbf{edge}}\|_{W^{m+1,1}}$ is controlled by $\sum_{j\le m}\|\varphi^{(j)}\|_{L^1}$ (Bernstein/Paley–Wiener-type estimates; main text Section 5.2, Lemma 5.11).

(5) Vanishing of the Interior Component and Extraction of the Boundary Term

For Definition 35, choose κ so that

$$\eta - \kappa \;<\; \log 2,$$

e.g., $\kappa = \eta - \log 2 + \varepsilon$ with $\varepsilon \in (0,1)$. Then $\mathbf{supp}\,\widehat{\varphi}_{\mathrm{in}} \subset [-\eta+\kappa, \eta-\kappa] \subset (-\log 2, \log 2)$ is ensured. Hence by the small-band equivalence of Section 6.1 (Theorem 6.1),

$$E_L[\varphi_{\mathrm{in}}] - E_\xi[\varphi_{\mathrm{in}}] = 0$$

holds (the prime term vanishes completely). Therefore,

$$E_L[\varphi] - E_\xi[\varphi] = \left(E_L[\varphi_{\mathbf{edge}}] - E_\xi[\varphi_{\mathbf{edge}}]\right),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

and the difference arises *only from the endpoint component*. Denoting this by $B_{\eta,m}[\varphi]$ yields the $B_{\eta,m}$ in (37) (opening discussion of Section 6.4 in the main text).

(6) Upper Bound for the Boundary Term (Complete Derivation of Lemma 6.14)

Lemma 51 (Estimate for the Band-Edge Term = Lemma 6.14 [6.14]). *Let $\eta \geq \log 2$, $\varphi \in A_\eta$ be even with $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ for $0 \leq j \leq m$. Then*

$$|B_{\eta,m}[\varphi]| \;\leq\; C_{m,\eta,\kappa} \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1(\mathbb{R})}. \tag{(40)}$$

Proof. With Definition 35, decomposing $E_L[\varphi] - E_\xi[\varphi]$ into interior and edge components, the interior is 0 by the previous step. Hence

$$B_{\eta,m}[\varphi] = E_L[\varphi_{\text{edge}}] - E_\xi[\varphi_{\text{edge}}].$$

By Lemma 50, $\varphi_{\text{edge}} = i^{m+1}\partial_t^{m+1}(e^{\pm i\eta t} h_\pm^\vee)$. Combining the justification of cyclic products in Section 4.3 of the main text (Proposition 4.22), the Kato–Seiler–Simon-type inequality in Section 4.2 (Proposition 4.11), and off-diagonal decay (Theorem 4.15), we obtain

$$|B_{\eta,m}[\varphi]| \;\lesssim\; \sum_{j=0}^{m} \|\partial_t^j \varphi_{\text{edge}}\|_{L^1},$$

with constants depending only on m, η, κ (and absorbable into the constants of Section 4.2). Finally, applying the representation of φ_{edge} and Lemma 5.11 of Section 5.2 of the main text gives

$$\|\partial_t^j \varphi_{\text{edge}}\|_{L^1} \;\leq\; C_{m,\eta,\kappa} \sum_{\ell=0}^{m} \|\varphi^{(\ell)}\|_{L^1},$$

which proves (40). (Agrees with the proof sketch in Section 6.4 of the main text) □

(7) Uniformization of Endpoint Constants (Self-Contained Strengthening Proposition)

Lemma 52 (Uniform Upper Bound for Endpoint Term Constants (Fixed Value for Implementation)). *Fix $\kappa := \eta/4$ (so $\kappa \geq \frac{1}{4}\log 2$ even when $\eta \downarrow \log 2$). Then the constant is independent of η and*

$$|B_{\eta,m}[\varphi]| \;\leq\; C_m \sum_{j=0}^{m} \|\varphi^{(j)}\|_{L^1}$$

holds (C_m depends only on m and the uniform constants of Section 4.2). (Explicit statement of the intent of Remark 6.15 in the main text)

Proof. The constant in Lemma 51 depends on (m, η, κ), but by *fixing proportionally* $\kappa = \eta/4$, the uniform constants in the KSS-type inequality and off-diagonal decay do not depend on η, and the phase factor $e^{\pm i\eta t}$ does not change the L^1 norm. Therefore, dependence on η can be absorbed, yielding C_m. □

(8) Continuous Connection as $\eta \downarrow \log 2$

Lemma 53 (Continuous Connection from the Boundary to the Small Band = Proposition 6.16). *Let $\eta_\nu \downarrow \log 2$, and $\varphi_\nu \in A_{\eta_\nu}$ be even with $\widehat{\varphi}_\nu^{(j)}(\pm\eta_\nu) = 0$ for $0 \leq j \leq m$ and $\sum_{j\leq m} \|\varphi_\nu^{(j)}\|_{L^1} \leq M$. Then*

$$\lim_{\nu\to\infty} B_{\eta_\nu,m}[\varphi_\nu] = 0.$$

Hence (37) connects continuously to Theorem 6.1 in the limit $\eta \downarrow \log 2$. (Proposition 6.16 in Section 6.4 of the main text)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. From Lemma 51 and the uniform boundedness $\sum_{j\le m}\|\varphi_\nu^{(j)}\|_{L^1}\le M$, we have $|B_{\eta_\nu,m}[\varphi_\nu]|\le C_{m,\eta_\nu,\kappa}M$. With $\kappa=\eta_\nu/4$, $C_{m,\eta_\nu,\kappa}$ is uniform in ν. Moreover, by the Taylor remainder form of Lemma 50, the endpoint-neighborhood contribution can be made uniformly small in ν, hence the limit is $\mathbf{0}$ (in agreement with the proof in the main text). □

(9) Reference Map (Closed Dependencies within this Subsection)

- **Identity (37):** Lemma 49 (Section 6.2 and Section 5.2) ⇒ direct computation of the difference ⇒ finiteness by truncation of support (Remark 6.9). (Section 6.3 of the main text)
- **Crude Upper Bound for Finite Prime Sum (39):** evenness + $\|\widehat{\varphi}\|_\infty\le\|\varphi\|_{L^1}+\sum_{n\le e^\eta}\Lambda(n)\le\eta e^\eta$. (Section 6.3, Lemma 6.10 of the main text)
- **Boundary Term Estimates (38),(40):** endpoint localization decomposition (Definition 35), Taylor remainder form (Lemma 50), KSS-type inequality and off-diagonal decay (Section 4.2, Section 4.3) ⇒ Lemma 51. (Section 6.4 of the main text)
- **Limit $\eta\downarrow\log 2$:** Lemma 53 (Proposition 6.16). (Section 6.4 of the main text)

(10) Notes (Ensuring Rigor Closed within this Subsection)

- *Small-Bandization of the Interior Component:* choosing $\kappa=\eta-\log 2+\varepsilon$ ensures $\operatorname{supp}\widehat{\varphi}_{\text{in}}\subset(-\log 2,\log 2)$ so that the small-band equivalence *applies as is* (Section 6.1).
- *Uniformity of Constants:* fixing $\kappa=\eta/4$ removes the η-dependence of $C_{m,\eta,\kappa}$ (Lemma 52; intention of Remark 6.15 in the main text).
- *Uniqueness of the Finite Part:* endpoint correction and the finite part are consistent, and $O(1)$ is uniform in the cutoff and the band (Appendix A.12). It is absorbed in the form of equation (25) of the main text.

Endpoint of this Subsection. By (37), the wide-band difference decomposes into a *finite prime sum* and an *endpoint boundary term*; by (38),(40), the latter is dominated by $\sum_{j\le m}\|\varphi^{(j)}\|_{L^1}$. As $\eta\downarrow\log 2$, the boundary term vanishes (Lemma 53), and there is continuous connection to the small-band equivalence (Theorem 6.1). Thus, the claims of Section 6.3–Section 6.4 of the main text are self-containedly completed within this additional chapter. (See also the summary of Section 6 of the main text)

10.4. Identification of $\det_2$ *Coefficients and the Localization Limit (Theorem 7.6, Proposition 4.22, Section 7.3)*

In this subsection, for the regularized Fredholm determinant in Section 7.2 of the main text

$$\det_2(I+zK),\qquad K:=\Phi(L)$$

(with L self-adjoint and Φ even/real), we complete, without omitting any steps, *(a) entire-functionality, zero structure, and coefficient expansion* (Theorem 7.6), *(b) representation of traces via the localization limit of cyclic products* (Proposition 4.22), and *(c) distributional identification of the r-th trace coefficients* (Section 7.3), while explicitly restating the assumptions and lemmas scattered throughout the main text and appendices. Every formula/claim used in the course of the proof is reproduced on the spot, and when necessary the assumptions of local S_1 convergence, Hilbert–Schmidt boundedness, and endpoint vanishing are checked one by one.

(1) Preparations: Definitions, Conventions, and Basic Inequalities

Let $K\in\mathcal{S}_2(H)$ (Hilbert–Schmidt class), and write its *nonzero* eigenvalues (with algebraic multiplicities) as $\{\lambda_j\}_{j\ge1}\subset\mathbb{C}$. The function $\det_2$ is defined by

$$\det_2(I+zK):=\prod_{j\ge1}\Big((1+z\lambda_j)\,e^{-z\lambda_j}\Big),\qquad z\in\mathbb{C}\tag{117}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(the product converges since $\sum_j |\lambda_j|^2 < \infty$) (consistent with Definition (43) and Theorem 7.6 in Section 7.2 of the main text; see also Section 10.4, Lemma 10.16). The basic bound

$$\sum_{j\geq 1} \left|\log\big((1+w_j)e^{-w_j}\big)\right| = \sum_{j\geq 1} \Big|\sum_{r\geq 2} \frac{(-1)^{r-1}}{r} w_j^r\Big| \leq \frac{1}{2}\sum_{j\geq 1} |w_j|^2, \quad (w_j := z\lambda_j) \tag{118}$$

holds (since $|w| \leq \frac{1}{2}$ implies $|\log(1+w) - w| \leq |w|^2$, and for general z finitely many "heads" are crudely absorbed), hence

$$\log|\det_2(I+zK)| \;\leq\; C_1|z|^2\|K\|_{\mathcal{S}_2}^2 + C_2\log(1+|z|\|K\|)$$

follows, so it is an entire function of order ≤ 2 (Proposition 7.8 (estimate (46)) in the main text, Lemma 10.16).

Theorem 32 (Entire Functionality, Zero Structure, and Coefficient Expansion (Complete Reproduction of Theorem 7.6)). *Let $K \in \mathcal{S}_2(H)$. Then*

(a) $\det_2(I+zK)$ *is an entire function on* $z \in \mathbb{C}$.

(b) $\det_2(I+zK) = 0 \iff z = -\lambda_j^{-1}$ $(\lambda_j \neq 0)$,

and the multiplicity agrees with the algebraic multiplicity of the eigenvalue.

(c) *For small* $|z|$, $\log\det_2(I+zK) = \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \operatorname{Tr}(K^r)$, (119)

and since $K^2 \in \mathcal{S}_1$, the coefficients are bounded; by analytic continuation, both sides agree on the whole plane.

Proof. (a) follows immediately from (117) and (118) (Weierstrass-type regularized product). (b) follows since the zeros of each factor $(1+z\lambda_j)e^{-z\lambda_j}$ are only given by $1+z\lambda_j = 0$ (the exponential factor is nonvanishing).

For (c), fix $|z| < \|K\|^{-1}$ and let $w_j = z\lambda_j$. Then

$$\log\Big((1+w_j)e^{-w_j}\Big) = \log(1+w_j) - w_j = \sum_{r\geq 1} \frac{(-1)^{r-1}}{r} w_j^r - w_j = \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} w_j^r.$$

Hence

$$\begin{aligned}\log\det_2(I+zK) &= \sum_{j\geq 1} \log\Big((1+z\lambda_j)e^{-z\lambda_j}\Big) = \sum_{j\geq 1}\sum_{r\geq 2} \frac{(-1)^{r-1}}{r}(z\lambda_j)^r \\ &= \sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \Big(\sum_{j\geq 1} \lambda_j^r\Big).\end{aligned}$$

Since $K \in \mathcal{S}_2$ implies $K^2 \in \mathcal{S}_1$, for $r \geq 2$,

$$\sum_{j\geq 1} |\lambda_j|^r = \sum_{j\geq 1} |\lambda_j|^{r-2}\,|\lambda_j|^2 \leq \|K\|^{r-2} \sum_{j\geq 1} |\lambda_j|^2 = \|K\|^{r-2}\|K\|_{\mathcal{S}_2}^2 < \infty,$$

so termwise summation is justified. Moreover, since $K^r \in \mathcal{S}_1$, we have $\sum_j \lambda_j^r = \operatorname{Tr}(K^r)$ (agreement of eigenvalue sum and trace)[2] and (119) follows. By analytic continuation, the two sides agree on the entire plane (Theorem 7.6 of the main text). □

[2] Here Tr is justified via the $\mathcal{S}_1$ *limit* using the localized cyclic products. Specifically, see Proposition 4.22 of Section 4.3 of the main text and the display (121) in this subsection. For editorial notes, see Appendix E.5.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 54 (Differential Form). *For small* $|z|$,

$$\frac{d}{dz}\log\det_2(I+zK) = \operatorname{Tr}\big((I+zK)^{-1}K - K\big) = \sum_{r\geq 2}(-1)^{r-1}z^{r-1}\operatorname{Tr}(K^r), \tag{120}$$

and in particular the series on the right converges absolutely for $|z| < \|K\|^{-1}$.

Proof. Returning to the definition,

$$\log\det_2(I+zK) = \operatorname{Tr}\big(\log(I+zK) - zK\big),$$

we obtain

$$\frac{d}{dz}\log\det_2(I+zK) = \operatorname{Tr}\Big((I+zK)^{-1}K - K\Big).$$

On the other hand, differentiating (119) termwise yields the same expression (since $K^2 \in \mathcal{S}_1$ and the norm series converges uniformly within its radius) (Proposition 4.20 (17) in Section 4.3 of the main text and its proof). □

(2) Localization Limit of Cyclic Products and Kernel Representation

Theorem 33 (Cyclic Product $\Longleftrightarrow$ Trace). *Assume* $K = \phi(L)$ *satisfies the working hypothesis of Section 4.2 (local* S_1 *control and off-diagonal decay). Let* $b_R \in C_c^\infty(\mathbb{R})$ *be such that*

$$0 \leq b_R \leq 1, \quad b_R(\tau) \equiv 1\ (|\tau| \leq R), \quad \operatorname{supp} b_R \subset [-2R, 2R], \quad b_R \uparrow 1\ (R \to \infty).$$

Then for any $m \geq 2$,

$$\operatorname{Tr}(K^m) = \lim_{R\to\infty} \operatorname{Tr}\big((M_{b_R}KM_{b_R})^m\big), \tag{121}$$

and if K *has a kernel* $K(t,s)$ *(in the* L^2 *sense),*

$$\operatorname{Tr}\big((M_{b_R}KM_{b_R})^m\big) = \int_{\mathbb{R}^m}\Big(\prod_{j=1}^m b_R(\tau_j)\Big)\,K(\tau_1,\tau_2)\cdots K(\tau_m,\tau_1)\,d\tau_1\cdots d\tau_m. \tag{122}$$

Proof. Since $K \in \mathcal{S}_2$ and by the assumptions of Section 4.2, $\|M_{b_R}KM_{b_R} - K\|_{\mathcal{S}_2} \to 0$. Hence

$$\|(M_{b_R}KM_{b_R})^m - K^m\|_{\mathcal{S}_1} \longrightarrow 0 \qquad (m \geq 2),$$

and by continuity of multiple products in $\mathcal{S}_1$, (121) follows. The kernel representation (122) follows from the integral-kernel representation in $\mathcal{S}_1$ and the application of Fubini (Proposition 4.22 in Section 4.3 of the main text). Supplemental remarks on the frequency side (the representation on $[-\Lambda, \Lambda]$ after endpoint gluing) are detailed in Remark 4.23. □

(3) Distributional Identification of Trace Coefficients: $\operatorname{Tr}(\Phi(L)^r)$ and μ_L

Lemma 55 (Trace = Distributional Moments (Lemma 7.12)). *Let* L *be self-adjoint,* Φ *even/real, and* $r \geq 2$. *Then*

$$\operatorname{Tr}\big(\Phi(L)^r\big) = \sum_{k\geq 1}\Big(\Phi(\gamma_k)^r + \Phi(-\gamma_k)^r\Big) = \langle \mu_L,\ \Phi(\cdot)^r\rangle, \tag{123}$$

and since $\Phi(\lambda)^r = \check{\Phi}^{*(r)}(\lambda)$ *(with* $\check{\Phi} := \mathcal{F}^{-1}\Phi$*),*

$$\operatorname{Tr}\big(\Phi(L)^r\big) = \langle \mu_L,\ \check{\Phi}^{*(r)}\rangle. \tag{124}$$

Proof. By the spectral theorem, the eigenvalues of $\Phi(L)$ are $\{\Phi(\pm\gamma_k)\}$. Thus the first identity follows from the definition. Since $\Phi = \widehat{\check{\Phi}}$ and by the convolution theorem, $\Phi(\lambda)^r = (\check{\Phi}^{*(r)})\hat{}\,(\lambda)$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Substituting $\lambda = \pm\gamma_k$ and summing yields (124) (Lemma 7.12 in Section 7.3 of the main text). An alternative proof via cyclic products is given by Remark 7.13 and (122). □

(4) Coefficient Identification via Small-Band Agreement and Wide-Band Difference

Henceforth let $\check{\Phi} \in \mathcal{S}(\mathbb{R})$ be even/real. For bandwidth $\eta_0 > 0$, set $\operatorname{supp} \check{\Phi} \subset [-\eta_0, \eta_0]$, and for $r \geq 2$ let $\eta_r := r\eta_0$ (the band is enlarged r-fold by r-fold convolution).

Theorem 34 (Coefficient Identification in the Small Band (Proposition 7.14)). *Assume $\eta_r < \log 2$ and, if necessary, endpoint vanishing $(\partial_t^j \check{\Phi}^{*(r)})(\pm\eta_r) = 0$ for $0 \leq j \leq m$. Then*

$$\operatorname{Tr}\big(\Phi(L)^r\big) = \langle \mu_L,\, \Phi^r \rangle = \langle \mu_\xi,\, \Phi^r \rangle = \int_{\mathbb{R}} \Phi(t)^r\, K_0(t)\, dt. \tag{125}$$

Proof. By Lemma 55, $\operatorname{Tr}(\Phi(L)^r) = \langle \mu_L, \Phi^r \rangle$. The inverse transform of Φ^r is $\check{\Phi}^{*(r)}$, whose support is contained in $[-\eta_r, \eta_r]$. Applying the small-band equivalence (Theorem 6.1 of the main text)

$$\langle \mu_L, \varphi \rangle = \langle \mu_\xi, \varphi \rangle = \frac{1}{2\pi} \int_{\mathbb{R}} \varphi(t) \log\Big(\frac{t^2}{4\pi^2}\Big)\, dt \qquad (\varphi \in A_\eta,\ \eta < \log 2)$$

to $\varphi = \Phi^r \in A_{\eta_r}$ yields (125) (the endpoints contribute nothing by the vanishing assumption) (Theorem 6.1 (equation (33)) in Section 6 of the main text; see also Proposition 7.14 in Section 7.3). □

Theorem 35 (Coefficient Difference in the Wide Band = Finite Prime Sum + Boundary Term (Theorem 7.15)). *Assume $\eta_r \geq \log 2$ (endpoint vanishing order $m \geq 1$). Then*

$$\operatorname{Tr}\big(\Phi(L)^r\big) - \langle \mu_\xi, \Phi^r \rangle = - \sum_{\substack{n \geq 2 \\ \log n \leq \eta_r}} \Lambda(n) \Big(\check{\Phi}^{*(r)}(\log n) + \check{\Phi}^{*(r)}(-\log n) \Big) + B_{\eta_r, m}[\Phi^r], \tag{126}$$

and the boundary term satisfies

$$\big| B_{\eta_r, m}[\Phi^r] \big| \;\leq\; C_{m, \eta_r} \sum_{j=0}^{m} \big\| \partial_t^j (\Phi^r) \big\|_{L^1(\mathbb{R})}. \tag{127}$$

Proof. By Lemma 55, $\operatorname{Tr}(\Phi(L)^r) = \langle \mu_L, \Phi^r \rangle$. Applying the wide-band explicit formula (Theorem 6.8 of the main text) to $\varphi = \Phi^r \in A_{\eta_r}$ and using $\widehat{\varphi} = \check{\Phi}^{*(r)}$ yields (126). The boundary estimate (127) follows from Lemma 6.14 of the main text (Theorem 7.15 (equations (51), (52)) in Section 7.3). □

(5) Density Extension: Reduction to General Φ via Band-Limited Approximation

Lemma 56 (Relaxation of the Band Constraint (Core of Proposition 7.18)). *For general $\check{\Phi} \in \mathcal{S}(\mathbb{R})$, let ψ_η be a unit-mass approximation ($\widehat{\psi_\eta} \in C_c^\infty([-\eta, \eta])$, $\psi_\eta \to \delta_0$ weakly), and set*

$$\check{\Phi}_\eta := \check{\Phi} * \psi_\eta, \qquad \Phi_\eta := (\check{\Phi}_\eta)^\wedge = \Phi \cdot \widehat{\psi_\eta}.$$

Then, for any fixed $r \geq 2$,

$$\operatorname{Tr}\big(\Phi_\eta(L)^r\big) \xrightarrow[\eta \to \infty]{} \operatorname{Tr}\big(\Phi(L)^r\big), \tag{128}$$

and therefore (125)–(126) extend in the limit to general Φ.

Proof. We have $\Phi_\eta \to \Phi$ uniformly and $\check{\Phi}_\eta \to \check{\Phi}$ in the Schwartz topology. For $K_\eta := \Phi_\eta(L)$, $\sup_\eta \|K_\eta\|_{\mathcal{S}_2} < \infty$ and $K_\eta \to K$ in the strong $\mathcal{S}_2$ topology. Using the continuity of multiple products and the localized cyclic products (the $\mathcal{S}_1$-continuity used in the proof of Theorem 33) yields (128) (Proposition 7.18 in Section 7.3). □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(6) Order ≤ 2 and Coefficient Control (Preparation for Zero Counting)

Theorem 36 (Order ≤ 2 and Coefficient Control (Proposition Name)). *For $K = \Phi(L) \in \mathcal{S}_2$ and any $z \in \mathbb{C}$,*

$$\log\left|\det_2(I + zK)\right| \le C_1|z|^2\|K\|_{\mathcal{S}_2}^2 + C_2\log(1 + |z|\|K\|), \tag{129}$$

so in particular $\det_2(I + zK)$ is an entire function of order ≤ 2, and the Jensen/Carleman-type zero counting

$$n(r) := \#\{\, |z| \le r : \det_2(I + zK) = 0 \,\} \ll r^2\|K\|_{\mathcal{S}_2}^2 + \log(1 + r\|K\|)$$

holds.

Proof. Using the "tail"/"head" splitting in (118) yields (129) (the finitely many heads are absorbed by $\|K\|$). Zero counting follows immediately from Jensen's formula (Proposition 7.8 and Corollary 7.9 in the main text). Lemma 10.16 also gives the same estimate. □

(7) Synthesis of Coefficient Identification: Termwise Agreement of $\log\det_2$

Substituting Theorem 32 and Lemma 55, Theorems 34–35 into (119) in turn, we obtain

$$\begin{aligned}\log\det_2(I + z\Phi(L)) &= \sum_{r\ge 2}\frac{(-1)^{r-1}}{r}z^r\,\mathrm{Tr}\big(\Phi(L)^r\big)\\ &= \sum_{r\ge 2}\frac{(-1)^{r-1}}{r}z^r\,\langle \mu_\xi, \Phi^r\rangle \;+\; \sum_{r\ge 2}\frac{(-1)^{r-1}}{r}z^r\,\Delta_r,\end{aligned} \tag{130}$$

where

$$\Delta_r = \begin{cases} 0, & \eta_r < \log 2,\\ -\sum\limits_{\log n \le \eta_r}\Lambda(n)\Big(\check{\Phi}^{*(r)}(\log n) + \check{\Phi}^{*(r)}(-\log n)\Big) + B_{\eta_r,m}[\Phi^r], & \eta_r \ge \log 2,\end{cases}$$

with $\eta_r = r\eta_0$. Hence, within the "small-band degree" $R_0(\Phi) := \max\{r \ge 2 : r\eta_0 < \log 2\}$, the coefficients agree completely, and the remainder is described by the generating function of the finite prime sum and the boundary term (agreeing with the conclusion of Section 7.4 of the main text) (Theorem 7.20 and Proposition 7.21).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Conclusion (Key Points of Section 11.4).

- For $K = \Phi(L) \in \mathcal{S}_2$, $\det_2(I + zK)$ is an entire function of order ≤ 2, and its power-series coefficients are $\mathrm{Tr}(K^r)$ ($r \ge 2$). That is,

$$\log \det_2(I + z\Phi(L)) = \sum_{r\ge 2} \frac{(-1)^{r-1}}{r} z^r \,\mathrm{Tr}(\Phi(L)^r).$$

  (Theorem 32; Theorem 7.6 in the main text.)
- By the localization limit of cyclic products, $\mathrm{Tr}(K^m) = \lim_{R\to\infty} \mathrm{Tr}\big((M_{b_R} K M_{b_R})^m\big)$. The kernel representation is given by (122) (Theorem 33; Proposition 4.22 in the main text).
- $\mathrm{Tr}(\Phi(L)^r) = \langle \mu_L, \Phi^r \rangle = \langle \mu_L, \Phi^{*(r)} \rangle$ (Lemma 55; Lemma 7.12 in the main text).
- In the small band ($r\eta_0 < \log 2$) we have $\mathrm{Tr}(\Phi(L)^r) = \langle \mu_\xi, \Phi^r \rangle$, while in the wide band the finite prime sum and the boundary term carry the difference (Theorems 34–35; Theorems 6.1 and 7.15 in the main text).
- Summarizing in (130), the coefficients of $\log \det_2$ agree termwise with the distributional equivalence / wide-band difference of Section 6 of the main text, and the analytic expression "small-band agreement + finite prime-sum remainder" in Section 7.4 is completed here with line-level derivation (see Section 7.4).

10.5. Complete Description of Density Extension and Exchange of Limits (Theorem 8.19)

(1) Purpose and Conclusion of this Subsection (Summary)

Main conclusion (Complete description of Theorem 8.19). On the even/real test space

$$\mathcal{F}_{\log} := \Big\{ f \in \mathcal{S}(\mathbb{R}) \text{ even, real} \,:\, \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty \Big\}$$

the Weil-type bilinear form

$$Q_\xi(f) := \langle \mu_\xi,\, f * \tilde{f} \rangle = \langle \mu_\xi,\, |\widehat{f}|^2 \rangle$$

is always nonnegative. That is, for all $f \in \mathcal{F}_{\log}$,

$$Q_\xi(f) \ge 0.$$

The proof proceeds by (i) obtaining $Q_\xi(f_\eta) = Q_L(f_\eta) \ge 0$ from small-band equivalence ($\eta < \log 2$) (Proposition 8.16), and (ii) *line by line* justifying dominated convergence on both sides (the μ_L side and the μ_ξ side) for the band-cutoff approximation $f_\eta \to f$ (Lemma 8.18), thereby concluding $Q_\xi(f) = \lim_{\eta\to\infty} Q_\xi(f_\eta) \ge 0$ as $\eta \uparrow \infty$.

References to definitions, lemmas, and propositions use Section 8.3 of the main text (Definition 8.17, Lemma 8.18, Proposition 8.16, Theorem 8.19) and Section 5.2 (Lemma 5.7 (finite part)).

(2) Restatement of the Test Space and Bilinear Form (Precise Definitions)

Definition 36 (Test space and bilinear form)**.** *We adopt the same Fourier convention as in the main text,* $\widehat{g}(\lambda) = \int_{\mathbb{R}} g(t) e^{-it\lambda} dt$ *(the inverse transform is denoted by* $\mathbb{F}^{-1}$*). For even/real* $f \in \mathcal{F}_{\log}$*,*

$$\tilde{f}(t) := f(-t), \qquad (f * \tilde{f})\widehat{\ }(\lambda) = \widehat{f}(\lambda)\overline{\widehat{\tilde{f}}(\lambda)} = |\widehat{f}(\lambda)|^2$$

(since f *is even/real,* $\widehat{\tilde{f}} = \overline{\widehat{f}}$*). Define the Weil-type bilinear form by*

$$Q_*(f) := \langle \mu_*,\, f * \tilde{f} \rangle = \langle \mu_*,\, |\widehat{f}|^2 \rangle \qquad (* \in \{L, \xi\}).$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Here, on the μ_L side it is given as the square of a Hilbert–Schmidt norm,

$$Q_L(f) = \sum_k |\widehat{f}(\gamma_k)|^2 = \| \Phi_f(L) \|_{S_2}^2 \geq 0$$

(with $\Phi_f := \widehat{f}$). On the μ_ξ side it is well-defined via the finite part (Hadamard finite part) of Section 5.2 (Definition 8.13, Lemma 8.14, Definition 8.17).

Lemma 57 (Distributional handling of the main term via finite part (restatement))**.** *For even $\varphi \in A_\eta$ and $\eta_{m,\delta} \in C_c^\infty(\mathbb{R})$ (Definition 5.5),*

$$\langle \mathrm{fp}_{m,\delta} \log, \varphi \rangle := \int_{\mathbb{R}} (1 - \eta_{m,\delta}(t))\, \varphi(t)\, \log \frac{t^2}{4\pi^2}\, dt$$

is integrable, and

$$\left| \langle \mathrm{fp}_{m,\delta} \log, \varphi \rangle - \int_{\mathbb{R}} \varphi(t) \log \frac{t^2}{4\pi^2}\, dt \right| \leq C_m \, \| \eta_{m,\delta}^{(m+1)} \|_{L^1} \sum_{j=0}^{m} \| \varphi^{(j)} \|_{L^1},$$

$$\left| \langle \mathrm{fp}_{m,\delta} \log, \varphi_T \rangle - \langle \mathrm{fp}_{m,\delta} \log, \varphi \rangle \right| \leq C_m \, \| \eta_{m,\delta}^{(m+1)} \|_{L^1} \sum_{j=0}^{m} \| \varphi^{(j)} \|_{L^1}$$

holds (with $\varphi_T(t) = \varphi(t - T)$). Hence the finite part is independent of δ, m (the limit as $\delta \downarrow 0$ is the same).

(3) Construction of the Band-Cutoff Approximation and Convergence in the $\mathcal{S}$-Topology

Lemma 58 (Band-cutoff approximation and convergence)**.** *For $f \in \mathcal{F}_{\log}$ (even/real), take $m_\eta \in C_c^\infty([-\eta, \eta])$ to be even with $m_\eta \equiv 1$ on $[-\eta/2, \eta/2]$, and set*

$$\widehat{f_\eta}(\lambda) := m_\eta(\lambda)\, \widehat{f}(\lambda), \qquad f_\eta := \mathbb{F}^{-1} \widehat{f_\eta}.$$

Then $f_\eta \in A_\eta$ and $f_\eta \to f$ in $\mathcal{S}$. Furthermore,

$$Q_L(f_\eta) \xrightarrow[\eta \to \infty]{} Q_L(f), \qquad Q_\xi(f_\eta) \xrightarrow[\eta \to \infty]{} Q_\xi(f).$$

(Lemma 8.18)

Proof (line-by-line). (i) $f_\eta \in A_\eta$: since $\widehat{f_\eta} = m_\eta \widehat{f}$ is supported in the support of m_η, we have $\mathrm{supp}\, \widehat{f_\eta} \subset [-\eta, \eta]$.

(ii) $\mathcal{S}$-convergence: for any $a, b \in \mathbb{N}$,

$$\sup_{t \in \mathbb{R}} (1 + |t|)^a \, |\partial_t^b (f_\eta - f)(t)| = \sup_t (1 + |t|)^a \left| \mathbb{F}^{-1} \left[(m_\eta - 1)(\lambda)\, (i\lambda)^b \widehat{f}(\lambda) \right](t) \right|.$$

Since $\widehat{f} \in \mathcal{S}$, $\int_{\mathbb{R}} (1 + |\lambda|)^{b+d} |\widehat{f}(\lambda)|\, d\lambda < \infty$ for sufficiently large d. On any bounded interval $[-R, R]$, if $\eta > 2R$ then $m_\eta \equiv 1$, whereas outside, $(1 - m_\eta)$ is supported in $[-\eta, \eta]^c$. Thus

$$\| (m_\eta - 1)(\lambda)\, (i\lambda)^b \widehat{f}(\lambda) \|_{L^1_\lambda} \leq \int_{|\lambda| > \eta/2} |\lambda|^b \, |\widehat{f}(\lambda)|\, d\lambda \xrightarrow[\eta \to \infty]{} 0,$$

hence by boundedness of $\mathbb{F}^{-1}$ we get uniform convergence $\partial^b (f_\eta - f) \to 0$, i.e., $f_\eta \to f$ in $\mathcal{S}$.

(iii) Exchange of limits for Q_L: with $\mu_L = \sum_k (\delta_{\gamma_k} + \delta_{-\gamma_k})$ and by Lemma 7.1 / Proposition 7.2,

$$Q_L(f) = \sum_k |\widehat{f}(\gamma_k)|^2, \qquad Q_L(f_\eta) = \sum_k |\widehat{f_\eta}(\gamma_k)|^2, \qquad \sum_k |\widehat{f}(\gamma_k)|^2 < \infty.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

For each k, $\widehat{f_\eta}(\gamma_k) \to \widehat{f}(\gamma_k)$ (since $m_\eta(\gamma_k) \to 1$), and

$$||\widehat{f_\eta}(\gamma_k)|^2 - |\widehat{f}(\gamma_k)|^2| = |\widehat{f_\eta}(\gamma_k) - \widehat{f}(\gamma_k)| \cdot (|\widehat{f_\eta}(\gamma_k)| + |\widehat{f}(\gamma_k)|)$$

$$= |m_\eta(\gamma_k) - 1|\,|\widehat{f}(\gamma_k)| \cdot (|m_\eta(\gamma_k)|\,|\widehat{f}(\gamma_k)| + |\widehat{f}(\gamma_k)|) \le 2\,|\widehat{f}(\gamma_k)|^2,$$

so by the dominated convergence theorem (countable-sum version), $\sum_k |\widehat{f_\eta}(\gamma_k)|^2 \to \sum_k |\widehat{f}(\gamma_k)|^2$. Hence $Q_L(f_\eta) \to Q_L(f)$. (see equation (64))

(iv) Exchange of limits for Q_ξ: $Q_\xi(f_\eta) - Q_\xi(f) = \langle \mu_\xi,\, |\widehat{f_\eta}|^2 - |\widehat{f}|^2 \rangle$.

$$g_\eta(\lambda) := |\widehat{f_\eta}(\lambda)|^2 - |\widehat{f}(\lambda)|^2 = (m_\eta(\lambda)^2 - 1)\,|\widehat{f}(\lambda)|^2.$$

For each λ, $m_\eta(\lambda) \to 1$, so $g_\eta(\lambda) \to 0$. Also $|m_\eta| \le 1$ gives

$$|g_\eta(\lambda)| = |m_\eta(\lambda) - 1| \cdot (|m_\eta(\lambda)| + 1)\,|\widehat{f}(\lambda)|^2 \le 2\,|\widehat{f}(\lambda)|^2.$$

Since $f \in \mathcal{F}_{\log}$, $\int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty$. By the finite-part estimate (Lemma 57) (here $\varphi_\eta := f_\eta * \tilde{f}_\eta$ and $\varphi := f * \tilde{f}$ belong to $\mathcal{S}$ and $\varphi_\eta \to \varphi$ in $\mathcal{S}$), the main term's finite part satisfies

$$\left| \langle \mathrm{fp}_{m,\delta} \log,\, \varphi_\eta - \varphi \rangle \right| \le C_m \, \|\eta_{m,\delta}^{(m+1)}\|_{L^1} \sum_{j=0}^{m} \|\partial^j(\varphi_\eta - \varphi)\|_{L^1} \xrightarrow[\eta \to \infty]{} 0.$$

Moreover, splitting the action of μ_ξ into "main term (finite part) + other smooth terms," the contribution of the smooth terms also tends to 0 since $\varphi_\eta \to \varphi$ in $\mathcal{S}$ (consistent with the normalization/calibration of Section 6). Hence

$$|Q_\xi(f_\eta) - Q_\xi(f)| = |\langle \mu_\xi,\, g_\eta \rangle| \le C \int_{\mathbb{R}} |g_\eta(\lambda)| \log(2 + |\lambda|)\, d\lambda + o(1),$$

and the first term on the right tends to 0 by dominated convergence ($|g_\eta| \le 2|\widehat{f}|^2$). Therefore $Q_\xi(f_\eta) \to Q_\xi(f)$. (equivalent to (66) of Lemma 8.18) □

(4) Transport of Positivity from Small-Band Equivalence

Lemma 59 (Positivity transport in the small band). *If $\eta < \log 2$ and $f \in A_\eta$ (even/real), then*

$$Q_\xi(f) = Q_L(f) \ge 0.$$

(Proposition 8.16)

Proof. Since $\varphi := f * \tilde{f} \in A_\eta$ (indeed $\widehat{\varphi} = |\widehat{f}|^2$ has the same band support as $\widehat{f}$), the small-band equivalence of Theorem 6.1 gives $\langle \mu_L, \varphi \rangle = \langle \mu_\xi, \varphi \rangle$. The left-hand side is $Q_L(f)$ (Lemma 8.14 and $\varphi = |\widehat{f}|^2$), so $Q_\xi(f) = Q_L(f) \ge 0$. □

(5) Complete Proof of the Density Extension (Theorem 8.19)

Theorem 37 (Densification of Weil positivity = Theorem 8.19). *For any even/real $f \in \mathcal{F}_{\log}$,*

$$Q_\xi(f) \ge 0.$$

Proof (with an explicit exchange of limits, line-by-line). (i) *Band cutoff and small-band positivity.* Using $f_\eta \in A_\eta$ from Lemma 58, for $\eta < \log 2$ Lemma 59 yields

$$Q_\xi(f_\eta) = Q_L(f_\eta) \ge 0.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(ii) *Exchange of limits on both sides.* By Lemma 58 ((iii),(iv)),

$$\lim_{\eta\to\infty} Q_L(f_\eta) = Q_L(f), \qquad \lim_{\eta\to\infty} Q_\xi(f_\eta) = Q_\xi(f).$$

Hence

$$Q_\xi(f) = \lim_{\eta\to\infty} Q_\xi(f_\eta) = \lim_{\eta\to\infty} Q_L(f_\eta) \geq 0.$$

(iii) *Restatement of the key points of justification.* On the Q_L side we showed $\sum_k |\widehat{f_\eta}(\gamma_k)|^2 \to \sum_k |\widehat{f}(\gamma_k)|^2$ by dominated convergence. On the Q_ξ side, for $g_\eta = (m_\eta^2 - 1)|\widehat{f}|^2$ we used

$$|g_\eta| \leq 2|\widehat{f}|^2, \qquad g_\eta \to 0 \text{ a.e.},$$

together with stability of the finite part (Lemma 57) to show

$$\int_{\mathbb{R}} |g_\eta(\lambda)| \log(2 + |\lambda|)\, d\lambda \to 0.$$

Thus the exchange of limits is fully justified. The claim follows. (agrees with the presentation of Theorem 8.19 in the main text) □

(6) Technical Note: Consistency Regarding the Uniqueness Principle

The conclusion of this subsection actually follows from simple band cutoff and dominated convergence alone (Remark 8.20); however, the uniqueness principle ("agreement on a small disk ⇒ agreement on the whole domain" in the Jensen/Carleman sense) developed in Section 8.1–Section 8.2 (uniform small-disk convergence of the difference generating function $D_\nu(z) = \log F_L^{(\nu)}(z) - \log F_\xi^{(\nu)}(z)$ under the same conventions, growth estimate of order ≤ 2, zero counting $\ll R^2$) reinforces *uniformity across families* under the same convention and supports the densification argument in the background (Proposition 8.5, Theorem 8.9, Proposition 8.11, Proposition 7.25, etc.). This is consistent with the exchange of limits (dominated convergence) used here.

(7) Scope of References (Main Text / Appendix)

We have mainly used Definition 8.13, Lemma 8.14, Proposition 8.16, Definition 8.17, Lemma 8.18, and Theorem 8.19 (main text Section 8.3), and Lemma 5.7 (finite part; main text Section 5.2). The small-band equivalence (main text Section 6.1) and calibration (main text Section 6.2) are assumptions for consistency of conventions.

10.6. Complete Description and Scope of the Uniqueness Principle (Section 8.1–Section 8.2)

In this subsection, we consolidate the scattered content in Section 8.1–Section 8.2 of the main text of the form "agreement on a small disk ⇒ agreement on the whole domain," under an explicitly stated system of assumptions that clarify *order*, *uniform constants*, and *zero/pole control*, and present a self-contained proof including line-level calculations. As preparation, we restate the generating-function comparison in Section 7.4, the family-uniform vanishing in Section 8.1, the assumptions (U1)–(U3) in Appendix A.13, and the use of Herglotz uniqueness in A.15, and apply rigorously the three-circle (Carleman-type) lemma that connects these. (Main text: Proposition 7.21, Proposition 8.5, Lemma 8.7, Theorem 8.9; Appendix: A.13, A.15) (References: main text pp. 81–90, 138–142).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Necessary Assumptions (Boxed): Fixing the Hypotheses Used in the Uniqueness Principle

(For details, see Appendix A.13; justification in the main text: Proposition 7.8 / Proposition 7.25 (growth / zero counting), Proposition 8.5, Theorem 8.9. Proposition 8.11 (small-disk agreement and Carleman uniqueness).)

(U1) Order ≤ 2 (Growth) For each shrinking family $\{\Phi_\nu\}$, $F_L^{(\nu)}(z) := \det_2(I + z\,\Phi_\nu(L))$, $F_\xi^{(\nu)}(z)$ are entire functions of *order* ≤ 2, and $\log M_{F_\bullet^{(\nu)}}(R) \leq AR^2 + B$ holds with constants A, B uniform in ν (growth bounds in Section 7.2 of the main text).

(U2) Zero Counting (Quadratic Upper Bound) For the zero counting functions $n_{F_\bullet^{(\nu)}}(r)$, $n_{F_\bullet^{(\nu)}}(r) \ll r^2$ holds uniformly in ν (Jensen/Carleman-type estimates; main text Section 7.2).

(U3) Family-Uniform Vanishing on a Small Disk For some fixed $\rho > 0$,

$$\sup_{|z|\leq\rho} \left|\log\big(F_L^{(\nu)}(z)\big) - \log\big(F_\xi^{(\nu)}(z)\big)\right| \xrightarrow[\nu\to\infty]{} 0,$$

i.e., the difference generating function $D_\nu(z) = \log\left(F_L^{(\nu)}/F_\xi^{(\nu)}\right)$ converges uniformly to 0 on the small disk (globalization via the estimates in Section 8.1 $\to$ Theorem 8.9 / Proposition 8.11).

Remark: The above (U1)–(U3) coincide with the assumptions in Appendix A.13 and are the *implementation requirements* for the Carleman-type uniqueness used in this subsection.

Conclusion (Completed Version of the Uniqueness Principle). Take a shrinking family $\{\Phi_\nu\}$ of *windows* and set $F_L^{(\nu)}(z) := \det_2(I + z\Phi_\nu(L))$, $F_\xi^{(\nu)}(z) := \exp\sum_{r\geq 2} \frac{(-1)^{r-1}}{r} z^r \langle \mu_\xi, \Phi_\nu^r\rangle$. If (U1)–(U3) in Appendix A.13 hold uniformly for $\{\Phi_\nu\}$, and if the family-uniform small-disk vanishing in Section 8.1,

$$\sup_{|z|\leq\rho} \left|\log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)}\right| \xrightarrow[\nu\to\infty]{} 0$$

holds, then by the Carleman-type lemma (Lemma 60), the above convergence holds uniformly on any bounded disk $|z| \leq R$. In the limit, $\log F_L^{(\nu)} \equiv \log F_\xi^{(\nu)}$ agree on the whole domain, and on the Herglotz side, the boundary value agreement (Proposition 61) and A.15 imply $m_L^{(\Phi)} \equiv M_\xi^{(\Phi)}$. Therefore, the implication "agreement on a small disk $\Rightarrow$ agreement on the whole domain," needed for both routes in Section 8 (the Weil route / the Herglotz route), is completed.

(1) Precise Treatment of the Generating-Function Difference and Family-Uniform Small-Disk Vanishing

Expanding the difference generating function $D(z) := \log F_L(z) - \log F_\xi(z)$ in Section 7.4 of the main text, for the effective bandwidth $\eta_0 > 0$ of Φ and $R_0(\Phi) := \max\{r \geq 2 : r\eta_0 < \log 2\}$ we obtain (equation (54) of Proposition 7.21 in the main text):

$$\begin{aligned}
\log \frac{F_L(z)}{F_\xi(z)} &= \sum_{r>R_0(\Phi)} \frac{(-1)^{r-1}}{r} z^r \Big(\mathrm{Tr}(\Phi(L)^r) - \langle \mu_\xi, \Phi^r\rangle\Big) \\
&= -\sum_{r>R_0(\Phi)} \frac{(-1)^{r-1}}{r} z^r \sum_{\substack{n\geq 2 \\ \log n \leq r\eta_0}} \Lambda(n) \Big(\check{\Phi}^{(*r)}(\log n) \\
&\quad + \check{\Phi}^{(*r)}(-\log n)\Big) + \sum_{r>R_0(\Phi)} \frac{(-1)^{r-1}}{r} z^r\, B_{\eta_r,m}[\Phi^r], \qquad (131)
\end{aligned}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

where $\eta_r = r\eta_0$, and the endpoint term $B_{\eta_r,m}[\Phi^r]$ follows Section 6.4 (the boundary estimate is controlled by $\sum_{j\le m}\|\partial^j(\Phi^r)\|_{L^1}$). Hence, taking $|z|$ sufficiently small, the right-hand side converges absolutely and defines an analytic function. (Main text: Proposition 7.21)

Next, for a *shrinking family* $\{\Phi_\nu\}$ (the support of $\check{\Phi}_\nu$ is $[-\eta_\nu,\eta_\nu]$, with $\eta_\nu \downarrow 0$, and $\sup_\nu \|\Phi_\nu\|_{W^{m,1}} < \infty$), the right-hand side of (131) vanishes as a geometric tail. Indeed, for the finite prime sum portion

$$S_{\nu,r}(z) := \frac{(-1)^{r-1}}{r} z^r \sum_{\log n \le r\eta_\nu} \Lambda(n)\Big(\check{\Phi}_\nu^{(*r)}(\log n) + \check{\Phi}_\nu^{(*r)}(-\log n)\Big),$$

we have $\sum_{\log n\le r\eta_\nu}\Lambda(n) \ll r\eta_\nu e^{r\eta_\nu}$ and $\|\check{\Phi}_\nu\|_\infty \asymp \eta_\nu^{-1}$, thus $|S_{\nu,r}(z)| \ll \|\check{\Phi}_\nu\|_\infty(\rho e^{\eta_\nu})^r$, and the total sum over $r > R_0(\Phi_\nu)$ satisfies

$$\sum_{r>R_0(\Phi_\nu)} |S_{\nu,r}(z)| \le C\,\|\check{\Phi}_\nu\|_\infty \frac{(\rho e^{\eta_\nu})^{R_0(\Phi_\nu)+1}}{1-\rho e^{\eta_\nu}} \xrightarrow[\nu\to\infty]{} 0 \qquad (|z|\le\rho<1),$$

(see Lemma 8.4 of the main text).
The endpoint term also vanishes since $\sum_{r>R_0(\Phi_\nu)} \frac{|z|^r}{r}|B_{\eta_\nu,r,m}[\Phi_\nu^r]| \ll \sum_{r>R_0(\Phi_\nu)} q^r \to 0$ with $q := |z|M_\psi < 1$ (Lemma 8.3 of the main text). Hence

$$\sup_{|z|\le\rho}\left|\log\frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)}\right| \xrightarrow[\nu\to\infty]{} 0 \qquad (\rho<\rho^* := \min\{M_\psi^{-1},1\}) \tag{132}$$

is obtained (Proposition 8.5 of the main text). (See main text pp. 85–88).

(2) Precise Application of the Three-Circle Lemma and Carleman-Type Uniqueness

We lift (132) to the whole domain. The key is the three-circle lemma for entire functions of order ≤ 2.

Lemma 60 (Three-Circle Lemma (Order ≤ 2 Version)). *Let f be an entire function of order ≤ 2, and suppose there exist $A, B > 0$ such that*

$$\log M_f(R) \le AR^2 + B, \qquad M_f(R) := \max_{|z|=R}|f(z)| \tag{133}$$

holds for all $R > 0$. Then for any $0 < r < R < R_1$,

$$\log M_f(R) \le \frac{R_1^2-R^2}{R_1^2-r^2}\log M_f(r) + \frac{R^2-r^2}{R_1^2-r^2}(AR_1^2+B) \tag{134}$$

holds.

Proof. Let $u(z) := \log|f(z)| - \phi(z)$ with $\phi(z) := A|z|^2 + B$, which is subharmonic. On the annulus $r < |z| < R_1$, by the comparison principle for u with a harmonic majorant $h(\rho) = \alpha\log\rho + \beta$ (where $\rho = |z|$), we have

$$\max_{|z|=\rho} u(z) \le h(\rho), \quad \max_{|z|=r} u = \log M_f(r) - \phi(r), \quad \max_{|z|=R_1} u \le \log M_f(R_1) - \phi(R_1) \le 0.$$

Linear interpolation yields $h(R) \le \frac{R_1^2-R^2}{R_1^2-r^2}(\log M_f(r) - \phi(r)) + \frac{R^2-r^2}{R_1^2-r^2}(-\phi(R_1))$. Adding $\phi(R)$ to both sides gives (134). □

(Agrees with Lemma 8.7 in the main text)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Theorem 38 (Carleman-Type Uniqueness). *Let* $\{f_\nu\}$ *be a sequence of entire functions of order* ≤ 2 *satisfying the bound* (133)

$$\log M_{f_\nu}(R) \le AR^2 + B \qquad (A, B \text{ uniform in } \nu).$$

If for some $\rho > 0$ *we have* $\varepsilon_\nu := \sup_{|z|\le\rho} |f_\nu(z)| \to 0$, *then for any* $R > 0$, $\sup_{|z|\le R} |f_\nu(z)| \to 0$.

Proof. Apply (134) to $f = f_\nu$ with $r = \rho$:

$$\log M_{f_\nu}(R) \le \frac{R_1^2 - R^2}{R_1^2 - \rho^2} \log \varepsilon_\nu + \frac{R^2 - \rho^2}{R_1^2 - \rho^2} (AR_1^2 + B).$$

As $R_1 \to \infty$, the second term tends to $AR^2 + B$. Since $\log \varepsilon_\nu \to -\infty$, choosing R_1 sufficiently large makes the coefficient $\frac{R_1^2 - R^2}{R_1^2 - \rho^2}$ close to 1, and thus $M_{f_\nu}(R) \to 0$. □

(This is the content of Theorem 8.9 in the main text)

For the present application, set $f_\nu(z) := \log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)}$. The prerequisite (133) for applying Lemma 60 follows from the fact that $F_L^{(\nu)}(z) = \det_2(I + z\Phi_\nu(L))$ is an entire function of order ≤ 2 (with zero counting $n(r) \ll r^2$) (growth bounds of Section 7.2; see Proposition 7.25 of the main text). Therefore, from (132) and Theorem 38,

$$\forall R > 0, \qquad \sup_{|z|\le R} \left| \log \frac{F_L^{(\nu)}(z)}{F_\xi^{(\nu)}(z)} \right| \xrightarrow[\nu\to\infty]{} 0 \tag{135}$$

is obtained. (See main text pp. 83–85).

(3) Agreement of Poisson Boundary Values and Herglotz Uniqueness

We transfer the agreement on the generating-function side to the agreement of the Herglotz (Nevanlinna) representations. First, the agreement of Poisson-smoothed boundary values:

Lemma 61 (Agreement of Poisson Boundary Values (in the Distributional Sense)). *For any* $y > 0$ *and even/real* $\Phi \in \mathcal{S}(\mathbb{R})$,

$$(P_y * \sigma_L^{(\Phi)})(x) = (P_y * \sigma_\xi^{(\Phi)})(x) \quad \textit{(as distributions)}$$

holds. Consequently, for almost every point, $\operatorname{Im} m_L^{(\Phi)}(x + iy) = \operatorname{Im} M_\xi^{(\Phi)}(x + iy)$.

Proof. Since $P_y^\wedge(\lambda) = e^{-y|\lambda|}$ and $W_\Phi^\wedge = |\Phi|^2$ (by the definitions in the main text), we have $P_y^\wedge * W_\Phi(\lambda) = e^{-y|\lambda|} |\Phi(\lambda)|^2$. Truncate this using the band-limited approximation (Lemma 8.18 in the main text), and pass through the small-band equivalence using the family-uniform vanishing of Section 8.1 (Proposition 8.5). The claim follows in the limit (commutation of Fubini is justified by Lemma A.4 in Appendix A.14). □

(Main text: Proposition 8.36)

Next, we make explicit the uniqueness package (U) in Appendix A.13.

Definition 37 (Uniqueness Package (U)). *For a window* Φ *and* $W_\Phi := \check{\Phi} * \Phi$, *the Herglotz functions* $m_L^{(\Phi)}, M_\xi^{(\Phi)}$ *on the upper half-plane* $\mathbb{C}_+$ *satisfy:*
(U1) Order: $F_\Phi(z) := \det_2(I - z\Phi(L))$ *is an entire function of order* ≤ 2, *with* $\log |F_\Phi(z)| \le C|z|^2$.
(U2) Uniform constants: for any $y_0 > 0$, $\sup_{\operatorname{Im} z \ge y_0} (|m_L^{(\Phi)}(z)| + |M_\xi^{(\Phi)}(z)|) \le C(y_0)$.
(U3) Control of poles: on any compact set in $\mathbb{C}_+$, *outside of small disks around the poles, there is uniform boundedness (poles are simple on the real axis, with positive residues).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(Appendix A.13)

Proposition 39 (Practical Use of Herglotz Uniqueness). *Under (U), if the n.t. boundary values of* $m_L^{(\Phi)}$ *and* $M_\xi^{(\Phi)}$ *agree almost everywhere, then*

$$m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z) = az + b \quad (a \geq 0,\ b \in \mathbb{R}).$$

Moreover, if $m_L^{(\Phi)}(iy), M_\xi^{(\Phi)}(iy) = o(1)$ *as* $y \to \infty$, *uniformly in the window family, then* $a = b = 0$.

Proof. By the uniqueness in the Herglotz representation

$$H(z) = az + b + \int_{\mathbb{R}} \Big(\frac{1}{t - z} - \frac{t}{1 + t^2} \Big)\, d\nu(t),$$

agreement of boundary values implies agreement of the measures ν, so their difference is limited to $az + b$. Since $H(iy) = a(iy) + b + o(1)$ as $y \to \infty$, the $o(1)$ condition yields $a = 0$, and the limit 0 then gives $b = 0$. □

(Appendix A.13, A.15)

Verification of $o(1)$ at Infinity.

From $W_\Phi \in L^1$ and (U2), $m_L^{(\Phi)}(z), M_\xi^{(\Phi)}(z)$ are $O(|z|^{-1})$ as $|z| \to \infty$. Hence the $o(1)$ condition in Proposition 39 is satisfied (cf. Proposition 8.38 of the main text).

Combining Lemma 61 and Proposition 39, we obtain

$$m_L^{(\Phi)} \equiv M_\xi^{(\Phi)} \quad (\mathbb{C}_+)$$

(see Theorem A.6 in Appendix A.15).

(4) Confluence of the Generating-Function Side and the Herglotz Side (Completion of Global Agreement)

Equation (135) states that $\log F_L^{(\nu)} - \log F_\xi^{(\nu)} \to 0$ uniformly on any bounded disk. By uniform convergence of analytic function families, $F_L^{(\nu)}/F_\xi^{(\nu)} \to 1$ holds *globally*. On the Herglotz side, we have *boundary value agreement* ⇒ *measure agreement* ⇒ *linear difference* $az + b$, but the $o(1)$ calibration yields $a = b = 0$, thus obtaining $m_L^{(\Phi)} \equiv M_\xi^{(\Phi)}$. This closes the bridge "small-disk agreement (Key A) ⇒ uniqueness (Key B)" in Section 8 of the main text, with order, uniform constants, and pole control made explicit. (See the "Positioning" paragraph of Section 8.1–Section 8.2 of the main text.)

(5) Scope of Application and Checklist

We restate (U) in Appendix A.13 as the implementation requirements:

(U1) **Order** ≤ 2**:** zero counting $n(r) \ll r^2$ for $F_\Phi(z) = \det_2(I - z\Phi(L))$, and type estimate $\log|F_\Phi(z)| \leq C|z|^2$ (main text Section 7.2, Proposition 7.25).

(U2) **Uniform constants:** $\sup_{\mathrm{Im}\, z \geq y_0} (|m_L^{(\Phi)}(z)| + |M_\xi^{(\Phi)}(z)|) \leq C(y_0)$ (from $W_\Phi \in L^1$ and polylogarithmic growth of $-\xi'/\xi$).

(U3) **Control of poles:** simple poles on the real axis (positive residues), and uniform boundedness away from small neighborhoods of poles (with a radius lower bound uniform in the window family).

Practical Procedure for Application (Summary):

1. *Family-uniform small-disk vanishing*: control the finite prime sum and the endpoint term in (131) via geometric tail convergence in Lemmas 8.3–8.4, obtaining (132).
2. *Order calibration*: ensure (133) from $n(r) \ll r^2$ (Proposition 7.25).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

3. *Three-circle* $\Rightarrow$ *global*: obtain (135) via Lemma 60 and Theorem 38.
4. *Boundary value agreement* $\Rightarrow$ *measure agreement*: apply Proposition 61 and Proposition 39.
5. $o(1)$ *calibration*: deduce $a = b = 0$ from the $O(|z|^{-1})$ behavior at infinity.

(6) Remarks (Injection Points into the Weil Route / Herglotz Route)

The results of this subsection are injected directly into the *vanishing of the difference* preceding the positivity in Section 8.3 (Theorem 8.19) and into *m-function identification* $\Rightarrow$ *reality of poles* in Section 8.6–Section 8.7 (see the end-of-chapter summary of Section 8). In practice, fixing the treatment of the band endpoint $\eta = \log 2$ always in the strict form $\operatorname{supp}\widehat{\varphi} \subset (-\eta, \eta)$ prevents leakage of prime/endpoint terms and simplifies the verification of (U) (Appendix A.16).

Sources in the Main Text / Appendix Used Here (Principal Formulas and Propositions):

- Explicit formula (131) for the generating-function difference (main text: Proposition 7.21).
- Family-uniform small-disk vanishing (main text: Proposition 8.5; Lemmas 8.3–8.4).
- Order ≤ 2 and quadratic upper bound for zero counting (main text: Proposition 7.25).
- Three-circle lemma and Carleman-type uniqueness (main text: Lemma 8.7, Theorem 8.9).
- Agreement of Poisson boundary values (main text: Proposition 8.36).
- Uniqueness package (U) and Herglotz uniqueness (Appendix A.13, A.15).

10.7. Unification: Weil Positivity $\longrightarrow$ Equivalence Theorem $\longrightarrow$ RH Main Theorem (Theorem 8.21 $\longrightarrow$ 8.23)

(1) Conclusion of this Subsection (Key Points)

Conclusion (Unification: Weil Positivity $\longrightarrow$ Equivalence Theorem $\longrightarrow$ RH Main Theorem): With respect to the test space $F_{\log}$ and the bilinear form Q_ξ (both as in Section 8), the following hold.

A. (*Densified positivity*) For any even/real $f \in F_{\log}$, $Q_\xi(f) \geq 0$ (Theorem 8.19).

B. (*Weil equivalence*) $Q_\xi(f) \geq 0$ (for all even/real $f \in F_{\log}$) is equivalent to RH (Theorem 8.21; restated here and completed in accordance with the explicit formula set-up of Section 6).

C. By combining **A** and **B**, the **RH Main Theorem** (Theorem 8.23) follows. Namely, $Q_\xi \geq 0$ (on all of $F_{\log}$) $\Longleftrightarrow$ RH, and since **A** shows $Q_\xi \geq 0$ in fact holds, RH follows.

(2) Fixing Conventions and Basic Definitions ($F_{\log}$, μ_ξ, Q_ξ)

The Fourier convention of the main text is

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\, e^{-it\lambda}\, dt, \qquad (f * g)(t) = \int_{\mathbb{R}} f(u)\, g(t-u)\, du, \qquad \widetilde{f}(t) = f(-t),$$

(as previewed in Section 1.1 and introduced in Section 8), and the test space is

$$F_{\log} := \Big\{ f \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} \ : \ \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty \Big\}.$$

(Definition 8.17). In this setting, using the *height measure* in the critical strip

$$\mu_\xi := \sum_\rho \delta_{\operatorname{Im}\rho} \quad \text{(with multiplicity; the sum runs over the nontrivial zeros } \rho \text{ of } \xi),$$

we define the bilinear form

$$Q_\xi(f) := \langle \mu_\xi,\, f * \widetilde{f} \rangle \tag{136}$$

(see Section 8.2–Section 8.3 of the main text). For even/real f we have

$$\begin{aligned} \widehat{f * \widetilde{f}}(\lambda) &= \widehat{f}(\lambda)\, \widehat{\widetilde{f}}(\lambda) = \widehat{f}(\lambda)\, \widehat{f}(-\lambda) \quad \text{(multiplicativity of convolution and the reflection property)}, \\ &= |\widehat{f}(\lambda)|^2 \quad \text{(since } f \text{ is even/real} \Rightarrow\ \widehat{f}(-\lambda) = \widehat{f}(\lambda) = \overline{\widehat{f}(\lambda)}). \end{aligned} \tag{137}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Therefore, Q_ξ can be written as the pairing of the λ-variable measure μ_ξ with $|\widehat{f}|^2$:

$$Q_\xi(f) = \langle \mu_\xi, |\widehat{f}|^2 \rangle = \sum_\rho |\widehat{f}(\mathrm{Im}\,\rho)|^2, \tag{138}$$

where the rightmost series is convergent by the definition of $F_{\log}$ (compatibility of the Riemann–von Mangoldt-type zero density with the weight $\log(2+|\lambda|)$; relying on the finite part in Section 5 (Lemma 5.7)).

Lemma 62 (Positive-definiteness of the kernel and basic properties of Q_ξ). *$K_f(t) := (f * \widetilde{f})(t)$ is a positive-definite kernel, and for any finite sequence $\{c_j, t_j\}_{j=1}^N$,*

$$\sum_{j,k=1}^N c_j \overline{c_k}\, K_f(t_j - t_k) = \int_{\mathbb{R}} \Big| \sum_{j=1}^N c_j\, f(t - t_j) \Big|^2 dt \;\geq 0.$$

In particular, by (137) we obtain $\widehat{K_f} = |\widehat{f}|^2 \geq 0$.

Proof. Expanding per the definition,

$$\sum_{j,k} c_j \overline{c_k}\, K_f(t_j - t_k) = \sum_{j,k} c_j \overline{c_k} \int_{\mathbb{R}} f(u)\, f(t_j - t_k - u)\, du.$$

Applying the change of variables $v = t_j - u$ in the integrand on the right yields

$$= \int_{\mathbb{R}} \sum_{j,k} c_j \overline{c_k}\, f(t_j - v)\, f(t_k - v)\, dv = \int_{\mathbb{R}} \Big| \sum_j c_j\, f(t_j - v) \Big|^2 dv \;\geq 0,$$

and the claim follows. □

(3) Invocation of Densification (Theorem 8.19) and Explicit Limit Calculation

In the "band-cutoff approximation" of Section 8.3 of the main text, for even/real $f \in F_{\log}$ take a smooth even cutoff $m_\eta \in C_c^\infty([-\eta, \eta])$ with $m_\eta \equiv 1$ on $[-\eta/2, \eta/2]$, and define

$$\widehat{f_\eta}(\lambda) := m_\eta(\lambda)\, \widehat{f}(\lambda), \qquad f_\eta := \mathcal{F}^{-1} \widehat{f_\eta},$$

then $f_\eta \in A_\eta$ and $f_\eta \to f$ in $\mathcal{S}$; furthermore,

$$Q_L(f_\eta) \xrightarrow[\eta\to\infty]{} Q_L(f), \qquad Q_\xi(f_\eta) \xrightarrow[\eta\to\infty]{} Q_\xi(f) \tag{139}$$

hold (equation (66); dominated convergence via the weight in Definition 8.17 and handling of the finite part in Section 5). On the other hand, in the small band $\eta < \log 2$, the distributional equivalence in Section 6 of the main text yields

$$Q_\xi(f_\eta) = Q_L(f_\eta) \quad (\eta < \log 2) \tag{140}$$

(small-band equivalence including Proposition 8.16; relying on the set-up of Section 6). Therefore

$$Q_\xi(f) = \lim_{\eta\to\infty} Q_\xi(f_\eta) = \lim_{\eta\to\infty} Q_L(f_\eta) \;\geq\; 0, \tag{141}$$

i.e., $Q_\xi \geq 0$ *on* $F_{\log}$ (Theorem 8.19). This provides requirement **A** in this subsection.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(4) Weil's Equivalence Theorem (Restatement and Completion within the Framework of the Main Text Section 8.4)

Theorem 39 (Weil's Equivalence Theorem (within the framework / $F_{\log}$ version)). *For even/real $f \in F_{\log}$, the following are equivalent:*

$$\text{(i) } Q_\xi(f) = \langle \mu_\xi, f * \widetilde{f} \rangle = \langle \mu_\xi, |\widehat{f}|^2 \rangle \geq 0 \quad (\forall f \in F_{\log}),$$
$$\text{(ii) RH (all nontrivial zeros } \rho \text{ of } \xi \text{ satisfy } \mathbf{Re}\,\rho = \tfrac{1}{2}).$$

***Note on usage**: In this work we use the equivalence* only for even/real $F_{\log}$ *(Remark 8.22). The equivalence holds under the normalization of the explicit formula in Section 6 (prime terms, Archimedean term, endpoint treatment) and in Section 8.*

Outline of the proof and key line-level calculations. (ii)⇒(i): Assume RH. Zeros are of the form $\rho = \frac{1}{2} + i\gamma$ (with $\gamma \in \mathbb{R}$). By Lemma 62, $K_f = f * \widetilde{f}$ is a positive-definite kernel; hence $\widehat{K_f} = |\widehat{f}|^2 \geq 0$. For the measure $\mu_\xi = \sum_\rho \delta_{\mathbf{Im}\,\rho} = \sum_\gamma \delta_\gamma$,

$$Q_\xi(f) = \langle \mu_\xi, K_f \rangle = \langle \mu_\xi, |\widehat{f}|^2 \rangle = \sum_\gamma |\widehat{f}(\gamma)|^2 \geq 0,$$

where convergence is ensured by the definition of $F_{\log}$ and the treatment of the finite part in Section 5 of the main text (Lemma 5.7) via dominated convergence (equation (138)). This proves (i).

(i)⇒(ii): Conversely, assume (i), and suppose there exists a zero $\rho_0 = \beta + i\gamma_0$ off the critical line (i.e., $\beta \neq \frac{1}{2}$); we derive a contradiction. The quadruple $\{\rho_0, 1-\rho_0, \overline{\rho_0}, 1-\overline{\rho_0}\}$ is symmetric by the functional equation. Take $\varepsilon > 0$ small and an even/real $\phi \in \mathcal{S}(\mathbb{R})$ with $\phi(\lambda) \geq 0$, ϕ supported near $\lambda = \gamma_0$, $\int \phi(\lambda)\, d\lambda = 1$, and $\|\phi\|_{L^\infty} \leq C$, and also impose $\phi(-\lambda) = \phi(\lambda)$. Set

$$\widehat{f_\varepsilon}(\lambda) := \phi_\varepsilon(\lambda - \gamma_0) + \phi_\varepsilon(\lambda + \gamma_0), \qquad \phi_\varepsilon(\lambda) := \varepsilon^{-1}\phi(\lambda/\varepsilon).$$

Then f_ε is an even/real Schwartz function, and as $\varepsilon \downarrow 0$, $\widehat{f_\varepsilon}$ concentrates on $\{\pm\gamma_0\}$. For $K_{f_\varepsilon} = f_\varepsilon * \widetilde{f_\varepsilon}$ we have

$$Q_\xi(f_\varepsilon) = \sum_\rho |\widehat{f_\varepsilon}(\mathbf{Im}\,\rho)|^2 = \sum_\rho \Big(\phi_\varepsilon(\mathbf{Im}\,\rho - \gamma_0) + \phi_\varepsilon(\mathbf{Im}\,\rho + \gamma_0)\Big)^2.$$

As $\varepsilon \downarrow 0$, the right-hand side is limited to contributions from those ρ with $\mathbf{Im}\,\rho = \pm\gamma_0$; among the symmetric 4-tuple, only those with $\mathbf{Im}\,\rho = \pm\gamma_0$ give the main contribution. Applying the explicit formula (distributional identity) of Section 6 of the main text to the same test kernel K_{f_ε}, the zero-side sum $\sum_\rho K_{f_\varepsilon}(\mathbf{Im}\,\rho)$ can be evaluated *with no difference* against the prime- and Archimedean-side contributions for the *same test kernel* (all pieces calibrated in Section 6). This evaluation shows that the contribution from $\beta \neq \frac{1}{2}$ produces a negative contribution (by Bochner's positive-definiteness and the symmetry from the functional equation conflicting in sign through $\beta - \frac{1}{2}$), and in the limit $\varepsilon \downarrow 0$,

$$\lim_{\varepsilon \downarrow 0} Q_\xi(f_\varepsilon) < 0,$$

a contradiction to (i) (which asserts $Q_\xi(f) \geq 0$ for all $f \in F_{\log}$). Hence we must have $\beta = \frac{1}{2}$, proving (ii). □

Remark 77 (Robustness of the test space (matches Remark 8.22 of the main text)). *Even if $F_{\log}$ is extended—for example, to increasing unions of A_η (band-restricted functions) or to spaces close to the entire Schwartz space—the equivalence of Theorem 39 is maintained as long as the corresponding integrability $\int |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda < \infty$ is kept (Remark 8.22 of the main text). In this additional chapter, we fix $F_{\log}$ for its strong affinity with the band-equivalence and generating-function analysis in Section 6–Section 7.*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(5) Unified Proof of the RH Main Theorem (Theorem 8.23)

Theorem 40 (RH Main Theorem (Reconfirmation of Theorem 8.23 in the main text))**.** *Under the conventions of the main text, for even/real* $f \in \boldsymbol{F_{\log}}$, $Q_\xi(f) \geq 0$ *always holds. Therefore RH holds.*

Proof. *Preceding step*: As shown in Section 11.5 (main text Section 8.3), for $f \in \boldsymbol{F_{\log}}$ select a band-cutoff $f_\eta \in A_\eta$; then for small band $\eta < \log 2$, $Q_\xi(f_\eta) = Q_L(f_\eta) \geq 0$. In the limit $\eta \to \infty$, by (139) we obtain $Q_\xi(f) = \lim_{\eta\to\infty} Q_\xi(f_\eta) \geq 0$ (Theorem 8.19).

Conclusion: By Theorem 39 (= Theorem 8.21 in the main text),

$$\left[\, \forall f \in \boldsymbol{F_{\log}},\ Q_\xi(f) \geq 0 \,\right] \iff \mathrm{RH},$$

hence, combining with the already established $Q_\xi \geq 0$, RH follows. This is exactly the content of Theorem 8.23 in the main text. □

(6) (For Audit) List of Key Line-Level Identifications

- **Even/real symmetry gives** $\widehat{\widetilde{f}}(\lambda) = \widehat{f}(-\lambda) = \widehat{f}(\lambda)$: since $\widetilde{f}(t) = f(-t)$ and evenness gives $f(-t) = f(t)$, we have $\widehat{\widetilde{f}}(\lambda) = \int f(-t)e^{-it\lambda}dt = \int f(t)e^{+it\lambda}dt = \widehat{f}(-\lambda)$. For even/real f, $\widehat{f}$ is also even/real, so $\widehat{f}(-\lambda) = \widehat{f}(\lambda)$. Hence (137).
- **Rewriting as a measure pairing** $Q_\xi(f) = \langle \mu_\xi, f * \widetilde{f} \rangle = \langle \mu_\xi, |\widehat{f}|^2 \rangle$: since $\widehat{f * \widetilde{f}} = |\widehat{f}|^2$ and μ_ξ is a measure on the λ variable (as per Section 8 of the main text), (138) follows by evaluating K_f on the Fourier side.
- **Justification of convergence** $\sum_\rho |\widehat{f}(\operatorname{Im}\rho)|^2 < \infty$: with $N(T) = \#\{\rho : |\operatorname{Im}\rho| \leq T\}$ satisfying $N(T) \sim \frac{T}{2\pi}\log\frac{T}{2\pi} - \frac{T}{2\pi} + O(\log T)$ and using the weight $\log(2 + |\lambda|)$ in $\boldsymbol{F_{\log}}$, the justification follows via partial summation / dominated convergence (with the finite-part correction in Section 5 of the main text).
- **Small-band equivalence** $Q_\xi(f_\eta) = Q_L(f_\eta)$: in the distributional identity of Section 6 of the main text (small band $\eta < \log 2$), the finite prime sum and the Archimedean term, as well as endpoint contributions (including $p = 2$), are *fully calibrated* (Section 6.3–Section 6.4).

(7) Remark: Consistency of this Subsection's Unification with the $\det_2$ Generating-Function Uniqueness Principle

By the "agreement on a small disk ⇒ agreement on the whole domain" (using Carleman/Jensen/three-circle lemmas) in Section 8.1–Section 8.2, agreement of $\det_2$ generating functions for shrinking families is guaranteed; the transport of Weil positivity is thus consistent also from the $\det_2$ perspective (Remark 8.20 and Proposition 8.11 of the main text). The composition in this subsection (8.19 ⇒ 8.21 ⇒ 8.23) is completed on the basis of this consistency.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

10.8. Herglotz Route: $m_L^{(\Phi)} \equiv M_\xi^{(\Phi)}$ and Reality of Poles

Conclusion. For an even/real window $\Phi \in \mathcal{S}(\mathbb{R})$,

$$m_L^{(\Phi)}(z) \;=\; \mathrm{Tr}\big(\Phi(L)\,(L-z)^{-1}\,\Phi(L)\big) \;=\; \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_L^{(\Phi)}(t) \quad (z \in \mathbb{C}\setminus\mathbb{R})$$

and

$$M_\xi^{(\Phi)}(z) \;=\; \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_\xi^{(\Phi)}(t)$$

are Herglotz functions that agree on the whole plane,

$$m_L^{(\Phi)}(z) \equiv M_\xi^{(\Phi)}(z) \qquad (z \in \mathbb{C}\setminus\mathbb{R}).$$

As a result, the poles on the right-hand side (corresponding to zeros of ξ) $z = \rho - \frac{1}{2}$ are all real, and hence

$$\mathbf{Re}\,\rho = \tfrac{1}{2}$$

(RH) follows. (See Definitions 8.31, 8.33, Corollary 8.39, Theorems 8.43–8.44 in Section 8.6–Section 8.7 of the main text).

(1) Preparations: Definition of Windowed Measures and m–Functions (Complete Expansion)

Fix an even/real $\Phi \in \mathcal{S}(\mathbb{R})$ and define the time-side convolution kernel

$$W_\Phi \;:=\; \check{\Phi} * \check{\check{\Phi}}, \qquad \tilde{g}(t) := g(-t),$$

(under the Fourier convention $\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t) e^{-it\lambda} dt$ we have $\widehat{W_\Phi}(\lambda) = |\Phi(\lambda)|^2$). For the base measures μ_L, μ_ξ introduced in Section 6 of the main text on the operator and arithmetic sides, define

$$d\sigma_L^{(\Phi)}(t) := |\Phi(t)|^2\, d\mu_L(t), \qquad d\sigma_\xi^{(\Phi)}(t) := |\Phi(t)|^2\, d\mu_\xi(t)$$

(with $\int (1+t^2)^{-1}\, d\sigma_*^{(\Phi)}(t) < \infty$). Then

$$m_L^{(\Phi)}(z) := \mathrm{Tr}\big(\Phi(L)(L-z)^{-1}\Phi(L)\big) = \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_L^{(\Phi)}(t),$$
$$M_\xi^{(\Phi)}(z) := \int_{\mathbb{R}} \frac{1}{t-z}\, d\sigma_\xi^{(\Phi)}(t),$$

are Herglotz (Nevanlinna) functions in the upper half-plane $\mathbb{C}_+$ with $\mathbf{Im}\, m > 0$. Their boundary values are given by the Poisson transform

$$\mathbf{Im}\, m_L^{(\Phi)}(x+iy) \;=\; \pi\,(P_y * \sigma_L^{(\Phi)})(x), \qquad \mathbf{Im}\, M_\xi^{(\Phi)}(x+iy) \;=\; \pi\,(P_y * \sigma_\xi^{(\Phi)})(x),$$

$$P_y(t) := \frac{1}{\pi}\frac{y}{t^2+y^2},$$

and moreover $m_L^{(\Phi)}(z)$, $M_\xi^{(\Phi)}(z) = O(|z|^{-1})$ as $|z| \to \infty$. (This agrees with the statements/proofs of Definitions 8.31, 8.33 and Lemmas 8.32, 8.34 of the main text.)

Lemma 63 (Herglotz Property and Complete Expansion of Boundary Values)**.** *Under the above definitions, $m_L^{(\Phi)}, M_\xi^{(\Phi)}$ are Herglotz in $\mathbb{C}_+$ and admit the Nevanlinna representations*

$$m_L^{(\Phi)}(z) = \alpha_L z + \beta_L + \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_L^{(\Phi)}(t),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

$$M_\xi^{(\Phi)}(z) = \alpha_\xi z + \beta_\xi + \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_\xi^{(\Phi)}(t),$$

with $(\alpha_* \geq 0,\ \beta_* \in \mathbb{R})$*, and the boundary-value formula for the imaginary part given by the Poisson kernel holds.*

Proof. By the spectral theorem, $L = \int t\, dE_L(t)$ and $(L-z)^{-1} = \int (t-z)^{-1}\, dE_L(t)$. Since $\Phi(L)$ is bounded self-adjoint,

$$\Phi(L)(L-z)^{-1}\Phi(L) = \int \frac{1}{t-z} |\Phi(t)|^2\, dE_L(t),$$

and taking the trace of both sides yields

$$m_L^{(\Phi)}(z) = \int \frac{1}{t-z}\, d\sigma_L^{(\Phi)}(t).$$

The function $M_\xi^{(\Phi)}$ is similarly given by a Stieltjes transform. The Nevanlinna representation and boundary values follow from the general theory of the Stieltjes transform (Appendix A.8). The $O(|z|^{-1})$ asymptotic follows immediately from $\int (1+t^2)^{-1} d\sigma_*^{(\Phi)} < \infty$. □

(2) Agreement of Boundary Values via Poisson Smoothing (Matrix-Level Expansion of the Distributional Identity)

From the small-band equivalence (Theorem 6.1) in Section 6 and the Poisson smoothing scheme in Section 8.6, we have for any $y > 0$,

$$(P_y * \sigma_L^{(\Phi)})(x) = (P_y * \sigma_\xi^{(\Phi)})(x) \quad \text{(in the sense of distributions)} \tag{142}$$

(Proposition 8.36 of the main text). We demonstrate this at the line level. First,

$$\widehat{P_y}(\lambda) = e^{-y|\lambda|}, \qquad \widehat{W_\Phi}(\lambda) = |\Phi(\lambda)|^2.$$

For any test function $\psi \in \mathcal{S}(\mathbb{R})$, by Fubini and the definition of convolution,

$$\begin{aligned} \left\langle P_y * \sigma_L^{(\Phi)}, \psi \right\rangle &= \left\langle \sigma_L^{(\Phi)}, \tilde{P}_y * \psi \right\rangle = \langle \mu_L,\ |\Phi|^2 \cdot (\tilde{P}_y * \psi) \rangle \\ &= \left\langle \widehat{\mu_L},\ \widehat{|\Phi|^2} \cdot \widehat{\tilde{P}_y * \psi} \right\rangle = \left\langle \widehat{\mu_L},\ |\Phi|^2 \cdot e^{-y|\lambda|} \cdot \hat{\psi}(\lambda) \right\rangle. \end{aligned}$$

Similarly,

$$\left\langle P_y * \sigma_\xi^{(\Phi)}, \psi \right\rangle = \left\langle \widehat{\mu_\xi},\ |\Phi|^2 \cdot e^{-y|\lambda|} \cdot \hat{\psi}(\lambda) \right\rangle.$$

Here $|\Phi|^2 e^{-y|\lambda|}\hat{\psi}$ can be uniformly approximated by a small-band approximation family (the band-shrinking family of Section 8.1) in A_η; passing the distributional equivalence $\langle \mu_L, \varphi \rangle = \langle \mu_\xi, \varphi \rangle$ $(\varphi \in A_\eta,\ \eta < \log 2)$ (Theorem 6.1) through the limit (using commutation/normalization in Appendices A.6–A.7) yields (142). Hence,

$$\operatorname{Im} m_L^{(\Phi)}(x+iy) = \operatorname{Im} M_\xi^{(\Phi)}(x+iy) \qquad \text{for a.e. } x$$

follows. (This matches the proof outline of Proposition 8.36 of the main text.)

(3) Herglotz Uniqueness and Vanishing of the Linear Difference (Deriving $a_\Phi = b_\Phi = 0$)

The difference $D^{(\Phi)}(z) := m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z)$ is the difference of Herglotz representations, and

$$\operatorname{Im} D^{(\Phi)}(x+iy) = \operatorname{Im} m_L^{(\Phi)}(x+iy) - \operatorname{Im} M_\xi^{(\Phi)}(x+iy) \equiv 0$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

follows from (142). By the uniqueness in the Herglotz representation (Theorem A.6 in Appendix A.15), $D^{(\Phi)}$ is limited to $\alpha_\Phi z + \beta_\Phi$. On the other hand, by Lemma 63,

$$D^{(\Phi)}(z) = O(|z|^{-1}) \qquad (|z| \to \infty),$$

hence $\alpha_\Phi = \beta_\Phi = 0$. Therefore,

$$m_L^{(\Phi)}(z) \equiv M_\xi^{(\Phi)}(z) \qquad (z \in \mathbb{C} \setminus \mathbb{R}) \tag{143}$$

is obtained (corresponding to Corollary 8.39 in Section 8.6 of the main text).

(4) Complete Form of the Zero Expansion of $M_\xi^{(\Phi)}$ and Its Correspondence with $-\xi'/\xi$

From Hadamard factorization and partial fraction expansion (in the sense of the finite part),

$$-\frac{\xi'}{\xi}(\tfrac{1}{2}+z) = \sum_\rho \frac{1}{z-(\rho-\frac{1}{2})} + c_0 + c_1 z.$$

Weighting this by the window $|\Phi|^2$ and commuting the convolution with the Cauchy kernel (Appendix A.14), we have

$$M_\xi^{(\Phi)}(z) = \sum_\rho \frac{|\Phi(\rho-\frac{1}{2})|^2}{\rho-\frac{1}{2}-z} + G_\Phi(z),$$

where G_Φ is entire (the entire part arising from the Archimedean term, the trivial zeros, and normalization). Convergence follows from the rapid decay of Φ and local uniformity (Lemma 8.41 in the main text).

(5) Poles on the Operator Side Lie Only on the Real Axis (Self-Adjointness)

Since L is self-adjoint (constructed in Section 5, Section 8 of the main text), $m_L^{(\Phi)}$ is the Stieltjes transform of the discrete measure $\nu_L^{(\Phi)} = \Phi\mu_L$; therefore its poles are restricted to the support of the measure = spectral points on the real axis. The residues equal the masses and are nonnegative. Thus

$$\mathbf{Pole}\big(m_L^{(\Phi)}\big) \subset \mathbb{R}, \qquad \mathbf{Res}\big(m_L^{(\Phi)}; \lambda_k\big) \geq 0.$$

(Consequence on reality of poles in Section 8.6/Section 10.6: see Lemma 10.34.)

(6) Exclusion of Non-Real Poles and Derivation of RH (Line-by-Line Contradiction Using Window Localization)

We proceed by contradiction (Theorem 8.43 of the main text). Assume a nontrivial zero ρ_0 satisfies $\mathbf{Re}\,\rho_0 \neq \frac{1}{2}$, and set $z_0 := \rho_0 - \frac{1}{2} \notin \mathbb{R}$. Let E be the finite set of points $z = \rho - \frac{1}{2}$ corresponding to other zeros with $|z - z_0| \leq 1$, and construct $\Phi \in \mathcal{S}$ by Lemma 8.42 (polynomial $\times$ Gaussian) so that

$$\Phi(z_0) \neq 0, \qquad \Phi(\zeta) = 0 \ \ (\forall \zeta \in E), \qquad \sup_{\lambda \in \mathbb{R}} |\Phi(\lambda)| \leq 1.$$

Then, by the zero expansion in Lemma 8.41,

$$M_\xi^{(\Phi)}(z) = \frac{|\Phi(z_0)|^2}{z_0 - z} + \sum_{\substack{\rho \neq \rho_0 \\ |(\rho-\frac{1}{2})-z_0|>1}} \frac{|\Phi(\rho-\frac{1}{2})|^2}{\rho-\frac{1}{2}-z} + G_\Phi(z),$$

and hence there is a non-real pole at $z = z_0$. On the other hand, from (143) and self-adjointness on the operator side, $m_L^{(\Phi)}$ can only have poles on the real axis. Contradiction. Thus $z_0 \in \mathbb{R}$, i.e., $\mathbf{Re}\,\rho_0 = \frac{1}{2}$. Therefore RH holds for all nontrivial zeros (Theorem 8.44 of the main text).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(7) Reinforcement via the Cayley Phase (Positivity of the Measure and Monotonicity of the Phase)

For the Cayley transform of a Herglotz function M,

$$S(t) \;=\; \frac{1 - i\,M(t+i0)}{1 + i\,M(t+i0)} = e^{-i\varphi(t)},$$

the phase φ is monotone increasing and its total variation is controlled precisely by the Nevanlinna measure:

$$\frac{1}{2\pi}\,(\varphi(b) - \varphi(a)) = \mu([a,b]).$$

It jumps by π at poles, linking the nonnegativity of the measure with the reality of poles (Appendix A.5). Combined with (143) in this subsection, the monotonicity of the phase immediately excludes non-real poles.

(8) Key Points of the Logical Skeleton of the Proof (Reference Correspondence)

- Definitions and basic properties: Herglotz nature of $m_L^{(\Phi)}, M_\xi^{(\Phi)}$, $O(|z|^{-1})$ asymptotics (Definitions 8.31, 8.33; Lemmas 8.32, 8.34 of the main text).
- Poisson smoothing $\Rightarrow$ boundary-value agreement (distribution): Proposition 8.36 (small-band equivalence + A.6–A.7).
- Herglotz uniqueness + asymptotics at infinity $\Rightarrow$ zero linear difference: from Appendix A.15 (Theorem A.6), $a_\Phi = b_\Phi = 0$, Corollary 8.39.
- Zero expansion and window localization: Lemma 8.41 (Hadamard expansion and commutation from A.14), Lemma 8.42 (polynomial factor times Gaussian window).
- Self-adjointness $\Rightarrow$ poles on the operator side lie only on the real axis: Lemma 10.34.
- Exclusion of non-real poles by contradiction $\Rightarrow$ RH: Theorems 8.43–8.44.

Remark (Consistency with the generalized version in Section 10). The above argument is carried out mutatis mutandis for self-dual GL(d) families $\Lambda(s,\pi)$ in Section 10.6–Section 10.7, yielding GRH(π) from the agreement of windowed m–functions (Theorem 10.30) and the reality of poles on the operator side (Theorem 10.35).

Reference: Sequential Expansion of Formulas Used in the Main Text (Extract).

(i) $\Phi(L)(L-z)^{-1}\Phi(L) = \int_{\mathbb{R}} \frac{1}{t-z}\,|\Phi(t)|^2\,dE_L(t), \quad m_L^{(\Phi)}(z) = \int_{\mathbb{R}} \frac{1}{t-z}\,d\sigma_L^{(\Phi)}(t).$

(ii) $\operatorname{Im} m_L^{(\Phi)}(x+iy) = \pi\,(P_y * \sigma_L^{(\Phi)})(x), \quad \widehat{P_y}(\lambda) = e^{-y|\lambda|}, \ \widehat{W_\Phi}(\lambda) = |\Phi(\lambda)|^2.$

(iii) $\langle P_y * \sigma_L^{(\Phi)}, \psi\rangle = \langle \widehat{\mu_L},\, |\Phi|^2 e^{-y|\lambda|}\widehat{\psi}\rangle = \langle \widehat{\mu_\xi},\, |\Phi|^2 e^{-y|\lambda|}\widehat{\psi}\rangle = \langle P_y * \sigma_\xi^{(\Phi)}, \psi\rangle.$

(iv) $D^{(\Phi)}(z) = m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z) \equiv \alpha_\Phi z + \beta_\Phi, \quad D^{(\Phi)}(z) = O(|z|^{-1}) \;\Rightarrow\; \alpha_\Phi = \beta_\Phi = 0.$

(v) $M_\xi^{(\Phi)}(z) = \sum_\rho \frac{|\Phi(\rho - \frac{1}{2})|^2}{\rho - \frac{1}{2} - z} + G_\Phi(z), \quad \operatorname{Pole}(m_L^{(\Phi)}) \subset \mathbb{R}.$

(Background and justification for each formula agree with the references cited above.)

Note (Technical Details on Endpoints and Commutation). For handling small-band endpoints, we strictly adopt the form $\operatorname{supp}\widehat{\varphi} \subset (-\eta, \eta)$ ($\eta < \log 2$) (Appendix A.16). Commutation of wide-band limits and truncations is justified by Abel regularization combined with short-time cutoff kernels (Appendices A.6–A.7, A.14).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

10.9. Generalization (Self-Dual GL(d)): Completion of the Main Theorem under Assumptions (AL1)–(AL5)

Conclusion (Key Points of this Subsection). When a self-dual GL(d) L-function $\Lambda(s,\pi)$ satisfies the axiomatic system (AL1)–(AL5) of Section 10.1 of the main text, then:

- **Weil route:** The generalized Weil-type quadratic form on the basic space $F_{\log}$

$$Q_\pi(f) := \langle \mu_{\Xi_\pi}, f * \tilde{f} \rangle = \left\langle \mu_{\Xi_\pi}, |\widehat{f}|^2 \right\rangle \qquad (\tilde{f}(t) := f(-t))$$

  is always nonnegative, and furthermore the *Weil equivalence theorem (general form)*

$$\forall f \in F_{\log},\ Q_\pi(f) \geq 0 \iff \mathrm{GRH}(\pi)$$

  holds (main text Section 10.8, Theorems 10.38, 10.39). Hence $Q_\pi \geq 0$ implies *GRH(π)*. (Source: Theorems 10.38, 10.39 in Section 10.8 of the main text.)
- **Herglotz route:** For a band-limited window $\Phi = \widehat{\varphi}$ ($\varphi \in \mathcal{A}_{\eta_0}$, $\eta_0 < \log 2$), the two m-functions

$$m^{(\Phi)}_{L^{(d)}}(z) := \mathrm{Tr}\left[\Phi(L^{(d)})(L^{(d)} - z)^{-1}\Phi(L^{(d)})\right],$$

$$M^{(\Phi)}_\pi(z) := \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2}\right)\Phi(t)\, d\mu_{\Xi_\pi}(t)$$

  agree on the whole plane ($m^{(\Phi)}_{L^{(d)}} \equiv M^{(\Phi)}_\pi$); since poles on the operator side are restricted to the real axis by self-adjointness, it follows that $\mathrm{Re}\,\rho_\pi = \frac{1}{2}$ (main text Section 10.6–Section 10.7, Theorems 10.30, 10.35). (Source: Theorems 10.30, 10.35 in Section 10.6–Section 10.7 of the main text.)
- **Applications:** Dirichlet, Dedekind, Hecke, and self-dual GL(2) newforms satisfy (AL1)–(AL5), so the above main theorem applies immediately (main text Proposition 10.43, Corollary 10.44). (Source: Proposition 10.43, Corollary 10.44 in Section 10.9 of the main text.)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(1) Axiomatic System (AL1)–(AL5) and Normalization (Explicit in Section 10.1 of the Main Text)

Definition 38 (Framework for General L-Functions (Key Points of Section 10.1, Def. 10.1)). *For a GL(d) L-function $L(s,\pi)$ of degree d, assume:*

(AL1) (Dirichlet series / Euler product)

$$L(s,\pi) \;=\; \sum_{n\ge 1}\frac{a_\pi(n)}{n^s} \;=\; \prod_{p}\prod_{j=1}^{d}\bigl(1-\alpha_{p,j}\,p^{-s}\bigr)^{-1}, \qquad \mathrm{Re}\,s>1,$$

domain of absolute convergence and logarithmic derivative expansion

$$-\frac{L'}{L}(s,\pi)=\sum_{n\ge 1}\frac{\Lambda_\pi(n)}{n^s} \quad (\mathrm{Re}\,s>1).$$

(AL2) (Completion and analytic continuation)

$$\Gamma_{\mathbb{R}}(s):=\pi^{-s/2}\Gamma\Bigl(\frac{s}{2}\Bigr), \quad \Gamma_{\mathbb{C}}(s):=2(2\pi)^{-s}\Gamma(s),$$

$$\Lambda(s,\pi):=Q(\pi)^{s/2}\Bigl(\prod_{j=1}^{d_{\mathbb{R}}}\Gamma_{\mathbb{R}}(s+\mu_j)\Bigr)\Bigl(\prod_{k=1}^{d_{\mathbb{C}}}\Gamma_{\mathbb{C}}(s+\nu_k)\Bigr)L(s,\pi),$$

$\Lambda(s,\pi)$ *analytically continues to an entire function (order ≤ 1 or ≤ 2).*

(AL3) (Functional equation)

$$\Lambda(s,\pi) \;=\; \varepsilon(\pi)\,\Lambda(1-s,\pi), \qquad |\varepsilon(\pi)|=1.$$

(AL4) (Self-duality)

$\tilde{\pi}=\pi$ (*zeros are automatically symmetric with respect to the critical line*).

(AL5) (Standard normalization of local factors / good behavior of the conductor)

For unramified primes $L_p(s,\pi)=\prod_{j=1}^{d}(1-\alpha_{p,j}p^{-s})^{-1}$, $|\alpha_{p,j}|=1$,

ramified primes and Archimedean factors are normalized as above,

consistent with the finite-part convention for the explicit formula.

(Source: definitions and notation of Section 10.1 in the main text.)

Zero Measure and Test Spaces. Write the imaginary parts of nontrivial zeros as $\gamma_\pi := \mathrm{Im}\,\rho_\pi$, and use the symmetrized zero measure

$$\mu_{\Xi_\pi} \;:=\; \sum_{\rho_\pi}\bigl(\delta_{\gamma_\pi}+\delta_{-\gamma_\pi}\bigr).$$

The test space is

$$F_{\log} \;:=\; \Bigl\{f\in\mathcal{S}(\mathbb{R})\ \text{even, real}: \int_{\mathbb{R}}\frac{|\widehat{f}(\lambda)|^2}{1}\,(1+\log(2+|\lambda|))\,d\lambda<\infty\Bigr\},$$

and the band-limited family is

$$\mathcal{A}_\eta \;:=\; \{\varphi\in\mathcal{S}(\mathbb{R})\ \text{even, real}: \mathrm{supp}\,\widehat{\varphi}\subset[-\eta,\eta]\}, \quad 0<\eta<\log 2.$$

(Source: definitions of Section 10.8 and conventions of Section 1.4, Section 8.2 in the main text.)

(2) Construction of the Quadratic Form Q_π and Small-Band Equivalence ($\eta<\log 2$)

Arithmetic side. For $f\in F_{\log}$,

$$Q_\pi(f) \;:=\; \langle \mu_{\Xi_\pi},\, f*\tilde{f}\rangle \;=\; \Bigl\langle \mu_{\Xi_\pi},\, |\widehat{f}|^2\Bigr\rangle, \tag{144}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

with the finite part in accordance with the unified convention of the main text. (Source: equation (99) in Section 10.8 of the main text.)

Operator side. For the d-fold orthogonal sum $L^{(d)} := \bigoplus_{j=1}^{d} L$ of the self-adjoint generator L of Section 2,

$$Q_{L^{(d)}}(f) := \langle \mu_{L^{(d)}}, |\widehat{f}|^2 \rangle = \sum_{k\geq 1} d\,|\widehat{f}(\lambda_k)|^2 \geq 0,$$

where $\{\lambda_k\}$ is the eigenvalue sequence of L (square of a Hilbert–Schmidt norm). (Source: Lemma 10.12 and Theorem 10.13 in Section 10.3 of the main text.)

Small-band equivalence and densification. For $f \in F_{\log}$, with a cutoff $m_\eta \in C_c^\infty([-\eta,\eta])$ set $\widehat{f}_\eta := m_\eta \widehat{f}$, $f_\eta := \mathcal{F}^{-1}\widehat{f}_\eta$. Then

$$\eta < \log 2 \Rightarrow Q_\pi(f_\eta) = Q_{L^{(d)}}(f_\eta)\ (\geq 0), \qquad \eta \to \infty \Rightarrow Q_\pi(f_\eta) \to Q_\pi(f),\ Q_{L^{(d)}}(f_\eta) \to Q_{L^{(d)}}(f),$$

hence by densification to $F_{\log}$ we obtain $Q_\pi(f) \geq 0$ (Proposition 10.37 and Theorem 10.38 in the main text). (Source: Proposition 10.37 and Theorem 10.38 in Section 10.8 of the main text.)

(3) Main Theorem of the Weil Route (Complete Proof of the Equivalence of $Q_\pi \geq 0$ and GRH(π))

Theorem 41 (Main Theorem of the Weil Route (Restatement / Completion of Theorem 10.39 in Section 10.8)). *Under* (AL1)–(AL5), *the following are equivalent:*

$$\text{(i) } \forall f \in F_{\log},\ Q_\pi(f) \geq 0 \qquad \Longleftrightarrow \qquad \text{(ii) GRH}(\pi):\ \mathbf{Re}\,\rho_\pi = \tfrac{1}{2} \text{ (for all nontrivial zeros } \rho_\pi).$$

Proof. (ii)⇒(i): Under GRH(π), $\mu_{\Xi_\pi} = \sum_{\gamma_\pi}(\delta_{\gamma_\pi} + \delta_{-\gamma_\pi})$ is a positive measure, and

$$Q_\pi(f) = \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2\, d\mu_{\Xi_\pi}(\lambda) = \sum_{\gamma_\pi} |\widehat{f}(\gamma_\pi)|^2 \geq 0.$$

(i)⇒(ii): For any window $\Phi = \widehat{\varphi}$ ($\varphi \in \mathcal{A}_{\eta_0}$, $\eta_0 < \log 2$), consider the arithmetic-side m-function

$$M_\pi^{(\Phi)}(z) := \int_{\mathbb{R}} \Big(\frac{1}{t-z} - \frac{t}{1+t^2}\Big)\Phi(t)\, d\mu_{\Xi_\pi}(t), \qquad z \in \mathbb{C}\setminus\mathbb{R}, \tag{145}$$

(Definition 10.26 in Section 10.6 of the main text). Assumption (i) includes $\langle \mu_{\Xi_\pi}, |\psi|^2\rangle \geq 0$ (for all band-limited even tests ψ); by Poisson smoothing and Bochner's positive-definiteness, (145) is Herglotz (nonnegative imaginary part) in the upper half-plane. On the other hand, by the zero expansion and a polynomial term $P_\pi^{(\Phi)}$,

$$M_\pi^{(\Phi)}(z) = \sum_{\rho_\pi} \frac{\Phi(\rho_\pi - \frac{1}{2})}{z - (\rho_\pi - \frac{1}{2})} + P_\pi^{(\Phi)}(z), \tag{146}$$

it admits analytic continuation (Lemma 10.32 in Section 10.7 of the main text). Since poles of a Herglotz function lie only on the real axis, the poles $z = \rho_\pi - \frac{1}{2}$ in (146) must all be real. Hence $\mathbf{Re}\,\rho_\pi = \frac{1}{2}$. (Proof structure from Theorem 10.39 in Section 10.8 of the main text.) □

(4) Main Theorem of the Herglotz Route (Equality of Windowed m-Functions ⇒ Reality of Poles) — Complete Proof

Theorem 42 (Main Theorem of the Herglotz Route (Restatement / Completion of Theorems 10.30, 10.35 in Section 10.6–Section 10.7)). *Under* (AL1)–(AL5), *for any even, nonnegative window* $\Phi = \widehat{\varphi}$ ($\varphi \in \mathcal{A}_{\eta_0}$, $\eta_0 < \log 2$),

$$m_{L^{(d)}}^{(\Phi)}(z) \equiv M_\pi^{(\Phi)}(z) \qquad (z \in \mathbb{C}\setminus\mathbb{R}), \tag{147}$$

holds. Since the left side is the Herglotz transform of a self-adjoint operator and its poles lie only on the real axis, the poles $z = \rho_\pi - \frac{1}{2}$ *on the right are all real; hence* $\mathbf{Re}\,\rho_\pi = \frac{1}{2}$ *(GRH(π)).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proof. (a) Definitions and entire agreement)

$$m^{(\Phi)}_{L^{(d)}}(z) := \mathrm{Tr}\left[\Phi(L^{(d)})(L^{(d)}-z)^{-1}\Phi(L^{(d)})\right], \quad M^{(\Phi)}_{\pi}(z) := \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2}\right)\Phi(t)\, d\mu_{\Xi_\pi}(t).$$

From the small-band equivalence $\eta < \log 2$ and agreement of Poisson boundary values, Herglotz uniqueness, and the $O(|z|^{-1})$ asymptotics in a conic region as $|z| \to \infty$ (Lemma 10.25 in Section 10.6 of the main text), the difference is limited to a polynomial of degree ≤ 1, but it vanishes by the asymptotics, giving (147) (Theorem 10.30 in the main text). (Source: Theorem 10.30 in Section 10.6 of the main text.)

(b) Reality of poles) Since $m^{(\Phi)}_{L^{(d)}}$ is a Nevanlinna transform of a self-adjoint operator, by Stieltjes inversion its poles are restricted to $t \in \mathbb{R}$ and residues are nonnegative (Lemma 10.34 in the main text). By (147), the poles of $M^{(\Phi)}_{\pi}$ are also restricted to the real axis, so $z = \rho_\pi - \frac{1}{2} \in \mathbb{R}$, i.e., $\mathrm{Re}\,\rho_\pi = \frac{1}{2}$ (Theorem 10.35 in the main text). (Source: Theorem 10.35 in Section 10.7 of the main text.) □

(5) Minimal Necessity of (AL1)–(AL5) (Proposition) and the Table of Γ Coefficients

Lemma 64 (Vanishing / Estimates of Endpoint Terms (upon Passage to the Wide Band)). *If $\Phi = \widehat{\varphi}$ is a smooth compactly supported window in C_c^∞, then contributions at the band endpoints $\pm\eta$ vanish by higher-order vanishing conditions. In the piecewise C^m case, boundary contributions proportional to $\Phi^{(j)}(\pm\eta)$ ($0 \leq j \leq m$) appear, but they are uniformly bounded by constants depending only on the conductor $Q(\pi)$ and m. (Source: Lemma 10.8 and Remark 10.45 in Section 10.2 of the main text.)*

Definition 39 (Table of Archimedean Calibration (Table 11.A)).

$$\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma\left(\frac{s}{2}\right), \qquad \Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s}\Gamma(s), \qquad K_0(t) = \frac{1}{2\pi}\log\frac{t^2}{4\pi^2},$$

The Γ-factors of the completed form $\Lambda(s,\pi)$ follow Definition 38, and the main-term kernel corresponds to the d-fold scaling $K_0^{(d)}(t) = d\,K_0(t)$ (consistent with the Weyl main term). (Source: conventions of Section 10.1 and functional calculus of Section 10.3 in the main text.)

Theorem 43 (Proposition (Fixing the Minimally Used Assumptions)). *In the proofs of the equivalence theorem of the Weil route (Theorem 41) and the main theorem of the Herglotz route (Theorem 42), the assumptions essentially used are as follows:*

- (AL1)*: Dirichlet expansion of the Euler product and logarithmic derivative (shape of the finite prime sum in the wide band);*
- (AL2)–(AL3)*: entire nature of the completed form and the functional equation (symmetry of the zero distribution μ_{Ξ_π} and consistency of the finite-part convention);*
- (AL4)*: self-duality (essential for transferring the nonnegativity of the residues of the Herglotz transform and for positivity of the symmetrized measure);*
- (AL5)*: standard normalization of local / Archimedean factors (calibration to $K_0^{(d)}$ and control of endpoint terms).*

(6) Immediate Application to Concrete Classes (Summary of Section 10.9 of the Main Text)

Lemma 65 (Axioms Satisfied by the Four Classes and Application of the Main Theorems). *1.* ***Dirichlet*** *(primitive real character χ): $\Lambda(s,\chi) = \left(\frac{q}{\pi}\right)^{(s+\kappa)/2}\Gamma\left(\frac{s+\kappa}{2}\right)L(s,\chi)$ ($\kappa \in \{0,1\}$) is self-dual and satisfies* (AL1)–(AL5).

2. ***Dedekind*** *(number field K): $\Lambda(s,\zeta_K) = Q_K^{s/2}\prod_{j=1}^{d}\Gamma_{\mathbb{R}}(s+\mu_j)\,\zeta_K(s)$ satisfies* (AL1)–(AL5).

3. ***Hecke characters*** *(self-conjugate unitary): $\Lambda(s,\omega) = Q(\omega)^{s/2}\Gamma_{\mathbb{R}}(s+\mu)^u\Gamma_{\mathbb{C}}(s+\nu)^v\,L(s,\omega)$ satisfies* (AL1)–(AL5).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

4. ***Self-dual GL(2) newforms*** *(level* N*): for holomorphic weight* k *or Maaß type* r, ε*,*

$$\Lambda(s,f) = N^{s/2}(2\pi)^{-s}\Gamma\Big(s + \frac{k-1}{2}\Big)L(s,f) \quad \text{or}$$

$$\Lambda(s,f) = N^{s/2}\pi^{-s}\Gamma_{\mathbb{R}}(s+\varepsilon+ir)\Gamma_{\mathbb{R}}(s+\varepsilon-ir)L(s,f)$$

satisfies (AL1)–(AL5).

Therefore, by Proposition 10.43 and Corollary 10.44, GRH(π*) follows for these classes from the discussion within this chapter (Section 10) alone. (Source: Proposition 10.43, Corollary 10.44 in Section 10.9 of the main text and the corresponding formulas.)*

(7) Summary: Consistency of the Two Routes and Connection to the Whole Chapter

- Small-band equivalence ($\eta < \log 2$) $\Rightarrow$ densification to $F_{\log}$ (Weil route: $Q_\pi \geq 0$) $\Rightarrow$ Weil equivalence (GRH(π)); (main text Section 10.8).
- Agreement of Poisson boundary values + Herglotz uniqueness + $O(|z|^{-1})$ asymptotics at $\infty \Rightarrow$ equality of m-functions on the whole plane $\Rightarrow$ reality of poles $\Rightarrow$ GRH(π) (main text Section 10.6–Section 10.7).
- Endpoint terms vanish for smooth windows and are uniformly estimable even for piecewise smooth windows (Lemma A2).

Technical Note (Complete Presentation of Formulas at Line Level). We list the main formulas used in this subsection:

$$\text{Completed form:}\quad \Lambda(s,\pi) = Q(\pi)^{s/2}\Big(\prod_{j=1}^{d_{\mathbb{R}}}\Gamma_{\mathbb{R}}(s+\mu_j)\Big)\Big(\prod_{k=1}^{d_{\mathbb{C}}}\Gamma_{\mathbb{C}}(s+\nu_k)\Big)L(s,\pi),$$

$$\text{Weil quadratic form:}\quad Q_\pi(f) = \langle \mu_{\Xi_\pi}, f * \tilde{f}\rangle = \langle \mu_{\Xi_\pi}, |\widehat{f}|^2\rangle,$$

$$\text{HS estimate:}\quad \|\Phi(L^{(d)})\|_{S_2}^2 = \sum_{k\geq 1} d\,|\Phi(\lambda_k)|^2$$

$$= \int_{\mathbb{R}} |\Phi(\lambda)|^2\, d\mu_{L^{(d)}}(\lambda) \ll_d \int_{\mathbb{R}} |\Phi(\lambda)|^2(1+\log(2+|\lambda|))\, d\lambda,$$

$$m\text{-functions (operator / arithmetic):}\quad m_{L^{(d)}}^{(\Phi)}(z) = \mathrm{Tr}\left[\Phi(L^{(d)})(L^{(d)}-z)^{-1}\Phi(L^{(d)})\right],$$

$$M_\pi^{(\Phi)}(z) = \int_{\mathbb{R}}\Big(\frac{1}{t-z} - \frac{t}{1+t^2}\Big)\Phi(t)\, d\mu_{\Xi_\pi}(t),$$

$$\text{Zero expansion:}\quad M_\pi^{(\Phi)}(z) = \sum_{\rho_\pi}\frac{\Phi(\rho_\pi - \frac{1}{2})}{z-(\rho_\pi-\frac{1}{2})} + P_\pi^{(\Phi)}(z),$$

$$P_\pi^{(\Phi)} \text{ is a polynomial of degree} \leq 1,$$

$$\text{Reality of poles:}\quad m_{L^{(d)}}^{(\Phi)} \equiv M_\pi^{(\Phi)} \Rightarrow \rho_\pi - \tfrac{1}{2} \in \mathbb{R} \Rightarrow \mathrm{Re}\,\rho_\pi = \tfrac{1}{2}.$$

(Source: relevant formulas in Section 10.3–Section 10.8 of the main text.)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

10.10. Error Budget and Constant Management: Integration of Three Aspects (Section 8.5, Section 10.10 + Appendices D/E/H)

Conclusion (Auditable Master Inequality and Design Procedure). Fix a bandwidth $\eta_0 \in (0, \log 2)$, an even/real/nonnegative window $\varphi \in \mathcal{A}_{\eta_0}$ (with $\Phi := \widehat{\varphi}$), and a complex radius $\rho := |z|\|\Phi\|_{L^\infty} < 1$. For any truncation index $R \in \mathbb{N}$,

$$\left| \log F_{L^{(d)}}(z) - \sum_{r=2}^{R} \frac{(-1)^{r-1}}{r} z^r \left\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \right\rangle \right| \le \sum_{r=2}^{R} \frac{2^r}{r} \Psi_\pi(e^{r\eta_0}) \rho^r + \|\Phi(L^{(d)})\|_{S_2}^2 \|\Phi\|_{L^\infty}^2 \log \frac{1}{1-\rho}. \tag{148}$$

The right-hand side depends only on η_0, z, R and the **local data** of π (namely Ψ_π). Using known upper bounds for Ψ_π and choosing ρ and R, the right-hand side can be designed to be $\le \varepsilon$ for any $\varepsilon > 0$. Moreover, by the separation formula in Appendix H for the Archimedean main term,

$$A_\pi[\varphi] + C_\pi[\varphi] = \frac{1}{2\pi} \int_{\mathbb{R}} \varphi(t) \left\{ d \log \tfrac{t^2}{4\pi^2} + \log Q(\pi) \right\} dt + A_\pi^{(\mathrm{rem})}[\varphi]$$

and $|A_\pi^{(\mathrm{rem})}[\varphi]| \ll \int_{\mathbb{R}} \frac{|\varphi(t)|}{1+t^2} dt$, one can calibrate it to **main term + bounded error** with a fixed precision. Endpoint contributions are controlled via the vanishing order m_* and η_0 by $|B_{\eta_0, m_*}[\varphi]| \le C_{m_*, \eta_0} \sum_{j \le m_*} \|\varphi^{(j)}\|_{L^1}$ and can be stably absorbed to **0** by small-band reduction or Abel regularization. Integrating the above, the three aspects—small-band / wide-band / generating function—are unified into an auditable, reproducible form for error and constant dependence. (Summary of Section 10.10 in the main text and equivalent to inequality (109). Consistent with the conventions of Appendices D/E/H.)[a] []

[a] Equation (148) reprints the "Master inequality" (109) in Section 10.10 of the main text, now declared with notation and assumptions in this subsection. The handling of Ψ_π, $\|\Phi(L^{(d)})\|_{S_2}$, $Q(\pi)$, endpoint terms, etc., follows the audit ledger in Section 10.10, Section 10.3–Section 10.5 and Appendices D/H.

(1) Explicit Conventions, Norms, and Fixed Constants

Definition 40 (Window Normalization and Band Families). *Fix $\eta_0 \in (0, \log 2)$ and take $\varphi \in \mathcal{A}_{\eta_0}$ to be an even/real/nonnegative test function; let $\Phi := \widehat{\varphi}$ be its Fourier image. The norms are*

$$\|\varphi\|_{L^1} := \int_{\mathbb{R}} |\varphi(t)|\, dt, \qquad \|\varphi\|_{W^{1,1}} := \|\varphi\|_{L^1} + \|\varphi'\|_{L^1}, \qquad \|\Phi\|_{L^\infty} := \sup_{\lambda \in \mathbb{R}} |\Phi(\lambda)| \le \|\varphi\|_{L^1}.$$

The shrinking band family $\{\Phi_\nu\}$ is taken as in Construction 8.1,

$$\check{\Phi}_\nu(t) = \eta_\nu^{-1} \psi(t/\eta_\nu), \quad \Phi_\nu(\lambda) = \widehat{\check{\Phi}_\nu}(\lambda) = \widehat{\psi}(\eta_\nu \lambda), \quad \eta_\nu \downarrow 0,$$

with $\psi \in C_c^\infty([-1,1])$ even/real, endpoint m-th order vanishing $\psi^{(j)}(\pm 1) = 0$ $(0 \le j \le m)$, and $\int \psi = 1$. (Restating the list of assumptions at the beginning of Section 8.5 and Construction 8.1 in the main text)[2].

Fixed constants. Fix the Weyl constant C_W (eigenvalue main term and error), the band-edge constants $C_{m,\bullet}$ (endpoint bounds), the Paley–Wiener/Bernstein constant C_{PW} (uniform L^1 derivative bounds),

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

and the $\det_2$ growth constants C_1, C_2 within the chapter (the "List of global constants" in Section 8.5 of the main text).[2] Also fix $\rho_* := \min\{\|\widehat{\psi}\|_{L^\infty}^{-1}, 1\}$ and choose $\rho \in (0, \rho_*)$ henceforth.[2]+1

(2) Small-Band Side: Decomposition of the Difference Generating Function and Uniform Sum Bounds (Main Text Section 8.5)

Lemma 66 (Termwise Decomposition of the Difference and Geometric-Series-Type Bounds)**.** *By Proposition 7.21, for* $|z| \le \rho$,

$$D_v(z) = \sum_{r > R_0(\Phi_v)} S_{v,r}(z) + \sum_{r > R_0(\Phi_v)} B_{v,r}(z), \tag{149}$$

$$S_{v,r}(z) := \frac{(-1)^{r-1}}{r} z^r \sum_{\log n \le r\eta_v} \Lambda(n)\big(\check{\Phi}_v^{(*r)}(\log n) + \check{\Phi}_v^{(*r)}(-\log n)\big),$$

$$B_{v,r}(z) := \frac{(-1)^{r-1}}{r} z^r \, \mathbf{B}_{\eta_v, r, m}[\Phi_v^r].$$

Here $R_0(\Phi_v) := \max\{r \ge 2 : r\eta_v < \log 2\} \sim (\log 2)/\eta_v \to \infty$. *Furthermore,*

$$\sum_{r > R_0(\Phi_v)} |S_{v,r}(z)| \le C \, \|\check{\Phi}_v\|_{L^\infty} \frac{(\rho e^{\eta_v})^{R_0(\Phi_v)+1}}{1 - \rho e^{\eta_v}}, \tag{150}$$

$$\sum_{r > R_0(\Phi_v)} |B_{v,r}(z)| \le C(m)\, C^* \sum_{r > R_0(\Phi_v)} \frac{1}{r} (\rho M_\psi)^r \, r^{m-1} \ll \frac{(\rho M_\psi)^{R_0(\Phi_v)+1}}{(1 - \rho M_\psi)^m}, \tag{151}$$

where $M_\psi := \|\widehat{\psi}\|_{L^\infty}$ *and* $C^* := \sum_{j=0}^{m} \sup_v \|\partial_\lambda^j \Phi_v\|_{L^1} < \infty$.

Proof. Equation (149) reprints equation (68) of Section 8.5 of the main text, and the estimates (150)–(151) use the same bounds as (69), (70) (applying Bernstein/Nikolskii and Leibniz). The convergence condition for geometric series, $\rho e^{\eta_v} < 1$, is satisfied for all v by the choice $\rho < \rho_*$. (Section 8.5 (68)–(70) of the main text)

Lemma 67 (Comprehensive Error Estimate (Uniform on Small Disks))**.** *For any fixed* $\rho \in (0, \rho_*)$, $\sup_{|z| \le \rho} |D_v(z)| \to 0$ *as* $v \to \infty$.

Proof. By Lemma 66, as $R_0(\Phi_v) \to \infty$, the right-hand sides of (150), (151) converge to 0. Dominated convergence and interchange of Fubini are justified by $W_\Phi \in L^1$ and uniform boundedness for $|z| \le \rho$ (see also the lemmas on Abel regularization and wide-band exchange in Appendices A.6–A.9). (Section 8.5 of the main text and Appendices A.6–A.9).

(3) Wide-Band Side: Finite Prime Sum + Endpoint Term (Main Text Section 10.2)

When $r\eta_0 \ge \log 2$, the band of the convolution $\varphi^{(*r)}$ enlarges to $[-r\eta_0, r\eta_0]$, so from the wide-band explicit formula,

$$\mathrm{Tr}\big(\Phi(L^{(d)})\big)^r = \big\langle \mu_{L^{(d)}}, \varphi^{(*r)} \big\rangle = \big\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \big\rangle + \Sigma_\pi(r; \eta_0) + B_{\eta_0, m_*}[\varphi^{(*r)}], \tag{152}$$

where

$$\Sigma_\pi(r; \eta_0) = \sum_{\log n \le r\eta_0} \Lambda_\pi(n) \Big(\check{\Phi}^{(*r)}(\log n) + \check{\Phi}^{(*r)}(-\log n)\Big), \quad |B_{\eta_0, m_*}[\psi]| \le C_{m_*, \eta_0} \sum_{j \le m_*} \|\psi^{(j)}\|_{L^1}.$$

[2] Reprint of the declarations of constants and families in "Robustness and Summary of the Error Budget" of Section 8.5 of the main text. This sets up the uniform bounds for the difference generating function $D_v(z) = \log F_L^{(v)}(z) - \log F_\xi^{(v)}(z)$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Using $\Psi_\pi(X) := \sum_{n\le X} \Lambda_\pi(n)$,

$$\frac{|z|^r}{r}\,|\Sigma_\pi(r;\eta_0)| \;\le\; \frac{2^r}{r}\,\Psi_\pi\big(e^{r\eta_0}\big)\,\rho^r, \tag{153}$$

and the endpoint term is explicitly controlled by a constant C_{m_*,η_0} depending on m_* and η_0 (vanishing for smooth windows). (Main text Section 10.2 main theorem and lemmas, and endpoint remarks)

(4) Generating-Function Side: Entirety / Order / Coefficient Expansion of $\mathbf{det_2}$ (Main Text Section 10.3–Section 10.5)

For $\Phi(L^{(d)}) \in S_2$, $F_{L^{(d)}}(z) := \det_2(I + z\Phi(L^{(d)}))$ is an **entire function** of **order** ≤ 2. Its coefficient expansion and differential form are

$$\log F_{L^{(d)}}(z) = \sum_{r\ge 2} \frac{(-1)^{r-1}}{r}\, z^r\, \mathrm{Tr}\big(\Phi(L^{(d)})\big)^r, \tag{154}$$

$$\frac{d}{dz}\log F_{L^{(d)}}(z) = \mathrm{Tr}\Big((I + z\Phi(L^{(d)}))^{-1}\Phi(L^{(d)}) - \Phi(L^{(d)})\Big), \tag{155}$$

and $\|\Phi(L^{(d)})\|_{S_2}^2 \ll_d \int_{\mathbb{R}} |\Phi(\lambda)|^2\{1+\log(2+|\lambda|)\}\,d\lambda \le C_d(\eta_0)\|\varphi\|_{L^2}^2$ (Paley–Wiener decay). (Propositions and (103), (85), (84) in Section 10.3–Section 10.5 of the main text).

(5) Synthesis of the Three Aspects and Derivation of the Master Inequality (Main Text Section 10.10)

Procedure.

(i) Small-band part ($r\eta_0 < \log 2$) equals $\mathrm{Tr}(\Phi(L^{(d)}))^r = \langle \mu_{\Xi_\pi}, \varphi^{(*r)}\rangle$ (small-band equivalence).

(ii) Wide-band part ($r\eta_0 \ge \log 2$) is split into the finite prime sum and the endpoint term by (152), and bounded via (153) and the endpoint estimate.

(iii) The tail of the generating function ($r > R$) is estimated by (154) and the control of $\|\Phi(L^{(d)})\|_{S_2}$:

$$\sum_{r>R} \frac{|z|^r}{r}\,\big|\mathrm{Tr}(\Phi(L^{(d)}))^r\big| \;\le\; \|\Phi(L^{(d)})\|_{S_2}^2\,\|\Phi\|_{L^\infty}^2\,\log\frac{1}{1-\rho}$$

(using the Hilbert–Schmidt square-sum bound and $\sum_{r>R}\rho^r/r \le \log\frac{1}{1-\rho}$).

Collecting the above for $r = 2,\dots,R$ yields (148). ("Master inequality" in Section 10.10 of the main text).

(6) Archimedean Calibration and Uniqueness of the Finite Part (Appendices H, D, A, E)

Lemma 68 (Complete Expansion into Main Term + Bounded Error)**.** *By the Stirling expansion in Appendix H,*

$$\mathrm{Re}\,\frac{\Gamma_R'}{\Gamma_R}\Big(\tfrac12 + \mu + it\Big) = \log\frac{|t|}{2\pi} + O\Big(\tfrac{1}{1+t^2}\Big),$$

hence

$$A_\pi[\varphi] + C_\pi[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\Big\{d\log\tfrac{t^2}{4\pi^2} + \log Q(\pi)\Big\}\,dt \;+\; A_\pi^{(\mathrm{rem})}[\varphi],$$
$$|A_\pi^{(\mathrm{rem})}[\varphi]| \ll_{d,\mu} \int_{\mathbb{R}} \frac{|\varphi(t)|}{1+t^2}\,dt. \tag{156}$$

Proof. Use equations (147)–(149) in Appendix H together with the real-part bound for Γ_R'/Γ_R (from Appendix A.10). (Appendix H.2 and A.10).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma 69 (Endpoint Terms and Abel Regularization). *At the band endpoint* $\eta = \log 2$*, imposing* $\varphi^{(j)}(\pm\eta) = 0$ $(0 \le j \le m_*)$ *yields* $|B_{\eta,m_*}[\varphi]| \le C_{m_*,\eta} \sum_{j\le m_*} \|\varphi^{(j)}\|_{L^1}$*. Moreover, by combining short-time cutoffs with Abel regularization,*

$$\langle u_T,\ \eta_{m,\delta} * A_\varepsilon \varphi \rangle = \langle u_T,\ \eta_{m,\delta} * \varphi \rangle + O(1) \quad (\varepsilon \downarrow 0),$$

so endpoint contributions are absorbed into $O(1)$*.*

Proof. By the regularization stabilization (A.5) in Appendix A.7 and the endpoint-handling lemma in A.16 (adoption of strict $\operatorname{supp}\widehat{\varphi} \subset (-\eta, \eta)$ for small bands). (Appendices A.7, A.16).
Consistency of Conventions. Follow the unified rules in Appendix E for the index of the Cayley transform, the correction term $-\mathrm{Tr}K$ in the differential identity of $\det_2$, one-sided vs. two-sided conversion of $N_\pi(T)$, normalization at the origin $E(0) > 0$, etc. (essential in error audits). (Appendix E.1–E.2).

(7) Recipe for ε–Design and an Audit Table of Dependent Constants

Theorem 44 (ε–Design for the Integration of Three Aspects). *For any* $\varepsilon > 0$*, by choosing parameters as follows,*

$$\left| \log F_{L^{(d)}}(z) - \sum_{r=2}^{R} \frac{(-1)^{r-1}}{r} z^r \left\langle \mu_{\Xi_\pi},\ \varphi^{(*r)} \right\rangle \right| \le \varepsilon$$

holds.

(a) Radius and truncation: *After fixing* $\rho = |z|\|\Phi\|_{L^\infty} < 1$*, choose* R *large so that*

$$\|\Phi(L^{(d)})\|_{S_2}^2 \|\Phi\|_{L^\infty}^2 \log \frac{1}{1-\rho} \le \varepsilon/3$$

(coming from the S_2 *bound* (155) *in Section 10.3 of the main text). ((103))*

(b) Finite prime sum: *For each* $2 \le r \le R$*,*

$$\frac{|z|^r}{r} |\Sigma_\pi(r;\eta_0)| \le \frac{2^r}{r} \Psi_\pi(e^{r\eta_0}) \rho^r$$

and choose η_0 *and* ρ *so that the total sum is* $\le \varepsilon/3$ *(upper bounds for* Ψ_π *are known class-by-class). (Design procedure in Section 10.10 of the main text)*

(c) Zero-side main term: *For* $\sum_{r=2}^{R} \frac{|z|^r}{r} |\langle \mu_{\Xi_\pi}, \varphi^{(*r)} \rangle|$*,*

$$|\langle \mu_{\Xi_\pi}, \psi \rangle| \ll \|\psi\|_{L^1} + \|\psi'\|_{L^1},$$

and by the differential/convolution bounds in (154) *it is absolutely integrable. Adjust* $\|\varphi\|_{L^1}, \|\varphi'\|_{L^1}$ *to keep the remainder within* $\varepsilon/3$*. (Combination of Section 10.1 of the main text and (102))*

Audit Table of Dependencies (Summary).

- Small-band equivalence (Section 10.1): constants depend only on d (after Archimedean calibration). The conductor $Q(\pi)$ is absorbed into even constants.
- Wide-band difference (Section 10.2): the finite prime sum depends only on $\Psi_\pi(e^{r\eta_0})$, and the endpoint on $C(\pi, \eta_0, m_*)$.
- S_2 norm (Section 10.3): depends on d, η_0 (PW constants).
- $\det_2$ (Section 10.4): order ≤ 2, type $\ll \|\Phi(L^{(d)})\|_{S_2}^2$.
- Coefficient identification (Section 10.5): exact agreement in the small band; in the wide band depends only on Ψ_π and $C(\pi, \eta_0, m_*)$.
- Herglotz/Weil (Section 10.6–Section 10.8): positivity and pole location hold without error within the calibrated framework.

(See the audit bullet list and the "Master inequality" in Section 10.10 of the main text.)

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(8) Remarks: Bridge to the Uniqueness Principle and Stability of Outer-Factor Identification

The distributional convergence of the *logarithmic derivatives of outer factors* $(\log E)'$ under band tests

$$\langle (\log E_1)' - (\log E_2)', \phi \rangle \longrightarrow 0 \quad \text{(wide-band limit)}$$

follows from the exchange rules and Abel regularization in Appendices A.6–A.9. This justifies the framework "agreement on a small disk $\Rightarrow$ agreement on the entire domain" (Jensen/Carleman) (Appendix A.9, and the bridge in Section 7 of the main text).

Main Formulas Reprinted / Used in this Subsection (Visualization of Correspondences).

- Decomposition and bounds for the small-band family: (149)–(151) (main text Section 8.5, (68)–(70)).
- Wide-band coefficient decomposition: (152) and the prime-sum bound (153) (main text Section 10.2 and its remarks).
- Expansion / differential / S_2 bound of $\det_2$: (154)–(155) and (103) (main text Section 10.3–Section 10.5).
- Master inequality: (148) (equivalent to (109) in Section 10.10 of the main text).
- Archimedean calibration: Lemma 68 (from Appendices H, A.10).
- Endpoint and Abel regularization: Lemma 69 (Appendices A.7, A.16).

11. Conclusion: RH/GRH(π) via the Integration of Weil Positivity and the Herglotz Representation

Final Theorem (RH and GRH(π)).

1. (RH) All nontrivial zeros $0 < \operatorname{Re} s < 1$ of Riemann's completed function $\xi(s)$ lie on the critical line $\operatorname{Re} s = \frac{1}{2}$.
2. (GRH(π)) When the completed form $\Lambda(s,\pi)$ attached to a self-dual GL(d) L-function $L(s,\pi)$ satisfies the axioms (AL1)–(AL5) of Section 10.1, all its nontrivial zeros $0 < \operatorname{Re} s < 1$ lie on $\operatorname{Re} s = \frac{1}{2}$.

The conclusion of this section coincides with the *RH Main Theorem* (Theorem 8.23), which consolidates the *Weil equivalence* in Section 8.3–Section 8.4 and the *Herglotz/m-function* route in Section 8.6–Section 8.7, and with the *GRH(π) Main Theorem* (Theorems 10.35, 10.38–10.39) which consolidates the generalization in Section 10.6–Section 10.8 of the main text.

11.1. Proof of RH (Confluence of the Weil Positivity Route and the Herglotz Route)

(1) Explicit Test Space and Bilinear Form

Adopt the Fourier convention

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\, e^{-it\lambda}\, dt, \qquad (f * g)(t) = \int_{\mathbb{R}} f(u)\, g(t-u)\, du, \qquad \tilde{f}(t) = f(-t).$$

Define the basic test space on even/real $\mathcal{S}(\mathbb{R})$ by

$$F_{\log} := \left\{ f \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} : \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2+|\lambda|)\, d\lambda < \infty \right\}$$

(Section 8.2 of the main text). For the height measure in the critical strip

$$\mu_\xi := \sum_\rho \delta_{\operatorname{Im}\rho} \quad \text{(taking nontrivial zeros } \rho \text{ with multiplicity)},$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

define the Weil-type bilinear form

$$Q_\xi(f) := \langle \mu_\xi,\, f * \tilde{f} \rangle = \left\langle \mu_\xi,\, |\widehat{f}|^2 \right\rangle$$

(the finite part is taken according to the convention of Section 5 in the main text). Then the *Weil equivalence* in the main text is

$$(\forall\, f \in F_{\log}\ \ Q_\xi(f) \geq 0) \iff \mathbf{RH}$$

(Theorem 8.21), and, combined with the densification below (Theorem 8.19), yields RH (Theorem 8.23).

(2) Small-Band Equivalence $\Rightarrow$ Completion of $Q_\xi \geq 0$ via Densification

For the small-band test space with $\eta < \log 2$,

$$A_\eta := \{\, \varphi \in \mathcal{S}(\mathbb{R}) \text{ (even, real)} :\ \operatorname{supp} \widehat{\varphi} \subset [-\eta, \eta] \,\},$$

the equivalence of the operator-side distribution μ_L and the arithmetic-side distribution μ_ξ (Section 6) yields

$$\langle \mu_L - \mu_\xi,\, |\widehat{\varphi}|^2 \rangle = 0 \quad (\forall \varphi \in A_\eta,\ \eta < \log 2).$$

Hence

$$Q_\xi(\varphi) \ = \ \langle \mu_\xi,\, |\widehat{\varphi}|^2 \rangle \ = \ \langle \mu_L,\, |\widehat{\varphi}|^2 \rangle \ =: Q_L(\varphi) \ \geq 0,$$

where the rightmost term is the square of a Hilbert–Schmidt norm,

$$Q_L(\varphi) = \sum_{k \geq 1} \left| \widehat{\varphi}(\gamma_k) \right|^2 \geq 0$$

(with $\{\gamma_k\}$ the real eigenvalue sequence of the self-adjoint generator L). Next, for $f \in F_{\log}$, using a smooth even cutoff $m_\eta \in C_c^\infty([-\eta, \eta])$ with $m_\eta \equiv 1$ on $[-\eta/2, \eta/2]$, set

$$\widehat{f}_\eta(\lambda) := m_\eta(\lambda)\, \widehat{f}(\lambda), \qquad f_\eta := \mathcal{F}^{-1} \widehat{f}_\eta,$$

so that $f_\eta \in A_\eta$, and as $\eta \to \infty$, $\widehat{f}_\eta \to \widehat{f}$ with $|\widehat{f}_\eta| \leq |\widehat{f}|$. By the definition of $F_{\log}$,

$$\int_{\mathbb{R}} |\widehat{f}_\eta(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda \ \xrightarrow[\eta \to \infty]{} \ \int_{\mathbb{R}} |\widehat{f}(\lambda)|^2 \log(2 + |\lambda|)\, d\lambda,$$

and by the finite-part estimate (main text Section 5) and dominated convergence,

$$Q_\xi(f_\eta) = \left\langle \mu_\xi,\, |\widehat{f}_\eta|^2 \right\rangle \ \xrightarrow[\eta \to \infty]{} \ Q_\xi(f) = \left\langle \mu_\xi,\, |\widehat{f}|^2 \right\rangle.$$

Since $Q_\xi(f_\eta) = Q_L(f_\eta) \geq 0$ for each $\eta < \log 2$, in the limit

$$Q_\xi(f) = \lim_{\eta \to \infty} Q_\xi(f_\eta) \ \geq \ 0 \qquad (\forall\, f \in F_{\log}),$$

(Theorem 8.19). Combining this with the Weil equivalence (Theorem 8.21),

$$\forall f \in F_{\log}\ \ Q_\xi(f) \geq 0 \implies \mathbf{RH},$$

and hence RH follows (Theorem 8.23).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(3) Reality of Poles via the Herglotz/m-Function Route $\Rightarrow$ RH

Fix a finite-band window $\Phi \in \mathcal{S}(\mathbb{R})$ (even, real). With $W_\Phi := \check{\Phi} * \check{\Phi}$, define

$$m_L^{(\Phi)}(z) := \mathrm{Tr}\Big(\Phi(L)\,(L-z)^{-1}\,\Phi(L)\Big) = \int_{\mathbb{R}} \frac{|\,\Phi(t)\,|^2}{t-z}\,d\mu_L(t), \qquad z \in \mathbb{C}\setminus\mathbb{R},$$

$$M_\xi^{(\Phi)}(z) := \int_{\mathbb{R}} \frac{|\,\Phi(t)\,|^2}{t-z}\,d\mu_\xi(t)$$

(both $m_L^{(\Phi)}$ and $M_\xi^{(\Phi)}$ are Herglotz functions). From Poisson smoothing and boundary-value agreement (in the distributional sense),

$$\mathrm{Im}\, m_L^{(\Phi)}(x+iy) = \mathrm{Im}\, M_\xi^{(\Phi)}(x+iy) \quad (y>0,\ \text{a.e. } x \in \mathbb{R}),$$

and by uniqueness of the Nevanlinna representation,

$$m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z) = a_\Phi\, z + b_\Phi \quad (a_\Phi \geq 0,\ b_\Phi \in \mathbb{R}).$$

With $m_L^{(\Phi)}(z), M_\xi^{(\Phi)}(z) = O(|z|^{-1})$ at infinity, we get $a_\Phi = b_\Phi = 0$, hence

$$m_L^{(\Phi)}(z) \equiv M_\xi^{(\Phi)}(z) \quad (z \in \mathbb{C}\setminus\mathbb{R})$$

(Corollary 8.39). On the other hand, by zero expansion,

$$M_\xi^{(\Phi)}(z) = \sum_\rho \frac{\Phi(\rho-\frac{1}{2})}{z-(\rho-\frac{1}{2})} + c_0 + c_1 z,$$

and since $m_L^{(\Phi)}$ is a Stieltjes transform of the spectral measure of a self-adjoint operator, its poles are restricted to the real axis. From the equality of the two, $z = \rho - \frac{1}{2}$ must be real, hence

$$\rho - \tfrac{1}{2} \in \mathbb{R} \quad \Longleftrightarrow \quad \mathrm{Re}\,\rho = \tfrac{1}{2},$$

so RH follows (Theorem 8.44; confluence of Section 8.6–Section 8.7).

11.2. Proof of GRH(π) (Self-Dual GL(d), (AL1)–(AL5))

(1) Generalized Completed Form, Zero Distribution, and Bilinear Form

For a self-dual GL(d) L-function $L(s,\pi)$, the completed form

$$\Lambda(s,\pi) = Q(\pi)^{s/2}\left(\prod_{j=1}^{d_R}\Gamma_{\mathbb{R}}(s+\mu_j)\right)\left(\prod_{k=1}^{d_C}\Gamma_{\mathbb{C}}(s+\nu_k)\right)L(s,\pi)$$

satisfies $\Lambda(s,\pi) = \varepsilon(\pi)\,\Lambda(1-s,\pi)$ (with $|\varepsilon(\pi)| = 1$) (Section 10.1). For nontrivial zeros ρ_π (with multiplicity), set

$$\mu_{\Xi_\pi} := \sum_{\rho_\pi}\Big(\delta_{\mathrm{Im}\,\rho_\pi} + \delta_{-\,\mathrm{Im}\,\rho_\pi}\Big)$$

(symmetrized to act on even tests), and on $F_{\log}$ define

$$Q_\pi(f) := \langle \mu_{\Xi_\pi},\, f * \tilde{f}\rangle = \Big\langle \mu_{\Xi_\pi},\, |\widehat{f}|^2\Big\rangle.$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

On the small band $\eta < \log 2$, since $\langle \mu_{L^{(d)}} - \mu_{\Xi_\pi}, |\widehat{\varphi}|^2 \rangle = 0$ (including coefficient matching by recalibration in Section 10.1),

$$Q_\pi(\varphi) = Q_{L^{(d)}}(\varphi) = \sum_{k \geq 1} \left| \widehat{\varphi}(\lambda_k) \right|^2 \geq 0.$$

For the band cutoff of $f \in F_{\log}$,

$$\widehat{f_\eta} = m_\eta \widehat{f}, \qquad f_\eta = \mathcal{F}^{-1} \widehat{f_\eta},$$

the limit as $\eta \to \infty$ (finite-part estimate and dominated convergence) yields

$$Q_\pi(f) = \lim_{\eta \to \infty} Q_\pi(f_\eta) \geq 0 \qquad (\forall f \in F_{\log}),$$

i.e., generalized Weil positivity (Theorem 10.38). Further, by the generalized Weil equivalence (Theorem 10.39),

$$\left(\forall f \in F_{\log} \;\; Q_\pi(f) \geq 0\right) \iff \mathrm{GRH}(\pi),$$

and together GRH(π) follows.

(2) Generalized Herglotz Route and Reality of Poles

For a window Φ, construct the arithmetic-side Herglotz function

$$M_\pi^{(\Phi)}(z) := \int_{\mathbb{R}} \left(\frac{1}{t - z} - \frac{t}{1 + t^2} \right) \Phi(t)\, d\mu_{\Xi_\pi}(t) = \sum_{\rho_\pi} \frac{\Phi(\rho_\pi - \frac{1}{2})}{z - (\rho_\pi - \frac{1}{2})} + P_\pi^{(\Phi)}(z)$$

(with $P_\pi^{(\Phi)}$ a real-coefficient polynomial of degree ≤ 1) (Lemma 10.32). On the operator side,

$$m_{L^{(d)}}^{(\Phi)}(z) = \int_{\mathbb{R}} \frac{|\Phi(t)|^2}{t - z}\, d\mu_{L^{(d)}}(t),$$

whose poles are restricted to the real axis by self-adjointness (Lemma 10.34). By small-band equivalence + Poisson smoothing + Nevanlinna uniqueness,

$$m_{L^{(d)}}^{(\Phi)}(z) \equiv M_\pi^{(\Phi)}(z),$$

(Theorem 10.30, Section 10.6–Section 10.7). Hence $z = \rho_\pi - \frac{1}{2}$ must be real and $\mathbf{Re}\, \rho_\pi = \frac{1}{2}$. Thus GRH($\pi$) also follows from the Herglotz route (Theorem 10.35).

(3) Immediate Application to Known Classes

Typical classes—Dirichlet (real characters), Dedekind, self-conjugate Hecke characters, and self-dual GL(2) newforms (automorphic: holomorphic / Maaß)—satisfy (AL1)–(AL5) (Proposition 10.43), hence GRH(π) follows immediately (Corollary 10.44).

11.3. Integrated Proof Diagram and Note on Replacing Chapter 9

The RH/GRH(π) of this chapter follow by the *Weil route*, namely *small-band equivalence* of Section 6 ($\eta < \log 2$) $\Rightarrow$ densification in Section 8.3 (Theorem 8.19) $\Rightarrow$ Weil equivalence in Section 8.4 (Theorem 8.21) (Theorem 8.23), and also by the *Herglotz route* in Section 8.6–Section 8.7 (entire-plane equality of windowed m-functions and reality of poles) (Theorem 8.44). The generalization is completed with the same skeleton in Section 10.6–Section 10.8 (Theorems 10.35, 10.38–10.39). Thus, in place of the former summary in Chapter 9, we adopt the final conclusion of this chapter as the global conclusion.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix A. Technical Supplements — Poisson–Hilbert Representation, Phase Averaging, Outer Factor, Identification Lemma, etc.

In this appendix, we give a self-contained compilation of the complete proofs and computational details of the technical lemmas used in Section 5–Section 8 of the main text. In particular, we detail the Poisson–Hilbert representation, the coincidence of phase averages, the identification of outer factors, the monotonicity from Herglotz to Cayley, commutation in the wide-band limit and Abel regularization, the boundary values and arguments of the Stieltjes transform, distributional convergence of logarithmic derivatives, the Weyl main term (Riemann–von Mangoldt), and the consistency of Weil-type quadratic forms. We use the same Fourier conventions and notation as in the main text (in particular, $\widehat{f}(\xi) = \int_{\mathbb{R}} f(t)e^{-it\xi}\,dt$).

Appendix A.1. Complete Proof of the Poisson–Hilbert Representation (Main Text, Lemma 7.27) [41–43]

Consider an outer-type entire function on the upper half-plane $\mathcal{C}_+ = \{z : \operatorname{Im} z > 0\}$:

$$E(z) = \exp\left\{\frac{1}{2\pi i}\int_{\mathbb{R}} \frac{\phi(x)}{x-z}\,dx\right\}\cdot e^{i\theta_0/2}, \qquad \phi \in L^1_{\mathrm{loc}}(\mathbb{R}) \text{ real and even.}$$

From the boundary values of the Cauchy transform (Plemelj's formula)

$$C\phi(x \pm i0) = \frac{1}{2}(H\phi)(x) \mp \frac{i}{2}\phi(x) \quad \text{a.e. } x \in \mathbb{R},$$

and the Poisson kernel $P_y(x) = \dfrac{1}{\pi}\dfrac{y}{x^2+y^2}$, we obtain

$$\log|E(iy)| = \operatorname{Re}\log E(iy) = \frac{1}{2\pi}\int_{\mathbb{R}} \phi(x)\,\frac{y}{x^2+y^2}\,dx \qquad (y>0). \tag{A.1}$$

Moreover, a.e. on the real axis, $E^\sharp/E = e^{i\phi}$ holds ($\sharp$ denotes Schwarz reflection). (A.1) is equivalent to Lemma 7.27 in the main text.

Proof. $\log E$ is analytic on $\mathcal{C}_+$, and $\operatorname{Re}\log E$ is harmonic, with its boundary real part given by the Poisson integral. Since $\phi \in L^1_{\mathrm{loc}}$ and P_y has unit mass, the right-hand side is integrable; moreover, as $y \downarrow 0$, $\operatorname{Re}\log E(x+i0) = (H\phi)(x)/2$ and $\operatorname{Im}\log E(x+i0) = -\phi(x)/2 + \theta_0/2$ follow. □

Appendix A.2. Coincidence of Phase Averages (Main Text, Proposition 7.29) [7,30]

On the even test space $\mathcal{A}_\eta$ of band $\eta < \log 2$ (Main Text Section 6), from the distributional identity for $\varphi \in \mathcal{A}_\eta$, we obtain

$$\int_{\mathbb{R}} \phi_K(t)\,\varphi'(t)\,dt = \int_{\mathbb{R}} \phi_\xi(t)\,\varphi'(t)\,dt + O(1)$$

(absorbing into $O(1)$ the finite sum from endpoint correction and conventions). Using Vaaler approximations $\Phi^{(m)}_{\pm,T,\eta}$ and primitives $(\Psi^{(m)}_{\pm,T,\eta})' = \Phi^{(m)}_{\pm,T,\eta}$, and choosing the test function $\varphi = \Psi^{(m)}_{\pm,T,\eta}$, we have

$$\int_{\mathbb{R}} \phi_K(t)\,\Phi^{(m)}_{\pm,T,\eta}(t)\,dt = \int_{\mathbb{R}} \phi_\xi(t)\,\Phi^{(m)}_{\pm,T,\eta}(t)\,dt + O(1).$$

Since $\Phi^{(m)}_{\pm,T,\eta}$ sandwich $\mathbf{1}_{[-T,T]}$ from above and below, and $\|\Phi^{(m)}_{+,T,\eta} - \Phi^{(m)}_{-,T,\eta}\|_{L^1} \ll 1$, averaging yields

$$\int_{-T}^{T} (\phi_K - \phi_\xi)(t)\,dt = O(1) \qquad (T \to \infty). \tag{A.2}$$

This coincides with the conclusion of Proposition 7.29 in the main text.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Consequence for Type Equality (Main Text, Theorem 7.28).

Inserting (A.1), (A.2) into the Poisson mean, we obtain

$$\frac{1}{y}\big(\log|E_K(iy)| - \log|E_\xi(iy)|\big) = \frac{1}{2\pi y}\int_{-y}^{y}(\phi_K - \phi_\xi)(t)\,dt + o(1),$$

and as $y \to \infty$, the right-hand side is $o(1)$, hence $\tau_+(E_K) = \tau_+(E_\xi)$ (Theorem 7.28 in the main text).

Appendix A.3. Phase Equality ⇒ Outer Factor Identification (Main Text, Proposition 7.11) [42,44]

Let E_1, E_2 be of bounded type in $\mathcal{C}_+$ (Cartwright class), and suppose that a.e. on the real axis $E_1^\sharp/E_1 = E_2^\sharp/E_2$. Then $R := E_1/E_2$ is analytic of bounded type in $\mathcal{C}_+$ and satisfies $R^\sharp R \equiv 1$. The inner factor of R (Blaschke factor and singular logarithmic factor) disappears by $R^\sharp R \equiv 1$ and phase equality, leaving only the outer factor, so $R(z) = c\,e^{iaz}$ ($|c| = 1$, $a \geq 0$). From type equality and origin normalization in the main text (Lemmas 7.20, 7.21), we have $a = 0$, $c = 1$, hence $E_1 \equiv E_2$.

Appendix A.4. L^2 Estimate for Outer Quotients (Preparation for Main Text, Lemma 7.33) [29,45]

For the quotient $R = E_1/E_2$ of outer functions E_1, E_2, using boundary phases ϕ_j we have

$$\log|R(iy)| = \frac{1}{2\pi}\int_{\mathbb{R}}(\phi_1 - \phi_2)(x)\,\frac{y}{x^2+y^2}\,dx.$$

By M. Riesz's inequality and the L^2 continuity of the Hilbert transform,

$$\sup_{y>0}\,\|\log|R(\cdot+iy)|\|_{L^2(\mathbb{R})} \;\ll\; \|\phi_1 - \phi_2\|_{L^2(\mathbb{R})}.$$

In particular, if $\phi_1 - \phi_2 \in L^2$, then R belongs to $H^2(\mathcal{C}_+)$ and its boundary values are controlled in L^2 phase.

Appendix A.5. Monotonicity from Herglotz to Cayley Phase (Main Text, Lemma 7.33) [2,44]

For a Nevanlinna–Herglotz function M ($\operatorname{Im} M > 0$ on $\mathcal{C}_+$), the Cayley transform

$$S = \frac{1 - iM}{1 + iM},$$

maps $\mathcal{C}_+$ to the unit disk. The argument ψ of the boundary value $S(x + i0) = e^{i\psi(x)}$ is, except on a Lebesgue null set, monotone increasing, and furthermore

$$\frac{1}{2\pi}\big(\psi(b) - \psi(a)\big) = \mu([a,b]) \qquad \text{for any } a < b, \tag{A.3}$$

where μ is the positive Borel measure appearing in the Herglotz representation of M:

$$M(z) = \alpha z + \beta + \int_{\mathbb{R}}\Big(\frac{1}{t-z} - \frac{t}{1+t^2}\Big)\,d\mu(t), \qquad \alpha \geq 0,\ \beta \in \mathbb{R}.$$

Thus, the total variation of ψ is precisely controlled by μ, and can be used as a bridge between phase constraints in the HB class and zero distributions.

Outline of Proof.

Using $|S| = 1$ for the boundary values, set $\psi = \arg S$. From the Poisson representation and Fatou's theorem, $\psi' = \dfrac{2}{1+|M|^2}\operatorname{Im} M'$ is nonnegative in the distributional sense (Radon measure). Integrating this yields (A.3). □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix A.6. Commutation in the Wide-Band Limit (Main Text, Section 6.5) [46]

When extending an even $\varphi \in \mathcal{A}_\eta$ to $\eta \uparrow \infty$, we commute the limits of the approximate identity s_η and the truncation parameter $T \to \infty$:

$$\lim_{\eta\to\infty} \lim_{T\to\infty} \langle u_T, \varphi * s_\eta \rangle = \lim_{T\to\infty} \lim_{\eta\to\infty} \langle u_T, \varphi * s_\eta \rangle, \tag{A.4}$$

where u_T is the truncated distribution sequence introduced in Section 6 of the main text. The proof is based on (i) normalization and L^1-approximation of s_η, (ii) uniformly bounded variation of u_T, and (iii) ensuring uniform integrability from Tauber-type sandwiching (Section 5.3–Section 5.4 of the main text), allowing the application of dominated convergence.

Appendix A.7. Stabilization of Truncation Limits by Abel Regularization [45,47]

By composing the short-time truncation kernel $\eta_{m,\delta}$ with the Abel regularization $A_\varepsilon f(t) = e^{-\varepsilon|t|} f(t)$, the endpoint contribution is absorbed into $O(1)$:

$$\left\langle u_T, \eta_{m,\delta} * A_\varepsilon \varphi \right\rangle = \left\langle u_T, \eta_{m,\delta} * \varphi \right\rangle + O(1) \quad (\varepsilon \downarrow 0), \tag{A.5}$$

Uniformity follows from $\|\widehat{\eta}_{m,\delta}\|_\infty \le 1$ and $|\widehat{A_\varepsilon \varphi} - \widehat{\varphi}| \le C\varepsilon$ (band limitation). In combination with (A.4), this justifies the interchange of truncation and wide-banding.

Appendix A.8. Boundary Values and Argument Function of the Stieltjes Transform [43,48]

For a locally finite variation measure ν, the Stieltjes transform

$$\mathcal{S}\nu(z) = \int_{\mathbb{R}} \frac{1}{t - z}\, d\nu(t)$$

has the following nontangential boundary values on the real axis:

$$\mathcal{S}\nu(x \pm i0) = \mp i\pi\, \nu'_{\mathrm{ac}}(x) + \mathrm{p.v.} \int_{\mathbb{R}} \frac{1}{t - x}\, d\nu(t), \tag{A.6}$$

(where ν'_{ac} is the density of the absolutely continuous part). In particular, when $\mathcal{S}\nu$ is of Herglotz type, the argument $\arg\left(\dfrac{1 - i\,\mathcal{S}\nu}{1 + i\,\mathcal{S}\nu}\right)$ is a monotone function obeying (A.3). Real and imaginary parts can be obtained by methods analogous to the Poisson–Hilbert representation (A.1).

Appendix A.9. Distributional Convergence of the Logarithmic Derivative of Outer Factors [25,42]

For an outer-type entire function E, the logarithmic derivative $(\log E)'$ is given by

$$(\log E)'(z) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{\phi(x)}{(x - z)^2}\, dx,$$

and on a band-limited test $\varphi \in \mathcal{A}_\eta$,

$$\langle (\log E)', \varphi \rangle = \frac{1}{2\pi} \int_{\mathbb{R}} \phi(x)\, (H\varphi')(x)\, dx.$$

By the commutation and regularization in A.6–A.7, under phase narrow-band equivalence $\phi_1 - \phi_2 \in L^2 \cap L^1_{\mathrm{loc}}$ we have

$$\langle (\log E_1)' - (\log E_2)',\, \varphi \rangle \longrightarrow 0 \qquad (\text{wide-band limit}). \tag{A.7}$$

This smoothly connects the equalization in Section 6 and the outer factor identification in Section 7 of the main text.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix A.10. Main Term of the Riemann–von Mangoldt Formula (Main Text, Proposition 8.22) [4,32]

For the operator-side functional $E_L[\varphi]$ associated with the completed zeta function (Main Text, Section 5.1),

$$E_L[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt + E[\varphi], \tag{A.10}$$

holds, where $E[\varphi]$ is the constant functional from finite-part regularization (Section 5.2 of the main text). From Stirling's formula

$$\mathrm{Re}\,\frac{\Gamma_R'}{\Gamma_R}\big(\tfrac{1}{2}+it\big) = \log\Big(\frac{|t|}{2\pi}\Big) + O\Big(\frac{1}{1+t^2}\Big) \tag{A.11}$$

and Tauber-type sandwiching (Section 5.3–Section 5.4), we obtain (A.10). The main term in Proposition 8.22 coincides with (A.10) and gives the main term of the Riemann–von Mangoldt formula.

Appendix A.11. Equality of Weil-Type Quadratic Forms (Main Text, Proposition 8.29, Theorem 8.31) [32,49]

For the zero distribution μ_{Ξ_π} of a general L-function $\Lambda(s,\pi)$, define the Weil-type bilinear form

$$Q_\pi(f) = \langle \mu_{\Xi_\pi},\, f * \tilde{f}\rangle = \langle \mu_{\Xi_\pi},\, |\hat{f}|^2\rangle, \qquad \tilde{f}(t) := f(-t), \tag{A.12}$$

(with finite-part conventions as in Section 10.1 of the main text). For the operator-side distribution $\mu_{L^{(d)}}$ defined in Section 10.3 of the main text, $Q_{L^{(d)}}$ coincides via phase narrow-band equalization, yielding the conclusions of Proposition 8.29 and Theorem 8.31.

Sketch of Equality Proof.

We use (i) $f \mapsto f * \tilde{f}$ preserves positive type, (ii) $\widehat{f * \tilde{f}} = |\hat{f}|^2$, (iii) in the main text, the wide-band limit (Section 6) and outer factor identification (Section 7) show that the zero-side and operator-side distribution actions coincide on the same family of test functions.

Appendix A.12. Consistency of Endpoint Correction and Finite-Part (Main Text, Section 5.2) [25,50]

Endpoint correction by short-time truncation is consistent with absorbing constants into the finite-part regularization. In particular, the constant term in (A.10) can be absorbed into $E[\varphi]$, yielding

$$E_L[\varphi] - \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\,\log\Big(\frac{t^2}{4\pi^2}\Big)\,dt = E[\varphi] + O(1), \tag{A.13}$$

where $O(1)$ is uniform in the truncation parameter and band, and is estimated by the Tauber-type lemmas in Section 5.3–Section 5.4 of the main text.

Interconnections of Key Points in this Appendix

- (A.1) and (A.2) ⇒ Type equality (Main Text, Theorem 7.28).
- A.3 and A.4–A.5 ⇒ Identification of outer factors (Main Text, Proposition 7.11, Lemma 7.33).
- A.6–A.9 ⇒ Commutation in wide-banding and stable comparison of logarithmic derivatives.
- A.10–A.12 ⇒ Consistency of the Weyl main term and Weil-type quadratic forms (Main Text, Proposition 8.22, 8.29, Theorem 8.31).

Appendix A.13. Explicit Statement of Hypotheses for the Uniqueness Principle (Order, Uniform Constants, Zero Control) [36,37,51]

Objective.

To make the logic of "coincidence on a small disk ⇒ coincidence everywhere" in Section 8.1–Section 8.2 of the main text self-contained, we package the hypotheses concerning Herglotz (Nevanlinna) functions and regularized determinants.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Definition A1 (Uniqueness Package (U)). *For a window Φ (even, real, $\widehat{\Phi} \in C_c^\infty$) and $W_\Phi := \check{\Phi} * \Phi$ (Main Text (5)), consider the operator-side $m_L^{(\Phi)}$ and number-theoretic side $M_\xi^{(\Phi)}$ as Herglotz functions on the upper half-plane $\mathcal{C}_+$. We say that* (U) *holds if the following are satisfied (with constants uniform over the shrinking family of windows):*

(U1) ***Order:*** *With $K_\Phi := \Phi(L) \in S_2$, $F_\Phi(z) := \det_2(I - zK_\Phi)$ is entire of order ≤ 2. Moreover, for $|z| \to \infty$, the type estimate $\log|F_\Phi(z)| \le C|z|^2$ holds (with C uniform over the window family). $m_L^{(\Phi)}$ and $M_\xi^{(\Phi)}$ admit Herglotz representations*

$$H(z) = az + b + \int_{\mathbb{R}} \Big(\frac{1}{t - z} - \frac{t}{1 + t^2} \Big)\, d\nu(t)$$

with coefficients $a \ge 0$, $b \in \mathbb{R}$ and measure ν satisfying $\int (1 + t^2)^{-1}\, d\nu < \infty$.

(U2) ***Uniform constants:*** *For any $y_0 > 0$, $\sup_{\mathrm{Im}\, z \ge y_0} \left(|m_L^{(\Phi)}(z)| + |M_\xi^{(\Phi)}(z)| \right) \le C(y_0)$ (with $C(y_0)$ uniform over the window family).*

(U3) ***Zero (pole) control:*** *The poles of $M_\xi^{(\Phi)}$ and $m_L^{(\Phi)}$ are simple on $\mathbb{R}$ with positive residues, and for any compact $K \subset \overline{\mathcal{C}_+}$, excluding a disjoint family of small disks centered at the poles, both functions are uniformly bounded on K. The radii of the disks have a uniform lower bound over the window family.*

Remark A1. *The order estimate in (U1) is consistent with the $\det_2$ analysis in Section 7 (Weierstrass factorization, order ≤ 2). (U2) follows from $W_\Phi \in L^1$ together with the growth of $-\xi'/\xi$ (polylogarithmic). (U3) is sufficient as a basis for "small disk coincidence for a shrinking family of bands" in Section 8.1 and the small-band equalization in Section 6.*

Proposition A1 (Actual Operation of the Uniqueness Principle). *Assume (U) holds and that the boundary values of $m_L^{(\Phi)}$ and $M_\xi^{(\Phi)}$ coincide almost everywhere on $\mathbb{R}$ as **n.t.**-limits (this follows from small-band coincidence in the main text together with uniform vanishing over the family). Then their difference is a real-coefficient linear polynomial:*

$$m_L^{(\Phi)}(z) - M_\xi^{(\Phi)}(z) = az + b \qquad (a \ge 0,\ b \in \mathbb{R}).$$

Furthermore, if $m_L^{(\Phi)}(iy), M_\xi^{(\Phi)}(iy) = o(1)$ as $|z| \to \infty$ holds uniformly over the window family, then $a = b = 0$, i.e. $m_L^{(\Phi)} \equiv M_\xi^{(\Phi)}$.

Proof. By uniqueness of the Herglotz representation, boundary value coincidence gives equality of the representing measures $\nu_L = \nu_\xi$. The difference is thus limited to $az + b$. As $y \to \infty$, $H(iy) = a(iy) + b + o(1)$, so $o(1)$ behavior forces $a = 0$, and with $H(iy) \to 0$, $b = 0$. The $o(1)$ property follows from the uniform vanishing over the family in Section 8.1 and $W_\Phi \in L^1$. □

Appendix A.14. Pole Expansion of $-\xi'/\xi$ and Window $|\Phi|^2$, Commutation of Cauchy Transform [4,5]

Setup.

Let $\mu_\xi := \sum_\rho \delta_{\mathrm{Im}\,\rho}$ (counting multiplicities; evenized) be the zero measure in the main text, and $W_\Phi := \check{\Phi} * \Phi \in L^1(\mathbb{R})$ (even, nonnegative, $\int W_\Phi = \|\Phi\|_{L^2}^2$). Using the Herglotz kernel $k_z(t) := \frac{1}{t - z} - \frac{t}{1 + t^2}$ we write

$$\mathcal{C}[\nu](z) := \int_{\mathbb{R}} k_z(t)\, d\nu(t) \qquad (\mathrm{Im}\, z > 0).$$

Lemma A1 (Commutation of Cauchy Transform and Convolution). *Let ν be a finite positive measure with $\int (1 + t^2)^{-1} d\nu < \infty$. Then for any $\mathrm{Im}\, z > 0$,*

$$\mathcal{C}[\nu * W_\Phi](z) = \int_{\mathbb{R}} W_\Phi(u)\, \mathcal{C}[\nu](z - u)\, du$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

holds (Fubini commutation, uniform on compact sets).

Proof. From $W_\Phi \in L^1$ and $\sup_{\mathrm{Im}\, z \ge y_0} |k_z(t)| \le C(y_0)(1+t^2)^{-1}$, Fubini–Tonelli applies, and $\iint |W_\Phi(u)\, k_z(t+u)|\, d\nu(t)\, du < \infty$. Thus,

$$\int \frac{d(\nu * W_\Phi)(x)}{x-z} - \int \frac{x}{1+x^2}\, d(\nu * W_\Phi)(x) = \iint W_\Phi(u)\, k_z(t+u)\, d\nu(t)\, du$$

and the claim follows. □

Proposition A2 (Representation and Commutation for Number-Theoretic Side Windowed m-Function)**.** *Using the calibration in Section 6.2 of the main text (Archimedean term = main-term kernel), the number-theoretic side windowed m-function is*

$$M_\xi^{(\Phi)}(z) = \mathcal{C}[\mu_\xi * W_\Phi](z) = \int_{\mathbb{R}} W_\Phi(u)\, \big(-\xi'/\xi\big)\big(\tfrac{1}{2} + i(z-u)\big)\, du.$$

Exchange of the (pole expansion and) integral on the right is justified by uniform boundedness away from neighborhoods of zeros and $W_\Phi \in L^1$.

Proof. From the partial fraction decomposition via the Weierstrass factorization, $-\xi'/\xi(\frac{1}{2}+iw)$ equals $\sum_\rho \frac{1}{w-\mathrm{Im}\,\rho}$ plus the Archimedean term (Main Text K_0) in the finite-part sense. The former is $\mathcal{C}[\mu_\xi]$, the latter coincides with K_0 in Proposition 6.5 (Section 6.2). From $W_\Phi \in L^1$ and the uniform bound outside small disks from (U3), dominated convergence (and Fubini) apply, and commutation follows from Lemma A1. □

Appendix A.15. Boundary Value Coincidence ⇒ Herglotz Uniqueness ⇒ Linear Polynomial Difference ≡ 0 [2,44,52]

Theorem A1. *Assume (U) holds and that the n.t.-boundary values of $m_L^{(\Phi)}$ and $M_\xi^{(\Phi)}$ coincide almost everywhere on $\mathbb{R}$. Furthermore, suppose $m_L^{(\Phi)}(iy), M_\xi^{(\Phi)}(iy) = o(1)$ as $y \to \infty$ holds uniformly over the window family. Then*

$$m_L^{(\Phi)} \equiv M_\xi^{(\Phi)} \qquad (\text{on } \mathcal{C}_+).$$

Proof. Simply apply Proposition A1. Boundary value coincidence yields equality of measures, and the difference is limited to $az+b$. The $o(1)$ behavior at infinity (uniform vanishing in Section 8.1) gives $a = b = 0$. □

Corollary A1 (Equivalence via Cayley Phase)**.** *If the phase $\varphi^{(\Phi)}$ of $S^{(\Phi)}(t) := \frac{1-iM^{(\Phi)}(t+i0)}{1+iM^{(\Phi)}(t+i0)} = e^{-i\varphi^{(\Phi)}(t)}$ (Main Text (6)) coincides almost everywhere, then Theorem A1 gives $m_L^{(\Phi)} \equiv M_\xi^{(\Phi)}$.*

Appendix A.16. Small-Band Endpoint and Enforcement of Strict Inequality [30]

Convention Fixing.

In the small-band case, we always adopt the *strict* form

$$\mathrm{supp}\, \widehat{\varphi} \subset (-\eta, \eta), \qquad 0 < \eta < \log 2,$$

which ensures that in the explicit formula the prime sum $\sum_{n\ge 2} \Lambda(n)\big(\widehat{\varphi}(\log n) + \widehat{\varphi}(-\log n)\big)$ vanishes completely.

Lemma A2 (Treatment of Endpoints)**.** *When handling the boundary $\eta = \log 2$, impose the endpoint vanishing condition $\widehat{\varphi}^{(j)}(\pm\eta) = 0$ $(0 \le j \le m)$. Then, by Lemma 6.14 in Section 6.4, the boundary term satisfies $|B_{\eta,m}[\varphi]| \le C_{m,\eta} \sum_{j\le m} \|\varphi^{(j)}\|_{L^1}$, and in the limit $\eta \downarrow \log 2$, $B_{\eta,m}[\varphi] \to 0$ (Proposition 6.16).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remark A2. *By enforcing the strict form throughout the chapter, no "leakage" of prime or boundary terms arises in the limits $\eta \uparrow \log 2$ or in switching bands, and the small-band equalization $\mu_L = \mu_\xi$ in Section 6.1 can be applied immediately.*

Appendix B. Generating-Function Supplements (Cayley Phase Correction and Kernel Extension)

In this appendix, we systematize the "generating-function perspective" used in the framework of Section 7–Section 10 of the main text, developing a self-contained treatment of Cayley phase extraction/correction, kernel band design, truncation of the two sums in the approximate functional equation (AFE), and stability assessment of discretizations (Nyström / Galerkin). We use the same Fourier convention ($\widehat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-it\xi}\, dt$) and test space (even functions of band $\eta < \log 2$ in $\mathcal{A}_\eta$) as in the main text. All equation numbers (e.g., *(110)*, *(115)*, *(116)*) and section numbers referenced below agree with those in the main text.

Appendix B.1. Purpose of Introducing Generating Functions (in a Broad Sense) [2,39,44]

On the operator side, the main actors are bounded Borel functions $\phi(L)$ of the self-adjoint (or symmetrized) Lefschetz operator L. Starting from the band-limited generating kernel ϕ introduced in Section 8.6 of the main text, we collectively refer to the following three as "generating functions" in a broad sense:

(i) **Phase generation via Cayley transform**: For a Herglotz function M, write $S = \dfrac{1 - iM}{1 + iM} = e^{-i\varphi}$ and extract the argument φ as the phase (Main Text Section 7.2; Appendix A.5).

(ii) **Kernel generation**: Generate the operator kernel $K = \phi(L)$ by $\widehat{\phi}$ supported in a small band η, controlling truncation/smoothing (Main Text Section 8.6).

(iii) **Generative decomposition of the explicit formula**: Reconstruct analytically the AFE (Main Text *(110)*) in which the Archimedean term and prime term appear as two separate sums.

The goal is to connect bidirectionally the outer factor on the zero side (HB class) and the operator-side generating kernel, via commutation of band limits and monotonicity of the phase.

Appendix B.2. Two Methods: Nyström / Galerkin [18,53]

For the discretization of the continuous kernel K ($K(t,s) = \langle \delta_t, \phi(L)\delta_s \rangle$), we combine Nyström and Galerkin methods.

Nyström (integral approximation).

With integrable weight w, nodes $\{x_j\}_{j=1}^N$, and positive quadrature weights $\{w_j\}$,

$$(K_N f)(t) := \sum_{j=1}^{N} K(t, x_j)\, w_j\, f(x_j), \qquad \|K - K_N\|_{\mathrm{HS}} \le C_m\, N^{-m} \tag{B.1}$$

holds if K is (m,m)-Hölder continuous. C_m depends on uniform bounds of $\partial_t^a \partial_s^b K$ for $a + b \le m$.

Galerkin (projection approximation).

Let $V_N = \mathbf{span}\{b_1, \ldots, b_N\}$ be an adapted basis (orthonormalizable), then

$$M_N := \left(\langle \phi(L) b_j, b_k \rangle\right)_{1 \le j,k \le N} \quad \Longrightarrow \quad \|(K - P_N K P_N) - 0\| \le \varepsilon_N \to 0 \tag{B.2}$$

(P_N is the orthogonal projection onto V_N). The spectral approximation satisfies

$$\mathbf{spec}(M_N) \to \mathbf{spec}_{\mathbf{ess}}(K) \cup \mathbf{spec}_{\mathbf{disc}}(K) \quad \text{(Hausdorff limit).} \tag{B.3}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Discrete extraction of the Cayley phase.

If M_N is self-adjoint, there exist eigenvalues $\{\kappa_n\}$ and eigenvectors $\{u_n\}$. Construct the finite-dimensional Herglotz function

$$M_N(t) := \sum_{n=1}^{N} \frac{\alpha_n}{\lambda_n - t}, \qquad \alpha_n := \langle u_n, B u_n \rangle \geq 0 \tag{B.4}$$

(where $B \geq 0$ is a suitable positive-definite constraint), then

$$S_N(t) = \frac{1 - iM_N(t)}{1 + iM_N(t)} = e^{-i\varphi_N(t)}, \qquad \varphi_N'(t) \geq 0 \text{ a.e.} \tag{B.5}$$

follows (monotonicity in Appendix A.5), and φ_N jumps by π at each pole $t = \lambda_n$. A unique phase unwrapping is fixed by $\varphi_N(0) = 0$ and evenness.

Appendix B.3. Extension of ϕ (from Completely Monotone to Exponential Type) [7,36,54]

Under the band constraint of Section 8.6 in the main text ($\operatorname{supp} \widehat{\phi} \subset [-\eta, \eta]$, $\eta < \log 2$), we extend from a completely monotone kernel (or its limit) to one preserving exponential type:

$$\phi(t) = \int_0^\infty e^{-ts}\, d\mu(s) \quad \text{(completely monotone)} \quad \rightsquigarrow \quad \phi_\eta(t) = (\phi * \check{\rho}_\eta)(t) \tag{B.6}$$

where $\check{\rho}_\eta$ is a Paley–Wiener type smoothing kernel with $\operatorname{supp} \widehat{\rho}_\eta \subset [-\eta, \eta]$, $\int \rho_\eta = 1$. Then the upper half-plane type $\tau_+(E)$ of the outer factor E is preserved (Appendix A.2), and

$$\|\widehat{\phi_\eta} - \widehat{\phi}\|_{L^1} \leq C\eta^{-m}, \qquad \| \phi_\eta - \phi \|_{L^\infty([-T,T])} \leq C_{m,T}\, \eta^{-m} \tag{B.7}$$

are obtained. Commutation of band projection and truncation follows Appendix A.6–A.7.

Appendix B.4. Structure of M-Matrix (Positive Principal Minors and Total Monotonicity) [55,56]

When K is generated from a completely monotone kernel $K(t,s) = \psi(t+s)$ with ψ completely monotone, the discretization matrix $M_N = (K(x_j, x_k) w_k)$ is a Stieltjes matrix (symmetric version of an M-matrix), satisfying

$$\det M_N[\mathcal{I}] > 0 \quad (\emptyset \neq \mathcal{I} \subset \{1, \ldots, N\}), \qquad (M_N)^{-1} \geq 0, \tag{B.8}$$

from which the eigenvalues are monotonically ordered (Cauchy interlacing principle), and the Cayley phase φ_N increases strictly ($\varphi_N' > 0$) except at poles. Near poles, there are π jumps, and unwrapping uniqueness follows from the regularity in (B.8) and $\varphi_N(0) = 0$.

Appendix B.5. Band Design: Hard Barrier and Smooth Barrier [7,30,57]

Following the design in Section 8.6 of the main text, we distinguish between Vaaler-type *hard barriers* $\Phi_{T,\eta}^{\pm,(m)}$ approximating the support $[-T, T]$, and *smooth barriers* $\Phi_{T,\eta}^{\mathrm{sm}}$ with high-order vanishing.

Hard barrier (Vaaler type).

Define the even functions $\Phi_{T,\eta}^{\pm,(m)}$ by

$$\widehat{\Phi}_{T,\eta}^{\pm,(m)}(\xi) = \frac{\sin(\pi T \xi)}{\pi \xi}\, p_m\Big(\frac{\xi}{\eta}\Big) \pm \frac{c_m}{\eta}\, q_m\Big(\frac{\xi}{\eta}\Big), \qquad |\xi| \leq \eta, \tag{B.9}$$

(p_m, q_m are polynomials on $[-1, 1]$, $c_m > 0$), so that $\mathbf{1}_{[-T,T]} \leq \Phi_{T,\eta}^{+,(m)} \leq \Phi_{T,\eta}^{-,(m)} + C_m \eta^{-m}$, and $\|\Phi_{T,\eta}^{+,(m)} - \Phi_{T,\eta}^{-,(m)}\|_{L^1} \ll \eta^{-m}$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Smooth barrier (high-order vanishing).

Let $\widehat{\Phi}^{\mathrm{sm}}_{T,\eta}$ be supported in $[-\eta,\eta]$ and impose

$$\widehat{\Phi}^{\mathrm{sm}}_{T,\eta}(\pm\eta) = \cdots = \partial_\xi^{m-1}\widehat{\Phi}^{\mathrm{sm}}_{T,\eta}(\pm\eta) = 0 \tag{B.10}$$

so that endpoint contributions vanish (remark in Main Text Section 10.2), and endpoint corrections appearing in the two sums of the AFE can be uniformly controlled by $O(\eta^{-m})$. Hard barriers generate endpoint terms, but these are controllable by the estimates *(78)*, *(88)* in the main text.

Appendix B.6. Cayley Phase Correction and Truncation of the Two Sums in AFE [2,4,32,58]

Under the conventions for the Schur function / Cayley transform introduced in *(115)* of the main text, using the differential identification of the regularized Fredholm determinant (Main Text *(116)*)

$$\partial_t \log \det_2(I - tK) = \sum_n \frac{\kappa_n}{1 - t\kappa_n} - \operatorname{Tr} K, \qquad |t| < \|K\|^{-1}, \tag{B.11}$$

we have in finite-dimensional approximation

$$\varphi_N(t) = 2\arctan\Big(\partial_t \log D_N(t)\Big), \qquad D_N(t) := \det_2(I - tM_N), \tag{B.12}$$

which gives a natural phase extraction formula (including the $-\operatorname{Tr} M_N$ normalization). Near a pole $t = \kappa_n^{-1}$, φ_N jumps by π, so the phase correction is given by

$$\varphi_N^{\mathrm{unw}}(t) = \varphi_N(t) + \pi \sum_{\kappa_n^{-1} \le t} 1 \quad \text{(normalized by evenness and } \varphi_N^{\mathrm{unw}}(0) = 0) \tag{B.13}$$

For the AFE (two sums; Main Text *(110)*), according to the analytic conductor $C_\pi(t)$, balanced truncation lengths on each side are taken as

$$N_\pm(t) \asymp C_\pi(t)^{1/2} \tag{B.14}$$

and large imbalance causes significant cancellation in the difference of real and imaginary parts. In implementation, use Kahan summation or double precision support to ensure

$$\left|\mathrm{AFE}_{N_+,N_-}(t) - \mathrm{AFE}_{\infty,\infty}(t)\right| \ll N_+^{-m} + N_-^{-m} + e^{-c\eta} \tag{B.15}$$

(m is the smoothness order of the barrier, $c > 0$ is the band margin). Endpoint terms vanish for smooth barriers satisfying (B.10); if using hard barriers, apply the main text's estimates *(78)*, *(88)*.

Appendix B.7. Summary: Role of the Generating-Function Perspective

From the above, the generating-function perspective provides the following unifying principles:

- **Phase**: By monotonicity of the Cayley transform (Appendix A.5) and (B.12)–(B.13), read both zero constraints and operator spectra in the same phase.
- **Band**: By (B.6)–(B.7) and (B.9)–(B.10), commute truncation, smoothing, and endpoint correction (Appendix A.6–A.7).
- **Discretization**: Ensure numerical stability and uniqueness of unwrapping via (B.1)–(B.3) and the M-matrix structure (B.8).
- **AFE**: Control the truncation error of the two sums uniformly via the band parameter using (B.14)–(B.15).

Through this integration, the logical line in Section 8–Section 10 of the main text (outer factor = operator kernel = Weil form) is non-circularly connected via the two design variables of phase and band.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Remarks on Numerical Implementation (excerpt). When poles cluster in the observation window, use separate left/right grids and apply the correction (B.13) avoiding neighborhoods of poles (Main Text Section 8.6B). Strictly observe $\eta \le \log 2 - \delta$ for the band, and when $\delta \to 0$, use a smooth barrier to suppress endpoint terms (remark in Main Text Section 10.2). Automatically adjust AFE truncations according to (B.14), and use compensated summation to prevent loss of significance.

Appendix C. List of Symbols and Abbreviations

This appendix presents in one place the principal symbols, abbreviations, and conventions used in the main text and other appendices. Definitions, representative occurrences (sections in the main text), and minimal notes are given so that it can be referenced independently. The Fourier convention is the same as in the main text:

$$\widehat{f}(\xi) := \int_{\mathbb{R}} f(t)\, e^{-i\xi t}\, dt, \qquad f(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{f}(\xi)\, e^{i\xi t}\, d\xi,$$

Convolution is $(f * g)(t) = \int_{\mathbb{R}} f(t-s)g(s)\, ds$, and reflection is $\tilde{f}(t) = f(-t)$.

Legend (how to read and notational cautions)

- Representative sections are indicated in square brackets after the symbol (e.g., "[§7.2]").
- "a.e." means almost everywhere; "p.v." means principal value (Cauchy principal value).
- The origin convention for outer factors is $E(0) > 0$ (fixing the phase freedom); the phase is unwrapped as an even function.

Appendix C.1. Basic Sets, Function Spaces, and Transforms [7,16,29]

$\mathbb{R}$, $\mathcal{C}$, $\mathcal{Z}$, $\mathbb{N}$
: Field of real numbers, complex numbers, ring of integers, natural numbers.

$\mathcal{S}(\mathbb{R})$ Schwartz space (rapidly decreasing C^∞ functions).

$\mathcal{S}_{\text{even}}$ Even-function subspace: $\{f \in \mathcal{S}(\mathbb{R}) : f(t) = f(-t)\}$.

$L^p(\mathbb{R})$ Lebesgue-measurable function norm spaces ($1 \le p \le \infty$). Plancherel: $\|f\|_{L^2} = (2\pi)^{-1/2}\|\widehat{f}\|_{L^2}$.

PW_η Paley–Wiener space (band $\eta > 0$): L^2 functions with $\operatorname{supp} \widehat{f} \subset [-\eta, \eta]$.

A_η Band-limited, even test family: $A_\eta := \{\, \phi \in \mathcal{S}_{\text{even}} : \operatorname{supp} \widehat{\phi} \subset [-\eta, \eta] \,\}$ [Section 6,Section 8].

Hf Hilbert transform: $(Hf)(x) = \frac{1}{\pi}\,\text{p.v.}\int_{\mathbb{R}} \frac{f(t)}{x-t}\, dt$ [$H : L^2 \to L^2$ bounded].

$P_y(x)$ Poisson kernel: $P_y(x) = \frac{1}{\pi}\,\frac{y}{x^2+y^2}$ ($y > 0$).

$C\phi$ Cauchy transform: $C\phi(z) = \frac{1}{2\pi i}\int_{\mathbb{R}} \frac{\phi(x)}{x-z}\, dx$ (used with Plemelj's boundary value formula).

$\mathbf{1}_E$ Indicator function: $\mathbf{1}_E(x) = 1$ if $x \in E$, else 0.

Appendix C.2. Operators, Herglotz/Cayley, de Branges [2,42,59]

$L^{[d]}$ Direct sum of self-adjoint operators (degree d). [Section 8.1]

$K := \phi(L)$
: Bounded operator giving a positive-semidefinite kernel (ϕ in the admissible class of the main text). [§7.1–§7.3]

M_K Herglotz function ($\operatorname{Im} M_K > 0$ on $\mathcal{C}_+$): associated with the boundary value structure of K. [§7.1–§7.3]

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

S Schur function (Cayley transform):

$$S(t) = \frac{1 - iM_K(t)}{1 + iM_K(t)} = e^{-i\varphi(t)} \quad \text{(real-axis boundary value; phase } \varphi \text{ is even) [§7.2, Section 8.5].}$$

E de Branges function (HB class): $|E(z)| > |E^{\#}(z)|$ for $\mathbf{Im}\, z > 0$.

$E^{\#}(z)$ Schwarz reflection: $E^{\#}(z) := \overline{E(\bar{z})}$.

$\mathbf{det_2}(I - zK)$
Carleman–Fredholm second-regularized determinant. [§7.2]

$\tau_+(F)$ Upper half-plane exponential type: $\tau_+(F) = \limsup_{y \to +\infty} \frac{1}{y} \log |F(iy)|$. [§7.2]

Appendix C.3. L-Functions, Completions, and Phases [10,11,32]

$\Gamma_{\mathbb{R}}(s)$ Real gamma factor: $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(\frac{s}{2})$.

$\Gamma_{\mathcal{C}}(s)$ Complex gamma factor: $\Gamma_{\mathcal{C}}(s) = 2(2\pi)^{-s}\Gamma(s)$ (used as needed).

$\Lambda(s, \pi)$
Completed L-function: $\Lambda(s, \pi) = Q_\pi^{s/2} \prod_{j=1}^{d} \Gamma_{\mathbb{R}}(s + \mu_j)\, L(s, \pi)$ [Section 8].

Q_π Analytic conductor (positive real).

μ_j Archimedean parameters (usually assumed $\mathbf{Re}\, \mu_j > -1/2$).

ρ Nontrivial zero of $\Lambda(s, \pi)$ (symmetry $\rho \leftrightarrow 1 - \rho$).

$a_\pi(n)$ Dirichlet coefficients (coefficients of the Euler product).

$\Lambda_\pi(n)$ von Mangoldt-type coefficients: $-\frac{L'}{L}(s, \pi) = \sum_{n \geq 1} \frac{\Lambda_\pi(n)}{n^s}$.

$S_\pi(t)$ Schur function:

$$S_\pi(t) = \varepsilon(\pi)^{-1} \frac{\Lambda(\frac{1}{2} - it, \tilde{\pi})}{\Lambda(\frac{1}{2} + it, \pi)} = e^{-i\varphi_\pi(t)}.$$

$\varphi_\pi(t)$ Real-axis boundary phase (even). Derivative: $\varphi'_\pi(t) = 2\, \mathbf{Re}\, \frac{\Lambda'}{\Lambda}(\frac{1}{2} + it, \pi)$ [Section 8.5].

$\theta_\Gamma(t)$ Phase of gamma factor: $\theta_\Gamma(t) = \mathbf{Arg} \prod_j \Gamma_{\mathbb{R}}(\frac{1}{2} + \mu_j + it)$.

$\varepsilon(\pi)$ Root number ($|\varepsilon(\pi)| = 1$).

Appendix C.4. Prime Kernel, Band Tests, Normalization [1,7,30]

$P_{\eta,\pi}(t)$
Prime kernel (GL(d)):

$$P_{\eta,\pi}(t) := -2 \sum_p \sum_{k \leq e^\eta} (\log p)\, p^{-k/2}\, \mathbf{Re}\Big(\mathbf{tr}\, A_p(\pi)^k\Big)\, e^{itk \log p} \quad [\textit{Section } 8.2].$$

($\mathbf{tr}\, A_p(\pi)^k = \sum_{j=1}^{d} \alpha_{p,j}(\pi)^k$ is the kth power sum of Satake parameters.)

θ_π Ramanujan exponent (upper bound exponent for $|\alpha_{p,j}| \leq p^{\theta_\pi}$).

$\Phi_{T,\eta}^{\pm,(m)}$
Vaaler-type band tests: $\mathbf{supp}\, \widehat{\Phi} \subset [-\eta, \eta]$ bracketing $[-T, T]$ [Section 8.6].

$A\text{-}\mathbf{lim}$ Abel regularization limit (weighted limit as $\varepsilon \downarrow 0$) [Section 6.5,Section 8.2].

$E(0) > 0$
Origin normalization of outer factor (fixing phase freedom; consistent with $\varphi(0) = 0$).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix C.5. Distributions, Measures, and Fourier-Side Support [25,29,50]

$\mu_{L^{[d]}}$ Operator-side spectral measure (evenized).

μ_π Measure corresponding to zero distribution on the L-side (evenized).

Δ Difference distribution: $\Delta := \mu_{L^{[d]}} - \mu_\pi$.

$\operatorname{supp} \widehat{\Delta}$
Support on the Fourier side (small-band equalization $\Longleftrightarrow$ vanishing of low frequencies) [Section 8.7A].

Appendix C.6. Analytic Classes and Function-Theoretic Abbreviations [36,37,42]

HB class
Hermite–Biehler class: class of entire functions satisfying $|E(z)| > |E^{\#}(z)|$ for $\operatorname{Im} z > 0$.

Cartwright class
Entire functions with real-axis consistency, bounded type, and finite type.

outer/inner
Outer factor / inner factor (log-integrable boundary values / Blaschke factor).

bounded type
Functions having a harmonic majorant of $\log^+ |f|$ in the upper half-plane.

a.e., a.s.
almost everywhere / almost surely.

Appendix C.7. Numerical and Algorithmic Abbreviations [58,60]

AFE Approximate Functional Equation (see equation (110) in the main text) [Section 8.6].

FFT Fast Fourier Transform.

SDP Semidefinite Programming (used to bound Weil-type quadratic forms) [Section 8.7C].

Nyström method
Quadrature-point discretization of integral kernels (positive weights and nodes) [Appendix B].

Galerkin
Projection approximation on an adapted basis.

Kahan sum
Compensated summation (loss-of-significance suppression; recommended for the two sums in AFE).

Appendix C.8. Conventions for Parameters and Scalars

d Degree of **GL** (number of direct-sum components).

η Bandwidth (small-band means $\eta < \log 2$).

T Observation window radius (for phase integration/visualization).

y Radius of Poisson smoothing (P_y).

Q_π, μ_j, θ_π
Analytic conductor, Archimedean parameters, Ramanujan exponent.

Appendix C.9. Notation Conventions (Dominance Symbols, Uniformity, Boundary Values) [29,43,48]

$O(\cdot)$, $\ll$
$A = O(B)$ or $A \ll B$ means $|A| \le C\,B$, where C is an absolute constant depending on the context.

$O_\alpha(\cdot)$ Explicitly allows constants uniform in parameter α (e.g., $O_{\eta,m}(1)$).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Boundary value

$F(x \pm i0)$ denotes the nontangential boundary value to the real axis, with real/imaginary parts described by the Plemelj formula.

Appendix C.10. Reference Formulas (Poisson–Hilbert, Phase Derivative, Determinant Identification) [6,26,41]

We restate here fundamental formulas that frequently appear in this paper (derivations in Appendix A or relevant sections in the main text):

$$\text{(PH)} \quad \mathbf{Re}\log E(iy) = \frac{1}{2\pi}\int_{\mathbb{R}} \phi(x)\,\frac{y}{x^2+y^2}\,dx, \qquad (y>0) \quad [\S 7.2], \tag{A1}$$

$$\text{(Phase)} \quad \varphi'_\pi(t) = 2\ \mathbf{Re}\,\frac{\Lambda'}{\Lambda}\left(\tfrac{1}{2}+it,\pi\right) \quad [\textit{Section } 8.5], \tag{A2}$$

$$(\det_2) \quad \partial_t \log \det{}_2(I-tK) = \sum_n \frac{\kappa_n}{1-t\kappa_n} - \mathrm{Tr}\,K \quad (|t|<\|K\|^{-1}) \quad [\S 7.2]. \tag{A3}$$

Usage Notes

(1) The Fourier convention is the same as in the main text, and the Poisson–Hilbert formula (A1), phase derivative (A2), and determinant identification (A3) are all consistent with this convention.

(2) The phase φ is even, and is monotonically extended by origin normalization $\varphi(0) = 0$ and unwrapping (with π jumps at poles).

(3) In principle, the band is $\eta < \log 2$, with endpoint contributions vanishing for smooth windows, and controlled by the evaluation formulas (Main Text Section 10.2) for hard windows.

Appendix D. List of Assumptions and Normalizations (Audit Ledger)

This appendix compiles in one place the conventions, assumptions, and normalizations used throughout the entire paper, edited in an *auditable* form so that one can cross-reference which results depend on which premises. Symbols and references all conform to the section and equation numbering in the main text. Focusing on the Fourier convention, boundary value conventions, outer factors (HB class) and phase conventions for Schur functions, small-band equalization, and Abel regularization, we explicitly state the *consistency* between the operator side and the L-function side (completed form).

Appendix D.1. Global Conventions (Fourier, Phase, Boundary Values) [41,44,52,61,62]

- **Fourier convention** (Main Text Section 6):

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(t)e^{-i\xi t}\,dt, \qquad f(t) = \frac{1}{2\pi}\int_{\mathbb{R}} \widehat{f}(\xi)e^{i\xi t}\,d\xi.$$

Even-function test space $\mathcal{S}_{\mathbf{even}} = \{f \in \mathcal{S}(\mathbb{R}) : f(t) = f(-t)\}$, band-limited, even test family

$$A_\eta := \{\phi \in \mathcal{S}_{\mathbf{even}} : \ \mathbf{supp}\,\widehat{\phi} \subset [-\eta,\eta]\}, \qquad 0 < \eta < \log 2.$$

- **Hilbert transform and Poisson kernel** (Main Text Section 7.1):

$$(Hf)(x) = \frac{1}{\pi}\,\mathrm{p.v.}\int_{\mathbb{R}} \frac{f(t)}{x-t}\,dt, \qquad P_y(x) = \frac{1}{\pi}\frac{y}{x^2+y^2}\ (y>0).$$

- **Cauchy transform and Plemelj**: For $C\phi(z) = \frac{1}{2\pi i}\int_{\mathbb{R}} \frac{\phi(x)}{x-z}\,dx$, $C\phi(x \pm i0) = \frac{1}{2}(H\phi)(x) \mp \frac{i}{2}\phi(x)$ holds a.e.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

- **Cayley transform and phase** (Main Text Section 7.2): For a Herglotz function M,

$$S(t) = \frac{1 - iM(t)}{1 + iM(t)} = e^{-i\varphi(t)} \quad \text{(Schur function)}.$$

  The phase φ is even, with origin convention $\varphi(0) = 0$, and is uniquely determined by continuous connection (unwrapping) including π-jumps at poles.
- **Boundary value convention**: $F(x + i0)$ denotes the nontangential limit from the upper half-plane. $|S(x + i0)| = 1$ holds.

Appendix D.2. Normalization of Operators, Kernels, and de Branges Functions [2,6,41,59]

- **Basic operator** (Main Text Section 8.1): L is self-adjoint, $L^{[d]} := L^{\oplus d}$ is its direct sum.
- **Kernel construction** (Main Text Section 7.1–Section 7.3): For even, real-valued ϕ, set $K := \phi(L)$. When necessary, assume $K \geq 0$ (Hilbert–Schmidt).
- **Herglotz function and Schur function**: $M_K(z) = \mathrm{Tr}\big((I - zK)^{-1}K\big)$ satisfies $\mathrm{Im}\, M_K > 0$ on $\mathcal{C}_+$. $S_K = (1 - iM_K)/(1 + iM_K) = e^{-i\varphi_K}$.
- **Second regularized determinant** (Main Text Section 7.2):

$$\det{}_2(I - zK) = \det\big((I - zK)\, e^{zK}\big), \qquad \partial_z \log \det{}_2(I - zK) = \sum_n \frac{\kappa_n}{1 - z\kappa_n} - \mathrm{Tr}\, K.$$

- **de Branges (HB class) and origin normalization** (Main Text Section 7.5): The outer factor E_K is HB class with the convention $E_K(0) > 0$. The upper half-plane exponential type is $\tau_+(E_K) := \limsup_{y\to\infty} y^{-1} \log |E_K(iy)|$.

Appendix D.3. L-Function Side Conventions and Completed Form [10,11,32,35]

- **Completed form and functional equation** (Main Text Section 10.2): With Archimedean factors $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(\frac{s}{2})$, and if needed $\Gamma_{\mathcal{C}}(s) = 2(2\pi)^{-s}\Gamma(s)$,

$$\Lambda(s, \pi) = Q(\pi)^{s/2} \prod_{j=1}^{d_R} \Gamma_{\mathbb{R}}(s + \mu_j) \prod_{k=1}^{d_C} \Gamma_{\mathcal{C}}(s + \nu_k)\, L(s, \pi),$$

  $\Lambda(s, \pi) = \varepsilon(\pi)\, \Lambda(1 - s, \pi)$, $|\varepsilon(\pi)| = 1$.
- **Schur function and phase** (Main Text Section 8.5):

$$S_\pi(t) = \varepsilon(\pi)^{-1} \frac{\Lambda(\frac{1}{2} - it, \tilde{\pi})}{\Lambda(\frac{1}{2} + it, \pi)} = e^{-i\varphi_\pi(t)}, \quad \varphi_\pi'(t) = 2\,\mathrm{Re}\, \frac{\Lambda'}{\Lambda}(\tfrac{1}{2} + it, \pi).$$

- **Archimedean calibration** (calibration formula in Main Text Section 6.2; e.g., around equation (26)):

$$\mathrm{Re}\, \frac{\Gamma_R'}{\Gamma_R}(\tfrac{1}{2} + it) = \log\Big(\tfrac{|t|}{2\pi}\Big) + O\big((1 + t^2)^{-1}\big),$$

  hence $\varphi_\pi'(t) = d \log |t| - d \log(2\pi) + \log Q(\pi) + O\big((1 + t^2)^{-1}\big)$ $(d = d_R + 2d_C)$.
- **Origin normalization**: Outer factor E_π satisfies $E_\pi(0) > 0$.

Appendix D.4. Small-Band Equalization, Abel Regularization, Low-Frequency Vanishing [7,25,45,49,63]

- **Small bandwidth**: Fix $\eta_0 \in (0, \log 2)$, and henceforth $0 < \eta < \eta_0$.
- **Main statement of equalization** (Main Text Section 6.1–Section 6.3): For the evenized difference distribution

$$\Delta := \mu_{L^{[d]}} - \mu_{\Xi_\pi}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

we have $\langle \Delta, \phi \rangle = 0$ for all $\phi \in A_\eta$.

- **Low-frequency vanishing** (Main Text Section 8.7A): $\langle \Delta, \phi \rangle = 0$ $(\forall \phi \in A_\eta)$ is equivalent to $\operatorname{supp} \widehat{\Delta} \subset \mathbb{R} \setminus (-\eta_0, \eta_0)$.
- **Abel regularization** (Main Text Section 6.5, Section 8.2): $A\text{-}\lim_{\varepsilon \downarrow 0} \langle e^{-\varepsilon|t|} \Delta, \phi \rangle = \langle \Delta, \phi \rangle$. Endpoint terms are absorbed into $O(1)$ (see Appendix A).

Appendix D.5. Assumptions for the Main Theorem (GL(d)) (H1′–H3′) [41,44,51]

(H1′) **Positivity and HB positivity**: Choose ϕ even and real so that $K^{[d]} = \phi(L^{[d]}) \geq 0$ (Hilbert–Schmidt). Associated $E_{K^{[d]}}$ is HB class (Main Text Section 7.5).

(H2′) **Small-band equalization**: For $\eta < \eta_0 < \log 2$, $\langle \mu_{L^{[d]}} - \mu_{\Xi_\pi}, \phi \rangle = 0$ (Main Text Section 6.1–Section 6.3).

(H3′) **Origin normalization**: $E_{K^{[d]}}(0) > 0$ and $E_\pi(0) > 0$ (consistent with phase origin convention).

Conclusion (Main Theorem group in Section 8):

(H1′)–(H3′) $\Longrightarrow$ Phase equality $\Longrightarrow$ Type equality $\Longrightarrow$ Outer factor identification $\Longrightarrow$ Critical line constraint.

Phase equality $\Rightarrow$ type equality follows from the Poisson–Hilbert representation (Main Text Section 7.2). Outer factor identification follows from the Cartwright/inner–outer decomposition and origin convention (Main Text Section 7.3).

Appendix D.6. Cross-Reference Table (Theorem/Proposition ↔ Assumptions)

Result in Main Text	Directly Used Assumptions/Conventions	Auxiliary References (Main Text)
Phase equality (Section 8: Theorem)	(H2′), Abel regularization, $\eta < \log 2$	Section 6.5 (regularization), Section 7.2 (PH representation)
Type equality (Section 8: Theorem)	Phase equality, PH representation	Section 7.2 (Poisson–Hilbert)
Outer factor identification (Section 8: Theorem)	Phase/Type equality, (H3′)	Section 7.3 (outer factor identification lemma)
GRH-type constraint (Section 8: Corollary)	(H1′), outer factor identification	Section 7.5 (phase monotonicity)
RvM main term (Section 6: Proposition)	Archimedean calibration	Section 6.2 (calibration formula), Section 5.1 (main representation)
Prime kernel bound (Section 10: Proposition)	Upper bound on Ramanujan exponent	Section 10.2 (endpoint term handling)

Appendix D.7. Audit Checklist (Theory and Implementation)

Theoretical check (paper verification)

- Fourier convention and signs match the main text (ϕ is even).
- Adoption of $\eta_0 < \log 2$ pushes prime terms outside the band (Main Text Section 6.1–Section 6.3).
- Small-band equalization (H2′) holds for all A_η (check coverage of applicable range).
- Origin normalization (H3′): $E_{K^{[d]}}(0) > 0$, $E_\pi(0) > 0$ consistently set.
- HB positivity (H1′): ϕ constructed as approximation (or limit) of a completely monotone family (Main Text Section 7.5).
- PH representation yields phase average $\int_{-T}^{T} (\phi_K - \phi_\pi)\, dt = O(1)$ (Main Text Section 7.2).

Numerical check (implementation; per Appendix B)

- Nyström convergence: existence of N satisfying $\sup_{t \in I} |\varphi^{(N)}(t) - \varphi^{(N/2)}(t)| < \varepsilon$.
- AFE internal error: difference for two auxiliary functions G is below threshold (Main Text Section 8.6).
- Phase monotonicity: $\varphi'(t) > 0$ except at poles, π-jumps at poles; consistency of unwrapping (Main Text Section 7.5).
- Band projection residual: bound $|R_{T,\eta}| \lesssim \eta^{-m} + e^{-c\eta}$ (Main Text Section 8.6).
- Endpoint terms: vanish for smooth barriers, explicit correction for hard barriers (remark in Main Text Section 10.2).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix D.8. Table of Typical Normalizations [35]

Object	Normalization Content	Reference
Outer factor E_K, E_π	$E(0) > 0$ (origin)	Section 7.2, Section 8.3
Phase φ_K, φ_π	$\varphi(0) = 0$, even, continuous connection (unwrapping)	Section 7.2
Schur function S	$\lvert S\rvert = 1$ (calibrate by $\varepsilon(\pi)$ if needed)	Section 8.5
Fourier	$\widehat{f}(\xi) = \int f(t)e^{-i\xi t}\,dt$ (inverse transform also given)	Section 6
Band	$0 < \eta < \eta_0 < \log 2$ (small band)	Section 6.1–Section 6.3
Second determinant	$\partial_t \log \det_2(I - tK) = \sum \frac{\kappa_n}{1-t\kappa_n} - \operatorname{Tr} K$	Section 7.2

Appendix D.9. Acceptable Range of Variants (Interchangeable Conventions)

- **Fourier sign**: Using $\widehat{f}(\xi) = \int f(t)e^{+i\xi t}\,dt$ is equivalent if one reverses the overall sign consistently (explicit formula, phase direction), absorbable into the main text's notation.
- **Change of reference point**: Normalization $E(z_0) > 0$ with $z_0 \in i\mathbb{R}_{>0}$ is also possible. Adjust the constant term in the PH representation to match (Main Text Section 7.2).
- **Treatment of band endpoint**: For $\eta = \log 2$, prime-term endpoint contributions appear; for auditing, always take $\eta < \log 2$ (Main Text Section 6.3, Section 10.2).

Summary: We have listed (H1′)–(H3′), the Fourier/phase/boundary value conventions, small-band equalization, and Abel regularization in an integrated way, and organized in Table D.6 the premises and auxiliary references on which each main result depends. With this ledger, theory (on paper) and numerics (implementation) can be independently reproduced and audited.

Appendix E. Typos, Notational Inconsistencies, and Editorial Notes (Errata Candidates)

This appendix compiles *possible typos, notational inconsistencies, and convention discrepancies* for the main text (Section 1–Section 10) and Appendices A–D, indicating the correct formula, recommended unified notation, and justification. References conform to the sections and equation numbers in the main text.

Appendix E.1. Overview: Locations Sensitive to Convention Differences [2,4,6,35,44]

- **Direction of Cayley transform and sign of phase** (Main Text Section 7.2):

$$S(t) = \frac{1 - iM(t)}{1 + iM(t)} = e^{-i\varphi(t)} \quad \text{(unified)}$$

  (On the real axis, $\lvert S\rvert = 1$, φ is even with origin convention $\varphi(0) = 0$).
- **Form of phase derivative** (Appendix A.5):

$$\varphi'(t) = \frac{2M'(t)}{1 + M(t)^2} > 0 \quad \text{(a.e. except at poles)}$$

  (unified).
- **Asymptotic main term of $\Gamma_{\mathbb{R}}$** (Main Text Section 6.2):

$$\operatorname{Re}\Big(\frac{\Gamma_{\mathbb{R}}'}{\Gamma_{\mathbb{R}}}\Big)\big(\tfrac{1}{2} + \mu + it\big) = \log\Big(\frac{|t|}{2\pi}\Big) + O\big((1+t^2)^{-1}\big)\,.$$

- **Constant in Riemann–von Mangoldt main term** (Main Text Section 6.2, Section 6.5): Pay attention to the position of $-d$ and $\log(2\pi)$ in the main term of $N_\pi(T)$.
- **Differential identity for $\det_2$** (Main Text Section 7.2): Always include the correction term $-\operatorname{Tr} K$ (equation (A9)).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

- **Treatment of small-band endpoint** (Main Text Section 6.1–Section 6.3): Impose $\eta_0 < \log 2$ *strictly*.
- **Conversion between one-sided and two-sided counts** (Main Text Section 8.5): $N_\pi(T)$ is one-sided. Pay attention to the conversion factor **1/2** for total two-sided sums.
- **Origin normalization** (Main Text Section 7.2, Section 8.3): Always state $E(0) > 0$ together with $\varphi(0) = 0$.

Appendix E.2. Equation-Level Correction Candidates (with Justification) [1,2,4,6,32,35,44]

(E2-1) Sign of Cayley transform

The correct convention in the main text is

$$S_K(t) = \frac{1 - i\,\mathcal{M}_K(t)}{1 + i\,\mathcal{M}_K(t)} = e^{-i\phi_K(t)} \qquad \text{(a.e. } t \in \mathbb{R}\text{).} \tag{A4}$$

Occurrences of the form $S_K = \frac{1+i\mathcal{M}_K}{1-i\mathcal{M}_K} = e^{+i\phi_K}$ should be unified to (A4) (adjust $\phi_K \mapsto -\phi_K$ accordingly).

Justification: $M = \mathcal{M}_K$ is Herglotz ($\operatorname{Im} M > 0$ on $\mathcal{C}_+$). The Cayley transform $S = (1 - iM)/(1 + iM)$ maps to the unit disk and satisfies $|S| = 1$ at the boundary. Thus $S = e^{-i\phi}$ with ϕ real (even).

(E2-2) Formula for phase derivative

The correct formula is

$$\phi_K'(t) = \frac{2\,\mathcal{M}_K'(t)}{1 + \mathcal{M}_K(t)^2}, \qquad \mathcal{M}_K'(t) = \sum_n \frac{\kappa_n^2}{(1 - t\,\kappa_n)^2} \geq 0. \tag{A5}$$

Misprints with denominator $1 - \mathcal{M}_K(t)^2$ should be corrected to (A5).

Derivation: From $S = (1 - iM)/(1 + iM) = e^{-i\phi}$,

$$\frac{S'}{S} = \frac{-iM'}{1 - iM} - \frac{iM'}{1 + iM} = -i\,\frac{2M'}{1 + M^2} = -i\,\phi',$$

hence $\phi' = 2M'/(1 + M^2)$. Since $K \geq 0$ and self-adjoint, $\mathcal{M}_K(t) = \operatorname{Tr}\big((I - tK)^{-1}K\big)$ and $\mathcal{M}_K'(t) = \operatorname{Tr}\big((I - tK)^{-1}K(I - tK)^{-1}K\big) \geq 0$, with the RHS sum given by the eigenvalue expansion.

(E2-3) Asymptotic of $\Gamma_{\mathbb{R}}$

For $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma\big(\frac{s}{2}\big)$,

$$\operatorname{Re}\frac{\Gamma_{\mathbb{R}}'}{\Gamma_{\mathbb{R}}}\big(\tfrac{1}{2} + \mu + it\big) = \log\Big(\frac{|t|}{2\pi}\Big) + O\Big(\frac{1}{1 + t^2}\Big) \qquad (\mu \text{ fixed}). \tag{A6}$$

Derivation: Apply Stirling's formula $\log\Gamma(z) = (z - \frac{1}{2})\log z - z + O(1)$ ($|\arg z| \leq \pi - \delta$) to $z = \frac{1}{2}(\frac{1}{2} + \mu + it)$, take the real part to get $\operatorname{Re}(\Gamma'/\Gamma)(z) = \log|z| + O(|z|^{-1})$. Since $\log|z| = \log\Big(\frac{|t|}{2}\Big) + O((1 + t^2)^{-1})$ and $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$, we obtain (A6).

(E2-4) Constant in Riemann–von Mangoldt main term

For the general $\mathrm{GL}(d)$ completed form, the main term of zero count $N_\pi(T)$ is

$$N_\pi(T) = \frac{T}{2\pi}\Big(d\log T - d\log(2\pi) + \log Q_\pi - d\Big) + O(\log T). \tag{A7}$$

Places with missing $-d$ or incorrect $\log(2\pi)$ should match (A7).

Outline of derivation: Localizing the explicit formula with even tests, the gamma factor contribution from (A6) integrates to $\frac{T}{2\pi}(d\log T - d\log(2\pi)) - \frac{d}{2\pi}T$. The conductor Q_π gives $\frac{T}{2\pi}\log Q_\pi$, and the prime term contributes nothing in the small-band setting (Main Text Section 6.2, Section 6.5).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(E2-5) Phase of Schur function

$$S_\pi(t) = \varepsilon(\pi)^{-1} \frac{\Lambda(\frac{1}{2} - it, \tilde{\pi})}{\Lambda(\frac{1}{2} + it, \pi)} = e^{-i\phi_\pi(t)}. \tag{A8}$$

Occurrences with RHS $e^{+i\phi_\pi}$ should be corrected to (A8). ϕ_π is even, normalized by $\phi_\pi(0) = 0$.

(E2-6) Differential of regularized determinant and phase

For the second regularized determinant,

$$\partial_t \log \det_2(I - tK) = \operatorname{Tr}\Big(\frac{K}{I - tK}\Big) - \operatorname{Tr} K = \sum_n \frac{\kappa_n}{1 - t\kappa_n} - \sum_n \kappa_n, \tag{A9}$$

where $\{\kappa_n\}$ are the eigenvalues of K (with multiplicity). Forms missing the $-\operatorname{Tr} K$ term should be corrected to (A9).
Derivation: From $\det_2(I - tK) = \det\big((I - tK)\, e^{tK}\big)$, $\partial_t \log \det_2 = \operatorname{Tr}\big(-(I - tK)^{-1}K + K\big) = \operatorname{Tr}\big(K(I - tK)^{-1}\big) - \operatorname{Tr} K$. Justify via finite-rank approximation and take limits (Main Text Section 7.2).

(E2-7) Treatment of small-band endpoint

$\eta_0 < \log 2$ is required *strictly*. If $\eta_0 = \log 2$ appears, replace by

$$\eta_0 = \log 2 - \delta \qquad (\delta > 0)$$

and absorb endpoint contributions by Abel regularization or a smooth window (Main Text Section 6.1–Section 6.3, Section 10.2).

(E2-8) Conversion between one-sided and two-sided counts

$N_\pi(T)$ counts one-sided $\gamma \in (0, T]$. For places using the two-sided sum $\sum_\rho \varphi(\operatorname{Im}\rho)$ (with $\varphi \approx \mathbf{1}_{[-T,T]}$),

$$\sum_{0<\gamma\le T} 1 = \tfrac{1}{2} \sum_{|\gamma|\le T} 1 + O(1)$$

should be made explicit (Main Text Section 8.5).

Appendix E.3. Rules for Unifying Notation [32,35]

- **Schwarz reflection**: unify as $f^\sharp(z) := \overline{f(\bar{z})}$ (do not mix with $f^\#$).
- **Direct sum notation**: $L^{[d]} := \underbrace{L \oplus \cdots \oplus L}_{d \text{ times}}$.
- **Conductor notation**: analytic conductor is Q_π; $C_\pi(t)$ is reserved for the *t-dependent version* of the analytic conductor (AFE truncation length).
- **Phase origin normalization**: always state $E(0) > 0$ and $\phi(0) = 0$ together (avoid stating only one).
- **Fourier convention**: unify as $\widehat{f}(\xi) = \int_{\mathbb{R}} f(t)\, e^{-i\xi t}\, dt$; when quoting other conventions, note the conversion.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix E.4. Correction of Typical Confusions: Checklist [1,4,64]

Item	Common Mistake	Correct Form / Justification
Fourier convention	$\widehat{f}(\xi) = \int f e^{+i\xi t} dt$	$\widehat{f}(\xi) = \int f e^{-i\xi t} dt$ (Main Text Section 6)
Cayley sign	$S = \frac{1+iM}{1-iM} = e^{+i\varphi}$	$S = \frac{1-iM}{1+iM} = e^{-i\varphi}$ ((A4))
Phase derivative	$\varphi' = \frac{2M'}{1-M^2}$	$\varphi' = \frac{2M'}{1+M^2} > 0$ ((A5))
$\mathbf{\Gamma}_{\mathbb{R}}$ asymptotic	$\operatorname{Re}\psi_{\mathbb{R}}(\frac{1}{2}+\mu+it) = \log\lvert t\rvert - \log\pi$	$\log\frac{\lvert t\rvert}{2\pi}$ ((A6))
RvM main term	$\frac{T}{2\pi}(d\log T + \log Q_\pi) - \cdots$	(A7) (Main Text Section 6.2, Section 6.5)
$\mathbf{det_2}$ differential	$\partial_t \log \det_2 = \sum \frac{\kappa}{1-t\kappa}$	(A9) (include $-\operatorname{Tr} K$)
Band endpoint	$\eta_0 = \log 2$	$\eta_0 < \log 2$ (Main Text Section 6.1–Section 6.3)
Count conversion	Missing $1/2$ factor	State factor $1/2$ explicitly (Main Text Section 8.5)
Origin convention	Only $E(0) > 0$ stated	State both $E(0) > 0$ and $\varphi(0) = 0$ (Main Text Section 7.2, Section 8.3)

Appendix E.5. Editorial Notes (Pitfalls in Implementation/Review) [4,6,30,32,35]

- **Phase *unwrapping***: In figures and tables, always use the continuous phase, absorbing principal value jumps ($\pm\pi$) by $\pm 2\pi$ correction (Main Text Section 7.2, Appendix A.5).
- **AFE left/right balance** (Main Text Section 8.6): Choose truncation length based on $N \asymp C_\pi(t)^{1/2}$ to avoid loss of significance due to imbalance.
- **Management of endpoint contributions** (Main Text Section 6.5, Section 10.2): For hard windows, residual endpoint terms remain; use Abel regularization or smooth windows with high-order vanishing.
- **$\det_2$ and cyclic products** (Main Text Section 4.3, Section 7.2): Justify exchange of traces/localization under boundedness/integrable kernel assumptions, and do not omit $-\operatorname{Tr} K$.
- **Thorough origin normalization**: Set $E(0) > 0$ and $\varphi(0) = 0$ together, and verify repeatedly in the text.

Summary: We have listed potentially *critical* typo candidates for Cayley transform, phase derivative, $\mathbf{\Gamma}_{\mathbb{R}}$ asymptotic, RvM main term, $\det_2$ differential, band endpoint, and count conversion, and indicated the correct forms. Following this table for a comprehensive check and correction of the manuscript ensures stable consistency between theory and numerics.

Appendix F. Taxonomy of Alternative Kernels and Positivity (Recipe for Completely Monotone Families)

This appendix presents a practical recipe for systematically designing the operator-side kernel $K = \phi(L)$ from *completely monotone* (CM) families and their extensions. We organize as follows: (1) *Positivity* based on CM functions and Hilbert–Schmidt (HS)/Schatten class criteria, (2) the influence on phase/type and *design guidelines* (monotonicity of Cayley phase; Main Text Section 7.2), (3) *parameterization and differentiation* suitable for numerical implementation (optimization and sensitivity analysis). We use the same Fourier convention as in the main text, $\widehat{f}(\xi) = \int_{\mathbb{R}} f(t) e^{-i\xi t}\, dt$, and assume L is self-adjoint and nonnegative ($L \geq 0$).

Appendix F.1. Completely Monotone Functions and Positivity: Bernstein–Bochner Viewpoint [38,54]

Definition A2 (Completely Monotone (CM)). *A C^∞ function ϕ on $[0,\infty)$ is* completely monotone *if, for all $k \in \mathbb{N} \cup \{0\}$ and $\lambda > 0$,*

$$(-1)^k \phi^{(k)}(\lambda) \geq 0.$$

Theorem A2 (Bernstein Representation). *ϕ is CM if and only if there exists a positive (possibly σ-finite) Borel measure ν such that*

$$\phi(\lambda) = \int_0^\infty e^{-t\lambda}\, \nu(dt) \qquad (\lambda \geq 0). \tag{A10}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Pushing (A10) through the spectral theorem $L = \int_{[0,\infty)} \lambda \, dE_L(\lambda)$ gives

$$\phi(L) = \int_0^\infty e^{-tL} \, \nu(dt) \qquad \text{(strong limit; } e^{-tL} \geq 0\text{).} \tag{A11}$$

Thus, if ϕ is CM, then $K = \phi(L)$ is automatically $K \geq 0$, directly connecting to the Herglotz/Cayley framework in Main Text Section 7.1–Section 7.2. In general, $\phi \geq 0$ (even if not CM) still gives $\phi(L) \geq 0$, but CM property is *advantageous in design* since it guarantees monotonicity, an integral representation, and a stable differentiation structure.

Remark A3 (Monotonicity and Tail Control)**.** *For CM families,* $\phi(0) = \nu([0,\infty)) \geq 0$, $\phi'(\lambda) \leq 0$ *automatically, and the eigenvalue sequence* $\kappa_n = \phi(\lambda_n)$ *of* $K = \phi(L)$ *has a monotonically decaying tail. The monotonicity of Cayley phase* φ *(Appendix A.5; Main Text Section 7.2) depends on* $K \geq 0$*, so CM design ensures it is satisfied.*

Appendix F.2. Spectral Growth and Schatten Classes (Including HS/Trace) [6,14]

Assume a rough Weyl-type bound

$$N_L(\Lambda) := \operatorname{Tr} E_L([0,\Lambda]) \ll \Lambda^\alpha \qquad (\Lambda \to \infty, \ \alpha > 0) \tag{A12}$$

(Main Text Section 6.2 calibration). Let $\mathbf{S}_p$ denote the Schatten class ($p \in [1,\infty)$).

Proposition A3 (Sufficient Condition for Schatten Membership (Integral Test))**.** *Let* $\phi : [0,\infty) \to [0,\infty)$ *be nonincreasing. For any* $p \in [1,\infty)$*,*

$$\int_1^\infty \phi(\lambda)^p \, dN_L(\lambda) \asymp \sum_{\lambda \in \sigma(L)} \phi(\lambda)^p,$$

and in particular,

$$\phi(\lambda) \ll (1+\lambda)^{-s} \ (\lambda \to \infty) \implies \phi(L) \in \mathbf{S}_p \ \text{if} \ sp > \alpha. \tag{A13}$$

At the boundary $sp = \alpha$, $\phi(\lambda) \ll (1+\lambda)^{-\alpha/p} (\log(2+\lambda))^{-1-\varepsilon}$ *ensures* $\phi(L) \in \mathbf{S}_p$ *for any* $\varepsilon > 0$*.*

Proof. By partial integration and (A12), $\sum \phi(\lambda)^p \asymp \int_1^\infty \phi(\lambda)^p \, dN_L(\lambda) \ll \int_1^\infty \lambda^{\alpha-1} (1+\lambda)^{-sp} \, d\lambda$. The convergence condition is $sp > \alpha$; at the boundary, assuming logarithmic decay gives convergence. □

Important special cases.

For $p = 2$ (HS) require $s > \alpha/2$; for $p = 1$ (Trace) require $s > \alpha$. Example: $\phi(\lambda) = (c+\lambda)^{-\sigma}$ gives "HS: $\sigma > \alpha/2$, Trace: $\sigma > \alpha$."

Appendix F.3. CBF Composition and Representative Examples (Matérn/Rational/ Heat-type) [65–67]

Definition A3 (Bernstein Function (BF), Complete Bernstein Function (CBF))**.** *A function* $g : (0,\infty) \to (0,\infty)$ *is a* Bernstein function *if* $g \in C^\infty$ *and* g' *is CM. In this case,*

$$g(\lambda) = a + b\lambda + \int_{(0,\infty)} \frac{\lambda}{\lambda + t} \, \mu(dt) = a + b\lambda + \int_{(0,\infty)} (1 - e^{-t\lambda}) \, \frac{\mu(dt)}{t}.$$

Furthermore, g *is CBF if it admits the Stieltjes representation* $g(\lambda) = c + d\lambda + \int_{(0,\infty)} \frac{\lambda}{\lambda + t} \, \sigma(dt)$ *(*$c, d \geq 0$*).*

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Proposition A4 (Composition Closure of CM with BF). *If ϕ is CM and g is BF, then*

$$\phi \circ g \text{ is CM.}$$

Proof. If $\phi(\lambda) = \int_0^\infty e^{-s\lambda}\, \nu(ds)$ (Bernstein representation) and g is BF, then for fixed $s > 0$, $e^{-sg(\lambda)}$ is CM in λ (Bochner subordination). Hence $\phi(g(\lambda)) = \int e^{-sg(\lambda)}\, \nu(ds)$ is CM. □

Representative examples and CM property.

- **Rational type**: $\phi(\lambda) = (c+\lambda)^{-\sigma}$ $(c > 0,\ \sigma > 0)$ is CM.
- **Exponential type**: $\phi(\lambda) = e^{-a\lambda}$ $(a > 0)$ is CM.
- **Matérn (first order)**: $\phi(\lambda) = (1+\alpha\lambda)^{-\nu}$ $(\alpha, \nu > 0)$ is CM (take $g(\lambda) = \alpha\lambda$ as BF in Proposition A4).
- **Heat-type (squared semigroup)**: $\phi(\lambda) = \int_0^\infty e^{-t\lambda^2}\, w(dt)$ is also CM as an average of e^{-tL^2} $(L^2 \geq 0)$, yielding $K = \phi(L) = \int e^{-tL^2} w(dt) \geq 0$. In this framework, $\phi(\lambda) = (1+\alpha\lambda^2)^{-\nu}$ can be represented as

$$(1+\alpha\lambda^2)^{-\nu} = \frac{1}{\Gamma(\nu)} \int_0^\infty t^{\nu-1} e^{-t}\, e^{-\alpha t\, \lambda^2}\, dt.$$

Corollary A2 (HS/Trace Thresholds (Representative Families)).

1. ***Rational type*** $\phi(\lambda) = (c+\lambda)^{-\sigma}$*: HS if* $\sigma > \alpha/2$*, Trace if* $\sigma > \alpha$*.*
2. ***Matérn (first order)*** $\phi(\lambda) = (1+\alpha\lambda)^{-\nu}$*: HS if* $\nu > \alpha/2$*, Trace if* $\nu > \alpha$*.*
3. ***Heat-type Matérn*** $\phi(\lambda) = (1+\alpha\lambda^2)^{-\nu}$*: since* $\phi(\lambda) \asymp \lambda^{-2\nu}$*, HS if* $\nu > \alpha/4$*, Trace if* $\nu > \alpha/2$*.*

Appendix F.4. Hilbert–Schmidt Norm and Phase: Expansion Near the Origin [6,15]

For the Cayley phase in Main Text Section 7.2,

$$S(t) = \frac{1 - iM(t)}{1 + iM(t)} = e^{-i\varphi(t)}, \qquad M(t) = \operatorname{Tr}\big((I - tK)^{-1}K\big),$$

we have the formal power series

$$(I - tK)^{-1}K = \sum_{m\geq 0} t^m K^{m+1} \quad \Longrightarrow \quad M(t) = \sum_{m\geq 0} t^m \operatorname{Tr} K^{m+1} \tag{A14}$$

converging for $|t| < \|K\|^{-1}$. Thus,

$$\varphi'(t) = \frac{2M'(t)}{1 + M(t)^2}, \qquad \varphi''(t) = \frac{2\big(M''(t)(1+M^2) - 2M(M')^2\big)}{(1+M^2)^2}. \tag{A15}$$

In particular, at the origin,

$$M(0) = \operatorname{Tr} K, \quad M'(0) = \operatorname{Tr} K^2, \quad M''(0) = 2!\, \operatorname{Tr} K^3, \tag{A16}$$

hence

$$\varphi'(0) = \frac{2\, \operatorname{Tr} K^2}{1 + (\operatorname{Tr} K)^2}, \tag{A17}$$

$$\varphi''(0) = \frac{4\big(\operatorname{Tr} K^3\big)\big(1 + (\operatorname{Tr} K)^2\big) - 4(\operatorname{Tr} K)\big(\operatorname{Tr} K^2\big)^2}{\big(1 + (\operatorname{Tr} K)^2\big)^2}. \tag{A18}$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Design guideline: the HS norm $\|K\|_{S_2}^2 = \mathbf{Tr}\, K^2$ is the dominant factor of phase slope $\varphi'(0)$, and $\mathbf{Tr}\, K^3$ appears in the curvature at the origin. By adjusting CM parameters (e.g., σ, ν, α), $\mathbf{Tr}\, K^2$ and $\mathbf{Tr}\, K^3$ can be varied to tune the phase rise.

Appendix F.5. Consistency with Bandwidth, Endpoints, and Normalization [7,30]

To align with the small-band equalization in Main Text Section 6.1–Section 6.3, use $\mathbf{supp}\, \widehat{\phi} \subset [-\eta, \eta]$ ($\eta < \log 2$) for the observational test side. Endpoint contributions are treated per Main Text Section 10.2: eliminated by smooth windows (high-order vanishing at endpoints), or absorbed by Abel regularization for hard windows. The *design* of the kernel itself (this appendix) and the *observation* (band projection) do not commute, but by the limit interchange in Appendix A.6–A.7,

$$\langle \mu,\ \phi * s_\eta \rangle \xrightarrow[\eta \uparrow \infty]{} \langle \mu,\ \phi \rangle$$

is justified (under the same conditions as Main Text Section 6.5).

Appendix F.6. Parameterization and Differentiation (Optimization and Sensitivity Analysis) [6,15,68]

For representative families, parametric derivatives are given in closed form. Let $\lambda \geq 0$ be the spectral variable, and $K(\theta) = \phi(\cdot\,; \theta)(L)$.

Rational type $\phi(\lambda) = (c + \lambda)^{-\sigma}$:

$$\partial_c \phi(\lambda) = -\sigma\, (c + \lambda)^{-\sigma - 1}, \qquad \partial_\sigma \phi(\lambda) = -\log(c + \lambda)\, (c + \lambda)^{-\sigma}. \tag{A19}$$

Matérn (first order) $\phi(\lambda) = (1 + \alpha\lambda)^{-\nu}$:

$$\partial_\alpha \phi(\lambda) = -\nu\, \lambda\, (1 + \alpha\lambda)^{-\nu - 1}, \qquad \partial_\nu \phi(\lambda) = -\log(1 + \alpha\lambda)\, (1 + \alpha\lambda)^{-\nu}. \tag{A20}$$

Heat-type Matérn $\phi(\lambda) = (1 + \alpha\lambda^2)^{-\nu}$:

$$\partial_\alpha \phi(\lambda) = -\nu\, \lambda^2\, (1 + \alpha\lambda^2)^{-\nu - 1}, \qquad \partial_\nu \phi(\lambda) = -\log(1 + \alpha\lambda^2)\, (1 + \alpha\lambda^2)^{-\nu}. \tag{A21}$$

The operator derivative follows from the functional-analytic chain rule:

$$\partial_\theta K(\theta) = (\partial_\theta \phi)(L; \theta).$$

Sensitivity of Cayley phase:

$$\partial_\theta \varphi'(t) = 2\, \frac{\partial_\theta M'(t)}{1 + M(t)^2} - 4\, \frac{M(t) M'(t)\, \partial_\theta M(t)}{(1 + M(t)^2)^2},$$

with $M^{(m)}(0) = m!\, \mathbf{Tr}\, K^{m+1}$ and $\partial_\theta M^{(m)}(0) = m! \sum_{j=0}^{m} \mathbf{Tr}\big(K^{m-j} (\partial_\theta K) K^j\big)$ allowing design of slope/curvature at the origin. In optimization, use trace cyclicity and the $\mathbf{S}_p$ duality $\|\partial_\theta K\|_{\mathbf{S}_1} \leq \|\partial_\theta \phi(L)\|_{\mathbf{S}_1}$.

Appendix F.7. Implementation Guidelines and Numerical Stability [58,69]

- **Phase correction near poles**: φ jumps by π at $t = \kappa_n^{-1}$. Choose initial condition $T_{\max} < 1/\kappa_{\max}$ so that the observation window $[-T, T]$ does not intersect poles (safe side), and correct near poles by splitting (left/right grids) plus unwrapping (Main Text Section 8.6B).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

- **Bandwidth design**: strictly enforce $\eta \le \log 2 - \delta$ ($\delta > 0$). For hard windows (Vaaler type), use *explicit estimates* of endpoint terms; for smooth windows, advantage in endpoint elimination (see Main Text Section 10.2).
- **AFE (two-sum) truncation**: match left/right truncation lengths to analytic conductor $C_\pi(t)$ with $N_\pm(t) \asymp C_\pi(t)^{1/2}$ (Main Text Section 8.6). Use compensated summation (Kahan) to mitigate loss of significance from imbalance.
- **Schatten management**: at design stage, control parameters to satisfy threshold $sp > \alpha$ in Proposition A3 (HS means $p = 2$).

Appendix F.8. Summary (Design Guidelines)

- **Ensure positivity**: CM families or heat-type mixtures (averages of e^{-tL}/e^{-tL^2}) guarantee $K = \phi(L) \ge 0$ (Main Text Section 7.1–Section 7.2).
- **Schatten thresholds**: under $N_L(\Lambda) \ll \Lambda^\alpha$, design $\phi(\lambda) \ll (1+\lambda)^{-s}$ to satisfy $sp > \alpha$ (HS: $s > \alpha/2$, Trace: $s > \alpha$).
- **Phase design**: origin slope controlled by $\mathbf{Tr}\, K^2$, curvature by $\mathbf{Tr}\, K^3$ ((A17), (A18)); tune $\mathbf{Tr}\, K^m$ via parameters (σ, ν, α).
- **Implementation consistency**: $\eta < \log 2$, eliminate endpoints with smooth window or absorb via Abel regularization, base AFE truncation on $N_\pm \asymp C_\pi^{1/2}$ (Main Text Section 6.5, Section 8.6, Section 10.2).

Appendix G. Concrete Construction of Window Functions / Bandwidth Approximation and Error Control

In this appendix, we provide explicit constructions and precise estimates for *band-limited test functions (windows)* used in the main text for small-band equalization (Section 6), regularized determinants and phase extraction (Section 7), densification and AFE design (Section 8), and endpoint term control (Section 10). We follow the Fourier convention from Main Text Section 6:

$$\widehat{f}(\lambda) = \int_{\mathbb{R}} f(t)\, e^{-it\lambda}\, dt, \qquad f(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{f}(\lambda)\, e^{i\lambda t}\, d\lambda,$$

and convolution $(f * g)(t) = \int_{\mathbb{R}} f(u)\, g(t-u)\, du$. The bandwidth is always $0 < \eta < \log 2$ (strict), and we write the even/real band-limited test space as

$$A_\eta := \{\phi \in \mathcal{S}(\mathbb{R}) :\ \phi(t) = \phi(-t),\ \mathrm{supp}\, \widehat{\phi} \subset [-\eta, \eta]\}$$

(see Main Text Section 6.1–Section 6.3, Section 8.6).

Appendix G.1. Beurling–Selberg / Vaaler-type Approximation of Indicator Functions [30,70–72]

Basic kernel and bandwidth preservation.

For $A > 0$, define the normalized sinc

$$s_A(t) := \frac{\sin(At)}{\pi t} \quad \Rightarrow \quad \widehat{s_A}(\lambda) = \mathbf{1}_{[-A,A]}(\lambda)$$

(with the convention that the endpoint value is $1/2$; we will use only the a.e. equality). Let

$$\mathcal{V}_\eta(t) := s_{(\eta+\pi)/2}(t)\, s_{(\eta-\pi)/2}(t).$$

Then, by the fact that *the Fourier transform of a product is the convolution*:

$$\widehat{\mathcal{V}_\eta}(\lambda) = \frac{1}{2\pi} \mathbf{1}_{[-(\eta+\pi)/2,(\eta+\pi)/2]} * \mathbf{1}_{[-(\eta-\pi)/2,(\eta-\pi)/2]}(\lambda),$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

so $\widehat{\mathcal{V}_\eta}$ is a *trapezoidal mask supported on* $[-\eta,\eta]$ (flat in the central region $[-(\eta-\pi),\eta-\pi]$).

Vaaler-type upper and lower approximations.

Approximate the interval indicator $\mathbf{1}_{[-T,T]}$ from above and below within bandwidth η by

$$\Phi^{\pm}_{T,\eta}(t) \;:=\; (\mathbf{1}_{[-T,T]} \pm \delta_{T,\eta}) * \mathcal{V}_\eta(t) \tag{A22}$$

($\delta_{T,\eta}$ is an endpoint correction; see below (A25)). $\Phi^{\pm}_{T,\eta}$ are even/real with $\widehat{\Phi^{\pm}_{T,\eta}} = \widehat{\mathbf{1}_{[-T,T]}}\cdot\widehat{\mathcal{V}_\eta} \pm \widehat{\delta_{T,\eta}}\cdot\widehat{\mathcal{V}_\eta}$, and $\operatorname{supp}\widehat{\Phi^{\pm}_{T,\eta}} \subset [-\eta,\eta]$, hence $\Phi^{\pm}_{T,\eta} \in A_\eta$.

Proposition A5 (L^1 gap and interior approximation)**.** *Let* $T \geq 1$*. There exists* $C > 0$ *such that*

$$0 \;\leq\; \int_{\mathbb{R}} (\Phi^{+}_{T,\eta} - \mathbf{1}_{[-T,T]})\,dt \;\leq\; \frac{C}{\eta}, \qquad 0 \;\leq\; \int_{\mathbb{R}} (\mathbf{1}_{[-T,T]} - \Phi^{-}_{T,\eta})\,dt \;\leq\; \frac{C}{\eta}. \tag{A23}$$

Furthermore, for $t \in (-T,T)$*,*

$$\Phi^{\pm}_{T,\eta}(t) \;=\; 1 \;+\; O\Big(\frac{1}{1+\eta\,\mathrm{dist}(t,\{\pm T\})}\Big), \tag{A24}$$

and once t *leaves the endpoints,* $|\Phi^{\pm}_{T,\eta}(t)| \ll (1+\eta\,||t|-T|)^{-2}$*.*

Sketch of proof. *(i)* L^1 *gap.* Since $\int f = \widehat{f}(0)$, $\int(\Phi^{+}_{T,\eta} - \mathbf{1}_{[-T,T]}) = \widehat{\mathcal{V}_\eta}(0)\widehat{\mathbf{1}_{[-T,T]}}(0) - \widehat{\mathbf{1}_{[-T,T]}}(0) + \int \delta_{T,\eta} * \mathcal{V}_\eta$. $\widehat{\mathcal{V}_\eta}(0) = \frac{1}{2\pi}\left((\eta+\pi)/2 + (\eta-\pi)/2\right) = \frac{\eta}{2\pi}$ is proportional to a positive constant, and the optimal Vaaler approximation (optimized endpoint correction) keeps the difference at $O(\eta^{-1})$. The lower case is analogous.

(ii) Interior/exterior estimates. $\mathcal{V}_\eta$ is an L^1 approximate identity; if t is inside by distance d from an endpoint, the L^1-concentration of the convolution deteriorates by $O((1+\eta d)^{-1})$. Outside, integration by parts yields $O((1+\eta\,||t|-T|)^{-2})$. □

Appendix G.2. Endpoint Correction Structure and Vanishing Order [30,64,71]

For hard barriers, $\widehat{\Phi^{\pm}_{T,\eta}}$ has corners at $\pm\eta$, producing *endpoint terms* in Main Text Section 10.2 (estimates (78), (88)). To suppress them, set in the Fourier side:

$$\widehat{\delta_{T,\eta}}(\lambda) \;=\; \sum_{j=0}^{m} c_j(T,\eta)\,\Big(\delta(\lambda-\eta) + \delta(\lambda+\eta)\Big)^{(j)} \tag{A25}$$

($m \in \mathbb{N}$ is the vanishing order), and choose c_j so that

$$\widehat{\Phi^{\pm}_{T,\eta}}^{(k)}(\pm\eta) = 0 \qquad (k = 0,1,\dots,m) \tag{A26}$$

holds; then the endpoint contributions vanish to high order.

Proposition A6 (Upper bound on endpoint term)**.** *If* (A26) *holds, the endpoint term in Main Text Section 10.2 satisfies*

$$\mathbf{End}_\eta[\Phi^{\pm}_{T,\eta}] \;=\; O_m\Big((1+\eta T)^{-m}\Big) \tag{A27}$$

(with constant depending only on m*).*

Sketch of proof. Integrate by parts m times the local Fourier integral near the endpoint, and use (A26) to kill the boundary term. Each integration yields $(\eta T)^{-m}$. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Appendix G.3. Smooth Windows and High-Order Vanishing [25,46,64]

A smooth window uses an even, smooth, compactly supported frequency mask $m_\eta \in C_c^\infty([-\eta,\eta])$:

$$\widehat{\Phi^{\mathrm{sm}}_{T,\eta}}(\lambda) := m_\eta(\lambda)\cdot\widehat{\mathbf{1}_{[-T,T]}}(\lambda) = m_\eta(\lambda)\cdot\frac{2\sin(\lambda T)}{\lambda} \tag{A28}$$

(e.g., $m_\eta(\lambda)=\vartheta(\lambda/\eta)$, $\vartheta\in C_c^\infty([-1,1])$, $\vartheta\equiv 1$ on $[-1+\delta,1-\delta]$). Then $\Phi^{\mathrm{sm}}_{T,\eta}=\check{m}_\eta * \mathbf{1}_{[-T,T]}$ and $\check{m}_\eta$ is rapidly decaying:

$$|\check{m}_\eta(t)| \le C_{m,\vartheta}\,(1+\eta|t|)^{-m} \qquad (\forall m\in\mathbb{N}),$$

so the exterior decays faster than any polynomial. If m_η vanishes to order m at the endpoints, (A26) applies and $\mathbf{End}_\eta[\Phi^{\mathrm{sm}}_{T,\eta}] = O_m((1+\eta T)^{-m})$ (matching Main Text Section 10.2 Remark 10.22).

Appendix G.4. Norm Control and Paley–Wiener [7,51,73]

Lemma A3 (Basic norms). *For $T\ge 1$, $0<\eta<\log 2$, and $\Phi\in\{\Phi^{\pm}_{T,\eta},\Phi^{\mathrm{sm}}_{T,\eta}\}$,*

$$\|\Phi\|_{L^1} = 2T+O(\eta^{-1}), \qquad \|\Phi\|_{L^\infty}\le 1+O(\eta^{-1}), \qquad \|\Phi'\|_{L^1}\ll 1+\log(1+\eta T).$$

Proof. $\int\Phi=\widehat{\Phi}(0)=\widehat{\mathcal{V}_\eta}(0)\widehat{1}(0)+\cdots=2T+O(\eta^{-1})$. L^∞ follows from Young's inequality and $\|\mathcal{V}_\eta\|_{L^1}=1+O(\eta^{-1})$. For the derivative, $\widehat{\Phi'}(\lambda)=i\lambda\,\widehat{\Phi}(\lambda)$ and the support $[-\eta,\eta]$ give the estimate. □

Lemma A4 (Paley–Wiener type). *If $\Phi\in A_\eta$, then Φ is an entire function of exponential type $\le\eta$, and*

$$|\Phi(x+iy)| \le C_\Phi\, e^{\eta|y|} \qquad (x,y\in\mathbb{R}).$$

Proof. Immediate from $\Phi(z)=\frac{1}{2\pi}\int_{-\eta}^{\eta}\widehat{\Phi}(\lambda)e^{i\lambda z}\,d\lambda$. □

Appendix G.5. Commutation of Poisson Smoothing and Abel Regularization [45,48,74]

The Abel regularization in Main Text Section 6.5 ($A\text{-}\lim_{\varepsilon\downarrow 0}$) applies *cut-off* and *smoothing* first so the limits commute.

Proposition A7 (Commutation). *Let F be a bounded variation measure (or a tempered distribution of finite order). For any of the above windows $\Phi_{T,\eta}$,*

$$\lim_{\varepsilon\downarrow 0}\lim_{T\to\infty}\langle F,\, e^{-\varepsilon|\cdot|}\Phi_{T,\eta}\rangle = \lim_{T\to\infty}\langle F,\,\Phi_{T,\eta}\rangle \tag{A29}$$

holds.

Proof. $e^{-\varepsilon|t|}\Phi_{T,\eta}(t)$ is uniformly L^1-integrable, and as $\varepsilon\downarrow 0$, $T\to\infty$, the integrand converges F-a.e. to $\Phi_{T,\eta}(t)$. Apply dominated convergence (for measures) / continuity of bounded linear functionals (for distributions). Endpoint terms are uniformly controlled by Proposition A6 at $O_m((1+\eta T)^{-m})$. □

Appendix G.6. Phase Average Testing and Cayley Phase Extraction [2,44,59]

For the Cayley transform $S=(1-iM)/(1+iM)=e^{-i\varphi}$ in Main Text Section 7.2, compare/identify phases using the phase average

$$\int_{\mathbb{R}}\varphi(t)\,\Phi_{T,\eta}(t)\,dt$$

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

(the phase is even, normalized by $\varphi(0) = 0$). On the determinant side,

$$\partial_t \log \det_2(I - tK) \;=\; \sum_n \frac{\kappa_n}{1 - t\kappa_n} \;-\; \sum_n \kappa_n \tag{A30}$$

(Main Text (116)) gives

$$\varphi(t) \;=\; 2\,\arctan\Big(\partial_t \log \det_2(I - tK) + \operatorname{Tr} K\Big), \tag{A31}$$

so

$$\int \varphi\, \Phi_{T,\eta} = 2 \int \arctan\Big(\partial_t \log \det_2(I - tK) + \operatorname{Tr} K\Big)\, \Phi_{T,\eta}(t)\, dt.$$

By choosing $\Phi_{T,\eta}$ hard/smooth, endpoint contributions are absorbed via (78), (88) or Proposition A6 into $O_m((1+\eta T)^{-m}) + O(\eta^{-1})$.

Appendix G.7. AFE (Two-Sum) Truncation and Bandwidth Matching [4,8,32]

In the approximate functional equation (AFE; equation (110) in Main Text Section 8.6), adjust the left/right truncation lengths according to the analytic conductor $C_\pi(t)$:

$$N_\pm(t) \;\asymp\; C_\pi(t)^{1/2}.$$

To match with window $\Phi_{T,\eta}$, choose T in the range

$$T \;\asymp\; \eta^{-1} \log C_\pi(t),$$

so that the combined error of the Vaaler gap (A23) $O(\eta^{-1})$ and endpoint term (A27) is below the digit accuracy of the AFE. Numerically, use compensated summation (Kahan) to suppress cancellation loss from imbalance (Main Text Section 8.6D).

Appendix G.8. Testing Small-Band Equalization [5,49,63,75]

The statement in Main Text Section 6.1–Section 6.3 is that for the distribution difference $\Delta = \mu_{L^{[d]}} - \mu_{\Xi_\pi}$,

$$\langle \Delta,\, \phi \rangle = 0 \qquad (\forall\, \phi \in A_\eta).$$

Using the explicit window family $\Phi_{T,\eta} \in A_\eta$ above,

$$\lim_{T\to\infty} \langle \Delta,\, \Phi_{T,\eta} \rangle \;=\; 0, \tag{A32}$$

and via Fourier transform,

$$\langle \widehat{\Delta},\, \widehat{\Phi_{T,\eta}} \rangle = 0 \qquad \big(\operatorname{supp} \widehat{\Phi_{T,\eta}} \subset [-\eta, \eta]\big)$$

implies $\operatorname{supp} \widehat{\Delta} \subset \mathbb{R} \setminus (-\eta, \eta)$ (Main Text Section 6.3, Section 8.7A). Even with hard windows, (A26) and Proposition A6 ensure endpoint terms vanish as $T \to \infty$.

Appendix G.9. Recipe: Window Family Selection and Error Budget [30,32,57,71,76]

- **Theoretical testing** (for proofs): adopt the smooth window (A28) with arbitrary endpoint vanishing order m. Endpoint terms vanish automatically (Main Text Section 10.2 Remark 10.22), and Paley–Wiener (Lemma A4) ensures exponential type $\leq \eta$.
- **Visualization/experiments** (clear view): use hard barrier (A22) to create a window close to a step function, adding endpoint correction (A25) with $m = 1, 2$ to fit (78), (88). Total error is $O(\eta^{-1}) + O_m((1+\eta T)^{-m})$.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

- **AFE matching**: choose $T \asymp \eta^{-1} \log C_\pi$ in accordance with $N_\pm \asymp C_\pi^{1/2}$ (Main Text (110)). Adjust η, m so that the Vaaler gap (A23) is much smaller than the AFE error.
- **Phase extraction**: extract φ using (A30)–(A31), monitor origin normalization $\varphi(0) = 0$ and monotonicity (Main Text Section 7.5 Lemma).
- **Small-band equalization**: use (A32) as the test, strictly enforce $\eta < \log 2$ to push prime terms outside the band (Main Text Section 6.1–Section 6.3).

In summary, we have given a self-contained *explicit design, error control, and commutation* of band-limited test functions consistent with Main Text (equations (78), (88), (110), (116) and Remark in Section 10.2).

Appendix H. Complete Expansion of Endpoint Terms and Constants in the Explicit Formula

In this appendix, we rigorously decompose the *Weil-type explicit formula* for the completed GL(d) L-function

$$\Lambda(s,\pi) = Q_\pi^{s/2} \prod_{j=1}^{d} \Gamma_{\mathbb{R}}(s+\mu_j)\, L(s,\pi), \qquad \Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma\Big(\frac{s}{2}\Big)$$

into (i) *prime sum (discrete frequencies)*, (ii) *Archimedean term*, (iii) *conductor term*, and (iv) *endpoint contributions (half rule)*, and fully expand the handling of constants and endpoints. The Fourier convention follows Main Text Section 6:

$$\widehat{\varphi}(\xi) = \int_{\mathbb{R}} \varphi(t) e^{-i\xi t}\, dt, \qquad \varphi \in \mathcal{S}_{\mathrm{even}}(\mathbb{R})$$

(evenness automatically symmetrizes under $t \mapsto -t$).

Appendix H.1. Skeleton and Decomposition of the GL(d) Explicit Formula [1,11,32]

Theorem A3 (Weil-type explicit formula (t-side representation)). *For $\varphi \in \mathcal{S}_{\mathrm{even}}$, the sum over imaginary parts γ of nontrivial zeros $\rho = \frac{1}{2} + i\gamma$ satisfies*

$$\sum_\rho \varphi(\operatorname{Im}\rho) \;=\; \mathsf{A}_\pi[\varphi] \;+\; \mathsf{C}_\pi[\varphi] \;+\; \mathsf{P}_\pi[\varphi] \;+\; \mathsf{E}_\pi[\varphi], \tag{A33}$$

where

$$\mathsf{A}_\pi[\varphi] := \frac{1}{\pi} \sum_{j=1}^{d} \int_{\mathbb{R}} \varphi(t)\, \operatorname{Re} \frac{\Gamma_{\mathbb{R}}'}{\Gamma_{\mathbb{R}}}\Big(\frac{1}{2} + \mu_j + it\Big)\, dt, \tag{A34}$$

$$\mathsf{C}_\pi[\varphi] := \frac{\log Q_\pi}{2\pi} \int_{\mathbb{R}} \varphi(t)\, dt, \tag{A35}$$

$$\mathsf{P}_\pi[\varphi] := -2 \sum_{n\ge 2} \frac{\Lambda_\pi(n)}{\sqrt{n}}\, \operatorname{Re}\, \widehat{\varphi}(\log n), \tag{A36}$$

with $\Lambda_\pi(n)$ the von Mangoldt-type coefficients of $-L'/L(s,\pi) = \sum_{n\ge 1} \Lambda_\pi(n) n^{-s}$, and the endpoint/pole contribution is

$$\mathsf{E}_\pi[\varphi] \;:=\; \frac{m_\pi}{2}\, \varphi(0) \;+\; \mathsf{End}_\eta[\varphi]. \tag{A37}$$

Here $m_\pi \in \{0,1\}$ is the total order of simple poles at $s=1$ and $s=0$ of $\Lambda(s,\pi)$ (for cusp forms $m_\pi = 0$), and $\mathsf{End}_\eta[\varphi]$ is the finite sum from the half rule *at the band endpoint (see H.3).*

Proof sketch. The standard Weilian transformation: regard $\Phi(s) := \int_{\mathbb{R}} \varphi(t)\, n^{it}\, dt = \widehat{\varphi}(\log n)$ as a Mellin kernel, take the vertical line integral of $\frac{\Lambda'}{\Lambda}(s,\pi)$ along $\operatorname{Re} s = 2$, and reflect to $\operatorname{Re} s = -1$ via the functional equation. Residue calculus recovers residues at nontrivial zeros (ρ) and poles ($s = 1, 0$), and the Euler product logarithmic derivative yields (A36). Evenness makes the complex conjugate

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

terms match, allowing combination via **Re**. The Archimedean factors appear as linear combinations of $\Gamma'_{\mathbb{R}}/\Gamma_{\mathbb{R}}$ in (A34). □

Remark A4 (Agreement with main term in the text). *Applying Stirling's expansion (Appendix A.10) to* (A34) *gives* $\mathbf{Re}\,\frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}}(\frac{1}{2}+\mu_j+it) = \log\frac{|t|}{2\pi} + O((1+t^2)^{-1})$, *and combining* (A34) *with* (A35) *yields*

$$\frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\Big\{d\log(t^2) - d\log(4\pi^2) + \log Q_\pi\Big\}\,dt \;+\; O\Big(\|\varphi\|_{L^1(\langle t\rangle^{-2}dt)}\Big),$$

matching the calibration in Main Text Section 6.2.

Appendix H.2. Complete Expansion of the Archimedean Term into "Main Term + Constant" [4,35]

Separation of constants via Stirling.

For fixed $\mu \in \mathcal{C}$,

$$\frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}}\Big(\frac{1}{2}+\mu+it\Big) = \log\frac{|t|}{2\pi} \;+\; R_\mu(t), \qquad R_\mu(t) = O\Big(\frac{1}{1+t^2}\Big),$$

based on the asymptotic $\psi(z) = \Gamma'(z)/\Gamma(z) = \log z + O(1/z)$ for $\mathbf{Re}\,z > 0$. Hence

$$\mathbf{A}_\pi[\varphi] \;=\; \frac{d}{2\pi}\int_{\mathbb{R}} \varphi(t)\log|t|\,dt \;-\; \frac{d\log(2\pi)}{2\pi}\int_{\mathbb{R}} \varphi(t)\,dt \;+\; \mathbf{A}_\pi^{(\mathrm{rem})}[\varphi], \tag{A38}$$

$$\mathbf{A}_\pi^{(\mathrm{rem})}[\varphi] := \frac{1}{\pi}\sum_{j=1}^{d}\int_{\mathbb{R}} \varphi(t)\;\mathbf{Re}\,R_{\mu_j}(t)\,dt.$$

Since $\mathbf{Re}\,R_{\mu_j}(t) = O((1+t^2)^{-1})$,

$$|\mathbf{A}_\pi^{(\mathrm{rem})}[\varphi]| \;\ll_{d,\mu}\; \int_{\mathbb{R}} \frac{|\varphi(t)|}{1+t^2}\,dt. \tag{A39}$$

Combination with conductor term.

Combining (A38) with (A35) gives

$$\mathbf{A}_\pi[\varphi] + \mathbf{C}_\pi[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\Big\{d\log(t^2) - d\log(4\pi^2) + \log Q_\pi\Big\}\,dt \;+\; \mathbf{A}_\pi^{(\mathrm{rem})}[\varphi], \tag{A40}$$

exactly matching the main term kernel in Main Text Section 6.2.

Appendix H.3. Refinement of Prime Sum and Endpoint Contribution (Half Rule) [1,7,32,63]

Standard form of the prime sum.

Equation (A36) becomes

$$\mathbf{P}_\pi[\varphi] = -2\sum_{p}\sum_{k\ge 1}\frac{\log p}{p^{k/2}}\;\mathbf{Re}\Big(\mathrm{tr}\,A_p(\pi)^k\,\widehat{\varphi}(k\log p)\Big), \tag{A41}$$

where $A_p(\pi)$ is the Satake matrix, $\mathrm{tr}\,A_p(\pi)^k = \sum_{j=1}^{d}\alpha_{p,j}^k$.

Proposition A8 (Vanishing in small band). *If* $\mathbf{supp}\,\widehat{\varphi} \subset [-\eta,\eta]$ *with* $\eta < \log 2$, *then* $\mathbf{P}_\pi[\varphi] = 0$.

Proof. Nonzero $\widehat{\varphi}(k\log p)$ requires $k\log p \le \eta$. The smallest case $k=1,\ p=2$ gives $\log 2 > \eta$, a contradiction; hence all vanish. □

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Lemma A5 (Half rule at band endpoint). *If* $\operatorname{supp}\widehat{\varphi} \subset [-\eta,\eta]$ *and* $\widehat{\varphi}$ *is discontinuous at* $\lambda = \pm\eta$ *but has left/right limits, the Riemann–Stieltjes theory gives*

$$\int_{[-\eta,\eta]} \widehat{\varphi}(\lambda)\, dF(\lambda) \;=\; \int_{(-\eta,\eta)} \widehat{\varphi}\, dF \;+\; \frac{1}{2}\big(\widehat{\varphi}(+\eta)+\widehat{\varphi}(-\eta)\big)\,\Delta F(\eta),$$

($\Delta F(\eta)$ *is the point mass). Applying this to* (A41) *yields*

$$\mathbf{P}_\pi[\varphi] = -\operatorname{Re}\big(\widehat{\varphi}(\eta)\, w_{2,\pi}\big) \;+\; O_\eta(0), \qquad w_{2,\pi} := \log 2\, \operatorname{tr} A_2(\pi), \tag{A42}$$

where $O_\eta(0)$ *vanishes if* $\eta < \log 3$.

Remark A5 (Suppressing endpoint contribution). *If a smooth window (Appendix G) satisfies* $\widehat{\varphi}(\pm\eta) = 0$, *the endpoint term in* (A42) *vanishes. For a hard window, Appendix G's endpoint correction (vanishing order* m*) absorbs it as* $\mathbf{End}_\eta[\varphi] = O_m\big((1+\eta T)^{-m}\big)$.

Appendix H.4. Reduction to Concrete Families: ζ*, Dirichlet, GL(2) [4,8,13,32]*

(i) Riemann ζ.

$d = 1$, $\mu_1 = 0$, $Q_\pi = 1$. From (A40) and (A41),

$$\sum_\rho \varphi(\operatorname{Im}\rho) = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\{\log(t^2) - \log(4\pi^2)\}\, dt \;-\; 2\sum_{n\ge 2}\frac{\Lambda(n)}{\sqrt{n}}\operatorname{Re}\,\widehat{\varphi}(\log n) + \mathbf{A}_\zeta^{(\mathrm{rem})}[\varphi] \;+\; \frac{1}{2}\varphi(0),$$

the last $\frac{1}{2}\varphi(0)$ coming from the simple poles at $s = 1, 0$ of $\Lambda(s) = \pi^{-s/2}\Gamma(s/2)\zeta(s)$ (half rule for simple poles). $\mathbf{A}_\zeta^{(\mathrm{rem})}$ is bounded as in (A39).

(ii) Dirichlet $L(s,\chi)$ (primitive character).

For parity $a \in \{0,1\}$, $\Lambda(s,\chi) = (q/\pi)^{(s+a)/2}\Gamma\big((s+a)/2\big)L(s,\chi)$. Here $m_\pi = 0$ (primitive nontrivial), $Q_\pi = q$. Apply (A40) and (A41) directly.

(iii) GL(2) cusp forms.

For Maass forms $\{\mu_j\} = \{\frac{1+\varepsilon}{2} \pm ir\}$, for holomorphic weight k $\{\mu_j\} = \{\frac{k-1}{2}, \frac{k+1}{2}\}$. In both cases $m_\pi = 0$; substitute into (A40) and (A41) to evaluate main and remainder kernels.

Appendix H.5. Rigorous Handling of Band Tests Including Endpoints [7,30,46]

Test design principle.

If $\operatorname{supp}\widehat{\varphi} \subset [-\eta,\eta]$ and $\widehat{\varphi}(\pm\eta) = 0$ (e.g. smooth window vanishing to high order at endpoints), the endpoint term in (A42) vanishes. For hard windows, Appendix G's endpoint correction (moment vanishing $\widehat{\varphi}^{(k)}(\pm\eta) = 0$) ensures $\mathbf{End}_\eta[\varphi] = O_m\big((1+\eta T)^{-m}\big)$.

General form of the Stieltjes half rule.

For a function F of bounded variation and ψ with left/right limits,

$$\int_{[a,b]} \psi\, dF \;=\; \int_{(a,b)} \psi\, dF \;+\; \frac{1}{2}\psi(a)\,\Delta F(a) \;+\; \frac{1}{2}\psi(b)\,\Delta F(b).$$

Applied to prime truncation $F(x) = \sum_{k\log p \le x}\cdots$, this yields (A42).

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

Commutation with Abel regularization.

Appendix G (Proposition G.5) shows that the limits $\varphi \mapsto e^{-\varepsilon|t|}\varphi$ and $T \to \infty$ commute, keeping endpoint term handling regular.

Appendix H.6. Summary

- The Archimedean term $\mathbf{A}_\pi$ and conductor term $\mathbf{C}_\pi$ decompose completely as

$$\mathbf{A}_\pi[\varphi] + \mathbf{C}_\pi[\varphi] = \frac{1}{2\pi}\int_{\mathbb{R}} \varphi(t)\Big\{d\log(t^2) - d\log(4\pi^2) + \log Q_\pi\Big\}dt \; + \; \mathbf{A}_\pi^{(\mathrm{rem})}[\varphi],$$

  with $\mathbf{A}_\pi^{(\mathrm{rem})}$ controlled by the bounded kernel in (A39).
- The prime sum $\mathbf{P}_\pi$ *vanishes* for small band $\eta < \log 2$ (Proposition A8); when touching the endpoint $\eta = \log 2$, it reduces to the finite correction from the *half rule* (A42) (vanishes for smooth windows).
- Poles of Λ at $s = 1, 0$ are aggregated as $\frac{m_\pi}{2}\varphi(0)$ ((A37)); for cusp forms this is zero.

Acknowledgments: In the pre-review of the initial version of this manuscript (September 2025, by Anik Chakraborty, University of Delhi, India), we received constructive and concrete comments concerning small-band equivalence, the uniqueness principle, rigorous treatment of band-edge terms, the regularized Fredholm determinant and coefficient identification, and the justification of densification and exchange of limits. In light of this advice, we created Section 11 and systematically reinforced these points to enhance the self-contained nature and auditability of the proofs. We express our sincere gratitude here. Any remaining inadequacies or errors in this manuscript are solely the responsibility of the author.

References

1. Weil, A. Sur les 'formules explicites' de la théorie des nombres. In Proceedings of the Acta Universitatis Lundensis, Nova Series, Sectio II, Lund, 1952; pp. 252–265. Tome supplémentaire, dédié à Marcel Riesz.
2. William F. Donoghue, J. *Monotone Matrix Functions and Analytic Continuation*; Vol. 207, *Grundlehren der Mathematischen Wissenschaften*, Springer: Berlin, 1974.
3. Iwaniec, H.; Kowalski, E. *Analytic Number Theory*; Vol. 53, *American Mathematical Society Colloquium Publications*, American Mathematical Society: Providence, RI, 2004.
4. Titchmarsh, E.C. *The Theory of the Riemann Zeta-Function*, 2nd ed.; Oxford University Press: Oxford, 1986. Revised by D. R. Heath-Brown.
5. Edwards, H.M. *Riemann's Zeta Function*; Academic Press: New York, 1974.
6. Simon, B. *Trace Ideals and Their Applications*, 2nd ed.; Vol. 120, *Mathematical Surveys and Monographs*, American Mathematical Society: Providence, RI, 2005.
7. Paley, R.E.A.C.; Wiener, N. *Fourier Transforms in the Complex Domain*; Vol. 19, *American Mathematical Society Colloquium Publications*, American Mathematical Society: New York, 1934.
8. Davenport, H. *Multiplicative Number Theory*, 3rd ed.; Vol. 74, *Graduate Texts in Mathematics*, Springer: New York, 2000. Revised by Hugh L. Montgomery.
9. Neukirch, J. *Algebraic Number Theory*; Springer: Berlin, 1999.
10. Tate, J.T. Fourier Analysis in Number Fields and Hecke's Zeta-Functions. In *Algebraic Number Theory*; Cassels, J.W.S.; Fröhlich, A., Eds.; Academic Press: London, 1967; pp. 305–347. Originally Ph.D. thesis, Princeton University, 1950.
11. Godement, R.; Jacquet, H. *Zeta Functions of Simple Algebras*; Vol. 260, *Lecture Notes in Mathematics*, Springer: Berlin, 1972.
12. Gelbart, S. *Automorphic Forms on Adele Groups*; Vol. 83, *Annals of Mathematics Studies*, Princeton University Press: Princeton, NJ, 1975.
13. Goldfeld, D. *Automorphic Forms and L-Functions for the Group* $\mathbf{GL}(2)$; Vol. I, *Cambridge Studies in Advanced Mathematics*, Cambridge University Press: Cambridge, 2006.
14. Reed, M.; Simon, B. *Methods of Modern Mathematical Physics, Vol. I: Functional Analysis*; Academic Press: New York, 1972.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

15. Kato, T. *Perturbation Theory for Linear Operators*; Classics in Mathematics, Springer: Berlin, 1995. Reprint of the 1980 edition.
16. Brezis, H. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*; Springer: New York, 2010.
17. Adams, R.A.; Fournier, J.J.F. *Sobolev Spaces*, 2nd ed.; Pure and Applied Mathematics, Academic Press: Amsterdam, 2003.
18. Kress, R. *Linear Integral Equations*, 2nd ed.; Vol. 82, *Applied Mathematical Sciences*, Springer, 1999.
19. Reed, M.; Simon, B. *Methods of Modern Mathematical Physics, Vol. IV: Analysis of Operators*; Academic Press: New York, 1978.
20. Cycon, M.; Froese, R.; Kirsch, W.; Simon, B. *Schrödinger Operators*; Springer Study Edition, Springer: Berlin, 1987.
21. Lieb, E.H.; Loss, M. *Analysis*, 2nd ed.; Vol. 14, *Graduate Studies in Mathematics*, American Mathematical Society, 2001.
22. Riemann, B. *Über die Anzahl der Primzahlen unter einer gegebenen Größe*; Akademie der Wissenschaften: Berlin, 1859.
23. Ingham, A.E. *The Distribution of Prime Numbers*; Cambridge University Press: Cambridge, 1932.
24. Seiler, E.; Simon, B. An Inequality Among Determinants. *Proceedings of the National Academy of Sciences USA* **1975**, *72*, 3277–3278. https://doi.org/10.1073/pnas.72.9.3277.
25. Hörmander, L. *The Analysis of Linear Partial Differential Operators I: Distribution Theory and Fourier Analysis*; Springer: Berlin, 1983.
26. Gohberg, I.; Krein, M. *Introduction to the Theory of Linear Nonselfadjoint Operators*; Vol. 18, *Translations of Mathematical Monographs*, American Mathematical Society: Providence, RI, 1969.
27. Weyl, H. Über die asymptotische Verteilung der Eigenwerte. *Nachrichten von der Königlichen Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* **1911**, pp. 110–117.
28. Hadamard, J. *Lectures on Cauchy's Problem in Linear Partial Differential Equations*; Yale University Press: New Haven, 1923.
29. Grafakos, L. *Classical Fourier Analysis*, 3rd ed.; Vol. 249, *Graduate Texts in Mathematics*, Springer: New York, 2014.
30. Vaaler, J.D. Some Extremal Functions in Fourier Analysis. *Bulletin of the American Mathematical Society (N.S.)* **1985**, *12*, 183–216.
31. Montgomery, H.L. *Ten Lectures on the Interface Between Analytic Number Theory and Harmonic Analysis*; Number 84 in CBMS Regional Conference Series in Mathematics, American Mathematical Society: Providence, RI, 1994.
32. Iwaniec, H.; Kowalski, E. *Analytic Number Theory*; Vol. 53, *American Mathematical Society Colloquium Publications*, American Mathematical Society: Providence, RI, 2004.
33. Carneiro, E.; Vaaler, J.D. Some Extremal Functions in Fourier Analysis. II. *Transactions of the American Mathematical Society* **2010**, *362*, 5803–5843.
34. Ikehara, S. An Extension of Landau's Theorem in the Analytic Theory of Numbers. *Journal of Mathematics and Physics (MIT)* **1931**, *10*, 1–12.
35. Olver, F.W.J.; Lozier, D.W.; Boisvert, R.F.; Clark, C.W., Eds. *NIST Digital Library of Mathematical Functions*; National Institute of Standards and Technology: Gaithersburg, MD, 2010. Online companion.
36. R. P. Boas, J. *Entire Functions*; Academic Press: New York, 1954.
37. Conway, J.B. *Functions of One Complex Variable I*, 2nd ed.; Vol. 11, *Graduate Texts in Mathematics*, Springer: New York, 1978.
38. Bochner, S. Monotone Funktionen, Stieltjessche Integrale und harmonische Analyse. *Mathematische Annalen* **1933**, *108*, 378–410.
39. Nevanlinna, R. Zur Theorie der meromorphen Funktionen. *Acta Mathematica* **1925**, *46*, 1–99.
40. Nikol'skiĭ, S.M. *Approximation of Functions of Several Variables and Imbedding Theorems*; Springer: New York, 1975.
41. Duren, P.L. *Theory of H^p Spaces*; Pure and Applied Mathematics, Academic Press: New York, 1970.
42. Koosis, P. *The Logarithmic Integral. I*; Number 12 in Cambridge Studies in Advanced Mathematics, Cambridge University Press: Cambridge, 1988.
43. Muskhelishvili, N.I. *Singular Integral Equations*; Dover Publications: New York, 1992. Reprint of the 1953 edition.
44. Garnett, J.B. *Bounded Analytic Functions*, revised 1st ed.; Vol. 236, *Graduate Texts in Mathematics*, Springer: New York, 2007.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.

Preprints.org (www.preprints.org) | NOT PEER-REVIEWED | Posted: 20 October 2025 doi:10.20944/preprints202505.2110.v3

45. Zygmund, A. *Trigonometric Series*, 3rd ed.; Cambridge University Press: Cambridge, 2002.
46. Folland, G.B. *Real Analysis: Modern Techniques and Their Applications*, 2nd ed.; Wiley: New York, 1999.
47. Katznelson, Y. *An Introduction to Harmonic Analysis*, 3rd ed.; Cambridge University Press: Cambridge, 2004.
48. Titchmarsh, E.C. *Introduction to the Theory of Fourier Integrals*, 2nd ed.; Oxford University Press: Oxford, 1948.
49. Weil, A. *Sur les 'formules explicites' de la théorie des nombres*; Lunds Universitet, 1952.
50. Gel'fand, I.M.; Shilov, G.E. *Generalized Functions, Vol. 1: Properties and Operations*; Academic Press: New York, 1964.
51. Levin, B.Y. *Distribution of Zeros of Entire Functions*, revised ed.; American Mathematical Society: Providence, RI, 1996.
52. Rudin, W. *Real and Complex Analysis*, 3rd ed.; McGraw–Hill: New York, 1987.
53. Atkinson, K.E. *The Numerical Solution of Integral Equations of the Second Kind*; Cambridge University Press: Cambridge, 1997.
54. Widder, D.V. *The Laplace Transform*; Princeton University Press: Princeton, 1946.
55. Berman, A.; Plemmons, R.J. *Nonnegative Matrices in the Mathematical Sciences*; Classics in Applied Mathematics, Academic Press: New York, 1979.
56. Karlin, S. *Total Positivity, Vol. I*; Stanford University Press: Stanford, CA, 1968.
57. Slepian, D.; Pollak, H.O. Prolate Spheroidal Wave Functions, Fourier Analysis and Uncertainty—I. *Bell System Technical Journal* **1961**, *40*, 43–63.
58. Higham, N.J. *Accuracy and Stability of Numerical Algorithms*, 2nd ed.; SIAM: Philadelphia, PA, 2002.
59. de Branges, L. *Hilbert Spaces of Entire Functions*; Prentice–Hall: Englewood Cliffs, NJ, 1968.
60. Golub, G.H.; Loan, C.F.V. *Matrix Computations*, 4th ed.; Johns Hopkins University Press: Baltimore, 2013.
61. Stein, E.M.; Shakarchi, R. *Fourier Analysis: An Introduction*; Vol. 1, *Princeton Lectures in Analysis*, Princeton University Press: Princeton, NJ, 2003.
62. Grafakos, L. *Classical Fourier Analysis*, 3rd ed.; Vol. 249, *Graduate Texts in Mathematics*, Springer, 2014.
63. Guinand, A.P. A Summation Formula in the Theory of Prime Numbers. *Proceedings of the London Mathematical Society* **1948**, *50*, 107–119. Often cited in connection with the 1955 literature on explicit formulas.
64. Stein, E.M.; Shakarchi, R. *Fourier Analysis: An Introduction*; Vol. 1, *Princeton Lectures in Analysis*, Princeton University Press, 2003.
65. Schilling, R.L.; Song, R.; Vondraček, Z. *Bernstein Functions: Theory and Applications*, 2nd ed.; Vol. 37, *de Gruyter Studies in Mathematics*, de Gruyter: Berlin, 2012.
66. Stein, E.M. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*; Vol. 43, *Princeton Mathematical Series*, Princeton University Press: Princeton, NJ, 1993.
67. Davies, E.B. *Heat Kernels and Spectral Theory*; Vol. 92, *Cambridge Tracts in Mathematics*, Cambridge University Press: Cambridge, 1989.
68. Higham, N.J. *Functions of Matrices: Theory and Computation*; SIAM: Philadelphia, PA, 2008.
69. Golub, G.H.; Loan, C.F.V. *Matrix Computations*, 4th ed.; Johns Hopkins University Press, 2013.
70. Graham, S.W.; Vaaler, J.D. A Class of Extremal Functions for the Fourier Transform. *Transactions of the American Mathematical Society* **1981**, *265*, 283–302.
71. Carneiro, E.; Littmann, F.; Vaaler, J.D. Gaussian Subordination for the Beurling–Selberg Extremal Problem. *Transactions of the American Mathematical Society* **2013**, *365*, 3493–3534.
72. Selberg, A. *Collected Papers, Vol. I*; Springer: Berlin, 1989.
73. Katznelson, Y. *An Introduction to Harmonic Analysis*, 3rd ed.; Cambridge University Press, 2004.
74. Hardy, G.H. *Divergent Series*; Oxford University Press: Oxford, 1949.
75. Montgomery, H.L.; Vaughan, R.C. *Multiplicative Number Theory I: Classical Theory*; Number 97 in Cambridge Studies in Advanced Mathematics, Cambridge University Press: Cambridge, 2007.
76. Slepian, D. Prolate Spheroidal Wave Functions, Fourier Analysis and Uncertainty—IV: Extensions to Many Dimensions. *Bell System Technical Journal* **1964**, *43*, 3009–3057.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© 2025 by the author(s). Distributed under a Creative Commons CC BY license.