

Article

Not peer-reviewed version

---

# Modified MOND Inertia and Covariant Vacuum State

---

[Yuanxin Li](#) \*

Posted Date: 20 May 2025

doi: 10.20944/preprints202504.2469.v4

Keywords: modified MOND inertia; nonlocal dynamics; covariant vacuum state



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

## Article

# Modified MOND Inertia and Covariant Vacuum State

Yuanxin Li

Independent Researcher; yuanxinli31@gmail.com

**Abstract:** In this paper, we investigate the inertia of macroscopic objects using a mean field approach. The dynamics are given by the equation of the Kuramoto-like form. The equilibrium solution of the equation is consistent with MOND theory. As a result, the state of the universe at large scales enters local dynamics in small systems. We also show that the vacuum energy density arising from zero-point fluctuations and symmetry breakings is acceleration-dependent. By requiring that the vacuum state is covariant, all acceleration-dependent vacuum energy should be regarded as non-physical.

**Keywords:** modified MOND inertia; nonlocal dynamics; covariant vacuum state

## 1. Introduction

The universe in its vast complexity harbors profound mysteries that challenge our understanding of fundamental physics. Observations of the flat rotation curves of galaxies reveal a striking discrepancy that galaxies rotate faster than can be accounted for by the visible matter alone, suggesting the presence of an invisible mass, known as dark matter. It is also hypothesized to explain a range of gravitational anomalies observed in gravitational lensing and the cosmic microwave background (CMB) [1]. Despite its success in accounting for these phenomena, the dark matter paradigm faces significant challenges. The lack of direct detection raises questions about its fundamental validity. The most widely accepted Cold Dark Matter (CDM) model struggles to account for observational discrepancies at smaller scales, such as the “missing satellites problem,” where the predicted number of dwarf galaxies around massive galaxies exceeds observations [2–4], and the “cusp-core problem,” where simulated dark matter halos exhibit central density cusps inconsistent with the flatter density profiles observed in some galaxies [5]. In addition, some observational data including rapid galaxy growth [6] and flat velocity curves extending beyond the expected virial radii of dark matter halos [7] also suggest inconsistencies with the CDM model. These issues imply that the CDM framework may need refinement or that our understanding of gravitational dynamics requires revision.

One such alternative theory is the Modified Newtonian Dynamics (MOND) proposed by Milgrom as a modification to Newtonian gravity at extremely low accelerations [8–10]. Although MOND struggles to account for phenomena at cosmological scales, such as the cosmic microwave background and large-scale structure formation, the simplicity and predictive power of MOND make it a valuable theoretical tool, prompting ongoing research into its foundations and potential extensions. Unlike the dark matter hypothesis, the MOND paradigm stipulates that the observed gravitational effects arise not from unseen mass but from a deviation in the law of gravity when accelerations fall below a critical threshold  $a_0 \approx 1.2 \times 10^{-10} \text{ m/s}^2$ . Milgrom’s law can be written as  $\mu(a/a_0)a = a_N$ , where  $a_N$  is the Newtonian acceleration produced by the visible matter,  $a$  is the true gravitational acceleration and the interpolating function  $\mu(x)$  satisfies  $\mu(x) \approx 1$  when  $x \gg 1$ , and  $\mu(x) \approx x$  when  $x \ll 1$ . In the deep-MOND regime ( $a \ll a_0$ ), MOND modifies the gravitational force law to scale inversely with distance rather than the square of the distance, effectively reproducing the flat rotation curves of galaxies and Tully-Fisher relation without invoking dark matter. MOND can be interpreted as a modification of gravity or inertia. In the modified gravity interpretation, Newton’s law of gravity should be modified at low accelerations, leading to a stronger gravitational effect than predicted by the usual inverse-square law. This approach has been formalized in theories such as the Tensor-Vector-Scalar (TeVeS) gravity [11] and bimetric theories [12,13], which attempts to provide a relativistic

extension of MOND [10]. Alternatively, the modified inertia interpretation suggests that the response of a body to a given force depends on the acceleration regime.

Given the empirical validations and elegant mathematical structure of general relativity, there is more potential in modified inertia as the basis for MOND because it seems to be less drastic. Inertia governs how objects respond to applied forces. However, the origin of inertia remains an open question in modern physics. Understanding the origin of inertia could provide critical insights into the foundations of the MOND theory. In classical mechanics, inertia is an intrinsic property of mass, yet its microscopic basis is not well understood. Some theories propose that inertia arises from interactions with the vacuum [14]. Einstein's theory of general relativity offers a partial explanation, suggesting that mass influences and is influenced by the local geometry of spacetime. Alternative perspectives, like Mach's principle, suggest that inertia is a relational property, emerging from an object's interaction with the global distribution of matter in the universe.

Clearly, the inertial mass of elementary particles primarily arises from the interaction with the Higgs field. For a charged particle, the interaction of particles with the electromagnetic field also contributes to inertia, known as electromagnetic mass. In general, we should consider all possible contributions. On the other hand, for macroscopic objects, various intricate interactions may contribute to the emergence of the inertial mass. A very noteworthy coincidence of MOND is that the value of  $a_0 \approx 1.2 \times 10^{-10} \text{ m/s}^2$  determined from galaxy dynamics is of the order of some acceleration constants of cosmological significance. It is of the same order as  $H_0$  ( $H_0$  is the Hubble constant) and  $(\Lambda/3)^{1/2}$  ( $\Lambda$  is the cosmological constant). This mysterious "cosmic coincidence" raises questions about its fundamental significance and potential cosmic connections. In this paper, we argue that the "cosmic coincidence" can be understood as a "synchronization" phenomenon at cosmological scales similar to the Kuramoto model [15,16]. A gravitational force that decays with distance is insufficient, as it would only result in local dynamics. However, considering the gravity including the cosmological constant, the fact that the interaction strength increasing with distance inevitably leads to the state of the universe at large scales entering local dynamics in small systems, such as galaxies. This nonlocal behavior is common in nonlinear dynamical systems. We will investigate the inertia of macroscopic objects and derive the modified acceleration formula. Upon using a mean field theory approach, it is possible to study a system with a large or infinite number of degrees of freedom. Thus we define a mean field characterized by the Unruh temperature to represent the average effect of all the contents of the universe. It may suggest a connection between gravity and thermodynamics. In some studies, gravity is regarded as an entropic force [17–20]. In my personal view, this only means that gravity can be described in another framework at the macroscopic level, whereas at microscopic scales gravity may still be quantum. Furthermore, emergent phenomena at the cluster and cosmological scales arise from the interactions, primarily gravity, of the universe's components. Thus, it is gravity that leads to emergent phenomena at larger scales rather than gravity itself originating from emergence.

Another concern of this study is the cosmological constant problem, which remain a theoretical puzzle that bridges cosmology and quantum field theory [21]. The problem originates from the interpretation of the cosmological constant  $\Lambda$  as the vacuum energy density. There are at least two sources for the vacuum energy. In the cosmological context, spontaneous symmetry breakings in the early universe may have induced phase transitions, potentially contributing to the vacuum energy density associated with the cosmological constant. Although one can always adjust the vacuum energy today to zero by tuning the parameter of the potential, it is not a very satisfactory method because the vacuum energy cannot be zero before and after the phase transition. In addition, in quantum field theory the vacuum is filled with quantum fluctuations contributing to a zero-point energy (ZPE). Since all energy gravitates, it is expected that the ZPE contribute to the cosmological constant. However, the theoretical prediction of the cosmological constant from quantum field theory contrasts with its observed value, giving rise to a discrepancy that spans over 120 orders of magnitude. One promising avenue for addressing the cosmological constant problem is supersymmetry (SUSY), a theoretical framework that posits a symmetry between fermions and bosons [22]. But experimental searches

at the Large Hadron Collider have yet to detect supersymmetric particles. In addition, the precise mechanism by which SUSY could resolve the cosmological constant problem remains elusive, as the required cancellations demand an extraordinary degree of fine-tuning in the SUSY-breaking sector. In this paper, we will calculate the vacuum energy for an accelerating observer. An interesting thing happens when we consider the Unruh effect which predicts that an accelerating observer in a vacuum will detect a thermal bath at a temperature proportional to its proper acceleration [23]. In Ref. [24], it has been shown that the electroweak phase transition (EWPT) occurs and the electroweak gauge symmetry can be restored when the acceleration exceeds the critical value as seen from the point of view of an accelerating observer. This means that the Unruh temperature is not just a formal artifact but a real temperature which can give rise to non-trivial thermal effects. We note that requiring that the vacuum state is covariant drives the theoretical value of the vacuum energy density to zero. Our analysis is based on the conservative assumption of maximal validity of quantum field theory and general relativity.

This paper is organized as follows. Section 2 is dedicated to the inertial mass of macroscopic objects. In Sec.3, we calculate the vacuum energy density by requiring that the vacuum state is covariant. Finally, in Sec.4 we summarize the main results obtained. For convenience, we use natural units with  $c = \hbar = k = 1$ .

## 2. Inertia of Macroscopic Objects and MOND

Let us consider a simplified model of the universe consisting of  $N$  masses coupled by gravity including a cosmological constant  $\Lambda$ . The fact that the couple strength increasing with distance inevitably leads to the state of the entire system at large scales entering local dynamics in small systems. Similar to the Kuramoto model [15,16], we can sum over the acceleration of all contents in the universe to obtain an average acceleration in a specific direction to describe the overall behavior of the universe. However, assuming the universe is isotropic, it is more appropriate to describe the dynamics of the system using a scalar related to acceleration, such as the magnitude or the square of the acceleration. In addition, 4-dimensional de Sitter spacetime can be embedded into a 5-dimensional flat Minkowski spacetime to generate a geometrical symmetry group  $SO(4,1)$  of the de Sitter spacetime. Here, we choose the Unruh temperature associated with acceleration as the degree of freedom. The dynamics expressed in terms of the temperature are given by the equation of the following Kuramoto-like form:

$$\dot{T}_i = \frac{1}{2\pi} F_i + \frac{\varepsilon}{N} \sum_{j=1}^N \frac{\Lambda r_{ij}}{3} \Gamma(T_j - T_i), \quad (1)$$

where the dot denotes the derivative with respect to time  $t$ ,  $T_i = \frac{1}{2\pi} (a^2 + H_0^2)^{1/2}$  is the Unruh temperature of the  $i$ -th mass seen by a local comoving accelerating observer in de Sitter spacetime,  $\varepsilon$  is the positive coupling strength,  $F_i$  represents the local external force applied to the  $i$ -th mass,  $r_{ij}$  is the distance and  $\Gamma(T_j - T_i)$  is a general coupling function for the interaction between the  $i$ -th and  $j$ -th masses. Here, we choose the coupling function  $\Gamma(T_j - T_i) = T_j - T_i$  because it naturally captures the acceleration differences between coupled masses in a simple and physically meaningful way. In addition, this linear term ensures that Equation (1) holds in the deep-MOND regime ( $a \ll a_0$ ). Let us define a mean field that represents the average effect of all the contents of the universe. Inertial observers in our universe with a positive cosmological constant detect a Gibbons–Hawking radiation with the temperature  $T_{\text{GH}} = H_0/2\pi$  [25]. Therefore, the mean temperature is  $T_{\text{GH}}$ . Every accelerated object appears to be subjected to a force that drives its acceleration toward zero. It is a “synchronization” phenomenon at cosmological scales and the overall behavior of the universe encoded in  $H_0$  enters local dynamics in small systems. Then Equation (1) becomes

$$\dot{T}_i = \frac{1}{2\pi} F_i + m(T_{\text{GH}} - T_i), \quad (2)$$

where  $m$  is the inertial mass. Since we have defined the inertial mass, external forces can be expressed in the form of Newton's second law, namely  $F_i = ma_N$  with  $a_N$  being the Newtonian expression for the acceleration. Thus Equation (2) can be written in terms of the acceleration as

$$\frac{a\dot{a}}{\sqrt{a^2 + H_0^2}} = m \left[ H_0 - (a^2 + H_0^2)^{1/2} \right] + ma_N. \quad (3)$$

The fixed point given by  $\dot{a} = 0$  represents stable solutions of Equation (3) and one arrives at

$$a_N = (a^2 + H_0^2)^{1/2} - H_0, \quad (4)$$

which leads to  $\mu(a/a_0)a = a_N$  with  $a_0 = 2H_0$ . The result is consistent with Milgrom's hypothesis [14]. It was interestingly noted that the local dynamics of small systems depend on the state of the universe at large scales and the Hubble constant varies during the evolution of the universe. Therefore, we should replace  $H_0$  with the varying Hubble parameter  $H(t)$  and Equation (4) becomes

$$\frac{H(t)\dot{H}(t)}{\sqrt{a^2 + H(t)^2}} = m \left[ H(t) - (a^2 + H(t)^2)^{1/2} \right] + ma_N. \quad (5)$$

One immediately obtains

$$a = a_N - \frac{H(t)\dot{H}(t)}{m\sqrt{a^2 + H(t)^2}}, \quad \text{for } a \gg 2H(t), \quad (6)$$

$$\frac{a^2}{2H(t)} = a_N - \frac{H(t)\dot{H}(t)}{m\sqrt{a^2 + H(t)^2}}, \quad \text{for } a \ll 2H(t). \quad (7)$$

It is expected that MOND has a different impact on the evolution of the universe during the period of decelerated expansion compared to the present universe. When considering the dynamics on short time scales, we can simply assume  $\dot{H}(t) = 0$ . The key point is that the current state at cosmic scales influences the behavior of local systems.

### 3. Acceleration-Dependent Vacuum Energy

For an accelerating observer the electroweak  $SU(2) \times U(1)$  gauge symmetry in the Standard Model is restored for acceleration larger than a critical value. The VEV  $v$  is given by [24]

$$v(a) = v_0 \sqrt{1 - \frac{a^2}{a_{EW}^2}}. \quad (8)$$

where  $v_0$  is the VEV for the inertial observer,  $a$  is the proper acceleration and  $a_{EW}$  is the critical proper acceleration of the EWPT. The second-order phase transition of the restoration of electroweak symmetry occurs at  $a_{EW}$  and for  $a > a_{EW}$ , we have  $v = 0$ . The elementary particles therefore acquire an acceleration-dependent mass which is

$$m(a) = m_0 \sqrt{1 - \frac{a^2}{a_{EW}^2}}, \quad (9)$$

where  $m_0$  is the mass of the elementary particle for the inertial observer. By introducing the Unruh-like temperature:

$$T_{EW} = \frac{a_{EW}}{2\pi} \quad (10)$$

and

$$T(a) = \frac{a}{2\pi}, \quad (11)$$



Equation (9) can be also written as

$$m(T) = m_0 \sqrt{1 - \frac{T^2}{T_{EW}^2}}, \quad (12)$$

where  $T_{EW} \sim 10^2$  GeV is the critical temperature of the EWPT. It turns out that all massive particles of Standard Model become massless for the local accelerating observer when the acceleration, or equivalently the temperature, exceeds the critical value.

The vacuum energy receives contributions from both zero-point fluctuations and symmetry breakings. We first calculate the ZPE. The ZPE density of a real free scalar field is given by

$$\rho_Z = \frac{1}{(2\pi)^3} \frac{1}{2} \int d^3k \omega(k) \quad (13)$$

with

$$\omega(k) = \sqrt{|k|^2 + m_0^2}, \quad (14)$$

where  $(\omega, k)$  is the four-dimensional momentum and  $m_0$  is the mass of the scalar field. Obviously, the integral is divergent in the ultraviolet region. The common method is to introduce an ultraviolet cut-off  $\Lambda_{UV}$  at the Planck scale, then one obtains  $\rho_Z \sim 10^{76}$  GeV<sup>4</sup>, which is larger than the observed value of vacuum energy density by a factor of  $10^{123}$ . But a straightforward but lengthy calculation leads to

$$\rho_Z = \frac{\Lambda_{UV}^4}{16\pi^2} + \frac{m_0^2 \Lambda_{UV}^2}{16\pi^2} + \frac{m_0^4}{64\pi^2} \ln \left( \frac{m_0^2 e^{\frac{1}{2}}}{4\Lambda_{UV}^2} \right) + \dots, \quad (15)$$

$$p = \frac{\Lambda_{UV}^4}{48\pi^2} - \frac{m_0^2 \Lambda_{UV}^2}{48\pi^2} - \frac{m_0^4}{64\pi^2} \ln \left( \frac{m_0^2 e^{\frac{7}{6}}}{4\Lambda_{UV}^2} \right) + \dots, \quad (16)$$

where  $p$  is the pressure. The Lorentz symmetry of the vacuum requires that the energy density and pressure satisfy the equation of state  $p = -\rho_Z$ . Notice that the first two terms of Equations (15) and (16) break Lorentz invariance and can be removed by local counterterms. Therefore, upon using a regularization scheme that preserves Lorentz symmetry of the vacuum, for any quantum field one arrives at the following expression for the ZPE density [26]

$$\rho_Z = \pm \frac{s m_0^4}{64\pi^2} \ln \left( \frac{m_0^2}{\mu^2} \right), \quad (17)$$

where  $\mu$  is the renormalization scale,  $s$  represents the number of polarization states and the signs  $\pm$  are associated with bosons and fermions respectively. The result can be generalized to any other interacting fields by simply replacing  $m_0$  with the renormalized mass  $m_R$ . We see that the expression is proportional to the mass of the particle to the power four and the massless particles do not contribute to the ZPE. This result is very different from the result obtained by imposing a Planck cut-off.

Another contribution to the cosmological constant comes from the symmetry breakings. We now calculate the vacuum energy produced by the EWPT at the classical level. We should also consider the QCD symmetry breaking ( $\sim 10^{-1}$  GeV) and other symmetry breakings at higher energy scales (e.g., the grand unification scale at  $10^{14}$  GeV and the Planck scale at  $10^{19}$  GeV). However, all these expressions take a similar form and the analysis parallels the electroweak case. The Higgs field consists of two complex scalar fields arranged into a doublet. After the EWPT, the field acquires a VEV and the corresponding vacuum energy density is  $\rho_{EW} = \lambda v^4 \sim 10^8$  GeV<sup>4</sup> with  $\lambda$  being the coupling constant describing the self-interaction of Higgs fields. In addition to being inconsistent with observational data, such a large vacuum energy density corresponding to a large cosmological constant would also produce a high Gibbons–Hawking temperature, thereby triggering a phase transition. From Equation (8), we see that the vacuum energy density of the EWPT must satisfy the equation:

$$\rho_{EW} = \rho_0 \left( 1 - \frac{T_h^2}{T_{EW}^2} \right)^2 \theta(T_{EW} - T_h), \quad (18)$$

where  $\rho_0$  is the vacuum energy density in the absence of Gibbons–Hawking radiation,  $\theta(x)$  is the Heaviside step function,  $T_{EW} \sim 10^2$  GeV is the EWPT temperature and  $T_h = \frac{1}{2\pi}(8\pi G\rho_s/3)^{1/2}$  is the Gibbons–Hawking temperature produced by the huge vacuum energy density  $\rho_s$  of symmetry breakings. When the vacuum energy density exceeds  $10^{10}$  GeV<sup>4</sup>, the broken electroweak symmetry is restored and  $\rho_{EW}$  vanishes. It can drive  $\Lambda$  back to zero, even when the vacuum energy density experiences large disturbances up to the Planck scale because large disturbances will lead to a restoration of the symmetry.

It is worth noticing that all the vacuum energy is acceleration-dependent and hence observers with different accelerations will measure different vacuum energy when quantum effects enter the stage. The key point is that the vacuum energy derived from zero-point fluctuations and symmetry breakings is not real if we require the vacuum state to be covariant. Therefore, for a potential covariant theory of quantum gravity, we should define a vacuum that is invariant for all observers and arbitrary coordinate transformations. Although the inertial mass arising from the interaction between the particle and the Higgs field depends on the acceleration and vanishes for an observer exceeding the acceleration threshold, we can extract the covariant part of the mass arising from other interactions, such as the electromagnetic interaction. If we interpret the current cosmological constant  $\Lambda$  as the vacuum energy density  $\rho_{vac} \sim 10^{-47}$  GeV<sup>4</sup> and assume that  $\Lambda$  arises from the covariant part of the electromagnetic mass, using Equation (17), the electromagnetic mass of a charged particle such as an electron for an inertial observer is approximately  $10^{-3}$  eV.

#### 4. Conclusions

In this paper, we investigate the inertia of macroscopic objects with a huge number of degrees of freedom. The system is drastically simpler by using a mean field theory approach. Equation (2) can be generalized to the framework of general relativity, where the system is characterized by the curvature rather than temperature. The LHS of Equation (2) represents the derivative of the connection and can thus be replaced by a tensor  $K_{\mu\nu}$  related to the true curvature of spacetime. The external force  $F_i$  is replaced by the usual Einstein tensor  $G_{\mu\nu}$  related to the visible matter content. The second term on the RHS of Equation (2) is replaced by a tensor  $C_{\mu\nu}$  representing nonlocal effects. As in bimetric theory,  $C_{\mu\nu}$  can be constructed by introducing an auxiliary metric. The rank-2 tensor  $C_{\mu\nu}$  plays the role of dark matter. It is a geometric version of the dynamics describing “synchronization” phenomena. We then calculate the vacuum energy density based on quantum field theory. It turns out that the vacuum energy is acceleration-dependent when quantum effects enter the stage. Here, we calculate the vacuum energy density arising from symmetry breaking at the tree level. If quantum corrections are considered, one only needs to replace it with the Coleman–Weinberg effective potential [27]. However, the result is the same because quantum corrections also depend on acceleration. Since we are considering gravitational effects, by requiring a covariant vacuum as demanded by general relativity, all acceleration-dependent energy should be regarded as non-physical. An intriguing conjecture is that the current cosmological constant may arise from the covariant electromagnetic mass and hence the electromagnetic mass of a charged particle such as an electron is approximately  $10^{-3}$  eV. However, this guess needs a further exploration.

#### References

1. Gianfranco Bertone and Dan Hooper. History of dark matter. *Reviews of Modern Physics*, 90(4):045002, 2018.
2. Guinevere Kauffmann, Simon DM White, and Bruno Guiderdoni. The formation and evolution of galaxies within merging dark matter haloes. *Monthly Notices of the Royal Astronomical Society*, 264(1):201–218, 1993.
3. Anatoly Klypin, Andrey V Kravtsov, Octavio Valenzuela, and Francisco Prada. Where are the missing galactic satellites? *The Astrophysical Journal*, 522(1):82, 1999.

4. Ben Moore, Sebastiano Ghigna, Fabio Governato, George Lake, Thomas Quinn, Joachim Stadel, and Paolo Tozzi. Dark matter substructure within galactic halos. *The Astrophysical Journal*, 524(1):L19, 1999.
5. David J Sand, Tommaso Treu, and Richard S Ellis. The dark matter density profile of the lensing cluster ms 2137–23: a test of the cold dark matter paradigm. *The Astrophysical Journal*, 574(2):L129, 2002.
6. Stacy S McGaugh, James M Schombert, Federico Lelli, and Jay Franck. Accelerated structure formation: The early emergence of massive galaxies and clusters of galaxies. *The Astrophysical Journal*, 976(1):13, 2024.
7. Tobias Misteale, Stacy McGaugh, Federico Lelli, James Schombert, and Pengfei Li. Indefinitely flat circular velocities and the baryonic tully–fisher relation from weak lensing. *The Astrophysical Journal Letters*, 969(1):L3, 2024.
8. Mordehai Milgrom. Mond—theoretical aspects. *New Astronomy Reviews*, 46(12):741–753, 2002.
9. Mordehai Milgrom. Mond—a pedagogical review. *arXiv preprint astro-ph/0112069*, 2001.
10. Benoit Famaey and Stacy S McGaugh. Modified newtonian dynamics (mond): observational phenomenology and relativistic extensions. *Living reviews in relativity*, 15:1–159, 2012.
11. Jacob D Bekenstein. Relativistic gravitation theory for the modified newtonian dynamics paradigm. *Physical Review D—Particles, Fields, Gravitation, and Cosmology*, 70(8):083509, 2004.
12. Christopher J Isham and D Storey. Exact spherically symmetric classical solutions for the f- g theory of gravity. *Physical Review D*, 18(4):1047, 1978.
13. Mordehai Milgrom. Bimetric mond gravity. *Physical Review D—Particles, Fields, Gravitation, and Cosmology*, 80(12):123536, 2009.
14. Mordehai Milgrom. The modified dynamics as a vacuum effect. *Physics Letters A*, 253(5-6):273–279, 1999.
15. Juan A Acebrón, Luis L Bonilla, Conrad J Pérez Vicente, Félix Ritort, and Renato Spigler. The kuramoto model: A simple paradigm for synchronization phenomena. *Reviews of modern physics*, 77(1):137–185, 2005.
16. ME Morrison. A quantum kuramoto model. *Bachelor thesis, University of Otago, Dunedin, New Zealand*, 2013.
17. Erik Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):1–27, 2011.
18. Chiu Man Ho, Djordje Minic, and Y Jack Ng. Cold dark matter with mond scaling. *Physics Letters B*, 693(5):567–570, 2010.
19. FR Klinkhamer and M Kopp. Entropic gravity, minimum temperature, and modified newtonian dynamics. *Modern Physics Letters A*, 26(37):2783–2791, 2011.
20. Chiu Man Ho, Djordje Minic, and Y Jack Ng. Quantum gravity and dark matter. *International Journal of Modern Physics D*, 20(14):2887–2893, 2011.
21. Jerome Martin. Everything you always wanted to know about the cosmological constant problem (but were afraid to ask). *Comptes Rendus Physique*, 13(6-7):566–665, 2012.
22. Stephen P Martin. A supersymmetry primer. In *Perspectives on supersymmetry II*, pages 1–153. World Scientific, 2010.
23. William G Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870, 1976.
24. Antonio Dobado. Brout-englert-higgs mechanism for accelerating observers. *Physical Review D*, 96(8):085009, 2017.
25. Gary W Gibbons and Stephen W Hawking. Cosmological event horizons, thermodynamics, and particle creation. *Physical Review D*, 15(10):2738, 1977.
26. Jurjen F Koksma and Tomislav Prokopec. The cosmological constant and lorentz invariance of the vacuum state. *arXiv preprint arXiv:1105.6296*, 2011.
27. Sidney Coleman and Erick Weinberg. Radiative corrections as the origin of spontaneous symmetry breaking. *Physical Review D*, 7(6):1888, 1973.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.