

Article

Not peer-reviewed version

---

# Variation of Fatty Acids, Lignans, and Tocols of Sesame (*Sesamum indicum*) Genotypes

---

[Cemal Kurt](#) \*

Posted Date: 24 January 2025

doi: 10.20944/preprints202501.1861.v1

Keywords: sesame; fatty acid; sesamol; sesamin; tocols



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

## Article

# Variation of Fatty Acids, Lignans, and Tocols of Sesame (*Sesamum indicum*) Genotypes

Cemal KURT

Department of Field Crops, Faculty of Agriculture, Çukurova University, Adana, Türkiye; ckurt@cu.edu.tr

**Abstract:** Sesame is one of the neglected crops. As a result, there are not many studies on the fatty acid, tocol and lignan composition of sesame genotypes. This study aimed to evaluate the diversity in fatty acid, lignan and tocol composition among different sesame accessions from Türkiye and 12 other countries. Fifty sesame genotypes were investigated under field conditions in Adana, Türkiye. The sesame genotypes varied widely in their fatty acid, lignan and tocol compositions. The content of palmitic acid, stearic acid, oleic acid, linoleic acid, linolenic acid,  $\alpha$ -tocopherol,  $\gamma$ -tocopherol,  $\alpha$ -tocotrienol, total tocopherol, sesamin, sesamol, sesamol varied between 8.39-10.11%, 4.19-5.36%, 35.83-43.49%, 40.64-48.8%, 0.28-0.49%, 1.08-2.79  $\mu\text{g/g}$ , 41.5-75.74  $\mu\text{g/g}$ , 0.96-1.8  $\mu\text{g/g}$ , 42.93-76.92  $\mu\text{g/g}$ , 2.34-14.31 mg/g, 0.23-1.81 mg/g, 0-0.59 mg/g respectively. While the sesamin content of the sesame genotypes used in the study was higher than previously reported values, it was determined that the sesamol content was at very low levels. The sesame genotypes are rich in sesamin content and can be used as parents in breeding programs. Sesamol was not detected in almost all of the cultivars from Türkiye. In addition, it was determined that the  $\gamma$ -tocopherol contents of these cultivars were below the average value for Turkish cultivars. The same applies to sesamin and sesamol. The results obtained from the study showed that the lignan and tocol composition of the cultivars produced in Türkiye were not at the desired levels. Therefore, improving the lignan and tocol composition of the cultivars in the breeding programs designed to obtain new cultivars will benefit consumers in Türkiye, which consumes sesame extensively.

**Keywords:** sesame; fatty acid; sesamol; sesamin; tocols

## 1. Introduction

Sesame is one of the oldest and most important oil crops known. The cultivation of sesame It dates back to 3500 years. During the last period, sesame cultivation was carried out in more than 60 countries, but it is consumed in different amounts in almost all countries [1]. The most important reason for the increasing interest in sesame seeds and oil is the nutritional properties of sesame and its positive effects on human health. Sesame seeds are used in bakery products and also made into tahini and halva. Also, it is used in industries such as pesticides, paint, medicine, cosmetics, perfumery, and soap. Sesame has anti-carcinogenic, anti-tumor, anti-aging, and anti-hypertension properties. Sesame seeds contain 40-60% oil, 18-25% protein, 13.5% carbohydrates and 5% ash. In addition, sesame seeds are rich in elements such as Ca, Fe, Zn, and Se [1]. It also contains sesame-specific lignans such as sesamin, sesamol and sesaminol. In addition to lignans, sesame seeds, which have antioxidative and health-promoting properties, also contain multiple tocopherol homologs such as  $\alpha$ ,  $\delta$ ,  $\gamma$ , and tocotrienols. The sesame seeds are commonly known as the 'Queen of oil seeds', probably due to their high oil yield, pleasant taste, high nutritional and therapeutic value, and resistance to oxidation and rancidity [2]

Unsaturated fatty acids make up about 85% of the oil in sesame seeds. However, sesame oil is resistant to oxidative degradation when compared to other vegetable oils. Its superior anti-oxidative property is not only due to its high  $\gamma$ -tocopherol content but also due to the sesame-specific lignans it contains such as sesamol, sesamin, and sesamol. These lignans have numerous health benefits

such as anti-carcinogenic, anti-inflammatory, anti-aging, anti-neuropathic, antihypertensive, cardioprotective and hypocholesterolemia [3].

The richness of unsaturated fatty acids in sesame oil makes it a very high-quality oil in terms of nutrition and health. It is directly related to cardio-protective, hypo-lipidemic, anti-atherogenic and anti-inflammatory effects [4].

Tocopherols are natural antioxidants that hamper oil oxidation. In addition to having an important nutritional function for humans as a source of vitamin E, tocopherols absorb free radicals and can prevent disease [5]. Vitamin E has been recognized as an important dietary component with anti-aging function. It is known that this feature of vitamin E is due to the  $\alpha$ -tocopherol in it. The highest tocopherol content in sesame seeds belongs to  $\gamma$ -tocopherol, and  $\alpha$ -tocopherol is present in tiny amounts. However, studies in rats have shown that  $\gamma$ -tocopherol in sesame seed exhibits vitamin E activity equal to that of  $\alpha$ -tocopherol through a synergistic interaction with sesame seed lignans [6].

Although sesame is generally grown in developing countries, it is consumed in different amounts in almost every country. Therefore, determining the nutritional content of sesame genotypes is important in providing necessary nutritional information to consumers. In addition, the characterization of genetic resources in terms of these features is of great importance in providing genetic material that breeders can use in the development of varieties with higher nutritional content. [1,7,8]. Previous studies on sesame were generally limited to the quantification of lignans alone or oil or tocopherols alone, lignans with fatty acids, or tocopherols with fatty acids [9–15]. Regarding the fatty acids, tocopherols and lignan contents of sesame, no reports have been found examining these three oil traits together. Hence, the present study aimed to evaluate the extent of variability for fatty acids, lignans tocopherols, and tocotrienols (tocopherols) in the 50 different sesame genotypes originating from Türkiye and 12 different countries.

## 2. Material and Methods

### 2.1. Plant Material

Fifty sesame genotypes were used as plant material in this study. All sesame genotypes were sown at the Experimental Farm of Cukurova University, in Adana (Türkiye). Before planting, 200 kg/ha of diammonium phosphate (36 kg/ha N, 92 kg/ha P) fertilizer was applied, while ammonium nitrate (33 %N) at the rate of 200 kg/ha was applied before the first irrigation. Sesame seeds were sown in the third week of June. The accessions were grown in two-row plots of 3 m row length with a row spacing of 70 cm and intra-row spacing of 15 cm. Thinning was carried out after 25 days of sowing to secure one plant per 15 cm. Sprinkler irrigation was established immediately after sowing and thereafter used as needed. Weeding was carried out by hand and no herbicides were applied during the experiment. All the plants were harvested by hand in the last week of September, 2020 [1].

### 2.2. Oil Extraction and GC Analysis

The samples of 50 sesame genotypes were subjected to oil samples extracted by cold press (Karaerler NF 500). An oil sample of 500 mg was dissolved in 2 ml isooctane followed by 1.5 ml of 0.5 M methanolic NaOH (99.6%, Sigma). The tube was then vortexed and held in boiling water for 7 min and allowed to cool to room temperature. Two ml of BF<sub>3</sub> (99.99% Boron trifluoride, Sigma) were added, vortexed, and held in boiling water for 5 min and allowed to come down to room temperature. The tube was vortexed after adding 5 ml NaCl (Merck), centrifuge at 4,000 rpm for 10 min. The supernatant was used for GC analyses (AOAC 1984).

The fatty acid (FA) composition was analyzed using a GC Clarus 500 with auto sampler (Perkin Elmer, USA) equipped with a flame ionization detector and a fused silica capillary SGE column (30 m • 0.32 mm, ID • 0.25  $\mu$ m, BP20 0.25  $\mu$ m, USA). The oven temperature was brought to 140 °C for 5 min, then raised to 200 °C at a rate of 4 °C/min and to 220 °C at a rate of 1 °C/min, while the injector and the detector temperatures were set at 220 °C and 280 °C, respectively [16].

2.3. Analysis of Tocols and Lignans

Tocols and lignans were analysed in the oil samples extracted by cold press (Karaerler NF 500) with a High Pressure Liquid Chromatography (HPLC) device (Agilent Technologies 1200) with UV detector. HPLC conditions in the analysis of tocols; Column: Inertsil NH2 250mm x 4.6mm - 5µm, Mobile phase: n-hexane/acetic acid/IPA (1000mL/5mL/6mL), Column temperature: 20°C, Wavelength: 296 nm, Flow rate: 1 mL, Injection volume: 10 µL (TS ISO, 2016). HPLC conditions in the analysis of lignans; Column: Prodigy ODS3 100A 250mm x 4.6mm - 5µm, Mobile phase: methanol/water (70mL/30mL), Column temperature: 30°C, Wavelength: 290 nm, Flow rate: 1 mL, Injection volume: 10 µL ([17]Shi et al. 2018).

All chemicals used in the analyses are Merk brand, and the standards are Sigma Aldrich (Sesamolin Product Number: SMB00701, Sesamin Product Number: 59867, Sesamol Product Number: S3003) and ChromaDex (Tocotrienol and Tocopherol Mixture Lot Number: 00020329-00569).

2.4. Statistical Analysis

The statistical software XLSTAT (www.xlstat.com) was employed to examine the mean, maximum, minimum, standard deviation, and correlation pertaining to all seven traits under investigation. Additionally, Principal Component Analysis (PCA) was conducted utilizing the same software. A biplot graph was generated using XLSTAT.

3. Results and Discussion

3.1. Fatty Acid Composition

Fatty acid composition is one of the important features used to indicate the nutritional value of oils. The sesame seed oil contains high amounts of monounsaturated fatty acids such as oleic acid (35-54%) and polyunsaturated fatty acids such as linoleic acid (39-59%) ([18] Langyan et al, 2022). The fatty acid compositions varied significantly among the tested sesame accessions (Table 1). USA 2 genotype at 43.49% exhibited the highest oleic acid (C18:1) content while Pakistan 2 at 35.83% exhibited the lowest oleic acid content in the study. The linoleic acid (C18:2) contents varied between 40.64-48.8% with a mean of 43.29%. When the average oleic (40.61%) and linoleic acid (43.29%) contents of the studied sesame genotypes were examined, it was found that the content of linoleic acid was higher. Linoleic acid content in earlier reports by Morris et al., (2021) [19], and Kurt (2018) [16] was higher, while lower oleic acid contents than that found in the present study. Teklu et al., (2022) [20] found oleic acid percentage ranged between 36.76-42.85%, while linoleic acid percentage varied from 36.56-41.69%. Diyarbakir-Bismil genotype at 0.49% exhibited the highest linolenic acid content, while Diyarbakir-Bismil- Bakacak genotype at 0.28% had the lowest linolenic acid content.

Palmitic acid was the predominant fatty acid with stearic acid in sesame oil and varied between 8.39-10.11% and 4.19-5.36% with averages of 9.37 and 4.72% respectively. Denizli-Acipayam and Diyarbakir Ergani Ziyaret Köy have higher palmitic and stearic acid content than averages. Kurt (2015) [21] reported that palmitic and stearic acid contents varied between 8.63-9.77% and 4.93-5.94% respectively. Were et al. (2006) [22] reported that average palmitic acid content (8.24%) was lower, while stearic acid content (4.89%) was higher than the present study.

Table 1. Fatty acids composition of sesame genotypes.

Genotypes		C16:0	C16:1	C18:0	C18:1	C18:2(ω-6)	C18:3(ω-3)	C20:0
Sanliurfa-Siverek-	Siverek-	9.34	0.14	4.89	41.58	42.59	0.31	0.52
Basdegirmen 1 <sup>T</sup>								
Sanliurfa-Siverek-		9.47	0.14	4.96	41.32	42.43	0.33	0.06
Basdegirmen 2 <sup>T</sup>								

Denizli-Dazkiri <sup>T</sup>	9.72	0.13	4.70	38.34	45.56	0.32	0.55
Denizli-Acipayam <sup>T</sup>	9.55	0.13	5.08	42.95	<b>40.64</b>	0.32	0.57
Sanliurfa- Bozova-Hasmersar <sup>T</sup>	9.69	0.14	4.69	39.84	44.05	0.33	0.53
Antalya-Finike <sup>T</sup>	9.90	0.14	4.53	38.59	45.31	0.32	0.51
Sanliurfa-Akziyaret <sup>T</sup>	9.40	0.14	4.55	42.27	42.12	0.35	0.50
Aydın-Çine <sup>T</sup>	9.48	0.13	4.81	41.05	43.02	0.34	0.55
Adıyaman-Gölbaşı <sup>T</sup>	9.27	0.14	4.91	39.71	44.52	0.28	0.55
Antalya-Konyaaltı <sup>T</sup>	9.50	0.13	4.55	41.03	43.33	0.33	0.51
Adana-Ceyhan <sup>T</sup>	9.61	0.14	4.66	41.66	42.45	0.31	0.50
Mardin <sup>T</sup>	9.93	0.14	4.60	41.08	42.70	0.36	0.51
Diyarbakir-Bismil- Bakacak <sup>T</sup>	10.07	0.14	4.55	40.01	43.82	<b>0.28</b>	0.50
Diyarbakir- Bismil Yukari Harim Köy <sup>T</sup>	9.82	0.14	4.49	38.90	45.19	0.31	0.51
Adana-Yumurtalik <sup>T</sup>	<b>10.11</b>	<b>0.15</b>	4.48	40.13	43.66	0.34	0.51
Batman <sup>T</sup>	9.80	0.13	5.06	42.29	41.31	0.31	0.55
İcel-Tarsus <sup>T</sup>	9.50	0.13	4.60	38.82	45.48	0.30	0.52
Diyarbakir- Ergani- Dağarası Köyü <sup>T</sup>	9.47	0.14	4.68	41.83	42.46	0.29	0.51
Diyarbakir- Ergani Ziyaret Köy <sup>T</sup>	10.08	0.14	4.89	41.29	42.13	0.28	0.56
Greece 1	9.12	0.11	4.78	40.93	43.47	0.33	0.50
Pakistan 1	9.53	0.11	<b>4.19</b>	37.88	46.90	0.34	0.47
Pakistan 2	9.74	0.12	4.25	<b>35.83</b>	48.69	0.34	0.48
Iran	9.56	0.12	4.23	35.97	<b>48.80</b>	0.33	0.44
Diyarbakir-Bismil <sup>T</sup>	9.69	0.13	4.51	39.72	44.27	<b>0.49</b>	0.51
Baydar-2001 <sup>C</sup>	9.32	0.12	4.77	42.12	41.91	0.44	0.53
Batem-Uzun <sup>C</sup>	9.27	0.13	4.64	41.65	42.48	0.40	0.49
Batem-Aksu <sup>C</sup>	8.86	0.12	4.60	42.56	41.97	0.45	0.47
Boydak <sup>C</sup>	9.09	0.11	4.88	41.90	41.61	0.38	0.49
Gölmarmara <sup>C</sup>	9.11	0.12	4.76	38.99	43.22	0.37	0.50
Tanas <sup>C</sup>	9.48	0.12	4.86	40.53	42.71	0.44	0.52
Hatipoğlu <sup>C</sup>	9.06	0.11	4.78	42.05	41.35	0.44	0.48
Tan-99 <sup>C</sup>	9.30	0.12	4.66	40.63	41.19	0.42	0.50
Özberk <sup>C</sup>	9.22	0.12	4.68	40.95	42.58	0.40	0.50
Mozambik	9.05	0.11	4.74	40.61	43.33	0.36	0.51
Israel	9.17	0.11	4.88	38.89	44.28	0.35	0.54
Pakistan 3	9.21	0.11	<b>5.36</b>	41.94	41.69	0.34	0.58
Greece 2	9.20	0.11	4.90	41.97	40.89	0.37	0.51
Russia 1	9.03	0.12	4.72	39.43	43.26	0.34	0.51
Russia 2	8.78	0.12	4.62	41.27	43.24	0.36	0.52
Libya	8.75	0.11	4.75	39.75	45.01	0.32	0.53



Egypt	8.91	0.11	5.11	39.75	43.76	0.32	0.58
Iraq	8.81	0.11	4.81	40.89	43.70	0.35	0.55
S. America	10.08	0.12	4.63	39.32	44.25	0.35	0.54
USA 1	9.04	0.12	4.63	41.98	42.33	0.41	0.55
Afghanistan	8.39	0.11	4.88	42.29	42.60	0.37	0.55
Manisa-Selendi <sup>T</sup>	8.75	0.12	4.85	42.92	41.45	0.39	0.54
USA 2	8.90	0.12	4.61	43.49	40.99	0.43	0.53
Kepsut-99 <sup>C</sup>	9.32	0.12	4.60	41.67	42.38	0.48	0.54
Sarisu <sup>C</sup>	9.47	0.13	4.67	40.48	43.36	0.45	0.53
Osmanlı-99 <sup>C</sup>	9.75	0.13	4.82	39.57	43.82	0.46	0.55
Max.	10.11	0.15	5.36	43.49	48.8	0.49	0.58
Min.	8.39	0.11	4.19	35.83	40.64	0.28	0.06
Ave.	9.37	0.12	4.72	40.61	43.29	0.36	0.51

<sup>C</sup>: Cultivar; <sup>T</sup>: Türkiye

3.2. Tocols and Lignans Composition

Edible oils mainly consist of fatty acids in the form of triacylglycerols, which produce energy for the human body during metabolism. In addition, edible oils are sources of minor compounds such as tocopherols, tocotrienols, or both (tocols), carotenoids, and phytosterols [23]. Interest in tocol studies has increased dramatically in recent years, most likely to increase awareness of the health effects of individual food items and diets [24]. Tocol-related compounds, tocopherols ( $\alpha$ ,  $\beta$ ,  $\delta$ ,  $\gamma$ ) and tocotrienols, belonging to the vitamin E family, are important bioactive components in vegetable oils, mainly due to their antioxidative effects [25].

Sesame is one of the neglected crops. As a result, there are not many studies on the tocol content of sesame genotypes. Sesame seed total tocol contents vary from one germplasm to another. As can be seen in Table 2, the mean of  $\alpha$ -tocopherol content ranged across studied sesame genotypes was 1.63  $\mu\text{g/g}$ , varying from 1.08 to 2.79  $\mu\text{g/g}$ . The highest value belongs to the Turkish cultivar, Batem-Aksu, and Adıyaman-Gölbaşı exhibited the lowest  $\alpha$ -tocopherol content.  $\alpha$ -tocotrienol content varied between 0.96-1.8  $\mu\text{g/g}$ , Türkiye-Aydın-Çine and Türkiye-İçel-Tarsus respectively.  $\gamma$ -tocopherol shows a higher antioxidant capacity compared to  $\alpha$ -tocopherol [26], which is important for the industrial use of sesame oil. According to Jiang et al., (2001) [27],  $\gamma$ -tocopherol is the most abundant form in sesame seeds, the predominant form of vitamin E in human and animal tissues, and may be important for human health compared to  $\alpha$ -tocopherol, which is the main form in supplements.  $\gamma$ -tocopherol content ranged from 41.5 to 75.74  $\mu\text{g/g}$  with an average of 52.43  $\mu\text{g/g}$  in studied sesame genotypes seeds.

Table 2. Tocol composition of sesame genotypes.

Genotypes		$\alpha$ - Tocopherol ( $\mu\text{g/g}$ oil)	$\alpha$ -Tocotrienol ( $\mu\text{g/g}$ oil)	$\gamma$ -Tocopherol ( $\mu\text{g/g}$ oil)	Total Tocopherol ( $\mu\text{g/g}$ oil)
Sanliurfa- Basdegirmen 1 <sup>T</sup>	Siverek-	1.52	1.40	58.68	60.20
Sanliurfa-Siverek- Basdegirmen 2 <sup>T</sup>		1.43	1.34	41.50	42.93
Denizli-Dazkiri <sup>T</sup>		1.34	1.20	53.16	54.50

Denizli-Acipayam <sup>T</sup>	1.55	1.35	50.43	51.97
Sanliurfa- Bozova-Hasmersar <sup>T</sup>	1.32	1.03	48.51	49.83
Antalya-Finike <sup>T</sup>	1.30	1.02	52.93	54.23
Sanliurfa-Akziyaret <sup>T</sup>	1.58	1.16	51.24	52.82
Aydın-Çine <sup>T</sup>	1.18	<b>0.96</b>	46.65	47.82
Adıyaman-Gölbaşı <sup>T</sup>	<b>1.08</b>	0.98	51.17	52.25
Antalya-Konyaaltı <sup>T</sup>	1.41	1.04	53.44	54.84
Adana-Ceyhan <sup>T</sup>	1.53	1.33	50.74	52.26
Mardin <sup>T</sup>	1.26	1.18	52.11	53.37
Diyarbakir-Bismil- Bakacak <sup>T</sup>	1.21	1.26	53.02	54.22
Diyarbakir- Bismil Yukari Harim Köy <sup>T</sup>	1.15	1.02	58.45	59.60
Adana-Yumurtalik <sup>T</sup>	1.37	1.36	58.41	59.78
Batman <sup>T</sup>	1.43	1.18	59.10	60.52
İcel-Tarsus <sup>T</sup>	1.18	<b>1.80</b>	<b>75.74</b>	<b>76.92</b>
Diyarbakir- Ergani- Dağarası Köyü <sup>T</sup>	1.47	1.48	73.34	74.81
Diyarbakir- Ergani Ziyaret Köy <sup>T</sup>	1.35	1.33	66.45	67.80
Greece 1	2.07	1.63	53.03	55.10
Pakistan 1	1.73	1.31	62.16	63.89
Pakistan 2	1.27	1.48	60.66	61.93
Iran	1.55	1.37	56.64	58.19
Diyarbakir-Bismil <sup>T</sup>	2.49	1.46	52.87	55.37
Baydar-2001 <sup>C</sup>	2.14	1.26	46.15	48.29
Batem-Uzun <sup>C</sup>	1.82	1.31	47.37	49.18
Batem-Aksu <sup>C</sup>	<b>2.79</b>	1.24	42.44	45.23
Boydak <sup>C</sup>	1.79	1.36	47.68	49.47
Gölmarmara <sup>C</sup>	1.66	1.32	48.90	50.56
Tanas <sup>C</sup>	1.57	1.29	44.45	46.01
Hatipoğlu <sup>C</sup>	2.31	1.39	45.58	47.89
Tan-99 <sup>C</sup>	1.73	1.47	48.49	50.22
Özberk <sup>C</sup>	1.73	1.30	48.20	49.94
Mozambik	1.56	1.30	49.79	51.35
Israel	1.56	1.25	53.64	55.20
Pakistan 3	1.54	1.35	47.43	48.97
Greece 2	1.63	1.31	49.32	50.95
Russia 1	1.75	1.39	49.22	50.96
Russia 2	1.57	1.37	50.48	52.05
Libya	1.66	1.25	47.78	49.43
Egypt	1.37	1.27	49.47	50.83
Iraq	1.84	1.44	59.27	61.11
S. America	1.56	1.24	56.11	57.67
USA 1	1.55	1.33	54.43	55.97

<b>Afghanistan</b>	1.58	1.40	49.83	51.40
<b>Manisa-Selendi<sup>T</sup></b>	1.50	1.32	45.64	47.14
<b>USA 2</b>	1.86	1.29	48.90	50.76
<b>Kepsut-99<sup>C</sup></b>	2.63	1.47	46.65	49.28
<b>Sarisu<sup>C</sup></b>	1.93	1.35	50.97	52.90
<b>Osmanlı-99<sup>C</sup></b>	2.18	1.53	52.76	54.94
<b>Max.</b>	2.79	1.63	75.74	76.92
<b>Min.</b>	1.08	0.96	42.44	42.93
<b>Ave.</b>	1.63	1.31	52.43	54.06

<sup>C</sup>: Cultivar; <sup>T</sup>: Türkiye.

Lignans in sesame seeds have attracted the attention of nutritional scientists and healthcare professionals for their health-promoting activities such as lowering blood sugar and cholesterol levels, preventing cardiovascular diseases, Alzheimer's and cancer, and relieving post-menopausal syndrome [28,29]. The ability of sesamin to suppress tumor growth [30]. Yokota et al., 2007 [31] suggests that sesamin could even be developed as a therapeutic agent. Sesame oil and lignans are components of creams and body oils [32,33]. In addition to nutritional, cosmetic, and health-promoting use, sesame lignans (especially sesamol) are used as an additive in insecticides due to their synergistic effect and low-grade oil is used for the manufacture of soaps, perfumes, and paints in the industry. There was significant variation in lignan contents among the sesame genotypes as shown in Table 3. The mean of sesamin content ranged across studied sesame genotypes was 6.65 mg/g, varying from 2.34 to 14.31 mg/g. The highest value belongs to the Sanliurfa-Siverek-Basdegirmen 1, the Turkish cultivar Osmanlı-99 exhibited the lowest sesamin content. Sesamin content values range in the present study were higher than those reported by Kim et al. 2006 [34] (0.38-5.12 mg/g), Wu et al. 2017 (1.7-5.1 mg/g) [35], Zhu et al. 2018 (2.16-7.37 mg/g) [36], Dar et al. 2019 (2.1-5.98 mg/g) [37] and Xu et al. 2021 (0.33-7.52 mg/g) [38]. While the highest value of sesamol content, another major lignan, was obtained from the Israel genotype (1.81 mg/g), the lowest value was obtained from the Turkish variety Osmanlı-99 (0.23 mg/g). The range of sesamol content values in this study was found to be considerably lower than the values in previously reported studies [Moazzami et al. 2007 (1.67-8.04 mg/g) [39]; Wang et al. 2012 (0.93-6.96 mg/g) [40]; Dar et al. 2019 (1.52-3.76 mg/g) [37]]. While the highest sesamol value was obtained from the Turkish genotype Denizli-Acipayam, sesamol could not be detected in 15 of the 50 genotypes studied. These are 8 of the 15 genotypes whose sesamol could not be detected are Turkish cultivars. Other Turkish cultivars used in the study, Tanas, contain 0.01 µg/g and Kepsut-99 contains 0.04 mg/g sesamol. When the present study is compared with previous studies, the sesamol ratio we detected in our results is at very low levels [14,37,41]. Reports on the variability of the sesame lignans contents in sesame seeds indicated that these lignans are affected by genetics, cultivars, origin, agronomic conditions and practices, environmental stresses, and other seed traits such as seed color [37,40–43].

**Table 3.** Lignans composition of sesame genotypes.

<b>Genotypes</b>	<b>Sesamol (mg/g oil)</b>	<b>Sesamin (mg/g oil)</b>	<b>Sesamol (mg/g oil)</b>
<b>Sanliurfa- Siverek- Basdegirmen 1<sup>T</sup></b>	0.28	<b>14.31</b>	1.45
<b>Sanliurfa-Siverek- Basdegirmen 2<sup>T</sup></b>	0.24	6.97	0.57
<b>Denizli-Dazkiri<sup>T</sup></b>	0.24	7.82	0.55
<b>Denizli-Acipayam<sup>T</sup></b>	<b>0.59</b>	11.42	1.49
<b>Sanliurfa- Bozova-Hasmersar<sup>T</sup></b>	0.33	8.03	0.65
<b>Antalya-Finike<sup>T</sup></b>	0.19	6.19	0.44



Sanliurfa-Akziyaret <sup>T</sup>	0.18	8.06	0.71
Aydın-Çine <sup>T</sup>	0.20	8.65	1.22
Adıyaman-Gölbaşı <sup>T</sup>	0.14	9.22	1.50
Antalya-Konyaaltı <sup>T</sup>	0.22	5.29	0.68
Adana-Ceyhan <sup>T</sup>	0.17	7.45	0.61
Mardin <sup>T</sup>	0.18	7.93	0.60
Diyarbakir-Bismil- Bakacak <sup>T</sup>	0.26	9.54	0.82
Diyarbakir- Bismil Yukari Harim Köy <sup>T</sup>	0.46	9.25	0.75
Adana-Yumurtalik <sup>T</sup>	0.23	9.15	0.57
Batman <sup>T</sup>	0.15	10.35	1.65
İcel-Tarsus <sup>T</sup>	0.13	10.47	0.99
Diyarbakir- Ergani- Dağarası Köyü <sup>T</sup>	0.19	6.10	0.75
Diyarbakir- Ergani Ziyaret Köy <sup>T</sup>	0.14	9.34	1.50
Greece 1	ND	9.40	0.78
Pakistan 1	ND	5.53	0.99
Pakistan 2	0.09	5.38	1.40
Iran	ND	5.77	1.59
Diyarbakir-Bismil <sup>T</sup>	0.01	3.59	0.60
Baydar-2001 <sup>C</sup>	ND	3.38	0.39
Batem-Uzun <sup>C</sup>	ND	3.71	0.41
Batem-Aksu <sup>C</sup>	ND	3.82	0.42
Boydak <sup>C</sup>	ND	6.77	0.74
Gölmarmara <sup>C</sup>	0.01	5.45	0.78
Tanas <sup>C</sup>	0.01	6.73	0.48
Hatipoğlu <sup>C</sup>	ND	5.32	0.60
Tan-99 <sup>C</sup>	0.01	4.36	0.42
Özberk <sup>C</sup>	ND	4.20	0.37
Mozambik	0.02	8.83	1.51
Israel	0.05	11.38	1.81
Pakistan 3	0.03	4.65	1.10
Greece 2	0.03	4.63	0.68
Russia 1	0.01	9.35	1.76
Russia 2	ND	8.52	1.46
Libya	ND	3.03	0.52
Egypt	0.01	7.81	0.73
Iraq	ND	5.09	1.21
S. America	0.11	2.61	1.09
USA 1	0.09	2.90	0.88
Afghanistan	0.02	9.10	1.22
Manisa-Selendi <sup>T</sup>	0.01	4.47	0.65

USA 2	ND	3.21	0.47
Kepsut-99 <sup>c</sup>	0.04	2.96	0.37
Sarisu <sup>c</sup>	ND	2.91	0.32
Osmanlı-99 <sup>c</sup>	ND	<b>2.34</b>	<b>0.23</b>
Max.	0.59	14.31	1.81
Min.	0	2.34	0.23
Ave.	0.1	6.65	0.87

<sup>c</sup>: Cultivar; <sup>T</sup>: Türkiye; ND: Not Determined.

3.3. Principal Component Analysis

Principal component analysis (PCA) was performed to determine the contribution of each trait to the overall variation observed in fatty acid, lignan, and tocol compositions. Using PCA based on the correlation matrix, we determined eigenvalues, the percentage of variability explained by a single eigenvector, and the cumulative variations explained by the first five eigenvectors (Table 4). These five PCs accounted for a total of 91.25% of the overall variations. Maximum variations were contributed by PC1 which accounted for a total of 33.344% variations and linoleic acid was the main variations contributor in this PC. PC2 accounted for a total of 18.72% variations and stearic acid was main contributor in this PC (Table 4). PC3 and PC4 accounted for a total of 14.611% and 10.508% variations, while sesamol and  $\alpha$ -tocotrienol were chief variation contributors in these PCs.

**Table 4.** Eigenvectors, eigenvalues, individual and cumulative percentages of variation explained by the first five principal components (PC) of sesame genotypes.

	<u>PC1</u>	<u>PC2</u>	<u>PC3</u>	<u>PC4</u>	<u>PC5</u>
<u>Palmitic</u>	-0.632	-0.043	-0.535	0.270	0.092
<u>Palmitoleic</u>	-0.567	0.308	-0.580	0.344	-0.046
<u>Stearic</u>	0.269	0.689	0.417	0.026	-0.040
<u>Oleic</u>	0.530	0.629	0.121	0.457	-0.022
<u>Linoleic</u>	-0.548	-0.661	-0.103	-0.442	0.025
<u>Linolenic</u>	0.808	-0.306	-0.211	0.186	0.107
<u>Arachidic</u>	-0.030	0.027	0.394	0.162	0.876
<u><math>\alpha</math>-Tocopherol</u>	0.747	-0.336	-0.025	0.271	-0.048
<u><math>\alpha</math>-Tocotrienol</u>	0.149	-0.485	0.463	0.461	-0.436
<u><math>\gamma</math>-Tocopherol</u>	-0.743	-0.346	0.309	0.434	0.039
<u>Total Tocopherol</u>	-0.716	-0.371	0.314	0.457	0.037
<u>Sesamol</u>	-0.629	0.522	-0.309	0.147	-0.018
<u>Sesamin</u>	-0.612	0.476	0.342	-0.078	-0.205
<u>Sesamolin</u>	-0.474	0.139	0.633	-0.322	-0.057
<u>Eigenvalue</u>	4.668	2.621	2.046	1.471	1.034
<u>Variability (%)</u>	<b>33.344</b>	18.720	14.611	10.508	7.387
<u>Cumulative %</u>	33.344	52.064	66.675	77.184	84.571

3.4. Correlations Among Fatty Acid, Tocol, and Lignan Composition

The results of the correlation analyses of fatty acids, lignans and tocols in the 50 sesame genotypes examined are shown in Table 5. Among the studied oil parameters, palmitic acid stands out as the parameter with the highest correlation with other oil parameters. While there was a positive and significant correlation with palmitoleic, linoleic,  $\alpha$ -tocopherol, total tocopherol and sesamol, it has been found to have a negative and significant correlation with stearic acid, oleic acid, linolenic acid and  $\alpha$ -tocopherol. A similar relationship between palmitic acid and stearic acid, palmitic acid and sesamol has been reported in sesame seeds ([37] Dar et al., 2019). While there was a positive correlation between stearic acid and oleic acid, it was determined that there was a positive correlation between  $\gamma$ -tocopherol and total tocopherol. Such a correlation between these fatty acids were also reported by Were et al., (2006) [22], Kurt (2018) [16], and Teklu et al., (2022) [20]. Morris et al., (2021) [19] also reported a similar relationship between stearic acid and  $\gamma$ -tocopherol. The strong negative correlation between oleic acid and linoleic acid has been reported by many researchers [16,22,44] and the results obtained from the present study are also consistent. Additionally, oleic acid was found to have a positive correlation with  $\alpha$ -tocopherol and a negative correlation with  $\gamma$ -tocopherol and total tocopherol. Unlike oleic acid, linoleic acid has been found to have a positive correlation with  $\gamma$ -tocopherol and total tocopherol.  $\alpha$ -tocopherol, on the other hand, stands out as an oil quality feature that has the most negative correlation with the others among the studied features. It was determined that there was a positive correlation between  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol, and a negative correlation between  $\gamma$ -tocopherol, total tocopherol, sesamol, sesamin and sesamolin. In the study, it was found that there was a positive correlation between sesamin and sesamolin, and between sesamin and sesamol. In the light of this information obtained, it can be said that increasing the amount of sesamin in breeding programs may contribute to the increase of the sesamol and sesamolin [15,45,46].

Table 5. Correlation among fatty acids, lignans and tocols in the 50 sesame genotypes.

Variables	Palmitic	Palmitoleic	Stearic	Oleic	Linoleic	Linolenic	Arachidic	$\alpha$ -Tocopherol	$\alpha$ -Tocotrienol	$\gamma$ -Tocopherol	Total Tocopherol	Sesamol	Sesamin
Palmitoleic	<b>0,706</b>												
Stearic	<b>-0,301</b>	-0,178											
Oleic	<b>-0,376</b>	-0,026	<b>0,523</b>										
Linoleic	<b>0,293</b>	0,029	<b>-0,614</b>	<b>-0,934</b>									
Linolenic	<b>-0,279</b>	<b>-0,340</b>	-0,092	0,260	<b>-0,294</b>								
Arachidic	-0,062	-0,169	0,137	0,072	-0,071	0,026							
$\alpha$ -Tocopherol	<b>-0,332</b>	<b>-0,396</b>	-0,058	<b>0,288</b>	-0,272	<b>0,795</b>	-0,016						
$\alpha$ -Tocotrienol	-0,193	<b>-0,321</b>	-0,025	-0,011	0,000	0,227	-0,078	<b>0,374</b>					
$\gamma$ -Tocopherol	<b>0,398</b>	0,270	<b>-0,297</b>	<b>-0,348</b>	<b>0,403</b>	<b>-0,500</b>	0,198	<b>-0,362</b>	<b>0,324</b>				
Total Tocopherol	<b>0,387</b>	0,253	<b>-0,306</b>	<b>-0,339</b>	<b>0,396</b>	<b>-0,465</b>	0,201	<b>-0,314</b>	<b>0,351</b>	<b>0,999</b>			
Sesamol	<b>0,534</b>	<b>0,679</b>	0,028	0,013	0,006	<b>-0,540</b>	-0,050	<b>-0,519</b>	<b>-0,371</b>	0,231	0,207		
Sesamin	0,151	<b>0,310</b>	0,226	-0,044	0,034	<b>-0,657</b>	0,039	<b>-0,526</b>	-0,079	<b>0,302</b>	<b>0,279</b>	<b>0,540</b>	
Sesamolin	-0,053	-0,116	0,159	-0,209	0,215	<b>-0,519</b>	0,152	<b>-0,380</b>	-0,005	<b>0,336</b>	<b>0,321</b>	0,139	<b>0,614</b>

Values in bold are different from 0 with a significance level alpha=0.05.

4. Conclusion

Sesame, a valuable vegetable oil, is known to have numerous beneficial properties for applications in the food industry. We identified considerable variation in studied oil parameters of 50 sesame genotypes. While the sesamin content of the sesame genotypes used in the study was higher than previously reported values, it was determined that the sesamol content was at very low levels. The sesame genotypes are rich in sesamin content and can be used as parents in breeding programs. Sesamol was not detected in almost all of the cultivars from Türkiye. In addition, it was determined that the gamma-tocopherol contents of these cultivars were below the average value. The same applies to sesamin and sesamolin. The only exception here is that the Osmanlı-99 cultivar has the highest sesamin content. The results obtained from the study showed that the lignan and tocol composition of the cultivars produced in Türkiye were not at the desired levels. Therefore, aiming to improve the lignan and tocol composition of the cultivars in the breeding programs designed to obtain new cultivars will benefit consumers in Türkiye, which consumes sesame extensively. In this

context, determining the genetic diversity of parameters in sesame oil that have positive effects on human nutrition and health is of great importance for determining the parents to be selected for these characteristics. Besides, these sesame genotypes are valuable for developing QTL mapping populations due to wide variation.

Among the oil parameters examined, palmitic acid stands out as the parameter with the highest correlation with other oil parameters. In contrast,  $\alpha$ -tocopherol stands out as the oil quality feature with the highest negative correlation among other parameters. Studying a large number of sesame genotypes and different traits such as fatty acids, tocopherols and lignans will contribute to supporting the correlations we detected between these traits.

Among the genotypes, Sanliurfa-Siverek-Basdegirmen 1 genotype has a total unsaturated fatty acids rate of 85%, sesamin 14.31 mg/g, sesamol 1.45,  $\alpha$ -tocopherol 1.52 mg/g,  $\alpha$ -tocotrienol 1.4 mg/g and  $\gamma$ -tocopherol 58.68 mg/g. was the prominent genotype.

## References

1. Kurt, C.; Demirbas, A.; Nawaz, M. A.; Chung, G.; Baloch, F. S.; Altunay, N. Determination of Se content of 78 sesame accessions with different geographical origin. *Journal of Food Composition and Analysis* **2020**, 94, Article 103621. <https://doi.org/10.1016/j.tifs.2024.104740>
2. Sukumar, D.; R. Arimboor ; A. C. Rumughan. HPTLC fingerprinting and quantification of lignans as markers in sesame oil and its polyherbal formulations. *Journal Pharmaceutical and Biomedical Analysis* **2008**, 47:795-801. <https://doi.org/10.1016/j.jpba.2008.03.018>
3. Arab, R, Casal S, Pinho T, Cruz R, Freidja ML, Lorenzo M, 2022. Effects of seed roasting temperature on sesame oil fatty acid composition Lignan sterol and tocopherol contents oxidative stability and antioxidant potential for food applications. *Molecules*. (2022) 27:4508. <https://doi:10.3390/molecules27144508>
4. Orsavova J, Misurcova L, Ambrozova JV, Vicha R, Mlcek J, 2015. Fatty acids composition of vegetable oils and its contribution to dietary energy intake and dependence of cardiovascular mortality on dietary intake of fatty acids. *Int J Mol Sci* 16:12871–12890. <https://doi.org/10.3390/ijms160612871>
5. Brigelius Flohe, R.; Kelly, F. J.; Salonen, J. T.; Neuzil, J.; Zingg, J. M.; Azzì, A. The European perspective on vitamin E: current knowledge and future research. *Am. J. Clin. Nutr.* . **2002**, 76:703 –716. <https://doi.org/10.1093/ajcn/76.4.703>
6. Yamashita, K.; Nohara, Y.; Katayama, K.; Namiki, M. Sesame Seed Lignans and  $\gamma$ -Tocopherol Act Synergistically to Produce Vitamin E Activity in Rats. *The Journal of Nutrition*, **1992**, 122(12), 2440-2446. <https://doi.org/10.1093/jn/122.12.2440>
7. Martins, S.R.; Vences, F.J.; Miera, L.E.S.; Barroso, M.R.; Carnide, V. RAPD analysis of genetic diversity among and within Portuguese landraces of common white bean (*Phaseolus vulgaris* L.). *Sci. Hortic.* **2006**, 108, 133–142. <https://doi.org/10.1016/j.scienta.2006.01.031>
8. Nadeem, M.A.; Karaköy, T.; Yeken, M.Z.; Habyarimana, E.; Hatipoğlu, R.; Çiftçi, V.; Nawaz, M.A.; Sönmez, F.; Shahid, M.Q.; Yang, S.H.; Chung, G.; Baloch, F.S. Phenotypic characterization of 183 Turkish common bean accessions for agronomic, trading, and consumer-preferred plant characteristics for breeding purposes. *Agronomy*, **2020**, 10, 272. <https://doi.org/10.3390/agronomy10020272>
9. Bhunia, R.K.; Chakraborty, A.; Kaur, R.; Gayatri, T.; Bhat, K.V.; Basu, A.; Maiti, M.K.; Sen, S.K. Analysis of Fatty Acid and Lignan Composition of Indian Germplasm of Sesame to Evaluate Their Nutritional Merits. *J. Am. Oil Chem. Soc.* **2015**, 92, 65–76. <https://doi.org/10.1007/s11746-014-2566-3>
10. Dar, A.A.; Kancharla, P.K.; Chandra, K. Assessment of variability in lignan and fatty acid content in the germplasm of *Sesamum indicum* L.. *J Food Sci Technol*, **2019**, 56, 976–986 <https://doi.org/10.1007/s13197-018-03564-x>
11. Hemalatha, S.; Ghafoorunissa. Lignans and tocopherols in Indian sesame cultivars. *Journal of American oil Chemists' Society*, **2004**, 81, 467–470.
12. Matthäus, Bertrand.; Mehmet Musa Özcan. Fatty acid composition and tocopherol contents of some sesame seed oils." *Iranian Journal of Chemistry and Chemical Engineering*, **2018**, 37.5: 151-155. <https://doi.org/10.30492/ijcce.2018.35427>

13. Rangkadilok, N.; Pholphana, N.; Mahidol, C.; Wongyai, M.; Saengsooksree, K.; Nookabkaew, S.; Satayavivad, J. Variation of sesamin, sesamol and tocopherols in Sesame (*Sesamum indicum* L) seeds and oil products in Thailand. *Food Chem.* **2010**, 122:724-730. <https://doi.org/10.1016/j.foodchem.2010.03.044>
14. Shi, L.; Liu, R.; Jin, Q.; Wang, X. The contents of lignans in sesame seeds and commercial sesame oils of China. *J Am Oil Chem Soc.* **2017**. <https://doi.org/10.1007/s11746-017-3018-7>
15. Wang, L.; Zhang, Y.; Li, P.; Wang, X.; Zhang, W.; Wei, W.; Zhang, X. HPLC analysis of seed sesamin and sesamol variation in a sesame germplasm collection in China. *J Am Oil Chem Soc.* **2012**, 89:1011–1020. <https://doi.org/10.1007/s11746-011-2005-7>
16. Kurt, C., Variation in oil content and fatty acid composition of sesame accessions from different origins. *Grasas Y Aceites* **2018** 69 (January–March (1)), 1–9 e241. <https://doi.org/10.3989/gya.0997171>
17. Shi, A.; Xian, Y.F.; Liu, R.J.; Chang, M.; Jin, Q.Z.; Wang, X.G. A Rapid Method for Simultaneous Analysis of Lignan and  $\gamma$ -Tocopherol in Sesame Oil by Using Normal- Phase Liquid Chromatography. *Oil Chem Soc.* **2018**, 95: 13–19. <https://doi.org/10.1002/aocs.12010>
18. Langyan S, Yadava P, Sharma S, Gupta NC, Bansal R, Yadav R, Kumar A. Food and nutraceutical functions of sesame oil: An underutilized crop for nutritional and health benefits. *Food Chem.* **2022** 389, 132990 <https://doi.org/10.1016/j.foodchem.2022.132990>
19. Morris JB, Wang ML, Tonniss BD. Variability for oil, protein, lignan, tocopherol, and fatty acid concentrations in eight sesame (*Sesamum indicum* L.) genotypes. *Ind. Crops Prod.* **2021**, 164, 113355 <https://doi.org/10.1016/j.indcrop.2021.113355>
20. Teklu DH, Shimelis H, Abady S. Genetic improvement in sesame (*Sesamum indicum* L.) progress and outlook: A review. *Agronomy.* **2022**, 12 (9), 2144. <https://doi.org/10.3390/agronomy12092144>
21. Kurt, C. Characterization of Some Turkish Sesame Populations and Cultivars Using Agronomical, Quality Traits and Issr Molecular Markers. Master thesis.
22. Were BA, Onkware AO, Gudu S, Welander M, Carlsson AS. Seed oil content and fatty acid composition in East African sesame (*Sesamum indicum* L.) accessions evaluated over 3 years. *Field Crops Res.* **2006** 97 (2-3), 254–260. <https://doi.org/10.1016/j.fcr.2005.10.009>
23. Shahidi, F.; Ambigaipalan, P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects - A review," *Journal of Functional Foods*, **2015**, 18 (Part B): 820-897. <https://doi.org/10.1016/j.jff.2015.06.018>
24. Delgado, A.; Al-Hamimi, S.; Ramadan, M.F.; De Wit, M.; Durazzo, A.; Nyam, K.L.; Issaoui, M. Contribution of tocopherols to food sensorial properties, stability, and overall quality, *J. Food Qual.* **2020**. <https://doi.org/10.1155/2020/8885865>
25. Shahidi, F.; Ambigaipalan, P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects - A review," *Journal of Functional Foods*, **2015**, 18 (Part B): 820-897. <https://doi.org/10.1016/j.jff.2015.06.018>
26. Fatnassi, S.; Chatti, S.; Nehdi, I. E.; Zarrouk, H. Chemical Composition, Phenolic Profile and Antioxidant Activity of *Maclura pomifera* (Rafin.) Schneider Seeds oil. In *International Symposium on Medicinal and Aromatic Plants-SIPAM* **2009**, 853, 383-390.
27. Jiang, Q.; Christen, S.; Shigenaga, M.K.; Ames, B.N.  $\gamma$ -Tocopherol the major form of vitamin E in the US diet deserves more attention. *Am. J. Clin. Nutr.* **2001**, 74 (6), 714–722. <https://doi.org/10.1093/ajcn/74.6.714>
28. Wu, D.; Wang, X.-P.; Zhang, W. Sesamol Exerts Anti-proliferative and apoptotic effect on human colorectal cancer cells via inhibition of JAK2/STAT3 signaling pathway. *Cell Mol. Biol.* **2019**, 65, 96–100. <https://doi.org/10.14715/cmb/2019.65.6.16>
29. Andargie, M.; Vinas, M.; Rathgeb, A.; Möller, E.; Karlovsky, P. Lignans of Sesame (*Sesamum indicum* L.): A Comprehensive Review. *Molecules.* **2021**, Feb 7;26(4):883. <https://doi.org/10.3390/molecules26040883>
30. Yokota, T.; Matsuzaki, Y.; Koyama, M.; Hitomi, T.; Kawanaka, M.; Enoki-Konishi, M.; Okuyama, Y.; Takayasu, J.; Nishino, H.; Nishikawa, A. Sesamin, a lignan of sesame, down-regulates cyclin D1 protein expression in human tumor cells. *Cancer Sci.* **2007**, 98, 1447–1453. <https://doi.org/10.1111/j.1349-7006.2007.00560.x>
31. Harikumar, K.B.; Sung, B.; Tharakan, S.T.; Pandey, M.K.; Joy, B.; Guha, S.; Krishnan, S.; Aggarwal, B.B. Sesamin manifests chemopreventive effects through the suppression of NF-kappa B-regulated cell



- survival, proliferation, invasion, and angiogenic gene products. *Mol. Cancer Res.* **2010**, *8*, 751–761. <https://doi.org/10.1158/1541-7786.MCR-09-0565>
32. Pecquet, C.; Leynadier, F.; Saiag, P. Immediate hypersensitivity to sesame in foods and cosmetics. *Contact Derm.* **1998**, *39*, 313. <https://doi.org/10.1111/j.1600-0536.1998.tb05948.x>
  33. Srisayam, M.; Weerapreeyakul, N.; Kanokmedhakul, K. Inhibition of two stages of melanin synthesis by sesamol, sesamin and sesamolin. *Asian Pac. J. Trop. Biomed.* **2017**, *7*, 886–895. <https://doi.org/10.1016/j.apjtb.2017.09.013>
  34. Kim, K.S.; Lee, J.R.; Lee, J.S. Determination of sesamin and sesamolin in sesame (*Sesamum indicum* L.) seeds using UV spectrophotometer and HPLC. *Korean J. Crop Sci.* **2006**, *51*, 95–100.
  35. Wu, K.; Wu, W.-X.; Yang, M.-M.; Liu, H.-Y.; Hao, G.-C.; Zhao, Y.-Z. QTL Mapping for Oil, Protein and Sesamin Contents in Seeds of White Sesame. *Acta Agron. Sin.* **2017**, *43*, 1003–1011. <https://doi.org/10.3724/SP.J.1006.2017.01003>
  36. Zhu, X.; Dai, Q.; Mu, J.; Cai, W.; Guo, Y.; Xu, Z.; Zhang, Z.; Wang, Z. Comparative assessment of lignan and total phenolic contents in sesame seeds and antioxidant capacity of sesame extracts from different producing areas in Anhui Province. *Farm Products Processing*, **2018**, *8*, 461. (in Chinese)
  37. Dar AA, Kancharla PK, Chandra K. Assessment of variability in lignan and fatty acid content in the germplasm of *Sesamum indicum* L.. *J Food Sci Technol.* **2019** *56*, 976–986. <https://doi.org/10.1007/s13197-018-03564-x>
  38. Xu, F.; Zhou, R.; Dossou, S.S.K.; Song, S.; Wang, L. Fine mapping of a major pleiotropic QTL associated with sesamin and sesamolin variation in sesame (*Sesamum indicum* L.). *Plants*, **2021**, *10*, 1–14. <https://doi.org/10.3390/plants10071343>
  39. Moazzami, A.A.; Haese, S.L.; Kamal-Eldin, A. Lignan contents in sesame seeds and products. *Eur J Lipid Sci Technol.* **2007**, *109*, 1022–1027
  40. Wang, L.; Zhang, Y.; Li, P.; Wang, X.; Zhang, W.; Wei, W.; Zhang, X. HPLC analysis of seed sesamin and sesamolin variation in a sesame germplasm collection in China. *J Am Oil Chem Soc.* **2012**, *89*, 1011–1020. <https://doi.org/10.1007/s11746-011-2005-7>
  41. Sadeghi N, Oveisi MR, Hajimahmoodi M, Jannat B, Mazaheri M, Mansouri S. The content of sesamol in Iranian sesame seeds. *Iran. J. Pharm. Res.* **2009** *8*: 101-105. <https://doi.org/10.22037/ijpr.2010.796>
  42. Kumazaki T, Yamada Y, Karaya S, Kawamura M, Hirano T, Yasumoto S, Katsuta M, Michiyama H. Effects of day length and air and soil temperatures on sesamin and sesamolin contents of sesame seed. *Plant Production Science*, **2009** *12*, 481–491. <https://doi.org/10.1626/pps.12.481>
  43. Kim JH, Seo WD, Lee SK, Lee YB, Park CH, Ryu HW, Lee JH, 2014. Comparative assessment of compositional components, antioxidant effects, and lignan extractions from Korean white and black sesame (*Sesamum indicum* L.) seeds for different crop years. *J Funct Foods* *7*:495–505. <https://doi.org/10.1016/j.jff.2014.01.006>
  44. Yol E, Toker R, Golukcu M, Uzun B, 2015. Oil Content and Fatty Acid Characteristics in Mediterranean Sesame Core Collection. *Crop Sci.* *55*, September–October. <https://doi.org/10.2135/cropsci2014.11.0771>
  45. Kancharla, P.K.; Arumugam, N. Variation of oil, sesamin and sesamolin content in the germplasm of the ancient oilseed crop *Sesamum indicum* L. *J Am Oil Chem Soc.* **2020**, *97*(5):475-83. <https://doi.org/10.1002/aocs.12346>
  46. Khuimphukhieo, I.; Khaengkhan, P. Combining ability and heterosis of sesamin and sesamolin content in sesame. *SABRAO Journal of Breeding and Genetics*, **2018** *50*:180–191.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.