

Article

Not peer-reviewed version

Identifying the Biological Potential of Western Balkan Polypore Mushroom Species to Mitigate the Negative Effects of Global Mushroom Cultivation

[Maja Kozarski](#) , [Anita Klaus](#) , [Bojana Špirović-Trifunović](#) , [Srdjan Miletić](#) , [Vesna Lazić](#) , Željko Žižak ,
[Jovana Vunduk](#) *

Posted Date: 29 November 2023

doi: 10.20944/preprints202311.1765.v1

Keywords: biological activity; bioprospecting; Corioulus; Fomitopsis; Ganoderma; mushroom cultivation; oxidative damage prevention; Polypore



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Identifying the Biological Potential of Western Balkan Polypore mushroom Species to Mitigate the Negative Effects of Global Mushroom Cultivation

Maja Kozarski ¹, Anita Klaus ¹, Bojana Špirović-Trifunović ², Srdjan Miletić ³, Vesna Lazić ¹, Željko Žižak ⁴ and Jovana Vunduk ^{5,*}

¹ Institute for Food Technology and Biochemistry, Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11080 Belgrade, Serbia; maja@agrif.bg.ac.rs (M.K.); aklaus@agrif.bg.ac.rs (A.K.); vlazic93@gmail.com (V.L.)

² Institute for Phytomedicine, Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11080 Belgrade, Serbia; spirovic@agrif.bg.ac.rs (B.Š.-T.)

³ Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia; srdjan.miletic@ihtm.bg.ac.rs (S.M.)

⁴ Institute of Oncology and Radiology of Serbia, Paterova 14, 11000 Belgrade, Serbia; zizakz@ncrc.ac.rs (Ž.Ž.)

⁵ Institute of General and Physical Chemistry, Studentski trg 12/V, 11158 Belgrade, Serbia; jvunduk@iofh.bg.ac.rs (J.V.)

* Correspondence: jvunduk@iofh.bg.ac.rs; Tel.: +381 64 020 9819

Abstract: Growing mushrooms is meeting challenges while aiming for sustainability and circularity. Wherever the producer is located, commercial strains are the same originating from several producers. Following the harvest, enormous quantities of spent mushroom substrate containing spores are disposed presenting a type of foreign material pressure on the ecosystem in the form of the loss of genetic diversity in wild mushroom populations. This challenge can be mitigated by bioprospecting local strains and using them to generate commercial inoculum. Accordingly, the mycoceutical potential of five polypore mushroom species from Serbia was evaluated: secondary metabolite composition, oxidative damage prevention, anti-tyrosinase, and anti-angiotensin converting enzyme (ACE) properties. The phenolic pattern was comparable in all samples, but the amounts of specific chemicals varied. Hydroxybenzoic acids were the primary components. All samples had varying quantities of ascorbic acid, carotene, and lycopene, and showed a pronounced inhibition of lipid peroxidation (LPx) and ability to scavenge HO[•]. Extracts were more potent tyrosinase inhibitors but unsuccessful when faced with ACE. *Fomitopsis pinicola* had the strongest anti-tumor efficacy while *Ganoderma lucidum* demonstrated strong selectivity in anti-tumor effect in comparison to healthy cells. The evaluated species provided a solid foundation for commercial development while keeping the local ecology in mind.

Keywords: biological activity; bioprospecting; *Coriolus*; *Fomitopsis*; *Ganoderma*; mushroom cultivation; oxidative damage prevention; Polypore

1. Introduction

Mushroom cultivation is speeding up in Europe driven by the widespread popularity of the vegan diet and investments in research and development of novel mushroom-based products. [1,2] Once mycophobic, the population of Western Europe is accepting mushrooms as food but also as dietary supplements as a consequence of scientific and user-based findings of fungal biological activity with few or no side effects. [3] Popular magazines, scientific journals, and specialized mushroom conferences like the International Medicinal Mushroom Conference (IMMC) are also contributing to this paradigm shift and “mushroomisation” of Europe and the Western World. However, the more they know the less satisfied with the range of mushroom products the Europeans are. [4] The biggest mushroom producers in Europe are in the Netherlands and Poland with intensive and highly industrialized facilities. They are focused on a few commercial and edible species with

the highest market share, like white button mushroom, shiitake, Oyster, and Enoki mushrooms. [5] Such a large-scale production uses standardized and certified mycelium supplied by only a few companies. This is also detrimental to the local environment. Spent substrate and spores of standardized mushroom species are discarded and dispersed in the surroundings of the production facility which decreases the diversity of the fungal genetic pool. [6,7] Moreover, small producers use the same mycelium (spawn) as the big ones, and its cost can count up to 20% of the total production expenses while not being customized for very diverse growing conditions and equipment. This can affect the final product's quality. When it comes to mushroom-based nutraceuticals the goal is to have high-quality raw materials rich in bioactive compounds, possibly standardized. [3] However, with the combination of small-scale diverse production and the lack of commercial spawn diversity, the quality is often compromised. Using wild strains can be a solution. Campi et al. reported that wild fruiting bodies showed higher concentrations of phenolic compounds than the cultivated ones when screening bioactive compounds from medicinal mushrooms from Paraguay. [8] On the other hand, mushroom producers and mushroom-based nutraceutical producers need to adhere to and follow the legislation requirements. One of them is to provide proof of the fungal strain origin (cannot be a wild strain), thus wild fruiting bodies are prohibited which severely limits the options. The situation in this sector is further complicated due to the borderline nature of edible/medicinal mushrooms (legislation bodies and peers are still unfamiliar with the material) which affects the product's legislation. [9]

Polypore mushrooms are well-known folk remedies and part of traditional medicine in Asia and Europe. Specimens of *Fomitopsis betulina* and *Fomes fomentarius* are the oldest examples of polypore mushrooms found in the bag of around 5000-years-old mummy known as the Iceman probably used as multifunctional remedies. [10] In the meantime, the biological activity (anticancer, antioxidant, anti-nematode, antidiabetes, hepatoprotective, antibiofilm, and antimicrobial activity, against hyperlipidemia, anti-inflammatory and cardioprotective activity and cytotoxic toward selected cancer cell lines, etc.) of these and some other polypore species has been reported by different authors. [11–16] The most promising ones were even further tested through clinical trials and polysaccharide K, known also as krestin or PSK, from *Coriolus versicolor*, has been officially used in Japan as an adjuvant for cancer treatment. [17,18] The cultivation of polypore species is challenging as in the case of *F. betulina* (Chaga). [19,20] If not, collecting the most promising species from the wild can be properly arranged. Only by knowing which species, when to collect, grow on which host, and how to preserve, transport, store, and process we can avoid natural environment devastation while profiting from polypore power to harness our wellbeing.

Under the frame of the aforementioned several genera of polypore mushrooms often occurring in Serbia, the Western Balkan region, were selected and tested for their biological activities. Namely, five species (*Fomitopsis pinicola*, *F. betulina*, *Ganoderma applanatum*, *Ganoderma lucidum* and *C. versicolor*) were evaluated for their secondary metabolites, including phenolic compounds (quantity and quality), ascorbic acid, β -carotene and lycopene content, and biological activities like antioxidant potential, anti-tyrosinase activity, inhibition of ACE, and cytotoxic/antitumor activity toward selected tumor cell lines. The activities of the examined species were compared with the same species from other regions of the world and evaluated as potential sources of compounds of nutraceutical value. Finally, the emphasis is given on the importance of local resources to preserve fungal diversity and fit into the circular and blue economy frame of production which goes in line with several of the United Nations Sustainable Development Goals, especially responsible consumption and production and climate action. [21]

2. Results

2.1. Extraction yields and secondary metabolite content

There were statistically significant differences in the yields of all extracts, with *F. pinicola* extract being the most abundant ().

Table 1. Extraction yields and secondary metabolite contents in methanol extracts of the five lignicolous mushroom species.

Extract	Yield (mg g ⁻¹) ^α	Total content (mg g ⁻¹) ^β			
		Phenols ^δ	Vitamin C	β-carotene	lycopene
<i>F. betulina</i>	260.5±15.3 ^{By}	32.6±2.1 ^C	1.9±0.1 ^B	51.7±6.2 ^E	23.8±2.6 ^E
<i>F. pinicola</i>	360.3±20.1 ^A	133.1±6.2 ^A	2.0±0.2 ^B	541.5±21.1 ^A	265.7±13.1 ^A
<i>G. applanatum</i>	120.2±10.3 ^C	39.4±1.8 ^B	1.2±0.1 ^C	188.6±13.5 ^C	57.5±4.7 ^D
<i>G. lucidum</i>	100.8±7.9 ^D	38.6±2.6 ^B	3.0±0.1 ^A	372.2±18.9 ^B	224.6±17.3 ^B
<i>C. versicolor</i>	60.7±5.8 ^E	10.5±0.9 ^D	3.1±0.2 ^A	158.7±10.0 ^D	109.2±8.2 ^C

^αAll values are on a dry weight (DW) basis of the fruiting body; ^βAll values of metabolites are on a DW basis of extract; data represent mean ± standard deviation (n ±3); ^γ within the same column, means followed by different letters are significantly different at p ≤ 0.05; ^δ gallic acid equivalents (GAE);.

Similarly, statistically significant differences in the content of examined secondary metabolites were identified across the tested samples. The results showed that phenol concentrations were the highest among the examined compounds. TPC levels were from 60.7 to 360.3 mg GAE g⁻¹ of extract DW, with *F. pinicola* having the highest level. In comparison to this study, high phenol content (278-313 mg GAE g⁻¹) was found in 70% and pure ethanol extracts of *F. pinicola* from Belgrad Forest, Istanbul, and 70% ethanol extract of *F. pinicola* (264 mg GAE g⁻¹) from Czech forests. [22,23] Reis et al. revealed the highest phenolics content (388 mgGAE g⁻¹) in *F. pinicola* methanol extract among ten wild mushroom species collected in Bragança (Northeast Portugal). [24] Janjusevic et al. reported 71.55 mg and 64.76 mg respectively. [25] In our study, TPC in methanol extracts of *C. versicolor* fruiting bodies obtained from Fruška Gora, a national park in Serbia, was lower (10 mg GAE g⁻¹) than reported by other authors. Macakova et al. discovered substantially higher phenol content (30.7 mg GAE g⁻¹) in a 70% ethanol extract of *C. versicolor* gathered in the Czech Republic's woodlands, while Matijášević et al. reported 25.8 mg GAE g⁻¹ in methanol extract of *C. versicolor* obtained from Serbia. [22,26]

Vitamin C was found in high concentrations, in all of the extracts examined, ranging from 1.2 mg g⁻¹ (*G. applanatum*) to 3.1 mg g⁻¹ (*C. versicolor*) (

). Besides being at the lowest concentration in *G. applanatum* extract, it was significantly higher than the vitamin C content of methanol extract (0.016 mg g⁻¹) of *G. applanatum* fruiting bodies collected from Gaurishankar Conservation Area, Central Nepal. [27] Reis et al reported 1.08 mg g⁻¹ of Vitamin C in methanol extract of *F. pinicola* collected in Bragança (Northeast Portugal. It was almost 50% lower than the vitamin C content measured in our study for the *F. pinicola* extract, Table 1. Moreover, the vitamin C content of investigated polypore extracts was higher than its content in some fruits e.g. apples, pears, and plums 3–5 mg 100g⁻¹. [28]

There were statistically significant differences in the quantity of carotenoids, lycopene, and its metabolite β-carotene, among the studied extracts (

). To our knowledge β-carotene and lycopene content was detected for the first time in *F. pinicola*. There were the highest concentrations among the tested extracts (541.5 μg g⁻¹ and 265.7 μg g⁻¹, Table 1.

2.1.1. Phenolic profile of the methanol extracts

Table 2 shows the polyphenol composition of methanol extracts of five lignicolous mushroom species. The extracts included nine of the total thirteen analyzed phenolic acids and their derivatives. The most abundant, according to the results, were hydroxybenzoic acids (HBA). Gallic acid was the primary phenolic element in *F. pinicola* extract, while *C. versicolor* extract had the least amount, nearly 99% less than *F. pinicola* extract (Table 2). Furthermore, Onar et al. discovered that gallic acid was one of the primary constituents (5.5333 ± 0.0181 ppm) of *F. pinicola* ethanol extract determined by HPLC. [23] The most abundant phenolic acid in *F. betulina* extract was p-hydroxybenzoic acid (94.27 μg g⁻¹ DW), while it was the least abundant in *C. versicolor* extract (0.24 μg g⁻¹ DW). p-Hydroxybenzoic acid and its esters, sometimes known as parabens, commonly occur in bound form and are typically a component of a complex structure in mushrooms such as lignins and hydrolyzable tannins. [25]

Protocatechuic acid, a p-hydroxybenzoic acid derivative, was found in all of the extracts tested. The minimum and maximum protocatechuic acid values in *G. lucidum* and *F. betulina* were $5.26 \mu\text{g g}^{-1}$ and $29.94 \mu\text{g g}^{-1}$, respectively. Ellagic acid was detected in methanol extracts of *F. pinicola* and *G. applanatum* ($31.23 \mu\text{g g}^{-1}$ and $22.03 \mu\text{g g}^{-1}$). Ellagic acid is derived from ellagitannins, and for comparison, it is widely found in a variety of plants and fruits (berries and walnuts). [29] Vanillin was found in extracts of *G. lucidum*, *F. pinicola*, and *G. applanatum* at concentrations ranging from 0.81 to $1.81 \mu\text{g g}^{-1}$. Vanillic acid, an oxidized form of vanillin, was found in extracts of *F. betulina* and *G. applanatum*. Vanillic acid is widely used as a flavoring agent in various food products and drinks, pharmaceutical, and cosmetics industries. [30] Syringic acid was detected in extracts of *F. pinicola*, *F. betulina*, and *G. applanatum* in the range of 3.07 – $9.34 \mu\text{g g}^{-1}$. Among the hydroxycinnamic acids (HCA), chlorogenic acid was found in low amounts in extracts of *F. betulina*, *F. pinicola*, *G. applanatum*, and *G. lucidum* (Table 2). Caffeic acid was exclusively found in *G. lucidum*. It can be found in foods like coffee, wine, and tea, as well as popular medications like propolis. [31] p-Coumaric acid was detected in *F. betulina*, *F. pinicola*, and *G. lucidum* in concentrations ranging from 0.49 to $18.24 \mu\text{g g}^{-1}$, respectively. This acid was not present in the extracts of *F. pinicola* and *C. versicolor*. On the contrary, Janjusevic et al. (2017) discovered p-coumaric acid ($1.28 \mu\text{g g}^{-1}$) in *C. versicolor* methanol extract. [25] Furthermore, Nowacka et al. reported p-coumaric acid ($0.56 \mu\text{g g}^{-1}$ DW) in an ethanol extract of *F. pinicola* gathered in the forests of Lublin Province, Poland. [32] All parallels and discrepancies in the presence and concentration of phenolic acids found in this study and the literature can be explained by the variety of growing circumstances, extraction methods, and growing environments, as well as stressors like UV radiation, virus, and parasite infection, damaging air pollution, and exposure to severe temperatures/altitude above sea level [33].

In this investigation, the flavonoid profile was compared to five standards (Table 2). Different mushroom species have been found to contain quercetin, rutin, catechin, and kaempferol. [34] In contrast to these findings, LC-MS/MS analysis of the studied methanol extracts in Table 2 did not show these flavonoids. Furthermore, the presence of stilbenes, such as resveratrol, was not detected.

Table 2. Identification and quantification of phenolic compounds in methanol extracts of the five lignicolous mushroom species by LC-MS/MS.

Compounds	<i>F. betulina</i>	<i>F. pinicola</i>	<i>G. applanatum</i>	<i>G. lucidum</i>	<i>C. versicolor</i>
Simple phenols/ derivative ($\mu\text{g g}^{-1}$)					
Vanillin	n.d.	$1.06 \pm 0.02^{\text{B}\gamma}$	$1.81 \pm 0.07^{\text{A}}$	$0.81 \pm 0.01^{\text{C}}$	$0.83 \pm 0.02^{\text{C}}$
Hydroxybenzoic acids ($\mu\text{g g}^{-1}$)					
Gallic acid	$1.48 \pm 0.05^{\text{C}}$	$951.12 \pm 6.30^{\text{A}}$	$1.93 \pm 0.06^{\text{B}}$	$0.94 \pm 0.01^{\text{D}}$	$0.77 \pm 0.01^{\text{E}}$
Ellagic acid	n.d.	$31.23 \pm 1.22^{\text{A}}$	$22.03 \pm 1.01^{\text{B}}$	n.d.	n.d.
Protocatechuic acid	$29.94 \pm 0.09^{\text{A}}$	$11.62 \pm 0.62^{\text{B}}$	$8.92 \pm 0.31^{\text{C}}$	$5.26 \pm 0.11^{\text{E}}$	$6.72 \pm 0.13^{\text{D}}$
p-Hydroxybenzoic acid	$94.27 \pm 1.11^{\text{A}}$	$7.59 \pm 0.13^{\text{D}}$	$12.76 \pm 0.08^{\text{B}}$	$11.42 \pm 0.07^{\text{C}}$	$0.24 \pm 0.01^{\text{E}}$
Vanillic acid	$11.20 \pm 0.07^{\text{B}}$	n.d.	$19.80 \pm 0.10^{\text{A}}$	n.d.	n.d.
Syringic acid	$6.37 \pm 0.22^{\text{B}}$	$3.07 \pm 0.09^{\text{C}}$	$9.34 \pm 0.12^{\text{A}}$	n.d.	n.d.
Hydroxycinnamic acids ($\mu\text{g g}^{-1}$)					
trans-cinnamic acid	n.d.	n.d.	n.d.	n.d.	n.d.
Chlorogenic acid	$1.33 \pm 0.02^{\text{A}}$	$0.76 \pm 0.01^{\text{B}}$	$0.75 \pm 0.02^{\text{B}}$	$0.36 \pm 0.01^{\text{C}}$	n.d.
Caffeic acid	n.d.	n.d.	n.d.	3.73 ± 0.10	n.d.
p-coumaric acid	$0.49 \pm 0.01^{\text{C}}$	n.d.	$0.79 \pm 0.01^{\text{B}}$	$18.24 \pm 0.92^{\text{A}}$	n.d.
Ferulic acid	LOD	LOD	LOD	LOD	LOD
Sinapic acid	LOD	LOD	LOD	LOD	LOD
Flavonoids					
Quercetin	n.d.	n.d.	LOD	n.d.	n.d.

Rutin	LOD	LOD	LOD	LOD	LOD
Catechin	n.d.	n.d.	n.d.	n.d.	n.d.
Kaempferol	n.d.	n.d.	LOD	n.d.	n.d.
Stilbenes					
($\mu\text{g g}^{-1}$)					
Resveratrol	LOD	LOD	LOD	LOD	LOD

nd: not detected; LOD: limit of detection-the smallest amount or concentration of the analyte in the test sample that can be reliably distinguished from zero; All values of metabolites are on a dry weight (DW) basis of extract; data represent mean \pm standard deviation ($n \pm 3$); γ within the same row, means followed by different letters are significantly different at $p \leq 0.05$.

2.2. Antioxidant potential

Five complementary experiments were performed in the current investigation to gain a comprehensive understanding of the antioxidant activities of the methanolic extracts of the five polypore mushroom species, and the findings are shown in Figure 1 and Figure 2, as well as in Table 3.




Figure 1. Antioxidant activity of polypore mushroom methanol extracts; inhibition of LPx evaluated in the linoleic acid model system (a), scavenging ability on DPPH radicals (SA•DPPH) (b), ferric-reducing antioxidant power (FRAP) (c), and Fe²⁺ chelating ability (d). Each value is expressed as mean \pm SEM ($n = 3$).




Figure 2. •OH scavenging activity (SA•OH) of polypore mushroom methanol extracts. The inserted figure represents the ESR spectrum of DMPO-OH spin adducts (error bars are too small and not visible).

Table 3. EC₅₀ and IC₅₀ values of polypore mushroom methanol extracts in the antioxidant potential and enzyme inhibition.

property	<i>F. betulina</i>	<i>F. pinicola</i>	<i>G. applanatum</i>	<i>G. lucidum</i>	<i>C. versicolor</i>
Antioxidant potential					
EC ₅₀ (mg mL ⁻¹)					
Inhibition of LPx	0.091±0.003 ^{C γ}	0.012±0.001 ^A	0.018±0.001 ^A	0.036±0.004 ^B	0.096±0.002 ^C
SA•DPPH	1.51±0.11 ^E	0.49±0.05 ^A	0.048±0.006 ^B	0.88±0.08 ^C	0.68±0.09 ^D
SA•OH	0.23±0.01 ^B	0.05±0.005 ^A	0.06±0.005 ^A	0.29±0.02 ^C	0.36±0.03 ^C
FRAP	3.51±0.16 ^E	2.37±0.13 ^D	0.30±0.02 ^A	1.36±0.09 ^B	1.92±0.12 ^C
Fe ²⁺ chelating ability	4.78±0.31 ^D	0.99±0.09 ^A	4.13±0.10 ^C	1.17±0.07 ^B	1.02±0.09 ^{A,B}
Enzyme inhibition					
IC ₅₀ (mg mL ⁻¹)					
Tyrosinase inhibitory activity	0.69±0.05 ^C	0.10±0.04 ^A	0.51±0.08 ^B	0.81±0.07 ^C	1.16±0.09 ^E
ACE inhibitory activity	1.01±0.12 ^B	0.65±0.04 ^A	0.89±0.09 ^B	1.28±0.11 ^C	1.96±0.18 ^D

data represent mean ± standard deviation (n ±3); γ within the same row, means followed by different letters are significantly different at p ≤ 0.05; EC₅₀ value represents the effective concentration of extract required to show 50% antioxidant activities; IC₅₀ value represents the effective concentration of extract required for 50 % *in vitro* enzyme inhibition.

2.2.1. Scavenging ability

The •DPPH and •OH tests were used to measure the scavenging capacity. The •DPPH assay can be used to determine the antioxidant capacity of complex samples in which HAT (hydrogen atom transfer), ET (electron transfer), and proton-coupled electron transfer (PCET) mechanisms may play different roles in different proportions depending on the corresponding reaction conditions (such as pH and solvent). [35] Each extract's percentage of •DPPH inhibition was found to be high (Figure 1). At 5 mg mL⁻¹, the greatest percentages of inhibition were 92.18, 92.77, 100, 89.24, and 94.44 in *F. betulina*, *F. pinicola*, *G. applanatum*, *G. lucidum*, and *C. versicolor* methanol extracts, respectively (Figure 1). Among the extracts tested, *G. applanatum* had the highest •DPPH scavenging capacity (IC₅₀) value

of $48 \mu\text{g mL}^{-1}$ (Table 1) when compared to the conventional additions of ascorbic acid (IC_{50} $56 \mu\text{g mL}^{-1}$) and alpha-tocopherol acetate ($39 \mu\text{g mL}^{-1}$). In the study of Karaman et al. eight fungal species originating from Serbia were evaluated for $\bullet\text{DPPH}$ scavenging capacity. [36] *G. applanatum* methanol extract was also the most promising with 82.80% of $\bullet\text{DPPH}$ inhibition at the $12.5 \mu\text{g mL}^{-1}$. $\bullet\text{OH}$ scavenging activity of the examined methanol extracts are presented in Figure 2 and Table 3. $\bullet\text{OH}$ may react with organic compounds in two ways: hydrogen atom (or electron) abstraction and (slightly faster) addition to double bonds. [37] This assay reveals not only a scavenger's ability to react with $\bullet\text{OH}$ but also its ability to form complexes with Fe^{2+} , influencing its redox behavior in the Fenton reaction. *F. pinicola* extract was the most effective antioxidant in scavenging $\bullet\text{OH}$, exhibiting 83% activity as early as 0.01 mg mL^{-1} . For the comparison, at 0.5 mg mL^{-1} , *F. betulina* and *G. lucidum* extracts had the same impact ($\text{SA}\bullet\text{OH}$: 85% and 82%, respectively).

2.2.2. LPx Inhibition

As a result and the fact that linoleic acid is among the most prevalent polyunsaturated fatty acids (PUFA) in the cell membranes [38], in this research, the total antioxidant activity in the protection of LPx in the linoleic acid model system was studied. The mean EC_{50} values for *F. pinicola* and *G. applanatum* suggested that they were the most powerful antioxidants in the protection of linoleic acid peroxidation, with values of 12 and $18 \mu\text{g mL}^{-1}$, respectively (Table 3). In addition, *F. betulina*, *G. lucidum*, and *C. versicolor* extracts were effective linoleic acid oxidation inhibitors. They had lower EC_{50} values (Table 3) than ascorbic acid ($1.65 \pm 0.08 \text{ mg mL}^{-1}$), a typical food ingredient utilized at mg levels. One of the most effective antioxidants extensively employed in industry is alpha-tocopherol acetate, with an EC_{50} value of $36 \mu\text{g mL}^{-1}$.

2.2.3. FRAP assay

Reductones can exert an antioxidant effect by interrupting the free radical chain reaction by donating a hydrogen atom. [39] In the FRAP experiment, newly generated electron donors provided by antioxidants can decrease Fe^{3+} to Fe^{2+} . This reducing capacity of substances could be used to predict possible antioxidant capabilities, and increased absorbance indicates a rise in reducing power. [40] Concentration-dependent, the reducing capabilities of mushroom methanol extracts increased with increasing concentration (Figure 2). *G. applanatum* extract had the highest increasing absorbance (2.48 at 2.5 mg mL^{-1}) and the lowest EC_{50} value (0.30 mg mL^{-1}) of the five extracts (Figure 2 and Table 3). In the FRAP assay, ascorbic acid had a lower EC_{50} value ($0.056 \pm 0.007 \text{ mg mL}^{-1}$) and stronger reducing power than the examined extracts.

2.2.4. Capability to chelate Fe^{2+}

Testing chelating ability on ferrous ions, methanol extracts in this study showed different patterns of chelating ability at 0.01 - 5 mg mL^{-1} (Figure 2 and Table 3). *F. pinicola* and *C. versicolor* extracts displayed the best Fe^{2+} chelating activity among the extracts at 5 mg mL^{-1} , 96.3%, and 90.7%, respectively (Figure 2). These samples also had the lowest EC_{50} values (0.99 and 1.02 mg mL^{-1}) (Table 3). Ascorbic acid had an EC_{50} value of 0.49 mg mL^{-1} . At 0.1 - 5 mg mL^{-1} , the chelating effect of the synthetic metal chelator EDTA was between 91% and 100% while citric acid was not an effective chelating agent. At 5 mg mL^{-1} , its chelating ability was 3.4%.

2.2.5. Correlation between antioxidant activity and components of methanol extracts

To better understand the mechanism of antioxidant action, the correlation coefficients (r) among results obtained with antioxidant assays, and antioxidant assays with the content of bioactive components of methanol mushroom extracts were determined (Figure 3, Table 1S).

Correlations among results obtained with antioxidant assays were positively very strong for $\text{SA}\bullet\text{DPPH}$ and FRAP ($r=0.85$); positively strong for inhibition of LPx and $\text{SA}\bullet\text{DPPH}$ ($r=0.68$), inhibition of LPx and $\text{SA}\bullet\text{OH}$ ($r=0.76$), and $\text{SA}\bullet\text{DPPH}$ and $\text{SA}\bullet\text{OH}$ ($r=0.62$); moderate, positive correlation was observed between inhibition of LPx and FRAP ($r=0.57$). There was no significant correlation between

the chelating ability of Fe²⁺ and other assays (detailed information is given in Supplementary material - Table 1S).

As a confirmation, the correlations between antioxidant assays and the content of bioactive components inferred, Figure 3: 1. TPC - strong and negative correlation with SA•OH and inhibition of LPx ($r=-0.70$ and $r=-0.68$), and weak with chelating ability of Fe²⁺ and SA•DPPH ($r=-0.30$ and $r=-0.25$); 2. Vitamin C - strong and negative correlation with chelating ability of Fe²⁺ ($r=-0.71$); 3. β -carotene - strong correlation with inhibition of LPx and chelating ability of Fe²⁺ ($r=-0.74$ and $r=-0.69$), moderate with SA•OH ($r=-0.40$), and weak with SA•DPPH ($r=-0.37$); 4. Lycopene - very strong correlation with chelating ability of Fe²⁺ ($r=-0.85$), moderate with inhibition of LPx ($r=-0.56$), and weak with SA•DPPH ($r=-0.21$).




Figure 3. Relationship between EC₅₀ values in antioxidant activities and analyzed secondary metabolite content.

2.3. Enzyme inhibition

2.3.1. Tyrosinase inhibitory potential

The inhibitory potential for tyrosinase exhibited by investigated methanol extracts is presented in Table 3; all samples had the ability to inhibit this enzyme. *F. pinicola* extract displayed the strongest inhibitory activity with the lowest IC₅₀ values (0.10 mg mL⁻¹) and exhibited a slightly weaker potential when compared with kojic acid (0.079 ± 0.009 mg mL⁻¹) commonly used as a standard inhibitor of tyrosinase. It is important to emphasize that in the literature, the IC₅₀ values for the tyrosinase inhibitors are commonly incomparable due to the variations in assay conditions, including different substrate concentrations, varied incubation times, and different batches of commercial tyrosinases. [41] Thus, most studies utilized a well-known tyrosinase inhibitor, such as kojic acid as a positive control in an effort to normalize the inhibitory activities of their inhibitors.

In addition, a significant and very strong and negative correlation ($r=-0.88$) between the TPC of methanol extracts and tyrosinase inhibition was reported (Figure 4, detailed information is given in Supplementary material - Table 2S). Furthermore, a strong correlation ($r=-0.60$) was observed between the total content of β -carotene and tyrosinase inhibition (Figure 4).




Figure 4. Relationship between IC₅₀ values in enzyme inhibition and analyzed secondary metabolite content.

2.3.2. Anti-ACE potential

As shown in Table 3 the results revealed varying levels of ACE inhibitory potential among tested samples. *F. pinicola* exhibited the highest ACE inhibitory activity with a IC₅₀ value of 0.65mg mL⁻¹. *G. lucidum*, *F. betulina*, and *G. applanatum* also demonstrated notable inhibitory activity, registering values of 1.28, 1.01, and 0.89 mg/mL respectively. On the other hand, *C. versicolor* exhibited the lowest ACE inhibition, with an IC₅₀ value of 1.96 mg/mL. Captopril the first oral ACE inhibitor found and commercially applied for the treatment of hypertension showed an IC₅₀ value of 0.006 ± 0.000 mg mL⁻¹. TPC had the most effect on ACE inhibition (Figure 3, Table 2S). Regression analysis revealed a strong correlation ($r=-0.71$) between the TPC of methanol extracts and ACE inhibition.

2.4. In vitro cytotoxic activity

The decrease in the survival of target cells induced by methanol extracts is presented in Table 4. In general, all extracts exerted statistically significant differences in the cytotoxic activity against the target malignant cell lines after 72 h of incubation. *F. pinicola* extract possessed the highest activity with IC₅₀ values in the range of 23-29 µg mL⁻¹ against the malignant cell lines. *C. versicolor* extract displayed less pronounced cytotoxicity, with IC₅₀ values from 220-251 µg mL⁻¹. Itharat et al. indicated that the criteria of cytotoxicity for crude plant extracts, as established by the American National Cancer Institute (NCI), is an IC₅₀ < 30 µg mL⁻¹ in the preliminary assay. [42] IC₅₀ value of cisplatin, which served as a positive control, in the cytotoxicity testing against K562 and HeLa cells was <15.6 µg mL⁻¹, and against MDA-MB-453 was 19.2 µg mL⁻¹, respectively.

Table 4. IC₅₀ values of lignicolous mushroom methanol extracts in the cytotoxicity against human malignant cells and normal-human cells.

Cytotoxicity IC ₅₀ (mg mL ⁻¹)	<i>F. betulina</i>	<i>F. pinicola</i>	<i>G. applanatum</i>	<i>G.lucidum</i>	<i>C. versicolor</i>
-human malignant cells-					
HeLa	0.042±0.004 ^{B γ}	0.024±0.005 ^A	0.060±0.012 ^C	0.164±0.020 ^D	0.232±0.023 ^E
K562	0.040±0.002 ^B	0.029±0.003 ^A	0.076±0.010 ^C	0.148±0.064 ^D	0.220±0.029 ^D
MDA-MB-453	0.042±0.001 ^B	0.023±0.002 ^A	0.066±0.013 ^C	0.170±0.002 ^D	0.251±0.021 ^E
-normal-human cells-					
MRC-5	0.043±0.005 ^A	0.032±0.006 ^A	0.083±0.006 ^B	0.360±0.010 ^C	0.402±0.021 ^D

BEAS-2B	0.040±0.007 ^A	0.029±0.004 ^A	0.079±0.006 ^B	0.230±0.020 ^C	0.280 ±0.013 ^D
---------	--------------------------	--------------------------	--------------------------	--------------------------	---------------------------

^{A-D}within the same row, means followed by different letters are significantly different at $p \leq 0.05$; IC₅₀ value representing the effective concentration of extract required for 50 % *in vitro* inhibition.

2.4.1. Selectivity coefficient in the cytotoxicity against tumor cells

Considering the possible effects of antitumor drugs on normal healthy cells, which are a normal part of the tumor microenvironment, the activities of the investigated extracts were evaluated against healthy MRC-5 and BEAS-2B cells (Figure 5, detailed information is given in Supplementary material - Table 3S). Extracts exhibited different cytotoxic effects against healthy cells. Generally, MRC-5, normal lung fibroblasts, were less sensible to the tested extracts than BEAS-2B, bronchial epithelial cells (Figure 5). MRC-5 cells contribute to the formation of connective tissue and secrete collagen, elastin, and other extracellular matrix proteins that help maintain the structural framework of tissues. Likewise, it was observed that *G. lucidum* extract exhibited significantly higher selectivity in antitumor action compared to other extracts (Figure 5).




Figure 5. Selectivity of methanol extracts in cytotoxicity against tumor cells.

3. Discussion

Three different types/classes of secondary metabolites of polypore mushroom methanol extracts were examined for their total content: polyphenols, vitamin C, and carotenoids lycopene and its metabolite β -carotene. These metabolites have a crucial role as antioxidants in keeping cells in a decreased environment, shielding them from oxidative damage, and ensuring correct cell function. [25,26] Moreover, as signaling pathway modulators, can influence cell regulation e.g. enzyme activity, cytokine generation, and gene expression. [34,43,44] Concerning these activities, selected polypore extracts rich in secondary metabolites were assessed for their antioxidant properties, potential for enzyme inhibition, and cytotoxicity against tumor cell lines. Selection of the most potent polypore species (often occurring in Serbia, the Western Balkan region) provides guidelines for its commercialization and potential application as a food additive and nutraceuticals formulation.

3.1. Polyphenols

In recent years polyphenol-rich mushroom extracts have been commercially available and according to their widespread structural diversity have been increasingly popular in the healthcare industry and cosmeceutical formulations. [25] Concerning the possible dietary application, it is

crucial to note that polyphenol content varies depending on the location and environmental conditions of growth as secondary metabolites with a protective effect against radiation, mechanical damage, and microbial infection. [45–47] Besides, mushrooms possess the ability to absorb polyphenols from the substrate on which they grow or from neighboring plants. [25] This could explain the differences in content and composition across investigated polypore species gathered from various locations, Table 2 and Table 3. At the same time, this emphasizes the significance of identifying the most potent species and developing appropriate growing techniques as well as spawn suitable for the local conditions.

The most abundant phenolic compounds, according to the results, were phenolic acids, Table 3. Phenolic acids might be used in direct interactions with radical species (RS) as enhancers of cell antioxidant defenses and to reestablish redox balance through primary (chain breaking) and secondary mechanism, e.g. chelating/deactivation of metals, inhibition or breakdown of lipid hydroperoxides. [37,46] The importance of these antioxidant compounds is related to both their concentration and reactivity. [26] In this regard, different RS can react via different primary and secondary mechanisms in different physiological media. Five assays based on different stages of the oxidation process and on different mechanisms were performed in the current investigation to gain a comprehensive understanding of the antioxidant activities of the methanolic extracts of the five polypore mushrooms, Table 3. FRAP measures the ability to reduce metals while $\bullet\text{OH}$ and $\bullet\text{DPPH}$ assays measure a sample's free radical scavenging capacity. [37] In FRAP, there is a SET reaction mechanism, while $\bullet\text{DPPH}$ and $\bullet\text{OH}$ combine, a HAT reaction and SET reaction mechanism, as well as protoncoupled electron transfer (PCET). Besides, $\bullet\text{OH}$ can act via radical adduct formation (RAF) to double bonds. [37] Chelating ability measures the potential of investigated metabolites to bind ferrous ions and form complex structures. Inhibition of LPx includes all chain-breaking mechanisms that can protect lipids from oxidative deterioration (HAT, SET, PCET, RAF). [37] Results of regression analysis indicated that TPC exerted the most significant impact on the ability of analyzed samples to primarily act as $\bullet\text{OH}$ scavengers and to reduce LPx in the linoleic model system (Figure 3, Table 1S) This could be explained by the high ability of the phenolic acids to donate hydrogen and electron and to react to several mechanisms of the scavenging actions of free radicals, e.g. HAT, ET, PCET as well as RAF. [37,48] In addition, most of the identified phenolic acids in the present study were known for their high antioxidant activity. [49] It is important to mention that as antioxidants phenolic acids have also a wide range of industrial applications. [47,50] Gallic acid is employed as a chelating agent and preservative in food and drinks and in the skin and leather industries. [51] p-Hydroxybenzoic acid is also utilized as a preservative in a variety of medications, cosmetics, pharmaceuticals, foods, and beverages. [51] Moreover, in *ex vivo* studies, Salau et al. confirmed that vanillin and vanillic acid protect against oxidative brain damage by increasing ATPase activity while inhibiting cholinergic enzymatic activity. [52]

Besides being well-known antioxidants, polyphenols may influence multiple immunomodulatory processes involved in the inflammatory response and carcinogenesis. [33,46,51] Each type of polyphenol e.g. phenolic acids can target and bind to one or more proteins and/or receptors on cells and thus trigger intracellular signaling pathways that ultimately regulate the host response, from central metabolism to signaling events. [46,47,53] This binding, mainly non-covalent, includes hydrophobic and van der Waals interactions, ionic interactions, and H bridge/bonding, and it is always reversible. [53] In this manner, cytotoxicity against tumor cells was observed in this study, Table 4. For example, the presence of methoxy groups in the aromatic ring at positions 3 and 5 is ascribed to syringic acid's medicinal action. It has the ability to influence enzyme activity, protein dynamics, and the activity of many transcription factors implicated in inflammation, cancer, and angiogenesis. [29] Furthermore, it was reported that kinases are among the major targets of polyphenols. [47] Protein kinases are important therapeutic targets in cancer because of their critical role in signaling mechanisms that drive malignant cell characteristics. [54] They have the potential to bind to ATP-binding sites thereby modulating the action of mitogen-activated protein kinase (MAPK), phosphoinositide 3-kinase (PI3K), Akt/protein kinase B (Akt/PKB), tyrosine kinases, and protein kinase C (PKC) pathways. Inhibiting or stimulating these pathways influences

phosphorylation events and modulation of gene expression. [47,54] The fact that various polyphenols interact with cyclin-dependent kinases (CDKs) highlights their potential control of cell cycle events, including cell proliferation and cancer cells development. [47] Moreover, it has been observed that compounds with gallate groups present in mushrooms can target/upregulate different cellular receptors e.g. 67 kDa laminin receptor (67LR) and zeta chain-associated 70 kDa protein (ZAP-70) which are abnormally expressed in multiple cancers. [46,55] Additionally, it has been shown to regulate multiple crucial cellular signaling pathways, including those mediated by epidermal growth factor receptor (EGFR), janus kinase/signal transducers and activators of transcription (JAK/STAT), MAPKs, nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B), phosphoinositide 3 kinase/Akt/mammalian target of rapamycin pathway (PI3K-AKT-mTOR), and others. [56] Deregulation of the abovementioned pathways is involved in the pathophysiology of cancer. In our study, *F. pinicola* extract with the highest TPC and gallic acid content (Table 1, Table 2 and Table 4) possessed the best anti-tumor activity against the malignant cell lines (K562, HeLa, and MDA-MB-453).

Due to the ability of phenols to react with proteins, in this study potential of inhibition of polypore methanol extracts against tyrosinase and ACE was investigated. Tyrosinase is widely present in fruits, mushrooms, and vegetables [50], and is responsible for enzymatic browning reactions in damaged foodstuff during post-harvest handling and processing. Accordingly, its inhibitors act as anti-browning compounds and have an important role in the control of food quality. Also in the cosmetic industry inhibitors of tyrosinase have important applications as skin-lightening agents. [57] ACE inhibition is a big challenge today. ACE inhibitors are used for the prevention of hypertension, cardiovascular diseases, and congestive heart failure. Hypertension stands as the primary cause of premature death worldwide. [57] *F. pinicola* extract displayed the strongest inhibitory activity almost comparable with kojic acid commonly used as a standard inhibitor of tyrosinase, Table 3. A very strong and significant correlation between TPC and tyrosinase inhibition was observed (Figure 4, detailed information is given in Supplementary material - Table 2S). Gallic acid was found to be the main polyphenol ingredient of *F. pinicola* extract. It has already been confirmed that this acid displays tyrosinase inhibitory activity. [58] Gallic acid can act as a substrate of tyrosinase and also can reduce dopaquinone back to levodopa (L-DOPA) through redox cycling, similar to ascorbic acid. [43] In addition, the antityrosinase activity exhibited by investigated methanol extracts could be also attributed to the presence of other phenolic acids like protocatechuic, p-hydroxybenzoic, chlorogenic, vanillic, p-coumaric, and caffeic acid. [59] An example is a confirmation that protocatechuic acid may significantly suppress the melanin content and cellular tyrosinase activity in α -melanocyte stimulating hormone-stimulated mouse melanoma cells. [59] Furthermore, TPC had the highest influence on ACE inhibition (Figure 4, Table 2S). However, *F. pinicola* extract which was the best in ACE inhibition expressed almost 100 times weaker inhibition potential compared to the captopril, often applied for the treatment of hypertension.

The presence of flavonoids was not observed among methanol extracts, Table 3. Various findings and conclusions have been published by scientists on the metabolic entry site for flavonoid biosynthesis in mushrooms and the presence or absence of the enzymes chalcone synthase (CHS) and chalcone isomerase (CHI), as well as other enzymes involved in flavonoid biosynthesis. [34] A couple of studies suggest that the abundance of flavonoids in mushrooms is related to these organisms' ability to absorb various nutrients and chemicals from the substrate on which they grow or from surrounding plants by spreading their hyphae or creating mycorrhizae. [60] In contrast to this result, Mohanta. [61] reports, for example, the presence of chalcone-flavanone isomerase (CHF1), which is related to flavonoid production in the lignicolous mushroom *Lentinula edodes* in his newest report on genome-wide research throughout the genomic sequences of the fungal kingdom. [61] The author did note, however, that the existence of genes involved in the flavonoid production pathway is not uniform in fungi. Shao et al. investigated secondary metabolites in six *Sanghuangporus sanghuang* strains which are employed for large-scale commercial manufacturing, and indicated the presence of seven flavonoid biosynthesis-related genes. [60] Furthermore, it was discovered that *S. sanghuang* produces more flavonoids in mycelial stages than in fruiting body stages. [62]

3.2. Lycopene and β -carotene

Lycopene and β -carotene contents in the investigated polypore extracts were in a wide range of concentrations, from several tens to several hundred $\mu\text{g g}^{-1}$, Table 1. Carotenoid levels and profiles in non-green tissues e.g. mushroom tissues can vary greatly and are regulated by a variety of factors, including developmental stage, environment, stress, or a combination of these. [45] As metabolites with a protective effect, they are among the most efficient physical quenchers of singlet oxygen ($^1\text{O}_2$), a highly reactive species, due to their energy levels being near to that of $^1\text{O}_2$. [37,63] Furthermore, as a potential $^1\text{O}_2$ quencher, lycopene was demonstrated to pass the blood-brain barrier and be present in the central nervous system in low amounts. [64] As food additives, lycopene and β -carotene may also be used as a chain-breaking antioxidant in a lipid environment, especially under low oxygen partial pressure. The extensive systems of double bonds make them susceptible to attack $\bullet\text{LOO}$, resulting in the formation of inactive products. [37] Among investigated polypore extracts, *F. pinicola* and *G. lucidum* can be considered excellent sources for β -carotene supplementation, i.e. vitamin A. According to data from the Mayo Clinic if β -carotene is applied as a supplement daily dose for teenagers and adults is between 6 to 15 mg (the equivalent of 10,000 to 25,000 Units of vitamin A activity) per day. [65,66] Moreover, a strong correlation between β -carotene content and inhibition of LPx of investigated extracts confirmed their potential as a chain-breaking antioxidant in a lipid environment. Additionally very strong correlation was observed among lycopene content and the chelating ability of Fe^{2+} as well as a strong correlation between β -carotene content and the chelating ability of Fe^{2+} (Figure 3 and Table 1S). Because Fe^{2+} are the most powerful pro-oxidants in the food system [67], the investigated polypore mushroom extracts might be applied to improve food quality.

Furthermore, a strong correlation was observed between the total content of β -carotene and tyrosinase inhibition (Figure 4, Table 2S). Anantharaman et al. indicated that apocarotenoids may inhibit tyrosinase activity in a dose-dependent manner. [68] Moreover, their molecular docking results implied that apocarotenoids were allosterically bound to tyrosinase through hydrophobic interactions [68].

There is not much research related to the anti-tumor activities of carotenoids. Preclinical data suggest that carotenoids might reduce growth or induce apoptosis in cancer cells. [69,70] An example is the research of Gloria et al. on the effect of carotenoids on the cell cycle and cell viability in human breast adenocarcinoma cancer cell lines (MCF-7, MDA-MB-231 and MDA-MB-235). [69] This research indicated that the effect of lycopene and β -carotene on cancer cells was cell type-dependent. In opposite to these findings, our study confirmed that β -carotene and lycopene were in the highest concentrations in *F. pinicola* extract which possessed the highest cytotoxicity against breast carcinoma MDA-MB-453 cells, as well as against K562 and HeLa cells, Table 4. It should also be taken into account that these molecules were in a significantly lower concentration compared to the TPC in *F. pinicola* extract (Table 1). Accordingly, potential synergistic and/or agonistic effect among these molecules should be investigated in expression of the anti-tumor activities. [71]

3.3. Vitamin C

Vitamin C was found in all of the extracts examined (

). The influence of location and environmental conditions of growth on the content of vitamin C in mushrooms has been insufficiently investigated so far. It can be speculated that the factors affecting the content of vitamin C in mushrooms may be the same as for plants such as genotypic differences, and climatic conditions e.g. the higher the intensity of light during the growing season- the greater vitamin C content in tissues, maturity and harvesting methods, and postharvest handling procedures. [72] It is an essential nutrient for humans. [73] Likewise, vitamin C is a normal skin constituent found at high levels in both the dermis and epidermis. [73] Vitamin C has been shown as an effective RS scavenger. It may also act as secondary antioxidant, e.g. recycle vitamin E, the main lipid-soluble antioxidant. [26,37] In opposite to these findings regression analysis in our study did not reveal a significant correlation between vitamin C content and $\text{SA}_{\bullet\text{DPPH}}$, $\text{SA}_{\bullet\text{OH}}$, FRAP, and LPx assays (Figure 3, Table 1S). A strong and negative correlation was observed with EC_{50} values in the chelating ability of Fe^{2+} (Figure 3). It was in opposition to the other constituents of the extracts,

polyphenols, and carotenoids, which primarily react as radical scavengers and as effective inhibitors of LPx (Figure 3, Table 1S). These results were consistent with the results obtained from correlation analysis among the antioxidant assays (Table 1S). Accordingly, it could be concluded that polypore methanol extracts may act as antioxidants in mixed mode. As primary antioxidants (chain breaking, RS scavengers) they can directly scavenge RS by direct reduction via electron transfers or by radical quenching via H atom transfer or by acting in the mixed mechanism. As secondary antioxidants, they may be involved in the deactivation of metals (Fe^{2+} chelating ability), and the inhibition or breakdown of lipid hydroperoxides. The significance of vitamin C in the expression of the chelating ability of Fe^{2+} may be important in the explanation of some cytotoxicity mechanisms against malignant cell lines in this study. Tumor cells contain more iron than other normal tissues. [74,75] Thus, it is reasonable to expect that vitamin C may influence on apoptosis *in vitro* in this study. [74–76]

From regression analysis, it can also be concluded that vitamin C content in this study has no significant influence on the inhibition of tyrosinase and ACE, Figure 4, Table 2S.

4. Materials and Methods

4.1. Collection, identification, and drying of mushroom fruiting bodies

Fresh wild-growing fruiting bodies of the five Polypore mushroom species, namely *F. betulina*, *F. pinicola*, *G. applanatum*, *G. lucidum*, and *C. versicolor* were collected from several sites/locations in the different regions of the Republic of Serbia (Figure 6, detailed data in Supplementary material - Table 4S). Carpophores of each mushroom species were identified according to the methods of classical herbarium taxonomy to confirm the correct species. [77,78] Representative voucher specimens were deposited in the herbarium of the Department for Industrial Microbiology at the Faculty of Agriculture, University of Belgrade, together with their mycelia cultures. For further analysis, collected mushrooms were cut into thin slices and lyophilized (Christ BETA 2–8 LD plus freeze dryer, Osterode, Germany), powdered, and subjected to methanol extraction. The solvent of choice (methanol) favors the extraction of compounds analyzed in this study [79].



Figure 6. Mushrooms sampling sites; (1) Divčibare, (2) Avala, (3) Babe, (4) Kopaonik, (5) Košutnjak.

4.2. Preparation of extracts

The dry sample (5 g) was extracted by stirring with 100 mL of methanol at 120 rpm for 24 h and filtered through the Whatman No. 4 paper. The residue was then extracted with two additional 100

mL portions of methanol, as described earlier. [80] the combined methanol extracts were evaporated at 40 °C to dryness (Buchi Rotavapor®R II) and stored at 4 °C for further use.

4.3. Secondary metabolites analysis

4.3.1. Total polyphenol content (TPC)

The Folin-Ciocalteu reaction method adapted for a 96-well microplate reader (Hanna Instruments EC 215 Conductivity Meter, Chelmsford, UK) was used to determine the TPC in methanol extracts of mushrooms [81]. Gallic acid was used as standard (0.015–0.25 mg mL⁻¹) and results were expressed as mg of gallic acid equivalent (GAE) per g of extract dry weight (DW).

4.3.2. Vitamin C content

The method adjusted for a 96-well microplate assay was previously described [79]. All samples were analyzed in triplicates and calculated based on the standard curve of vitamin C (concentration range: 0.03125–2 mg mL⁻¹). The results are expressed as mg g⁻¹ of ascorbic acid (based on extract DW).

4.3.3. Determination of β-carotene and lycopene content

β-Carotene and lycopene were determined according to the method of Nagata & Yamashita [82]. The dried methanol extract (100 mg) was vigorously shaken with 10 mL of an acetone–hexane mixture (4:6) for 1 min and filtered through Whatman No. 4 filter paper. The absorbance of the filtrate was measured at 453, 505, and 663 nm. Contents of β-carotene and lycopene were calculated according to the following equations: lycopene (mg/100 mL) = $-0.0458A_{663} + 0.372A_{505} - 0.0806A_{453}$; β-carotene (mg/100 mL) = $0.216A_{663} - 0.304A_{505} + 0.452A_{453}$. The results are expressed as μg of carotenoid per g of the extract DW.

4.3.4. LC-MS/MS of phenolics

Samples and standards were analyzed using a 1260 Series liquid chromatography system (Agilent Technologies, Santa Clara, USA) coupled with a 6460A Triple Quad tandem mass spectrometer with an electrospray ion source (Agilent Technologies) by the analytical conditions reported in an earlier study. [83] This coupled system was controlled using MassHunter software (version B.06.00, Agilent Technologies). Authentic polyphenol standards were: gallic acid, protocatechuic acid, p-hydroxybenzoic acid, chlorogenic acid, syringic acid, vanillic acid, caffeic acid, p-coumaric acid, ellagic acid, trans-cinnamic acid, ferulic acid, vanillin, quercetin, rutin, catechin, kaempferol, and resveratrol.

4.4. Determination of antioxidant potential

Five methods were used to evaluate radical-blocking capacities: 1,1-diphenyl-2-picrylhydrazyl free radical (•DPPH) scavenging ability (SA_{•DPPH}) [40,80], inhibition of lipid peroxidation (LPx) in a linoleic acid model system [40,80], ferric-reducing antioxidant power (FRAP) assay [40,80], ferrous-ion chelating ability [40,80], and hydroxyl radical (•OH) scavenging ability (SA_{•OH}) [37]. •OH were generated in the Fenton reaction system by mixing 200 μL of 112 mM DMPO (5,5-dimethyl-1-pyrroline-N-oxide), 200 μL of DMF (N, N- dimethylformamide), 200 μL of 2 mM H₂O₂, and 2 μL of 30 mM Fe²⁺ (control). The stabilization of •OH was investigated by the ESR spin trapping method [37,84]. The ESR spectra were recorded after 2.5 min, with the following spectrometer settings: field modulation 100 kHz, modulation amplitude 0.226 G, receiver gain 5 × 10⁵, time constant 80.72 ms, conversion time 327.68 ms, center field 3440.00 G, sweep width 100.00 G, x-band frequency 9.64 GHz, power 20 mW and temperature 23°C. SA_{•OH} of mushroom extracts dissolved in DMF was defined as SA_{•OH} (%) = $100 \times (h_0 - h_x)/h_0$, where h₀ and h_x are the heights of the second peak in the ESR spectrum of 5,5-dimethyl-1-pyrroline-N-oxide (DMPO)-OH spin adduct of the control and the sample, respectively.

In all assays, results were expressed as EC_{50} ($mg\ mL^{-1}$) values, representing the effective concentrations of each extract required to show 50% antioxidant activities. Extracts were analyzed at the concentration range of 0.01-0.5 $mg\ mL^{-1}$ for $\bullet OH$ scavenging ability, 0.001-0.1 $mg\ mL^{-1}$ for inhibition of LPx in the linoleic acid model system, and 0.01-5 $mg\ mL^{-1}$ for all other antioxidant assays. Ascorbic acid, α -tocopherol acetate, citric acid, and ethylenediaminetetraacetic acid (EDTA) were used as positive controls. They are common additives in food formulations and are typically used at mg levels.

4.5. Enzyme inhibition activities

Tyrosinase inhibitory potential of the tested extracts was determined in the reaction solution of 46 units/mL tyrosinase (EC 1.14.18.1, Merck) and 2.5 mM of L-DOPA in a 96-well plate using an absorbance microplate reader (ELx808, BioTek Instruments, Inc., USA) controlled by Gen5™ Software. [57] Extracts were analyzed at the concentration range of 0.031-4.0 $mg\ mL^{-1}$, using kojic acid as a reference. The inhibition rate of tyrosinase was calculated according to the following formula:

$$Inhibition (\%) = [1 - (C - D) / (A - B)] \times 100 \quad (1)$$

where A is the absorbance of each well without testing extracts, B is blank (the absorbance without testing extract and L-DOPA), C is the absorbance of wells with testing extracts, and D is the absorbance with testing extracts but without tyrosinase.

The Angiotensin-converting enzyme (ACE) activity was analyzed using the method described in a previous investigation. [10] Briefly, the inhibition percentage of extracts was determined by replacing 30 μL of water with the same volume of the sample to be studied in the concentration range 0.005–5.0 $mg\ mL^{-1}$ in the reaction solution of 26 $u\ mL^{-1}$ ACE (EC 3.4.15.1, Merck). After the addition of 10 mM hippuryl-His-Leu (HHL) (200 μL) dissolved in 0.2 M phosphate buffer (pH 8.3), the reaction solution was incubated at 37 °C for 80 min. ACE activity was stopped by a decrease in pH by the addition of 1 M HCl (150 μL). The hippuric acid formed in the enzymatic process was extracted with 1 ml of ethyl acetate, shaken, and then centrifuged at 3000 g for 10 min. The aliquot of the organic layer was taken and evaporated under airflow at 37 °C. The residue, hippuric acid, was redissolved in 1 ml of MQ water, and the absorbance was measured at 228 nm. A solution of captopril (1 $mg\ mL^{-1}$) was used as the positive control. One tablet containing 25 mg of captopril was pulverized in a mortar and extracted with 25 mL of distilled water in an ultrasonic bath. Water extract was filtered through a filter with a pore size of 0.45 μm . Extracts contained substances that could interfere, and it was necessary to determine their absorbance and apply the formula provided in this study. [10] The sample blank was prepared in the same way as the reaction blank, replacing the volume of water with the sample. The inhibition rate of ACE was calculated using equation 1, as for the calculation of anti-tyrosinase activity, where A represents absorbance in the presence of ACE, B is the absorbance of the reaction blank, C is the absorbance in the presence of ACE and inhibitor (captopril or extract), and D is the absorbance of the sample blank.

The enzyme inhibitory activities in both assays were expressed as inhibition percentage and IC_{50} values, which were calculated using linear regression analyses, as the concentration of extract required for 50 % *in vitro* inhibition.

4.6. *In vitro* cytotoxic activity

Stock solutions of methanol mushroom extracts were prepared in dimethyl sulfoxide (DMSO) and afterward diluted with a complete nutrient medium (RPMI-1640) as described in our previous report. [80] Human cervix adenocarcinoma (HeLa), breast carcinoma MDA-MB-453, normal lung fibroblast MRC-5, and normal human bronchial epithelial (BEAS-2B) cells were cultured as monolayers in the nutrient medium, while myelogenous leukemia K562 cells were maintained as a suspension culture. The cells were grown at 37 °C in a 5% CO₂ and humidified air atmosphere. Cell lines were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA). For the cell sensitivity analysis HeLa (2.000 cells per well), MDA-MB-453 (3.000 c/w), MRC-5, and BEAS-2B (5.000 c/w) cells were seeded into 96-well microtiter plates and 20 h later different concentrations

(0.0156–2 mg mL⁻¹) of investigated extracts were appended to the wells. Investigated compounds were added to a suspension of K562 cells (5.000 c/w) 2 h after cell seeding, in the same final concentrations applied to other cells. Cell survival was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) test according to the method of Ohno and Abe (1991), 72 h after the investigated extracts were added. Concentrations of the extracts that induced a 50% decrease in malignant and normal cell survival (IC₅₀ values) were calculated from a dose-response growth curve. Doxorubicin was used as a control.

4.6.1. Selectivity coefficient in the cytotoxic/antitumor action

To evaluate further the anticancer potential of the extracts, selectivity in cytotoxic/antitumor action against specific malignant cell lines, in comparison to healthy MRC-5 and BEAS-2B cells, was also determined. The selectivity coefficient (SC) was calculated by the following equation: SC = IC₅₀ healthy cells/IC₅₀ cancer cells.

4.7. Statistical analysis

All experiments were carried out in triplicate and expressed as the mean ± standard deviation (SD). Statistical analyses were performed with the Statistica 12.0 software package (StatSoft Inc., Tulsa, OK, USA), using a one-way analysis of variance (ANOVA) for all collected data. Differences between the means for each treatment were determined using Duncan's multiple range tests ($P < 0.05$). The correlation coefficients (r) between antioxidant activities and methanol extracts as well as enzyme inhibition activities and methanol extracts were determined.

5. Conclusion

Due to the high levels of phenolic compounds, vitamin C, and carotenoids, polypore mushroom methanol extracts of the five lignicolous species *F. betulina*, *F. pinicola*, *G. applanatum*, *G. lucidum*, and *C. versicolor* from various regions of Serbia, the Western Balkan region, have a high biological/industrial potential and should be considered for spawn development. Furthermore, in the functional food market, the commercialization of wild mushroom species with potential biological activity is a growing trend. The extracts tested positive for suppression of LPx in the linoleic acid model system as well as inhibition of the detrimental propagation of radical chain reactions. All extracts had strong anti-tyrosinase activity and modest ACE inhibition. The most promising actions were identified for *F. pinicola* extract, which corresponded to the highest values of the total amount of secondary metabolites. Furthermore, *F. pinicola* had the best activity against the selected malignant cell lines that met the NCI cytotoxicity criteria for crude extracts. Otherwise, *G. lucidum* extract displayed considerably stronger selectivity in antitumor action against malignant cell lines, mainly MRC-5, which are involved in connective tissue production, when compared to healthy cells. The content of bioactive compounds like phenolics is frequently determined by the location and environmental conditions under which they grow. Thus, screening and adopting local strains can offer higher quality material to use in the food, pharmaceutical, and agricultural sectors. At the same time, this practice supports the preservation of the local pool of fungi species.

As observed and emphasized in this study, the environmental conditions, substrate, climate, stage of development, and species affect bioactive molecules content, diversity, and biological activity directing future studies to consider these factors and determine the exact species-selected factors relations.

One should keep in mind that there are considerable changes in the activities of the metabolic form of bioactive molecules and their form in the mushroom. Thus, potential candidates should be isolated and tested in real systems like food or dietary supplements and further strengthened by clinical studies. In this way, promising activities like the inhibition of tyrosinase, antihypertensive, or antioxidant activity may or may not be confirmed which will strengthen the market position of this material and build trust among consumers.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Table 1S: Relationship between EC₅₀ values in antioxidant activities and analyzed secondary metabolite content; Table 2S: Relationship between IC₅₀ values in enzyme inhibition and analyzed secondary metabolite content; Table 3S: Selectivity of methanol extracts in antitumor action; Table 4S: Mushroom species collected with corresponding family, habitat, sampling locations and usability. According to the map of Figure 1.

Author Contributions: Conceptualization, J.V., A.K. and M.K.; Methodology, J.V., A.K., M.K., B.Š-T. and Ž.Ž.; Software and Validation, S.M. and M.K.; Formal analysis, J.V., M.K., A.K., B.Š-T., Ž.Ž. and V.L.; Investigation, M.K., J.V. and A.K.; Resources, A.K., M.K. and J.V.; Data curation, M.K. and S.M.; writing – original draft preparation, M.K., J.V., B.Š-T. and S.M.; writing – review and editing, M.K., J.V. and S.M.; Visualization, M.K. and S.M.; Supervision, J.V. and A.K.; Project administration, A.K.; Funding acquisition, A.K., M.K. and J.V. All authors have read and agreed to the published version of the manuscript.

Funding: This research has been financially supported by the Ministry of Science, Technological Development and Innovation of Republic of Serbia (Contracts No: 451-03-47/2023-01/200051 and 451-03-47/2023-01/200026).

Acknowledgments: The authors are grateful to Ph.D. Snežana Spasić (Principal Research Fellow at the Institute of Chemistry, Technology and Metallurgy, University of Belgrade,) for the valuable comments and help during the preparation of original draft.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Giusti, A.; Tinacci, L.; Verdigi, F.; Narducci, R.; Gasperetti, L.; Armani, A. Safety and Commercial Issues in Fresh Mushrooms and Mushroom-Based Products Sold at Retail in Tuscany Region. *Ital. J. Food Saf.* **2022**, *11*, doi:10.4081/ijfs.2022.10044.
- Regulation (EU) 2015/2283 of the European Parliament and of the Council of 25 November 2015 on Novel Foods, Amending Regulation (EU) No 1169/2011 of the European Parliament and of the Council and Repealing Regulation (EC) No 258/97 of the European Parliament. *Off. J. Eur. Union* **2015**, *L 327*, 1–22.
- Vunduk, J.; Tura, D.; Biketova, A.Y. Medicinal Mushroom Nutraceutical Commercialization: Two Sides of a Coin. In *Wild Mushrooms Characteristics, Nutrition, and Processing*; Dhull, S.B., Bains, A., Chawla, P., Sadh, P.K., Eds.; CRC Press Taylor and Francis Group, Boca Raton, 2022; p. 43 ISBN 9781003152583.
- Almádi, B. The Producer and Consumer Aspects of the Mushroom Sector in Hungary, Szent István University, Gödöllő, Hungary, 2021.
- Bringye, B.; Fekete-Farkas, M.; Vinogradov, S. An Analysis of Mushroom Consumption in Hungary in the International Context. *Agriculture* **2021**, *11*, 677, doi:10.3390/agriculture11070677.
- Obatake, Y.; Murakami, S.; Matsumoto, T.; Fukumasa-Nakai, Y. Isolation and Characterization of a Sporeless Mutant in *Pleurotus Eryngii*. *Mycoscience* **2003**, *44*, 33–40, doi:10.1007/S10267-002-0074-Z.
- Okuda, Y. Sustainability Perspectives for Future Continuity of Mushroom Production: The Bright and Dark Sides. *Front. Sustain. Food Syst.* **2022**, *6*, doi:10.3389/fsufs.2022.1026508.
- Campi, M.; Mancuella, C.; Ferreira, F.; Maubet, Y.; Cristaldo, E.; Gayoso, E.; Robledo, G. Does the Source Matter? Phenolic Compounds and Antioxidant Activity from Mycelium in Liquid Medium, Wild and Cultivated Fruiting Bodies of the Neotropical Species *Ganoderma Tuberculosis*. *J. Microbiol. Biotechnol. food Sci.* **2023**, e6148, doi:10.55251/jmbfs.6148.
- Molitorisová, A.; Monaco, A.; Purnhagen, K.P. An Analysis of the Regulatory Framework Applicable to Products Obtained from Mushroom and Mycelium. *SSRN Electron. J.* **2021**, doi:10.2139/ssrn.3955899.
- Vunduk, J.; Klaus, A.; Kozarski, M.; Petrovic, P.; Zizak, Z.; Niksic, M.; Van Griensven, L.J.L.D. Did the Iceman Know Better? Screening of the Medicinal Properties of the Birch Polypore Medicinal Mushroom, *Piptoporus Betulinus* (Higher Basidiomycetes). *Int. J. Med. Mushrooms* **2015**, *17*, 1113–1125, doi:10.1615/intjmedmushrooms.v17.i12.10.
- Sułkowska-Ziaja, K.; Szewczyk, A.; Galanty, A.; Gdula-Argasińska, J.; Muszyńska, B. Chemical Composition and Biological Activity of Extracts from Fruiting Bodies and Mycelial Cultures of *Fomitopsis Betulina*. *Mol. Biol. Rep.* **2018**, *45*, 2535–2544, doi:10.1007/s11033-018-4420-4.

12. Bishop, K.S. Characterisation of Extracts and Anti-Cancer Activities of *Fomitopsis Pinicola*. *Nutrients* **2020**, *12*, 609, doi:10.3390/nu12030609.
13. Darkal, A.K.; Zuraik, M.M.; Ney, Y.; Nasim, M.J.; Jacob, C. Unleashing the Biological Potential of *Fomes Fomentarius* via Dry and Wet Milling. *Antioxidants* **2021**, *10*, 303, doi:10.3390/antiox10020303.
14. Hossain, M.S.; Barua, A.; Tanim, M.A.H.; Hasan, M.S.; Islam, M.J.; Hossain, M.R.; Emon, N.U.; Hossen, S.M.M. *Ganoderma Applanatum* Mushroom Provides New Insights into the Management of Diabetes Mellitus, Hyperlipidemia, and Hepatic Degeneration: A Comprehensive Analysis. *Food Sci. Nutr.* **2021**, *9*, 4364–4374, doi:10.1002/fsn3.2407.
15. Nikolic, M.; Lazarevic, N.; Novakovic, J.; Jeremic, N.; Jakovljevic, V.; Zivkovic, V.; Bradic, J.; Pecarski, D.; Tel-Çayan, G.; Glamocija, J.; et al. Characterization, In Vitro Biological Activity and In Vivo Cardioprotective Properties of *Trametes Versicolor* (L.:Fr.) Quél. Heteropolysaccharides in a Rat Model of Metabolic Syndrome. *Pharmaceutics* **2023**, *16*, 787, doi:10.3390/ph16060787.
16. Vunduk, J.; Klaus, A.; Lazić, V.; Kozarski, M.; Radić, D.; Šovljanski, O.; Pezo, L. Artificial Neural Network Prediction of Antiadhesion and Antibiofilm-Forming Effects of Antimicrobial Active Mushroom Extracts on Food-Borne Pathogens. *Antibiotics* **2023**, *12*, 627, doi:10.3390/antibiotics12030627.
17. Panda, S.K.; Luyten, W. Medicinal Mushrooms: Clinical Perspective and Challenges. *Drug Discov. Today* **2022**, *27*, 636–651, doi:10.1016/j.drudis.2021.11.017.
18. PDQ Integrative - Alternative and Complementary Therapies Editorial Board *Medicinal Mushrooms (PDQ®): Health Professional Version*; 2002;
19. Pleszczyńska, M.; Wiater, A.; Siwulski, M.; Lemieszek, M.K.; Kunaszewska, J.; Kaczor, J.; Rzeski, W.; Janusz, G.; Szczodrak, J. Cultivation and Utility of *Piptoporus Betulinus* Fruiting Bodies as a Source of Anticancer Agents. *World J. Microbiol. Biotechnol.* **2016**, *32*, 151, doi:10.1007/s11274-016-2114-4.
20. Den Herder, M. Chaga Cultivation Available online: https://efi.int/sites/default/files/files/knowledge/projects/EFI_Chaga_Cultivation_EN.pdf (accessed on 25 June 2023).
21. Do You Know All 17 SDGs? Available online: <https://sdgs.un.org/goals> (accessed on 25 June 2023).
22. Macáková, K.; Opletal, L.; Polásek, M.; Samková, V. Free-Radical Scavenging Activity of Some European Polyporales. *Nat. Prod. Commun.* **2010**, *5*, 923–926.
23. Onar, O.; Akata, I.; Celep, G.S.; Yildirim, O. Antioxidant Activity of Extracts from the Red-Belt Conk Medicinal Mushroom, *Fomitopsis Pinicola* (Agaricomycetes), and Its Modulatory Effects on Antioxidant Enzymes. *Int. J. Med. Mushrooms* **2016**, *18*, 501–508, doi:10.1615/IntJMedMushrooms.v18.i6.40.
24. Reis, F.S.; Pereira, E.; Barros, L.; Sousa, M.J.; Martins, A.; Ferreira, I.C.F.R. Biomolecule Profiles in Inedible Wild Mushrooms with Antioxidant Value. *Molecules* **2011**, *16*, 4328–4338, doi:10.3390/molecules16064328.
25. Janjušević, L.; Karaman, M.; Šibul, F.; Tommonaro, G.; Iodice, C.; Jakovljević, D.; Pejin, B. The Lignicolous Fungus *Trametes Versicolor* (L.) Lloyd (1920): A Promising Natural Source of Antiradical and AChE Inhibitory Agents. *J. Enzyme Inhib. Med. Chem.* **2017**, *32*, 355–362, doi:10.1080/14756366.2016.1252759.
26. Matijašević, D.; Pantić, M.; Rašković, B.; Pavlović, V.; Duvnjak, D.; Sknepnek, A.; Nikšić, M. The Antibacterial Activity of *Coriolus Versicolor* Methanol Extract and Its Effect on Ultrastructural Changes of *Staphylococcus Aureus* and *Salmonella Enteritidis*. *Front. Microbiol.* **2016**, *7*, doi:10.3389/fmicb.2016.01226.
27. Adhikari, M.; Bhusal, S.; Pandey, M.; Raut, J.; Bhatt, L. Mycochemical and Nutritional Analysis of Selected Wild Mushrooms from Gaurishankar Conservation Area. *Int. J. Pharmacogn. Chinese Med.* **2019**, *3*, 000169.
28. Davey, M.W.; Montagu, M. Van; Inze, D.; Sanmartin, M.; Kanellis, A.; Smirnoff, N.; Benzie, I.J.; Strain, J.J.; Favell, D.; Fletcher, J. PlantL-Ascorbic Acid: Chemistry, Function, Metabolism, Bioavailability and Effects of Processing. *J. Sci. Food Agric.* **2000**, *80*, 825–860, doi:10.1002/(SICI)1097-0010(20000515)80:7<825::AID-JSFA598>3.0.CO;2-6.
29. Srinivasulu, C.; Ramgopal, M.; Ramanjaneyulu, G.; Anuradha, C.M.; Suresh Kumar, C. Syringic Acid (SA) – A Review of Its Occurrence, Biosynthesis, Pharmacological and Industrial Importance. *Biomed. Pharmacother.* **2018**, *108*, 547–557, doi:10.1016/j.biopha.2018.09.069.
30. Kaur, J.; Gulati, M.; Singh, S.K.; Kuppasamy, G.; Kapoor, B.; Mishra, V.; Gupta, S.; Arshad, M.F.; Porwal, O.; Jha, N.K.; et al. Discovering Multifaceted Role of Vanillic Acid beyond Flavours: Nutraceutical and Therapeutic Potential. *Trends Food Sci. Technol.* **2022**, *122*, 187–200, doi:10.1016/j.tifs.2022.02.023.
31. Espíndola, K.M.M.; Ferreira, R.G.; Narvaez, L.E.M.; Silva Rosario, A.C.R.; da Silva, A.H.M.; Silva, A.G.B.; Vieira, A.P.O.; Monteiro, M.C. Chemical and Pharmacological Aspects of Caffeic Acid and Its Activity in Hepatocarcinoma. *Front. Oncol.* **2019**, *9*, doi:10.3389/fonc.2019.00541.
32. Nowacka, N.; Nowak, R.; Drozd, M.; Olech, M.; Los, R.; Malm, A. Antibacterial, Antiradical Potential and Phenolic Compounds of Thirty-One Polish Mushrooms. *PLoS One* **2015**, *10*, e0140355, doi:10.1371/journal.pone.0140355.
33. Çayan, F.; Deveci, E.; Tel-Çayan, G.; Duru, M.E. Identification and Quantification of Phenolic Acid Compounds of Twenty-Six Mushrooms by HPLC–DAD. *J. Food Meas. Charact.* **2020**, *14*, 1690–1698, doi:10.1007/s11694-020-00417-0.

34. Kozarski, M.; Klaus, A.; van Griensven, L.; Jakovljevic, D.; Todorovic, N.; Wan-Mohtar, W.A.A.Q.I.; Vunduk, J. Mushroom β -Glucan and Polyphenol Formulations as Natural Immunity Boosters and Balancers: Nature of the Application. *Food Sci. Hum. Wellness* **2023**, *12*, 378–396, doi:10.1016/j.fshw.2022.07.040.
35. Siddeeg, A.; AlKehayez, N.M.; Abu-Hiamed, H.A.; Al-Sanea, E.A.; AL-Farga, A.M. Mode of Action and Determination of Antioxidant Activity in the Dietary Sources: An Overview. *Saudi J. Biol. Sci.* **2021**, *28*, 1633–1644, doi:10.1016/j.sjbs.2020.11.064.
36. Karaman, M.; Jovin, E.; Malbasa, R.; Matavuly, M.; Popović, M. Medicinal and Edible Lignicolous Fungi as Natural Sources of Antioxidative and Antibacterial Agents. *Phytother. Res.* **2010**, *24*, 1473–1481, doi:10.1002/ptr.2969.
37. Tumbas Šaponjac, V.; Čanadanović-Brunet, J.; Četković, G.; Djilas, S.; Četojević-Simin, D. Dried Bilberry (*Vaccinium Myrtillus* L.) Extract Fractions as Antioxidants and Cancer Cell Growth Inhibitors. *LWT - Food Sci. Technol.* **2015**, *61*, 615–621, doi:10.1016/j.lwt.2014.04.021.
38. Poli, A.; Agostoni, C.; Visioli, F. Dietary Fatty Acids and Inflammation: Focus on the n-6 Series. *Int. J. Mol. Sci.* **2023**, *24*, 4567, doi:10.3390/ijms24054567.
39. Gulcin, İ. Antioxidants and Antioxidant Methods: An Updated Overview. *Arch. Toxicol.* **2020**, *94*, 651–715, doi:10.1007/s00204-020-02689-3.
40. Kozarski, M.; Klaus, A.; Niksic, M.; Jakovljevic, D.; Helsper, J.P.F.G.; Van Griensven, L.J.L.D. Antioxidative and Immunomodulating Activities of Polysaccharide Extracts of the Medicinal Mushrooms *Agaricus Bisporus*, *Agaricus Brasiliensis*, *Ganoderma Lucidum* and *Phellinus Linteus*. *Food Chem.* **2011**, *129*, 1667–1675, doi:10.1016/j.foodchem.2011.06.029.
41. Lee, S.Y.; Baek, N.; Nam, T. Natural, Semisynthetic and Synthetic Tyrosinase Inhibitors. *J. Enzyme Inhib. Med. Chem.* **2016**, *31*, 1–13, doi:10.3109/14756366.2015.1004058.
42. Itharat, A.; Houghton, P.J.; Eno-Amooquaye, E.; Burke, P.; Sampson, J.H.; Raman, A. In Vitro Cytotoxic Activity of Thai Medicinal Plants Used Traditionally to Treat Cancer. *J. Ethnopharmacol.* **2004**, *90*, 33–38, doi:10.1016/j.jep.2003.09.014.
43. Kozarski, M.; Klaus, A.; Jakovljevic, D.; Todorovic, N.; Vunduk, J.; Petrović, P.; Niksic, M.; Vrvic, M.; van Griensven, L. Antioxidants of Edible Mushrooms. *Molecules* **2015**, *20*, 19489–19525, doi:10.3390/molecules201019489.
44. Chew, B.P.; Park, J.S. Carotenoid Action on the Immune Response. *J. Nutr.* **2004**, *134*, 257S–261S, doi:10.1093/jn/134.1.257S.
45. Carr, A.C.; Maggini, S. Vitamin C and Immune Function. *Nutrients* **2017**, *9*, doi:10.3390/nu9112121.
46. Ding, S.; Jiang, H.; Fang, J. Regulation of Immune Function by Polyphenols. *J. Immunol. Res.* **2018**, *2018*, 1–8, doi:10.1155/2018/1264074.
47. Lacroix, S.; Klicic Badoux, J.; Scott-Boyer, M.-P.; Parolo, S.; Matone, A.; Priami, C.; Morine, M.J.; Kaput, J.; Moco, S. A Computationally Driven Analysis of the Polyphenol-Protein Interactome. *Sci. Rep.* **2018**, *8*, 2232, doi:10.1038/s41598-018-20625-5.
48. Kozarski, M.; van Griensven, L. Oxidative Stress Prevention by Edible Mushrooms and Their Role in Cellular Longevity. In *Wild Mushrooms Characteristics, Nutrition, and Processing*; Dhull, S.B., Bains, A., Chawla, P., Sadh, P.K., Eds.; CRC Press, 2022; pp. 319–348 ISBN 9781003152583.
49. Grienke, U.; Zöll, M.; Peintner, U.; Rollinger, J.M. European Medicinal Polypores – A Modern View on Traditional Uses. *J. Ethnopharmacol.* **2014**, *154*, 564–583, doi:10.1016/j.jep.2014.04.030.
50. Wang, G.; He, M.; Huang, Y.; Peng, Z. Synthesis and Biological Evaluation of New Kojic Acid-1,3,4-Oxadiazole Hybrids as Tyrosinase Inhibitors and Their Application in the Anti-Browning of Fresh-Cut Mushrooms. *Food Chem.* **2023**, *409*, 135275, doi:10.1016/j.foodchem.2022.135275.
51. Fernandes, F.H.A.; Salgado, H.R.N. Gallic Acid: Review of the Methods of Determination and Quantification. *Crit. Rev. Anal. Chem.* **2016**, *46*, 257–265, doi:10.1080/10408347.2015.1095064.
52. Salau, V.F.; Erukainure, O.L.; Ibeji, C.U.; Olasehinde, T.A.; Koorbanally, N.A.; Islam, M.S. Vanillin and Vanillic Acid Modulate Antioxidant Defense System via Amelioration of Metabolic Complications Linked to Fe²⁺-Induced Brain Tissues Damage. *Metab. Brain Dis.* **2020**, *35*, 727–738, doi:10.1007/s11011-020-00545-y.
53. Jakobek, L. Interactions of Polyphenols with Carbohydrates, Lipids and Proteins. *Food Chem.* **2015**, *175*, 556–567, doi:10.1016/j.foodchem.2014.12.013.
54. Theivendren, P.; Kunjiappan, S.; Mariappa Hegde, Y.; Vellaichamy, S.; Gopal, M.; Rajan Dhramalingam, S.; Kumar, S. Importance of Protein Kinase and Its Inhibitor: A Review. In *Protein Kinases - Promising Targets for Anticancer Drug Research*; Kumar Singh, R., Ed.; IntechOpen, 2021.
55. Ferreira, I.; Barros, L.; Abreu, R. Antioxidants in Wild Mushrooms. *Curr. Med. Chem.* **2009**, *16*, 1543–1560, doi:10.2174/092986709787909587.
56. Kciuk, M.; Alam, M.; Ali, N.; Rashid, S.; Głowacka, P.; Sundaraj, R.; Celik, I.; Yahya, E.B.; Dubey, A.; Zerroug, E.; et al. Epigallocatechin-3-Gallate Therapeutic Potential in Cancer: Mechanism of Action and Clinical Implications. *Molecules* **2023**, *28*, 5246, doi:10.3390/molecules28135246.

57. Kozarski, M.; Klaus, A.; Jakovljevic, D.; Todorovic, N.; Wan-Mohtar, W.; Niksic, M. Ganoderma Lucidumas a Cosmeceutical: Antiradical Potential and Inhibitory Effect on Hyperpigmentation and Skin Extracellular Matrix Degradation Enzymes. *Arch. Biol. Sci.* **2019**, *71*, 253–264, doi:10.2298/ABS181217007K.
58. Srisuksomwong, P.; Kaenhin, L.; Mungmai, L. Collagenase and Tyrosinase Inhibitory Activities and Stability of Facial Cream Formulation Containing Cashew Leaf Extract. *Cosmetics* **2023**, *10*, 17, doi:10.3390/cosmetics10010017.
59. Zolghadri, S.; Bahrami, A.; Hassan Khan, M.T.; Munoz-Munoz, J.; Garcia-Molina, F.; Garcia-Canovas, F.; Saboury, A.A. A Comprehensive Review on Tyrosinase Inhibitors. *J. Enzyme Inhib. Med. Chem.* **2019**, *34*, 279–309, doi:10.1080/14756366.2018.1545767.
60. Cateni, F.; Gargano, M.L.; Procida, G.; Venturella, G.; Cirlincione, F.; Ferraro, V. Mycochemicals in Wild and Cultivated Mushrooms: Nutrition and Health. *Phytochem. Rev.* **2022**, *21*, 339–383, doi:10.1007/s11101-021-09748-2.
61. Mohanta, T.K. Fungi Contain Genes Associated with Flavonoid Biosynthesis Pathway. *J. Funct. Foods* **2020**, *68*, 103910, doi:10.1016/j.jff.2020.103910.
62. Shao, Y.; Guo, H.; Zhang, J.; Liu, H.; Wang, K.; Zuo, S.; Xu, P.; Xia, Z.; Zhou, Q.; Zhang, H.; et al. The Genome of the Medicinal Macrofungus Sanghuang Provides Insights Into the Synthesis of Diverse Secondary Metabolites. *Front. Microbiol.* **2020**, *10*, doi:10.3389/fmicb.2019.03035.
63. Keller, N.P. Fungal Secondary Metabolism: Regulation, Function and Drug Discovery. *Nat. Rev. Microbiol.* **2019**, *17*, 167–180, doi:10.1038/s41579-018-0121-1.
64. Rao, A. V.; Rao, L.G. Carotenoids and Human Health. *Pharmacol. Res.* **2007**, *55*, 207–216, doi:10.1016/j.phrs.2007.01.012.
65. Beta Carotene (Oral Route) Proper Use - Mayo Clinic Available online: <https://www.mayoclinic.org/drugs-supplements/beta-carotene-oral-route/proper-use/drg-20066795> (accessed on 15 November 2023).
66. Nutrients: Carotene, Beta (Mg) Available online: <https://ods.od.nih.gov/pubs/usdandb/VitA-betaCarotene-Content.pdf> (accessed on 15 November 2023).
67. Einafshar, S.; Poorazrang, H.; Farhoosh, R.; Seiedi, S.M. Antioxidant Activity of the Essential Oil and Methanolic Extract of Cumin Seed (Cuminum Cyminum). *Eur. J. Lipid Sci. Technol.* **2012**, *114*, 168–174, doi:10.1002/ejlt.201100127.
68. Anantharaman, A.; Hemachandran, H.; Priya, R.R.; Sankari, M.; Gopalakrishnan, M.; Palanisami, N.; Siva, R. Inhibitory Effect of Apocarotenoids on the Activity of Tyrosinase: Multi-Spectroscopic and Docking Studies. *J. Biosci. Bioeng.* **2016**, *121*, 13–20, doi:10.1016/j.jbiosc.2015.05.007.
69. Gloria, N.F.; Soares, N.; Brand, C.; Oliveira, F.L.; Borojevic, R.; Teodoro, A.J. Lycopene and Beta-Carotene Induce Cell-Cycle Arrest and Apoptosis in Human Breast Cancer Cell Lines. *Anticancer Res.* **2014**, *34*, 1377–1386.
70. Rowles, J.L.; Erdman, J.W. Carotenoids and Their Role in Cancer Prevention. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **2020**, *1865*, 158613, doi:10.1016/j.bbalip.2020.158613.
71. Liu, R.H. Potential Synergy of Phytochemicals in Cancer Prevention: Mechanism of Action. *J. Nutr.* **2004**, *134*, 3479S–3485S, doi:10.1093/jn/134.12.3479S.
72. Lee, S.K.; Kader, A.A. Preharvest and Postharvest Factors Influencing Vitamin C Content of Horticultural Crops. *Postharvest Biol. Technol.* **2000**, *20*, 207–220, doi:10.1016/S0925-5214(00)00133-2.
73. Gonzalez, M.J.; Miranda-Massari, J.R.; Olalde, J. Vitamin C and Mitochondrial Function in Health and Exercise. In *Molecular Nutrition and Mitochondria*; Elsevier, 2023; pp. 225–242.
74. Shterman, N.; Kupfer, B.; Moroz, C. Comparison of Transferrin Receptors, Iron Content and Isoferritin Profile in Normal and Malignant Human Breast Cell Lines. *Pathobiology* **1991**, *59*, 19–25, doi:10.1159/000163611.
75. Lui, G.Y.L.; Kovacevic, Z.; Richardson, V.; Merlot, A.M.; Kalinowski, D.S.; Richardson, D.R. Targeting Cancer by Binding Iron: Dissecting Cellular Signaling Pathways. *Oncotarget* **2015**, *6*, 18748–18779, doi:10.18632/oncotarget.4349.
76. Lim, J.Y.; Kim, D.; Kim, B.R.; Jun, J.S.; Yeom, J.S.; Park, J.S.; Seo, J.-H.; Park, C.H.; Woo, H.O.; Youn, H.-S.; et al. Vitamin C Induces Apoptosis in AGS Cells via Production of ROS of Mitochondria. *Oncol. Lett.* **2016**, *12*, 4270–4276, doi:10.3892/ol.2016.5212.
77. Uzelac, B. *Gljive Srbije i Zapadnog Balkana (in Serbian)*; BGV logik, 2009; ISBN 9788691267704.
78. Chen, B.; Ke, B.; Ye, L.; Jin, S.; Jie, F.; Zhao, L.; Wu, X. Isolation and Varietal Characterization of Ganoderma Resinaceum from Areas of Ganoderma Lucidum Production in China. *Sci. Hortic. (Amsterdam)* **2017**, *224*, 109–114, doi:10.1016/j.scienta.2017.06.002.
79. Vunduk, J.; Kozarski, M.; Djekic, I.; Tomašević, I.; Klaus, A. Effect of Modified Atmosphere Packaging on Selected Functional Characteristics of Agaricus Bisporus. *Eur. Food Res. Technol.* **2021**, *247*, 829–838, doi:10.1007/s00217-020-03666-x.
80. Kozarski, M.; Klaus, A.; Vunduk, J.; Zizak, Z.; Niksic, M.; Jakovljevic, D.; Vrvic, M.M.; Van Griensven, L.J.L.D. Nutraceutical Properties of the Methanolic Extract of Edible Mushroom Cantharellus Cibarius (Fries): Primary Mechanisms. *Food Funct.* **2015**, *6*, 1875–1886, doi:10.1039/C5FO00312A.

81. Kozarski, M.; Klaus, A.; Vunduk, J.; Nikšić, M. The Influence of Mushroom *Coriolus Versicolor* and Hazelnuts Enrichment on Antioxidant Activities and Bioactive Content of Dark Chocolate. *Food Feed Res.* **2020**, *47*, 23–32, doi:10.5937/FFR2001023K.
82. Nagata, M.; Yamashita, I. Simple Method for Simultaneous Determination of Chlorophyll and Carotenoids in Tomato Fruit. *Nippon Shokuhin Kogyo Gakkaishi* **1992**, *39*, 925–928, doi:10.3136/nskkk1962.39.925.
83. Špirović-Trifunović, B.; Vuković, G.; Čurić, A.; Bursić, V.; Petrović, A.; Marinković, D. Influence of Climate Change on Fumonizine Production. *Biljn. Lek.* **2022**, *50*, 127–139, doi:10.5937/BiljLek2202127S.
84. Šćepanović, M.; Sarić-Krsmanović, M.; Šoštarčić, V.; Brijačak, E.; Lakić, J.; Špirović Trifunović, B.; Gajić Umiljendić, J.; Radivojević, L. Inhibitory Effects of Brassicaceae Cover Crop on *Ambrosia Artemisiifolia* Germination and Early Growth. *Plants* **2021**, *10*, 794, doi:10.3390/plants10040794.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.