

TNFSF14/LIGHT, a non-canonical NF- κ B stimulus, induces the HIF pathway

Laura D'Ignazio¹, Michael Batie², and Sonia Rocha^{1,2 *}

¹Center for Gene Regulation and Expression, School of Life Sciences, University of Dundee.
DD15EH. UK.

²Department of Biochemistry, Institute of Integrative Biology, University of Liverpool.
L697ZB, UK.

*Corresponding author: srocha@liverpool.ac.uk

Abstract

Non-canonical NF- κ B signalling plays important roles in development and function of the immune system but also is deregulated in a number of inflammatory diseases. Although, NF- κ B and HIF crosstalk has been documented, this has only been described following canonical NF- κ B stimulation, involving RelA/p50 and the HIF-1 dimer. Here we report that the non-canonical inducer TNFSF14/LIGHT leads to HIF induction and activation in cancer cells. We demonstrate that only HIF-2 α is induced at the transcriptional level following non-canonical NF- κ B activation, via a mechanism dependent on the p52 subunit. Furthermore, we demonstrate that p52 can bind to the HIF-2 α gene in cells. These results indicate that non-canonical NF- κ B can lead to HIF signalling implicating HIF-2 α as one of the downstream effectors of this pathway in cells.

Keywords: HIF, NF-kappaB, EPAS1, LIGHT, TNFSF14, p52, NIK, ChIP

Introduction

NF- κ B is the name of a family of transcription factors mostly known for their function in regulation of the cellular response to infection and inflammation [1], but it also has pleiotropic function in responses to a number of cellular stresses, such as DNA damage and hypoxia [2]. Given its role in the control of inflammatory responses, it has also been widely associated with cancer, where it can act as both as tumour suppressor but also tumour promoter [3,4].

NF- κ B encompasses five genes which give rise to seven proteins, RelA/p65, RelB, cRel, NFKB1/p105 and p50, and NFKB2/p100 and p52 [5]. Due to its importance, it is kept under several strict control mechanisms, such cellular localisation, binding to inhibitor proteins and numerous negative feedback loops [6]. Activation of NF- κ B follows, in a simplified view, three main pathways: canonical, non-canonical and atypical activation pathways, which are dependent on the involvement of the upstream kinases called Inhibitor of κ B (I κ B) kinases (IKK) [2]. The canonical pathway, responding to stimuli such as TNF- α and IL-1 β , activates the IKK complex, leading to phosphorylation and inactivation of I κ B proteins, and consequent release of the NF- κ B dimers, mainly containing RelA and p50 [5]. Once translocated into the nucleus, NF- κ B subunits can receive additional control mechanisms, such as post-translational modifications [7], to induce or repress the specific and appropriate complement of target genes. In the non-canonical pathway, stimuli such B-cell activating Factor, or Lymphotoxin- β , activate the NF- κ B activating Kinase (NIK), which phosphorylates and activates IKK α , leading, in turn, to the phosphorylation and processing of p100 into p52 [8]. This event releases primarily the RelB/p52 dimers, now able to translocate into the nucleus, and together with additional stimulus-specific inputs activate or repress specific genes in cells. Atypical pathways, usual activate NF- κ B independent of IKKs, or act on nuclear NF- κ B subunits [2].

NF- κ B transcription factors are also known to crosstalk to other transcription factors such as p53, E2F1 and HIF [9-12], amongst others. HIF is an important family of transcription factors (HIF-1 α , HIF-2 α , or HIF-3 α partnered with HIF-1 β), known for their role in the response to reduction in oxygen availability or hypoxia [13]. Although HIF is mainly controlled by the activity of the oxygen sensor prolyl-hydroxylases enzymes (PHDs) [14], other mechanisms are also important to control the expression and activity of HIF [10,12,15]. We

and others have shown that in response to canonical pathway activating stimuli, NF- κ B is able to induce HIF-1 α [16-19] and HIF-1 β [18] directly, and some reports indicate that HIF-3 α is also controlled by NF- κ B [20,21]. However, so far, no direct control of HIF-2 α gene by NF- κ B subunits has been reported, although we had previously identified an indirect role for NF- κ B to induce HIF-2 α in response to TNF- α [18].

TNFSF14 or LIGHT is a non-canonical inducer of NF- κ B, analogous to Lymphotoxin- β [22], which also is able to induce the canonical pathway in parallel [23]. It has been shown to control inflammation resolution in the intestine in mice [24], but also enhance immune-mediated reduction of metastasis in colon cancer [25]. On the other hand, it increases adipose tissue inflammation [26]. Here we demonstrate that LIGHT can induce the non-canonical NF- κ B pathway in two types of cancer cells. In addition, we show that LIGHT can induce the expression and activity of HIF-1 α and HIF-2 α . Mechanistically, LIGHT induces the mRNA expression of HIF-2 α but not HIF-1 α or HIF-1 β . LIGHT-induced p52 is required for the elevation of HIF-2 α mRNA and protein. Finally, we identify specific binding sites for p52 at the HIF-2 α promoter, indicating a direct control of p52 over the HIF-2 α gene.

Material and Methods

Cell lines and Growth Conditions

All cells were maintained in Dulbecco's modified Eagle's medium (DMEM, Lonza) supplemented with 10% fetal bovine serum (FBS, Invitrogen), 1% penicillin-streptomycin (Lonza), and 1% L-glutamine (Lonza) at 5% CO₂ and 37°C for no more than 30 passages. Cells were routinely tested for mycoplasma contamination using MycoAlert Mycoplasma Detection Kit (Lonza).

Human cervix carcinoma HeLa, and human lung carcinoma A549 cell lines were obtained from the American Type Culture Collection (ATCC).

HeLa-HRE Luciferase cells were previously created in the lab, by co-transfection of 3xHRE-luciferase construct (a kind gift from G. Melillo, National Cancer Institute, National Institutes of Health, Bethesda, MD) with a puromycin-resistant construct [27].

siRNA Cell Transfection

Small interfering RNA oligonucleotides were purchased from Eurofins Genomics and used in a final concentration of 27 nM. siRNA transfections in all cell lines were performed using Interferin (Polyplus) according to manufacturer's instructions. Oligonucleotide sequences used for siRNA knockdown are: control, 5'- CAGUCGCGUUUGCGACUGGTT-3'; p100/p52, 5'- CUACGAGGGACCAGCCAAG-3'; RelB, 5'- AAUUGGAGAUCACGACGAGU-3'.

Cell Treatments

To stimulate the inflammatory response, human recombinant LIGHT (TNFSF14, Peprotech) was dissolved in sterile PBS and used at a final concentration of 100 ng/mL.

Luciferase assay

2 x 10⁵ cells stably transfected with a luciferase reporter gene were seeded in 6-well plates, stimulated with LIGHT for the indicated times, and harvested with 400 µL of Passive Lysis Buffer (Promega). Luciferase activity was measured according to manufacturer's instructions (Luciferase Assay System, Promega), and normalised to protein concentration (Bradford, BioRad). All experiments were performed a minimum of three times before calculating average and standard error of the means.

RNA extraction and Real Time quantitative PCR analysis

PeqGOLD total RNA kit (Pepqlab) or PureLink RNA Mini Kit (Life Technologies) were used to extract total RNA from cells according to the manufacturer's instructions. RNA was converted to cDNA using Quantitect Reverse Transcription Kit (Qiagen) or First Strand cDNA Synthesis kit (Thermo Fisher Scientific). For quantitative PCR, Brilliant II Sybr green kit (Stratagene/Agilent), and recommended MX3005P 96-well skirted plates were used to analyse samples on the Mx3005P qPCR platform (Stratagene/Agilent). Alternatively, PerfeCTa Sybr Green FastMix (Quanta Bio) with ROX dye added in 1:250 ratio, and recommended Microamp Optical 96-well reaction plates were used to analyse samples on the QuantStudio 6 Flex qPCR platform (Applied Biosystem). Actin was used as normalising gene. RT-PCR results were analysed by the $\Delta\Delta C_t$ method. The primers used for gene expression analysis by RT-PCR are:

Actin, For: 5'- CTGGGAGTGGGTGGAGGC-3', Rev: 5'- TCAACTGGTCTCAAGTCAGTG-3'.
HIF-1 α , For: 5'- ATAAAGTCTGCAACATGGAAGGT-3', Rev: 5'- TTTGATGGGTGAGGAATGGGTT-3'.
HIF-1 β , For: 5'- CAAGCCCCTTGAGAAGTCAG-3', Rev: 5'- GAGGGGCTAGGCCACTATTC-3'.
HIF-2 α , For: 5'- TTTGATGTGGAAACGGATGA-3', Rev: 5'- GGAACCTGCTCTTGCTGTTC-3'.
PHD2, For: 5'- GAAAGCCATGGTTGCTTGTT-3', Rev: 5'- TGTCTTCTGGAAAAATTTCG-3'.
RANTES, For: 5'- GTCGTCTTTGTCACCCGAAAG-3', Rev: 5'- TCCCGAACCCATTTCTTCTCT-3'.

Protein Lysis

Cells were lysed using 100 μ L of whole cell protein lysis buffer (20 mM Tris pH 7.6, 150 mM NaCl, 0.75% NP-40, 5 mM NaF, 500 μ M Na₃VO₄, and 1 Pierce Protease Inhibitor Mini Tablet [EDTA Free, ThermoFisher Scientific] per 10 mL of buffer). Upon collection, cells were kept on ice for 30 minutes before centrifugation at 16060 x g at 4°C for 15 minutes. The supernatant was collected and stored at -80°C.

Western Blotting

Protein concentration was determined using Bradford (BioRad) method. 20-30 μ g of protein was prepared in 2x SDS loading buffer (100 mM Tris-HCl pH 6.8, 20% glycerol, 4%

SDS, 200 mM DTT, and Bromophenol Blue), and incubated for 5-10 minutes at 105°C. Western blotting was performed as described in [18]. Briefly, samples were loaded into an SDS-page gel (Tris-HCl poly-acrylamide gel) previously prepared and run at 80-120 volts in Running Buffer (25 mM Tris, 0.195 M glycine, and 0.1% SDS). The gel was then transferred in a semi-dry transfer (BioRad) into a PVDF membrane (Millipore) for 1.5-2 hours at 15 volts/0.80 mA in Transfer Buffer (50 mM Tris, 40 mM glycine, 0.001% SDS and 10% methanol). Then, the membrane was blocked with 10% Milk in TBS-tween buffer (20 mM Tris pH 7.6, 150 mM NaCl, 0.1% Tween) for 10 minutes, followed by three 5 minutes washes with TBS-tween buffer. Membranes were incubated with primary antibodies for 1 hour at room temperature or overnight at 4°C, in accordance with primary antibodies' manufacturer instructions. Membranes were then washed three times with TBS-tween and incubated with the appropriate secondary HRP antibody. After washes, membranes were developed using ECL solution (Pierce). Primary antibodies purchased from Cell signalling were: HIF-1 α -OH Pro564 (#3434), HIF-1 β (#5537), HIF-2 α (#7096), NDRG1 (#5196), NIK (#4994), RelB (#10544). Primary antibodies against HIF-1 α (#610958) were from BD Biosciences, anti-p100/p52 from Millipore (#05-361), anti-PHD2 from Bethyl (#A300-322A), anti-Cyclin D1 from Abcam (ab137875), anti-CITED-2 from R&D System (#MAB5005), and anti- β -Actin from Proteintech (#66009-1-1g).

ChIP-Sequencing data analysis

L1236 p52 ChIP sequencing data [28] was downloaded from the NCBI GEO database (GSE63736). Coverage tracks were generated using the R Bioconductor package, Gviz [29].

Chromatin Immunoprecipitation

Chromatin Immunoprecipitation (ChIP) was performed adapting the protocol described in [30]. Cells were plated and grown to 70-80% confluency on 150 mm plates in 16 mL of appropriate culturing media. When necessary cells were treated with LIGHT for 4 hours. Then, proteins and chromatin were cross-linked with 1% formaldehyde at 37°C for 10 minutes. To quench the cross-linking, glycine was added to a final concentration of 0.125 M for 5 minutes at 37°C. Cells were washed twice with ice-cold PBS, then scraped and centrifuged at 1000 rpm in a Beckman Coulter's Allegra X-12 benchtop centrifuge for 5 minutes. The supernatant was removed and the pellet resuspended in 400 μ L of ChIP lysis buffer (1% SDS, 10 mM EDTA,

50 mM Tris-HCl pH 8.1, and 1 Pierce Protease Inhibitor Mini Tablet [EDTA Free, ThermoFisher Scientific], up to 10 mL of buffer), before being snap-frozen in dry ice and stored at -80°C. Once thawed on ice, 200 µL aliquots of each sample were transferred into 1.5 mL TPX Polymethylpentene (PMP) tubes (Diagenode) to improve sonication and shearing efficiency. Samples were sonicated in Bioruptor NGS (Diagenode) at 4°C for five cycles of 30 seconds ON/30 seconds OFF, at high intensity amplitude. This sonication procedure was repeated four times. Supernatants were recovered by centrifugation at maximum speed in a benchtop centrifuge for 10 minutes at 4°C prior storage of 10% of each sample as input. Remaining samples were split into 120 µL aliquots before being diluted 10 fold in ChIP Dilution Buffer (1% Triton X-100, 2 mM EDTA, 150 mM NaCl, 20 mM Tris-HCl pH 8.1). Diluted samples were pre-cleared for 2 hours at 4°C by incubation with 2 µg of sheared salmon sperm DNA and 20 µL of protein G-Sepharose (50% slurry), previously washed in cold PBS. Immunoprecipitation was performed overnight on the remaining samples with 2 µg of anti-p52 (santa cruz, #sc-848) or Normal Rabbit IgG (Sigma, #I5381) as negative control, and 0.1% of BRIJ-35 detergent. The following day, immune complexes were captured by incubation with 30 µL of protein G-Sepharose (50% slurry, previously washed with cold PBS) and 2 µg of salmon sperm DNA for 2.5 hours at 4°C. The immunoprecipitates were washed sequentially for 5 minutes at 4°C in 1 mL of cold Wash Buffer 1 (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.1, 150 mM NaCl), 1mL of cold Wash Buffer 2 (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.1, 500 mM NaCl), and 1 mL of cold Wash Buffer 3 (0.25 M LiCl, 1% Nonidet P-40, 1% deoxycholate, 1 mM EDTA, 10 mM Tris-HCl pH 8.1). Next, beads were washed twice with 500 µL of Tris-EDTA (TE) buffer.

Chromatin reverse cross-linking and DNA elution were performed using the IPure kit v2 (Diagenode) following manufacturer's instructions. Briefly, 10% inputs and beads were resuspended in 90 µL and 100 µL of Elution Buffer mix, respectively. The elution buffer mix was prepared with 115.4 µL of Buffer A and 4.6 µL of Buffer B per sample. All samples were incubated overnight at 65°C on a thermomixer with continuous shaking at 300rpm. The following day, supernatants were recovered by centrifugation at 1000rpm for 1 minute in a benchtop centrifuge. 2 µL of carrier, 100 µL of 100% isopropanol and 10 µL of magnetic beads were added to each input and immunoprecipitate sample, then incubated on a rotating wheel (40rpm) for 10 minutes at room temperature. After a quick centrifugation, tubes were placed into a magnetic rack for separation of buffer, then discarded, and beads. Captured beads were gently mixed to 100 µL of Wash Buffer 1 (previously diluted with 100% isopropanol with 1:1 ratio), prior incubation on rotating wheel (40rpm) for 5 minutes at room temperature.

Following another step of buffer separation, beads were gently mixed to 100 μ L of Wash Buffer 2 (previously diluted with 100% isopropanol with 1:1 ratio), and incubated on rotating wheel (40rpm) for 5 minutes at room temperature. Finally, DNA elution was performed adding to captured beads 25 μ L of Buffer C and incubating tubes on a rotating wheel (40rpm) for 15 minutes at room temperature. After separation on magnetic rack, the first fraction of eluted DNA was transferred into a new storage tube, while captured beads were subjected to a second step of DNA elution by adding other 25 μ L of Buffer C, to obtain a final volume of 50 μ L of purified DNA.

3 μ L DNA were used for RT-PCR analysis with primers specifically designed on promoter regions of interest. Specifically, for HIF-2 α -2683 κ B site, For: 5'-AAGGTGCGGTGGCTTAT-3', Rev: 5'-GAACTCCTGGCCTCATGT-3'; for HIF-2 α -2534 κ B site, For: 5'-GGTGGTGCACATCTGTAGT-3', Rev: 5'-CAGGTTGGAGCGGAGTG-3'; for HIF-2 α -1278 κ B site, For: 5'-CCCAACCCTTTCTGTGTA-3', Rev: 5'-CAGAAGCGATTTGAAGAGAAG-3'.

Statistical Analysis

Means, standard deviations and standard error means were calculated prior performing Student t tests on a minimum of three independent experiments and calculating p-values. * = $p < 0.050$; ** = $p < 0.010$ and *** = $p < 0.001$.

Results

To determine if LIGHT was able to activate NF- κ B in cancer cells, we treated HeLa cells with 100 ng/mL LIGHT for different time periods and analysed markers of non-canonical NF- κ B pathway activation (Figure 1A-B). LIGHT treatment readily results in accumulation of NIK, with concomitant increases in the levels of p52 and RelB, as well as one of p52 known targets in cancer cells, Cyclin D1 [30,31]. We could also demonstrate that RANTES (CCL5), a known target of non-canonical NF- κ B [32], was robustly induced in HeLa cells following LIGHT treatment (Figure 1B).

Given our interest in how NF- κ B and HIF crosstalk, we next determined if LIGHT could induce a HIF transcriptional response in HeLa cells in normoxia (Figure 1C). To this end, we used a generic HIF-dependent gene reporter assay in HeLa cells, treated with LIGHT (Figure 1C). This analysis revealed a small but consistent induction of HIF-dependent luciferase activity upon treatment with LIGHT (Figure 1C). To validate this finding, we analysed levels of HIF proteins and some of its targets in two different cancer cell lines, following treatment with LIGHT (Figure 1D-E). Levels of HIF-1 α , HIF-1 β and HIF-2 α were induced by LIGHT treatment in both cell lines (Figure 1D). This was not a result of PHD inactivation since we did not detect any reduction in the levels of hydroxylated HIF-1 α (Figure 1D). Importantly, we could demonstrate that increased HIF levels resulted in increased levels of some of its targets following LIGHT treatment (Figure 1E-F). These included HIF-1 targets such as PHD2 and HIF-2 genes such as CITED-2 and NDRG1. These results demonstrate the LIGHT is able to activate both NF- κ B and HIF transcription factors.

To determine the mechanism behind LIGHT-mediated induction of HIF levels and activity, we analysed levels of HIF subunit mRNA in HeLa cells treated with LIGHT for different time periods (Figure 2). Although we could not detect any consistent or statistically significant increases in HIF-1 α (Figure 2A) or HIF-1 β (Figure 2B) mRNA levels in HeLa cells treated with LIGHT, HIF-2 α mRNA levels were induced in a statistically significant manner in such conditions (Figure 2C). This suggests that LIGHT was inducing HIF-2 α transcriptionally, while inducing HIF-1 β and HIF-1 α via an indirect mechanism.

Given that p52 is strongly induced by LIGHT in our cells, we next investigated if p52 was contributing to LIGHT-induced HIF levels and activity under conditions of LIGHT treatment. To address this question, we depleted p52 in HeLa cells by siRNA and repeated the treatment with LIGHT (Figure 3A-B). mRNA analysis revealed a strict requirement for p52 in

the LIGHT-mediated induction of HIF-2 α levels (Figure 3B, Sup. Figure S1A). p52 is also required for LIGHT-induced HIF-2 α protein and activity in HeLa cells (Figure 3B). Reduced levels of HIF-2 α targets CITED-2 and NDRG1 were observed in the absence of p52 despite treatment with LIGHT (Figure 3B). Interestingly, HIF-1 β levels were also decreased in the absence of p52, while HIF-1 α levels were elevated (Figure 3B). Given that p52 activity is partly dependent on its partner RelB, we extended our analysis to investigate RelB's contribution to LIGHT-induced HIF levels and activity (Figure 3C). In this case, RelB depletion resulted in reduced levels of all HIF subunits and also targets we investigated. This suggests that p52 and RelB have joint functions but also independent functions of each other in the control of different HIF subunits.

In view of the strong effect on HIF-2 α mRNA we had observed with p52 depletion, we next investigated if there was a direct link between these two genes. Due to the availability of an increasing number of publicly available ChIP-Sequencing datasets, we interrogated the NCBI GEO database for p52 binding sites at the HIF-2 α gene. Despite the limited availability for p52 ChIP-sequencing results, we were able to determine that in lymphoma cells, p52 has been shown to bind to multiple sites in the HIF-2 α gene[28]. These binding sites are downstream of the HIF-2 α promoter, and whilst one binding site is located near the transcription start site (TSS) of isoform 9 (EPAS1-209(ensembl transcript name)), this isoform does not result in a protein product (Figure 4A). As such, we used bioinformatics analysis utilising different software tools, which revealed a variety of putative κ B binding sites present at the HIF-2 α promoter (Figure 4B). To determine if indeed p52 is able to bind to the HIF-2 α promoter, we designed several primers to encompass some of the predicted binding sites. As an additional control, we also included siRNA depletion of p52, to really demonstrate specificity of the assay. Using this approach, we could see that analysis of -2683 site on the HIF-2 α promoter did not identify any specific and significant binding by p52, as the signal obtained was not eliminated following p52 depletion (Sup. Figure S2A). Analysis of -2534 site revealed a specific binding of p52 which was induced upon LIGHT treatment (Figure 4C, top panels). Similar results were also obtained when we analysed the -1278 site, where we can detect a specific signal for p52 binding which is present following treatment with LIGHT (Figure 4C, lower panels). Taken together these results show a direct binding of p52 to the HIF-2 α promoter, which we have demonstrated is required for HIF-2 α mRNA and protein expression following LIGHT stimulation.

Discussion

In this report, we demonstrate that HIF-2 α gene is responsive to the activation of non-canonical NF- κ B subunits, p52 and RelB, when LIGHT is used as a stimulus. Although other HIF subunits have been shown to be directly controlled by NF- κ B [16,18], these have been as a consequence of activation of the canonical NF- κ B pathways and have not shown any direct effect on HIF-2 α .

Our results demonstrate that non-canonical activation of NF- κ B is also able to induce expression and activity of HIF transcription factors, but in this case, the effect is mediated by p52 on the HIF-2 α gene. This has implications for situations where non-canonical signalling is high, such as inflammatory conditions and pathogenic infections [33], since this would result in a “pseudo-hypoxic” environment, with HIF transcription factors being activated. In fact, studies in patients with *Helicobacter pylori*, a pathogen that can activate non-canonical NF- κ B [33], have shown increased levels of HIF-2 α in patient biopsies [34]. However, these studies did not determine the mechanism behind the increased level of HIF-2 α expression, and as such more work is necessary to determine, if the mechanism we identified in our study is present under such conditions.

Non-canonical NF- κ B is important for B cell development and expression of Germinal Centres [33], however, whether HIF-2 α has any involvement in these functions has not been reported or formally addressed. Interestingly, a recent study revealed that the HIF system, including VHL and HIF-1 α and HIF-2 α , are involved in germinal centre functional regulation, in particular regarding antibody quality [35]. These findings, together with the results presented here suggest a potential functional crosstalk occurring between non-canonical NF- κ B signalling and HIF. In addition, it is known that HIF-2 α is expressed in B cell lymphoma [36,37]. As such, our findings raise two possibilities, one that HIF-2 α contributes to non-canonical NF- κ B signalling in normal B cells, or alternatively, HIF-2 α is expressed under pathological conditions of non-canonical NF- κ B activation.

Very little is known about HIF-2 α gene regulation. Our own studies, trying to address this lack of knowledge, have found that HIF-2 α is regulated by E2F1 [36] and more recently by SP1 and the transcriptional repressor complex SIN3A-HDAC, via the action of SINHCAF [38]. Although no direct interaction between E2F1 and p52 has been described so far, both of them are required for cell cycle progression [30,36], and as such are thought to be part of similar

network. Understanding, the differential regulation of the HIF genes, will help to determine how the same transcription factor can have opposing roles in different tissues. For example, while it is clear that HIF-2 α acts as an oncogene in renal cancer [39], it can act as a tumour suppressor in sarcoma [40] and possibly other tissues as well [38]. Investigating alternative regulatory mechanisms to classic oxygen regulation will help identify mechanisms responsible for tissue specific actions of these transcription factors.

Supplementary Material

This file contains two supplementary figures and legends. Figure S1. Validation of p52 siRNA-mediated knockdown. Figure S2. Validation of NF- κ B subunit, p52, ChIP specificity at the HIF-2 α promoter.

Acknowledgments

This work was funded by CRUK Senior Fellowship to SR (C99667/A12918), a Wellcome Trust Strategic Award (097945/B/11/Z), a Wellcome Trust PhD studentship to LDI, and by the University of Liverpool.

Author contributions statement

LDI performed all the experiments and help SR to design the project. LDI, MB and SR analysed the data and wrote the manuscript.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Park, M.H.; Hong, J.T. Roles of nf-kappab in cancer and inflammatory diseases and their therapeutic approaches. *Cells* **2016**, *5*.
2. Perkins, N.D. The diverse and complex roles of nf-kappab subunits in cancer. *Nature reviews. Cancer* **2012**, *12*, 121-132.
3. Hanahan, D.; Weinberg, R.A. Hallmarks of cancer: The next generation. *Cell* **2011**, *144*, 646-674.
4. Xia, Y.; Shen, S.; Verma, I.M. Nf-kappab, an active player in human cancers. *Cancer immunology research* **2014**, *2*, 823-830.
5. Gilmore, T.D. Introduction to nf-kappab: Players, pathways, perspectives. *Oncogene* **2006**, *25*, 6680-6684.
6. Renner, F.; Schmitz, M.L. Autoregulatory feedback loops terminating the nf-kappab response. *Trends in biochemical sciences* **2009**, *34*, 128-135.
7. Perkins, N.D. Post-translational modifications regulating the activity and function of the nuclear factor kappa b pathway. *Oncogene* **2006**, *25*, 6717-6730.
8. Sun, S.C. The noncanonical nf-kappab pathway. *Immunological reviews* **2012**, *246*, 125-140.
9. Cummins, E.P.; Taylor, C.T. Hypoxia-responsive transcription factors. *Pflugers Arch* **2005**, *450*, 363-371.
10. Kenneth, N.S.; Rocha, S. Regulation of gene expression by hypoxia. *The Biochemical journal* **2008**, *414*, 19-29.
11. D'Ignazio, L.; Bandarra, D.; Rocha, S. Nf-kappab and hif crosstalk in immune responses. *The FEBS journal* **2016**, *283*, 413-424.
12. D'Ignazio, L.; Batie, M.; Rocha, S. Hypoxia and inflammation in cancer, focus on hif and nf-kappab. *Biomedicines* **2017**, *5*.
13. Rocha, S. Gene regulation under low oxygen: Holding your breath for transcription. *Trends in biochemical sciences* **2007**, *32*, 389-397.
14. Kaelin, W.G., Jr. The von hippel-lindau protein, hif hydroxylation, and oxygen sensing. *Biochemical and biophysical research communications* **2005**, *338*, 627-638.
15. Bremm, A.; Moniz, S.; Mader, J.; Rocha, S.; Komander, D. Cezanne (otud7b) regulates hif-1alpha homeostasis in a proteasome-independent manner. *EMBO reports* **2014**, *15*, 1268-1277.
16. van Uden, P.; Kenneth, N.S.; Rocha, S. Regulation of hypoxia-inducible factor-1alpha by nf-kappab. *The Biochemical journal* **2008**, *412*, 477-484.

17. Rius, J.; Guma, M.; Schachtrup, C.; Akassoglou, K.; Zinkernagel, A.S.; Nizet, V.; Johnson, R.S.; Haddad, G.G.; Karin, M. Nf-kappab links innate immunity to the hypoxic response through transcriptional regulation of hif-1alpha. *Nature* **2008**, *453*, 807-811.
18. van Uden, P.; Kenneth, N.S.; Webster, R.; Muller, H.A.; Mudie, S.; Rocha, S. Evolutionary conserved regulation of hif-1beta by nf-kappab. *PLoS genetics* **2011**, *7*, e1001285.
19. Bandarra, D.; Biddlestone, J.; Mudie, S.; Muller, H.A.; Rocha, S. Hif-1alpha restricts nf-kappab-dependent gene expression to control innate immunity signals. *Disease models & mechanisms* **2015**, *8*, 169-181.
20. Augstein, A.; Poitz, D.M.; Braun-Dullaeus, R.C.; Strasser, R.H.; Schmeisser, A. Cell-specific and hypoxia-dependent regulation of human hif-3alpha: Inhibition of the expression of hif target genes in vascular cells. *Cellular and molecular life sciences : CMLS* **2011**, *68*, 2627-2642.
21. Kumar, H.; Lim, J.H.; Kim, I.S.; Choi, D.K. Differential regulation of hif-3alpha in lps-induced bv-2 microglial cells: Comparison and characterization with hif-1alpha. *Brain Res* **2015**, *1610*, 33-41.
22. Mauri, D.N.; Ebner, R.; Montgomery, R.I.; Kochel, K.D.; Cheung, T.C.; Yu, G.L.; Ruben, S.; Murphy, M.; Eisenberg, R.J.; Cohen, G.H., *et al.* Light, a new member of the tn timer superfamily, and lymphotoxin alpha are ligands for herpesvirus entry mediator. *Immunity* **1998**, *8*, 21-30.
23. Jin, H.R.; Jin, X.; Lee, J.J. Zinc-finger protein 91 plays a key role in light-induced activation of non-canonical nf-kappab pathway. *Biochemical and biophysical research communications* **2010**, *400*, 581-586.
24. Krause, P.; Zahner, S.P.; Kim, G.; Shaikh, R.B.; Steinberg, M.W.; Kronenberg, M. The tumor necrosis factor family member tn timer 14 (light) is required for resolution of intestinal inflammation in mice. *Gastroenterology* **2014**, *146*, 1752-1762 e1754.
25. Qiao, G.; Qin, J.; Kunda, N.; Calata, J.F.; Mahmud, D.L.; Gann, P.; Fu, Y.X.; Rosenberg, S.A.; Prabhakar, B.S.; Maker, A.V. Light elevation enhances immune eradication of colon cancer metastases. *Cancer research* **2017**, *77*, 1880-1891.
26. Kim, H.M.; Jeong, C.S.; Choi, H.S.; Kawada, T.; Yu, R. Light/timer 14 enhances adipose tissue inflammatory responses through its interaction with hvem. *FEBS letters* **2011**, *585*, 579-584.
27. Melvin, A.; Mudie, S.; Rocha, S. Further insights into the mechanism of hypoxia-induced nf-kappab. [corrected]. *Cell cycle* **2011**, *10*, 879-882.
28. de Oliveira, K.A.; Kaergel, E.; Heinig, M.; Fontaine, J.F.; Patone, G.; Muro, E.M.; Mathas, S.; Hummel, M.; Andrade-Navarro, M.A.; Hubner, N., *et al.* A roadmap of constitutive nf-kappab activity in hodgkin lymphoma: Dominant roles of p50 and p52 revealed by genome-wide analyses. *Genome Med* **2016**, *8*, 28.

29. Hahne, F.; Ivanek, R. Visualizing genomic data using gviz and bioconductor. *Methods Mol Biol* **2016**, *1418*, 335-351.
30. Schumm, K.; Rocha, S.; Caamano, J.; Perkins, N.D. Regulation of p53 tumour suppressor target gene expression by the p52 nf-kappab subunit. *The EMBO journal* **2006**, *25*, 4820-4832.
31. Rocha, S.; Martin, A.M.; Meek, D.W.; Perkins, N.D. P53 represses cyclin d1 transcription through down regulation of bcl-3 and inducing increased association of the p52 nf-kappab subunit with histone deacetylase 1. *Molecular and cellular biology* **2003**, *23*, 4713-4727.
32. Schneider, K.; Potter, K.G.; Ware, C.F. Lymphotoxin and light signaling pathways and target genes. *Immunological reviews* **2004**, *202*, 49-66.
33. Sun, S.C. The non-canonical nf-kappab pathway in immunity and inflammation. *Nature reviews. Immunology* **2017**, *17*, 545-558.
34. Griffiths, E.A.; Pritchard, S.A.; McGrath, S.M.; Valentine, H.R.; Price, P.M.; Welch, I.M.; West, C.M. Hypoxia-associated markers in gastric carcinogenesis and hif-2alpha in gastric and gastro-oesophageal cancer prognosis. *British journal of cancer* **2008**, *98*, 965-973.
35. Cho, S.H.; Raybuck, A.L.; Stengel, K.; Wei, M.; Beck, T.C.; Volanakis, E.; Thomas, J.W.; Hiebert, S.; Haase, V.H.; Boothby, M.R. Germinal centre hypoxia and regulation of antibody qualities by a hypoxia response system. *Nature* **2016**, *537*, 234-238.
36. Moniz, S.; Bandarra, D.; Biddlestone, J.; Campbell, K.J.; Komander, D.; Bremm, A.; Rocha, S. Cezanne regulates e2f1-dependent hif2alpha expression. *Journal of cell science* **2015**, *128*, 3082-3093.
37. Burrows, N.; Maxwell, P.H. Hypoxia and b cells. *Exp Cell Res* **2017**, *356*, 197-203.
38. Biddlestone, J.; Batie, M.; Bandarra, D.; Munoz, I.; Rocha, S. Sin3a/fam60a and sin3a specifically repress hif 2alpha expression. *The Biochemical journal* **2018**.
39. Martinez-Saez, O.; Gajate Borau, P.; Alonso-Gordoa, T.; Molina-Cerrillo, J.; Grande, E. Targeting hif-2 alpha in clear cell renal cell carcinoma: A promising therapeutic strategy. *Crit Rev Oncol Hematol* **2017**, *111*, 117-123.
40. Nakazawa, M.S.; Eisinger-Mathason, T.S.; Sadri, N.; Ochocki, J.D.; Gade, T.P.; Amin, R.K.; Simon, M.C. Epigenetic re-expression of hif-2alpha suppresses soft tissue sarcoma growth. *Nature communications* **2016**, *7*, 10539.


Figure Legends:**Figure 1. LIGHT, a non-canonical NF- κ B inducer, induces HIF expression and activity.**

A) HeLa cells were treated with 100 ng/mL LIGHT for 0, 4 and 24 hours. Whole cell lysates were prepared and analysed by western blot for the indicated non-canonical NF- κ B pathway regulators, subunits and target genes. β -Actin was used as loading control. **B)** HeLa cells were treated with 100 ng/mL LIGHT for 0, 4 and 24 hours prior mRNA extraction and RT-qPCR analysis for RANTES transcript, normalised to Actin mRNA levels. All the values were normalised to the untreated sample. The graphs depicts mean and SEM determined from at least three independent experiments. Student t-test analysis was performed and significance determined as follows: * $p \leq 0.05$, *** $p \leq 0.001$. **C)** HeLa cells, stably transfected with HRE luciferase reporter, were treated with 100 ng/mL LIGHT for 8 and 24 hours prior to luciferase measurements. All the values were normalised to the untreated sample. Graph depicts mean and SEM of a minimum of three independent experiments. Student t-test analysis was performed and significance determined as follows: ns = not significant, * $p \leq 0.05$, *** $p \leq 0.001$. **D)** HeLa and A549 cells were treated with 100 ng/mL LIGHT for 0, 4 and 24 hours prior collection of whole cell lysates and western blot analysis for the depicted proteins. β -Actin was used as loading control. **E)** HeLa and A549 cells were treated with 100 ng/mL LIGHT for 0, 4 and 24 hours. Then, whole cell lysates were collected and western blot analysis was performed for a subset of HIF- α specific targets. β -Actin was used as loading control. **F)** HeLa cells were treated with 100 ng/mL LIGHT for 0, 4 and 24 hours, prior mRNA extraction and RT-qPCR analysis for PHD2 gene transcript, normalised to Actin mRNA levels. All the values were normalised to the untreated samples. The graph depicts mean and SEM determined from at least three independent experiments. Student t-test analysis was performed and significance determined as follows: ** $p \leq 0.01$.


Figure 2. LIGHT induces the expression of HIF-2 α mRNA. mRNA was extracted from HeLa cells, previously treated with 100 ng/mL LIGHT for 0, 4 and 24 hours, and RT-qPCR analysis was performed for **A)** HIF-1 α , **B)** HIF-1 β and **C)** HIF-2 α , normalising to Actin mRNA levels. All the values were normalised to the untreated samples. All graphs indicate mean and SEM determined from at least three independent experiments. Student t-test analysis was performed and significance determined as follows: ns = not significant, ** $p \leq 0.01$, *** $p \leq 0.001$.

Figure 3. LIGHT-induced HIF-2 α expression is p52 dependent. **A)** HeLa cells were transfected with siRNA control and p52 oligonucleotides and treated with 100 ng/mL LIGHT for the 0, 4, and 24 hours prior mRNA extraction. RT-qPCR analysis for HIF-2 α gene transcript was performed, normalising to Actin mRNA levels. All the values were normalised to the control treated with LIGHT for 4 hours. The graph shows mean and SEM determined from at least three independent experiments. Student t-test analysis was performed and significance determined as follows: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$. **B-C)** HeLa cells were transfected with siRNA control and p52 (**B**) or RelB (**C**) oligonucleotides and treated with 100 ng/mL LIGHT for 0, 4 and 24 hours. Whole cell lysates were prepared and analysed by western blot to investigate the protein expression of the HIF subunits and a subset of HIF specific targets following treatment. β -Actin was used as loading control.


Figure 4. The NF- κ B subunit, p52, specifically binds to multiple sites on the HIF-2 α promoter. **A)** Coverage tracks and peaks from L1236 p52 ChIP-seq data at the HIF-2 α gene. Peak start and end distances from the HIF-2 α transcription start site (TSS) are also shown. (*Numbers in brackets*) = distances from HIF-2 α isoform 9 (EPAS-209 (ensembl transcript name)) TSS. **B)** Schematic diagram showing annotations of putative κ B binding sites (dashed line pentagons) identified on the promoter region (\pm 3000 bp from TSS) of HIF-2 α gene through bioinformatics analysis conducted using ALGGEN PROMO software. 4 out of 19 site were subjected to further analysis, to be confirmed as binding sites for the NF- κ B subunit p52. **C)** HeLa cells were transfected with siRNA control and p52 oligonucleotides for 72 hours prior to cross-linking and lysis (left panels) or treated with 100 ng/mL LIGHT for the 0 and 4 hours prior to cross-linking and lysis (right panels). Chromatin immunoprecipitations (ChIPs) were performed for the levels of p52 present at the indicated putative κ B sites on the HIF-2 α promoter. Rabbit IgG was used as antibody control. All the graphs depict mean and SEM of a minimum of three independent experiments. Student t test was performed and significance determined as follows: ns = not significant, * $p \leq 0.05$, *** $p \leq 0.001$.



A



B



C

