

Article

Not peer-reviewed version

Some Results in Classes Of Neutrosophic Graphs

[Mohammadesmail Nikfar](#)*

Posted Date: 17 March 2022

doi: 10.20944/preprints202203.0248.v1

Keywords: Neutrosophic Quasi-Order; Neutrosophic Quasi-Size; Neutrosophic Quasi-Number; Neutrosophic Quasi-Co-Number; Neutrosophic Co-t-Neighborhood



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Some Results in Classes Of Neutrosophic Graphs

Mohammadesmail Nikfar

Independent Researcher

DrHenryGarrett@gmail.com

Twitter's ID: @DrHenryGarrett | ©DrHenryGarrett.wordpress.com

Abstract

New setting is introduced to study co-neighborhood, neutrosophic t-neighborhood, neutrosophic quasi-vertex set, neutrosophic quasi-order, neutrosophic neighborhood, neutrosophic co-t-neighborhood, neutrosophic quasi-edge set, neutrosophic quasi-size, Neutrosophic number, neutrosophic co-neighborhood, co-neutrosophic number, quasi-number and quasi-co-number. Some classes of neutrosophic graphs are investigated.

Keywords: Neutrosophic Quasi-Order, Neutrosophic Quasi-Size, Neutrosophic Quasi-Number, Neutrosophic Quasi-Co-Number, Neutrosophic Co-t-Neighborhood

AMS Subject Classification: 05C17, 05C22, 05E45

1 Background

Fuzzy set in **Ref.** [16], neutrosophic set in **Ref.** [3], related definitions of other sets in **Refs.** [3, 13, 15], graphs and new notions on them in **Refs.** [1, 4, 8–11, 14, 17], neutrosophic graphs in **Ref.** [5], studies on neutrosophic graphs in **Ref.** [2], relevant definitions of other graphs based on fuzzy graphs in **Ref.** [12], related definitions of other graphs based on neutrosophic graphs in **Ref.** [6], are proposed. Also, some studies and researches about neutrosophic graphs, are proposed as a book in **Ref.** [7].

2 Preliminaries

In this subsection, basic material which is used in this article, is presented. Also, new ideas and their clarifications are elicited.

Basic idea is about the model which is used. First definition introduces basic model.

Definition 2.1. (Graph).

$G = (V, E)$ is called a **graph** if V is a set of objects and E is a subset of $V \times V$ (E is a set of 2-subsets of V) where V is called **vertex set** and E is called **edge set**. Every two vertices have been corresponded to at most one edge.

Neutrosophic graph is the foundation of results in this paper which is defined as follows. Also, some related notions are demonstrated.

Definition 2.2. (Neutrosophic Graph And Its Special Case).

$NTG = (V, E, \sigma = (\sigma_1, \sigma_2, \sigma_3), \mu = (\mu_1, \mu_2, \mu_3))$ is called a **neutrosophic graph** if it's graph, $\sigma_i : V \rightarrow [0, 1]$, and $\mu_i : E \rightarrow [0, 1]$. We add one condition on it and we use **special case** of neutrosophic graph but with same name. The added condition is as follows, for every $v_i v_j \in E$,

$$\mu(v_i v_j) \leq \sigma(v_i) \wedge \sigma(v_j).$$

(i) : σ is called **neutrosophic vertex set**.

(ii) : μ is called **neutrosophic edge set**.

(iii) : $|V|$ is called **order** of NTG and it's denoted by $\mathcal{O}(NTG)$.

(iv) : $\sum_{v \in V} \sigma(v)$ is called **neutrosophic order** of NTG and it's denoted by $\mathcal{O}_n(NTG)$.

(v) : $|E|$ is called **size** of NTG and it's denoted by $\mathcal{S}(NTG)$.

(vi) : $\sum_{e \in E} \sum_{i=1}^3 \mu_i(e)$ is called **neutrosophic size** of NTG and it's denoted by $\mathcal{S}_n(NTG)$.

Some classes of well-known neutrosophic graphs are defined. These classes of neutrosophic graphs are used to form this study and the most results are about them.

Definition 2.3. Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then

(i) : a sequence of vertices $P : x_0, x_1, \dots, x_n$ is called **path** where $x_i x_{i+1} \in E$, $i = 0, 1, \dots, n-1$;

(ii) : **strength** of path $P : x_0, x_1, \dots, x_n$ is $\bigwedge_{i=0, \dots, n-1} \mu(x_i x_{i+1})$;

(iii) : **connectedness** amid vertices x_0 and x_n is

$$\mu^\infty(x, y) = \bigwedge_{P: x_0, x_1, \dots, x_n} \bigwedge_{i=0, \dots, n-1} \mu(x_i x_{i+1});$$

(iv) : a sequence of vertices $P : x_0, x_1, \dots, x_n$ is called **cycle** where $x_i x_{i+1} \in E$, $i = 0, 1, \dots, n-1$ and there are two edges xy and uv such that $\mu(xy) = \mu(uv) = \bigwedge_{i=0, 1, \dots, n-1} \mu(v_i v_{i+1})$;

(v) : it's **t-partite** where V is partitioned to t parts, V_1, V_2, \dots, V_t and the edge xy implies $x \in V_i$ and $y \in V_j$ where $i \neq j$. If it's complete, then it's denoted by $K_{\sigma_1, \sigma_2, \dots, \sigma_t}$ where σ_i is σ on V_i instead V which mean $x \notin V_i$ induces $\sigma_i(x) = 0$;

(vi) : t-partite is **complete bipartite** if $t = 2$, and it's denoted by K_{σ_1, σ_2} ;

(vii) : complete bipartite is **star** if $|V_1| = 1$, and it's denoted by S_{1, σ_2} ;

(viii) : a vertex in V is **center** if the vertex joins to all vertices of a cycle. Then it's **wheel** and it's denoted by W_{1, σ_2} ;

(ix) : it's **complete** where $\forall uv \in E$, $\mu(uv) = \sigma(u) \wedge \sigma(v)$;

(x) : it's **strong** where $\forall uv \in E$, $\mu(uv) = \sigma(u) \wedge \sigma(v)$.

The notions of neighbor and neighborhood are about some vertices which have one edge with a fixed vertex. These notions present vertices which are close to a fixed vertex as possible. Based on strong edge, it's possible to define different neighborhood as follows.

Definition 2.4. (Neighborhood).

Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Suppose $x \in V$. Then

$$N(x) = \{y \in V \mid xy \in E\}.$$

Definition 2.5. (Co-Neighborhood).

Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph and $t \in \mathbb{N}$. Suppose $x_i \in V$, $i = 1, 2, \dots, t$. Then

(i)

$$N_t(x_1, x_2, \dots, x_t) = \{y \in V \mid x_i y \in E, i = 1, 2, \dots, t\}.$$

If $\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V$, then **neutrosophic t-neighborhood** is called **neutrosophic quasi-vertex set** and t is called **neutrosophic quasi-order**. Generally, when t isn't fixed, it's called **neutrosophic neighborhood**. **Neutrosophic number** is

$$\sum_{\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V} \sum_{j=1}^3 \sigma_j(x_i).$$

(ii)

$$N_t(x_1, x_2, \dots, x_t) = \{e \in E \mid e = x_i y \in E, i = 1, 2, \dots, t\}.$$

If $\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E$, then **neutrosophic co-t-neighborhood** is called **neutrosophic quasi-edge set** and t is called **neutrosophic quasi-size**. Generally, when t isn't fixed, it's called **neutrosophic co-neighborhood**. **Co-neutrosophic number** is

$$\sum_{\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E} \sum_{j=1}^3 \sigma_j(x_i).$$

3 Neutrosophic Quasi-Order

Proposition 3.1. Let $NTG : (V, E, \sigma, \mu)$ be an odd path. Then $\{x_2, x_4, \dots, x_{n-1}\}$ is related to quasi-order $\lfloor \frac{n}{2} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an odd path. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n$ where n and 1 has same parity. There are two sets. $\{x_1, x_3, \dots, x_n\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor + 1$ but $\{x_2, x_4, \dots, x_{n-1}\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ which is minimum number amid these two sets. So $\{x_2, x_4, \dots, x_{n-1}\}$ is related to quasi-order $\lfloor \frac{n}{2} \rfloor$. $\{x_2, x_4, \dots, x_{n-1}\}$ is quasi-vertex set which is optimal. \square

Proposition 3.2. Let $NTG : (V, E, \sigma, \mu)$ be an even path. Then $\{x_2, x_4, \dots, x_n\}$ is related to quasi-order $\frac{n}{2}$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an even path. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n$ where n and 1 has different parity. There are two sets. $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ and $\{x_2, x_4, \dots, x_n\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ which is minimum number like first set. So $\{x_2, x_4, \dots, x_n\}$ and $\{x_1, x_3, \dots, x_{n-1}\}$ are related to quasi-order $\lfloor \frac{n}{2} \rfloor$. $\{x_2, x_4, \dots, x_n\}$ and $\{x_1, x_3, \dots, x_{n-1}\}$ are quasi-vertex sets which are optimal. \square

Proposition 3.3. Let $NTG : (V, E, \sigma, \mu)$ be an odd cycle. Then $\{x_2, x_4, \dots, x_{n-1}\}$ is related to quasi-order $\lfloor \frac{n}{2} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an odd cycle. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n, x_1$ where n and 1 has same parity. There are two sets. $\{x_1, x_3, \dots, x_n\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor + 1$ but $\{x_2, x_4, \dots, x_{n-1}\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ which is minimum number amid these two sets. So $\{x_2, x_4, \dots, x_{n-1}\}$ is related to quasi-order $\lfloor \frac{n}{2} \rfloor$. $\{x_2, x_4, \dots, x_{n-1}\}$ is quasi-vertex set which is optimal. \square

Proposition 3.4. Let $NTG : (V, E, \sigma, \mu)$ be an even cycle. Then $\{x_2, x_4, \dots, x_n\}$ and $\{x_1, x_3, \dots, x_{n-1}\}$ are related to quasi-order $\frac{n}{2}$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an even cycle. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n, x_1$ where n and 1 has different parity. There are two sets. $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ and $\{x_2, x_4, \dots, x_n\}$ has quasi-order $\lfloor \frac{n}{2} \rfloor$ which is minimum number like first set. So $\{x_2, x_4, \dots, x_n\}$ and $\{x_1, x_3, \dots, x_{n-1}\}$ are related to quasi-order $\lfloor \frac{n}{2} \rfloor$. $\{x_2, x_4, \dots, x_n\}$ and $\{x_1, x_3, \dots, x_{n-1}\}$ are quasi-vertex sets which are optimal. \square

Proposition 3.5. Let $NTG : (V, E, \sigma, \mu)$ be complete. Then $\{x\}$ is related to quasi-order 1.

Proof. Let $NTG : (V, E, \sigma, \mu)$ be complete. Suppose x is a given vertex. Thus

$$N[x] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V.$$

It implies the set $\{x\}$ which is 1-neighborhood, is quasi-vertex set. Then $\{x\}$ is related to quasi-order 1. \square

Proposition 3.6. Let $NTG : (V, E, \sigma, \mu)$ be star. Then $\{x\}$ is related to quasi-order 1.

Proof. Let $NTG : (V, E, \sigma, \mu)$ be star. Suppose c is the center. Thus

$$N[c] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V.$$

It implies the set $\{c\}$ which is 1-neighborhood, is quasi-vertex set. Then $\{c\}$ is related to quasi-order 1. \square

4 Neutrosophic Quasi-Size

Proposition 4.1. Let $NTG : (V, E, \sigma, \mu)$ be an odd path. Then either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ are related to quasi-size $\lfloor \frac{n}{3} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an odd path. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n$ where n and 1 has same parity. There is two sets. Either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ which are minimum number amid these all sets. So either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$. Either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ are quasi-vertex set which are optimal. \square

Proposition 4.2. Let $NTG : (V, E, \sigma, \mu)$ be an even path. Then $\{x_2, x_4, \dots, x_n\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an even path. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n$ where n and 1 has different parity. There are two sets. Either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$. $\{x_1, x_3, \dots, x_n\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ and $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ which is minimum number like first set. So either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$. Either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ is quasi-vertex set which is optimal. \square

Proposition 4.3. Let $NTG : (V, E, \sigma, \mu)$ be an odd cycle. Then $\{x_1, x_4, \dots, x_{n-3}\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an odd cycle. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n, x_1$ where n and 1 has same parity. There are two sets. $\{x_1, x_3, \dots, x_n\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ and $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ which are minimum numbers amid all sets. So either $\{x_1, x_3, \dots, x_{n-1}\}$ or $\{x_1, x_3, \dots, x_n\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$. Either $\{x_1, x_3, \dots, x_{n-1}\}$ or $\{x_1, x_3, \dots, x_n\}$ is quasi-edge set which is optimal. \square

Proposition 4.4. Let $NTG : (V, E, \sigma, \mu)$ be an even cycle. Then $\{x_1, x_4, \dots, x_{n-3}\}$ is related to quasi-size $\lfloor \frac{n}{3} \rfloor$.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ be an even cycle. Thus $NTG : (V, E, \sigma, \mu)$ is $P : x_1, x_2, \dots, x_n, x_1$ where n and 1 has different parity. There are two sets. $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ and $\{x_1, x_3, \dots, x_{n-1}\}$ has quasi-size $\lfloor \frac{n}{3} \rfloor$ which is minimum number like first set. So either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ are related to quasi-size $\lfloor \frac{n}{3} \rfloor$ which this number is optimal. Either $\{x_1, x_3, \dots, x_n\}$ or $\{x_1, x_3, \dots, x_{n-1}\}$ is quasi-edge set which are optimal. \square

Proposition 4.5. Let $NTG : (V, E, \sigma, \mu)$ be complete. Then $\{x_1, x_2, \dots, x_{n-1}\}$ is related to quasi-size $n - 1$.

Proof. Let $NTG : (V, E, \sigma, \mu)$ be complete. Suppose x is a given vertex. Thus

$$N[x_1, x_2, \dots, x_{n-1}] = \min_{x_1, x_2, \dots, x_{n-1} \in V} N_t[x_1, x_2, \dots, x_{n-1}] = E.$$

It implies the set $\{x_1, x_2, \dots, x_{n-1}\}$ which is co- $(n - 1)$ -neighborhood, is only quasi-vertex set as optimal set. Then $\{x_1, x_2, \dots, x_{n-1}\}$ is related to quasi-size $n - 1$. \square

Proposition 4.6. Let $NTG : (V, E, \sigma, \mu)$ be star. Then $\{x\}$ is related to quasi-size 1.

Proof. Let $NTG : (V, E, \sigma, \mu)$ be star. Suppose c is the center. Thus

$$N[c] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E.$$

It implies the set $\{c\}$ which is 1-neighborhood, is quasi-vertex set. Then $\{c\}$ is related to quasi-order 1. \square

5 Setting of Neutrosophic Quasi-Number and Quasi-Co-Number

Definition 5.1. (Quasi-Number & Quasi-Co-Number).

Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then

(i) **quasi-number** for x_0 is

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1});$$

(ii) **quasi-co-number** for x_0 is

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\}.$$

Proposition 5.2. Let $NTG : (V, E, \sigma, \mu)$ be a complete-neutrosophic graph. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph and $x_0 x'_0 \in E$. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

Since

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) = \mu(x_0 x'_0).$$

and

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \leq \mu(x_0 x'_0)$$

hold if $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph. Also,

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \geq \mu(x_0 x'_0)$$

holds since $x_0 x'_0$ is a path from x_0 to x'_0 . □ 135

Proposition 5.3. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0 z \in E$. Then □ 136

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0 z \in E$. An arbitrary path P has consecutive vertices x_0, z as their ends from a given vertex to leaf z . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

It implies for every given vertex $x_t \in V$,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z). □ 137$$

Proposition 5.4. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0 z \in E$ is weakest edge. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0z \in E$ is a weakest edge. An arbitrary path P has consecutive vertices x_0, z as their ends from a given vertex to leaf x_0 . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

By $x_0z \in E$ is a weakest edge,

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

It implies for every given vertex $x_t \in V$,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

□ 138

Proposition 5.5. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 has a weakest edge $x_0z \in E$. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 has a weakest edge $x_0z \in E$. For every vertex, there are only two ways to form a path. By $x_0z \in E$ is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

$P : x_0, z$ is a path from x_0 to z . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

□ 139

Proposition 5.6. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 is incident to one edge $x_0z \in E$ which isn't a weakest edge. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 has an edge $x_0z \in E$ which isn't a weakest edge. For every vertex, there are only two ways to form a path. By $x_0z \in E$ isn't a weakest edge and $P : x_0, z$ is a path from x_0 to z . It implies

$$\bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

□ 140

Proposition 5.7. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 is incident to two edges $x_0z, x_0z' \in E$ which aren't weakest edges. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if $\mu(x_0 z) < \mu(x_0 z')$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if $\mu(x_0 z') < \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if $\mu(x_0 z') = \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 is incident to two edges $x_0z, x_0z' \in E$ which aren't weakest edges. For every vertex, there are only two ways to form a path which are with edges either x_0z or x_0z' . $P : x_0, z$ and $P' : x_0, z'$ are paths from x_0 to z and z' . It implies

$$\bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if $\mu(x_0 z) < \mu(x_0 z')$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if $\mu(x_0 z') < \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if $\mu(x_0 z') = \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

□ 141

Proposition 5.8. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph. Suppose x_0 is center. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 is center. All paths from x_0 has a form $x_0 x_i$. Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

□ 142

Proposition 5.9. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph. Suppose x_0 isn't center and c is a center. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 isn't center and c is a center. There's only one path from x_0 has a form $x_0 c$. Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

□ 143

Proposition 5.10. Let $NTG : (V, E, \sigma, \mu)$ be a star-strong-neutrosophic graph. Suppose x_0 is center such that $\bigwedge_{i=0}^{\mathcal{O}-1} = \sigma(x_0)$. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 is center such that $\bigwedge_{i=0}^{\mathcal{O}-1} \mu(x_0 x_i) = \sigma(x_0)$. All paths from x_0 has a form $x_0 x_i$. Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i) = \sigma(x_0).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

□ 144

Proposition 5.11. Let $NTG : (V, E, \sigma, \mu)$ be a complete-neutrosophic graph. Then

145

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph and $x_0 x'_0 \in E$. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

Since

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) = \mu(x_0 x'_0).$$

and

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \leq \mu(x_0 x'_0)$$

hold if $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph. Also,

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \geq \mu(x_0 x'_0)$$

holds since $x_0 x'_0$ is a path from x_0 to x'_0 . So

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

□ 146

Proposition 5.12. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0z \in E$. Then

$$\mathcal{C}_c(x_0) = \{x_zx_{z'} \mid \mu(x_zx_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \{x_zx_{z'} \mid \mu(x_zx_{z'}) \leq \mu(x_0z)\}.$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0z \in E$. An arbitrary path P has consecutive vertices x_0, z as their ends from a given vertex to leaf z . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0z).$$

It implies for every given vertex $x_t \in V$,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0z).$$

Thus

$$\mathcal{C}_c(x_0) = \{x_zx_{z'} \mid \mu(x_zx_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \{x_zx_{z'} \mid \mu(x_zx_{z'}) \leq \mu(x_0z)\}.$$

□ 147

Proposition 5.13. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0z \in E$ is a weakest edge. Then

$$\mathcal{C}_c(x_0) = \{x_zx_{z'} \mid \mu(x_zx_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \{x_zx_{z'} \mid \mu(x_zx_{z'}) = \mu(x_0z)\}.$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Suppose x_0 is a leaf and $x_0z \in E$ is a weakest edge. An arbitrary path P has consecutive vertices x_0, z as their ends from a given vertex to leaf x_0 . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0z).$$

By $x_0z \in E$ is a weakest edge,

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0z).$$

It implies for every given vertex $x_t \in V$,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z)\}. \end{aligned}$$

□ 148

Proposition 5.14. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 has two weakest edges $x_0 z, x_0 z' \in E$. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z) \vee \mu(x_0 z') = \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\} = \{x_0 z, x_0 z'\}. \end{aligned}$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 has a weakest edge $x_0 z \in E$. For every vertex, there are only two ways to form a path. By $x_0 z \in E$ is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

$P : x_0, z$ is a path from x_0 to z . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

By $x_0 z' \in E$ is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

$P : x_0, z'$ is a path from x_0 to z' . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

By for every vertex, there are only two ways to form a path,

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z) \vee \mu(x_0 z') = \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\} = \{x_0 z, x_0 z'\}. \end{aligned}$$

□ 149

Proposition 5.15. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 is incident to one edge $x_0 z \in E$ which isn't a weakest edge. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \geq \mu(x_0 z), x_z x_{z'} \in E\}. \end{aligned}$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 has an edge $x_0 z \in E$ which isn't a weakest edge. For every vertex, there are only two ways to form a path. By $x_0 z \in E$ isn't a weakest edge and $P : x_0, z$ is a path from x_0 to z . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \geq \mu(x_0 z), x_z x_{z'} \in E\}. \end{aligned}$$

□ 150

Proposition 5.16. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Suppose x_0 is incident to two edges $x_0 z, x_0 z' \in E$ which aren't weakest edges. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) \vee \mu(x_0 z'), x_z x_{z'} \in E\}. \end{aligned}$$

Furthermore, if $\mu(x_0 z) < \mu(x_0 z')$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z'), x_z x_{z'} \in E\};$$

if $\mu(x_0 z') < \mu(x_0 z)$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z), x_z x_{z'} \in E\};$$

if $\mu(x_0 z') = \mu(x_0 z)$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\}.$$

Proof. Assume $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Suppose x_0 is incident to two edges $x_0z, x_0z' \in E$ which aren't weakest edges. For every vertex, there are only two ways to form a path which are with edges either x_0z or x_0z' . $P : x_0, z$ and $P' : x_0, z'$ are paths from x_0 to z and z' . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if $\mu(x_0 z) < \mu(x_0 z')$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if $\mu(x_0 z') < \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if $\mu(x_0 z') = \mu(x_0 z)$, then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) \vee \mu(x_0 z'), x_z x_{z'} \in E\}. \end{aligned}$$

Furthermore, if $\mu(x_0 z) < \mu(x_0 z')$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z'), x_z x_{z'} \in E\};$$

if $\mu(x_0 z') < \mu(x_0 z)$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z), x_z x_{z'} \in E\};$$

if $\mu(x_0 z') = \mu(x_0 z)$, then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\}.$$

□ 151

Proposition 5.17. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph. Suppose x_0 is center. Then

152

153

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 is center. All paths from x_0 has a form x_0x_i . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

□ 154

Proposition 5.18. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph. Suppose x_0 isn't center and c is a center. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(cx_0)\} = \{cx_0\}. \end{aligned}$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 isn't center and c is a center. There's only one path from x_0 has a form x_0c . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(cx_0)\} = \{cx_0\}. \end{aligned}$$

□ 155

Proposition 5.19. Let $NTG : (V, E, \sigma, \mu)$ be a star-strong-neutrosophic graph. Suppose x_0 is center such that $\bigwedge_{i=0}^{\mathcal{O}-1} \mu(x_0 x_i) = \sigma(x_0)$. And x isn't center. Then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

And

$$\mathcal{C}_c(x) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

Proof. Consider $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Suppose x_0 is center such that $\bigwedge_{i=0}^{\mathcal{O}-1} \mu(x_0 x_i) = \sigma(x_0)$. All paths from x_0 has a form $x_0 x_i$. Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i) = \sigma(x_0).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

Thus

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

And

$$\mathcal{C}_c(x) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

□ 156

References

1. N.H. Abughazalah, and M. Khan, “Selection of Optimum Nonlinear Confusion Component of Information Confidentiality Mechanism Using Grey Theory Based Decision-Making Technique.” Preprints 2021, 2021120310 (doi: 10.20944/preprints202112.0310.v1). 157
2. M. Akram, and G. Shahzadi, “Operations on Single-Valued Neutrosophic Graphs”, Journal of uncertain systems 11 (1) (2017) 1-26. 158
3. K. Atanassov, “Intuitionistic fuzzy sets”, Fuzzy Sets Syst. 20 (1986) 87-96. 159

160

161

162

163

164

4. S. Boumova, P. Boyvalenkov, and M. Stoyanova, “*Bounds for the Minimum Distance and Covering Radius of Orthogonal Arrays via Their Distance Distributions.*” Preprints 2021, 2021120293 (doi: 10.20944/preprints202112.0293.v1).

165
166
167
168

5. S. Broumi, M. Talea, A. Bakali and F. Smarandache, “*Single-valued neutrosophic graphs*”, Journal of New Theory 10 (2016) 86-101.

169
170

6. N. Shah, and A. Hussain, “*Neutrosophic soft graphs*”, Neutrosophic Set and Systems 11 (2016) 31-44.

171
172

7. Henry Garrett, (2022). “*Beyond Neutrosophic Graphs*”, Ohio: E-publishing: Educational Publisher 1091 West 1st Ave Grandview Heights, Ohio 43212 United States. ISBN: 978-1-59973-725-6 (http://fs.unm.edu/BeyondNeutrosophicGraphs.pdf).

173
174
175
176

8. Henry Garrett, “*Co-degree and Degree of classes of Neutrosophic Hypergraphs*”, Preprints 2022, 2022010027 (doi: 10.20944/preprints202201.0027.v1).

177
178

9. D. Jean, and S. Fault-Tolerant Seo, “*Detection Systems on the King’s Grid.*” Preprints 2022, 2022010249 (doi: 10.20944/preprints202201.0249.v1).

179
180

10. H. Oliveira, “*Synthesis of Strategic Games With Multiple Pre-set Nash Equilibria - An Artificial Inference Approach Using Fuzzy ASA.*” Preprints 2019, 2019080128 (doi: 10.20944/preprints201908.0128.v2).

181
182
183

11. R.E. Perez, F.N. Gonzalez, M.C. Molina, and J.M. Moreno “*Methodological Analysis of Damage Estimation in Hydraulic Infrastructures.*” Preprints 2022, 2022010376 (doi: 10.20944/preprints202201.0376.v1).

184
185
186

12. A. Shannon and K.T. Atanassov, “*A first step to a theory of the intuitionistic fuzzy graphs*”, Proceeding of FUBEST (Lakov, D., Ed.) Sofia (1994) 59-61.

187
188

13. F. Smarandache, “*A Unifying field in logics neutrosophy: Neutrosophic probability, set and logic, Rehoboth:* ” American Research Press (1998).

189
190

14. F. Tang, X. Zhu, J. Hu, J. Tie, J. Zhou, and Y. Fu, “*Generative Adversarial Unsupervised Image Restoration in Hybrid Degradation Scenes.*” Preprints 2022, 2022020159 (doi: 10.20944/preprints202202.0159.v1).

191
192
193

15. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, “*Single-valued neutrosophic sets*”, Multispace and Multistructure 4 (2010) 410-413.

194
195

16. L. A. Zadeh, “*Fuzzy sets*”, Information and Control 8 (1965) 338-354.

196

17. S. Zuev, Z. Hussain, and P. Kabalyants, “*Nanoparticle Based Water Treatment Model in Squeezing Channel under Magnetic Field.*” Preprints 2021, 2021120295 (doi: 10.20944/preprints202112.0295.v1).

197
198
199