

Article

Not peer-reviewed version

Diversity of Weed Flora Survey as a Starting Point for the Development of a Weed Management Strategy for Medicinal Crops

[Dragana Božić](#) , [Ana Dragumilo](#) , [Tatjana Marković](#) , [Urban Šilc](#) , [Svetlana Ačić](#) , [Teodora Tojić](#) , [Miloš Rajković](#) , [Sava Vrbničanić](#) *

Posted Date: 25 June 2025

doi: 10.20944/preprints202506.2111.v1

Keywords: medicinal crops; weed flora; diversity; abundance; frequency; control strategy



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Diversity of Weed Flora Survey as a Starting Point for the Development of a Weed Management Strategy for Medicinal Crops

Dragana Božić¹, Ana Dragumilo², Tatjana Marković², Urban Šilc³, Svetlana Aćić¹, Teodora Tojić¹, Miloš Rajković² and Sava Vrbničanin^{1*}

¹ University of Belgrade, Faculty of Agriculture, Belgrade-Zemun, Serbia

² Institute for Medicinal Plant Research "Dr Josif Pančić", Belgrade, Serbia

³ Jovan Hadži Institute of Biology, Research Centre of the Slovenian Academy of Sciences and Arts, Ljubljana, Slovenia

* Correspondence: sava@agrif.bg.ac.rs

Abstract

Similar to conventional field crops, weeds often pose significant problems in the cultivation of medicinal plants. To date, no comprehensive documentation exists regarding weed infestation levels in these crops in Serbia. Objective of this study was to provide a valuable foundation for developing effective, site-specific weed management strategies in medicinal crop production. Weeds in five medicinal crops (lemon balm, fennel, peppermint, ribwort plantain, German chamomile), were surveyed based on the agro-phytosociological method between 2019 and 2024, and across 59 plots. A total of 109 weed species were recorded, belonging to 29 families and 88 genera. Among them, 75 were annuals and 34 perennials, including 93 broadleaved species, 10 grasses, and one parasitic species. The most important ecological factors influencing the composition of weed vegetation in investigated medicinal crops were temperature and light for fennel and peppermint plots, soil reaction for lemon balm and ribwort plantain plots and nutrient content for German chamomile plots. Weed control strategies should consider such information, targeting the control of the most frequent, abundant and dominant species existing in a crops or locality.

Keywords: medicinal crops; weed flora; diversity; abundance; frequency; control strategy

1. Introduction

Medicinal crops are most often grown on small areas around the world and as such are classified as minor crops. The areas under cultivated medicinal, aromatic and spicy plants in Serbia vary from 2,000-3,000 ha per year (<0.15% arable lands), with 20-25 species being cultivated the most: chamomile, peppermint, fennel, lemon balm, ribwort plantain, thyme, marshmallow, parsley, marigold, dill, basil, oregano, etc. [1]. Medicinal crop production has an important and beneficial role in the economic, social, cultural and ecological aspects of human communities in the worldwide. Those so-called minor crops in Serbia, which are typically cultivated on small farms averaging less than 0.3 ha per plot, can help smallholder farmers directly improve their livelihoods by generating income through trade and by contributing to health-related industries [2,3]. Also, the production of medicinal crops is a benefit for everyone, whether those herbs are used as food, in hygiene, personal care, in folk medicine, etc. [4].

As with conventional field crops, various constraints pose serious problem for the cultivation of medicinal crops, among which weeds are particularly significant. According to the definition adopted by the European Weed Research Society "Weeds are any plant or vegetation, excluding fungi, that interferes with the objectives or requirements of people". Similarly, the Weed Science Society of America defines a weed as "a plant growing where it is not desired". Unlike plant

pathogens and pests, weed species represent a permanent and major challenge in the cultivation of medicinal crops. It is generally estimated that weeds cause yield losses of about 10% in developed countries and up to 25% in less developed countries in conventional crop production [5]. Further, Kashem et al. [6] estimate that unmanaged weed growth during the early stages of crop establishment can reduce yields by 25% to 80%. Globally, it is estimated that approximately 1,800 weed species are responsible for a 31.5% decline in crop production, causing annual economic losses of about 32 billion USD [7]. However, there are still limited studies on weed diversity and their harmfulness in medicinal crop production systems [8-10]. The abundance and composition of weed species in medicinal crops, in addition to the type of crop and cultivation practices, largely depend on the soil type and quality, as well as on climatic and meteorological conditions in a given season [11-13].

The harmfulness of weeds in different crops is reflected in their ability to successfully compete with cultivated plants for natural resources such as light, nutrients, water, and space. Weeds negatively impact both the quantity and quality of crop yield, increase crop production costs due to the necessity of weed control measures, and can serve as hosts to numerous plant pests and diseases [14, 15]. Globally, approximately 1,570 weed species have been identified [16], with more recent estimates suggesting around 1,800 species [7]. In Serbia, about 300 weed species have been confirmed in crop fields, while the total number of weed species, including those in non-crop areas, is estimated at around 1,000 species [17]. However, there are still limited studies addressing weed diversity and their harmfulness specifically in medicinal plant production systems [9, 10]. Interestingly, weed-induced stress conditions may even enhance the synthesis of biologically active secondary metabolites in medicinal plants [18]. In addition, some weed species such as *Adonis aestivalis*, *Aristolochia clematidis*, *Cirsium arvense*, *Carduus* spp., *Conium maculatum*, *Datura stramonium*, *Lolium temulentum*, *Solanum nigrum*, can cause significant problems during mechanized harvesting, potentially contaminating the harvested agricultural or medicinal plants [19, 20].

The reasons for weed control in medicinal crops are numerous: increasing crop productivity, enhancing product quality, reducing production costs, limiting the spread of other pests, improving animal health, supporting human activities, reducing impacts on transportation, and minimizing health risks. However, the primary reason for weed management in medicinal crops remains the prevention of yield and quality losses. A fundamental prerequisite for timely and effective weed control in medicinal crops is precise knowledge of weed species composition and their abundance in the fields. Detailed information on weed density, frequency, and community structure is essential for predicting potential yield losses and for defining cost-effective weed management thresholds [21].

Therefore, the aim of this research was to conduct a multi-year floristic survey across different seasons and years in order to provide reliable data on weed biodiversity and abundance, as a solid starting point for the development of an effective weed management strategy in medicinal crop plantations in Serbia. The data obtained can now be used to identify major weed problems that require targeted research or the implementation of improved weed management practices.

2. Materials and Methods

Study area. The analysis of weed flora diversity of selected medicinal crops, lemon balm (*Melissa officinalis* L.), fennel (*Foeniculum vulgare* Mill.), peppermint (*Mentha piperita* L.), ribwort plantain (*Plantago lanceolata* L.), and German chamomile (*Chamomilla recutita* L.), was conducted between 2019 and 2024 on commercial farms of Institute for Medicinal Plant Research "Dr Josif Pančić" located in Pančevo, South Banat, Republic of Serbia, Southeast Europe (44°52'20.0"N, 20°42'04.7"E). The research was based on the agro-phytosociological method, according to Braun-Blanquet [22].

Basic data on crop characteristics (i.e. life cycle, crop type, sowing time, seeding rate, and harvest period) are provided in Table 1. In these fields, medicinal crops are traditionally grown in crop rotation with small grain cereals.

The soil type was chernozem, containing 2.3% humus and 0.19% total nitrogen while the available phosphorus (P_2O_5) and potassium (K_2O), extracted using ammonium lactate method, were 36 mg kg^{-1} and 362 mg kg^{-1} , respectively. Fertilizer was applied three times per growing season,

adapted to soil nutrient status. The first fertilization was carried out following the deep autumn plowing using a solid NPK mineral fertilizer (NPK 15–15–15) at rate 0.6 t ha⁻¹. The subsequent two fertilizations were performed with calcium ammonium nitrate (KAN, 27% N) at rate of 0.2 t ha⁻¹. The first KAN application was performed in early spring, after the development of two to four true leaves in all studied species. The timing of the second KAN application varied depending on the crop. In lemon balm and peppermint, the second application was carried out immediately after the first harvest to support regrowth and ensure sufficient nutrient availability for the subsequent growth cycle. In fennel and ribwort plantain, typically harvested once, the second application was made during peak vegetative growth, about 4–6 weeks after the initial fertilization and prior to flowering. In German chamomile, the second KAN application was timed at the beginning of the flowering stage to promote optimal flower development and essential oil yield.

Table 1. Overview of key cultivation practices for selected medicinal crops.

Crop	Lemon balm	Fennel	Peppermint	Ribwort plantain	German chamomile
Life cycle	Perennial (5-6 years)	Perennial (5-10 years)	Perennial (1 -3 years)	Perennial (1 -3 years)	Annual
Crop type	Row crop	Row crop	Row crop	Row crop	Dense stand crop
Sowing/ planting	Transplant seedlings: during autumn or early spring; 60-70 x 25-30 cm	Direct sowing: end of March to beginning of April; 60-70 x 20- 30 cm	Planting stolon: mid-September to end of November or early spring; 60-70 x 70 cm	Direct sowing: end of March - beginning of April; 50-60 x 30 cm	Direct sowing: during September
Seeding rate (plants ha⁻¹)	47.600-66.600	120 -150.000	30.000	55.500- 66.600	200.000
Harvest time	Twice per year: first harvest in July; second harvest in beginning of October	Second half of September	Twice per year: first harvest in July; second harvest in September/October	June (first year) - May (other years)	May

Meteorological data, including average monthly temperatures and precipitation amounts for the period 2019–2024, were obtained from the local Meteorological Stations of the Republic of Serbia for

the Pančevo locality (Table 2). According to these data, weather conditions varied notably between seasons and years. Generally, July or August were the warmest months with average temperatures ranging from 23.7 to 26.9 °C, while January was typically the coldest month, with averages between 0.8 and 5.7 °C. Precipitation patterns showed no consistent seasonal regularity throughout the study period; however, the highest precipitation occurred in June 2020 (158.5 mm) or December 2021 (157.8 mm), depending on the year.

Table 2. Meteorological conditions in Pančevo during the period 2019–2024.

Month	2019		2020		2021		2022		2023		2024	
	T (°C)	P (mm)										
January	0.8	81.8	2.0	22.1	4.3	68.6	2.4	45.7	5.7	79.3	3.6	50.6
February	5.6	33.7	7.6	55.9	6.5	34.4	6.9	22.2	4.9	62.9	10.6	9.6
March	11.4	11.5	9.1	48.0	7.2	49.3	7.1	10.5	10.1	37.6	11.5	51.1
April	14.2	76.8	14.3	8.9	10.6	50.7	12.3	80.1	11.3	79.0	15.6	38.8
May	15.6	142.3	16.6	70.9	17.4	93.4	20.3	32.2	17.5	92.8	18.2	132.9
June	24.2	138.7	20.9	158.5	24.3	34.2	24.8	43.3	21.7	75.6	24.1	77.2
July	24.3	43.0	23.7	37.6	26.6	63.1	25.9	63.9	26.0	46.8	26.6	114.4
August	26.2	39.7	24.9	89.6	24.0	38.2	25.0	89.7	24.7	87.7	26.9	0.8
September	20.2	26.1	21.1	22.1	19.4	9.4	18.0	98.0	22.2	71.2	19.8	87.9
October	16.1	13.3	14.3	93.3	11.7	73.4	15.5	13.1	17.6	13.0	14.3	38.2
November	12.4	54.3	7.2	12.6	8.9	122.8	9.3	64.9	9.5	107.4	6.0	70.9
December	6.0	55.3	5.6	34.8	4.1	157.8	7.0	76.1	7.0	33.3	3.7	105.7

T- average temperature (°C), P- precipitation amounts (mm).

Survey of weed diversity and abundance. The qualitative and quantitative evaluation of weed flora was conducted over six years each spring, following crop emergence and prior to any weed management measures, on chamomile (18 plots), ribwort plantain (11), peppermint (10), fennel (10), and lemon balm (10). Each plot measured approximately 0.50 ha. Plant species abundance and coverage were assessed using the scale by Westhoff and van der Maarel [23], while weed frequency was calculated following Braun-Blanquet [22] principles, and expressed on a scale from I to V, where I indicates presence in 1–20% of plots, II in 21–40%, III in 41–60%, IV in 61–80%, and V in 81–100%. The vouchers of weed species were collected and are kept in the Laboratory of Botany and Laboratory of Weed Science at the Faculty of Agriculture, University of Belgrade. Species nomenclature was standardized according to Euro+Med PlantBase [24].

Data analysis. Each plant species was classified by its life form according to Raunkiaer [25] as annuals (therophytes, T), biennials (thero-hemicryptophytes, TH), and perennials, with the latter group including geophytes (G), hemicryptophytes (H), chamaephytes (Ch), and phanerophytes (P). Diagnostic species (Dg) for each group were identified using JUICE 7.1 software, with the phi coefficient as a measure of fidelity [26, 27]. Species with phi coefficient values greater than 0.15 were considered diagnostic. Species with a cover exceeding 25% and present in at least 50% of the total plots were considered dominant (Dm) in each group, while species occurring in at least 50% of the total plots were classified as constant (C).

To assess the influence of environmental factors on the weed species composition across different communities, non-metric multidimensional scaling (NMDS) was applied using the JUICE 7.1 software, the R-project platform [28], and the vegan package [29]. Species cover was changed into

percentages and transformed using the Hellinger transformation [30]. For ecological gradient analysis, Ecological indicator values (EIVs) according to Pignatti [31] were used. Mean unweighted EIVs for light, continentality, temperature, nutrients, moisture, and soil reaction were used in NMDS as passively projected explanatory variables.

3. Results

3.1. Weed flora

We confirmed presence of 109 distinct weed species across 59 plots of five different medicinal crops, including 75 annuals and 34 perennials. The weed flora comprised 93 broadleaved species, 10 grasses, and 1 parasitic species. These species belonged to 29 families and 88 genera. Among the families, Asteraceae had the highest number of weed species (26), followed by Poaceae (10), Brassicaceae (7), Fabaceae (7), Polygonaceae (6), Apiaceae (4), Caryophyllaceae (4), Chenopodiaceae (4), Lamiaceae (4), Scrophulariaceae (4), Solonaceae (3), Boraginaceae (3), Geraniaceae (3), Plantaginaceae (3), Malvaceae (3), Amaranthaceae (2), Convolvulaceae (2), Ranunculaceae (2), Rosaceae (2), and ten families represented by a single species (Table 3).

Biological spectrum of life forms in the weed communities of medicinal crops was dominated by therophytes (T=64), followed by thero-hemicryptophytes (TH=11), hemicryptophytes (H=17), geophytes (G=14), while chamaephytes (Ch) and phanerophytes (P) were represented by 1 and 2 species, respectively (Figure 1).

All surveyed plots of medicinal crops contained perennial weed species, including *Elymus repens*, *Artemisia vulgaris*, *Cirsium arvense*, *Convolvulus arvensis*, *Lepidium draba*, *Rumex crispus*, *Sonchus arvensis*, *Sorghum halepense*, and *Taraxacum officinale*, among others. These perennials are particularly problematic due to their persistence and difficulty of control. Most of these species reproduce vegetatively through underground organs such as rhizomes, roots and root shoots, complicating soil cultivation practices.




Figure 1. The participation of the life forms of weed flora in medicinal crops.

In addition to perennials, a significant number of annual weeds were present with high abundance and frequency, including *Amaranthus retroflexus* (abundance 2–7; frequency I–IV), *Chenopodium album* (ab. 1–7; fr. II–IV), *Galium aparine* (ab. 2–5; fr. II–IV), *Lactuca serriola* (ab. 1–7; fr. II–IV), *Lamium amplexicaule* (ab. 2–8; fr. III–IV), *L. purpureum* (ab. 2–9; fr. III–IV), *Papaver rhoeas* (ab. 1–9; fr. I–V), *Stellaria media* (ab. 1–8; fr. III–V), *Veronica hederifolia* (ab. 2–9; fr. III–V), *V. persica* (ab. 2–9; fr. III–V), etc. (Table 3).

In addition to typical weeds, volunteer medicinal crops frequently emerged as weeds within the plantation fields. For example, *Foeniculum vulgare* was often found in ribwort plantain plots, *Cynara*

cardunculus appeared in fennel plots, and *Linum usitatissimum* was commonly present in fennel, peppermint and ribwort plantain plots. Moreover, overall weed flora of medicinal crops showed a high abundance of alien invasive species, notably *Ambrosia artemisiifolia* (ab. 1–9) and *Ailanthus altissima* (ab. 2). The presence of weed species known for their high toxicity in vegetative and generative organs was also confirmed, including members of the Solanaceae (*Datura stramonium*, *Solanum nigrum*, *S. dulcamara*) and Apiaceae family (*Bifora radians*, *Conium maculatum*) (Table 3).

Table 3. Survey of weed flora in studied medicinal crops.

Weed species	Life form	Medicinal crops									
		Lemon balm		Fennel		Peppermint		Ribwort plantain		German Chamomile	
		Ab.	Fr.	Ab.	Fr.	Ab.	Fr.	Ab.	Fr.	Ab.	Fr.
Amaranthaceae											
<i>Amaranthus blitoides</i> S.Wat.	T	2-5	II			2-5	III	2	I		
<i>Amaranthus retroflexus</i> L.	T	3	I	2	II	2-7	IV	2	I		
Apiaceae											
<i>Bifora radians</i> M.B.	T							2	I		
<i>Conium maculatum</i> L.	T			1-2	IV						
<i>Daucus carota</i> L.	TH	3	I								
<i>Foeniculum vulgare</i> Mill.	H							2	I		
Asteraceae											
<i>Achillea millefolium</i> L.	G	2	I								
<i>Ambrosia artemisiifolia</i> L.	T	1-7	II	3	I	2-5	III	5-9	II		
<i>Anthemis arvensis</i> L.	T							3	II	2	I
<i>Anthemis cotula</i> L.	T							2	II	3-5	II
<i>Arctium lappa</i> L.	TH			1	III					2	I
<i>Artemisia vulgaris</i> L.	H			2-3	V	1-2	IV	2-3	IV		
<i>Carduus acanthoides</i> L.	TH	1-2	III	2	IV			1-5	III	2	II
<i>Chamomilla recutita</i> (L.) Rausch.	T	2-5	III	7	I	2-3	IV	2-7	III		
<i>Cichorium intybus</i> L.	H			1	I	1	I				
<i>Cirsium arvense</i> (L.) Scop.	G	2-5	III	2-3	III	1-2	IV	3-8	IV	2-8	IV
<i>Crepis biennis</i> L.	TH	2	II								
<i>Cynara cardunculus</i> L.	H			2	II						
<i>Erigeron canadensis</i> L.	T	2-5	II	1	II			2	II		
<i>Helianthus annuus</i> L.	T			1	II			1			
<i>Lactuca serriola</i> L.	T	1-2	II	2	III	1	III	3-7	IV	2-3	III
<i>Lactuca viminea</i> (L.) J.P. & C.P.	T	2	I	2	III	1	III	2	II	3	I
<i>Picris hieracioides</i> L.	TH	2	III	3	II	2	I			2	I

<i>Senecio vernalis</i> W. et K.	T	2	II								
<i>Senecio vulgaris</i> L.	T	1-2	III			1-2	III	2-3	III	2-3	V
<i>Sonchus arvensis</i> L.	G	2	II	2	III	1	IV	2	IV	2	II
<i>Sonchus asper</i> (L.) Hill.	TH	2	I								
<i>Sonchus oleraceus</i> (L.) Gou.	T	2	I			2	I			2	I
<i>Erigeron annuus</i> (L.) Desf.	T	2	III	3	II	2	III	3	II	2	I
<i>Taraxacum officinale</i> Weber.	H	2-3	V	3	IV	2	II	3	V	2	III
<i>Tragopogon dubius</i> Scop.	H										
<i>Xanthium strumarium</i> L.	T			2	I			1	I		
Boraginaceae											
<i>Heliotropium europeum</i> L.	T					2	II				
<i>Lithospermum arvense</i> L.	TH									2-3	II
<i>Myosotis arvensis</i> (L.) Hill.	T									7-8	I
Brassicaceae											
<i>Brassica nigra</i> L.	T							1	II		
<i>Camelina microcarpa</i> Andr. ex DC	TH	3	II					3	II		
<i>Capsella bursa-pastoris</i> (L.) Medic.	T	2-8	III	3	II	2	I	2-7	II	2-5	V
<i>Lepidium draba</i> L.	G	2-3	IV			1-2	III	3-7	III	2-3	II
<i>Roripa sylvestris</i> (L.) Bess.	G	1	I								
<i>Sinapis arvensis</i> L.	T	2	I	2-5	III			1	I	2-7	V
<i>Thlaspi arvense</i> L.	T	2	I								
Caryophyllaceae											
<i>Arenaria serpyllifolia</i> L.	T							3	I		
<i>Cerastium arvense</i> L.	ZC	3	II			2	III			7	I
<i>Silene alba</i> (Mill.) Krause.	H	5	II	2	II						
<i>Stellaria media</i> (L.) Vill.	T	1-8	V			2	III	2-7	III	5-8	V
Chenopodiaceae											
<i>Atriplex patula</i> L.	T	2	I					2	I		
<i>Chenopodium album</i> L.	T	1-3	III	2	IV	2-7	IV	2-3	III	2-3	II
<i>Chenopodium hybridum</i> L.	T	3	III			1		2	I		
<i>Chenopodium polyspermum</i> L.	T					2	II				
Convolvulaceae											
<i>Calystegia sepium</i> (L.) R.Br.	G			1	III	2	III			2	I
<i>Convolvulus arvensis</i> L.	G	1-8	IV	2-7	V	2-3	III	2-5	IV	2-3	II
Cuscutaceae											

<i>Cuscuta campestris</i> Yunck.	T			3	I						
Euphorbiaceae											
<i>Euphorbia virgata</i> L.	G			1	I						
Fabaceae											
<i>Lathyrus tuberosus</i> L.	G	2	I	1	I	1-2	II	1-2	II		
<i>Medicago lupulina</i> L.	T	1	I								
<i>Medicago sativa</i> L.	H							2	II		
<i>Trifolium repens</i> L.	H	2	II								
<i>Vicia cracca</i> L.	G							2	II	3	I
<i>Vicia hirsuta</i> (L.) S.F.Gray.	T							1	II		
<i>Vicia sativa</i> L.	T	2	I			2	I	2	II		
Fumariaceae											
<i>Fumaria officinalis</i> L.	T			2	II	1-2	V			2	II
Geraniaceae											
<i>Erodium cicutarium</i> (L.) L'Herit.	T			2	III			3	IV	5	I
<i>Geranium dissectum</i> Jusl.	T	1	I							2	I
<i>Geranium molle</i> L.	TH							2	II		
Lamiaceae											
<i>Lamium amplexicaule</i> L.	T	2-7	III	2	III	3	IV	3-8	III	2-8	IV
<i>Lamium purpureum</i> L.	T	2-5	III	2-7	III	5	IV	2-9	III	2-8	IV
<i>Melissa officinalis</i> L.	G					2	I				
<i>Stachys annua</i> L.	T			2	I						
Linaceae											
<i>Linum usitatissimum</i> L.	T			2	I	1	I	1	I		
Malvaceae											
<i>Abutilon theophrasti</i> Medik.	T			2	I	2	I	2	I		
<i>Hibiscus trionum</i> L.	T	2	I	3-5	I	2-3	III	1	I		
<i>Malva silvestris</i> L.	TH							1	II		
Papaveraceae											
<i>Papaver rhoeas</i> L.	T	1	I	1-3	V	1-2	IV	3-9	V	2-8	V
Plantaginaceae											
<i>Plantago lanceolata</i> L.	H									2	I
<i>Plantago major</i> L.	H			3	II					3	I
<i>Plantago media</i> L.	H							2	II		
Poaceae											
<i>Elymus repens</i> (L.) Beauv.	G	2-7	IV	2-7	V	2	III	2-3	III	5	I
<i>Avena fatua</i> L.	T	2-3	III	3-5	III	2-3	IV	2-5	II	2-9	V

<i>Bromus tectorum</i> L.	T	2	I	3	III				2-8	III
<i>Cynodon dactylon</i> (L.) Pers.	G	3	I							
<i>Dactylis glomerata</i> L.	H							1-2	II	
<i>Echinochloa crus-galli</i> (L.) P.B.	T	2	I			2-7	II	2-3	III	
<i>Hordeum murinum</i> L.	T	2	II							
<i>Setaria glauca</i> (L.) P.B.	T					3	I	2	I	
<i>Setaria viridis</i> (L.) P.B.	T	2	II	2	I					
<i>Sorghum halepense</i> (L.) Pers.	G	1-3	IV	3	IV	3-7	III	3	II	8 I
Polygonaceae										
<i>Bilderdykia convolvulus</i> (L.) Dum.	T	2-3	III	2	I	2-7	IV	2-7	III	2-3 II
<i>Polygonum aviculare</i> L.	T	1-2	III	3	I	2-3	IV	1	III	2-3 II
<i>Polygonum lapathifolium</i> L.	T			2	I	2-8	IV			
<i>Polygonum persicaria</i> L.	T	2	II	3	II			2-3	III	
<i>Rumex crispus</i> L.	H	2-7	V	3-8	V	2-3	IV	2-5	V	2-7 V
<i>Rumex obtusifolius</i> L.	H			2-3	II	2	III			2 I
Portulacaceae										
<i>Portulaca oleracea</i> L.	T	5	I			3-5	II			
Ranunculaceae										
<i>Consolida orientalis</i> (Gaz.) Schr.	T	1	I	1-2	III	3	III	5	II	5 II
<i>Consolida regalis</i> S.F.Gray.	T	3	II			3	III	3	II	3-5 II
Resedaceae										
<i>Reseda lutea</i> L.	TH							1-2	III	
Rosaceae										
<i>Potentilla anserina</i> L.	H			2	II					
<i>Rubus caesius</i> L.	H					1-2	II	1	I	I
Rubiaceae										
<i>Galium aparine</i> L.	T			2-5	IV			2	II	5 III
Scrophulariaceae										
<i>Kickxia elatine</i> (L.) Dumort.	T			2	I					
<i>Veronica hederifolia</i> L.	T	3-8	III	2	III	8	III	2-5	III	3-9 V
<i>Veronica persica</i> Poir.	T	2-3	IV	2-3	V	2-8	IV	2-9	V	5-8 III
<i>Veronica polita</i> Fr.	T					2-5	III	2-5	I	5 I
Simaroubaceae										

<i>Ailanthus altissima</i> (Mill.) Swingle	P	2	I								
Solanaceae											
<i>Datura stramonium</i> L.	T					2-3	II	2	I		
<i>Solanum dulcamara</i> L.	nP	1	I								
<i>Solanum nigrum</i> L.	T	2-5	III	2-3	II	3-5	III	2	I		
Violaceae											
<i>Viola arvensis</i> Murr.	T					2-3	IV	3	II	2-5	IV

Ab.– abundance, Fr.– frequency.

The total number of weed species recorded across studied medicinal crops was 109, while the mean number of weed species per specific medicinal crop ranged between 13 and 15, except in *Chamomilla recutita* plots, where significantly fewer weed species were observed (Figure 2).




Figure 2. Mean number of weed species in different medicinal crops.

3.2. Weed Vegetation Characteristics of Medicinal Crops

Cluster analysis of weed vegetation grouped plots into five groups and diagnostic, constant and dominant species with their fidelity values are showed (Figure 3).

Group 1: *Melissa officinalis* (Lemon balm) plots

Number of plots: 10

Diagnostic species: *Crepis biennis* 18.5, *Senecio vernalis* 18.5, *Trifolium repens* 18.5

Constant species: *Convolvulus arvensis* 28, *Lamium amplexicaule* 21, *Stellaria media* 25

Group 2: *Foeniculum vulgare* (Fennel) plots

Number of plots: 10

Diagnostic species: *Arctium lappa* 16.2, *Artemisia vulgaris* 16.0, *Conium maculatum* 26.8, *Cynara cardunculus* 18.5

Constant species: *Convolvulus arvensis* 24

Group 3: *Mentha piperita* (Peppermint) plots

Number of plots: 10

Diagnostic species: *Amaranthus blitoides* 19.9, *Amaranthus retroflexus* (C, Dm) 35.2, *Chenopodium polyspermum* 18.5, *Fumaria officinalis* 18.7, *Heliotropium europeum* 21.4

Constant species: *Amaranthus retroflexus* (Dg, Dm) 24, *Chenopodium album* (Dm) 24, *Lamium purpureum* 25

Dominant species: *Amaranthus retroflexus* (Dg, C) 30, *Chenopodium album* (C) 30

Group 4: *Plantago lanceolata* (Ribwort plantain) plots

Number of plots: 11

Diagnostic species: *Brassica nigra* 17.1, *Geranium molle* 17.6, *Medicago sativa* 17.6

Constant species: *Ambrosia artemisiifolia* 24, *Chamomilla recutita* (Dm) 24, *Cirsium arvense* (Dm) 35, *Lactuca serriola* (Dm) 31, *Lamium amplexicaule* 21, *Lamium purpureum* 24, *Lepidium draba* (Dm) 22

Dominant species: *Chamomilla recutita* (C) 36, *Cirsium arvense* (C) 27, *Lactuca serriola* (C) 27, *Lepidium draba* (C) 27

Group 5: *Chamomilla recutita* (German Chamomile) plots

Number of plots: 18

Diagnostic species: *Lithospermum arvense* 20.6, *Sinapis arvensis* (C) 27.3

Constant species: *Lamium amplexicaule* 26, *Lamium purpureum* (Dm) 31, *Sinapis arvensis* (Dg) 21, *Stellaria media* (Dm) 53

Dominant species: *Lamium purpureum* (C) 33, *Stellaria media* (C) 67

3.3. Weed Vegetation-Environment Relationships

The Non-metric Multidimensional Scaling (NMDS) ordination, based on species composition and ecological indicator values (Figure 3), revealed that the most important ecological factors influencing the diversity and variability of weed vegetation in investigated medicinal crops were temperature and light in *Foeniculum vulgare* and *Mentha piperita* plots, soil reaction in *Melissa officinalis* and *Plantago lanceolata* plots, and nutrient content in *Chamomilla recutita* plots. The first ordination axis was positively correlated with light ($r=0.99392$; $P=0.001$) and negatively with moisture ($r=-0.87779$; $P=ns$). The second NMDS axis was positively correlated with temperature ($r=0.95785$; $P=0.001$) and nutrients ($r=0.96740$; $P=ns$), and negatively with soil reaction ($r=-0.99685$; $P=0.002$).




Figure 3. NMDS ordination analysis of weed vegetation in different medicinal crop plots. Group 1: *Melissa officinalis* (lemon balm) plots; Group 2: *Foeniculum vulgare* (fennel) plots; Group 3: *Mentha piperita* (peppermint) plots; Group 4: *Plantago lanceolata* (ribwort plantain) plots; Group 5: *Chamomilla recutita* (German chamomile) plots.

4. Discussion

The diversity, abundance, and frequency of weed flora in the surveyed medicinal crops (lemon balm, fennel, peppermint, ribwort plantain, German chamomile) are influenced by agro-ecological conditions, crop species, and cultivation practices. These findings are consistent with previous studies [4, 9, 12, 13, 32, 33, 34]. The high weed diversity recorded (109 weed species across 29 families and 87 genera) in a relatively small area, such as the Pančevo farm, suggests that the environmental conditions are favorable for weed flora development. However, this also indicates challenges in weed management practices [10, 34].

Grouping of the surveyed plots by crop species revealed that specific diagnostic weed species developed in lemon balm and ribwort plantain plots. Annual weeds such as *Crepis biennis*, *Senecio vernalis*, and *Brassica nigra* were predominant in lemon balm plots, while *Geranium molle* was characteristic of ribwort plantain plots. Weed vegetation in lemon balm and ribwort plantain was primarily influenced by soil reaction, whereas the weed composition in German chamomile plots was more strongly affected by nutrient availability. The relatively small overall influence of soil properties on weed composition may be due to the artificial adaptation of soil conditions through fertilization, mechanical weed control, and herbicide use [35].

The absolute dominance of annual weed species (68.8% therophytes and 10% hemitherophytes) in studied medicinal crops, which are part of arable lands, further confirms that these areas are under continuous agricultural pressure [15]. If these annual weed species, particularly those with high abundance, are not controlled promptly and effectively, they can severely reduce both the yield and quality of medicinal crops [9, 36]. Some annuals develop large canopies, such as: *Abutilon theophrasti*,

Amaranthus retroflexus, *Ambrosia artemisiifolia*, *Brassica nigra*, *Chenopodium* spp., *Datura stramonium*, and *Xanthium strumarium*, making them strong competitors for aboveground resources. Additionally, many weeds germinate and establish earlier (winter and early spring species) such as *Avena fatua*, *Bifora radians*, *Brassica nigra*, *Consolida* spp., *Capsella bursa-pastoris*, *Galium aparine*, *Lamium* spp., *Papaver rhoeas*, *Stellaria media*, *Veronica* spp., and *Viola arvensis*, giving them a competitive advantage for vital resources such as water, nutrients, and light.

Medicinal crops such as lemon balm, fennel, peppermint, ribwort plantain, and German chamomile, generally exhibit lower competitiveness against weeds, especially during their early growth stages, which contributes to high levels of weed infestation [9]. This competition can prolong flowering and fruiting phases and significantly decrease final yields [37]. Some weeds, like *Stellaria media* (observed in medicinal crops with an abundance of 1–8 and a frequency of III–V), are particularly problematic due to their ephemeral life cycle and ability to produce multiple generations per year [38]. Although classified as an annual, *S. media* can also propagate vegetatively in the short term. It germinates throughout the year under favorable conditions and effectively competes with neighboring plants [39].

Perennial weed species also pose substantial management challenges. Perennials such as *Artemisia vulgaris*, *Rumex crispus*, *Taraxacum officinale*, *Elymus repens*, *Cirsium arvense*, *Convolvulus arvensis*, *Lepidium draba*, *Sonchus arvensis*, and *Sorghum halepense*, were found with high frequency in the surveyed medicinal crops at Pančevo farm. The situation is more complex when their abundance is also significant, as seen with *Elymus repens* (abundance= 2–7), *Cirsium arvense* (ab.= 1–8), *Rumex crispus* (ab.= 2–8), *Sorghum halepense* (ab.= 1–8), *Taraxacum officinale* (ab.= 2–3), all of which were present in every surveyed field with medicinal crops. According to Miller [40], the combined application of multiple weed control strategies is typically more effective for managing perennial weeds.

Since medicinal crops generally allow only limited herbicide use [36, 41], repeated mechanical or other non-chemical measures are essential to deplete perennial weed populations [8] and reduce both the seed bank and vegetative reproductive structures in the soil. Rhizomatous grasses, particularly *Elymus repens* and *Sorghum halepense*, present persistent control challenges [42]. Furthermore, species like *Cirsium arvense* (which forms clean oases) and *Convolvulus arvensis* and *Calystegia sepium* (which twine around crop plants), with their substantial aboveground biomass, are aggressive competitors and can significantly reduce yields if not controlled in a timely manner [43, 44].

For sustainable and effective management of perennial weeds in medicinal crops, an integrated weed management approach is recommended, combining cultural and chemical methods with a focus on prevention and early intervention. Agronomic practices such as tillage, mowing, and grazing can help suppress rhizome spread and seed production. When herbicides are used, targeting weeds at their early, most susceptible stages are crucial for effective control.

To further mitigate weed issues, it is important to ensure that the plots are thoroughly cleaned after harvest, removing any residual crop plants such as in our cases, *Foeniculum vulgare*, *Cynara cardunculus*, and *Linum usitatissimum*, as these can emerge as volunteer crops in subsequent seasons and are often difficult to control chemically. Special attention should also be given to toxic weed species, such as *Bifora radians*, *Conium maculatum*, *Datura stramonium*, and *Solanum* spp., which contain various harmful alkaloids (atropine, coniine, conhydrine, gamma-coniceine, hyoscyne, hyoscyamine, N-methylconiine, pseudotropine, pseudoconhydrine, scopolamine, solasonine, solamargine, etc.). These compounds can significantly compromise the quality and safety of medicinal crops [45-47].

Ultimately, knowledge of weed diversity, abundance, and frequency in medicinal crops provides a reliable framework for accurately assessing weed infestation and serves as a foundation for strategic weed management planning. Weed control measures should be tailored to the most frequent, abundant, and dominant species specific to each cropland or locality. So far, no comprehensive documentation exists regarding the relative importance and quantitative weed

infestation levels in medicinal crop fields in Serbia. This study represents a valuable starting point for developing targeted weed management strategies for medicinal crop plantations in the region.

5. Conclusions

The diversity, abundance and frequency of weed species in medicinal crops are influenced by multiple factors, including crop type and genotype, agro-ecological conditions, and applied weed management strategies. This study confirmed the presence of 109 weed species across five different medicinal crops, comprising 75 annuals and 34 perennials, with a predominance of broadleaved weeds. The recorded weed flora spanned 29 families and 88 genera. The most frequent and abundant species included annual weeds such as *Amaranthus retroflexus*, *Chenopodium album*, *Galium aparine*, *Lactuca serriola*, *Lamium amplexicaule*, *Lamium purpureum*, *Papaver rhoeas*, *Stellaria media*, *Veronica hederifolia*, *Veronica persica*, as well as perennial weeds like *Elymus repens*, *Artemisia vulgaris*, *Cirsium arvense*, *Convolvulus arvensis*, *Lepidium draba*, *Rumex crispus*, *Sonchus arvensis*, *Sorghum halepense*, and *Taraxacum officinale*.

These findings provide a solid foundation for accurately assessing weed infestation levels in medicinal crop fields and highlight the need for species-specific and locally adapted weed management strategies. Considering the limited herbicide options available in medicinal crop production, integrating cultural, mechanical, and targeted chemical control measures is essential for sustainable weed management. The results of this study contribute to filling the existing knowledge gap regarding weed flora in medicinal crop plantations in Serbia and can serve as a valuable reference for the development of effective weed control programs tailored to this specific cropping system.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, S.V., D.B. and A.D.; methodology, S.V. and D.B.; software, S.A. and U.Š.; validation, S.V., D.B. and T.M.; formal analysis, S.V. and T.T.; investigation, D.B. and A.D.; resources, T.M., A.D. and M.R.; data curation, S.A. U.Š. and S.V.; writing—original draft preparation, D.B. and S.V.; writing—review and editing, T.M. and S.V.; visualization, A.D., M.R. and T.T.; supervision, S.V.; project administration, A.D. and T.M.; funding acquisition, S.V., D.B., T.M. and A.D. All authors have read and agreed to the published version of the manuscript.”

Funding: This research was funded by Ministry of Science, Technological Development and Innovation of the Republic of Serbia, grant numbers 451-03-136/2025-03/200003 and 451-03-137/2025-03/200116 and the APC was funded by Ministry of Science, Technological Development and Innovation of the Republic of Serbia, grant number 451-03-137/2025-03/200116.

Data Availability Statement: We encourage all authors of articles published in MDPI journals to share their research data. In this section, please provide details regarding where data supporting reported results can be found, including links to publicly archived datasets analyzed or generated during the study. Where no new data were created, or where data is unavailable due to privacy or ethical restrictions, a statement is still required. Suggested Data Availability Statements are available in section “MDPI Research Data Policies” at <https://www.mdpi.com/ethics>.

Acknowledgments: In this section, you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments). Where GenAI has been used for purposes such as generating text, data, or graphics, or for study design, data collection, analysis, or interpretation of data, please add “During the preparation of this manuscript/study, the author(s) used [tool name, version information] for the purposes of [description of use]. The authors have reviewed and edited the output and take full responsibility for the content of this publication.”

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

KAN	calcium ammonium nitrate
T	therophytes
TH	thero-hemicryptophytes
G	geophytes
H	hemicryptophytes
Ch	chamaephytes
P	phanerophytes
Dg	diagnostic species
Dm	dominant species
C	constant species
NMDS	non-metric multidimensional scaling
EIVs	ecological indicator values

References

1. www.stat.gov.rs (accessed on 2 May 2025)
2. World Bank. Exploring the Potential of Agriculture in the Western Balkans, Washington, DC, 2018.
3. FAO. Small Family Farms Country Factsheet: Serbia. Food and Agriculture Organization of the United Nations, available at <https://www.fao.org/3/cb3863en/cb3863en.pdf>
4. Bayisa, N.G.; Hundesa, N. Assessment and identification of weed flora associated to medicinal and aromatic plants at Wondo Genet District, Ethiopia. *Int. J. Agric. Biosci.* 2017, 6 (3), 136-140. <https://doi.org/10.11648/j.ajpb.20220702.12>
5. Tomado, T.; Milberg, P. Weed flora in arable fields of eastern Ethiopia with emphasis on the occurrence of *Parthenium hysterophorus* L. *Weed Res.* 2000, 40, 507-521. <https://doi.org/10.1046/j.1365-3180.2000.00208.x>
6. Kashem, M.A.; Faroque, M.A.; Bilkis, S.E. Weed management in Bangladesh: Policy issues to better way out. *J. Sci. Found.* 2009, 7, 59-67.
7. Kubiak, A.; Wolna-Maruwka, A.; Niewiadomska, A.; Pilarska, A.A. The Problem of Weed Infestation of Agricultural Plantations vs. the Assumptions of the European Biodiversity Strategy. *Agronomy* 2022, 12, 1808. <https://doi.org/10.3390/agronomy12081808>
8. Matković, A.; Božić, D.; Filipović, V.; Radanović, D.; Vrbničanin, S.; Marković, T. Mulching as a physical weed control method applicable in medicinal plants cultivations. *Lekovite sirovine* 2015, XXXV (35), 37-51. <https://doi.org/10.5937/leksir1535037M>
9. Hendawy, S.F.; Abouzienna, H.F.; Abd El-Razik, T.M.; Amer, H.M.; Hussein, M.S. Winter weeds and its control in the medicinal plants in Egypt: a survey study. *Egypt. Pharm. J.* 2019, 18(1), 16–26. https://doi.org/10.4103/epj.epj_13_18
10. Pala, F.; Mennan, H. Weed problem in medicinal plants. In *New development on medicinal and aromatic plants*, Özyazici G., Ed.; Iksad Publishing hous: Ankara, Turkey, 2021, Chapter 3, pp. 67-89.
11. Pinke, G.; Blazsek, K.; Magyar, L.; Nagy, K.; Karácsony, P.; Czúcz, B.; Botta-Dukát, Z. Weed species composition of conventional soybean crops in Hungary is determined by environmental, cultural, weed management and site variables. *Weed Res.* 2016, 56 (6), 470–481. <https://doi.org/10.1111/wre.12225>
12. Bisht, A.S. Weed Floral Diversity of Medicinal Value in Terraces of Horticulture Crop Fields in Bharsar, Uttarakhand, India. *IJPGR* 2017, 30 (2), 153-161. <https://doi.org/10.5958/0976-1926.2017.00022.5>
13. Ljevnaić-Mašić, B.; Brdar-Jokanović, M.; Džigurski, D.; Nikolić, Lj.; Meseldžija, M. Weed Composition in conventionally and organically grown medical and aromatic plants. *Acta Sci. Pol. Hortorum Cultus* 2022, 21(4), 115-126. <https://doi.org/10.24326/asphc.2022.4.12>
14. Reddy, C. A study on crop weed competition in field crops. *J Pharmacogn Phytochem* 2018, 7(4), 3235-3240.
15. Vrbničanin, S.; Božić, D. Korovi. [Weeds]. Univerzitet u Beogradu, Poljoprivrdni fakultet: Beograd; Serbia, 2021; pp. 1-432. [in Serbian]
16. Wiersema, J.H.; León, B. *World Economic Plants: A Standard Reference*. CRC Press: New York, USA, 1999, pp. 792. <https://doi.org/10.1201/9781482274431>

17. Vrbničanin, S.; Kojić, M. Biološka i ekološka proučavanja korova na području Srbije: razvoj, sadašnje stanje i perspektive. [Biological and ecological research of weeds in Serbia: development, current status, perspectives]. *Acta herbologica* 2000, 9 (1), 41-59. [in Serbian]
18. Isah, T. Stress and defense responses in plant secondary metabolites production. *Biol. Res.* 2019, 52(1), 39. <https://doi.org/10.1186/s40659-019-0246-3>
19. Puschner, B.; Peters, A.; Woods, L. Toxic weeds and their impact on animals. In *Proceedings of 36th Western Alfalfa & Forage Symposium, Reno, USA, 11-13 December 2006*.
20. Carrubba, A. Weed and weeding effects on medicinal herbs. In *Medicinal Plants and Environmental Challenges*; Ghorbanpour, M., Varma, A., Eds.; Springer: Cham, Germany, 2017; pp. 295–327. https://doi.org/10.1007/978-3-319-68717-9_17
21. Kropff, M.J.; Spitters, C.J.T. A simple model of crop loss by weed competition from early observations on relative leaf area of the weeds. *Weed Res.* 1991, 31(2), 97-110.
22. Braun-Blanquet, J. *Pflanzensoziologie. Grundzuger der Vegetationskunde*, 3 Aufl.; Springer: Wien, Austria, 1964. [in German]
23. Westhoff, V.; van der Maarel, E. The Braun-Blanquet Approach. In: *Ordination and classification of communities*, Whittaker, R.H., Ed.; Dr. W. Junk: The Hague, Netherland, 1973; pp. 616-726.
24. Euro+Med PlantBase, Euro+Med PlantBase – the information resource for Euro-Mediterranean plant diversity, 2025. <https://www.europlusmed.org>. Accessed 12 Jan 2025.
25. Raunkier, C. *The life forms of plants and statistical plant geography*. Calderon Press: Oxford, UK, 1934.
26. Tichý, L. JUICE, software for vegetation classification. *J. Veg. Sci.* 2002, 13, 451-453. <https://doi.org/10.1111/j.1654-1103.2002.tb02069.x>
27. Chytrý, M.; Tichý, L.; Holt, J.; Botta-Dukat, Z. Determination of diagnostic species with statistical fidelity measures. *J Veg Sci.* 2002, 13, 79-90. <https://www.jstor.org/stable/3236655>
28. RStudio Team. *RStudio: Integrated Development for R*; RStudio, PBC: Boston, MA, USA, 2022; Available online: <http://www.rstudio.com> (accessed on 5 March 2025)
29. Oksanen, J.; Simpson, G.; Guillaume, B.; Kindt, R.; Legendre, P.; Minchin, P.; O'Hara, R.B.; Solymos, P.; Henry, M.; Stevens, M.; Szoecs, E.; Wagner, H.; Barbour, M.; Bedward, M.; Bolker, B.; Borcard, D.; Carvalho, G.; Chirico, M.; De Caceres, M.; Durand, S.; Evangelista, H.; FitzJohn, R.; Friendly, M.; Furneaux, B.; Hannigan, G. O.; Hill, M.; Lahti, L.; McGlenn, D.; Ouellette, M-H.; Cunha, E.; Smith, T.; Stier, A.; Ter Braak, C.; Weedon, J. *Cran.rproject*. Available from: <https://cran.r-project.org/web/packages/vegan/index.htm>
30. Hellinger E. Neue Begründung der Theorie quadratischer Formen von unendlichvielen Veränderlichen. *J. für Reine Angew. Math.*, 1909, 136, 210–271. <http://eudml.org/doc/149313>
31. Pignatti, S. Valori di bioindicazione delle piante vascolari della flora d'Italia. *Br.-Bl.* 2005, 39, 1-97.
32. Travlos, I.S.; Cheimona, N.; Roussis, I.; Bilalis, D.J. Weed-species abundance and diversity indices in relation to tillage systems and fertilization. *Front. Environ. Sci.* 2018, 6, 11. <https://doi.org/10.3389/fenvs.2018.00011>
33. Mallick, D.; Paul, P.; Mondal, S.; Pal, A.; Dasgupta, S.; Chowdhury, M. Survey and documentation of the Weed Flora in NBU Garden of Medicinal Plants. *NBU J. Plant Sci.* 2020, 12, 15-25. <https://ir.nbu.ac.in/handle/123456789/4545>
34. Kobir, M.S.; Paul, S.; Hajong, P.; Harun-Or-Rashid, M.; Rahman, M.H. Taxonomic diversity of weed flora in pulse crops growing field at south-western part of Bangladesh. *Arch. Agric. Environ. Sci.* 2021, 6(4), 436-440. <https://dx.doi.org/10.26832/24566632.2021.060404>
35. Cimalová, Š.; Lososová, Z. Arable weed vegetation of the northeastern part of the Czech Republic: effects of environmental factors on species composition. *Plant Ecol.* 2009, 203, 45–57. <https://doi.org/10.1007/s11258-008-9503-1>
36. Upadhyay, R.K.; Baksh, H.; Patra, D.D.; Tewari, S.K.; Sharma, S.K.; Katiyar, R.S. Integrated weed management of medicinal plants in India. *International Journal of Medicinal and Aromatic Plants* 2011, 1(2), 51-56.
37. Carrubba, A.; Militello, M. Nonchemical weeding of medicinal and aromatic plants. *ASD* 2013, 33, 551–561. <https://doi.10.1007/s13593-012-0122-9>

38. Roberts, H.A.; Feast, P.M. Seasonal distribution of emergence in some annual weeds. *Exp. Hortic.* 1970, 21, 36-41.
39. Pacanoski, Z. *Stellaria media* (L.) Vill. (common chickweed) – a strong or weak competitor in the autumn and early-spring sown crops? *Acta herbologica* 2024, 33(2), 77-89. <https://doi.org/10.5937/33ah-51194>
40. Miller, T.W. Integrated Strategies for Management of Perennial Weeds. *Invas. Plant Sci. and Mana.* 2017, 9(2), 148-158. <https://doi.org/10.1614/IPSM-D-15-00037.1>
41. Darre, A.; Novo, R.; Zumelzu, G.; Bracamonte, R. Chemical control of annual weeds in *Mentha piperita*. *Agriscientia* 2004, 21, 39-44.
42. Squires, C.C.; Walsh, M.J. *Sorghum halepense*. In *Biology and Management of Problematic Crop Weed Species*, Chauhan, B.S., Ed.; Elsevier: London, UK, 2021; Chapter 18, pp. 391-405.
43. Davis, S.; Mangold, J.; Menalled, F.; Orloff, N.; Miller, Z.; Lehnhoff, E. A Meta-Analysis of Canada Thistle (*Cirsium arvense*) Management. *Weed Sci.* 2018, 66(4), 548-557. <https://doi.org/10.1017/wsc.2018.6>
44. Al-Abbasi, S.H.A.; Mustafa, H.A.; Al-Naqib, A.T.H.; Adnan, A.; Al-Majmaei, M.; Hameed, A.M.; Alazzawi, H.A.H.; Muhammed aldouri, G.A.; Awad, A.A.; Majeed, Q.A.; Dyaah yasin, A. Weed of *Convolvulus arvensis* damage and control methods. *Chemical and Environmental Science Archives* 2021, 1(1), 28-31.
45. Berkov, S.; Zayed, R.; Doncheva, T. Alkaloid patterns in some varieties of *Datura stramonium*. *Fitoterapia* 2006, 77, 179-182. <https://doi.org/10.1016/j.fitote.2006.01.002>
46. John, H.; Rychlik, M.; Thiermann, H.; Schmidt, C. Simultaneous quantification of atropine and scopolamine in infusions of herbal tea and Solanaceae plant material by Matrix-Assisted Laser Desorption/Ionization Time-of-Flight (Tandem) Mass Spectrometry. *Rapid Commun. Mass Spectrom* 2018, 32, 1911–1921.
47. de Nijs, M.; Crews, C.; Dorgelo, F.; MacDonald, S.; Mulder, P.P.J. Emerging Issues on Tropane Alkaloid Contamination of Food in Europe. *Toxins* 2023, 15, 98. <https://doi.org/10.3390/toxins15020098>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.