

Review

Not peer-reviewed version

# Naphthoquinones and Their Importance in Facing the Challenge of Parkinson's Disease

Thaís Barreto Santos , Leonardo Gomes Cavaliere De Moraes , [Paulo Anastácio Furtado Pacheco](#) , Douglas Galdino Dos Santos , Rafaella Machado de Assis Cabral Ribeiro , Caroline dos Santos Moreira , [David Rodrigues Da Rocha](#) \*

Posted Date: 16 October 2023

doi: 10.20944/preprints202310.0928.v1

Keywords: naphthoquinones; Parkinson disease; P2X7 receptor



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Review

# Naphthoquinones and Their Importance in Facing the Challenge of Parkinson's Disease

Thaís Barreto Santos,<sup>a</sup> Leonardo Gomes Cavalieri de Moraes,<sup>a</sup> Paulo Anastácio Furtado Pacheco and<sup>a</sup> Douglas Galdino dos Santos,<sup>a</sup> Rafaella Machado de Assis Cabral Ribeiro,<sup>a</sup> Caroline dos Santos Moreira,<sup>a,b</sup> David Rodrigues da Rocha<sup>a</sup>

Universidade Federal Fluminense, Instituto de Química, Outeiro de São João Batista s/nº, CEP 24.020-141, Niterói-RJ, Brazil

Instituto Federal do Rio de Janeiro, Campus Paracambi, Rua Sebastião Lacerda s/nº, Fábrica, CEP 26.600-000, Paracambi-RJ, Brazil

**Abstract:** Parkinson's disease (PD) is a degenerative disease that affects approximately 6.1 million and is primarily caused by the loss of dopaminergic neurons. Naphthoquinones have several biological activities explored in the literature, including neuroprotective effects. Therefore, this review shows an overview of natural and synthetic naphthoquinones with neuroprotective effects and with potential for the development of molecules for the treatment of PD. Many examples of natural naphthoquinones, such as shikonin, plumbagin and vitamin K, exerted a neuroprotective effect by preventing oxidative stress, in addition to multiple mechanisms. Synthetic naphthoquinones with inhibitory activity on the P2X7 receptor were also found, leading to a neuroprotective effect on Neuro-2a cells. It was found that naphthazarin can act as inhibitors of the MAO-B enzyme. Vitamin K and synthetic naphthoquinones hybrids with tryptophan or dopamine showed inhibition of the aggregation of  $\alpha$ -synuclein. A library of synthetic derivatives of juglone and naphthazarin were able to protect Neuro-2a cells against neurodegenerative effects of neurotoxins. We report many examples of naphthoquinones that act in therapeutic pathways for the treatment of PD. In addition, routes for producing synthetic derivatives were also discussed. With the data presented, 1,4-naphthoquinones can be considered a promising class in the treatment of PD.

**Keywords:** naphthoquinones; Parkinson disease; P2X7 receptor

## 1. Introduction

Parkinson's disease (PD) is a pathology of neurological origin, degenerative of the region present in the central nervous system known as substantia nigra. It was described in 1817, when James Parkinson published his monograph entitled *An Essay on the Shaking Palsy* describing the cardinal symptoms of the disease that was later named after him [1].

It is a very common disease, with approximately 6.1 million people affected worldwide in 2016 and, for reasons that are still not fully understood, in the last two decades the incidence, prevalence and deaths caused by this disease have increased [2].

### 1.1. Epidemiology and risk factors

Worldwide, the disease occurs without notable epidemiological differences, the only exceptions being a rapid increase in cases in high-income countries in Europe and a disproportionately rapid increase in new cases in China. The number of PD-related deaths and disabilities has more than doubled in the past two decades [2].

Although PD is age-related, including being the biggest risk factor, where the incidence and prevalence regularly increase with age, it is a mistake to believe that the disease exclusively affects the elderly. For about 25% of affected individuals, age of onset is less than 65 years and for 5-10% it is less than 50 years. For affected individuals younger than 40 years, the term early-onset Parkinson's disease is used [3].

Although Parkinson's disease affects both sexes, women have some differences in relation to men, such as the fact that the incidence of the disease is lower, especially in individuals aged 50 to 59 years, and the age at which the onset of the disease is higher, however, they have a greater risk of developing dyskinesia (involuntary muscle movements), urinary problems, depression and fluctuations in motor and non-motor response, due to their low body weight, leading to relative overdose [4]. Men, on the other hand, live a greater number of years with the disabilities caused by PD and have greater cognitive loss [5]. It is important to emphasize that the health behavior is different between men and women with PD, with late or less frequent access to specialized medical care, which results in undertreatment, including less use of neurosurgical interventions [6].

Some environmental factors are linked to the risk of developing PD, such as exposure to certain pesticides and rural life [7]. Likewise, some substances such as 1-methyl-4-phenyl tetrahydropyridine (MPTP) and annonacin can cause nigrostriatal cell death and a form of atypical parkinsonism [8]. Furthermore,  $\beta$ 2 adrenergic receptor antagonists have been associated with an increased risk of PD, whereas  $\beta$ 2 adrenergic receptor agonists appear to reduce it [9]. On the other hand, there is an inverse association between the risk of PD and coffee consumption [10], smoking [7], statins [11] and calcium channel blockers [12], while conflicting evidence is available regarding the use of non-steroidal anti-inflammatory drugs [13].

Family history is a risk factor for PD and the relative risk in first-degree relatives of PD cases increases approximately two to three times compared to controls. Familial forms of PD represent 5% to 15% of cases [14].

### 1.2. Pathophysiology

The main pathological characteristics of PD are the loss of dopaminergic neurons with consecutive depigmentation of the substantia nigra and the presence of Lewy bodies. Lewy bodies are round, eosinophilic, intraneuronal inclusions with a hyaline nucleus and a pale peripheral halo that are composed of more than 90 proteins; its main components are  $\alpha$ -synuclein and ubiquitin [15].  $\alpha$ -Synuclein has the ability to misfold, become insoluble, and form amyloid-rich aggregates in  $\beta$  sheets that accumulate and form intracellular inclusions. Intermediates in this aggregation process are the toxic oligomeric and protofibrillar forms that impair mitochondrial [16]. Lysosomal and proteasomal function [17], damage biological membranes [18] and the cytoskeleton [19], alter synaptic function and cause neuronal degeneration [20]. At the time of diagnosis, it is presumed that up to 60% of dopaminergic neurons have been lost [21].

Therefore, a model of Lewy body formation and  $\alpha$ -synuclein deposition was proposed, which starts in the dorsal motor nucleus of the glossopharyngeal and vagus nerves and anterior olfactory nucleus, with progressive spread to the brainstem and, in subsequent phases, to the mesocortex and allocortex, and finally to the neocortex [22].  $\alpha$ -Synuclein tends to spread through neurons in a similar way to a prion, and this transmission mechanism is probably behind the progression of the pathological changes described above. Furthermore, some data suggest that  $\alpha$ -synuclein aggregation may start in the gut autonomic plexus and spread rostrally and that this may be influenced by the gut [23] microbiome.

### 1.3. Disease Mechanisms

The gene coding for  $\alpha$ -synuclein, SNCA, was the first linked to PD, and A53T, a mutation that gives  $\alpha$ -synuclein a greater predisposition to unfold and aggregate, was the first identified pathogenic SNCA mutation. The similar features of all SNCA mutations include early disease onset, rapid progression of motor signs, and notable presence of non-motor features, including cognitive impairment [24].

Likewise, impairment of mitochondrial function is another important mechanism for the disease and is usually linked to different genes involved in familial forms of PD [25]. The PINK1 and PRKN genes interact in a mitochondrial quality control pathway, their mutations being the main causes of autosomal recessive and early-onset PD, accounting for up to 77% of cases of juvenile Parkinson's, when the age of onset is below 20 years and 10-20% of early-onset Parkinson's [26], however its

progression is slower and responds well to treatment. The DJ-1 gene, on the other hand, plays an important role in regulating the flow of calcium in the mitochondria, which protects the cell from oxidative stress produced by the regulatory activity of dopaminergic neurons and from dopamine toxicity [27]. There is information of mitochondrial DNA mutations in the substantia nigra of PD brains [28].

Likewise, a growing body of evidence links PD to dysfunction in cellular clearance pathways and various genes linked to autophagy [29]. Mutations in the LRRK2 gene affect autophagy and delay the degradation of  $\alpha$ -synuclein, which favors its accumulation. These mutations are seen at lower rates in sporadic cases and represent 3-41% of familial PD cases [24,30].

Likewise, mutations in GBA1, which encodes the enzyme glucocerebrosidase (GCase) are highly prevalent in individuals with PD and currently constitute the most important genetic risk factor, where 5-25% of patients have GBA1 mutations. GBA-linked PD has an early onset and a severe course, particularly with rapid cognitive decline. The GBA contribution is complex and there are interactions with different pathways implicated in the pathogenesis of PD, such as a reciprocal relationship with  $\alpha$ -synuclein accumulation, endoplasmic reticulum stress and mitochondrial dysfunction [31].

Recently, rare variants of LRP10, a protein that travels between the trans-Golgi network, endosomes and plasma membrane, have been associated with familial PD, dementia from PD and dementia with Lewy bodies. Other proteins included in the same network were linked to PD, including VPS35 and GGA1. However, more research is needed to elucidate the pathogenetic role of alterations in these pathways in PD and other neurodegenerative disorders [32]. Why the dopaminergic neurons of the substantia nigra pars compacta are particularly susceptible to degeneration remains unknown, but it has been pointed out that the autonomous pace-making nature of the dopaminergic neurons of the substantia nigra and calcium homeostasis play an important role [33]. Finally, the importance of the microbiome in the pathogenesis of PD has attracted increasing interest and the pathogenetic mechanisms include alteration of dopamine synthesis and metabolism, dysregulation and inflammation of the immune system, and changes in enteral mucosal permeability [34].

#### 1.4. Clinical Manifestations

PD presents a set of motor and non-motor characteristics, whose expression may vary between individuals, however, all patients must exhibit the main clinical characteristics and respond to dopaminergic therapy to meet the parameters for the diagnosis of PD. Thus, cardinal motor symptoms include tremor, bradykinesia, hypokinesia, akinesia, rigidity, postural instability, gait and speech disorders, among others. Responsiveness of motor symptoms to levodopa administration is an important feature [35].

The clinical picture among patients can be quite heterogeneous, which allows the definition of different motor subtypes, such as “dominant tremor”, “postural instability and gait difficulty” or “undetermined”. The possible association of the subtype with etiological or prognostic aspects and with the response to treatment is what makes its definition interesting, for example, PD “dominant tremor” has been associated with a slower and less disabling progression compared to “instability posture and gait difficulty” [36].

Even though historically defined as a movement disorder, non-motor signs are important aspects of the clinical picture. Non-motor signs range from dysphagia and sialorrhea to autonomic, sensory, sleep, gastrointestinal, cognitive, and neuropsychiatric disorders; are little investigated by physicians; however, they have a great impact on the patient's health-related quality of life [37].

Some other symptoms, generally called prodromal or premotor symptoms, may manifest up to 10 years before diagnosis and the appearance of motor symptoms. The most recognized are constipation, hyposmia, depression, and rapid eye movement sleep disorder, but may include anxiety, visual changes, and other autonomic disorders. The prodromal phase of PD presents itself as a unique opportunity to identify those at high risk of developing PD before the occurrence of massive neurodegeneration, bringing important information about the mechanisms of the disease

and its progression, proving to be a promising therapeutic window for neuroprotective treatments; therefore, many efforts have been made to better recognize this phase [38].

## 2. Quinones as a privileged structure for the development of new derivatives for PD

Quinones are a class of organic compounds widely distributed in nature, whose structure has an unsaturated ring of six carbon atoms and two carbonyl groups, which can be situated in an *ortho* or *para* position. Furthermore, quinones are classified due to their aromatic system type as benzoquinones (1), naphthoquinones (2), anthraquinones (3) and phenanthrenequinones (4) (Figure 1) [39,40]. In this paper we will make an overview of natural and synthetic 1,4-naphthoquinones with neuroprotective effect and potential activity for the treatment of PD.




Figure 1. General structures of quinones types.

## 3. Naphthoquinones

Naphthoquinones are the most common type of quinones and compose an important series of natural distribution metabolites of plants, animals, fungi and bacteria. [41] Besides these substances have great importance in vital biochemical processes [42], there are many biological effects associated to 1,4-naphthoquinones, the isomer with the highest incidence [43], such as antioxidant/anti-inflammatory [44], antimalarial [45], antitumor [46], tripanocidal [47], antifungal [48] and antibacterial [49]. These effects are related to the formation of reactive oxygen species (ROS), inducing oxidative stress in cells [50], which causes irreversible damage to DNA and proteins leading to apoptosis [51].

However, another important biological effect is the neuroprotective, which occurs by the ability of some 1,4-naphthoquinones to maintain the redox potential of neuronal cells, acting on enzymes responsible for the balance of ROS. This is what protects cells from oxidative stress generated by neurotoxins, capturing free radicals [52].

In this sense, it is worth emphasizing some 1,4-naphthoquinones related to this neuroprotective effect. First, the lapachol (5) is isolated from *Tabebuia avellanedae* and extremely used in American folk medicine for the treatment of many diseases such as cancer, lupulus and infections [53]. The juglone (6) is found in some species of the family Jugladaceae, for example *Juglans regia*, *Juglans nigra* and *Juglans cineraria*, and your use is reported against ringworm, fungal, bacterial and viral infections and as cure for heat stroke [54]. Finally, the shikonin (7) is present in the root extract of *Lithospermum erythrorhizon*, named *Zicao*, and was used in traditional Chinese medicine for its anti-inflammatory effects, until studies revealed other interesting properties, like anticancer, antimicrobial and wound healing [55] (Figure 2).





Figure 2. Chemical structures of lapachol (5), juglone (6) and shikonin (7).



### 3.1. Neuroprotective effect of naphthoquinones


Several studies support a neuroprotective effect of naphthoquinones and their derivatives through different mechanism of action. Natural and synthetic naphthoquinones have been assessed for their antioxidant properties. Oxidative stress is a common pathological feature of traumatic brain injury [56]. Therefore, the use of antioxidant drugs can have a positive impact on the treatment of this neurological condition. It has been identified that the natural naphthoquinone thymoquinone (8) and, particularly, its reduced form, thymohydroquinone (9), present strong antioxidant activity (Figure 3) [57]. In this sense, Al-Majed et al. investigated the potential neuroprotective effect of this metabolite, that can be obtained from black cumim (*Nigella sativa*) essential oil, in transient forebrain ischemia-induced neuronal damage in the rat hippocampus [58].



**Figure 3.** Chemical structure of Thymoquinone (8) and its metabolite Thymohydroquinone (9).


Pretreatment with thymoquinone reduced ischemia-induced neuronal damage by preventing hippocampal neuronal cells death. In addition, thymoquinone also decreased levels of malondialdehyde and increased the levels of antioxidant enzymes in ischemic rats. Thymoquinone and thymohydroquinone avoided *in vitro* lipid peroxidation in hippocampal homogenate induced by iron-ascorbate with  $IC_{50}$  of 12 and 3  $\mu$ M, respectively. Thymoquinone has also showed neuroprotective effects on different chemical-induced models. In model of neuronal injury promoted by chronic exposure to toluene, treatment with thymoquinone (50 mg/kg body weight) for 12 weeks caused a significant morphologic improvement on neurogenerative status of frontal cortex in rats [59]. Treatment with thymoquinone also showed beneficial effects on lead-induced brain damage rats [60]. Rats treated with thymoquinone at a dose of 20 mg/kg body weight for one month exhibited similar brain histology to the control group. In an experimental model of arsenic-promoted neurotoxicity, thymoquinone (10 mg/kg body weight) mitigated deleterious effects by suppressing oxidative stress in the cerebral cortex, cerebellum, and brain stem [61].

Based on previous results that showed a neuronal anti-inflammatory role of Shikonin (7), the main bioactive compound isolated from the roots of *Lithospermum erythrorhizon* (Figure 4), Wang and coworkers studied the potential antioxidant activity of this naphthoquinone *in vitro* and in a murine cerebral/reperfusion injury. Treatment with shikonin at 50 mg/kg intragastric protected from neurodegeneration after ischemia/reperfusion induction as measured by neurological deficit, cerebral infarct volume and histopathological abnormalities. Shikonin also prevented oxidative stress markers and increased antioxidant defenses in the treated group [62]. Using computational tools and cellular assays, Wang *et al* identified among a group of 12 natural compounds that acetylshikonin (10) and its derivatives could inhibit the activity of acetylcholinesterase (Figure 4). In addition, these compounds exhibited antiapoptotic activity *in vitro* by preventing oxidative stress due loss of mitochondria membrane potential in mammal neuron cell lines [63].




**Figure 4.** Chemical structure of Shikonin (7) and Acetylshikonin (10).

It has been shown that plumbagin (11) (Figure 5), a naphthoquinone that can be obtained in the roots of *Plumbago zeylanica* L., exerts neuroprotective effects by modulating multiple essential pathways. Upon oxidative stress signals, the nuclear factor E2-related factor 2 (Nrf2) is activated and binds to antioxidant response element (ARE) upregulating the expression of genes coding for antioxidant defense mechanisms [64]. Incubation of human neuroblastoma cell line with plumbagin increased Nrf2 nuclear localization and expression of Nrf2/ARE-dependent genes. Plumbagin also protected against oxidative stress-induced cell death in neuroblastoma cell line and rat primary cortical neurons after oxidative and metabolic stimulus. This protective effect was reversed by RNA interference-mediated Nrf2 silencing. *In vivo* experiments showed that plumbagin treatment reduced brain damage and neurological deficit after ischemic injury [65]. In isoflurane-induced neurotoxicity model, plumbagin treatment reduced apoptosis and expression of pro-apoptotic factors in hippocampus. Moreover, plumbagin downregulated PI3K/Akt pathway while upregulating ERK1/2 levels [66]. In LPS-activated microglial BV-2 cell line, treatment with plumbagin inhibited nitric oxide production more potently than the selective iNOS inhibitor L-N6-(1-iminoethyl) lysine (L-NIL). Also, immunofluorescence assays showed that plumbagin inhibited inducible nitric oxide synthase (iNOS) expression and pro-inflammatory cytokines [67].




**Figure 5.** Chemical Structure of Plumbagin (11).

Corroborating the high potential of naphthoquinones as source of effective neuroprotectors, Choi *et al* identified the compound fusarubin (12), isolated from *Fusarium solanum*, an endophytic fungus present in leaves of the medicinal plant *Morus alba* (Figure 6). This naphthoquinone exhibited reactive oxygen species (ROS) scavenging activity, preventing oxidative stress-mediated cell death in murine hippocampal HT22 cell line. Using *in silico* approach, the authors proposed that the mechanism of action of this compound may be related to modulation of ubiquinone levels [68].



**Figure 6.** Chemical structure of Fusarubin (12).


Vitamin K (13) has also been investigated for neuroprotective actions against different neurological insults. It has been demonstrated that vitamin K, or more specifically menaquinone-4 (14) (MK-4), is present in brain tissues [69]. In experimental model of ischemia/reperfusion injury, MK-4 exerted its beneficial effects by inhibiting neuroinflammation markers and neurotoxicity. In addition, MK-4 treatment improved behavioral and cognitive parameters after ischemic insult. MK-4 administration resulted in the elevation of superoxide dismutase (SOD) activity, an essential antioxidant defense mechanism [70]. In agreement with these results, vitamin K prevented cell death mainly by avoiding intracellular glutathione (GSH) depletion in a chemical-induced neurotoxic model with methylmercury [71]. In a different neurotoxicity model induced by amyloid-(A $\beta$ ), vitamin K also suppressed death of neuronal cells, production of ROS and caspase-3 activation, suggesting that protective effect may occur by A $\beta$ -mediated apoptosis. In this work, it was identified that vitamin K protection mechanism might be related to phosphatidylinositol 3-kinase (PI3K) signaling pathway (Figure 7) [72].



**Figure 7.** Structure of Vitamin K1 (13) and K2 (14).


Considering these actions, Josey and collaborators designed a series of vitamin K derivatives and studied the neuroprotective effect of these compounds. They employed a chemical-based approach to modify and optimize the vitamin K pharmacophore to obtain new derivatives with improved pharmacological properties. Particularly, one derivative (17a) showed superior protective potency in vitro and higher safety index with low toxicity (Figure 8) [73].






**Figure 8.** Synthesis and in vitro activity of Vitamin K derivatives.

Other synthetic naphthoquinones have also been investigated in regard of neuroprotective effects. Nepovimova's group designed and synthesized a group of hybrid molecules containing a naphthoquinone core associated to tacrine, the first drug released for treatment of Alzheimer disease (Figure 9). The synthetic strategy had as starting point the compound memoquine (**18**), which exhibited inhibitory activity for acetylcholinesterase (AChE) and amyloid- $\beta$  aggregation, besides free-radical scavenging function. Following a structural simplification approach, they identified the compound **19** that retained the promising biological activity of the parental compound. For the synthesis of the hybrid compounds **20**, they employed a convergent approach and kept the 1,4-naphthoquinone core that was associated to the tacrine moiety through methylene linker with proper size chain. Notably, two compounds exhibited excellent performance in the inhibitory AChE and A $\beta$  aggregation assays. These compounds showed good safety profile on mouse cortical neurons Neuro2A (N2A) and primary rat cerebellar granule neurons. In addition, the compounds prevented A $\beta$ -induced neurotoxicity and tert-butyl hydroperoxide (TBH)-mediated ROS production [74].




**Figure 9.** Hybrid compounds containing a naphthoquinone and tacrine units.

Neurological diseases can be triggered by persistent activation of microglia and release of pro-inflammatory cytokine. Škandík and collaborators associated two electrophilic pharmacophores to generate a hybrid naphthoquinone-flavonoid derivative and studied its effect on modulation of the Nrf2 pathway in BV-2 microglial cells (Figure 10). This derivative at a non-toxic concentration reduced iNOS, COX-2 and TNF $\alpha$  levels on LPS-stimulated microglia. In addition, the authors also observed upregulation of antioxidant mechanisms upon treatment with this compound [75].



**Figure 10.** Structure of hybrid compound containing naphthoquinone-quercetin units.

Besides the neurotoxic impact of aggregated proteins, neuroinflammation stands out as a primary contributor to neurodegenerative diseases, including Alzheimer's disease (AD) and Parkinson's disease [76]. Upon activation, immune cells located within the central nervous system, such as microglia, generate a cascade of inflammatory mediators that harm neurons. Additionally, the release of extracellular ATP, primarily due to cell death, can trigger purinergic receptors on the surface of glial cells, ultimately prompting the synthesis of neurotoxic cytokines [77]. Among the cellular components involved in the activation of microglia and subsequent pathological changes associated with several neurodegenerative diseases, the P2X7 purinoreceptor plays a pivotal role [78]. Given its significance, Pislyagin et al reported the synthesis of a small library of 1,4-naphthoquinone derivatives and their antagonistic effects against mouse P2X7R. An initial screening with a  $\text{Ca}^{2+}$  influx assay, they identified four compounds with promising inhibitory activity (Figure 11). Subsequent biological evaluation of these compounds in Neuro-2a cells evidenced a potent blockade of dye uptake, decrease of ROS and NO production and protection of ATP-mediated toxicity. Molecular docking indicated that the naphthoquinone derivatives might exert their inhibitory activity due to interaction with the allosteric site located in the extracellular region of P2X7R [79].



**Figure 11.** Chemical structure of 1,4-naphthoquinone thioglucosides and their tetracyclic conjugates with inhibitory activity towards P2X7 receptor.


In conclusion, these reports reinforce the elevated potential of naphthoquinone derivatives as neuroprotectors with varied mechanism of action. In the next section, we will discuss the recent literature on synthetic strategies and biological evaluation naphthoquinone compounds targeting PD.

### 3.2. Naphthoquinones with activity against PD

At the end of the 20th century, epidemiological evidence emerged that smokers had a lower incidence of PD than non-smokers [80,81]. Then, this profile was associated with a lower activity of the enzyme monoamine oxidase (MAO) in the brain of smokers [82]. MAO is an important enzyme for metabolizing amines in the body via oxidative deamination [83]. MAO has two isoforms, MAO-A and MAO-B. The MAO-B isoform is predominant in the brain, where it metabolizes dopamine into 3,4-dihydroxyphenylacetic acid and homovanillic acid [84]. This enzyme is involved in the metabolism of neurotransmitters such as norepinephrine, epinephrine, dopamine and serotonin. The loss of dopamine, norepinephrine and serotonin are the basis for the degenerative process that leads to PD and there is an association between elevated levels of MAO-B and this neurodegenerative disease [85]. MAO-B inhibition prevents the breakdown of dopamine molecules, which makes them more available to act on their receptors. Its inhibition also decreases the formation of free radicals

from dopamine oxidation [86]. Thus, supplementation with L-DOPA and MAO inhibitors is one of the main ways to treat the disease [83].


With the desire to determine which substances were responsible for the MAO inhibitory activity, a screening of the substances present in burley tobacco leaves was carried out [87]. In this screening, naphthoquinone 2,3,6-trimethyl-1,4-naphthoquinone (**28**) was identified as the main substance responsible for MAO inhibition. Its inhibition was observed as reversible competitive, both against MAO-A and MAO-B, with a  $K_i$  of 3  $\mu\text{M}$  and 6  $\mu\text{M}$  respectively. Naphthoquinone **28** was also synthesized for structure confirmation and to provide greater mass for biological assays. The synthesis of this naphthoquinone started with the oxidation of 2,3-dimethylphenol (**29**) with sodium nitrite forming an oxime, followed by an oxidation with copper oxide I producing 2,3-dimethylbenzoquinone (**30**) [88]. Then, **30** reacted by cycloaddition with isoprene forming the Diels-Alder adduct **31**, which was not isolated. Finally, **31** underwent dehydrogenation forming **28** (Scheme 1).



**Scheme 1.** Synthetic route for production of **28**.

Then **28** was tested in C57BL/6 mice after neurotoxic MPTP treatment [89]. MAO is also the enzyme responsible for converting 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) to the neurotoxic metabolite, 1-methyl-4-phenylpyridinium ion (MPP<sup>+</sup>) [90]. Mice treated with **28** and MPTP exhibited a 50% increase in dopamine concentration in the striatum when compared to mice treated with MPTP alone. Thus, under experimental conditions, **28** exerted a neuroprotective effect on mice.

These works concluded that the naphthoquinone nucleus could be an important pharmacophore for MAO inhibition, which led to the beginning of studies investigating the molecular details of the mechanism of inhibition of 1,4-naphthoquinones. It was discovered that 1,4-naphthoquinone (**32**) and menadione (**33**) (Figure 12) showed strong inhibition of MAO-B, with  $K_i$  of 1.4  $\mu\text{M}$  and 0.4  $\mu\text{M}$  respectively [91]. Regarding MAO-A, a  $K_i$  of 7.7  $\mu\text{M}$  and 26  $\mu\text{M}$  was found for **32** and **33**, respectively, showing a lower susceptibility of MAO-A in relation to MAO-B with these substances. Both naphthoquinones showed characteristics of reversible and competitive inhibitors for MAO-B. Docking studies suggest that phenyl group interactions at amino acid residues Tyr407 and Tyr444 for MAO-A or Tyr398 and Tyr435 for MAO-B are important for the interaction of the naphthoquinone core and the active site of the enzyme.




**Figure 12.** Structure of naphthoquinones with MAO inhibitory activity initially studied.

Continuing with the exploration of this activity of naphthoquinones and contributing to expanding knowledge about structure activity relationship, the inhibitory capacity of fourteen natural and synthetic naphthoquinones was studied [92]. In this study, naphthoquinones were grouped according to the degree of hydroxylation of the aromatic ring into 1,4-naphthoquinone analogues (**32-39**), juglone analogues (**6, 40**), naphthazarine analogues (**41, 7**), in addition to lapachol

(5) and *nor*-lapachol (42) (Figure 13). Naphthoquinones naphthazarine (41) ( $IC_{50} = 0.860 \pm 0.241 \mu M$ ) and shikonin (7) ( $IC_{50} = 1.50 \pm 0.302 \mu M$ ) showed the greatest capacity to inhibit MAO-B and MAO-A respectively. Similar results were also found previously when identifying shikonin (7) as one of the main substances responsible for the MAO inhibitory effect in *Lithospermum erythrorhizon* extract [93]. An interesting aspect observed was that all naphthoquinones with a hydroxyl in the aromatic ring showed a higher MAO-B inhibitory activity than their non-hydroxylated analogues, as is the example of the comparison of 32 ( $IC_{50} = 8.40 \pm 1.20 \mu M$ ) with 6 ( $IC_{50} = 4.36 \pm 0.180 \mu M$ ) and 33 ( $IC_{50} = 3.02 \pm 0.638 \mu M$ ) with 40 ( $IC_{50} = 1.09 \pm 0.063 \mu M$ ). However, the presence of a C2 hydroxyl does not help this inhibitory activity, since, while 32 is an MAO inhibitor, its C2 hydroxylated analogue, 34, showed no activity. Interestingly, 34 and its tautomers extensively perform hydrogen bonding and  $\pi$  stacking according to docking experiments and therefore an explanation for this observation cannot yet be realized. It was also possible to note that the presence of a methyl in C2 is associated with an inversion of selectivity. The C2-methylated analogs tended to be more selective for MAO-B, while the demethylated ones for MAO-A. 2,3-dichloro-1,4-naphthoquinone (36) was also highlighted for its potency against MAO-A ( $IC_{50} = 2.00 \pm 0.878 \mu M$ ) and MAO-B ( $IC_{50} = 2.61 \pm 0.616 \mu M$ ). Naphthoquinones 5 and 42 were not identified as MAO inhibitors.

#### MAO-B inhibition activity comparison



**Figure 13.** Structure of naphthoquinones studied by Mostert *et al* (2016).

$\alpha$ -Synuclein is a 14 kDa cytosolic protein composed of 140 amino acids and divided into an N-terminal amphipathic  $\alpha$ -helix, a non-amyloid component and a C-terminal acidic tail [94]. This protein presents its aggregation and accumulation as one of the hallmarks of PD [95]. This accumulation produces proteinaceous inclusions known as Lewy bodies or Lewy neurites [96].  $\alpha$ -Synuclein aggregates cause damage and destabilize several processes within the neuronal cell, such as interrupting the formation of the synaptic vesicle release complex, its mobility and dopamine release [97]. Regarding its activity in mitochondria,  $\alpha$ -synuclein can inhibit complex I of the respiratory chain, increase the formation of reactive oxygen species (ROS), interfere with fusion, fission and potential of the mitochondrial membrane [96,98]. Accumulation of  $\alpha$ -synuclein in the endoplasmic reticulum (ER), an essential organelle for protein metabolism, interferes with protein folding, causing stress in the ER [99]. With the broad spectrum of important activities in neurodegenerative diseases of naphthoquinones, the activity of vitamin K analogues and 1,4-

naphthoquinone itself (**32**) (Figure 12) on  $\alpha$ -synuclein fibrillation was investigated. The results showed that these naphthoquinones, in addition to inhibiting MAO, also have the potential to inhibit  $\alpha$ -synuclein aggregation [100]. All naphthoquinones demonstrated dose-dependent inhibition of fibril formation by  $\alpha$ -synuclein with  $EC_{50}$  between 15 and 30  $\mu$ M. Even though many substances show inhibition of  $\alpha$ -synuclein fibrillation through non-specific hydrophobic interactions, the largest aliphatic chain of phytoquinone (**13**) ( $EC_{50}$  = 29  $\mu$ M) and menaquinone (**14**) ( $EC_{50}$  = 30  $\mu$ M) (Figure 7) in relation to menadione (**33**) ( $EC_{50}$  = 18  $\mu$ M) and **32** ( $EC_{50}$  = 15  $\mu$ M) (Figure 12), did not translate into a better inhibitory activity. Naphthoquinones **32** and **33** were able to destabilize preformed fibrils, which may be associated with their lower molecular volume allowing greater diffusion within the fibrillar structure.

The tau protein has an important activity in the growth and development of neurons. The aggregation of tau proteins causes destabilization of microtubules, which leads to severe consequences for the neuron, such as, for example, the disturbance of axonal transport and neurite outgrowth and leaving the DNA prone to suffer damage. These consequences can lead to cell death [101]. The deposition of tau protein aggregates is an important point in several neurodegenerative diseases, such as PD. The naphthoquinone shikonin (**7**) (Figure 2) was able to decrease the rate and extent of tau aggregation [102]. This substance showed an  $IC_{50}$  of  $1.2 \pm 0.06$  and  $1.0 \pm 0.3$   $\mu$ M for oligomerization and aggregation of tau 4R2N, respectively, in heparin-containing medium. The  $IC_{50}$  for inhibition of RNA- and arachidonic acid-induced tau aggregation were  $1.0 \pm 0.2$  and  $1.5 \pm 0.1$   $\mu$ M respectively. Therefore, **7** was able to inhibit tau aggregation in heparin, RNA and arachidonic acid-containing medium. Naphthoquinone **7** not only decreased tau oligomerization, but also decreased the average size of tau oligomers. Added to this, with an  $DC_{50}$  of  $6.3 \pm 0.4$   $\mu$ M, **7** was able to disaggregate preformed tau filaments. Tau oligomers were added to SH-SY5Y neuroblastoma cell culture medium pretreated with **7**. The viability of SH-SY5Y cells treated with 8  $\mu$ M of tau oligomers rose from  $34 \pm 2$  to  $71 \pm 6$  % when pretreated with 250 nM of **7**. The  $IC_{50}$  of **7** was significantly lower than other known inhibitors. Planar rings of **7** containing delocalized  $\pi$  electrons can interact with aromatic or polar side chains of tau, which can modify the protein's conformation. This conformational change would make tau less likely to form aggregates.

**7** is associated with several other activities that promote neuroprotective effects in PD models. This naphthoquinone is associated with a decrease in ROS levels attributed to an increase in the activity of the antioxidant enzyme glutathione peroxidase 1 (GPX-1) [103]. **7** at a concentration of 10  $\mu$ M induced increased expression of GPx-1 in PC12 cells, increasing cell survival by 70% against 6-hydroxydopamine (6-OHDA) toxicity [104]. **7** also increased the expression of the glutathione-independent antioxidant enzyme, superoxide dismutase 2 (SOD-2), increased the expression of Bcl-2 and decreased expression of Bax, which reduced the rate of nuclear morphology change that normally occurs in apoptosis. This was the first time that **7** was reported to protect dopaminergic neurons.

Pin1 is a peptidyl-prolyl isomerase which has a profound impact on the regulation of cell growth and proliferation, stress response, immune function, neuronal differentiation, cell motility and apoptosis [105]. Pin1 has already been found in Lewy bodies in PD and also facilitates the formation of  $\alpha$ -synuclein inclusions in a cellular model of  $\alpha$ -synuclein aggregation. It has also been found that Pin1 is expressed in 50-60% of Lewy bodies in PD patients [106]. Thus, Pin1 is a degenerative factor for dopaminergic neurons. Juglone (**6**) (Figure 2) is a naphthoquinone with recognized Pin1 inhibitory activity [107]. Naphthoquinone **13** (Figure 7) irreversibly inhibits the enzymatic activity of Pin1 by modifying the thiol groups of cysteine residues. Previously it had already been shown that **6** at a concentration of 5.7  $\mu$ M *in vitro* blocks the activity of Pin1 without impacting the activity of other isomerases [108]. Treatment with **6** attenuates Pin1 expression, protects the nigrostriatal axis and improves hypolocomotion in a preclinical model of rats treated with MPTP at a dose of 3 mg/kg, which is non-toxic [109]. In this work, it was found that **6** at 1  $\mu$ M concentration attenuates MPP<sup>+</sup>-induced Pin1 expression in MN9D cells. Substance **6** also restored dopamine reuptake in primary mesencephalic neurons and their neurites from MPP<sup>+</sup> toxicity. It was also found that **6** restored the behavioral activities as well as the levels of dopamine and its metabolites in the striatum of rats




treated with MPTP. Interestingly, even though juglone has been a recognized Pin1 inhibitor for many years, there is a large gap in the literature as to how structural modifications could affect its inhibitory activity.

Naphthazarine (**41**) (Figure 13) at subtoxic doses was investigated for its ability to protect the brain through activation of adaptive stress response pathways in an animal model of MPTP-induced PD [110]. In this work, **41** in 1 mg/kg significantly prevented the decrease in motor function caused by MPTP. Pretreatment with **41** effectively reduced the loss of dopaminergic neurons in the substantia nigra and loss of nerve endings in the striatum. It was identified that **41** also decreased the activation of astrocytes and microglia in the MPTP model by observing a decrease in GFAP and Iba-1 markers.

It has also been found that  $\beta$ -lapachone (**43**) (Figure 14) promotes a neuroprotective effect in rats in a PD model with the administration of MPTP [111]. Naphthoquinone **43** improved impaired movement in MPTP-administered rats through recovery of dopaminergic neurons in the substantia nigra and striatum and expression of Bcl-2. In this study, it was found that **43** increased the expression of the enzyme heme oxygenase-1 (HO-1), which has antioxidant activity, and of the p-AMPK and Nrf2 signaling molecules in astrocytes of MPTP administered rats. In a study of co-treatment of **43** with L-DOPA, **43** alleviated dyskinesia induced by chronic L-DOPA treatment [112]. Dyskinesia is a serious motor problem that occurs in about half of patients when L-DOPA is given to PD patients for a long period of time. Substance **43** decreased the abnormal involuntary movements. Co-treatment substantially decreased astrocyte activation in the striatum and substantia nigra in a unilateral 6-OHDA model.


It was verified that the treatment of rats with 20 mg/kg of rinacanthine C (**44**) (Figure 14) significantly decreased the cataleptic effect of haloperidol when compared to control groups [113]. Treatment with **44** also significantly increased levels of dopamine, norepinephrine and serotonin when compared to the diseased control group. The treatment also increased the rats' exploratory behavior, which had declined due to the effects of the disease. Improvement of these non-motor symptoms is associated with restoration of serotonin and dopamine levels in the rat brains. Treatment with **44** normalized brain antioxidant losses at all doses tested. No significant side effects were found in other organs of the body, which could mean it is a safer profile that should be investigated.



**Figure 14.** Other natural naphthoquinones with neuroprotective activity for PD.


Considering previous works that showed that naphthoquinones have the capacity to inhibit MAO, the conjugation of these substances with substrate groups for the enzyme could help in the inhibitory activity. Therefore, spermidine-conjugated naphthoquinones (**45**) derived from lapachol (**5**), lawsone (**34**) and *nor*-lapachol (**42**) were synthesized (Scheme 2) [114]. For the production of these molecules, initially the hydroxyl of the quinonoid ring was methylated. The methylated derivatives of **5** and **42**, **46a** and **46b** respectively, were produced by reacting these naphthoquinones with dimethylsulfate in acetone and potassium carbonate [115]. The methylated derivative of **34**, **46c** was synthesized from the sodium salt of 1,2-naphthoquinone-4-sulfonic acid (**47**) [116]. Next, the methoxy group underwent a nucleophilic displacement by the Boc-protected spermidine, producing **48**. The naphthoquinones **45** were then obtained with the removal of the protection group (Scheme 2). Naphthoquinones **45** showed  $K_i$  between 12 and 63  $\mu$ M with low selectivity between MAO-A and MAO-B and a kinetic study compatible with a competitive mechanism. Although **34** had the lowest  $K_i$  than the other naphthoquinones, all other derivatives with substituents on the quinonoid ring were

less potent than 1,4-naphthoquinone (32). All these naphthoquinones showed characteristics of reversible inhibitors. The greater activity of **42** compared to **5** may be associated with its closer proximity to the FAD portion as well as the orientation of the quinonoid ring in the aromatic stacking of Tyr435/Tyr398 residues. Lapachol (**5**) showed no inhibitory activity, while *nor*-lapachol (**34**) showed low activity against MAO-A, which may be due to the absence of hydrogen bonds between naphthoquinone and the catalytic site, as previously reported [89]. In the same reasoning, **45c** was the naphthoquinone-spermidine with the highest potency for MAO-B inhibition, which may be associated with the greater number of hydrogen bonds between naphthoquinone and the FAD moiety at the catalytic site.




**Scheme 2.** Synthesis of spermidine-conjugated naphthoquinones **45**; a) (CH<sub>3</sub>O)<sub>2</sub>SO<sub>2</sub>, acetone, K<sub>2</sub>CO<sub>3</sub>; b) MeOH, H<sub>2</sub>SO<sub>4</sub>, D; c) MeOH, 90 °C, Boc-spermidine; d) TFA/CH<sub>2</sub>Cl<sub>2</sub>, MeOH, 0 °C.

There are works in the literature that demonstrate the ability of hybrids of naphthoquinones and tryptophan (**49**) (Scheme 3) to inhibit the aggregation of the amyloid-beta peptide [117,118]. These naphthoquinones were also tested as potential inhibitors of  $\alpha$ -synuclein protein aggregation [119]. In the same context, dopamine can form adducts with  $\alpha$ -synuclein in vitro, inhibiting the formation of fibrils. This dopamine activity, along with the already known ability of naphthoquinones to inhibit  $\alpha$ -synuclein aggregation, stimulated the synthesis of dopamine and naphthoquinone hybrids (**50**) (Scheme 3) [120]. These naphthoquinones were produced by means of a Michael addition to the quinonoid ring by tryptophan or dopamine followed by replacement of the hydrogen or chlorine atom to produce **49a** and **49b** or **50a** and **50b** respectively (Scheme 3) [121]. The two naphthoquinones **49** demonstrated dose-dependent inhibition of fibril formation by  $\alpha$ -synuclein. Analog **49a** showed better inhibition capacity, which may be associated with lower steric hindrance. The synthesis of hybrids **50** occurred similarly to the production of hybrids **49**, however the reaction conditions and yields were not mentioned. Maximum inhibition was observed with a hybrid concentration 5 times higher than  $\alpha$ -synuclein by the ThT assay, being 88% for **50a** and 76% for **50b**. Analysis of data from experiments carried out with **32** and dopamine demonstrate a superior inhibitory activity of **50a** and that the presence of the naphthoquinone structure and dopamine in the same molecule has a synergistic effect.




**Scheme 3.** Synthesis of hybrids of naphthoquinones with tryptophan (**49**) and dopamine (**50**); a) L-tryptophan, EtOH/H<sub>2</sub>O, 23 °C, 24 h; b) L-tryptophan, KOH, MeOH/H<sub>2</sub>O, 24 h.

With the hypothesis that the conjugation of these naphthoquinones with mannitol could improve the ability of these molecules to cross the blood-brain barrier, analogues of naphthoquinone **49a** conjugated with mannitol were synthesized [122]. These naphthoquinones were synthesized by click reactions (**51**) or PEG connector (**52**). These molecules have been shown to reduce the kinetics and extent of  $\alpha$ -synuclein aggregation. Furthermore, the conjugates are more potent than **49a** or its mixture with mannitol. The synthetic route to produce naphthoquinone **51** starts from **49a** with an amide formation reaction with propargylamine, producing **52**. In the terminal alkyne group of **52** there is a click reaction with the azide **53** to form the triazole (**54**) and ends with hydrolysis of the acetate groups (Scheme 4).




**Scheme 4.** Synthesis of naphthoquinone conjugated with mannitol **51** by click reaction; a) propargylamine, EDCI, HOBt, DIEA, THF/DCM; b) **53**, sodium ascorbate, CuSO<sub>4</sub>; c) CH<sub>3</sub>ONa/CH<sub>3</sub>OH, pH = 8,5.

Naphthoquinone **52b** was produced by amide group formation from the PEG connector **55** with the carboxyl of **49a** (Scheme 5) and showed the best inhibitory activity. It is speculated that the longer PEG chain in **52b** may confer greater flexibility and, consequently, a more efficient interaction with  $\alpha$ -synuclein.




**Scheme 5.** Synthesis of naphthoquinones conjugated with mannitol (**52**) by PEG connector; a) **55**, DIEA, HOBT, EDC.Cl, DMSO, r.t., 12h.

Considering the already explored protective effect of naphthazarine and other structurally analogous naphthoquinones, Aminin and collaborators evaluated the neuroprotective activity in Neuro-2a neuroblastoma cells in a model of Parkinson's of numerous natural naphthoquinones and their synthetic derivative. The tested naphthoquinones were divided into four groups: naphthazarin (**41**) and its analogues; derivatives with acetylated *O*-glucoside groups (**56**); derivatives with acetylated *S*-glucoside groups (**57**); derivatives with deacetylated *S*-glucoside groups (**58**). The analogues of **41** were obtained by various methods. Naphthopurpurine (**59**), 2-hydroxy-3-methylnaphthazarine (**60**) and 2-hydroxy-3-ethylnaphthazarine (**61**) were obtained by a reductive dechlorination of **62** using iron in acetic acid followed by air oxidation [123]. One way to obtain the dichlorinated derivatives **62** is through the Friedel-Crafts cycloaddition between a hydroquinone and dichloromaleic anhydride (**63**), as is the case of the synthesis of **62a**, which uses hydroquinone **64** [124]. 2,3-Dihydroxynaphthazarine (**65**), 2,3-dihydroxy-6-methylnaphthazarine (**66**), 2,3-dihydroxy-6-ethylnaphthazarine (**67**) and spinochrome D (**68**) were synthesized using an efficient synthetic route starting from **64**. Naphthoquinone **64** reacted with sodium nitrite forming intermediate **69**, followed by a reduction producing 3-amino-2-hydroxynaphthazarines (**70**) [125]. Naphthoquinones **70** then undergo dimethylsulfoxide-mediated oxidation [126,127]. 2-hydroxy-6,7-dimethylnaphthazarine (**71**) was prepared by oxidation **64a** with MnO<sub>2</sub> in concentrated acid, while 6,7-dichloro-2-hydroxynaphthazarine (**72**) was produced by nucleophilic substitution of chlorine atoms by groups methyl, but the reaction conditions were not informed [128]. Finally, 2-hydroxy-7-methoxynaphthazarine (**73**) and 2-hydroxy-6,7-dimethoxynaphthazarine (**74**) were synthesized by a retro aldol disintegration from their 2-hydroxy-3-(1-hydroxy-ethyl) corresponding naphthazarins (**75**) (Scheme 6) [129].



**Scheme 6.** Synthesis of naphthazarine (**41**) analogues. a)  $\text{AlCl}_3$ ,  $\text{NaCl}$ ,  $190^\circ\text{C}$ ; b) For **62b-d**: 1.  $\text{Fe}$ ,  $\text{HOAc}$ ; 2.  $[\text{O}]$ ; c) For **70a-c**:  $\text{DMSO}$ ,  $\text{HCO}_2\text{H}$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{O}$ , reflux; d) For **70d**: 1.  $\text{DMSO}$ ,  $\text{HCO}_2\text{H}$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{O}$ , reflux; 2.  $\text{MeSO}_3\text{H}$ ,  $\text{HCO}_2\text{H}$ , reflux; e) For **62a,e,f**: 1.  $\text{NaNO}_2$ , acetone– $\text{MeOH}$ , reflux, 2 h; 2.  $\text{HCl}$  c.; f) For **62b**: 1. isoamyl alcohol,  $\text{H}_2\text{SO}_4$ , reflux; 2.  $\text{NaNO}_2$ , acetone– $\text{MeOH}$ , reflux, 2 h; 3.  $\text{HCl}$  c.; g) For **69a-b**:  $\text{Na}_2\text{S}_2\text{O}_4$ ,  $\text{H}_2\text{O}$ , 2 h; h) For **69d**:  $\text{Na}_2\text{S}$ ,  $\text{H}_2\text{O}$ , r.t.; i) For **62a**:  $\text{MnO}_2$ ,  $\text{H}_2\text{SO}_4$ , r.t., 2h; j)  $\text{NaOH}/\text{H}_2\text{O}$ , reflux.


These naphthoquinones were converted into their derivatives with acetylated *O*-glucoside groups (**56**) by means of a condensation with 3,4,6-tri-*O*-acetyl- $\alpha$ -d-glucopyranose 1,2-(tert-butoxy orthoacetate) (**76**) [130,131]. The amount used was equimolar to the amount of  $\beta$ -hydroxy groups. Thus, the glycosidation of 2-hydroxy-naphthazarines (**59-61**, **71**), 2,3-dihydroxy-naphthazarines (**65-67**) and echinochrome A (**77**) formed the derivatives mono- (**56a-d**), bis- (**56e-g**) and acetylated tris-*O*-glycosides (**56h**) respectively (Scheme 7).



**Scheme 7.** Synthesis of acetylated triglucosidyl derivatives (**56**); a) **76**, chlorobenzene, reflux.

It was found that deacetylation of derivatives **56** produced very unstable molecules. This led to the need for the synthesis of thioglycosylated derivatives with the moiety of **41** [131]. The 2-hydroxynaphthazarines (**59**, **71-74**) were used for the synthesis of derivatives with acetylated S-glucoside groups (**35**). Tetra-*O*-acetyl-1-mercapto-D-glucose (**78**) was used as a thiol source. The reaction starts with a Knoevenagel condensation between the naphthazarine analogue and paraformaldehyde, forming methide *o*-quinone. Next, there is a nucleophilic attack of **77** forming conjugates **57**. Another possibility is that the *o*-quinone methide reacts with another molecule of naphthazarine analogue, forming dimers. Two produced dimers were included in the biological tests, **79a-b**, which are spinochrome D (**68**) dimers linked by a methylene or ethylidene bridge. Conjugates **57** were then deacetylated in acid solution, forming derivatives with deacetylated S-glucoside groups (**58**), or they were methylated at the 2-position hydroxyl with a diazomethane solution yielding **57f-j**. These also underwent subsequent deacetylation, yielding **58f-j** (Scheme 8).





**Scheme 8.** Synthesis of thioglucosidyl derivatives **57** and **58**; a) **78**, paraformaldehyde, HCOOH, acetone, reflux; b) HCl/MeOH; c) 1.  $\text{CH}_2\text{N}_2$ ; 2. HCl/MeOH; b) HCl/MeOH.

Most naphthoquinones tested were able to protect Neuro-2a cells against the neurodegenerative effects of neurotoxins. The naphthoquinone with the greatest neuroprotective effect in the paraquat model in the MTT test was **74**, which at a concentration of 0.1  $\mu\text{M}$  increased the number of viable cells by 45.7% compared to the control. The neuroprotective effect was greater in the paraquat model than in the 6-OHDA model. In this model, spinochrome D (**68**) had the best result with an increase of only 22.3% of viable cells compared to the control. Naphthoquinones **65**, **66**, **74**, **79b** and **58h** had the best results in improving cell survival in MTT, non-specific esterase and PI fluorescent dye tests, both in paraquat and 6-OHDA models. Naphthoquinone **65** was also the substance with the greatest ability to decrease ROS production in the paraquat and 6-OHDA models, decreasing production by 30.2% (0.1  $\mu\text{M}$ ) and 31.6% (1.0  $\mu\text{M}$ ) respectively. Incubation with paraquat resulted in a charge shift in the mitochondrial membrane potential (MMP), of which **65** and **66**, at the 0.01  $\mu\text{M}$  concentration, were able to increase by 8.9 % and 21.2 %, respectively. Of all the tests, **74** was the most active in terms of its neuroprotective capacity, while naphthazarine (**41**) was the most inactive. The QSAR study suggests that greater activity of **74** is related to a greater number of hydrogen bond acceptor sites and a greater hydrophobic surface. Naphthoquinone **41**, on the other hand, showed a greater number of hydrogen bond donor sites and a significantly smaller hydrophobic region. These naphthoquinones demonstrated protection of non-specific esterase activity from the inhibitory activity of neurotoxins, normalized the cell cycle and inhibited the lytic activity of membranes. These activities are related to the ability of these naphthoquinones to decrease ROS production. For naphthoquinone **55**  $\beta$ -hydroxyl groups are essential for the antioxidant effect, which explains the lower neuroprotective activity in methoxylated derivatives [132]. Glycosidation with O- and S-glycoside groups, acetylated or not,

blocked  $\beta$ -hydroxyl groups and were responsible for the low neuroprotective activity of these derivatives.

The acetylated triglucosidic derivative of echinochrome A (**56h**) was used to activate the transcription factor HSF1 and increase the expression of the molecular chaperone HSP70 [133]. This naphthoquinone activity is already known, and it was used in this work to provide more evidence that the elevation of HSP70 levels is a possible therapeutic target for the treatment of PD. This activity reverses the neurodegeneration process, as HSP70 has the ability to prevent  $\alpha$ -synuclein aggregation and microglia activation. Naphthoquinone **56h** was also indicated as therapeutically effective, since in an animal model with lactacystin-induced PD it reduced the loss of dopaminergic neurons by two times when compared to the control, protected the activity of neurons in the substantia nigra pars compacta, eliminated motor dysfunction characteristic of PD, reduced microglial activation and decreased neuroinflammation. When tested in aged rats in a lactacystin-induced preclinical PD model, the **56h** administration led to a 130% increase in HSP70 compared to the control, which decreased neurodegeneration by 2.4 times [134].

#### 4. Conclusions

In this review, we describe the epidemiology of Parkinson's Disease worldwide, highlighting the main risk factors, clinical manifestations and also the pathophysiological mechanisms of the disease. As oxidative stress is a pathological manifestation of traumatic brain injuries such as PD, the use of antioxidant drugs, such as those belonging to the 1,4-naphthoquinone class, may be an effective alternative in the treatment of this disease. For this reason, we report many examples of 1,4-naphthoquinones that have shown neuroprotective effect. In addition, we also highlight the main biological targets related to the neurological disorders that cause PD and the naphthoquinones that showed promising results in assays directed towards these targets. The synthetic routes chosen for the production of new naphthoquinone derivatives were also highlighted in this review. Thus, in reason of to the various examples and expressive biological results presented, 1,4-naphthoquinones can be considered a promising class the treatment for PD.

#### References

1. a) Parkinson J. An Essay on the Shaking Palsy. London: Sherwood, Neely and Jones, **1817**. b) Kempster, P.A.; Hurwitz, B.; Lees, A.J. A new look at James Parkinson's Essay on the Shaking Palsy. *Neurology* **2007**, *69*, 482–485. doi:10.1212/01.wnl.0000266639.50620.d1 c) Goetz C.G. The history of Parkinson's disease: early clinical descriptions and neurological therapies. *Cold Spring Harb. Perspective Med.* **2011**, *1*, a008862. doi: 10.1101/cshperspec.a008862
2. a) GBD 2016 Neurology Collaborators. Global, regional, and national burden of neurological disorders, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* **2019**, *18*, 459–480. doi:10.1016/S1474-4422(18)30499-X b) Dorsey, E.R.; Sherer, T.; Okun, M.S.; Bloem, B.R. The Emerging Evidence of the Parkinson Pandemic. *J Parkinsons Dis.* **2018**, *8*, S3-S8. doi:10.3233/JPD-181474. c) Deuschl, G.; Beghi, E.; Fazekas, F.; Varga, T.; Christoforidi, K.A.; Sipido, E.; Bassetti, C.L.; Vos, T.; Feigin, V.L. The burden of neurological diseases in Europe: an analysis for the Global Burden of Disease Study 2017. *Lancet Public Health* **2020**, *5*, 551–567. doi:10.1016/S2468-2667(20)30190-0.
3. Pringsheim, T.; Jette, N.; Frolkis, A.; Steeves, T.D. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Move Discord.* **2014**, *13*, 1583–1590. doi:10.1002/mds.25945.
4. a) Bjornestad, A.; Forsaa, E.B.; Pedersen, K.F.; Tysnes, O.B.; Larsen, J.P.; Alves, G. Risk and course of motor complications in a population-based incident Parkinson's disease cohort. *Park. Relat. Disord.* **2016**, *22*, 48–53. doi:10.1016/j.parkreldis.2015.11.007. b) Picillo, M.; Palladino, R.; Moccia, M.; Erro, R.; Amboni, M.; Vitale, C.; Barone, P.; Pellecchia, M.T. Gender and non motor fluctuations in Parkinson's disease: A prospective study. *Park. Relat. Disord.* **2016**, *27*, 89–92. doi:10.1016/j.parkreldis.2016.04.001.
5. Nicoletti, A.; Vasta, R.; Mostile, G.; Nicoletti, G.; Arabia, G.; Iliceto, G.; Lamberti, P.; Marconi R.; Morgante, L.; Barone P.; Quattrone, A.; Zappia, M. Gender effect on non-motor symptoms in Parkinson's disease: are men more at risk? *Park. Relat. Disord.* **2017**, *35*, 69–74. doi:10.1016/j.parkreldis.2016.12.008.
6. a) Fullard, M.E.; Thibault, D.P.; Hill, A.; Fox, J.; Bhatti, D.E.; Burack, M.A.; Dahodwala, N.; Haberkfeld, E.; Kern, D.S.; Klepitskava, O.S.; Urrea-Mendoza, E.; Myers, P.; Nutt, J.; Rafferty, M.R.; Schwab, J.M.; Shulman, L.M.; Willis, A.W. Parkinson Study Group Healthcare Outcomes and Disparities Working Group. Utilization of rehabilitation therapy services in Parkinson disease in the United States. *Neurology*

- 2017, 89, 1162-1169. doi:10.1212/WNL.0000000000004355. b) Fullard, M.E.; Thibault, D.P.; Todaro, V.; Foster, S.; Katz, L.; Morgan, R.; Kern, D.S.; Schwalb, J.M.; Urrea-Mendoza, E.; Dahodwala, N.; Shulman, L.; Willis, A.W. Sex disparities in health and health care utilization after Parkinson diagnosis: Rethinking PD associated disability. *Park. Relat. Disord.* **2018**, *48*, 45-50. doi:10.1016/j.parkreldis.2017.12.012.
7. Breckenridge, C.B.; Berry, C.; Chang, E.T.; Sielken, R.L.; Mandel, J.S. Association between Parkinson's Disease and Cigarette Smoking, Rural Living, Well-Water Consumption, Farming and Pesticide Use: Systematic Review and Meta-Analysis. *PLoS One* **2016**, *11*, e0151841. doi:10.1371/journal.pone.0151841.
  8. a) Langston, J.W.; Ballard, P.; Tetrud, J.W.; Irwin, I. Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* **1983**, *219*, 979-980. doi:10.1126/science.6823561. b) Angibaud, G.; Gaultier, C.; Rascol, O. Atypical parkinsonism and Annonaceae consumption in New Caledonia. *Mov. Disord.* **2004**, *19*, 603-604. doi:10.1002/mds.20104. c) Höglinger, G.U.; Michel, P.P.; Champy, P.; Feger, J.; Hirsch, E.C.; Ruberg, M.; Lannuzel, A. Experimental evidence for a toxic etiology of tropical parkinsonism. *Mov. Disord.* **2005**, *20*, 118-119. doi:10.1002/mds.20300.
  9. Mittal, S.; Bjørnevik, K.; Im, D.S.; Flierl, A.; Dong, X.; Locascio, J.J.; Abo, K.M.; Long, E.; Jin, M.; Xu, B.; Xiang, Y.K.; Rochet, J.C.; Engeland, A.; Rizzu, P.; Heutink, P.; Bartels, T.; Selkoe, D.J.; Caldarone, B.J.; Glicksman, M.A.; Khurana, V.; Schule, B.; Park, D.S.; Riise, T.; Scherzer, C.R.  $\beta$ 2-Adrenoreceptor is a regulator of the  $\alpha$ -synuclein gene driving risk of Parkinson's disease. *Science*. **2017**, *357*, 891-898. doi:10.1126/science.aaf3934.
  10. Palacios, N.; Gao X.; McCullough, M.L.; Schwarzschild, M.A.; Shah, R.; Gapstur S.; Ascherio, A. Caffeine and risk of Parkinson's disease in a large cohort of men and women. *Mov. Disord.* **2012**, *27*, 1276-1282. doi:10.1002/mds.25076.
  11. Bai, S.; Song, Y.; Huang, X.; Peng, L.; Jia, J.; Liu, Y.; Lu, H.; Statin Use and the Risk of Parkinson's Disease: An Updated Meta-Analysis. *PLoS One* **2016**, *11*, e0152564. doi:10.1371/journal.pone.0152564.
  12. Gudala, K.; Kanukula, R.; Bansal, D. Reduced Risk of Parkinson's Disease in Users of Calcium Channel Blockers: A Meta-Analysis. *Int. J. Chronic Dis.* **2015**, *2015*, 697404. doi:10.1155/2015/697404.
  13. a) Gagne, J.J.; Power, M.C. Anti-inflammatory drugs and risk of Parkinson disease: a meta-analysis. *Neurology*. **2010**, *74*, 995-1002. doi:10.1212/WNL.0b013e3181d5a4a3. b) Becker, C.; Jick, S.S.; Meier, C.R. NSAID use and risk of Parkinson disease: a population-based case-control study. *Eur. J. Neurol.* **2011**, *18*, 1336-1342. doi:10.1111/j.1468-1331.2011.03399.x.
  14. Gasser, T. Genetics of Parkinson's disease. *Curr. Opin. Neurol.* **2005**, *18*, 363-369. doi:10.1097/01.wco.0000170951.08924.3d.
  15. Wakabayashi, K.; Tanji, K.; Odagiri, S.; Miki, Y.; Mori, F.; Takahashi, H. The Lewy body in Parkinson's disease and related neurodegenerative disorders. *Mol. Neurobiol.* **2013**, *47*, 495-508. doi:10.1007/s12035-012-8280-y.
  16. Hsu, L.J.; Sagara, Y.; Arroyo, A.; Rockenstein, E.; Sisk, A.; Mallory, M.; Wong, J.; Takenouchi, T.; Hashimoto, M.; Masliah, E.  $\alpha$ -Synuclein promotes mitochondrial deficit and oxidative stress. *Am. J. Clin. Pathol.* **2000**, *157*, 401-410. doi:10.1016/s0002-9440(10)64553-1.
  17. Snyder, H.; Mensah, K.; Theisler, C.; Lee, J.; Matouschek, A.; Wolozin, B. Aggregated and monomeric  $\alpha$ -synuclein bind to the S6' proteasomal protein and inhibit proteasomal function. *J. Biol. Chem.* **2003**, *278*, 11753-11759. doi:10.1074/jbc.M208641200.
  18. Danzer, K.M.; Haasen, D.; Karow, A.R.; Moussaud, S.; Habeck, M.; Giese, A.; Kretschmar, H.; Hengerer, B.; Kostka, M. Different species of alpha-synuclein oligomers induce calcium influx and seeding. *J. Neurosci.* **2007**, *27*, 9220-9232. doi:10.1523/JNEUROSCI.2617-07.2007.
  19. Alim, M.A.; Ma, Q.L.; Takeda, K.; Aizawa, T.; Matsubara, M.; Nakamura, M.; Asada, A.; Saito, T.; Kaji, H.; Yoshii, M.; Hisanaga, S.; Ueda, K. Demonstration of a role for  $\alpha$ -synuclein as a functional microtubule-associated protein. *J. Alzheimer's Dis.* **2004**, *6*, 435-442. doi:10.3233/jad-2004-6412.
  20. Scott, D.A.; Tabarean, I.; Tang, Y.; Cartier, A.; Masliah, E.; Roy, S. A pathologic cascade leading to synaptic dysfunction in alpha-synuclein-induced neurodegeneration. *J. Neurosci.* **2010**, *30*, 8083-8095. doi:10.1523/JNEUROSCI.1091-10.2010.
  21. Marsden, C.D. Parkinson's disease. *Lancet* **1990**, *335*, 948-952. doi:10.1016/0140-6736(90)91006-v.
  22. Braak, H.; Del-Tredici, K.; Rüb, U.; de Vos, R.A.; Jansen-Steur, E.N.; Braak, E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging* **2003**, *24*, 197-211. doi:10.1016/s0197-4580(02)00065-9.
  23. Sampson, T.R.; Debelius, J.W.; Thron, T.; Janssen, S.; Shastri, G.G.; Ilhan, Z.E.; Challis, C.; Schretter, C.E.; Rocha, S.; Gradinaru, V.; Chesselet, M.F.; Keshavarzian, A.; Shannon, K.M.; Krajmalnik-Brown, R.; Wittung-Stafshede, P.; Knight, R.; Mazmanian, S.K. Gut Microbiota Regulate Motor Deficits and Neuroinflammation in a Model of Parkinson's Disease. *Cell* **2016**, *167*, 1469-1480. e12.doi:10.1016/j.cell.2016.11.018.
  24. a) Polymeropoulos, M.H.; Lavedan, C.; Leroy, E.; Ide, S.E.; Dehejia, A.; Dutra, A.; Pike, B.; Root, H.; Rubenstein, J.; Boyer, R.; Stenroos, E.S.; Chandrasekharappa, S.; Athanassiadou, A.; Papapetropoulos, T.;

- Johnson, W.G.; Lazzarini, A.M.; Duvoisin, R.C.; Di-Iorio, G.; Golbe, L.I.; Nussbaum, R.L. Mutation in the  $\alpha$ -synuclein gene identified in families with Parkinson's disease. *Science*. **1997**, *276*, 2045-207. doi:10.1126/science.276.5321.2045. b) Trinh, J.; Zeldenrust, F.M.J.; Huang, J.; Kasten, M.; Schaake, S.; Petkovic, S.; Madoev, H.; Grünewald, A.; Almuammar, S.; König, I.R.; Lill, C.M.; Lohmann, K.; Klein, C.; Marras, C. Genotype-phenotype relations for the Parkinson's disease genes SNCA, LRRK2, VPS35: MDSGene systematic review. *Mov. Disord.* **2018**, *33*, 1857-1870. doi:10.1002/mds.27527.
25. Schapira, A.H.; Cooper, J.M.; Dexter, D.; Jenner, P.; Clark, J.B.; Marsden, C.D. Mitochondrial complex I deficiency in Parkinson's disease. *Lancet* **1989**, *1*, 1269. doi:10.1016/s0140-6736(89)92366-0.
  26. a) Kasten, M.; Hartmann, C.; Hampf, J.; Schaake, S.; Westenberger, A.; Vollstedt, E.J.; Black, A.; Domingo, A.; Vulinovic, F.; Dulovic, M.; Zorn, I.; Madoev, H.; Zehnle, H.; Lembeck, C.M.; Schawe, L.; Reginold, J.; Huang, J.; König, I.R.; Bertram, L.; Marras, C.; Lohmann, K.; Lill, C.M.; Klein, C. Genotype-Phenotype Relations for the Parkinson's Disease Genes Parkin, PINK1, DJ1: MDSGene Systematic Review. *Mov. Disord.* **2018**, *33*, 730-741. doi:10.1002/mds.27352. b) Valente, E.M.; Abou-Sleiman, P.M.; Caputo, V.; Muqit, M.M.; Harvey, K.; Gispert, S.; Ali, Z.; Del-Turco, D.; Bentivoglio, A.R.; Healy, D.G.; Albanese, A.; Nussbaum, R.; González-Maldonado, R.; Deller, T.; Salvi, S.; Cortelli, P.; Gilks, W.P.; Latchman, D.S.; Harvey, R.J.; Dallapiccola, B.; Auburger, G.; Wood, N.W. Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* **2004**, *304*, 1158-60. doi:10.1126/science.1096284. c) Kitada, T.; Asakawa, S.; Hattori, N.; Matsumine, H.; Yamamura, Y.; Minoshima, S.; Yokochi, M.; Mizuno, Y.; Shimizu, N. Mutations in the parkin gene causes autosomal recessive juvenile parkinsonism. *Nature* **1998**, *392*, 605-608. doi:10.1038/33416.
  27. Bonifati, V.; Rizzu, P.; van Baren, M.J.; Schaap, O.; Breedveld, G.J.; Krieger, E.; Dekker, M.C.; Squitieri, F.; Ibanez, P.; Joosse, M.; van Dongen, J.W.; Vanacore, N.; van Swieten, J.C.; Brice, A.; Meco, G.; van Duijn, C.M.; Oostra, B.A.; Heutink, P. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science*. **2003**, *299*, 256-259. doi:10.1126/science.1077209.
  28. Bender, A.; Krishnan, K.J.; Morris, C.M.; Taylor, G.A.; Reeve, A.K.; Perry, R.H.; Jaros, E.; Hersheson, J.S.; Betts, J.; Klopstock, T.; Taylor, R.W.; Turnbull, D.M. High levels of mitochondrial DNA deletions in substantia nigra neurons in aging and Parkinson disease. *Nat. Genet.* **2006**, *38*, 515-517. doi:10.1038/ng1769.
  29. Gan-Or, Z.; Dion, P.A.; Rouleau, G.A. Genetic perspective on the role of the autophagy-lysosome pathway in Parkinson disease. *Autophagy* **2015**, *11*, 1443-1457. doi:10.1080/15548627.2015.1067364.
  30. a) Funayama, M.; Hasegawa, K.; Kowa, H.; Saito, M.; Tsuji, S.; Obata, F. A new locus for Parkinson's disease (PARK8) maps to chromosome 12p11.2-q13.1. *Ann. Neurol.* **2002**, *51*, 296-301. doi:10.1002/ana.10113. b) Yue, Z.; Yang, X.W. Dangerous duet: LRRK2 and  $\alpha$ -synuclein jam at CMA. *Nat. Neurosci.* **2013**, *16*, 375-377. doi:10.1038/nn.3361.
  31. a) Balestrino, R.; Schapira, A.H.V. Glucocerebrosidase and Parkinson Disease: Molecular, Clinical, and Therapeutic Implications. *Neuroscientist* **2018**, *24*, 540-559. doi:10.1177/1073858417748875. b) Ryan, E.; Seehra, G.; Sharma, P.; Sidransky, E. GBA1-associated parkinsonism: new insights and therapeutic opportunities. *Curr. Opin. Neurol.* **2019**, *32*, 589-596. doi:10.1097/WCO.0000000000000715.
  32. a) Quadri, M.; Mandemakers, W.; Grochowska, M.M.; Masius, R.; Geut, H.; Fabrizio, E.; Breedveld, G.J.; Kuipers, D.; Minneboo, M.; Vergouw, L.J.M.; Carreras-Mascaro, A.; Yonova-Doing, E.; Simons, E.; Zhao, T.; Di Fonzo, A.B.; Chang, H.C.; Parchi, P.; Melis, M.; Correia-Guedes, L.; Criscuolo, C.; Thomas, A.; Brouwer, R.W.W.; Heijmans, D.; Ingrassia, A.M.T.; Calandra-Buonaura, G.; Rood, J.P.; Capellari, S.; Rozemuller, A.J.; Sarchioto, M.; Fen, C.H.; Vanacore, N.; Olgiati, S.; Wu-Chou, Y.H.; Yeh, T.H.; Boon, A.J.W.; Hoogers, S.E.; Ghazvini, M.; IJpma, A.S.; van IJcken, W.F.J.; Onofri, M.; Barone, P.; Nicholl, D.J.; Puschmann, A.; De Mari, M.; Kievit, A.J.; Barbosa, E.; De Michele, G.; Majoor-Krakauer, D.; van Swieten, J.C.; de Jong, F.J.; Ferreira, J.J.; Cossu, G.; Lu, C.S.; Meco, G.; Cortelli, P.; van de Berg, W.D.J.; Bonifati, V. International Parkinsonism Genetics Network. LRP10 genetic variants in familial Parkinson's disease and dementia with Lewy bodies: a genome-wide linkage and sequencing study. *Lancet Neurol.* **2018**, *17*, 597-608. doi:10.1016/S1474-4422(18)30179-0. Erratum in: *Lancet Neurol.* **2020**, *19*, e2. b) Williams, E.T.; Chen, X.; Moore, D.J. VPS35, the Retromer Complex and Parkinson's Disease. *J Parkinsons Dis.* **2017**, *7*, 219-233. doi:10.3233/JPD-161020.
  33. Cali, T.; Ottolini, D.; Brini, M. Mitochondria, calcium, and endoplasmic reticulum stress in Parkinson's disease. *Biofactors* **2011**, *37*, 228-240. doi:10.1002/biof.159.
  34. Spielman, L.J.; Gibson, D.L.; Klegeris, A. Unhealthy gut, unhealthy brain: The role of the intestinal microbiota in neurodegenerative diseases. *Neurochem. Int.* **2018**, *120*, 149-163. doi:10.1016/j.neuint.2018.08.005.
  35. Bloem, B.R.; Okun, M.S.; Klein, C. Parkinson's disease. *Lancet* **2021**, *397*, 2284-2303. doi:10.1016/S0140-6736(21)00218-X.
  36. a) Fereshtehnejad, S.M.; Postuma, R.B. Subtypes of Parkinson's Disease: What Do They Tell Us About Disease Progression? *Curr. Neurol. Neurosci. Rep.* **2017**, *17*, 34. doi:10.1007/s11910-017-0738-x. b) Stebbins, G.T.; Goetz, C.G.; Burn, D.J.; Jankovic, J.; Khoo, T.K.; Tilley, B.C. How to identify dominant tremor and



- postural instability/gait difficulty groups with the movement disorder society unified Parkinson's disease rating scale: comparison with the unified Parkinson's disease rating scale. *Mov. Disord.* **2013**, *28*, 668-670. doi:10.1002/mds.25383.
37. a) Schapira, A.H.V.; Chaudhuri, K.R.; Jenner, P. Non-motor features of Parkinson disease. *Nat. Rev. Neurosci.* **2017**, *18*, 435-450. doi:10.1038/nrn.2017.62. Erratum in: *Nat. Rev. Neurosci.* **2017**, *18*, 509. b) Balestrino, R.; Martinez-Martin, P. Neuropsychiatric symptoms, behavioral disorders, and quality of life in Parkinson's disease. *J. Neurol. Sci.* **2017**, *373*, 173-178. doi:10.1016/j.jns.2016.12.060.
  38. Postuma, R.B.; Aarsland, D.; Barone, P.; Burn, D.J.; Hawkes, C.H.; Oertel, W.; Ziemssen, T. Identifying prodromal Parkinson's disease: pre-motor disorders in Parkinson's disease. *Mov. Disord.* **2012**, *27*, 617-626. doi:10.1002/mds.24996.
  39. Sousa, E.T.; Lopes, W.A.; Andrade, J.B. Fontes, formação, reatividade e determinação de quinonas na atmosfera. *Quim. Nova* **2016**, *39*, 486-495. doi: 10.5935/0100-4042.20160034
  40. Futuro, D.O.; Ferreira, P.G.; Nicoletti, C.D.; Borba-Santos, L.P.; da Silva, F.C.; Rozental, S.; Ferreira, V.F. The Antifungal Activity of Naphthoquinones: An Integrative Review. *An. Acad. Bras. Cienc.* **2018**, *90*, 1187-1214. doi: 10.1590/0001-3765201820170815
  41. Aminin, D.; Polonik, S. 1,4-Naphthoquinones: Some Biological Properties and Application. *Chem. Pharm. Bull.* **2020**, *68*, 46-57. DOI: 10.1248/cpb.c19-00911
  42. da Silva, M.N.; Ferreira, V.F.; de Souza, M.C.B.V. Um panorama atual da química e da farmacologia de naftoquinonas, com ênfase na beta-lapachona e derivados. *Quim. Nova*, **2003**, *26*, 407-416. DOI:10.1590/S0100-40422003000300019
  43. Ramos-Peralta, L.; López-López, L.I.; Silva-Belmares, S.Y.; Zugasti-Cruz, A.; Rodríguez-Herrera, R.; Aguilar-González, C.N. Naphthoquinone: Bioactivity and Green Synthesis. In *The Battle Against Microbial Pathogens: Basic Science, Technological Advances and Educational Programs* (A. Mendez-Vilas, Ed.) **2015**, 542-550.
  44. Nicolaidis, D.N.; Gautam, D.R.; Litinas, K.E.; Litina, D.J.H.; Fylaktakidou, K.C. Synthesis and evaluation of the antioxidant and antiinflammatory activities of some benzo[1]kellactone derivatives and analogues. *Eur. J. Med. Chem.* **2004**, *39*, 323-332. DOI: 10.1016/j.ejmech.2004.01.003
  45. Carneiro, P.F.; Pinto, M.C.R.F.; Marra, R.K.F.; Silva, F.C.; Resende, J.A.L.C.; Rocha e Silva, L.F.; Alves, H.G.; Barbosa, G.S.; Vasconcellos, M.C.; Lima, E.S.; Pohlit, A.M.; Ferreira, V.F. Synthesis and antimalarial activity of quinones and structurally-related oxirane derivatives. *Eur. J. Med. Chem.* **2016**, *108*, 134-140. DOI: 10.1016/j.ejmech.2015.11.020
  46. Lamberti, M.J.; Rumie, V.N.B.; Silva, F.C.; Ferreira, V.F.; Rivarola, V.A. Synergistic enhancement of antitumor effect of  $\beta$ -Lapachone by photodynamic induction of quinone oxidoreductase (NQO1). *Phytomedicine*, **2013**, *20*, 1007-1012. DOI: 10.1016/j.phymed.2013.04.018
  47. Pinto, C.N.; Dantas, A.P.; de Moura, K.C.G.; Emery, F.S.; Polequevitch, P.F.; Pinto, M.C.F.R.; de Castro, S.L.; Pinto, A.V. Chemical reactivity studies with naphthoquinones from tabebuia with antitrypanosomal efficacy. *Arzneim. Forsch. Drug. Res.* **2000**, *50*, 1120-1128. DOI: 10.1055/s-0031-1300337
  48. Ferreira, M.P.; Cardoso, M.F.C.; da Silva, F.C.; Ferreira, V.F.; Lima, E.S.; Souza, B.V.J. Antifungal activity of synthetic naphthoquinones against dermatophytes and opportunistic fungi: preliminary mechanism-of-action tests. *Ann. Clin. Microbiol. Antimicrob.* **2014**, *13*, 26-32. DOI: 10.1186/1476-0711-13-26
  49. Ravichandiran, P.; Sheet, S.; Premnath, D.; Kim, A.R.; Yoo, D.J. 1,4-naphthoquinone analogues: potent antibacterial agents and mode of action evaluation. *Molecules* **2019**, *24*, 1437. DOI: 10.3390/molecules24071437
  50. Ferraz, P.A.L.; de Abreu, F.C.; Pinto, A.V.; Glezer, V.; Tonholo, J. Goulart, M.O.F.J. Electrochemical aspects of the reduction of biologically active 2-hydroxy-3-alkyl-1,4-naphthoquinones. *Electroanal. Chem.* **2001**, *507*, 275. DOI: 10.1016/S0022-0728(01)00439-9
  51. Rover Júnior, L.; Höehr, N.F.; Vellasco, E.A.P.; Kubota, L.T. Sistema antioxidante envolvendo o ciclo metabólico da glutatona associado a métodos eletroanalíticos na avaliação do estresse oxidativo. *Quim. Nova* **2001**, *24*, 112. DOI: 10.1590/S0100-40422001000100019
  52. Menchinskaya, E.; Chingizova, E.; Pisyagin, E.; Likhatskaya, G.; Sabutski, Y.; Pelageev, D.; Polonik, S.; Aminin, D. Neuroprotective Effect of 1,4-Naphthoquinones in an In Vitro Model of Paraquat and 6-OHDA-Induced Neurotoxicity. *Int. J. Mol. Sci.* **2021**, *22*, 9933. DOI: 10.3390/ijms22189933
  53. Hussain, H.; Krohn, K.; Ahmad, V.U.; Miana, G.A.; Green, I.R. Lapachol: An overview. *Arkivoc*, **2007**, *2*, 145. DOI:10.3998/ark.5550190.0008.204
  54. Thakur, A. Juglone: A therapeutic phytochemical from Juglans regia L. *J. Med. Plants Res.* **2011**, *5*, 5324-5330.
  55. Andújar, I.; Ríos, J.L.; Giner, R.M.; Recio, M.C. Pharmacological Properties of Shikonin - A Review of Literature since 2002. *Planta Med.* **2013**, *79*, 1685-1697.
  56. DOI: 10.1055/s-0033-1350934

57. Rodriguez-Rodriguez, A.; Egea-Guerrero, J.; Murillo-Cabezas, F.; Carrillo-Vico, A. Oxidative Stress in Traumatic Brain Injury. *Curr. Med. Chem.* **2014**, *21*, 1201-1211. doi:10.2174/0929867321666131217153310.
58. Staniek, K.; Gille, L. Is thymoquinone an antioxidant? *BMC Pharmacol.* **2010**, *10*. doi:10.1186/1471-2210-10-s1-a9.
59. Al-Majed, A.A.; Al-Omar, F.A.; Nagi, M.N. Neuroprotective effects of thymoquinone against transient forebrain ischemia in the rat hippocampus. *Eur. J. Pharmacol.* **2006**, *543*, 40-47. doi:10.1016/j.ejphar.2006.05.046.
60. Kanter, M. Protective effects of thymoquinone on the neuronal injury in frontal cortex after chronic toluene exposure. *J. Mol. Histol.* **2011**, *42*, 39-46. doi:10.1007/s10735-010-9305-3.
61. Radad, K.; Hassanein, K.; Al-Shraim, M.; Moldzio, R.; Rausch, W.D. Thymoquinone ameliorates lead-induced brain damage in Sprague Dawley rats. *Exp. Toxicol. Pathol.* **2014**, *66*, 13-17. doi:10.1016/j.etp.2013.07.002.
62. Kassab, R.B.; El-Hennamy, R.E. The role of thymoquinone as a potent antioxidant in ameliorating the neurotoxic effect of sodium arsenate in female rat. *Egypt. J. Basic Appl. Sci.* **2017**, *4*, 160-167. doi:10.1016/j.ejbas.2017.07.002.
63. Wang, Z.; Liu, T.; Gan, L.; Wang, T.; Yuan, X.; Zhang, B.; Chen, H.; Zheng Qiusheng, Q. Shikonin protects mouse brain against cerebral ischemia/reperfusion injury through its antioxidant activity. *Eur. J. Pharmacol.* **2010**, *643*, 211-217. doi:10.1016/j.ejphar.2010.06.027.
64. Wang, Y.; Pan, W.L.; Liang, W.C.; Law, W.K.; Tsz-Ming Ip, D.; Ng, T.B.; Miu-Yee Waye, M.; Chi-Cheong Wan, D. Acetylshikonin, a novel ache inhibitor, inhibits apoptosis via upregulation of heme oxygenase-1 expression in sh-sy5y cells. *Evidence-based Complement. Altern. Med.* **2013**, *2013*. doi:10.1155/2013/937370.
65. Vomhof-DeKrey, E.E.; Picklo, M.J. The Nrf2-antioxidant response element pathway: A target for regulating energy metabolism. *J. Nutr. Biochem.* **2012**, *23*, 1201-1206.
66. Son, T.G.; Camandola, S.; Arumugam, T. V.; Cutler, R.G.; Telljohann, R.S.; Mughal, M.R.; Moore, T.A.; Luo, W.; Yu, Q.S.; Johnson, D.A.; et al. Plumbagin, a novel Nrf2/ARE activator, protects against cerebral ischemia. *J. Neurochem.* **2010**, *112*, 1316-1326. doi:10.1111/j.1471-4159.2009.06552.x.
67. Yuan, J.H.; Pan, F.; Chen, J.; Chen, C.E.; Xie, D.P.; Jiang, X.Z.; Guo, S.J.; Zhou, J. Neuroprotection by plumbagin involves BDNF-TrkB-PI3K/Akt and ERK1/2/JNK pathways in isoflurane-induced neonatal rats. *J. Pharm. Pharmacol.* **2017**, *69*, 896-906. doi:10.1111/jphp.12681.
68. Messeha, S.S.; Zarmouh, N.O.; Mendonca, P.; Kolta, M.G.; Soliman, K.F.A. The attenuating effects of plumbagin on pro-inflammatory cytokine expression in LPS-activated BV-2 microglial cells. *J. Neuroimmunol.* **2017**, *313*, 129-137. doi:10.1016/j.jneuroim.2017.09.007.
69. Choi, H.G.; Song, J.H.; Park, M.; Kim, S.; Kim, C.E.; Kang, K.S.; Shim, S.H. Neuroprotective  $\gamma$ -pyrones from *fusarium solani* js-0169: Cell-based identification of active compounds and an informatics approach to predict the mechanism of action. *Biomolecules* **2020**, *10*. doi:10.3390/biom10010091.
70. Thijssen, H.H.W.; Driktij-Reijnders, M.J. Vitamin K distribution in rat tissues: dietary phylloquinone is a source of tissue menaquinone-4. *Br. J. Nutr.* **1994**, *72*, 415-425. doi:10.1079/bjn19940043.
71. Moghadam, B.F.; Fereidoni, M. Neuroprotective effect of menaquinone-4 (MK-4) on transient global cerebral ischemia/ reperfusion injury in rat. *PLoS One* **2020**, *15*. doi:10.1371/journal.pone.0229769.
72. Sakaue, M.; Mori, N.; Okazaki, M.; Kadowaki, E.; Kaneko, T.; Hemmi, N.; Sekiguchi, H.; Maki, T.; Ozawa, A.; Hara, S.; et al. Vitamin K has the potential to protect neurons from methylmercury-induced cell death In Vitro. *J. Neurosci. Res.* **2011**, *89*, 1052-1058. doi:10.1002/jnr.22630.
73. Huang, S.H.; Fang, S.T.; Chen, Y.C. Molecular mechanism of vitamin k2 protection against amyloid- $\beta$ -induced cytotoxicity. *Biomolecules* **2021**, *11*, 1-22. doi:10.3390/biom11030423.
74. Josey, B.J.; Inks, E.S.; Wen, X.; Chou, C.J. Structure-activity relationship study of vitamin K derivatives yields highly potent neuroprotective agents. *J. Med. Chem.* **2013**, *56*, 1007-1022. doi:10.1021/jm301485d.
75. Nepovimova, E.; Uliassi, E.; Korabecny, J.; Peña-Altamira, L.E.; Samez, S.; Pesaresi, A.; Garcia, G.E.; Bartolini, M.; Andrisano, V.; Bergamini, C.; et al. Multitarget drug design strategy: Quinone-tacrine hybrids designed to block amyloid- $\beta$  aggregation and to exert anticholinesterase and antioxidant effects. *J. Med. Chem.* **2014**, *57*, 8576-8589. doi:10.1021/jm5010804.
76. Škandík, M.; Mrvová, N.; Bezek, Š.; Račková, L. Semisynthetic quercetin-quinone mitigates BV-2 microglia activation through modulation of Nrf2 pathway. *Free Radic. Biol. Med.* **2020**, *152*. doi:10.1016/j.freeradbiomed.2020.02.030.
77. Chen, W.W.; Zhang, X.; Huang, W.J. Role of neuroinflammation in neurodegenerative diseases (Review). *Mol. Med. Rep.* **2016**, *13*, 3391-3396. doi:10.3892/mmr.2016.4948
78. Inoue, K. Microglial activation by purines and pyrimidines. *Glia*. **2002**, *40*, 156-163. doi: 10.1002/glia.10150
79. a) Takenouchi, T.; Sekiyama, K.; Sekigawa, A.; Fujita, M.; Waragai, M.; Sugama, S.; Iwamaru, Y.; Kitani, H.; Hashimoto, M. P2X7 Receptor Signaling Pathway as a Therapeutic Target for Neurodegenerative Diseases. *Arch. Immunol. Ther. Exp.* **2010**, *58*, 91-96. DOI: 10.1007/s00005-010-0069-y b) Calzaferri, F.; Ruiz-Ruiz, C.; de



- Diego, A.M.G.; de Pascual, R.; Méndez-López, I.; Cano-Abad, M.F.; Maneu, V.; de los Ríos, C.; Gandía, L.; García, A.G. The purinergic P2X7 receptor as a potential drug target to combat neuroinflammation in neurodegenerative diseases. *Med. Res. Rev.* **2020**, *40*, 2427-2465. Doi: 10.1002/med.21710
80. Pislyagin, E.; Kozlovskiy, S.; Menchinskaya, E.; Chingizova, E.; Likhatskaya, G.; Gorpenchenko, T.; Sabutski, Y.; Polonik, S.; Aminin, D. Synthetic 1,4-Naphthoquinones inhibit P2X7 receptors in murine neuroblastoma cells. *Bioorganic Med. Chem.* **2021**, *31*. doi:10.1016/j.bmc.2020.115975.
  81. Li, X.; Li, W.; Liu, G.; Shen, X.; Tang, Y. Association between cigarette smoking and Parkinson's disease: A meta-analysis. *Arch. Gerontol. Geriatr.* **2015**, *61*, 510-516. doi: 10.1016/j.archger.2015.08.004
  82. Mappin-Kasirer, B.; Pan, H.; Lewington, S.; Kizza, J.; Gray, R.; Clarke, R.; Peto, R. Tobacco smoking and the risk of Parkinson disease - A 65-year follow-up of 30,000 male British doctors. *Neurology* **2020**, *94*, e2132-e2138. doi: 10.1212/WNL.00000000000009437
  83. Castagnoli, K.; Petzer, J.B.; Steyn, S.J.; van der Schyf, C.J.; Castagnoli Jr., N. Inhibition of human MAO-A and MAO-B by a compound isolated from flue-cured tobacco leaves and its neuroprotective properties in the MPTP mouse model of neurodegeneration. *Inflammopharmacology* **2003**, *11*, 183-188. doi: 10.1163/156856003765764353
  84. Sari, Y.; Khalil, A. Monoamine Oxidase Inhibitors Extracted from Tobacco Smoke as Neuroprotective Factors for Potential Treatment of Parkinson's Disease. *CNS Neurol. Disord. Drug Targets* **2015**, *14*, 777-785. doi: 10.2174/1871527314666150325235608
  85. Marti, J.S.; Kettler, R.; Da Prada, M.; Richards, J.G. Molecular neuroanatomy of MAO-A and MAO-B. *J. Neural Transm. Suppl.* **1990**, *32*, 49-53. doi: 10.1007/978-3-7091-9113-2\_5
  86. Hauser, D.N.; Hastings, T.G. Mitochondrial dysfunction and oxidative stress in Parkinson's disease and monogenic parkinsonism. *Neurobiol. Dis.* **2013**, *51*, 35-42. doi: 10.1016/j.nbd.2012.10.011
  87. Dezs, L.; Vecsei, L. Monoamine Oxidase B Inhibitors in Parkinson's Disease. *CNS Neurol. Disord. Drug Targets* **2017**, *16*, 425-439. doi: 10.2174/1871527316666170124165222
  88. Khalil, A.A.; Steyn, S.; Castagnoli Jr, N. Isolation and Characterization of a Monoamine Oxidase Inhibitor from Tobacco Leaves. *Chem. Res. Toxicol.* **2000**, *13*, 31-35, 2000. doi: 10.1021/tx990146f
  89. Norris, R.K.; Sternhell, S. Long-range spin-spin coupling in 1,4-benzoquinones and some related compounds. *Aust. J. Chem.* **1966**, *19*, 617-627. doi: 10.1071/CH9660617
  90. Castagnoli, K.P.; Steyn, S.J.; Petzer, J.P.; van der Schyf, C.J.; Castagnoli Jr, N. Neuroprotection in the MPTP Parkinsonian C57BL/6 Mouse Model by a Compound Isolated from Tobacco. *Chem. Res. Toxicol.* **2001**, *14*, 523-527. doi: 10.1021/tx000224v
  91. Chiba, K.; Trevor, A.; Castagnoli Jr, N. Metabolism of the neurotoxic tertiary amine, MPTP, by brain monoamine oxidase. *Biochem. Biophys. Res. Commun.* **1984**, *120*, 574-578. doi: 10.1016/0006-291x(84)91293-2
  92. Cerqueira, E.C.; Netz, P.A.; Diniz, C.; do Canto, V.P.; Follmer, C. Molecular insights into human monoamine oxidase (MAO) inhibition by 1,4-naphthoquinone: Evidences for menadione (vitamin K3) acting as a competitive and reversible inhibitor of MAO. *Bioorg. Med. Chem.* **2011**, *19*, 7416-7424. doi: 10.1016/j.bmc.2011.10.049
  93. Mostert, S.; Petzer, A.; Petzer, J.P. Evaluation of Natural and Synthetic 1,4-naphthoquinones as Inhibitors of Monoamine Oxidase. *Chem. Biol. Drug Des.* **2016**, *87*, 737-746. doi: 10.1111/cbdd.1270890. 93. Choi, W.H.; Hong, S.S.; Lee, S.A.; Han, X.H.; Lee, K.S.; Lee, M.K.; Hwang, B.Y.; Ro, J.S. Monoamine Oxidase Inhibitory Naphthoquinones from the Roots of *Lithospermum erythrorhizon*. *Arch. Pharm. Res.* **2005**, *28*, 400-404. doi: 10.1007/BF02977668
  94. Lashuel, H.A.; Overk, C.R.; Oueslati, A.; Masliah, E. The many faces of  $\alpha$ -synuclein: from structure and toxicity to therapeutic target. *Nat. Rev. Neurosci.* **2013**, *14*, 38-48. doi: 10.1038/nrn3406
  95. Bengoa-Vergniory, N.; Roberts, R.F.; Wade-Martins, R.; Alegre-Abarrategui, J. Alpha-synuclein oligomers: a new hope. *Acta Neuropathol.* **2017**, *134*, 819-838. doi: 10.1007/s00401-017-1755-1
  96. Fields, C.F.; Bengoa-Vergniory, N.; Wade-Martins, R. Targeting Alpha-Synuclein as a Therapy for Parkinson's Disease. *Front. Mol. Neurosci.* **2019**, *12*, 299. doi: 10.3389/fnmol.2019.00299
  97. Choi, B.; Choi, M.; Kim, J.; Yang, Y.; Lai, Y.; Kweon, D.; Lee, N.K.; Shin, Y. Large  $\alpha$ -synuclein oligomers inhibit neuronal SNARE-mediated vesicle docking. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 4087-4092. doi: 10.1073/pnas.1218424110
  98. Di Maio, R.; Barrett, P.J.; Hoffman, E.K.; Barrett, C.W.; Zharikov, A.; Borah, A.; Hu, X.; McCoy, J.; Chu, C.T.; Burton, E.A.; Hastings, T.G.; Greenamyre, J.T.  $\alpha$ -Synuclein binds to TOM20 and inhibits mitochondrial protein import in Parkinson's disease. *Sci. Transl. Med.* **2016**, *8*, 342ra78. doi: 10.1126/scitranslmed.aaf3634
  99. Colla, E.; Jensen, P.H.; Pletnikova, O.; Troncoso, J.C.; Glabe, C.; Lee, M.K. Accumulation of Toxic  $\alpha$ -Synuclein Oligomer within Endoplasmic Reticulum Occurs in  $\alpha$ -Synucleinopathy *In Vivo*. *J. Neurosci.* **2012**, *32*, 3301-3305. doi: 10.1523/JNEUROSCI.5368-11.2012
  100. da Silva, F.L.; Cerqueira, E.C.; de Freitas, M.S.; Gonçalves, D.L.; Costa, L.T.; Follmer, C. Vitamins K interact with N-terminus  $\alpha$ -synuclein and modulate the protein fibrillization *in vitro*. Exploring the interaction between quinones and  $\alpha$ -synuclein. *Neurochem. Int.* **2013**, *62*, 113-112.

101. doi: 10.1016/j.neuint.2012.10.001
102. Wang, Y.; Mandelkow, E. Tau in physiology and pathology. *Nat. Rev. Neurosci.* **2016**, *17*, 22-35. doi: 10.1038/nrn.2015.1
103. Venkatramani, A.; Mukherjee, S.; Kumari, A.; Panda, D. Shikonin impedes phase separation and aggregation of tau and protects SH-SY5Y cells from the toxic effects of tau oligomers. *Int. J. Biol. Macromol.* **2022**, *204*, 19-33. doi: 10.1016/j.ijbiomac.2022.01.172
104. Kourounakis, A.P.; Assimopoulou, A.N.; Papageorgiou, V.P.; Gavalas, A.; Kourounakis, P.N. Alkannin and shikonin: effect on free radical processes and on inflammation - a preliminary pharmacochemical investigation. *Arch. Pharm. (Weinheim)* **2002**, *335*, 262-266. doi: 10.1002/1521-4184(200208)335:6
105. Esmaeilzadeh, E.; Gardaneh, M.; Gharib, E.; Sabouni, F. Shikonin Protects Dopaminergic Cell Line PC12 Against 6-Hydroxydopamine-Mediated Neurotoxicity Via Both Glutathione-Dependent and Independent Pathways and by Inhibiting Apoptosis. *Neurochem. Res.* **2013**, *38*, 1590-1604. doi: 10.1007/s11064-013-1061-9
106. Li, J.; Mo, C.; Guo, Y.; Zhang, B.; Feng, X.; Si, Q.; Wu, X.; Zhao, Z.; Gong, L.; He, D.; Shao, J. *et al.* Roles of peptidyl-prolyl isomerase Pin1 in disease pathogenesis. *Theranostics* **2021**, *11*, 3348-3358. doi: 10.7150/thno.45889
107. Ryo, A.; Togo, T.; Nakai, T.; Hirai, A.; Nishi, M.; Yamaguchi, A.; Suzuki, K.; Hirayasu, Y.; Kobayashi, H.; Perrem, K.; Liou, Y.; Aoki, I. Prolyl-isomerase Pin1 Accumulates in Lewy Bodies of Parkinson Disease and Facilitates Formation of  $\alpha$ -Synuclein Inclusions. *J. Biol. Chem.* **2006**, *281*, 4117-4125. doi: 10.1074/jbc.M507026200
108. Tang, Y.T.; Li, Y.; Chu, P.; Ma, X.D.; Tang, Z.Y.; Sun, Z.L. Molecular biological mechanism of action in cancer therapies: Juglone and its derivatives, the future of development. *Biomed. Pharmacother.* **2022**, *148*, 112785. doi: 10.1016/j.biopha.2022.112785
109. Hennig, L.; Christner, C.; Kipping, M.; Schelbert, B.; Rücknagel, K.P.; Grabley, S.; Küllertz, G.; Fischer, G.A. Selective Inactivation of Parvulin-Like Peptidyl-Prolyl cis/trans Isomerases by Juglone. *Biochemistry* **1998**, *37*, 5953-5960. doi: 10.1021/bi973162p
110. Ghosh, A.; Saminathan, H.; Kanthasamy, A.; Anantharam, V.; Jin, H.; Sondarva, G.; Harischandra, D.S.; Qian, Z.; Rana, A.; Kanthasamy, A.G. The Peptidyl-prolyl Isomerase Pin1 Up-regulation and Proapoptotic Function in Dopaminergic Neurons: Relevance to the Pathogenesis of Parkinson Disease. *J. Biol. Chem.* **2013**, *288*, 21955-21971. doi: 10.1074/jbc.M112.444224
111. Choi, S.Y.; Son, T.G.; Park, H.R.; Jang, Y.J.; Oh, S.B.; Jin, B.; Lee, J. Naphthazarin Has a Protective Effect on the 1-Methyl-4-Phenyl-1,2,3,4-Tetrahydropyridine-Induced Parkinson's Disease Model. *J. of Neurosci. Res.* **2012**, *90*, 1842-1849. doi: 10.1002/jnr.23061
112. Park, J.; Leem, Y.; Park, J.; Kim, D.; Kim, H. Neuroprotective Effect of  $\beta$ -Lapachone in MPTP-Induced Parkinson's Disease Mouse Model: Involvement of Astroglial p-AMPK/Nrf2/HO-1 Signaling Pathways. *Biomol. Ther. (Seoul)* **2019**, *27*, 178-184. doi: 10.4062/biomolther.2018.234
113. 112 Ryu, Y.; Park, H.; Go, J.; Lee, I.; Choi, Y.; Lee, C.; Kim, K.  $\beta$ -Lapachone ameliorates L-DOPA-induced dyskinesia in a 6-OHDA-induced mouse model of Parkinson's disease. *Mol. Med. Rep.* **2021**, *23*, 217. doi: 10.3892/mmr.2021.11856
114. 113. Saleem, U.; Gull, Z.; Saleem, A.; Shah, M.A.; Akhtar, M.F.; Anwar, F.; Ahmad, B.; Panichayupakaranant, P. Appraisal of anti-Parkinson activity of rhinacanthin-C in haloperidol-induced parkinsonism in mice: A mechanistic approach. *J. Food Biochem.* **2021**, *45*, e13677. doi: 10.1111/jfbc.13677
115. 114. Cerqueira, E.C.; Netz, P.A.; do Canto, V.P.; Pinto, A.C.; Follmer, C. Beyond Topoisomerase Inhibition: Antitumor 1,4-Naphthoquinones as Potential Inhibitors of Human Monoamine Oxidase. *Chem. Biol. Drug Des.* **2014**, *84*, 401-410. doi: 10.1111/cbdd.12255
116. 115. Cunha, A.S.; Lima, E.L.S.; Pinto, A.C.; Esteves-Souza, A.; Echevarria, A.; Camara, C.A.; Vargas, M.D.; Torres, J.C. Synthesis of Novel Naphthoquinone-Spermidine Conjugates and their Effects on DNA-Topoisomerases I and II- $\alpha$ . *J. Braz. Chem. Soc.* **2006**, *17*, 439-442. doi: 10.1590/S0103-50532006000300002
117. 116. Fiezer, L.F.; Martin, E.L. 2-Hydroxy-1,4-naphthoquinone. *Org. Synth.* **1941**, *21*, 56. doi: 10.15227/orgsyn.021.0056
118. 117. Scherzer-Attali, R.; Convertino, M.; Pellarin, R.; Gazit, E.; Segal, D.; Caflic, A. Methylations of Tryptophan-Modified Naphthoquinone Affect Its Inhibitory Potential toward A $\beta$  Aggregation. *J. Phys. Chem. B* **2013**, *117*, 1780-1789. doi: 10.1021/jp309066p
119. 118. Paul, A.; Viswanathan, G.K.; Mahapatra, S.; Balboni, G.; Pacifico, S.; Gazit, E.; Segal, D. Antagonistic Activity of Naphthoquinone-Based Hybrids toward Amyloids Associated with Alzheimer's Disease and Type-2 Diabetes. *ACS Chem. Neurosci.* **2019**, *10*, 3510-3520. doi: 10.1021/acscchemneuro.9b00123
120. 119. Scherzer-Attali, R.; Shaltiel-Karyo, R.; Adalist, Y. H.; Segal, D.; Gazit, E. Generic inhibition of amyloidogenic proteins by two naphthoquinone-tryptophan hybrid molecules. *Proteins* **2012**, *80*, 1962-1973. doi: 10.1002/prot.24080

121. 120. Paul, A.; Huber, A.; Rand, D.; Gosselet, F.; Cooper, I.; Gazit, E.; Segal, D. Naphthoquinone–Dopamine Hybrids Inhibit  $\alpha$ -Synuclein Aggregation, Disrupt Preformed Fibrils, and Attenuate Aggregate-Induced Toxicity. *Chem. Eur. J.* **2020**, *26*, 16486–16496. doi: 10.1002/chem.202003374
122. 121. Shrestha-Dawadi, P.D.; Bittner, S.; Fridkin, M.; Rahimpour, S. On the synthesis of naphthoquinonyl heterocyclic amino acids. *Synthesis* **1996**, *12*, 1468–1472. doi: 10.1055/s-1996-4417
123. 122. Paul, A.; Zhang, B.; Mohapatra, S.; Li, G.; Li, Y.; Gazit, E.; Segal, D. Novel Mannitol-Based Small Molecules for Inhibiting Aggregation of  $\alpha$ -Synuclein Amyloids in Parkinson's Disease. *Front. Mol. Biosci.* **2019**, *6*, 16. doi: 10.3389/fmolb.2019.00016
124. 123. Anufriev, V.P.; Malinovskaya, G.V.; Novikov, V.L.; Balanyova, N.N.; Polonik, S.G. The Reductive Dehalogenation Of Halosubstituted Naphthazarins and Quinizarins As A Simple Route To Parent Compounds. *Synth. Commun.* **1998**, *28*, 2149–2157. doi: 10.1080/00397919808007029
125. 124. Huot, R.; Brassard, P. Friedel-Crafts Condensations with Maleic Anhydrides. III. The Synthesis of Polyhydroxylated Naphthoquinones. *Can. J. Chem.* **1974**, *52*, 838–842. doi: 10.1139/v74-132
126. 125. Polonik, N.S.; Polonik, S.G.; Denisenko, V.A.; Moiseenko, O.P. Reaction of Dichloronaphthazarins with Sodium Nitrite as a Route to Natural Pigments Echinamines A and B and Related Aminonaphthazarins. *Synthesis* **2011**, *2011*, 3350–3358. doi: 10.1055/s-0030-1260229
127. 126. Polonik, N.S.; Polonik, S.G. DMSO-mediated transformation of 3-amino-2-hydroxynaphthazarins to natural 2,3-dihydroxynaphthazarins and related compounds. *Tetrahedron Lett.* **2016**, *57*, 3303–3306. doi: 10.1016/j.tetlet.2016.06.056
128. 127. Sabutskii, Y.E.; Denisenko, V.A.; Polonik, S.G. The Acid-Catalyzed 2-O-Alkylation of Substituted 2-Hydroxy-1,4-naphthoquinones by Alcohols: Versatile Preparative Synthesis of Spinochrome D and Its 6-Alkoxy Derivatives. *Synthesis* **2018**, *50*, 3738–3748. doi: 10.1055/s-0037-1610415
129. 128. Anufriev, V.P.; Novikov, V.L.; Maximov, O.B.; Elyakov, G.B.; Levitsky, D.O.; Lebedev, A.V.; Sadretdinov, S.M.; Shvilkin, A.V.; Afonskaya, N.I.; Ruda, M.Y.; Cherpachenko, N.M. Synthesis Of Some Hydroxynaphthazarins And Their Cardioprotective Effects Under Ischemia-Reperfusion *In Vivo*. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 587–592. doi: 10.1016/s0960-894x(98)00075-4
130. 129. Pelageev, D.N.; Panchenko, M.N.; Pokhilo, N.D.; Anufriev, V. F. Synthesis of 2,2'-(ethane-1,1-diyl)bis(3,5,6,7,8-pentahydroxy-1,4-naphthoquinone), a metabolite of the sea urchins *Spatangus purpureus*, *Strongylocentrotus intermedius*, and *S. droebachiensis*. *Russ. Chem. Bull.* **2010**, *59*, 1472–1476. doi: 10.1007/s11172-010-0265-2
131. 130. Polonik S.G.; Tolkach A.M.; Uvarova N.I. Glycosidation of echinochrome and related hydroxynaphthazarins by orthoester method. *Russ. J. Org. Chem.* **1994**, *30*, 248–253.
132. 131. Polonik, S.; Likhatskaya, G.; Sabutski, Y.; Pelageev, D.; Denisenko, V.; Pisl'yagin, E.; Chingizova, E.; Menchinskaya, E.; Aminin, D. Synthesis, Cytotoxic Activity Evaluation and Quantitative Structure-Activity Analysis of Substituted 5,8-Dihydroxy-1,4-naphthoquinones and Their O- and S-Glycoside Derivatives Tested against Neuro-2a Cancer Cells. *Mar. Drugs* **2020**, *18*, 602. doi: 10.3390/md18120602
133. 132. Lebedev, A. V.; Ivanova, M. V.; Ruuge, E. K. How do calcium ions induce free radical oxidation of hydroxy-1,4-naphthoquinone?  $\text{Ca}^{2+}$  stabilizes the naphthosemiquinone anion-radical of echinochrome A. *Arch. Biochem. Biophys.* **2003**, *413*, 191–198. doi:10.1016/S0003-9861(03)00111-5
134. 133. Ekimova, I.V.; Plaksina, D.V.; Pastukhov, Y.F.; Lapshina, K.V.; Lazarev, V.F.; Mikhaylova, E.R.; Polonik, S.G.; Pani, B.; Margulis, B.A.; Guzhova, I.V.; Nudler, E. New HSF1 inducer as a therapeutic agent in a rodent model of Parkinson's disease. *Exp. Neurol.* **2018**, *306*, 199–208. doi: 10.1016/j.expneurol.2018.04.012
135. 134. Belan, D.V.; Polonik, S.G.; Ekimova, I.V. Assessment of the Efficacy of Preventive Therapy with Chaperone Inducer U133 in a Model of the Preclinical Stage of Parkinson's Disease in Elderly Rats. *Neurosci. Behav. Physiol.* **2021**, *51*, 673–680. doi: 10.1007/s11055-021-01120-3

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.