

# A Proof Of The Riemann Hypothesis Based On A New Expression Of The Completed Zeta Function

Weicun Zhang

**Abstract** Based on the Hadamard product  $\xi(s) = \xi(0) \prod_{\rho} (1 - \frac{s}{\rho})$ , a new absolute convergent expression of  $\xi(s)$  is obtained by paring  $\rho_i$  and  $\bar{\rho}_i$ , and putting all the  $\rho_i$  related multiple zeros together in one factor

$$\xi(s) = \xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i}$$

where  $\xi(0) = \frac{1}{2}$ ,  $\rho_i = \alpha_i + j\beta_i$  and  $\bar{\rho}_i = \alpha_i - j\beta_i$  are the complex conjugate zeros of  $\xi(s)$ ,  $0 < \alpha_i < 1$  and  $\beta_i \neq 0$  are real numbers,  $d_i \geq 1$  are the real multiplicities of  $\rho_i$ ,  $\beta_i$  are in order of increasing  $|\beta_i|$ , i.e.,  $|\beta_1| \leq |\beta_2| \leq |\beta_3| \leq \dots$ .

Then, by the functional equation  $\xi(s) = \xi(1-s)$ , we have

$$\xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i} = \xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(1-s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i}$$

i.e.,

$$\prod_{i=1}^{\infty} \left( 1 + \frac{(s - \alpha_i)^2}{\beta_i^2} \right)^{d_i} = \prod_{i=1}^{\infty} \left( 1 + \frac{(1-s - \alpha_i)^2}{\beta_i^2} \right)^{d_i}$$

which, by Lemma 3, is equivalent to

$$\alpha_i = \frac{1}{2}, i = 1, 2, 3, \dots, \infty$$

Thus, we conclude that the Riemann Hypothesis is true.

**Keywords** Riemann Hypothesis (RH) · Proof · Completed zeta function

**Mathematics Subject Classification (2020)** 11M26

Weicun Zhang

University of Science and Technology Beijing, Beijing 100083, China

ORCID: 0000-0003-0047-0558

E-mail: weicunzhang@ustb.edu.cn

## 1 Introduction

It has been almost 163 years since the Riemann Hypothesis (RH) was proposed in 1859 [1]. Many efforts and achievements have been made towards proving this celebrated hypothesis, but it is still an open problem [2–3].

The Riemann zeta function is the function of the complex variable  $s$ , defined in the half-plane  $\Re(s) > 1$  by the absolutely convergent series [2]

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \Re(s) > 1 \quad (1)$$

The connection between the Riemann zeta function and prime numbers can be established through the well-known Euler product, i.e.

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p (1 - p^{-s})^{-1}, \Re(s) > 1 \quad (2)$$

where  $p$  runs over the prime numbers.

Riemann showed how to extend zeta function to the whole complex plane  $\mathbb{C}$  by analytic continuation

$$\zeta(s) = \frac{\pi^{s/2}}{\Gamma(s/2)} \left\{ \frac{1}{s(s-1)} + \int_1^{\infty} (x^{\frac{s}{2}-1} + x^{-\frac{s}{2}-\frac{1}{2}}) \cdot \left( \frac{\theta(x)-1}{2} \right) dx \right\} \quad (3)$$

where  $\theta(x) = \sum_{-\infty}^{\infty} e^{-n^2 \pi x}$  being the Jacobi theta function,  $\Gamma$  being the Gamma function in the following Weierstrass expression (Meanwhile, there are also Gauss expression, Euler expression, and integral expression of the Gamma function.)

$$\frac{1}{\Gamma(s)} = s \cdot e^{\gamma s} \prod_{n=1}^{\infty} \left( 1 + \frac{s}{n} \right) e^{-s/n} \quad (4)$$

where  $\gamma$  is the Euler-Mascheroni constant.

As shown by Riemann,  $\zeta(s)$  extends to  $\mathbb{C}$  as a meromorphic function with only a simple pole at  $s = 1$ , with residue 1, and satisfies the following functional equation

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \zeta(1-s) \quad (5)$$

The Riemann zeta function  $\zeta(s)$  has zeros at the negative even integers:  $-2, -4, -6, -8, \dots$  and one refers to them as the **trivial zeros**. The other zeros of  $\zeta(s)$  are the complex numbers, i.e., **non-trivial zeros** [2].

In 1896, Hadamard [4] and Poussin [5] independently proved that no zeros could lie on the line  $\Re(s) = 1$ . Together with the functional equation and the fact that there are no zeros with real part greater than 1, this showed that all non-trivial zeros must lie in the interior of the **critical strip**  $0 < \Re(s) < 1$ .

This was a key step in their first proofs of the famous **Prime Number Theorem**.

Later on, Hardy (1914) [6], Hardy and Littlewood (1921) [7] showed that there are infinitely many zeros on the **critical line**  $\Re(s) = \frac{1}{2}$ , which was an astonishing result at that time.

As a summary, we have the following results on the properties of the non-trivial zeros of  $\zeta(s)$  [4–9].

**Lemma 1:** Non-trivial zeroes of  $\zeta(s)$ , noted as  $\rho = \alpha + j\beta$ , have the following properties

- 1) The number of non-trivial zeroes is infinity;
- 2)  $\beta \neq 0$ ;
- 3)  $0 < \alpha < 1$ ;
- 4)  $\rho, \bar{\rho}, 1 - \bar{\rho}, 1 - \rho$  are all non-trivial zeroes.

As further study, a completed zeta function  $\xi(s)$  is defined as

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\zeta(s) \quad (6)$$

It is well-known that  $\xi(s)$  is an entire function of order 1. This implies  $\xi(s)$  is analytic, and can be expressed as infinite polynomial, in the whole complex plane  $\mathbb{C}$ .

In addition, replacing  $s$  with  $1-s$  in Eq.(6), and combining Eq.(5), we have the following functional equation

$$\xi(s) = \xi(1-s) \quad (7)$$

Considering the definition of  $\xi(s)$ , and recalling Eq.(4), the trivial zeros of  $\zeta(s)$  are canceled by the poles of  $\Gamma(\frac{s}{2})$ . The zero of  $s-1$  and the pole of  $\zeta(s)$  cancel; the zero  $s=0$  and the pole of  $\Gamma(\frac{s}{2})$  cancel [9–10]. Thus, all the zeros of  $\xi(s)$  are exactly the nontrivial zeros of  $\zeta(s)$ . Then we have the following Lemma 2.

**Lemma 2:** The zeros of  $\xi(s)$  coincide with the non-trivial zeros of  $\zeta(s)$ .

According to Lemma 2, the following two statements for the RH are equivalent.

**Statement 1 of the RH:** All the non-trivial zeros of  $\zeta(s)$  have real part equal to  $\frac{1}{2}$ .

**Statement 2 of the RH:** All the zeros of  $\xi(s)$  have real part equal to  $\frac{1}{2}$ .

To prove the RH, a natural thinking is to estimate the numbers of non-trivial zeros of  $\zeta(s)$  inside or outside some areas according to Argument Principle. Along this train of thought, there are many research works. Let  $N(T)$

denote the number of zeros of  $\zeta(s)$  inside the rectangle:  $0 < \alpha < 1, 0 < \beta \leq T$ , and let  $N_0(T)$  denote the number of zeros of  $\zeta(s)$  on the line  $\alpha = \frac{1}{2}, 0 < \beta \leq T$ . Selberg proved that there exist positive constants  $c$  and  $T_0$ , such that  $N_0(T) > cN(T), (T > T_0)$  [11], later on, Levinson proved that  $c \geq \frac{1}{3}$  [12], Lou and Yao proved that  $c \geq 0.3484$  [13], Conrey proved that  $c \geq \frac{2}{5}$  [14], Bui, Conrey and Young proved that  $c \geq 0.41$  [15], Feng proved that  $c \geq 0.4128$  [16].

On the other hand, many zeros have been calculated by hand or by computer programs. Among others, Riemann found the first three non-trivial zeros [17]. Gram found the first 15 zeros based on Euler-Maclaurin summation [18]. Titchmarsh calculated the 138<sup>th</sup> to 195<sup>th</sup> zeros using the Riemann-Siegel formula [19–20]. Here are the first three (pairs of) zeros:  $\frac{1}{2} \pm j14.1347251$ ;  $\frac{1}{2} \pm j21.0220396$ ;  $\frac{1}{2} \pm j25.0108575$ .

The idea of this paper is originated from Euler's work on proving the following famous equality

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \cdots = \frac{\pi^2}{6} \quad (8)$$

This interesting result is deduced by comparing the like terms of two types of infinite expressions, i.e., infinite polynomial and infinite product, as shown in the following

$$\begin{aligned} \frac{\sin x}{x} &= 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \cdots \\ &= (1 - \frac{x^2}{\pi^2})(1 - \frac{x^2}{4\pi^2})(1 - \frac{x^2}{9\pi^2}) \cdots \end{aligned} \quad (9)$$

Then it is conjectured that  $\xi(s)$  should be factored into  $(1 + \frac{(s-\alpha_i)^2}{\beta_i^2})$  or something like that, which was verified by pairing  $\rho_i$  and  $\bar{\rho}_i$  in the Hadamard product of  $\xi(s)$  to obtain

$$(1 - \frac{s}{\rho_i})(1 - \frac{s}{\bar{\rho}_i}) = \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} (1 + \frac{(s - \alpha_i)^2}{\beta_i^2})$$

The Hadamard product of  $\xi(s)$  as shown in Eq.(10) was first proposed by Riemann, however, it was Hadamard [21] who showed the validity of this infinite product expansion.

$$\xi(s) = \xi(0) \prod_{\rho} (1 - \frac{s}{\rho}) \quad (10)$$

where  $\xi(0) = \frac{1}{2}$ ,  $\rho$  runs over all the non-trivial zeros of the Riemann zeta function  $\zeta(s)$ , or in another word,  $\rho$  runs over all the zeros of the completed zeta function  $\xi(s)$ .

To ensure the absolute convergence of the infinite product expansion,  $\rho$  and  $1 - \rho$  are paired. Later in Section 3, we will show that  $\rho$  and  $\bar{\rho}$  can also be paired to ensure the absolute convergence of the infinite product expansion.

## 2 Lemmas

In this section, we first explain the concepts of multiple zeros of  $\xi(s)$  with their real multiplicities. And then we give three lemmas to support the proof of the RH, in which Lemma 3 is the key lemma.

**Multiple zeros of  $\xi(s)$ :** As shown in Figure 1, the multiple zeros of  $\xi(s)$  are defined in terms of the quadruplet, i.e.,  $\rho, \bar{\rho}, 1 - \rho, 1 - \bar{\rho}$ .




Figure 1 Illustration of the multiple zeros of  $\xi(s)$

It should be noticed that the multiple zeros with their real multiplicities of  $\xi(s)$  are objective existence, but the expression of the corresponding factors of  $\xi(s)$  are optional to some extent. For example, the multiple zeros as shown in Figure 1 have two different expressions as factors of  $\xi(s)$  and  $\xi(1-s)$ , respectively, i.e.,  $\left[ \left(1 + \frac{(s-\alpha_1)^2}{\beta_1^2}\right)^2, \left(1 + \frac{(1-s-\alpha_1)^2}{\beta_1^2}\right)^2 \right]$ , or  $\left[ \left(1 + \frac{(s-\alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s-\alpha_2)^2}{\beta_2^2}\right), \left(1 + \frac{(1-s-\alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1-s-\alpha_2)^2}{\beta_2^2}\right), \alpha_1 + \alpha_2 = 1, \beta_1 = \beta_2 \right]$ .

To exclude the latter expression, we stipulate that the multiple zero  $\rho_i$  related factor of  $\xi(s)$  takes the form of  $\left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i}$ , where  $d_i \geq 1$  is the real multiplicity of  $\rho_i$ .

**Lemma 3:** Given two infinite products

$$f(s) = \prod_{i=1}^{\infty} \left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} \quad (11)$$

and

$$f(1-s) = \prod_{i=1}^{\infty} \left(1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} \quad (12)$$

where  $s$  is a complex variable,  $\rho_i = \alpha_i + j\beta_i$  and  $\bar{\rho}_i = \alpha_i - j\beta_i$  are the complex conjugate zeros of  $\xi(s)$ ,  $0 < \alpha_i < 1$  and  $\beta_i \neq 0$  are real numbers,  $d_i \geq 1$  are the real multiplicities of  $\rho_i$ ,  $i$  are natural numbers from 1 to infinity,  $\beta_i$  are in order of increasing  $|\beta_i|$ , i.e.,  $|\beta_1| \leq |\beta_2| \leq |\beta_3| \leq \dots$ .

Then we have

$$f(s) = f(1-s) \Leftrightarrow \alpha_i = \frac{1}{2}, i = 1, 2, 3, \dots, \infty \quad (13)$$

where " $\Leftrightarrow$ " is the equivalent sign.

**Proof:** First of all, we have the following fact:

$$\left(1 + \frac{(s-\alpha)^2}{\beta^2}\right)^d = \left(1 + \frac{(1-s-\alpha)^2}{\beta^2}\right)^d \Leftrightarrow \alpha = \frac{1}{2} \quad (14)$$

where  $d \geq 1$  is a natural number,  $\alpha \neq 0$  and  $\beta \neq 0$  are real numbers.

Next, the proof is based on Transfinite Induction.

Let  $P(n)$  be:

$$\begin{aligned} \prod_{i=1}^n \left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} &= \prod_{i=1}^n \left(1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \Leftrightarrow \\ \left\{ \begin{array}{l} \left(1 + \frac{(s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1-s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \dots \\ \left(1 + \frac{(s-\alpha_n)^2}{\beta_n^2}\right)^{d_n} = \left(1 + \frac{(1-s-\alpha_n)^2}{\beta_n^2}\right)^{d_n} \end{array} \right. & \quad (15) \\ \Leftrightarrow \alpha_i = \frac{1}{2}, i = 1 \dots n \end{aligned}$$

According to Eq.(14),  $P(1)$  is an obvious fact as the **Base Case**, i.e.,

$$\begin{aligned} \prod_{i=1}^1 \left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} &= \prod_{i=1}^1 \left(1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \Leftrightarrow \left(1 + \frac{(s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} &= \left(1 + \frac{(1-s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \Leftrightarrow \alpha_1 &= \frac{1}{2} \end{aligned} \quad (16)$$

To be more convincing, let's further check  $P(2)$ , i.e.,

$$\begin{aligned} \prod_{i=1}^2 \left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} &= \prod_{i=1}^2 \left(1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \Leftrightarrow \\ \left\{ \begin{array}{l} \left(1 + \frac{(s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1-s-\alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \left(1 + \frac{(s-\alpha_2)^2}{\beta_2^2}\right)^{d_2} = \left(1 + \frac{(1-s-\alpha_2)^2}{\beta_2^2}\right)^{d_2} \end{array} \right. & \quad (17) \\ \Leftrightarrow \alpha_1 = \alpha_2 &= \frac{1}{2} \end{aligned}$$

which is also an obvious fact according to Lemma 5 with another possibility  $\alpha_1 + \alpha_2 = 1, \beta_1 = \beta_2$  excluded, due to that means the second group of zeros  $(\alpha_2 \pm j\beta_2, 1 - \alpha_2 \pm j\beta_2)$  and the first group of zeros  $(\alpha_1 \pm j\beta_1, 1 - \alpha_1 \pm j\beta_1)$  are duplicated with each other in terms of quadruplet. This contradicts the fact that the real multiplicities of  $\rho_1$  and  $\rho_2$  are  $d_1$  and  $d_2$ , respectively.

As the **Successor Case**, i.e., we need to prove  $P(n) \Rightarrow P(n+1)$ .  
Actually, we have

$$\prod_{i=1}^{n+1} \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} = \prod_{i=1}^{n+1} \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i}$$

$$\Leftrightarrow$$

$$\prod_{i=1}^n \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \left(1 + \frac{(s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}} = \prod_{i=1}^n \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \left(1 + \frac{(1 - s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}}$$

$\Leftrightarrow$  (by Lemma 5 with the similar consideration as in proving  $P(2)$  to exclude the possibility that  $\beta_{n+1} = \beta_n, \alpha_{n+1} + \alpha_n = 1$ )

$$\left\{ \prod_{i=1}^n \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} = \prod_{i=1}^n \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \right. \\ \left. \left(1 + \frac{(s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}} = \left(1 + \frac{(1 - s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}} \right.$$

$\Leftrightarrow$  (by Eq.(15))

$$\left\{ \begin{array}{l} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \dots \\ \left(1 + \frac{(s - \alpha_n)^2}{\beta_n^2}\right)^{d_n} = \left(1 + \frac{(1 - s - \alpha_n)^2}{\beta_n^2}\right)^{d_n} \\ \left(1 + \frac{(s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}} = \left(1 + \frac{(1 - s - \alpha_{n+1})^2}{\beta_{n+1}^2}\right)^{d_{n+1}} \end{array} \right.$$

$\Leftrightarrow$  (by Eq.(14))

$$\alpha_i = \frac{1}{2}, i = 1, 2, 3, \dots, n, n+1 \quad (18)$$

Thus the **Successor Case** is true, i.e.,  $P(n) \Rightarrow P(n+1)$ .

Next, we prove that  $P(\infty)$  holds by considering well-ordered ordinal set  $A$  indexing the family of statements  $P(\gamma : \gamma \in A)$ ,  $A = \mathbb{N} \cup \{\omega\}$  with the ordering that  $n < \omega$  for all natural numbers  $n$ ,  $\omega$  is the first limit ordinal. It is well-known that  $\omega = \bigcup \{\gamma : \gamma < \omega\}$ .

To prove that  $P(\infty)$  holds, it suffices to prove the **Limit Case**, i.e.,  $P(\gamma < \omega) \Rightarrow P(\omega)$ .  
Actually, we have

$$\begin{aligned} \prod_{i=1}^{\omega} \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} &= \prod_{i=1}^{\omega} \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \Leftrightarrow \\ \prod_{i=1}^{\gamma < \omega} \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \left(1 + \frac{(s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} &= \prod_{i=1}^{\gamma < \omega} \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \left(1 + \frac{(1 - s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} \end{aligned}$$

$\Leftrightarrow$  (by Lemma 5 with the similar consideration as in proving  $P(2)$  to exclude the possibility that  $\beta_{\omega} = \beta_{\gamma < \omega}, \alpha_{\omega} + \alpha_{\gamma < \omega} = 1$ )

$$\left\{ \prod_{i=1}^{\gamma < \omega} \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} = \prod_{i=1}^{\gamma < \omega} \left(1 + \frac{(1 - s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \right. \\ \left. \left(1 + \frac{(s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} = \left(1 + \frac{(1 - s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} \right.$$

$\Leftrightarrow$  (by  $P(\gamma < \omega)$ )

$$\left\{ \begin{array}{l} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \dots \\ \left(1 + \frac{(s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} = \left(1 + \frac{(1 - s - \alpha_{\omega})^2}{\beta_{\omega}^2}\right)^{d_{\omega}} \end{array} \right.$$

$\Leftrightarrow$  (by Eq.(14))

$$\alpha_i = \frac{1}{2}, i = 1, 2, 3, \dots, \omega \quad (19)$$

Thus the **Limit Case** is true, i.e.,  $P(\gamma < \omega) \Rightarrow P(\omega)$ .



Hence we conclude by Transfinite Induction that  $P(\infty)$  holds, i.e.,

$$\begin{aligned} \prod_{i=1}^{\infty} \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} &= \prod_{i=1}^{\infty} \left(1 + \frac{(1-s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \Leftrightarrow \\ \begin{cases} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1-s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \\ \dots \\ \left(1 + \frac{(s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} = \left(1 + \frac{(1-s - \alpha_i)^2}{\beta_i^2}\right)^{d_i} \\ \dots \end{cases} & \quad (20) \\ \Leftrightarrow \alpha_i &= \frac{1}{2}, i = 1 \dots \infty \end{aligned}$$

i.e.,

$$f(s) = f(1-s) \Leftrightarrow \alpha_i = \frac{1}{2}, i = 1, 2, 3, \dots, \infty \quad (21)$$

That completes the proof of Lemma 3.

**Lemma 4:** Given

$$\left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1-s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1-s - \alpha_2)^2}{\beta_2^2}\right) \quad (22)$$

where  $s$  is a complex variable,  $0 < \alpha_i < 1$  and  $\beta_i \neq 0$  are real numbers,  $|\beta_1| \leq |\beta_2|$ .

Then we have

$$\begin{aligned} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) &= \left(1 + \frac{(1-s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1-s - \alpha_2)^2}{\beta_2^2}\right) \\ \Leftrightarrow \\ \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \\ \Leftrightarrow \\ \begin{cases} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) = \left(1 + \frac{(1-s - \alpha_1)^2}{\beta_1^2}\right), \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1-s - \alpha_2)^2}{\beta_2^2}\right), & \beta_1 \neq \beta_2 \\ \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) = \left(1 + \frac{(1-s - \alpha_2)^2}{\beta_2^2}\right), \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1-s - \alpha_1)^2}{\beta_1^2}\right), & \beta_1 = \beta_2 \end{cases} & \quad (23) \end{aligned}$$

**Proof:** Expanding both sides of Eq.(22), and comparing the coefficients of like terms, we obtain (details are omitted to save space)

$$\begin{aligned} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) &= \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right) \\ \Rightarrow \\ \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \end{aligned} \quad (24)$$

The inverse inference of Eq.(24) is also an obvious fact. i.e.,

$$\begin{aligned} \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \\ \Rightarrow \\ \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) &= \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right) \end{aligned}$$

Then we have

$$\begin{aligned} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) &= \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right) \\ \Leftrightarrow \\ \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \end{aligned} \quad (25)$$

Further, according to Eq.(14), i.e.,

$$\left(1 + \frac{(s - \alpha)^2}{\beta^2}\right)^d = \left(1 + \frac{(1 - s - \alpha)^2}{\beta^2}\right)^d \Leftrightarrow \alpha = \frac{1}{2}$$

and another similar fact

$$\left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) = \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right) \Leftrightarrow \alpha_1 + \alpha_2 = 1, \beta_1 = \beta_2$$

$$\left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right) \Leftrightarrow \alpha_1 + \alpha_2 = 1, \beta_1 = \beta_2$$

we have

$$\begin{aligned}
 & \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right) \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right) \\
 & \Leftrightarrow \\
 & \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \\
 & \Leftrightarrow \\
 & \begin{cases} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right), \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right), & \beta_1 \neq \beta_2 \\ \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right) = \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right), \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right) = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right), & \beta_1 = \beta_2 \end{cases} \\
 & \quad \quad \quad (26)
 \end{aligned}$$

That completes the proof of Lemma 4.

**Lemma 5:** Given

$$\left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} \quad (27)$$

where  $s$  is a complex variable,  $0 < \alpha_i < 1$  and  $\beta_i \neq 0$  are real numbers,  $d_1 \geq 1, d_2 \geq 1$  are natural numbers,  $|\beta_1| \leq |\beta_2|$ .

Then we have

$$\begin{aligned}
 & \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} \\
 & \Leftrightarrow \\
 & \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases} \\
 & \quad \quad \quad (28)
 \end{aligned}$$

**Proof:** Based on Lemma 4, and considering another possibility that  $\left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \mid \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \mid \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}$  (where " $\mid$ "

is the divisible sign), or vice versa, we have

$$\begin{aligned}
 & \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} \\
 & \Leftrightarrow \\
 & \begin{cases} \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1}, \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} = \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, & \beta_1 \neq \beta_2 \\ \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \mid \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} \mid \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, & \beta_1 = \beta_2, d_1 < d_2 \\ \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} \mid \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1}, \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2} \mid \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1}, & \beta_1 = \beta_2, d_1 > d_2 \\ \left(1 + \frac{(s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(1 - s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, \left(1 + \frac{(1 - s - \alpha_1)^2}{\beta_1^2}\right)^{d_1} = \left(1 + \frac{(s - \alpha_2)^2}{\beta_2^2}\right)^{d_2}, & \beta_1 = \beta_2, d_1 = d_2 \end{cases} \\
 & \Leftrightarrow \\
 & \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2, d_1 < d_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2, d_1 > d_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2, d_1 = d_2 \end{cases} \\
 & \Leftrightarrow \\
 & \begin{cases} \alpha_1 = \alpha_2 = \frac{1}{2}, & \beta_1 \neq \beta_2 \\ \alpha_1 + \alpha_2 = 1, & \beta_1 = \beta_2 \end{cases}
 \end{aligned} \tag{29}$$

That completes the proof of Lemma 5.

### 3 A Proof of the RH

This section is planned to present a proof of the Riemann Hypothesis. We first prove that Statement 2 of the RH is true, and then by Lemma 2, Statement 1 of the RH is also true.

**Proof of the RH:** The details are delivered in three steps as follows.

**Step 1:** It is well-known that all the zeros of  $\xi(s)$  always come in complex conjugate pairs. Then by pairing  $\rho_i = \alpha_i + j\beta_i$  and  $\bar{\rho}_i = \alpha_i - j\beta_i$  in the

Hadamard product as shown in Eq.(10), we have

$$\begin{aligned}
 \xi(s) &= \xi(0) \prod_{\rho} \left(1 - \frac{s}{\rho}\right) \\
 &= \xi(0) \prod_{i=1}^{\infty} \left(1 - \frac{s}{\rho_i}\right) \left(1 - \frac{s}{\bar{\rho}_i}\right) \\
 &= \xi(0) \prod_{i=1}^{\infty} \left(1 - \frac{s}{\alpha_i + j\beta_i}\right) \left(1 - \frac{s}{\alpha_i - j\beta_i}\right) \\
 &= \xi(0) \prod_{i=1}^{\infty} \left(\frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2}\right)
 \end{aligned} \tag{30}$$

where  $\xi(0) = \frac{1}{2}$ .

The absolute convergence of the infinite product in Eq.(30) in the form

$$\xi(s) = \xi(0) \prod_{i=1}^{\infty} \left(1 - \frac{s}{\rho_i}\right) \left(1 - \frac{s}{\bar{\rho}_i}\right) = \xi(0) \prod_{i=1}^{\infty} \left(1 - \frac{s(2\alpha_i - s)}{|\rho_i|^2}\right) \tag{31}$$

depends on the convergence of infinite series  $\sum_{i=1}^{\infty} \frac{1}{|\rho_i|^2}$ , or equivalently,  $\sum_{\rho} \frac{1}{|\rho|^2}$ , which is an obvious fact according to Theorem 2 in Section 2, Chapter IV of Ref.[22], as shown in the following.

**Theorem 2.**<sup>[22]</sup> The function  $\xi(s)$  is an entire function of order one that has infinitely many zeros  $\rho_n$  such that  $0 \leq \mathbf{Re} \rho_n \leq 1$ . The series  $\sum |\rho_n|^{-1}$  diverges, but the series  $\sum |\rho_n|^{-1-\varepsilon}$  converges for any  $\varepsilon > 0$ . The zeros of  $\xi(s)$  are the nontrivial zeros of  $\zeta(s)$ .

**Remark:** In the Theorem 2 of Ref.[22],  $\mathbf{Re}(\cdot)$  is identical to  $\Re(\cdot)$  in this paper, both  $\mathbf{Re}(\cdot)$  and  $\Re(\cdot)$  mean the real part of any complex number.

Further, considering the absolute convergence of

$$\xi(s) = \xi(0) \prod_{i=1}^{\infty} \left(1 - \frac{s(2\alpha_i - s)}{|\rho_i|^2}\right) = \xi(0) \prod_{i=1}^{\infty} \left(\frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2}\right) \tag{32}$$

we have the following new expression of  $\xi(s)$  by putting all the  $\rho_i$  related multiple factors (zeros) together in the above Eq.(32)

$$\xi(s) = \xi(0) \prod_{i=1}^{\infty} \left(\frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2}\right)^{d_i} \tag{33}$$

where  $d_i \geq 1$  are the real multiplicities of  $\rho_i$ ,  $i$  are natural numbers from 1 to infinity.

**Step 2:** Replacing  $s$  with  $1 - s$  in Eq.(33), we obtain the infinite product expression of  $\xi(1 - s)$ , i.e.,

$$\xi(1-s) = \xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(1-s-\alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i} \quad (34)$$

**Remark:** According to the new expressions of  $\xi(s)$  and  $\xi(1-s)$ , i.e., Eq.(33) and Eq.(34), we may conclude that all  $\rho_i$  and  $1-\rho_i$  related multiple zeros, i.e.,  $(\alpha_i \pm j\beta_i)^{d_i}$ ,  $(1-\alpha_i \pm j\beta_i)^{d_i}$  are included in the  $i^{th}$  group of factors,  $\left(1 + \frac{(s-\alpha_i)^2}{\beta_i^2}\right)^{d_i}$  and  $\left(1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2}\right)^{d_i}$ , respectively, or in another word, before or after the  $i^{th}$  group of factors of  $\xi(s)$  and  $\xi(1-s)$ , there are no  $\rho_i$  and  $1-\rho_i$  related multiple zeros.

Actually, with such arrangement of  $\rho_i$  and  $1-\rho_i$  related multiple factors of  $\xi(s)$  and  $\xi(1-s)$ , we set the reason to exclude, in the proof of the Lemma 3, the "abnormal" situation, i.e., the successor factor and its predecessor factor represent the same quadruplet of zeros.

**Step 3:** According to the functional equation  $\xi(s) = \xi(1-s)$ , and considering Eq.(33) and Eq.(34), we have

$$\xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s-\alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i} = \xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(1-s-\alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i} \quad (35)$$

which is equivalent to

$$\prod_{i=1}^{\infty} \left( 1 + \frac{(s-\alpha_i)^2}{\beta_i^2} \right)^{d_i} = \prod_{i=1}^{\infty} \left( 1 + \frac{(1-s-\alpha_i)^2}{\beta_i^2} \right)^{d_i} \quad (36)$$

And that  $\beta_i$  can be certainly arranged in order of increasing  $|\beta_i|$ , i.e.,  $|\beta_1| \leq |\beta_2| \leq |\beta_3| \leq \dots$ .

Then, according to Lemma 3, Eq.(36) is equivalent to  $\alpha_i = \frac{1}{2}$ , with  $i$  from 1 to infinity.

Thus, we conclude that all the zeros of the completed zeta function  $\xi(s)$  have real part equal to  $\frac{1}{2}$ , i.e., Statement 2 of the RH is true. According to Lemma 2, Statement 1 of the RH is also true, i.e., all the non-trivial zeros of the Riemann zeta function  $\zeta(s)$  have real part equal to  $\frac{1}{2}$ .

That completes the proof of the RH.

**Remark:** By Lemma 1, there are 2 pairs of complex zeros of  $\zeta(s)$  simultaneously, i.e.,  $\rho = \alpha + j\beta$ ,  $\bar{\rho} = \alpha - j\beta$ ,  $1-\rho = 1-\alpha-j\beta$ ,  $1-\bar{\rho} = 1-\alpha+j\beta$  are all the non-trivial zeroes of  $\zeta(s)$ . With the proof of the RH, i.e.,  $\alpha = \frac{1}{2}$ , these 2 pairs of zeros are actually only one pair, because  $\rho = 1-\bar{\rho} = \frac{1}{2} + j\beta$ ,  $\bar{\rho} = 1-\rho = \frac{1}{2} - j\beta$ . Thus Lemma 1 could be modified more precisely as follows.

**Lemma 1\*:** Non-trivial zeroes of  $\zeta(s)$ , noted as  $\rho = \alpha + j\beta$ , have the following properties

- 1) The number of non-trivial zeroes is infinity;
- 2)  $\beta \neq 0$ ;
- 3)  $0 < \alpha < 1$ ;
- 4)  $\rho = 1 - \bar{\rho}$ ,  $\bar{\rho} = 1 - \rho$  are all non-trivial zeroes.

#### 4 Conclusion

The celebrated Riemann Hypothesis is proved to be true based on a new expression of the completed zeta function  $\xi(s)$ , i.e.,

$$\xi(s) = \xi(0) \prod_{i=1}^{\infty} \left( \frac{\beta_i^2}{\alpha_i^2 + \beta_i^2} + \frac{(s - \alpha_i)^2}{\alpha_i^2 + \beta_i^2} \right)^{d_i}$$

where  $\xi(0) = \frac{1}{2}$ ,  $\rho_i = \alpha_i + j\beta_i$  and  $\bar{\rho}_i = \alpha_i - j\beta_i$  are the complex conjugate zeros of  $\xi(s)$ ,  $0 < \alpha_i < 1$  and  $\beta_i \neq 0$  are real numbers,  $d_i \geq 1$  are the real multiplicities of  $\rho_i$ ,  $i$  are natural numbers from 1 to infinity,  $\beta_i$  are in order of increasing  $|\beta_i|$ , i.e.,  $|\beta_1| \leq |\beta_2| \leq |\beta_3| \leq \dots$ .

**Acknowledgements** The author would like to gratefully acknowledge the help received from Prof. Tianguang Chu (Peking University) while preparing this article.

The author states that there is no conflict of interest.

My manuscript has no associated data.

#### References

1. Riemann B. (1859), Über die Anzahl der Primzahlen unter einer gegebenen Grösse. Monatsberichte der Deutschen Akademie der Wissenschaften zu Berlin, 2, 671-680.
2. Bombieri E. (2000), Problems of the millennium: The Riemann Hypothesis, CLAY
3. Peter Sarnak (2004), Problems of the Millennium: The Riemann Hypothesis, CLAY
4. Hadamard J. (1896), Sur la distribution des zéros de la fonction  $\zeta(s)$  et ses conséquences arithmétiques, Bulletin de la Société Mathématique de France, 14: 199-220, doi:10.24033/bmf.545 Reprinted in (Borwein et al. 2008).
5. de la Vallée-Poussin Ch. J. (1896), Recherches analytiques sur la théorie des nombres premiers, Ann. Soc. Sci. Bruxelles, 20: 183-256
6. Hardy G. H. (1914), Sur les Zéros de la Fonction  $\zeta(s)$  de Riemann, C. R. Acad. Sci. Paris, 158: 1012-1014, JFM 45.0716.04 Reprinted in (Borwein et al. 2008).
7. Hardy G. H., Littlewood J. E. (1921), The zeros of Riemann's zeta-function on the critical line, Math. Z., 10 (3-4): 283-317.
8. Tom M. Apostol (1998), Introduction to Analytic Number Theory, New York: Springer.
9. Chengdong Pan, Chengbiao Pan (2016), Basic Analytic Number Theory (in Chinese), 2nd Edition, Harbin Institute of Technology Press.
10. Reyes E. O. (2004), The Riemann zeta function, Master Thesis of California State University, San Bernardino, Theses Digitization Project. 2648. <https://scholarworks.lib.csusb.edu/etd-project/2648>
11. A. Selberg (1942), On the zeros of the zeta-function of Riemann, Der Kong. Norske Vidensk. Selsk. Forhand. 15, 59-62; also, Collected Papers, Springer-Verlag, Berlin - Heidelberg - New York 1989, Vol. I, 156-159.
12. N. Levinson (1974), More than one-third of the zeros of the Riemann zeta function are on  $\sigma = \frac{1}{2}$ , Adv. Math. 13, 383-436.

13. S. Lou and Q. Yao (1981), A lower bound for zeros of Riemanns zeta function on the line  $\sigma = \frac{1}{2}$ , Acta Mathematica Sinica (in chinese), 24, 390-400.
14. J. B. Conrey (1989), More than two fifths of the zeros of the Riemann zeta function are on the critical line, J. reine angew. Math. 399, 1-26.
15. H. M. Bui, J. B. Conrey and M. P. Young (2011), More than 41% of the zeros of the zeta function are on the critical line, <http://arxiv.org/abs/1002.4127v2>.
16. Feng S. (2012), Zeros of the Riemann zeta function on the critical line, Journal of Number Theory, 132(4), 511-542.
17. Siegel, C. L. (1932), Über Riemanns Nachlaß zur analytischen Zahlentheorie, Quellen Studien zur Geschichte der Math. Astron. Und Phys. Abt. B: Studien 2: 45-80, Reprinted in Gesammelte Abhandlungen, Vol. 1. Berlin: Springer-Verlag, 1966.
18. Gram, J. P. (1903), Note sur les zéros de la fonction  $\zeta(s)$  de Riemann, Acta Mathematica, 27: 289-304.
19. Titchmarsh E. C. (1935), The Zeros of the Riemann Zeta-Function, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, The Royal Society, 151 (873): 234-255.
20. Titchmarsh E. C. (1936), The Zeros of the Riemann Zeta-Function, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, The Royal Society, 157 (891): 261-263.
21. Hadamard J. (1893), Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann. Journal de mathématiques pures et appliquées, 9, 171-216.
22. Karatsuba A. A., Nathanson M. B. (1993), Basic Analytic Number Theory, Springer, Berlin, Heidelberg.