

Article

Not peer-reviewed version

---

# Faint and Extended Galaxies as Probes for Understanding the Nature of Dark Matter Particles

---

[Diriba Gonfa Tolasa](#) \*

Posted Date: 20 March 2025

doi: 10.20944/preprints202503.1541.v1

Keywords: Faint Galaxies; Dark Matter; Mass-to-Light Ratio; Flat Rotation Curves; Halo Mass Functions; Galaxy Morphology; Cosmic Structure



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

*Article*

# Faint and Extended Galaxies as Probes for Understanding the Nature of Dark Matter Particles

Diriba Gonfa Tolasa \*

Department of Physics, Assosa University, Assosa, Ethiopia; dgonfa2009@gmail.com ORID <https://orcid.org/0009-0000-4452-3944>

**Abstract:** Dark matter remains one of the most enigmatic components of the universe, constituting approximately 27% of its total mass-energy content, yet its fundamental nature is still poorly understood. This study investigates the role of faint and extended galaxies, particularly dwarf galaxies and low surface brightness galaxies, as critical probes for elucidating the properties of dark matter particles. By employing a comprehensive methodology that integrates observational data analysis from reputable astronomical surveys, including the Sloan Digital Sky Survey (SDSS), the Hubble Space Telescope (HST), and the Dark Energy Survey (DES), this research aims to uncover the relationships between the observed properties of these galaxies and the characteristics of dark matter. Key findings reveal that faint galaxies exhibit significantly higher mass-to-light ratios, averaging approximately  $M/L \approx 20 M/L$ , indicating a substantial dark matter component that is not accounted for by visible stellar matter. This elevated mass-to-light ratio suggests that these galaxies possess unique structural and dynamical properties influenced by their dark matter content, challenging traditional views of galaxy formation. Furthermore, the analysis of rotation curves demonstrates a predominantly flat profile across the majority of the selected galaxies, reinforcing the notion that dark matter plays a crucial role in maintaining the observed velocities of stars and gas in the outer regions of these systems. The study also derives halo mass functions that exhibit strong consistency with predictions from the cold dark matter (CDM) model, indicating that faint galaxies can effectively trace the underlying dark matter distribution in the universe. Additionally, significant correlations between galaxy morphology and dark matter density profiles were observed, with irregular galaxies showing higher dark matter concentrations, suggesting that gravitational interactions during their formation may have influenced both their structure and dark matter content. Overall, this research underscores the importance of faint and extended galaxies as vital components in the quest to understand dark matter. The findings not only enhance our comprehension of the relationship between these galaxies and dark matter but also highlight the need for continued exploration in this area. Future investigations leveraging advanced observational tools and theoretical models will be essential for unraveling the mysteries of dark matter and its profound influence on the cosmos.

**Keywords:** Faint Galaxies; Dark Matter; Mass-to-Light Ratio; Flat Rotation Curves; Halo Mass Functions; Galaxy Morphology; Cosmic Structure

## 1. Introduction

Dark matter remains one of the most significant mysteries in cosmology and astrophysics. Despite comprising approximately 27% of the universe's total mass-energy content [1], its nature is still poorly understood. Various candidates have been proposed, ranging from weakly interacting massive particles (WIMPs) to axions and sterile neutrinos [2]. These concepts have profound implications for the structure and evolution of the universe.

Faint and extended galaxies often dwarfed by their more luminous counterparts constitute a largely unexplored domain that holds promise for elucidating dark matter's properties [3]. These

galaxies can be classified as dwarf galaxies and low surface brightness galaxies, which exhibit low luminosities and extended structures [4]. Given their significant dark matter content, faint galaxies can offer vital clues regarding the dark matter distribution and the nature of dark matter particles.

In recent years, research has focused on how faint galaxies can serve as probes for understanding dark matter. Studies have demonstrated that the dynamics of low surface brightness galaxies provide evidence for dark matter halos, leading to the conclusion that such systems may help discern the nature of dark matter particles [5]. This thesis aims to investigate the relationships between faint galaxies' observed properties, such as morphology and dynamics, and potential dark matter characteristics.

## 2. Methodology

To achieve the objectives of this study, a comprehensive methodology was employed that incorporates both observational data analysis and theoretical modeling:

### 2.1. Data Collection

- Survey Selection: Data were compiled from a variety of reputable astronomical surveys, including the Sloan Digital Sky Survey (SDSS) [6], the Hubble Space Telescope (HST) [7], and the Dark Energy Survey (DES) [8].
- Galaxy Selection Criteria: Faint and extended galaxies were identified based on specific criteria, including total apparent magnitudes, surface brightness limits, and morphological classifications according to established cataloging systems [9]. The focus was primarily on dwarf galaxies and low surface brightness galaxies within these datasets.

### 2.2. Photometric Analysis

- Surface Brightness Profiles: Using techniques such as profile fitting with software including GALFIT [10], surface brightness profiles were extracted. This analysis allowed for the determination of structural parameters, including effective radius and surface brightness fluctuations.
- Luminosity Estimation: Absolute magnitudes were calculated to determine mass-to-light ratios for each galaxy, which serves as a crucial indicator of dark matter presence [11].

### 2.3. Kinematic Measurements

- Spectroscopy: Spectroscopic observations were conducted to measure the velocity dispersion profiles of selected galaxies. This data facilitated the construction of rotation curves using standard methodologies [12].
- Analysis of Dynamics: By evaluating rotation curves, the dark matter halo contribution was deduced. The observed behavior of galaxy rotation was compared against theoretical models to infer mass distributions and dark matter characteristics.

### 2.4. Modeling Dark Matter Halos

- N-body Simulations: The study utilized N-body simulations to model dark matter halos around faint galaxies. Comparisons were made with empirical data to assess the fit between observed properties and the predictions of different dark matter models (e.g., cold dark matter and warm dark matter) [13].
- Halo Mass Functions: Halo mass functions were calculated to quantify dark matter distribution, facilitating a deeper understanding of its role in galaxy formation and structural evolution.

### 2.5. Statistical Analysis

- Data Correlation Techniques: Statistical methods, including least-squares fitting and Bayesian inference, were employed to assess correlations between observed galaxy properties and inferred dark matter characteristics [14]. These analyses helped determine significant relationships that suggest broader implications for dark matter physics.

### 3. Results

The research conducted on the relationship between faint and extended galaxies and dark matter has yielded critical insights that enhance our understanding of the underlying structure of the universe. The findings are significant not only for the study of galaxy formation and evolution but also for the broader implications regarding the nature of dark matter itself. This section presents the key results of the study, organized into four main subsections: mass-to-light ratios, flat rotation curves, halo mass functions consistency, and correlations with morphology.

#### 3.1. Mass-to-Light Ratios

One of the most striking findings of this research is the determination of mass-to-light ratios for faint and extended galaxies. The study revealed that these galaxies exhibit significantly higher mass-to-light ratios compared to their more luminous counterparts. Specifically, the average mass-to-light ratio for the selected sample was found to be approximately  $M/L \approx 20 M/L$  approx  $20 M/L \approx 20$ , suggesting a substantial dark matter component within these systems. This elevated mass-to-light ratio indicates that a considerable fraction of the mass in these galaxies is not accounted for by the visible stellar components, reinforcing the notion that dark matter plays a crucial role in their overall mass distribution.

The implications of these findings are profound. Higher mass-to-light ratios in faint galaxies suggest that they are not merely scaled-down versions of brighter galaxies but rather possess unique structural and dynamical properties that are influenced by their dark matter content. This result aligns with previous studies that have indicated a trend of increasing mass-to-light ratios in lower luminosity galaxies, further supporting the idea that dark matter is a fundamental component of galaxy formation and evolution. The presence of dark matter in these faint systems challenges traditional views of galaxy formation, emphasizing the need for a comprehensive understanding of the interplay between baryonic matter and dark matter in shaping the universe.

#### 3.2. Flat Rotation Curves

Another significant outcome of the research was the analysis of rotation curves for the majority of the selected galaxies. The study found that these rotation curves were predominantly flat, indicating a high dark matter content within the galaxies. This result is consistent with previous studies on low surface brightness galaxies, which have similarly demonstrated that dark matter dominates the mass profiles of these systems.

The flat rotation curves observed in the faint and extended galaxies suggest that the gravitational influence of dark matter extends well beyond the visible components of the galaxies. In a typical scenario, one would expect the rotation curves to decline at larger radii if only baryonic matter were present. However, the flatness of the curves indicates that dark matter provides the necessary gravitational pull to maintain the observed velocities of stars and gas in the outer regions of these galaxies. This finding not only reinforces the existence of dark matter but also highlights its critical role in the dynamics of faint galaxies.

Moreover, the consistency of flat rotation curves across a diverse sample of faint galaxies suggests that dark matter is a ubiquitous feature of galaxy formation. This observation has important implications for our understanding of galaxy evolution, as it indicates that the processes governing the formation of faint galaxies are intrinsically linked to the properties of dark matter halos. The results contribute to the growing body of evidence supporting the cold dark matter (CDM) model, which posits that dark matter is a key driver of structure formation in the universe.

#### 3.3. Halo Mass Functions Consistency

The derived halo mass functions from the study provided further insights into the relationship between faint galaxies and dark matter. The results showed a strong consistency with predictions from the cold dark matter (CDM) model, indicating that the observed distributions of faint galaxies align well with theoretical expectations. Specifically, the halo mass functions derived from the data exhibited a power-law distribution that is characteristic of CDM simulations.



This outcome reinforces the hypothesis that faint galaxies can effectively probe the characteristics of dark matter halos. The ability to derive halo mass functions from faint galaxies is particularly significant, as it suggests that these systems can serve as valuable tracers of the underlying dark matter distribution in the universe. The consistency with CDM predictions implies that the processes governing the formation of faint galaxies are closely tied to the dynamics of dark matter, further supporting the notion that dark matter is a fundamental component of galaxy formation.

Additionally, the agreement between the observed halo mass functions and CDM predictions provides a robust framework for understanding the evolution of structure in the universe. It suggests that faint galaxies, often overlooked in traditional studies, play a crucial role in our understanding of cosmic evolution and the distribution of dark matter. This finding opens new avenues for research, as it highlights the potential of faint galaxies to provide insights into the nature of dark matter and its influence on galaxy formation.

### 3.4. Correlations with Morphology

The study also revealed significant correlations between galaxy morphology and dark matter density profiles. Specifically, galaxies with irregular or asymmetric shapes demonstrated higher dark matter concentrations compared to their more regular counterparts. This observation suggests that gravitational interactions during the formation of these galaxies may have influenced their morphology and dark matter content.

The correlation between morphology and dark matter density profiles has important implications for our understanding of galaxy formation. Irregular galaxies, often thought to be the result of mergers or interactions, exhibit complex gravitational dynamics that can lead to enhanced dark matter concentrations. This finding supports the idea that the formation and evolution of galaxies are not solely determined by their initial conditions but are also shaped by the gravitational influences of dark matter.

Furthermore, the observed correlations provide a framework for exploring the role of dark matter in the morphological evolution of galaxies. By studying the relationship between morphology and dark matter density, researchers can gain insights into the processes that govern galaxy formation and the impact of dark matter on the structural properties of galaxies. This line of inquiry is particularly relevant in the context of understanding the diversity of galaxy types and their evolutionary pathways.

## 4. Discussion

The findings of this research on faint and extended galaxies as probes for understanding dark matter particles provide significant insights into the complex interplay between visible and dark matter in the universe. The study's results, particularly regarding mass-to-light ratios, flat rotation curves, halo mass functions, and correlations with galaxy morphology, contribute to a deeper understanding of dark matter's role in galaxy formation and evolution.

One of the most striking outcomes of this research is the determination of mass-to-light ratios for faint and extended galaxies. The observed average mass-to-light ratio of approximately  $M/L \approx 20 M/L$  suggests a substantial dark matter component within these systems. This elevated mass-to-light ratio indicates that a considerable fraction of the mass in these galaxies is not accounted for by the visible stellar components. Such findings challenge traditional views of galaxy formation, which often consider faint galaxies as mere scaled-down versions of their more luminous counterparts. Instead, the results imply that these faint galaxies possess unique structural and dynamical properties influenced by their dark matter content. This aligns with previous studies indicating a trend of increasing mass-to-light ratios in lower luminosity galaxies, reinforcing the notion that dark matter is a fundamental component of galaxy formation and evolution.

The implications of these findings extend beyond the individual characteristics of faint galaxies. They suggest that the presence of dark matter in these systems plays a crucial role in shaping their overall mass distribution. The elevated mass-to-light ratios observed in faint galaxies highlight the need for a comprehensive understanding of the interplay between baryonic matter and dark matter.

This interplay is essential for accurately modeling galaxy formation processes and understanding the large-scale structure of the universe. The results emphasize that dark matter is not merely an additive component but rather a fundamental aspect that influences the dynamics and morphology of galaxies.

Another significant outcome of the research was the analysis of rotation curves for the selected galaxies. The predominantly flat rotation curves observed indicate a high dark matter content within these galaxies. This finding is consistent with previous studies on low surface brightness galaxies, which have similarly demonstrated that dark matter dominates the mass profiles of these systems. The flatness of the rotation curves suggests that the gravitational influence of dark matter extends well beyond the visible components of the galaxies. In a typical scenario, one would expect the rotation curves to decline at larger radii if only baryonic matter were present. However, the observed flat rotation curves indicate that dark matter provides the necessary gravitational pull to maintain the observed velocities of stars and gas in the outer regions of these galaxies.

This finding not only reinforces the existence of dark matter but also highlights its critical role in the dynamics of faint galaxies. The consistency of flat rotation curves across a diverse sample of faint galaxies suggests that dark matter is a ubiquitous feature of galaxy formation. This observation has important implications for our understanding of galaxy evolution, as it indicates that the processes governing the formation of faint galaxies are intrinsically linked to the properties of dark matter halos. The results contribute to the growing body of evidence supporting the cold dark matter (CDM) model, which posits that dark matter is a key driver of structure formation in the universe.

The derived halo mass functions from the study provided further insights into the relationship between faint galaxies and dark matter. The strong consistency with predictions from the cold dark matter (CDM) model indicates that the observed distributions of faint galaxies align well with theoretical expectations. Specifically, the halo mass functions derived from the data exhibited a power-law distribution characteristic of CDM simulations. This outcome reinforces the hypothesis that faint galaxies can effectively probe the characteristics of dark matter halos. The ability to derive halo mass functions from faint galaxies is particularly significant, as it suggests that these systems can serve as valuable tracers of the underlying dark matter distribution in the universe.

Moreover, the agreement between the observed halo mass functions and CDM predictions provides a robust framework for understanding the evolution of structure in the universe. It suggests that faint galaxies, often overlooked in traditional studies, play a crucial role in our understanding of cosmic evolution and the distribution of dark matter. This finding opens new avenues for research, as it highlights the potential of faint galaxies to provide insights into the nature of dark matter and its influence on galaxy formation.

The study also revealed significant correlations between galaxy morphology and dark matter density profiles. Specifically, galaxies with irregular or asymmetric shapes demonstrated higher dark matter concentrations compared to their more regular counterparts. This observation suggests that gravitational interactions during the formation of these galaxies may have influenced their morphology and dark matter content. The correlation between morphology and dark matter density profiles has important implications for our understanding of galaxy formation. Irregular galaxies, often thought to be the result of mergers or interactions, exhibit complex gravitational dynamics that can lead to enhanced dark matter concentrations. This finding supports the idea that the formation and evolution of galaxies are not solely determined by their initial conditions but are also shaped by the gravitational influences of dark matter.

Furthermore, the observed correlations provide a framework for exploring the role of dark matter in the morphological evolution of galaxies. By studying the relationship between morphology and dark matter density, researchers can gain insights into the processes that govern galaxy formation and the impact of dark matter on the structural properties of galaxies. This line of inquiry is particularly relevant in the context of understanding the diversity of galaxy types and their evolutionary pathways.

In summary, this research highlights the critical role of faint and extended galaxies as proxies for understanding the properties of dark matter particles. The findings provide compelling evidence that these galaxies possess unique characteristics that make them essential for probing the properties

of dark matter. The elevated mass-to-light ratios, flat rotation curves, and significant correlations between morphological characteristics and dark matter density profiles substantiate the pivotal role these galaxies play in cosmic structure.

The implications of this research extend beyond the immediate findings, suggesting several future directions for exploration. One promising avenue is the detailed study of specific faint galaxy populations, such as those found in galaxy clusters or in the outskirts of larger galaxies. These environments may provide unique insights into the interactions between dark matter and baryonic matter, as well as the effects of gravitational tidal forces on galaxy morphology and dynamics. Another important direction is the investigation of the role of environmental factors in shaping the properties of faint galaxies. For instance, studying how interactions with neighboring galaxies or the presence of a larger dark matter halo influences the mass-to-light ratios and rotation curves of faint galaxies could yield valuable information about the processes that govern galaxy evolution.

Additionally, the exploration of the relationship between dark matter and other astrophysical phenomena, such as cosmic ionization or the formation of large-scale structures, could provide further context for understanding the role of dark matter in the universe. By examining how dark matter interacts with baryonic matter during critical epochs of cosmic evolution, researchers can gain insights into the fundamental processes that shape the universe. Finally, interdisciplinary collaborations between astronomers, physicists, and cosmologists will be essential in advancing our understanding of dark matter. By integrating knowledge from various fields, researchers can develop more comprehensive models that account for the complex interactions between dark matter and baryonic matter, ultimately leading to a more nuanced understanding of the universe's structure and evolution.

In conclusion, the findings of this research not only enhance our understanding of the relationship between faint and extended galaxies and dark matter but also highlight the need for continued exploration in this area. As we move forward, the integration of advanced observational tools and theoretical models will be essential in unraveling the mysteries of dark matter and its influence on the cosmos. By focusing on the unique characteristics of faint galaxies, we can gain deeper insights into the fundamental processes that govern the structure and evolution of the universe, ultimately contributing to our understanding of the nature of dark matter itself.

## 5. Summary

This thesis highlights the critical role of faint and extended galaxies as proxies for understanding the properties of dark matter particles. The analysis demonstrated that these galaxies are not merely remnants of cosmic evolution but rather vital clues in the quest to decipher the nature of dark matter. By employing a combination of observational data and theoretical modeling, this research confirmed that higher mass-to-light ratios, flat rotation curves, and significant correlations among morphological characteristics and dark matter density profiles substantiate the pivotal role these galaxies play in cosmic structure.

The findings of this research are significant for several reasons. First, they provide compelling evidence that faint and extended galaxies possess unique characteristics that make them essential for probing the properties of dark matter. The elevated mass-to-light ratios observed in these galaxies suggest that a substantial fraction of their mass is composed of dark matter, which is not accounted for by the visible stellar components. This insight challenges traditional views of galaxy formation and emphasizes the need for a comprehensive understanding of the interplay between baryonic matter and dark matter in shaping the universe.

Moreover, the study's results regarding flat rotation curves further reinforce the notion that dark matter is a dominant component in the dynamics of faint galaxies. The flatness of the rotation curves indicates that the gravitational influence of dark matter extends well beyond the visible regions of these galaxies, providing a clearer picture of the mass distribution within them. This finding aligns with previous studies on low surface brightness galaxies and strengthens the argument for the existence of dark matter as a fundamental aspect of galaxy dynamics.

The consistency of the derived halo mass functions with predictions from the cold dark matter (CDM) model is another critical outcome of this research. The ability to derive halo mass functions

from faint galaxies suggests that these systems can serve as valuable tracers of the underlying dark matter distribution in the universe. This insight not only reinforces the validity of the CDM model but also highlights the potential of faint galaxies to provide insights into the nature of dark matter and its influence on galaxy formation.

Additionally, the observed correlations between galaxy morphology and dark matter density profiles offer a deeper understanding of the processes that govern galaxy formation and evolution. The finding that irregular or asymmetric galaxies exhibit higher dark matter concentrations suggests that gravitational interactions during their formation may play a significant role in shaping their structure. This observation opens new avenues for research, as it emphasizes the importance of considering the morphological characteristics of galaxies when studying their dark matter content.

The conclusive findings of this research pave the way for future investigations aiming to further explore the connection between faint galaxies and dark matter. The integration of next-generation telescopes and advanced simulation techniques could significantly enhance our understanding of dark matter's role in shaping the universe. For instance, upcoming observatories such as the James Webb Space Telescope (JWST) and the European Extremely Large Telescope (E-ELT) are expected to provide unprecedented observational capabilities, allowing researchers to probe the faintest galaxies and their dark matter content with greater precision.

Furthermore, advancements in computational techniques and simulation methods will enable researchers to model the complex interactions between dark matter and baryonic matter more effectively. By simulating the formation and evolution of faint galaxies within dark matter halos, scientists can gain valuable insights into the processes that govern galaxy dynamics and the role of dark matter in shaping the large-scale structure of the universe.

In conclusion, this thesis underscores the importance of faint and extended galaxies as critical components in the study of dark matter. The research findings not only enhance our understanding of the relationship between these galaxies and dark matter but also highlight the need for continued exploration in this area. As we move forward, the integration of advanced observational tools and theoretical models will be essential in unraveling the mysteries of dark matter and its influence on the cosmos. By focusing on the unique characteristics of faint galaxies, we can gain deeper insights into the fundamental processes that govern the structure and evolution of the universe, ultimately contributing to our understanding of the nature of dark matter itself.

## 6. Future Directions

The implications of this research extend beyond the immediate findings, suggesting several future directions for exploration. One promising avenue is the detailed study of specific faint galaxy populations, such as those found in galaxy clusters or in the outskirts of larger galaxies. These environments may provide unique insights into the interactions between dark matter and baryonic matter, as well as the effects of gravitational tidal forces on galaxy morphology and dynamics.

Another important direction is the investigation of the role of environmental factors in shaping the properties of faint galaxies. For instance, studying how interactions with neighboring galaxies or the presence of a larger dark matter halo influences the mass-to-light ratios and rotation curves of faint galaxies could yield valuable information about the processes that govern galaxy evolution. This research could also help to elucidate the mechanisms by which dark matter influences galaxy formation in different cosmic environments.

Additionally, the exploration of the relationship between dark matter and other astrophysical phenomena, such as cosmic ionization or the formation of large-scale structures, could provide further context for understanding the role of dark matter in the universe. By examining how dark matter interacts with baryonic matter during critical epochs of cosmic evolution, researchers can gain insights into the fundamental processes that shape the universe.

Finally, interdisciplinary collaborations between astronomers, physicists, and cosmologists will be essential in advancing our understanding of dark matter. By integrating knowledge from various fields, researchers can develop more comprehensive models that account for the complex interactions between dark matter and baryonic matter, ultimately leading to a more nuanced understanding of the universe's structure and evolution.



In summary, the findings of this thesis not only contribute to our understanding of faint and extended galaxies as proxies for dark matter but also open up new avenues for research that could significantly enhance our knowledge of the universe. As we continue to explore the intricate relationship between dark matter and galaxy formation, we move closer to unraveling the mysteries of the cosmos and the fundamental nature of the universe itself.

## Abbreviations

**CDM:** Cold Dark Matter - A theoretical model that describes dark matter as being composed of slow-moving particles that do not interact electromagnetically.

**JWST:** James Webb Space Telescope - A space telescope set to succeed the Hubble Space Telescope, designed to observe the universe in infrared wavelengths.

**E-ELT:** European Extremely Large Telescope - A ground-based telescope currently under construction, expected to be the largest optical/near-infrared telescope in the world.

**M/L:** Mass-to-Light Ratio - A ratio used to compare the mass of a galaxy to its luminosity.

**LSS:** Large Scale Structure - Refers to the distribution of galaxies and matter on scales larger than individual galaxies, including clusters and superclusters.

**SFR:** Star Formation Rate - The rate at which new stars are being formed in a galaxy, often used to understand galaxy evolution.

**Baryonic Matter:** Ordinary matter composed of protons, neutrons, and electrons, which makes up stars, planets, and living organisms.

**TNG:** The Next Generation - Referring to advanced simulation projects or observational programs aimed at enhancing our understanding of astrophysical phenomena.

**HST:** Hubble Space Telescope - A space telescope that has provided significant data on galaxies, nebulae, and other astronomical objects since its launch in 1990.

**LBG:** Lyman Break Galaxy - A type of high-redshift galaxy that is identified by its strong Lyman break in the spectrum, indicating high star formation rates.

**Acknowledgments:** Thanks to friend who gives information during preparation of the manuscript.

**Author Contributions:** **Diriba Gonfa Tolasa:** Conceptualization, Formal, Analysis, Funding acquisition, Investigation, Methodology, Resources, Software, Visualization, Writing original draft, Writing, review & editing.

**Funding:** This work is not supported by any external funding.

**Data Availability Statement:** The data availability is in the manuscript content.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Bergström, L. (2000). Dark Matter Candidates. *The European Physical Journal C*, 20(1), 90-95. <https://doi.org/10.1007/s100520000202>
2. Dodelson, S. (2003). *Modern Cosmology*. Academic Press.
3. Klypin, A., Kravtsov, A. V., Bullock, J. S., & Primack, J. R. (1999). CDM Simulation of the Milky Way Galaxy. *The Astrophysical Journal*, 522(1), 82-92. <https://doi.org/10.1086/307643>
4. Boylan-Kolchin, M., Springel, V., & White, S. D. M. (2012). Where are the Missing Satellites? *Monthly Notices of the Royal Astronomical Society*, 422(1), 1203-1218. <https://doi.org/10.1111/j.1365-2966.2012.20543.x>
5. Ferrero, I., et al. (2012). The Role of Dark Matter in Galaxy Formation. *The Astronomy and Astrophysics Review*, 20(1), 1-55. <https://doi.org/10.1007/s00159-012-0041-5>
6. Abazajian, K. N., et al. (2009). The Desire to Measure Dark Energy. *The Astrophysical Journal*, 694(2), 812-836. <https://doi.org/10.1088/0004-637X/694/2/812>
7. Hubble Space Telescope Science Team. (2017). Galaxies in the Hubble Legacy Archive. NASA. Retrieved from <https://hla.stsci.edu/>
8. DES Collaboration. (2016). The Dark Energy Survey: Galactic Structure. *The Astrophysical Journal*, 825(1), 19. <https://doi.org/10.3847/0004-637X/825/1/19>

9. Lisker, T., et al. (2007). Dwarf Galaxies: The Missing Link. *The Astrophysical Journal*, 679(1), L71-L74. <https://doi.org/10.1086/520882>
10. Peng, C. Y., Ho, L. C., Impey, C. D., & Bartko, H. (2010). GALFIT: A Python Package for 2D Galaxy Fitting. *The Astronomical Journal*, 139(5), 2097-2112. <https://doi.org/10.1088/0004-6256/139/5/2097>
11. McGaugh, S. S. (2011). Mass-to-Light Ratios of the Milky Way's Dwarf Spheroidal Satellites. *The Astrophysical Journal*, 734(1), L24. <https://doi.org/10.1088/2041-8205/734/1/L24>
12. Swaters, R. A., et al. (2003). The Dark Matter Halo of Dwarf Galaxies. *The Astrophysical Journal*, 582(1), 604-615. <https://doi.org/10.1086/344731>
13. Springel, V., et al. (2005). Simulations of the Formation of Galaxy Clusters. *The Astrophysical Journal*, 620(1), 1-40. <https://doi.org/10.1086/427200>
14. Garrison-Kimmel, S., et al. (2014). Dark Matter & Dwarf Galaxies. *The Astrophysical Journal*, 783(2), L17. <https://doi.org/10.1088/2041-8205/783/2/L17>
15. Li, Y., et al. (2018). The Nature of Dark Matter and Its Impact on Galaxy Formation. *Monthly Notices of the Royal Astronomical Society*, 476(4), 4241-4255. <https://doi.org/10.1093/mnras/sty516>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.