

Article

Not peer-reviewed version

Photochemical and Oxidative Degradation of Chamazulene Contained in Artemisia, Matricaria and Achillea Essential Oils and Setup of Protection Strategies

[Simone Gabbanini](#)^{*}, Jerome Ngwa Neba, [Riccardo Matera](#), [Luca Valgimigli](#)^{*}

Posted Date: 6 May 2024

doi: 10.20944/preprints202405.0280.v1

Keywords: Artemisia arborescence; blue color; analysis; UHPLC MS/MS; GC-MS; photodegradation; anti-oxidant; sunscreen



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Photochemical and Oxidative Degradation of Chamazulene Contained in *Artemisia*, *Matricaria* and *Achillea* Essential Oils and Setup of Protection Strategies

Simone Gabbanini ^{1,*} Jerome Ngwa Neba ², Riccardo Matera ¹ and Luca Valgimigli ^{2,3,*}

¹ R&D Department, BeC s.r.l., Via C. Monteverdi 49, 47122 Forlì, Italy

² Department of Chemistry "Ciamician", University of Bologna, Via Gobetti 85, 40129 Bologna, Italy

³ Tecnopolo di Rimini, Via D. Campana 71, 47922 Rimini, Italy

* Correspondence: laboratorio@bec-natura.com (S.G.); luca.valgimigli@unibo.it (L.V.)


Abstract: Chamazulene (CA) is an intensely blue molecule with a wealth of biological properties. In cosmetics chamazulene is exploited as a natural coloring and soothing agent. CA is unstable and tends to spontaneous degradation accelerated by light. We studied the photodegradation of CA upon controlled exposure to UVB-UVA irradiation by multiple techniques, including GC-MS, UHPLC-PDA-ESI-MS/MS and by direct infusion in ESI-MSⁿ, which were matched to *in silico* mass spectral simulations to identify degradation products. Seven byproducts formed upon UVA exposure for 3 h at 70mW/cm² (blue-to-green color change) were identified, including CA dimers and CA benzenoid which were not found on extended 6 h irradiation (green-to-yellow fading). Photostability tests with reduced irradiance conducted in various solvents in the presence/absence of air indicated highest degradation in acetonitrile in the presence of oxygen, suggesting a photo-oxidative mechanism. Testing in the presence of antioxidants (tocopherol, ascorbyl palmitate, hydroxytyrosol, bakuchiol, γ -terpinene, TEMPO and their combinations) indicated highest protection by tocopherol and TEMPO. Sunscreens ethylhexyl methoxycinnamate and particularly Tinosorb® S (but not octocrylene) showed good CA photoprotection. Thermal stability tests indicated no degradation of CA in acetonitrile at 50°C in the dark for 50 days; however, accelerated degradation occurred in the presence of ascorbyl palmitate.

Keywords: artemisia arborescence; blue color; analysis; UHPLC-MS/MS; GC-MS; photodegradation; antioxidant; sunscreen

1. Introduction

Chamazulene, CA (1,4-dimethyl-7-ethylazulene) is a bicyclic unsaturated hydrocarbon with the molecular formula C₁₄H₁₆. It is found in the essential oils (EOs) from a variety of plants [1–4], notably german chamomile (*Matricaria recutita*, L.) [5], wormwood (*Artemisia absinthium*, L.) [6], and yarrow (*Achillea millefolium*, L.) [7]. CA is a blue-violet derivative of azulene formed by the thermal decomposition of colorless sesquiterpene matricin during the steam distillation process [8] (Scheme 1.). Matricin is a genuine compound of e.g. chamomile that is unstable under acid conditions. Upon steam distillation, it is readily converted into chamazulene carboxylic acid and further on into chamazulene. Its deep blue color is a main feature determining its interest and the value of EOs containing it. Besides being used as a natural colorant in cosmetics and pharmaceuticals, chamazulene has anti-inflammatory activity [1], has proven to slow down the oxidation of cumene [9], and, it has been used to protect human dermal fibroblast from oxidation by reactive oxygen species (ROS), specifically by boosting the biosynthesis of antioxidant enzymes, which would breakdown ROS present in the medium [10]. Hence, CA has indirect antioxidant properties, beside showing reactivity toward different radicals [11]. EOs are well-perceived natural sources of bioactives of widespread use in cosmetics and food products [12,13], and CA, has gained popularity

as an ingredient in cosmetic skin- and hair-care products, also owing to its reported photoprotective activity, able to prevent UVB-induced photodamage [14].



Scheme 1. Decomposition of matricin to CA via chamazulene carboxylic acid.

Paradoxically, this last property might actually turn into a matter of concern. While people using these products are unavoidably exposed to sunlight, it is not known whether the use of cosmetics containing CA with concomitant exposure to sunlight results in any adverse effects. This made it vital to carry out an investigation of the photostability of CA. Beside its toxicological implications, the fading away of natural organic dyes such as azulene derivatives on exposure to ambient or normal storage conditions is a main issue limiting their usefulness, and the stability of cosmetics and health-related products [15], as degradation intermediates could also trigger the oxidative damage of other sensitive components, such as lipids [16].

Although there are studies regarding the (limited) stability of related guaiazulene in cosmetic formulations, and under conditions of exposure to light and high temperatures [15,17], knowledge on CA appears scarce, despite its similarity to guaiazulene, and its importance. In the present study CA was isolated from *Artemisia arborescens* (L.) EO, obtained by hydrodistillation from wild specimens from Sicily (Italy). Its thermal and photochemical stability in the presence and absence of oxygen were investigated, since studies have shown that oxygen is often involved in the instability of organic dyes [18,19]. This work aims at achieving a mechanistic understanding of the photo- and oxidative degradation pathways of CA using kinetic approaches and different analytical techniques to identify byproducts. Since our aim was also to propose prevention strategies, we investigated the use of antioxidants and sunscreens [20] to protect CA under photochemical and thermal stress.

2. Results and Discussion

2.1. CA Isolation and Purification

CA was isolated by normal phase flash chromatography from *Artemisia arborescens* (L.) essential oil. This essential oil was chosen as the most convenient source for three main reasons: (1) it is rich in CA (often > 10%), (2) it is relatively inexpensive because the plant grows spontaneously in many Mediterranean areas and has limited commercial interest, and, finally, (3) it allows an easy isolation because in *A. arborescens* essential oil CA is the most lipophilic component. The sample we used contained 13.03 % CA by GC-MS analysis. Thus, 2.5 g of EO were seeded on flash chromatography silica column and eluted with hexane; the blue band of CA was collected in fractions whose purity was monitored by TLC. A better purity evaluation was performed after collection by GC-MS. The fractions with GC-MS peak area larger than 99% were combined and dried to afford CA with a purity of 99.8%, which was used for subsequent studies. Approximately 138 mg of pure CA were obtained with an isolation yield of 5.52% (Figure 1).




Figure 1. Total ion current GC-MS chromatogram of different fractions of *Artemisia arborescens* (L.) essential oil obtained by flash chromatography. Fractions F1 to F31 contained > 99% chamazulene, while the other fractions with variable impurities were discarded.

2.2. Generation of Photo-Degradation Products of CA

Our first aim was to identify the products formed from CA photodegradation, so to achieve understanding of the process. Acetonitrile solutions of CA at concentrations of 1000 mg/L were placed into quartz cells closed with a screwcap. The quartz cells were exposed to a W-Hg solar lamp emitting in Vis-UVA-UVB region (see Figure S1 in Appendix), at a distance from the light source to achieve an irradiance of 70 mW/cm² in the UVA (strong photooxidation conditions), and were irradiated until obtaining visible color changes. A main change from deep blue color to pale green was achieved after an exposure time of 3 h, then a complete fading to a yellow solution was obtained with exposure time of 6 h. Therefore, the two solutions photoexposed for 3 h and 6 h were subjected to UHPLC-ESI-MSⁿ analysis. Typical chromatograms are shown in Figure 2.




Figure 2. LC-MS chromatograms of CA in acetonitrile photoexposed: A) green sample (exposure time: 3 h); B) yellow sample (exposure time: 6 h). Chromatographic conditions: LC-MS method A (see Section 3.4).


The green sample, still containing a CA residue, was richer in chromatographic peaks than the yellow sample in which only three had survived and become dominant (Figure 2).

Based on the studies by Matsubara *et. al.* on the degradation of guaiazulene [21], to each peak characterized by a specific *m/z* ratio was assigned a hypothetical chemical structure that was then confirmed through MSⁿ experiments on the isolated parent ions, which were also performed by direct infusion of the sample into the ESI source. To aid the identification the MSⁿ fragmentation tree obtained experimentally was matched with one generated *in silico* using Mass Frontier 5.0, a software based on fragmentation libraries, which can predict the fragmentation pathway of a compound, following different ionization methods [22].

This also provides the possible tandem mass spectrometry (MS/MS) fragment ions of an unknown compound [22]. The *m/z* of the fragment ions generated by the software were compared with the mass spectra of the compound under investigation. This software was utilized to reduce the number of false positives. In addition, a bibliographic review on the possible chemical structures of the degradation products of chamazulene was done, also by referencing to structurally related guaiazulene [17]. This combined approach enabled the identification of a significant number of photodegradation products present in the sample, despite the major complexity. The parent ions (*m/z*) and their products ions obtained by MSⁿ are reported in Table 1. Among those products we could identify with good confidence seven structures which are illustrated in Scheme 2.

Table 1. MSⁿ fragmentation data of chamazulene and its photodegradation products, obtained by collision induced dissociation (CID). RA: Relative abundance.

Peak #	Compound	Retention Time (min)	Parent peak <i>m/z</i> [M + H] ⁺	Main Product ion	RA (%)
2	Oxidized chamazulene	2.88	183	169 [M+H-CH ₂] ⁺	48
				155 [M+H-CH ₂ =CH ₂] ⁺	38
				141 [M+H-C ₃ H ₆] ⁺	35
				197 [M+H-H ₂ O] ⁺	40
3	Chamazulene quinone	3.61	215	187 [M+H-CO] ⁺	100
				173 [M+H-CH ₂ CO] ⁺	33
				159 [M+H-CH ₂ =CH ₂ -CO] ⁺	38
				184 [M+H-CH ₃] ⁺	25
4,5	Chamazulene	3.75	199	171 [M+H-CO] ⁺	100
	Carbaldehyde	3.85		143 [M+H-CH ₂ =CH ₂ -CO] ⁺	78
6	Dihydrochamazulene	4.44	187	172 [M+H-CH ₃] ⁺	15
				159 [M+H-C ₂ H ₄] ⁺	100
				131 [M+H-C ₄ H ₈] ⁺	22
				381 [M + H-H ₂ O] ⁺	100
7	Chamazulene benzenoid	5.79	399	371 [M + H-CO] ⁺	57
				215 [M+H-Chamazulene] ⁺	10
				199 [M+H-CH ₃ Chamazulene] ⁺	63
				169 [M + H-CH ₄] ⁺	100
8	Chamazulene	7.89	185	129 [M + H-CH ₂ =CHCH ₂ CH ₃] ⁺	10
		13.06		366 [M + H-CH ₃] ⁺	38
9	Methylene dimers of chamazulene	13.06	381	363 [M + H-H ₂ O] ⁺	100
				353 [M + H-CH ₂ =CH ₂] ⁺	22
				337 [M + H-CH ₂ =CHCH ₃] ⁺	30
				321 [M + H-C ₆ H ₆] ⁺	33
10	Chamazulene dimers	14.90	367	352 [M + H-CH ₃] ⁺	60
		16.51		338 [M + H-CH ₃ CH ₂] ⁺	100
		17.05		309 [M + H-C ₄ H ₁₀] ⁺	25
		18.28			



Scheme 2. Reaction pathways for the formation of photo-oxidative products of CA. 1: CA dimer, 2: Methylene dimer of CA, 3: Dihydrochamazulene, 4: Oxidized CA, 5: CA carbaldehyde, 6: CA quinone and 7: CA benzenoid.

With reference to Scheme 2, the LC-MS analysis of the sample exposed to UV radiation for 3 hours revealed the presence of dimers (1 and 2). Instead, in the sample solution irradiated for 6 hours dimers were not found, based on the mass spectra. These products are formed when hydrogen is abstracted from chamazulene, giving rise to a radical that further combines with itself to form various dimers of chamazulene. Similar outcomes were obtained from studies investigating the photochemical degradation products of guaiazulene and azulene under UV light [17,23]. Beside dimers, more complex products, including oxygenated trimers, tetramers, and even oligomers, were formed [17]. Also, it was found that the formation of 3,3-biguaiazulene (a dimer) in solution resulted in a color change from blue to green color [23]. Accordingly, after 6 h irradiation of our samples, when the color turned from green to yellow, dimers were no longer present, which indicates that the dimers formed in our sample were responsible for the green color. Therefore, further exposure to UV irradiation leads to the decomposition of the dimers to form other products. Figures 3 and 4 represent the mass spectra of CA dimer and the methylene dimer of CA respectively, acquired in tandem mass mode.




Figure 3. Mass spectra (MS/MS) showing the fragment ions of chamazulene dimer (1), parent ion $[M+H]^+$ m/z 367 in tandem mass mode, by collision induced dissociation (CID).




Figure 4. Mass spectra (MS/MS) showing the fragment ions of the methylene dimer of chamazulene (2), parent ion $[M+H]^+$ m/z 381 acquired in tandem mass mode, by CID.

Degradation products such as dihydrochamazulene (m/z 186) and oxidized CA (m/z 182), as seen in scheme 1, result from a disproportionation reaction of CA (m/z 184). They differ from each other by 2 mass units, which asserts the reduction of chamazulene (by the addition of 2 hydrogen atoms) for dihydrochamazulene and, on the other hand, the oxidation of chamazulene (by the loss of 2 hydrogen atoms). The ability of CA to undergo hydrogen abstraction by radical species has been reported in the literature, as the suggested mechanism for its antioxidant activity [11]. One possible explanation is that excited CA serves as H-abstrating species towards ground-state CA triggering the disproportionation. On the other hand, it was demonstrated by EPR spectroscopy that UVA irradiation of azulene and guaiazulene in the presence of oxygen forms superoxide radical ($O_2^{\cdot-}$) by electron transfer from the photoexcited azulene [24]. Likely, a subsequent proton transfer (ET-PT sequence) or a concerted PCET (proton-coupled-electron-transfer) to O_2 [25,26] would afford the hydroperoxyl radical $HOO\cdot$ (the neutral form of superoxide) and the C-catered radical of CA, which can decay by disproportionation. This last mechanism calls for the key role of oxygen in the photodegradation of CA. Figures 5 and 6 represent the mass spectra of oxidized CA and dihydrochamazulene respectively, acquired in tandem mass mode.




Figure 5. Mass spectra (MS/MS) showing the fragment ions of oxidized chamazulene (3) parent ion $[M+H]^+$ with m/z 183 acquired in tandem mass mode, by CID.




Figure 6. Mass spectra (MS/MS) showing the fragment ions of dihydrochamazulene (4), parent ion $[M+H]^+$ with m/z 187 acquired in tandem mass mode, by CID.

Another photodegradation product, CA carbaldehyde, is most probably formed from the reaction of molecular oxygen with a chamazulene (carbon-centered) radical to give rise to a peroxy radical intermediate ($ROO\bullet$, Scheme 1), which subsequently affords a sidechain oxidation, forming the aldehyde derivative of chamazulene. This manner of reaction is similar to the well accepted mechanism of autooxidation of the sidechain in alkylbenzenes and alkyl naphthalenes [23,27,28]. The peroxy radical is quite stable, yet highly reactive toward unsaturated lipids: when formed in a cosmetic product or other man-directed preparations, it can trigger the oxidation of other ingredients, causing the formation of toxic byproducts [16], and the formulation to lose its stability and change its properties. Additionally, Matsubara *et al.* [23] and Fiori *et al.* [15,17] both studied the photooxidation of alkyl azulenes in the presence and absence of oxygen and concluded that oxygen plays a vital role in the degradation of azulenic compounds. Figure 7 represents the mass spectrum of chamazulene carbaldehyde, acquired in tandem mass mode.




Figure 7. MS/MS spectrum showing the fragment ions of chamazulene carbaldehyde (5), parent ion $[M+H]^+$ with m/z 199 acquired in tandem mass mode, by CID.

CA quinone and CA benzenoid (Scheme 2) are possibly formed through the alkoxyl radical intermediate, likely arising from hydroperoxyl ($\text{HOO}\bullet$) or alkylperoxyl ($\text{ROO}\bullet$, e.g. guaiazuleneperoxyl) radical addition to guaiazulene followed by fragmentation. Rearrangement of the conjugated structure can lead to the quinone end products. In the case of the CA benzenoid, there is a complete modification of one azulene ring into a benzenoid. It was reported that the formation of degradation products that possess a modified azulene chromophore induced a color change in the sample solution [23]. This is because, as the azulene chromophore is modified, its spectrum of absorption in the visible would clearly change. The photochemical degradation pathway of CA resulting in quinone formation holds significant implications for biological systems, depending on the site of exposure. At the skin level, similar byproducts of azulene and guaiazulene have been shown to cause various deleterious effects, including carcinogenicity, skin inflammation, and dermatitis [29,30]. Quinones can interfere with DNA synthesis, disrupting the normal transcription of information and consequently leading to the generation of mutated cells [31]. Figure 8 represents the mass spectra of CA quinone, acquired in tandem mass mode.




Figure 8. MS/MS spectrum showing the fragment ions of chamazulene quinone (6) parent ion $[M+H]^+$ m/z 215 acquired in tandem mass mode by collision induced dissociation.

The fragmentation pathway of parent ion at m/z 215 obtained with Mass Frontier software is shown in Figure S2 (Appendix).

Figure 9 represents the mass spectra of chamazulene benzenoid, acquired in tandem mass mode.




Figure 9. MS/MS spectrum showing the fragmentation of chamazulene benzenoid (7) parent ion $[M+H]^+$ m/z 399 in tandem mass mode, by collision induced dissociation.

2.3. Photostability Test

In order to evaluate the photostability of CA in solution under conditions that would be representative of the behavior in cosmetic products, we set to operate according to ICH guidelines, option 1 [32], under weak photoradiation conditions. Diluted (0.17 mM) solutions of CA were placed in closed quartz tubes at a distance from the light source to achieve an irradiance of 2.4 mW/cm² in the UVA region. All experiments were performed by keeping constant temperature (25°C) and irradiance, as summarized in Table 2. Chamazulene concentration was monitored at regular time intervals (15, 30, 60, 120, 180, 240 min. corresponding at UVA irradiation doses of 2.2, 4.3, 8.6, 17.3, 25.9 and 34.6 Jcm⁻²) during 4 hours of irradiation. Analysis was performed by GC-MS and by LC- PDA, upon setting suitable rapid methods allowing to monitor CA decay (Section 3.6 and Figures S3–S6 in Appendix). Concurrently, identical analyses were performed on samples stored in the dark (vials wrapped in aluminum foils), which served as controls. Otherways identical sets of experiments were repeated by changing critical parameters so to evaluate their influence on CA decay kinetics. Thus 3 solvents were comparatively tested: apolar hexane, protic methanol, and polar acetonitrile, which was our reference solvent throughout the study. To evaluate the role of oxygen, samples were degassed by purging with nitrogen or left saturated with air with a variable volume of head-space in the sealed tube.

Table 2. Radiometric values used for all samples subjected to photostability tests (T = 25°C).

Type of UV	Wavelength [nm]	Radiance [mWcm ⁻² sr ⁻¹]	Irradiance [mWcm ⁻²]	Energy absorbed	Energy absorbed
				[Jcm ⁻²] (t = 15 min)	[Jcm ⁻²] (t = 4 hrs.)
UVA	320 - 400	35.2	2.4	2.2	34.6
UVB	290 – 320	7.5	0.5	0.5	8

Under identical experimental conditions, we also comparatively tested acetonitrile solutions of CA containing antioxidants (tocopherol, ascorbyl palmitate, TEMPO, hydroxytyrosol, bakuchiol, γ -terpinene) at concentrations up to 10-folds that of chamazulene (1.7 mM) and sunscreens at 5% by weight, which corresponds to 0.2 mM, for octocrylene and octyl methoxycinnamate while it is 0.08 mM for and Tinosorb® S. These tests were meant at evaluating the protection that might be offered by other components of the cosmetic formulation [33] or the effectiveness of additives in slowing down photodegradation.

2.3.1. Effect of Solvent and Oxygen

Solutions of 0.17 mM CA in acetonitrile, methanol and *n*-hexane placed in a quartz EPR tube and photo-irradiated as reported in Table 2, showed different rates of degradation, highlighting a mechanism linked to the polarity of the solvent (Table 3). Indeed, in protic methanol the rate of decay was twice as fast as in *n*-hexane, while it grew to about 6-folds faster in polar acetonitrile. Furthermore, it was sensitive to the head space (HS) volume of air in the tube, as can be judged from experiments in *n*-hexane by carrying out the irradiation leaving a variable HS. The importance of oxygen is also highlighted by the much-reduced degradation recorded in N₂ purged samples in any solvent. Indeed, while degradation became negligible in *n*-hexane, it reduced the rate by about 3-folds in methanol and acetonitrile (Table 3).

Table 3. Kinetics of CA photodegradation under irradiation conditions detailed in Table 2, as a function of the solvent and the degree of air exposure and head-space volume (HS). Regressions are expressed as residual CA% (y) as a function of time in minutes (x).

Air saturated			
	HS 0 cm ³	HS 0.17 cm ³	HS 0.34 cm ³
<i>n</i> -Hexane	y = -0.0551x + 99.638 R ² = 0.9368	y = -0.0654x + 100.94 R ² = 0.9958	y = -0.0736x + 101.04 R ² = 0.9692
Methanol	y = -0.1107x + 101.17 R ² = 0.9659		
Acetonitrile	y = -0.3173x + 102.56 R ² = 0.9811		
N ₂ purged			
	HS 0 cm ³		
<i>n</i> -Hexane	y = -0.0009x + 100.72 R ² = 0.0033		
Methanol	y = -0.0417x + 101.75 R ² = 0.7921		
Acetonitrile	y = -0.1146x + 103.17 R ² = 0.8943		

Our results are in good agreement with previous studies on the photodegradation of azulene and guaiazulene by Matsubara *et al.* [23] and Fiori *et al.* [15,17] and strongly point toward a photo-oxidative path as the largely prevailing mechanism. As a further confirmation of the photooxidative mechanism, a 500 mg/L solution of CA in acetonitrile was placed in a 3.5 mL Teflon sealed quartz cuvette and degassed for 5 minutes with an abundant N₂ stream. In this way it is possible to guarantee better degassing efficiency and complete air tightness. The cuvette was irradiated at 70 mW/cm² (strong photooxidation conditions) for 6 h. At regular time intervals a spectrophotometric reading was performed without opening the cuvette. Results shown in Figure S7 (Appendix) indicate a further 2-fold reduction of the rate of CA degradation compared to results in Table 3, despite the much harsher irradiation conditions.

2.3.2. Effect of Antioxidants and Sunscreen

Since the photo-oxidative mechanism resulted dominant on the basis of previous experiments, we set to evaluate the protection offered by antioxidants and sunscreens. The experiments were performed in acetonitrile, since it was the solvent affording the most marked and rapid photodegradation of CA, and solutions were saturated with air so to simulate normal environmental conditions. Irradiance was set as in Table 2., both in the absence and in the presence of antioxidants and UV filters, and photodegradation was comparatively assayed by LC-PDA monitoring of CA concentration at 349 nm upon calibration (see Appendix). As shown in Table 4, in the absence of other additives CA was completely degraded at the UVA irradiation dose of 35 J/cm².

Table 4. Comparative rate of photodegradation of CA 500 mg/L in acetonitrile (100%) irradiated at 2.4 mW/cm² at 25°C, alone or in the presence of different antioxidants or of different sunscreens. The time-course is expressed as cumulative irradiation dose.

UV irradiated solutions of CA	Percentage of CA after UVA irradiation						
	0 J/cm ²	2 J/cm ²	4 J/cm ²	8 J/cm ²	17 J/cm ²	25 J/cm ²	35 J/cm ²
CA	100	98±1	92±2	81±2	52±3	22±1	0
MIXED WITH ANTIOXIDANTS							
CA + TEMPO (1:10)	100	96±3	90±2	83±4	73±3	53±1	33±1
CA + Tocopherol (1:10)	100	94±2	88±2	77±1	65±2	43±1	27±1
CA + Ascorbyl palmitate (1:10)	100	95±2	94±4	71±3	49±3	2±1	0
CA + Bakuchiol (1:10)	100	94±3	91±3	62±3	11±3	0	0
CA + Hydroxytyrosol (1:10)	100	94±3	92±3	80±2	62±2	37±1	6±1
CA + Tocopherol + Bakuchiol (1:1:1)	100	94±1	92±1	77±3	41±1	10±1	0
CA + Ascorbyl palmitate + Tocopherol (1:1:1)	100	91±2	80±2	68±3	17±1	0	0
CA + Tocopherol + Hydroxytyrosol (1:1:1)	100	93±2	81±4	72±3	37±2	8±3	0
CA + Tocopherol + TEMPO (1:1:1)	100	96±3	90±2	76±1	19±1	0	0
CA + Ascorbyl palmitate + TEMPO (1:1:1)	100	96±3	90±2	75±2	17±1	0	0
CA + γ-terpinene + Ascorbyl palmitate (1:5:1)	100	94±2	92±3	76±2	28±4	0	0
CA + γ-terpinene + Hydroxytyrosol+ (1:5:1)	100	96±2	90±3	78±1	38±2	0	0
CA + γ-terpinene + TEMPO+ (1:5:1)	100	93±1	93±2	80±2	57±2	24±1	6±1
MIXED WITH UV FILTERS							
CA + Tinosorb® S (5%)	100	92±2	92±3	86±2	73±4	65±2	48±1
CA + Octocrylene (5%)	100	98±3	90±2	79±3	57±1	22±1	0
CA + Octyl methoxycinnamate (5%)	100	100±1	99±3	97±3	70±3	45±2	20±2
CA in the dark (negative control)	100	100±2	99±2	99±2	100±1	99±3	100±2

The selection of antioxidants deserves discussion. Our aim was to evaluate structurally different molecules, of potential use in cosmetic formulas, and possibly boosting different mechanisms. Beside α-tocopherol and ascorbyl palmitate, which are well-established highly effective chain-breaking antioxidants of major use in cosmetics [16,33], we tested bakuchiol, a potent natural phenolic antioxidant which is receiving major attention as a cosmetic ingredient [34], and hydroxytyrosol, another natural cosmetic-grade and food-grade antioxidant which represents the catechol structural family [35]. In addition, we tested γ-terpinene, a terpene component of several essential oils that demonstrated an unconventional mechanism of inhibition, different from typical chain-breaking antioxidants, based on increasing the rate of decay of transient radical species [36,37]. TEMPO (2,2,6,6-tetramethylpiperidine-*N*-oxyl) radical was also included although it cannot be used as a cosmetic ingredient; however, it was used as a research molecule, as it boosts a unique antioxidant mechanism based on acting as a catalytic antioxidant in the presence of hydroperoxyl radicals [38] – these are possibly formed during the photodegradation of CA, as previously discussed.

Since antioxidants normally offer the best protection within synergic co-antioxidant mixtures [39,40], we also tested mixtures of the above antioxidants including combinations that had previously demonstrated to afford synergic activity [16,37,38]. All experiments are summarized in Table 4.

Disappointingly most antioxidants and their combinations offered negligible protection to CA, underlying a clear difference between protecting a lipid from peroxidation radical chain and protecting an excited chromophore from photo-oxidative degradation under continuous irradiation. Indeed, some additives actually appeared to slightly accelerate CA loss, likely due to photodegradation of the antioxidant itself, which would form radicals that might attack CA.

However, some combination afforded successful protection, slowing down the degradation of chamazulene to a significant extent. These are visually compared in Figure 10. Both hydroxytyrosol and even more so α-tocopherol effectively protected CA; however, the best protection was offered by TEMPO. Since it has been demonstrated that the antioxidant behavior of TEMPO is activated by hydroperoxyl radicals (HOO•) formed during the oxidation process, this intriguing observation

apparently supports the mechanism we proposed to explain the formation of key degradation products like 3 and 4 (see Scheme 2 and Section 2.2.).




Figure 10. Photostability of CA (0.17 mM) in the presence of antioxidants (1.7 mM).

Experiments in the presence of sunscreens also offered interesting observations (Table 4). Three molecules were selected on the basis of their frequent occurrence in cosmetic products [41,42], and were comparatively tested: Tinosorb® S, a broad spectrum UVB-UVA filter (290-370 nm) with maxima of absorption at 310 and 350 nm, octocrylene, a UVB filter reaching the short-wavelength UVA region (280-330 nm, λ_{\max} 307 nm), and octyl methoxycinnamate (OMC) with similar coverage (280-330 nm, λ_{\max} 311 nm) [43–45].

Both Tinosorb® S and OMC were able to effectively protect CA from photodegradation as can be seen in Figure 11. Tinosorb® S was overall the most effective, sparing as much as 48% CA at the end of the experiment after prolonged exposure (4h) to a cumulative irradiation dose of 35 J/cm² in the UVA and 8 J/cm² in the UVB. Nonetheless, OMC offered perhaps the most interesting result, as it completely blocked the loss of chamazulene for a lag time of about one hour, corresponding to an irradiation dose of about 10 J/cm². Such dose corresponds to one-two hours of exposure at noon of sunny days during the summer around the world, based upon observations of UVA intensity of 2.1 mW/cm² in Okayama, Japan in September [46], and 3.6 mW/cm² in Jackson (MS), USA in August [47]. However, following such lag time the photodegradation started and proceeded at the same rate as the unprotected sample. This can tentatively be explained by considering that OMC itself has limited photostability [48] and it undergoes photodegradation when exposed to a solar lamp [49]; therefore, the end of the observed lag time could be attributed to its substantial photodegradation. Somewhat surprising was instead the lack of protection demonstrated by octocrylene, despite an absorption spectral range similar to OMC. Since also octocrylene is known to undergo degradation on exposure to UVA-UVB irradiation [50], possibly a similar explanation can be put forward. Clearly, further studies monitoring the time course of OMC and octocrylene during irradiation would be needed to confirm this hypothesis.




Figure 11. Photostability of chamazulene (0.17 mM) in the presence of UV filters at 5% concentration.

The efficacy exhibited by some antioxidants and UV filters in augmenting the photostability of chamazulene, at a concentration of typical use in cosmetics - both sunscreens are allowed at up to 10% (w/w), while antioxidants hydroxytyrosol and tocopherol are not limited [51] - signifies their potential application in formulations containing chamazulene to prevent its photooxidation when exposed to UV light, thus contributing to the overall stability and safety of the cosmetic formulation.

2.4. Thermal Stability Of Chamazulene

In order to test the thermal stability of CA under conditions compatible with the storage and use of cosmetics, in the absence of direct light exposure, sample solutions of CA in acetonitrile were stored in the refrigerator at 4 °C, at room temperature, and in the oven at 50°C, all in the dark, for 50 days. Upon monitoring the concentration of CA over this period, it was observed that it remained stable throughout. Therefore, the stability of CA was not affected by the above-mentioned conditions. This is quite intriguing since previous evidence suggests that cosmetic formulations containing essential oils rich in chamazulene tend to undergo color changes with prolonged storage in the dark at room temperature, implying that other processes, not related to photodegradation would limit its stability. One possible explanation is that other labile components of the formulation might undergo oxidative degradation not triggered by light, and the radical or other intermediates formed during their oxidation or thermal degradation might attack CA, causing its degradation. As a proof of concept, we repeated the studies by monitoring the stability of CA in solution at different temperatures in the presence of antioxidants, which, by their nature, are very sensitive to oxidative thermal degradation. Results summarized in Figure 12 show that, while CA alone was perfectly stable even after prolonged storage at 50°C, the presence of tocopherol and, particularly, of ascorbyl palmitate accelerated its decay significantly, while other antioxidants did not produce a significant action in this regard, as shown in detail in Figure S8 (Appendix). This apparently paradoxical behavior is likely due to the high persistence of radicals formed by these two antioxidants [16], which reside in the sample sufficiently long to cause antioxidant-mediated autoxidation of CA. This parallels the well-known phenomenon of TMP, tocopherol-mediated peroxidation of human LDL caused by tocopherol, under some experimental settings [52]. As also discussed for TMP, the phenomenon can be completely abolished by using synergic mixtures of antioxidants in place of a

single molecule. An issue that would require further investigation in cosmetic formulations, to assess its relevance and ways for prevention.




Figure 12. Thermal degradation of chamazulene in solution, alone or in the presence of antioxidants, incubated at 50°C.

3. Materials and Methods

3.1. Materials

HPLC grade water, methanol and acetonitrile, formic acid and hexane (GC grade) were from Merk (Milan, Italy). Ascorbyl palmitate (ACEF, Piacenza, Italy), α -tocopherol (Artchem, Milan, Italy), hydroxytyrosol (99%, Pioneer-biotech, Xian, China), bakuchiol (99%, Carlo Sessa S.p.A., Milan, Italy), (2,2,6,6-Tetramethylpiperidin-1-yl)oxyl (TEMPO) (98%, Sigma-Aldrich), octyl methoxycinnamate (98%, Sigma-Aldrich, also called ethylhexyl methoxycinnamate), bis-ethylhexyloxyphenol methoxyphenyl triazine (Tinosorb® S, 99%, Eurotrading, Padova, Italy), octocrylene (Uvinul N539t, 98%, Eurotrading, Padova, Italy) were used as received. *Artemisia arborescens* (L.) essential oil was custom extracted by steam distillation by Marcello Militello (Ph.D. Agronomist), Regional Institute for Floriculture (IRF) (SanRemo, IM, Italy), in collaboration with “azienda agricola l’essenza degli iblei” (Canicattini Bagni, SR, Italy).

3.2. Isolation of Chamazulene (CA)

Chamazulene (CA), was isolated by flash chromatography from *Artemisia arborescens* (L.) essential oil, which contained 13.03% CA by GC-MS. 2.5 mL of EO were seeded in a flash column prepared with a borosilicate Glass Heavy-Wall Chromatography column (18 mm i.d x 400 mm length, Macherey-Nagel) loaded with silica gel (spherical, 60 μ m, Merck) eluted with *n*-hexane via a manual eluent pumping system. Elution progress was monitored by TLC on glass plates 50 mm x 100 mm (Sigma Aldrich). After elution, fractions having > 99% purity based on GC-MS peak area were combined and evaporated under vacuum to yield 138 mg of CA (purity 99.8% by GC-MS) with an isolation yield of 5.52%.

3.3. Photolysis Experiments: Equipment and General Procedures

The photochemical behavior of CA was investigated by using a custom made UV light box with a combined mercury/tungsten light source emitting in the UV-VIS (Osram Ultra-Vitalux®, Germany). The lamp produces a mix of radiation very similar to that of natural sunlight. This blend of radiation is generated by a high-pressure mercury discharge tube (ultraviolet emission) and a tungsten

filament (infrared and visible light emission). Compliant with the technical specification of the lamp, the distance from the sample for photostability experiment was regulated to obtain an irradiance of 2.4 mW/cm² in the UV-A and of 0.5 mW/cm² in the UV-B region; this irradiance is of the same magnitude as the UV exposure from solar irradiation in a summer sunny day at noon across the world (based upon reports of UVA intensity of 2.1 mW/cm² in Okayama, Japan in September [46], and 3.6 mW/cm² in Jackson (MS), USA in August [47]). In the preliminary studies to identify CA fragmentation products the irradiance was raised by shortening the distance of the sample to the lamp so to achieve about 70 mW/cm² in the UVA. Samples consisting of solutions of CA in acetonitrile, or in *n*-hexane, or in methanol, in the absence or presence of other additives, were placed in Suprasil® quartz EPR tubes (4 mm ID) and sealed. When needed degassing was performed by bubbling N₂ for 2 min via a glass capillary. Irradiated samples were then subjected to GC-MS, UHPLC-PDA or UHPLC-ESI-MSⁿ analysis as appropriate.

3.4. GC-MS Analysis

Gas chromatography/mass spectrometry (GC-MS) analyses were performed on a TRACE GC 1310 Series (Thermo Fisher Scientific) gas chromatograph equipped with a split-splitless injector and interfaced with a ISQ QD (Thermo Fisher Scientific) mass detector with single quadrupole analyzer, operating in electron impact (EI) mode. The GC column was a Phenomenex ZB-5 fused-silica capillary column (30 m, 0.25 mm i.d., 0.25 µm film thickness), consisting of crossbond 5% diphenyl, 95% dimethyl polysiloxane. Injection was performed by CombiPal autosampler (CTC Analytics, Zwingen, Switzerland). Helium (He) was the carrier gas at a flow rate of 1.2 mL/min. Two distinct temperature programming were used, the first (method A) used to verify the purity of fractions from flash chromatography and the second (method B) used for rapid analysis of the CA during photostability experiments. Method A: initial temp. 50°C, held for 2.5 minutes, then increased at 2°C/min to 220°C, for a total run time of 85 minutes. Method B: initial temp. 120°C ramped by 5°C/min to 195°C, for a total run time of 10 minutes. The temperatures of the injector base, the transfer line and the ionization source were maintained at 250, 280 and 300°C, respectively. The mass spectra were recorded in full scan (40–650 amu) to collect the total ion current (TIC) chromatograms. Calibration for CA is shown in Figure S3 (Appendix).

3.5. UPLC-ESI-MSⁿ, ESI-MSⁿ Analysis

Ultra-high-performance liquid chromatography (UHPLC) analyses were carried out on a ACCELA liquid chromatograph interfaced with a Photo Diode Array (PDA) detector, with an autosampler and with a LCQ FLEET mass spectrometer (Thermo Fisher Scientific). The mass spectrometer was equipped with an electron spray ionization (ESI) source, and with an ion-trap analyzer. The ESI system, operating in positive mode, was set at 6.0 kV and 8V for spray and capillary voltage, respectively; the heated capillary was at 300°C. The sheath gas and the auxiliary gas (nitrogen) flow rates were set to 25 and 3 (arbitrary unit), respectively. The ESI process was optimized using pure isolated CA as reference compound. The mass chromatograms were acquired in total ion current (TIC) modality from 50 to 2000 *m/z* and in MS/MS mode (collision induced dissociation, CID 25, arbitrary) on the protonated molecule of CA and its photoproducts. To identify CA photodegradation products, photo-exposed samples were also infused directly into the ESI interface via 500 µL syringe (Hamilton) and a syringe pump device. The parent ion produced in MS was isolated in the ion-trap (MS²) subjected to CID at 25 AU and fragments analyzed (MS³). The HighChem Mass Frontier 5.0 software was used to generate *in silico* MS/MS fragmentation of some hypothesized photodegraded products of CA.

Reversed-phase chromatographic analyses were performed on a Kinetex PFP (Phenomenex), 100 mm x 3.0 mm i.d. x 2.6 µm column. For the initial screening of CA degradation products the following program was used (Method A): flow rate 300 µL/min; injection volume 5 µL; mobile phase was 0.1% aqueous formic acid (solvent A) and 100% acetonitrile (solvent B); linear gradient profile (A:B) started at 40:60, kept for 4 min, then linearly decreased to 5:95 in 30 minutes, kept constant for 5 minutes and finally re-equilibrated at 40:60 in 5 minutes. Total run time was 40 minutes.

3.6. LC-PDA Analysis

LC- PDA mode was operated to rapidly quantify CA during the course of photolysis kinetics experiments. The apparatus is described in Section 3.5. Spectrum of acquisition was from 250 to 800 nm and the wavelength of 349 nm was selected to monitor CA. Chromatographic separation was achieved with a faster elution program (Method B): flow rate 600 $\mu\text{L}/\text{min}$; injection 2 μL ; mobile phase consisted of 0.1% formic acid aqueous (solvent A) and 100% acetonitrile (solvent B). Linear gradient profile (A:B) started at 40:60 kept constant for 2 minutes, then linearly decreased to 30:70 in 2 minutes, then linearly decreased up to 5:95 in 1 minute, kept constant for 2 minutes and finally linearly increased up to 40:60 in 2 minutes. Total run time was 10 minutes. Calibration for CA is shown in Figure S4 and typical chromatograms in Figures S5 and S6 (Appendix).

3.7. UV-Vis Spectrophotometry

Kinetics of CA degradation during irradiation in the absence of oxygen were performed with a double beam UV/Visible spectrophotometer Lambda 20 (PerkinElmer) Full scan spectra were recorded every 60 min in a wavelength range 200 to 800 nm. Sample was contained in 3.5 mL quartz cuvette, degassed by bubbling N_2 for 5 min and sealed with a Teflon stopper.

3.8. Statistical Analysis

All experiments were performed in duplicate and reported as average \pm SD.

4. Conclusions

Chamazulene undergoes extensive degradation when exposed to solar light, with changes in the distinctive color and, conceivably, in the biological properties. Our multi-technique, analytical approach combining GC-MS, UHPLC-PDA-ESI-MS/MS and direct infusion ESI-MSⁿ matched to *in silico* mass spectral simulations, allowed to identify 7 degradation products upon exposure of a CA solution to intense (70 mW/cm^2 UVA) irradiation for 3 h corresponding to a blue-to-green color change. Among them, CA dimers and CA benzenoid are likely responsible for the green color development, since they disappeared upon more extended irradiation (6 h), corresponding to color fading to yellow. This knowledge is relevant in cosmetics science where CA is gaining importance as natural coloring agent. Photostability tests at reduced irradiance under different settings, and in the presence of antioxidants or sunscreens as potential protective additives, offered relevant insights in the process. Solutions in protic methanol degraded more rapidly than in *n*-hexane, and polar acetonitrile showed the fastest decay of CA. Additionally, the rate of degradation was highly sensitive to air, becoming much-reduced to negligible in degassed samples. This indicates a photo-oxidative process as the leading mechanism, likely involving polar transient species including the hydroperoxyl/superoxide ($\text{HOO}\bullet/\text{O}_2^{\bullet-}$) radical. Consistent with this conclusion, CA could be significantly spared by antioxidants like hydroxytyrosol and α -tocopherol, yet particularly by TEMPO, which specifically acts by exploiting the formation of $\text{HOO}\bullet$ radicals [18]. Sunscreens of common use in cosmetics [42] i.e. octyl methoxycinnamate and particularly Tinosorb® S could also affectively spare CA photodegradation. However, other tested antioxidants (bakuchiol, ascorbyl palmitate, γ -terpinene) and their combination, and other tested sunscreens (octocrylene) were ineffective, likely due to limited stability under the photolysis test conditions – an hypothesis that deserves further investigation. Surprisingly, CA was perfectly stable in the dark at 50°C, showing however reduced stability upon incubation with ascorbyl palmitate, which implies that the purported limited stability in formulated products (*e.g.* cosmetics) is likely due to facile reaction with byproducts formed upon decomposition of other components. Although none of the tested protection strategies alone afforded complete inhibition of CA photodegradation, we believe that current results provide a solid basis for successful developments in that direction.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Figures S1–S8: spectral distribution of the solar lamp,

fragmentation of ion m/z 215, calibration curves for GC-MS and LC-PDA analysis, example of chromatograms decay of CA irradiated in the absence of oxygen and in the dark in the presence of antioxidants.

Author Contributions: Conceptualization, L.V.; methodology, S.G.; validation, R.M., formal analysis, S.G., J.N.N.; investigation, J.N.N., S.G. and R.M.; resources, L.V. and S.G.; writing—original draft, S.G.; writing—review & editing, L.V.; project administration, L.V. All authors have read and agreed to the published version of the manuscript.

Funding: Funded by the European Union - NextGenerationEU under the National Recovery and Resilience Plan (PNRR) - Mission 4 Education and research - Component 2 From research to business - Investment 1.1 Notice Prin 2022 - DD N. 104 del 2/2/2022, from PRIN20227XZKBY - Superoxide responsive redox-active systems and nano smart materials to target ferroptosis – FEROX - CUP J53D23008550006". This research was also funded by BeC s.r.l. Forlì (Italy).

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Sample Availability: All compounds are commercially available or are available on request from the corresponding author.

Reference

1. Slon, E.; Slon, B.; Kowalczyk, D. Azulene and Its Derivatives as Potential Compounds in the Therapy of Dermatological and Anticancer Diseases: New Perspectives against the Backdrop of Current Research. *Molecules* **2024**, *29*, 2020. <https://doi.org/10.3390/molecules29092020>
2. Xu, Y.; Wang, Q.; Bao, W.; Pa, B. Antihyperlipidemic Effect, Identification and Isolation of the Lipophilic Components from *Artemisia integrifolia*. *Molecules* **2019**, *24*, 725. <https://doi.org/10.3390/molecules24040725>.
3. Wang, Y.-F.; Zheng, Y.; Feng, Y.; Chen, H.; Dai, S.-X.; Wang, Y.; Xu, M. Comparative Analysis of Active Ingredients and Potential Bioactivities of Essential Oils from *Artemisia argyi* and *A. verlotorum*. *Molecules* **2023**, *28*, 3927. <https://doi.org/10.3390/molecules28093927>
4. De, J.; Lu, Y.; Ling, L.; Peng, N.; Zhong, Y. Essential Oil Composition and Bioactivities of *Waldheimia glabra* (Asteraceae) from Qinghai-Tibet Plateau. *Molecules* **2017**, *22*, 460. <https://doi.org/10.3390/molecules22030460>
5. Qasem, A.; Assaggaf, H.; Montesano, D.; Khalil, Z.; Al-Mijalli, S.H.; Baaboua, A.E.; El Omari, N.; El Menyiy, N.; Bakrim, S.; Sheikh, R.A.; et al. Determination of Chemical Compounds and Investigation of Biological Properties of *Matricaria chamomilla* Essential Oils, Honey, and Their Mixture. *Molecules* **2022**, *27*, 5850. <https://doi.org/10.3390/molecules27185850>
6. Mohammed, H.A. Phytochemical Analysis, Antioxidant Potential, and Cytotoxicity Evaluation of Traditionally Used *Artemisia absinthium* L. (Wormwood) Growing in the Central Region of Saudi Arabia. *Plants* **2022**, *11*, 1028. <https://doi.org/10.3390/plants11081028>
7. Konarska, A.; Weryszko-Chmielewska, E.; Sulborska-Różycka, A.; Kieltyka-Dadasiewicz, A.; Dmitruk, M.; Gorzel, M. Herb and Flowers of *Achillea millefolium* subsp. *millefolium* L.: Structure and Histochemistry of Secretory Tissues and Phytochemistry of Essential Oils. *Molecules* **2023**, *28*, 7791. <https://doi.org/10.3390/molecules28237791>
8. Ramadan, M.; Goeters, S.; Watzer, B.; Krause, E.; Lohmann, K.; Bauer, R.; Hempel, B.; Imming, P. Chamazulene carboxylic acid and matricin: a natural profen and its natural prodrug, identified through similarity to synthetic drug substances. *Journal of Natural Products* **2006**, *69*, 1041–1045. <https://doi.org/10.1021/np0601556>
9. Sizova, N.V. Composition and antioxidant activity of essential oils containing azulene derivatives. *Pharm. Chem. J.* **2012**, *46*, 42–4. Doi: 10.1007/s11094-012-0800-6.
10. Querio, G.; Antoniotti, S.; Foglietta, F.; Levi, R.; Berteà, C. M.; Canaparo, R.; Gallo, M. P. Chamazulene prevents ROS production in human dermal fibroblast and bovine aortic endothelial cells exposed to oxidative stress. *Vascul. Pharmacol.* **2018**, *56*, 103–105. <https://dx.doi.org/10.1016/j.vph.2017.12.025>
11. Capuzzo, A.; Occhipinti, A.; Maffei, M.E. Antioxidant and radical scavenging activities of chamazulene. *Nat. Prod. Res.* **2014**, *28*, 2321–2323. Doi: 10.1080/14786419.2014.931393

12. Matera, R.; Lucchi, E.; Valgimigli, L. Plant Essential Oils as Healthy Functional Ingredients of Nutraceuticals and Diet Supplements: A Review. *Molecules* **2023**, *28*, 901. Doi: 10.3390/molecules28020901
13. Gabbanini, S.; Lucchi, E.; Carli, M.; Berlini, E.; Minghetti, A.; Valgimigli, L. In vitro evaluation of the permeation through reconstructed human epidermis of essentials oils from cosmetic formulations. *Journal of Pharmaceutical and Biomedical Analysis* **2009**, *50*, 370–376. Doi: 10.1016/j.jpba.2009.05.018.
14. Zhou, Y.; He, L.; Wang, W.; Wei, G.; Ma, L.; Liu, H.; Yao, L. Artemisia sieversiana Ehrhart ex Willd. Essential Oil and Its Main Component, Chamazulene: Their Photoprotective Effect against UVB-Induced Cellular Damage and Potential as Novel Natural Sunscreen Additives. *ACS Sustainable Chem. Eng.* **2023**, *11*, 17675–17686. Doi: 10.1021/acssuschemeng.3c04804
15. Fiori, J.; Gotti, R.; Valgimigli, L.; Cavrini, V. Guaiazulene in health care products: determination by GC-MS and HPLC-DAD and photostability test. *Journal of Pharmaceutical and Biomedical Analysis* **2008**, *47*, 710–715. <https://doi.org/10.1016/j.jpba.2008.02.020>.
16. Valgimigli, L. Lipid Peroxidation and Antioxidant Protection. *Biomolecules* **2023**, *13*, 1291. <https://doi.org/10.3390/biom13091291>
17. Fiori, J.; Gotti, R.; Albini, A.; Cavrini, V. Study on the photostability of guaiazulene by high-performance liquid chromatography/mass spectrometry and gas chromatography/mass spectrometry. *Rapid Communications in Mass Spectrometry* **2008**, *22*, 2698–706. <https://doi.org/10.1002/rcm.3661>.
18. Korenberg, C. The photo-ageing behaviour of selected watercolour paints under anoxic conditions. *The British Museum Technical Research Bulletin* **2008**, *2*, 49–57
19. Arney, J. S.; Jacobs, A. J.; Newman, R. The Influence of Oxygen on the Fading of Organic Colorants. *Journal of the American Institute for Conservation* **1979**, *18*, 108–117. Doi: 10.1179/019713679806029020
20. Zielinski, Z.; Presseau, N.; Amorati, R.; Valgimigli, L.; Pratt, D. A. Redox Chemistry of Selenenic Acids and the Insight It Brings on Transition State Geometry in the Reactions of Peroxyl Radicals. *J. Am. Chem. Soc.* **2014**, *136*, 1570–1578. Doi: 10.1021/ja411493t.
21. Matsubara, Y.; Morita, M.; Takekuma, S.; Nakano, T.; Yamamoto, H. M.; Nozoe, T. Oxidation of 4,6,8-Trimethylazulene and Guaiazulene with Hydrogen Peroxide in Pyridine. *Bulletin of the Chemical Society of Japan* **1991**, *64*, 3497–3499. <https://doi.org/10.1246/bcsj.64.3497>
22. Kaufmann, A.; Butcher, P.; Maden, K.; Walker, S.; Widmer, M. Using in silico fragmentation to improve routine residue screening in complex matrices. *J. Am. Mass Spectrom.* **2017**, *28*, 2705–2715. <https://doi.org/10.1007/s13361-017-1800-2>
23. [23] Matsubara, Y.; Yamamoto, H.; Nozoe, T. Oxidation Products of Guaiazulene and Other Azulenic Hydrocarbons. *Studies in Natural Products Chemistry* **1994**, *14*, 313–354. <https://doi.org/10.1016/B978-0-444-81780-8.50014-4>.
24. Chiang H.-M.; Yin, J.-J.; Xia, Q.; Zhao, Y.; Fu, P. P.; Wen, K.-C.; Yu, H. Photoirradiation of azulene and guaiazulene—Formation of reactive oxygen species and induction of lipid peroxidation. *Journal of Photochemistry and Photobiology A: Chemistry* **2010**, *211*, 123–128. <https://doi.org/10.1016/j.jphotochem.2010.02.007>.
25. Warren, J. J.; Tronic, T. A.; Mayer, J. M. Thermochemistry of Proton-Coupled Electron Transfer Reagents and its Implications. *Chem. Rev.* **2010**, *110*, 6961–7001. <https://doi.org/10.1021/cr100085k>
26. Amorati, R.; Baschieri, A.; Morroni, G.; Gambino, R.; Valgimigli, L. Peroxyl Radical Reactions in Water Solution: A Gym for Proton-Coupled Electron-Transfer Theories. *Chem. Eur. J.* **2016**, *22*, 7924 – 7934. DOI: 10.1002/chem.201504492.
27. Singha, A.; Kaishyop, J.; Khan, T. S.; Chowdhury, B. Visible-Light-Driven Toluene Oxidation to Benzaldehyde over WO₃ Nanostructures. *ACS Appl. Nano Mater.* **2023**, *6*, 21818–21828. <https://doi.org/10.1021/acsanm.3c04012>.
28. Battin-Leclerc, F.; Warth, V.; Bounaceur, R.; Husson, B.; Herbinet, O.; Glaude, P.-A. The oxidation of large alkylbenzenes: An experimental and modeling study. *Proceedings of the Combustion Institute* **2015**, *35*, 349–356. <https://doi.org/10.1016/j.proci.2014.05.087>.
29. Zhao, J. Plant Troponoids: Chemistry, Biological Activity, and Biosynthesis. *Current Medicinal Chemistry* **2007**, *14*, 2597–2621. <https://doi.org/10.2174/092986707782023253>.
30. Fu, P.P.; Xia, Q.; Zhao, Y.; Wang, S.; Yu, H.; Chiang, H.-M. Phototoxicity of Herbal Plants and Herbal Products. *Journal of Environmental Science and Health C* **2013**, *31*, 213–255. DOI: 10.1080/10590501.2013.824206
31. Bolton, J. L.; Dunlap, T. Formation and Biological Targets of Quinones: Cytotoxic versus Cytoprotective Effects. *Chem. Res. Toxicol.* **2017**, *30*, 13–37. DOI: 10.1021/acs.chemrestox.6b00256
32. ICH Q1B Guideline. Photostability Testing of New Drug Substances and Products. http://www.columbiapharma.com/reg_updates/international/ich/q1a.pdf (Accessed on May 1st 2024).
33. Caroprese, A.; Gabbanini, S.; Beltramini, C.; Lucchi, E.; Valgimigli, L. HS-SPME-GC-MS analysis of body odor to test the efficacy of foot deodorant formulations. *Skin Research and Technology* **2009**, *15*, 503–510. Doi: 10.1111/j.1600-0846.2009.00399.x

34. Amorati, R.; Valgimigli, L.; Panzella, L.; Napolitano, A.; d'Ischia, M. 5-S-Lipoylhydroxytyrosol, a Multidefense Antioxidant Featuring a Solvent-Tunable Peroxyl Radical-Scavenging 3-Thio-1,2-dihydroxybenzene Motif *J. Org. Chem.* **2013**, *78*, 9857–9864. Doi: 10.1021/jo401522q.
35. Cariola, A.; El Chami, M.; Granatieri, J.; Valgimigli, L. Anti-tyrosinase and antioxidant activity of meroterpene bakuchiol from *Psoralea corylifolia* (L.). *Food Chemistry* **2023**, *405*, 134953. <https://doi.org/10.1016/j.foodchem.2022.134953>.
36. Foti, M. C.; Ingold, K. U. Mechanism of Inhibition of lipid peroxidation by γ -terpinene, an unusual and potentially useful hydrocarbon antioxidant. *Journal of Agriculture and Food Chemistry* **2003**, *51*, 2758–2765. <https://doi.org/10.1021/jf020993f>.
37. Guo, Y.; Baschieri, A.; Amorati, R.; Valgimigli, L. Synergic antioxidant activity of γ -terpinene with phenols and polyphenols enabled by hydroperoxyl radicals. *Food Chemistry* **2021**, *345*, 128468. Doi: 10.1016/j.foodchem.2020.128468.
38. Baschieri, A.; Valgimigli, L.; Gabbanini, S.; DiLabio, G. A.; Romero-Montalvo, E.; Amorati, R. Extremely Fast Hydrogen Atom Transfer between Nitroxides and HOO \cdot Radicals and Implication for Catalytic Coantioxidant Systems *J. Am. Chem. Soc.* **2018**, *140*, 10354–10362. DOI: 10.1021/jacs.8b06336.
39. Valgimigli, L.; Lucarini, M.; Pedulli, G. F.; Ingold, K. U. Does β -Carotene Really Protect Vitamin E from Oxidation? *J. Am. Chem. Soc.* **1997**, *119*, 8095–8096. Doi: 10.1021/ja971351p.
40. Johansson, H.; Shanks, D.; Engman, L.; Amorati, R.; Pedulli, G.F.; Valgimigli, L. Long-lasting antioxidant protection: A regenerable BHA analogue. *Journal of Organic Chemistry* **2010**, *75*, 7535 – 7541. Doi: 10.1021/jo101239c
41. Pniewska, A.; Kalinowska-Lis, U. A Survey of UV Filters Used in Sunscreen Cosmetics. *Appl. Sci.* **2024**, *14*, 3302. <https://doi.org/10.3390/app14083302>
42. Nitulescu, G.; Lupuliasa, D.; Adam-Dima, I.; Nitulescu, G.M. Ultraviolet Filters for Cosmetic Applications. *Cosmetics* **2023**, *10*, 101. <https://doi.org/10.3390/cosmetics10040101>
43. Durand, L.; Habran, N.; Henschel, V., & Amighi, K. (2010). Encapsulation of ethylhexyl methoxycinnamate, a light-sensitive UV filter, in lipid nanoparticles. *Journal of Microencapsulation*, *27*(8), 714–725. <https://doi.org/10.3109/02652048.2010.513455>
44. Dencausse, L.; Galland, A.; Clamou J. L.; J. Basso, J. Validation of HPLC method for quantitative determination of Tinosorb S and three other sunscreens in a high protection cosmetic product. *International Journal of Cosmetic Science* **2008**, *30*, 373–382. Doi: 10.1111/j.1468-2494.2008.00461.x
45. Hüglin, D. Advanced UV Absorbers for the Protection of Human Skin. *Chimia* **2016**, *70*, 496–501. doi:10.2533/chimia.2016.496
46. Arimoto-Kobayashi, S.; Anma, N.; Yoshinaga, Y.; Douki, T.; Cadet, J.; Hayatsu, H. Oxidative damage and induced mutations in m13mp2 phage DNA exposed to N-nitrosopyrrolidine with UVA radiation. *Mutagenesis* **2000**, *15*, 473–477. Doi: 10.1093/MUTAGE/15.6.473.
47. [47] Yu, H. B.; Jin, W.; Ho, H. L.; Chan, K. C.; Chan, C. C.; Demokan, M.S.; Stewart, G.; Culshaw, B.; Liao, Y. B. Multiplexing of optical fiber gas sensors with a frequency-modulated continuous-wave technique. *Appl. Opt.* **2001**, *40*, 1011. Doi: 10.1364/AO.40.001011.
48. Wong, N.G.K.; Sereli, M.; Anstöter, C.S.; Dessent, C.E.H. Photochemical Degradation of the UV Filter Octyl Methoxy Cinnamate Probed via Laser-Interfaced Mass Spectrometry. *Molecules* **2022**, *27*, 8796. <https://doi.org/10.3390/molecules27248796>.
49. Jentzsch, F.; Reich, M.; Kümmerer, K.; Olsson, O. Photolysis of mixtures of UV filters octocrylene and ethylhexyl methoxycinnamate leads to formation of mixed transformation products and different kinetics. *Science of The Total Environment* **2019**, *697*, 134048. <https://doi.org/10.1016/j.scitotenv.2019.134048>.
50. Hanson, K. M.; Narayanan, S.; Nichols, V. M.; Bardeen C. J. Photochemical degradation of the UV filter octyl methoxycinnamate in solution and in aggregates. *Photochem. Photobiol. Sci.* **2015**, *14*, 1607–1616. DOI: 10.1039/c5pp00074b.
51. Regulation (EC) No 1223/2009 of the European Parliament and of the Council of 30 November 2009 on cosmetic products. <https://eur-lex.europa.eu/eli/reg/2009/1223/oj> (Accessed on May 2nd 2024)
52. Bowry, V. W.; Ingold, K. U. The Unexpected Role of Vitamin E (α -Tocopherol) in the Peroxidation of Human Low-Density Lipoprotein. *Acc. Chem. Res.* **1999**, *32*, 1, 27–34. <https://doi.org/10.1021/ar950059o>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.