

Evaluation of an Alternative Screening Method for Gestational Diabetes Diagnosis During the COVID-19 Pandemic (DIABECOVID STUDY): An Observational Cohort Study

Alba Casellas , Cristina Martinez , Judit Amigó , Roser Ferrer , Laia Marti , Carme Merced ,
M Carmen Medina , Istria Molinero , Marta Calveiro , Anna Maroto , [Ester del Barco](#) ^{*} , Elena Carreras ,
[Maria M. Goya](#)

Posted Date: 25 September 2024

doi: 10.20944/preprints202409.1985.v1

Keywords: Gestational diabetes; COVID-19; O'Sullivan; OGTT; glucose level; glycated haemoglobin; screening; macrosomia; large for gestational age (LGA); neonatal hypoglycemia



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Evaluation of an Alternative Screening Method for Gestational Diabetes Diagnosis during the COVID-19 Pandemic (DIABECOVID STUDY): An Observational Cohort Study

Alba Casellas ¹, Cristina Martínez ², Judit Amigó ³, Roser Ferrer ⁴, Laia Martí ⁵, Carme Merced ⁶, M Carmen Medina ⁷, Istria Molinero ⁸, Marta Calveiro ⁹, Anna Maroto ¹⁰, Ester del Barco ^{1,*}, Elena Carreras ^{1,†}, Maria Goya ^{1,†} and on behalf of the DIABECOVID STUDY Group [‡]

¹ Maternal-Foetal Medicine Unit, Department of Obstetrics and Gynecology, Hospital Universitari Vall d'Hebron. Universitat Autònoma de Barcelona, Barcelona, Spain

² Sexual and Reproductive Health Services. Catalan Institute of Health. Universitat de Barcelona (UB), Barcelona, Spain

³ Department of Endocrinology. Hospital Universitari Vall d'Hebron. Universitat Autònoma de Barcelona, Barcelona, Spain

⁴ Department of Clinical Biochemistry, Hospital Universitari Vall d'Hebron. Universitat Autònoma de Barcelona, Barcelona, Spain

⁵ Department of Obstetrics and Gynecology, Hospital Universitari Parc Taulí. Sabadell, Barcelona, Spain

⁶ Department of Obstetrics and Gynecology, Consorci Hospitalari de Vic, Barcelona, Spain

⁷ Department of Obstetrics and Gynecology, Hospital de la Creu i Sant Pau. Barcelona, Barcelona, Spain

⁸ Department of Obstetrics and Gynecology, Hospital De Igualada, Barcelona, Spain.

⁹ Atenció a la salut sexual i reproductiva (ASSIR) Muntanya Coordinator, Barcelona, Spain

¹⁰ Department of Obstetrics and Gynecology, Hospital Josep Trueta. Girona, Spain

¹¹ Servei d'Endocrinologia i Nutrició, Hospital de la Santa Creu i Sant Pau, Barcelona; Department of Medicine, Universitat Autònoma de Barcelona; CIBER-BBN, Madrid. Hospital Sant Pau. Barcelona, Barcelona, Spain

¹² Department of Endocrinology, Consorci Hospitalari de Vic, Barcelona, Spain

¹³ Midwife ASSIR Osona, Barcelona, Spain

* Correspondence: ester.delbarco@vallhebron.cat; Tel.: +34934893085

† Equal contribution.

‡ On behalf of the DIABECOVID TRIAL group: Andrea Maria Anleu Carranza ⁶, Clementina de Antonio ¹, Silvia Arévalo ¹, Marta Boqué ⁹, Laura Borrega Escalante ⁸, Laura Bou Sánchez ⁵, Anna Brox ¹, Berta Buldó Ibáñez ¹⁰, Núria Burgaya Guiu ⁵, Anna Casteras ³, Cristina Celada Castro ⁶, Rosa Corcoy ⁷, Marta Dalmau ¹, Núria Estopiña Cros ⁶, Patricia Díaz ¹, Luciana Gimenez ⁸, Marina Giralt ⁴, Santiago Lopez Vanega ¹¹, Laura Mañé, PhD ¹², Eva Martinez Garcia ⁸, Pilar Matencio Belchi ⁵, M Carmen Medina, PhD ¹¹, Jordina Munros, PhD ⁹, Ester Muñoz Sellés, PhD ¹³, Núria Pagès Campderrós ⁸, Eider Peñas Calvo ⁵, Ana Perestelo ¹, Judit Platero Diago ¹¹, Elisabeth Sanfeliu ¹, Marta Sanjosé ¹, Olga Simó-Servat ³, Rafael Simó ³, Anna Suy ¹, Judit Tarrés Rovira ¹³, Laura Toledo Mesa ⁵, Elvira Vallés ¹, Natàlia Visa Figueredo ¹⁰ and Ana Zaragoza ⁹.

Abstract: Aim: To evaluate the impact of applying alternative diagnostic criteria for gestational diabetes mellitus (GDM) during the COVID-19 pandemic on GDM prevalence, obstetrical and perinatal outcomes, and costs, as compared to the standard diagnostic method. Methods: A cohort of pregnant individuals undergoing GMD screening with the alternative GDM method, which uses plasma glucose (fasting or non-fasting) and HbA1c, was compared with a cohort of pregnant individuals undergoing the standard GDM screening method. Both cohorts were obtained from 6 hospitals across Catalonia, Spain, from April 2020 to April 2022. The primary outcome was large for gestational age rate at birth. Secondary outcomes were composite adverse outcomes including pregnancy complications, delivery complications, and neonatal complications. Cost differences between screening methods were also evaluated. A similar analysis was performed in the subgroup diagnosed with GDM. Results: Data were collected from 1,543 pregnant individuals in the standard screening group and 2,197 in the alternative screening group. The standard screening group had a higher GDM diagnostic rate than the alternative screening group (10.8% vs 6.9%, respectively; $p < 0.0001$). The primary outcome (large for

gestational age rate) was similar between groups: 200/1543 (13.0%) vs 303/2197 (13.8%). The adjusted OR for this outcome was 1.74 (95% CI: 0.74-4.10). An adjusted analysis showed no differences between groups in composite adverse outcomes for pregnancy complications (OR: 1.11; 95% CI: 0.91-1.36), delivery complications (OR: 0.95; 95% CI: 0.75-1.19), and neonatal complications (OR: 1.28; 95% CI: 0.94-1.75). Among individuals diagnosed with GDM, the large for gestational age rate was similar between groups: 13/166 (7.8 %) vs 15/151 (9.9 %). The OR adjusted for this outcome was 1.24 (95% CI: 0.51-3.09). An adjusted analysis showed no differences in composite adverse outcomes for pregnancy complications (OR: 1.57; 95% CI: 0.84-2.98), delivery complications (OR: 1.21; 95% CI: 0.63-2.35), and neonatal complications (OR: 1.35; 95% CI: 0.61-3.04). The mean cost (which included expenses for consumables, equipment and personnel) of the alternative screening method was 46.0 euros (22.3 SD), as compared to 85.6 euros (67.5 SD) for the standard screening method. Conclusions: In this Spanish population during the COVID-19 pandemic, GDM prevalence was lower in the alternative screening group than in the standard screening group. After adjusting for GDM risk factors, outcomes related to obstetrics, delivery, and neonatal complications were comparable between both groups. Finally, the alternative screening method was cheaper than the standard screening method.

Keywords: gestational diabetes; COVID-19; O'Sullivan; OGTT; glucose level; glycated haemoglobin; screening; macrosomia; large for gestational age (LGA); neonatal hypoglycemia

1. Introduction

The diagnostic challenges of gestational diabetes mellitus (GDM) have sparked extensive debates among experts. The primary areas of debate include selecting between universal and selective screening, two-step versus one-step diagnostic approaches, in particular, the oral glucose tolerance test (OGTT), selecting the most suitable diagnostic test (e.g., 100-g or 75-g OGTT), and establishing appropriate cut-off values for diagnosis [1,2]. To make matters worse, the COVID-19 pandemic has added an unexpected layer of complexity to this ongoing dilemma.

During the COVID-19 pandemic, it became apparent that pregnant individuals faced significant risks when attending healthcare sites for GDM screening following the usual two-step approach, including the 50-g O'Sullivan test (OGTT) with 1-h glucose measurement and the 100-g OGTT with 3-h glucose measurement. Consequently, several international societies, including the Spanish Group of Diabetes and Pregnancy (GEDE) and the Catalan Department of Health, developed an alternative approach to the standard GDM screening method [3,4,5,6,7]. This alternative method streamlines the screening process through a one-step process that includes plasma glucose (fasting or non-fasting) and hemoglobin A1c (HbA1c). This approach minimizes prolonged hospital stays [8,9,10,11,12,13]. Recently, several researchers have evaluated the consequences of implementing these alternative diagnostic criteria for gestational diabetes by comparison to a historical dataset. The suggested modifications to the diagnostic criteria for GDM in a number of countries have resulted in different GDM prevalences [16]. Some of these findings indicate an increased number of cases with undiagnosed GDM, of up to 30-50% [14,15]. On the other hand, when using the alternative diagnostic criteria, a Spanish cohort showed a GDM prevalence during the COVID-19 pandemic similar to that found in 2019, when the standard diagnostic criteria was being used. [17]

In response to the unique circumstances during the COVID-19 pandemic, different regions and healthcare domains decided to modify their GDM diagnostic criteria. In April 2020, some Catalan regions adjusted their standard criteria to adapt to the pandemic circumstances. Since then, the criteria for GDM screening adapted to COVID-19 circumstances based on Spanish recommendations has been used in those regions during the pandemic and post-pandemic, although currently most regions have returned to the standard screening method. By contrast, other regions did not see a reason to change their standard GDM screening method.

The primary aim of this study was to assess the impact of following alternative diagnostic criteria for gestational diabetes mellitus (GDM) during the COVID-19 pandemic on GDM prevalence, obstetrical and perinatal outcomes, and cost as compared to the standard diagnostic criteria.

2. Material and Methods

An observational cohort study was performed to investigate the impact of replacing the standard GDM screening method with an alternative screening method to adapt to the circumstances of the COVID-19 pandemic. This study was approved by the Vall d'Hebron Barcelona Hospital Campus Ethics Committee (VHEC) PR(AMI)381/2020 in October 2020. This study was also approved by the Ethics Committee of each site. The study period ranged from April 2020 to April 2022. When the COVID-19 pandemic started, an alternative Spanish method for GDM screening was developed and published in May 2020. This alternative method was implemented and performed from May 2020 to April 2022. The retrospective period of the study included a cohort of patients undergoing the standard GDM screening from May 2020 to November 2020. Patient consent was waived for this cohort due to its retrospective nature. Informed consent was obtained from all individuals included in the prospective period of the study (November 2020 to April 2022). This work complied with the guidelines for human studies and was conducted ethically following the World Medical Association Declaration of Helsinki.

A cohort of pregnant individuals undergoing an alternative GMD screening (based on plasma glucose and HbA1c, fasting or non-fasting) due to the COVID-19 pandemic was compared with a cohort of pregnant individuals undergoing the standard GMD screening. Participants for both cohorts were obtained from 6 hospitals across Catalonia, Spain, from April 2020 to April 2022. The participating hospitals were Hospital Vall d'Hebron, Hospital Universitari Parc Tauli, Consorci Hospitalari de Vic, Hospital de Igualada, Hospital de Sant Pau and Hospital Josep Trueta. Each participating hospital, following their internal protocols for the prevention of in-hospital COVID transmission, decided to either maintain the standard GDM screening protocol or switch to an alternative method proposed by different medical societies [13]. Thus, two cohorts were obtained: a group undergoing the standard GMD screening and a group undergoing the alternative GMD screening.

The standard GMD screening consists of a two-step format, and tests are performed in the first, second and third trimesters [18]. In the first trimester, screening is performed following the Catalan regional guidelines (20), only if risk factors are detected. Risk factors include age >35 years, obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$ or 27.5 kg/m^2 in Asian Americans), first-degree family history of diabetes, and certain ethnicities (African Americans, Latino Americans, Pacific Islanders, Asian Americans). The test performed in the first trimester consists of fasting glucose. If this is over 92 mg/dl, the 1-hour O'Sullivan test (OGTT), which involves ingesting 50 g of glucose, is performed. If glucose levels after one hour are $\geq 140 \text{ mg/dl}$, the 100/g OGTT is then conducted. Fasting glucose cut-off levels for GDM diagnosis are $\geq 105 \text{ mg/dl}$, 1-hour $\geq 190 \text{ mg/dl}$, 2-hour $\geq 165 \text{ mg/dl}$, and 3-hour $\geq 145 \text{ mg/dl}$ (at least two of this cut-offs). Universal GDM screening is conducted in the second trimester between 24 and 28 gestational weeks. Third-trimester screening is performed only in cases with large for gestational age (LGA) fetus or polyhydramnios, or if GDM screening has not been conducted during the first or second trimester.

The alternative screening method is also conducted in the first trimester only if risk factors are detected, in the second trimester for all patients, and in the third trimester only in some cases. Third-trimester screening is performed if cases with LGA fetus or polyhydramnios, or if GDM screening is not conducted during the first or second trimester. The alternative screening method involves measuring both HbA1c and plasma glucose levels (preferably non-fasting due to feasibility or if fasting is not possible). The established cut-off values for GDM diagnosis in the first trimester were as follows: HbA1c $\geq 5.9\%$, fasting PG $\geq 100 \text{ mg/dl}$, and non-fasting PG $\geq 165\text{-}199 \text{ mg/dl}$. The established cut-off values for GDM diagnosis in the second trimester were HbA1c $\geq 5.7\%$, fasting PG $\geq 95 \text{ mg/dl}$, and non-fasting PG $\geq 165\text{-}199 \text{ mg/dl}$ [13]

Inclusion criteria were being between 18 and 50 years old and availability of perinatal outcomes. Participants with multiple gestation or pre-existing type-1 or type-2 diabetes were excluded.

All pregnant individuals included adhered to the established prenatal care protocol of each site, which was based on national guidelines [19,20]. The group diagnosed with GDM underwent a

specific diet and glucose self-monitoring. Insulin was added as an additional treatment according to recommendations of Grupo Español de Diabetes y Embarazo (GEDE)[21].

Demographic, pregnancy, delivery and neonatal data were collected using an electronic form by RedCap, a web-based record specifically designed for this study.

Demographic data, including maternal age, ethnicity, smoking status, family history of DM, previous GDM, and parity, were collected.

Pregnancy outcomes were collected. Gestational hypertension was classified as worsening of chronic hypertension, new-onset gestational hypertension, and preeclampsia, up to 3 months postpartum; polyhydramnios was defined as amniotic fluid index ≥ 25 cm; LGA was defined as a birth weight above the 90th percentile according to customised growth curves; small for gestational age (SGA) was defined as a birth weight below the 10th percentile according to customised growth curves; gestational loss included miscarriage, antepartum death and intrapartum death; gestational age at delivery was defined in complete weeks based on the date of last menstrual period (LMP) or using the earliest ultrasound if there was a discrepancy with LMP.

Delivery data were also collected. Obstetric trauma was defined as shoulder dystocia, clavicle fracture, or brachial plexus injury; type of delivery was defined as vaginal delivery or caesarean section; maternal stay in the hospital was defined as the number of days after delivery at hospital for the mother; and macrosomia rate was defined as a birth weight >4000 g.

Neonatal outcomes were also recorded. Congenital anomalies were classified by EUROCAT codes references; perinatal death was defined as fetal death from week 28 up to 7 days postpartum; neonatal hypoglycemia was defined as plasma glucose levels <45 mg/dL within the first 24 hours of life and <50.5 mg/dL afterwards; neonatal hypocalcemia was defined as calcium <7 mg/dl; neonatal hyperbilirubinemia was defined as hyperbilirubinemia treated with phototherapy; neonatal polycythemia was defined as a hematocrit of peripheral venous sample $>65\%$; respiratory distress syndrome was defined as progressive respiratory failure shortly after birth, along with certain features in a chest X-ray; excluding other causes; hypertrophic cardiomyopathy was defined as left ventricular wall thickness ≥ 15 mm by echocardiography.

Cost data were also taken into consideration. Expenses for lab tests (50-g and 100-g glucose bottles, blood glucose test, and glycated hemoglobin test) and personnel for conventional analysis were collected and compared between groups.

The primary outcome for both groups was defined as large for gestational age (LGA) rate at birth. Secondary outcomes were composite adverse outcomes including pregnancy complications (macrosomia at birth, LGA at birth, hypertension during pregnancy, polyhydramnios, preterm birth <37 weeks, congenital anomalies, gestational loss in the second trimester, and induction delivery), delivery complications (caesarean section, obstetric trauma, intrapartum death, postpartum complications, and maternal admission), and neonatal complications (Apgar test <3 at 1 min, Apgar test <3 at 5 min, arterial pH <7 at birth, venous pH <7 at birth, neonatal death, neonatal intensive care unit (NICU) admission, neonatal hypoglycemia, neonatal hypocalcemia, neonatal hyperbilirubinemia, neonatal polycythemia, and perinatal death).

The sample size was estimated taking into account the impact of screening tests on GDM and large for gestational age prevalences. The GDM prevalence in our population is estimated at 12% with an error of up to 4%. With a 95% confidence level, the sample size would be 1015 participants per group (2030 cases in total). The large for gestational age fetal prevalence in our population is estimated at 16.5% with an error of up to 4%. With a 95% confidence level, the sample size would be 1324 participants per group (2648 cases in total). Therefore, the required target sample size is 3000 pregnant individuals, which allows for up to 10% of cases with incomplete or missing data.

Descriptive data are expressed as the median and interquartile range (IQR), or as absolute and relative frequencies, as appropriate. Variables were compared using the t-test or the Fisher's exact test. A multivariable logistic regression model was fitted for each composite variable and for each outcome variable, using the composite or outcome variables as the dependent variable and the screening group (standard screening vs alternative screening) as the independent variable, adjusted

for obesity, maternal age, metabolic control, ethnicity, smoking status, family history of diabetes, previous GDM, chronic hypertension, parity, and site of inclusion.

The analysis was conducted for the entire sample and only in the group diagnosed with GDM. The diagnostic rate for each trimester with each method was compared using a proportionality test [22].

The odds ratio, 95% confidence interval and p values are reported. All analyses were conducted using the software R, version 4.3 [and SPSS for Windows 19.0 statistical package].

3. Results

A total of 4,811 pregnant individuals were considered for this study. Of those, 1,071 were excluded: 274 due to pregestational diabetes, 182 due to multiple gestations, and 615 due to unavailability of perinatal outcomes. Therefore, a total of 3,740 pregnant individuals were included in this study: data were collected from 1,543 pregnant individuals in the standard screening group and from 2,197 in the alternative screening group.

3.1. Whole-Population Analysis

In the whole-population analysis, some differences in demographics were found (Table 1). Although maternal age, parity and number of preterm births were similar in both groups (p=0.220, p=0.199 and p=0.131, respectively), there were statistically significant differences in body mass index (BMI), with pregnant individuals having slightly higher BMI in the alternative screening group (p=0.002). We also found that individuals in the standard screening group were more frequently nulliparous (p=0.001), had a higher prevalence of tobacco use during pregnancy (p<0.0001), and chronic HTA prior to pregnancy (p=0.024). In addition, there were some statistically significant differences in family history of DM and prior GDM between both groups (p=0.007 and p<0.0001, respectively), probably due to the high number of missing data in the standard screening group. There were also statistically significant differences for ethnicity between both groups. A higher rate of GDM risk factors in the first trimester was observed in the alternative screening group (p=0.024). However, the standard screening group had a higher GDM diagnostic rate (10.8% vs 6.9%, p<0.0001).

The primary outcome (LGA rate) was similar between groups: 200/1543 (13.0%) vs 303/2197 (13.8%) (OR: 1.74; 95% CI: 0.74-4.10; p=0.672).

Table 1. Demographic and general characteristics of the whole population of the study, classified according to screening method.

	Standard screening method n=1543	Alternative screening method n=2197	P value
Maternal age (years)	32.2 (5.7)	32.2 (5.9)	0.220
Obesity [BMI (kg/m²) ≥30]			
No	1260 (81.7%)	1729 (78.7%)	0.002
Yes	237 (15.3%)	431 (19.6%)	
No data	46 (3.0%)	37 (1.7%)	
Previous term birth			
No	748 (48.5%)	997 (45.4%)	0.199
One	514 (33.3%)	732 (33.3%)	
Two	189 (12.2%)	314 (14.3%)	
Three	68 (4.4%)	104 (4.7%)	
Four	16 (1.0%)	32 (1.5%)	
Five or more	8 (0.5%)	18 (0.8%)	
Previous preterm birth			
No	1476 (95.7%)	2065 (93.9%)	0.131
One	63 (4.1%)	121 (5.5%)	

Two	4 (0.3%)	10 (0.5%)	
Three or more	0 (0%)	1 (0.1%)	
Parity			
Nulliparous	674 (43.7%)	845 (38.5%)	0.001
Multiparous	868 (56.3%)	1352 (61.5%)	
Ethnicity			
Caucasian	873 (56.8%)	1298 (59.1%)	
African-American	106 (6.9%)	59 (2.7%)	
Asiatic	35 (2.3%)	33 (1.5%)	
Maghreb	237 (15.4%)	231 (10.5%)	<0.0001
Hindu	35 (2.3%)	79 (3.6%)	
Latin-American	241 (15.7%)	422 (19.2%)	
Others	10 (0.7%)	67 (3.0%)	
No data	6 (0.4%)	8 (0.4%)	
Tobacco use during pregnancy			
No	1310 (84.9%)	1952 (88.9%)	<0.0001
Yes	233 (15.1%)	245 (11.1%)	
Drug abuse during pregnancy			
No	1512 (98.0%)	2157 (98.2%)	0.557
Yes	30 (1.9%)	37 (1.7%)	
No data	1 (0.1%)	3 (0.1%)	
Family history of DM			
No	803 (52.0%)	1725 (78.5%)	<0.0001
Yes	66 (4.3%)	177 (8.1%)	
No data	674 (43.7%)	295 (13.4%)	
Previous GDM			
No	1412 (91.5%)	2048 (93.2%)	0.007
Yes	46 (3.0%)	74 (3.4%)	
No data	85 (5.5%)	75 (3.4%)	
Risk factors for GMD in first trimester			
No	1267 (82.1%)	1689 (76.9%)	<0.0001
Yes	276 (17.9%)	508 (23.1%)	
Chronic HTA before pregnancy	24 (1.6%)	17 (0.8%)	0.024
GDM diagnosis rate	166 (10.8%)	152 (6.9%)	<0.0001

BMI: body mass index, GDM: gestational diabetes mellitus, HTA: hypertension arterial. *Data are shown as the median and standard deviation for quantitative variables; as number of cases and percentage for qualitative variables. The p value was considered statistically significant at <0.05. Tests used were the Chi-square test for qualitative variables and the Student's t test for quantitative variables.

An adjusted analysis considering maternal obesity (BMI $\geq$ 30 kg/m²), parity, ethnicity, smoking status, family history of DM, previous GDM, GMD risk factors in the first trimester, previous HTA, GDM screening method and site of inclusion was performed (Table 2A and 2B). There were no differences between groups for the composite adverse outcomes of pregnancy complications (OR: 1.11; 95% CI: 0.91-1.36; p=0.312), delivery complications (OR: 0.95 (95% CI: 0.75-1.19; p=0.642), and neonatal complications (OR: 1.28; 95% CI: 0.94-1.75; p=0.118).

Table 2. A. Pregnancy complications, perinatal events, and neonatal morbidities in the whole population of the study, classified according to screening method. B. Composite adverse outcomes including pregnancy complications, delivery complications, and neonatal complications classified according to screening method (reference category: standard screening method).

A				
Characteristic	Standard screening method n=1543	Alternative screening method n=2197	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Macrosomia at birth	96 (6.2 %)	122 (5.6%)	0.89 (0.67-1.17)	0.98 (0.65-1.48)
LGA at birth (>P90)	200 (13.0 %)	303 (13.8 %)	1.07 (0.89-1.30)	0.92 (0.68-1.23)
Hypertension during pregnancy	17 (1.1%)	25 (1.1%)	1.3 (0.98-1.74)	1.03 (0.47-2.26)
Polyhydramnios	16 (1.0 %)	16 (0.7%)	0.71 (0.35-1.43)	0.86 (0.27-2.78)
Preterm birth (PB) < 37 weeks	90 (5.8%)	201 (9.1%)	1.73 (1.34-2.25)	1.16 (0.78-1.73)
Spontaneous PB < 37 weeks	42 (2.7%)	98 (4.5%)	2.11 (1.46-3.09)	1.64 (0.96-2.82)
Gestational loss in 2nd trimester	9 (0.6 %)	17 (0.8%)	1.35 (0.61-3.17)	0.49 (0.16-1.47)
Induction delivery rate	524 (34.0%)	824 (37.5%)	1.17 (1.02-1.34)	0.91 (0.74-1.12)
Cesarean section rate	342 (22.3%)	520 (23.7%)	1.17 (1.00-1.36)	0.81 (0.63-1.03)
Obstetric trauma	15 (1 %)	19 (0.9%)	0.90 (0.46-1.80)	1.79 (0.81-3.95)
Intrapartum fetal death	2 (0.1 %)	2 (0.1%)	0.71 (0.09-5.93)	0.82 (0.05-14.08)
Postpartum complications	5 (0.3%)	7 (0.3%)	0.98 (0.31-3.33)	0.75 (0.55-1.07)
Maternal admission to hospital	125 (8.1%)	188 (9.1%)	1.13 (0.90-1.44)	1.07 (0.82-1.40)
Apgar test <3 at 1 min	20 (1.3 %)	26 (1.2 %)	0.97 (0.54-1.76)	0.86 (0.35-2.19)
Apgar test <3 at 5 min	13 (0.8 %)	12 (0.5 %)	0.69 (0.31-1.52)	0.34 (0.12-1.10)
Neonatal death	4 (0.3%)	6 (0.3%)	1.07 (0.3-4.18)	0.63 (0.09-4.62)
NICU admission	63 (4.1 %)	114 (5.2%)	1.33 (0.98-1.84)	0.78 (0.47-1.31)
Neonatal hypoglycemia	41 (2.7 %)	57 (2.6 %)	1.01 (0.67-1.53)	1.28 (0.73-2.26)
Perinatal death	50 (3.2%)	79 (3.6%)	1.15 (0.81-1.66)	0.95 (0.55-1.65)
B				
Characteristic	Unadjusted model		Adjusted model*	
Composite of pregnancy complications ¹	OR: 1.38 (95% CI: 1.17-1.62) p<0.001		OR: 1.11 (95% CI: 0.91-1.36) p=0.312	
Composite of delivery complications ²	OR: 1.15 (95% CI: 0.95-1.39) p=0.159		OR: 0.95 (95% CI: 0.75-1.19) p=0.642	
Composite of neonatal complications ³	1.05 (0.83-1.33) p=0.718		OR: 1.28 (95% CI: 0.94-1.75) p=0.118	

¹ Includes macrosomia at birth, large for gestational (LGA) at birth (P≥90), hypertension during pregnancy, polyhydramnios, preterm birth <37 weeks, gestational loss during second trimester and induction delivery rate.
² Includes caesarean section rate, obstetric trauma, intrapartum death, postpartum complications and maternal admission.
³ Includes Apgar test <3 at 1 min, Apgar test <3 at 5 min, neonatal death, NICU admission, neonatal hypoglycemia and perinatal death. *Logistic regression models adjusted for maternal obesity (BMI≥30), parity, ethnicity, smoking status, previous GDM, risk factors for GMD in first trimester, previous HTA and site of inclusion.

3.2. GDM Group Analysis

Demographic characteristics of individuals diagnosed with GDM were analyzed and compared (Table 3). There were no differences between groups for maternal age, maternal weight increase during pregnancy, previous preterm births, tobacco use and drug abuse during pregnancy, previous GDM, GDM risk factors in the first trimester, chronic HTA prior to pregnancy, and metformin treatment. However, there were some statistically significant differences in family history of DM between groups and there was a higher rate of nulliparity in the standard screening group ($p=0.003$). There were also statistically significant differences between groups for metabolic control, with a poorer metabolic control ($p<0.001$), higher need for basal insulin ($p=0.015$) and multiple insulin doses ($p=0.002$) in the alternative screening group.

Table 3. Demographic and general characteristics of individuals with diagnosis of gestational diabetes according to screening method.

	Standard screening method n=166	Alternative screening method n=151	P value
Maternal age (years)	33.9 (5.2)	34.0 (5.5)	0.929
Obesity [BMI (kg/m²) ≥30]			
No	111 (66.9%)	74 (49.0%)	0.002
Yes	54 (32.5%)	74 (49.0%)	
No data	1 (0.6%)	3 (2.0%)	
Maternal weight increase during pregnancy (kg)	6.4 (0.5)	6.6 (0.5)	0.131
Previous term birth			
No	74 (44.6%)	39 (25.8%)	0.016
One	45 (27.1%)	55 (36.4%)	
Two	30 (18.1%)	35 (23.2%)	
Three	12 (7.2%)	18 (11.9%)	
Four	3 (1.8%)	1 (0.7%)	
Five or more	2 (1.2%)	3 (2.0%)	
Previous preterm birth			
No	157 (94.6%)	141 (93.4%)	0.331
One	9 (5.4%)	8 (5.3%)	
Two	0 (0%)	2 (1.3%)	
Three or more	0 (0%)	0 (0%)	
Parity			
Multiparous	64 (38.6%)	35 (23.2%)	0.003
Nulliparous	102 (61.4%)	116 (76.8%)	
Ethnicity			
Caucasian	84 (50.6%)	58 (38.4%)	<0.0001
African-American	11 (6.6%)	7 (4.6%)	
Asiatic	4 (2.4%)	1 (0.7%)	
Maghreb	38 (22.9%)	21 (13.9%)	
Hindu	6 (3.6%)	17 (11.3%)	
Latin-American	21 (12.7%)	42 (27.2%)	
Others	2 (1.2%)	6 (4.0%)	
Tobacco use during pregnancy			
No	147 (88.6%)	135 (89.4%)	0.809
Yes	19 (11.4%)	16 (10.6%)	

Drug abuse during pregnancy			
No	165 (99.4%)	150 (99.3%)	0.946
Yes	1 (0.6%)	1 (0.7%)	
Family history of DM			
No	74 (44.6%)	97 (64.2%)	<0.0001
Yes	25 (15.1%)	37 (24.5%)	
No data	67 (40.4%)	17 (11.3%)	
Previous GDM			
No	131 (78.9%)	116 (76.8%)	0.716
Yes	31 (18.7%)	29 (19.2%)	
No data	4 (2.4%)	6 (4.0%)	
Risk factors of GDM in first trimester			
No	95 (57.2%)	84 (55.6%)	0.774
Yes	71 (42.8%)	67 (44.4%)	
Chronic HTA before pregnancy			
No	152 (91.6%)	132 (87.4%)	0.227
Yes	14 (8.4%)	19 (12.6%)	
Glycemic control			
No control	5 (3.1%)	16 (11%)	<0.0001
Inadequate control	10 (6.3%)	32 (21.9%)	
Suboptimal control	35 (21.9%)	30 (20.5%)	
Optimal control	110 (68.8%)	68 (46.6%)	
Basal insulin			
No	107 (64.5%)	77 (51%)	0.015
Yes	59 (35.5%)	74 (49%)	
Insulin treatment with multiple doses			
No	118 (71.1%)	82 (54.3%)	0.002
Yes	48 (28.9%)	69 (45.7%)	
Metformin treatment			
No	164 (98.8%)	146 (96.7%)	0.202
Yes	2 (1.2%)	5 (3.3%)	

*Data are shown as the median and standard deviation for quantitative variables; as the number of cases and percentage for qualitative variables. P value was considered statistically significant at <0.05. Tests used were the Chi-square test for qualitative variables and the Student’s t test for quantitative variables.

In individuals diagnosed with GDM, the primary outcome (LGA rate at birth) was similar between groups: 13/166 (7.8%) vs 15/151 (9.9%) (p=0.534) (OR: 1.24; 95% CI: 0.51-3.09; p=0.634). An analysis adjusting for maternal obesity (BMI≥30), parity, ethnicity, smoking status, family history of GDM, previous GDM, GMD risk factors in the first trimester, previous HTA, GDM screening method and site of inclusion was performed for composite outcomes (Tables 4A and 4B). When comparing the standard and alternative screening groups, there were no differences in composite adverse outcomes for pregnancy complications (OR: 1.57; 95% CI: 0.84-2.98; p=0.159), delivery complications (OR: 1.21; 95% CI: 0.63-2.35; p=0.565) and neonatal complications (OR: 1.35; 95% CI: 0.61-3.04; p=0.455).

Table 4. A. Pregnancy complications, perinatal events, and neonatal morbidities in individuals diagnosed with gestational diabetes, classified according to screening method. B. Composite adverse outcomes considering pregnancy complications, delivery complications, and neonatal complications in individuals diagnosed with gestational diabetes according to screening method (reference category: standard screening method).

A				
Characteristic	Standard screening method n=166	Alternative screening method n=152	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Macrosomia at birth	13 (7.8%)	15 (9.9%)	1.29 (0.59-2.84)	1.19 (0.35-4.09)
LGA at birth (>P90)	36 (21.7%)	48 (31.8%)	1.67 (1.01-2.77)	1.74 (0.74-4.10)
Hypertension during pregnancy	10 (6.0%)	17 (11.2%)	1.96 (0.88-4.59)	1.43 (0.40-5.09)
Polyhydramnios	7 (4.2%)	4 (2.6%)	0.61 (0.16-2.08)	0.50 (0.07-3.30)
Preterm birth PB < 37 weeks	11 (6.7%)	15 (10.4%)	1.63 (0.73-3.76)	1.17 (0.41-3.38)
Gestational loss on second trimester	1 (0.6%)	0 (0%)	-	-
Induction delivery rate	78 (47.0%)	94 (61.8%)	1.83 (1.17-2.87)	1.57 (0.88-2.82)
Caesarean section rate	42 (25.5%)	44 (30.1%)	1.26 (0.77-2.08)	0.72 (0.37-1.4)
Obstetric trauma	4 (2.4%)	4 (2.6%)	1.09 (0.25-4.7)	1.16 (0.24-5.60)
Intrapartum fetal death	0 (0%)	1 (0.7%)	-	-
Postpartum complications Maternal sepsis	12 (7.2%)	20 (13.2%)	0.51 (0.24-1.08)	0.38 (0.15-0.91)
Maternal Admission	15 (9.1%)	19 (13.3%)	1.53 (0.75-3.18)	1.71 (0.71-4.19)
Apgar test <3 at 1 min	3 (1.8%)	1 (0.7%)	0.38 (0.02-2.99)	-
Apgar test <3 at 5 min	2 (1.2%)	1 (0.7%)	0.57 (0.03-6.01)	-
Neonatal death	1 (0.6%)	2 (1.3%)	2.2 (0.21-47.62)	-
NICU admission	6 (3.6%)	8 (5.3%)	1.52 (0.52-4.73)	1.25 (0.27-5.76)
Neonatal hypoglycemia	12 (7.2%)	15 (9.9%)	1.45 (0.66-3.26)	1.27 (0.48-3.37)
Perinatal death	8 (4.8%)	5 (3.3%)	0.69 (0.2-2.1)	0.24 (0.4-1.34)
B				
Characteristic	Unadjusted model		Adjusted model*	
Composite of pregnancy complications ¹	OR: 1.41 (95% CI: 0.82-2.43) p=0.216		OR: 1.57 (95% CI: 0.84-2.98) p=0.159	
Composite of delivery complications ²	OR: 1.14 (95% CI: 0.65-2.01) p=0.640		OR: 1.21 (95% CI: 0.63-2.35) p=0.565	
Composite of neonatal complications ³	OR: 1.14 (95% CI: 0.65-2.01) p=0.640		OR: 1.35 (95% CI: 0.61-3.04) p=0.455	

¹Includes macrosomia at birth, large for gestational (LGA) at birth (P≥90), hypertension during pregnancy, polyhydramnios, preterm birth <37 weeks, gestational loss 2T and induction delivery rate. ²Includes caesarean section rate, obstetric trauma, intrapartum death, postpartum complications and maternal admission. ³Includes Apgar test <3 at 1 min, Apgar test <3 at 5 min, neonatal death, NICU admission, neonatal hypoglycemia and perinatal death. *Logistic

regression models adjusted for maternal Obesity (IMC>=30), parity, ethnicity, smoking status, previous GDM, risk factors for GMD in 1T, previous HTA and centre of inclusion.

In order to investigate the impact of metabolic control in pregnancy and neonatal outcomes in individuals diagnosed with GDM and show their impact on previous results, we studied adverse outcomes in terms of pregnancy complications, perinatal complications, and neonatal complications, adjusting for metabolic control and insulin treatment variables (number of individuals needing insulin and dose) (Table 5). After adjusting for confounders, no differences were found between the two groups (BMI, parity, ethnicity, smoking status, family history of DM, previous GDM, risk factors for GDM in the first trimester, previous HTA, GDM screening method for and site of inclusion) in composite adverse outcomes for pregnancy complications (OR: 0.51; 95% CI: 0.22-1.14; p=0.109), delivery complications (OR: 0.98; 95% CI: 0.49-1.99; p=0.955), and neonatal complications (OR: 0.83; 95% CI: 0.35-1.96; p=0.664). 0.83 (0.35-1.96)

Table 5. Composite adverse outcomes considering pregnancy complications, delivery complications, and neonatal complications correlated with metabolic control (and insulin variables) in individuals diagnosed with gestational diabetes.

Characteristic	Adjusted model (95% CI)*
Composite for pregnancy complications ¹	0.51 (0.22-1.14)
Composite for delivery complications ²	0.98 (0.49-1.99)
Composite for neonatal complications ³	(0.35-1.96)

¹Includes macrosomia at birth, large for gestational (LGA) at birth (P≥90), hypertension during pregnancy, polyhydramnios, preterm birth <37 weeks, gestational loss 2T and induction delivery rate. ²Includes caesarean section rate, obstetric trauma, intrapartum death, postpartum complications and maternal admission. ³Includes Apgar test <3 at 1 min, Apgar test <3 at 5 min, neonatal death, NICU admission, neonatal hypoglycemia and perinatal death. *Logistic regression models adjusted for maternal obesity (BMI≥30), parity, ethnicity, smoking status, previous GDM, risk factors for GMD in the first trimester, previous HTA, screening method for gestational diabetes and site of inclusion.

In order to study the financial impact of the alternative screening method, we analyzed the procedure and costs, including expenses for consumables, equipment and personnel, of each screening method (Table 6). There were statistically significant differences between the costs of the two screening methods, with the alternative screening method being cheaper. The mean cost of the alternative screening method was 46.0 euros (SD: 22.3 SD), as compared to 85.6 euros (SD: 67.5) for the standard screening method.

Table 6. Cost of each method (Euros per individual).

	Standard screening method	Alternative screening method	p-value
Total cost	n=1543	n=2197	
Mean (SD)	85.6 (67.5)	46.0 (22.3)	<0.001

*Three aspects were considered for the cost: consumables (catheter, gases, vacutainer, tubes), laboratory (equipment) and personnel (nurses or midwives performing the tests).

4. Discussion

4.1. Findings

The DIABECOVID study assessed the impact of using an alternative method for GDM screening during the COVID-19 pandemic as compared to the standard method. The research involved two cohorts of pregnant individuals from different healthcare sites across Catalonia, Spain, included between April 2020 and April 2022. One cohort underwent the standard screening method for GDM, while the other cohort underwent the alternative screening method based on plasma glucose and HbA1c levels (fasting or non-fasting). The aim was to assess the impact of using alternative diagnostic

criteria for gestational diabetes mellitus (GDM) during the COVID-19 pandemic on GDM prevalence and obstetrical and perinatal outcomes as compared with the standard screening method, as well as the financial impact.

The findings revealed several important insights. Firstly, the standard screening group had a higher GDM prevalence than the alternative screening group (10.8% vs 6.9%). Interestingly, the primary outcome (LGA rate) was similar between the two groups. An adjusted analysis showed no significant differences in the composite adverse outcomes for pregnancy, delivery, and neonatal complications between both groups.

Again, the sub-analysis of individuals diagnosed with GDM showed no significant differences in LGA rates or composite adverse outcomes for pregnancy, delivery, and neonatal complications between both groups. Although there was a higher need for insulin therapy in the alternative screening group, this did not translate into adverse outcomes. This suggests that the alternative screening method, being a less specific and less sensitive method, is not able to identify patients with less severe GDM.

Furthermore, a cost analysis revealed that the alternative screening method was cheaper than the standard screening method. The average cost per patient for the alternative screening was 46.0 euros, while the cost for the standard screening was 85.6 euros.

This study suggests that the alternative GDM screening criteria, implemented during the COVID-19 pandemic, has a lower GDM diagnostic rate than the standard method. However, the obstetrical, delivery and neonatal outcomes for both groups remained comparable. The alternative screening method had also a lower diagnostic sensitivity. This research highlights the adaptability of diagnostic screening approaches in response to unique circumstances, such as those of a pandemic, and the importance of assessing the impact of implementing a new screening method on both healthcare outcomes and costs.

4.2. Impact and Correlation with the Literature

Elevated fasting glucose levels in the first trimester are consistently linked to a higher risk of large-for-gestational-age babies, macrosomia, and cesarean section. This association persists even after excluding individuals with a basal fasting glucose higher than 105 mg/dL [23]. The rise in fasting plasma glucose during early pregnancy [24], driven by placental hormones inducing insulin resistance, challenges the appropriateness of performing GDM screening (normally performed at 24-28 weeks) before 20 weeks of gestation.

Moreover, some studies suggest that overtreatment may lead to small-for-gestational-age infants and adverse fetal metabolic programming, contributing to cardiovascular diseases later in life for both the mother and the infant [25,26]. Thus, a key challenge is establishing evidence-based glycemic cut-off values in the first trimester for early GDM diagnosis.

Few studies have explored the addition of HbA1c to the diagnostic criteria for patients with a high GDM risk. Amylidi et al. found that first-trimester HbA1c levels were predictive of GDM, similar to the standard screening method. Adding HbA1c to the GDM diagnostic criteria with a cut-off value of 5.4% in the first trimester may have identified 18% of GDM cases in the second trimester, and a cut-off value of 5.7% may have identified 32% of cases [27]. These authors suggest that a HbA1c >6% predicts GDM with 100% accuracy and warrants treatment. In the Roeder intervention study [28], the group with fasting glucose ≥ 5.1 mmol/l (92 mg/dl) or HbA1c $\geq 5.7\%$ had more individuals with high HbA1c than with high fasting glucose. However, using HbA1c alone as the only criterion for GDM diagnosis lacks sufficient sensitivity or specificity, although higher HbA1c in the first trimester is significantly associated with an increased risk of GDM [29].

In a prospective study conducted in a Spanish population, adding HbA1c to the diagnostic criteria in the first-trimester and second-trimester GDM screenings without additional criteria, a cut-off value of <4.7% for HbA1c had a 95.3% negative predictive value for GDM in the first trimester, potentially reducing screening costs by 5.7% [30]. However, this cut-off value had only a 14% positive predictive value in the second trimester. With a higher cut-off value for HbA1c, such as 5.6%, the positive predictive value in the second trimester increased to 31.6%, with a maintained negative

predictive value of 89.8%. The results of that study highlight the importance of optimizing first-trimester screening criteria for early GDM prediction, including glycemic criteria, to avoid unnecessary tests and costs.

Several studies show that glucose levels at different points of the OGTT test in the first trimester are significantly higher in subjects who later develop GDM in the second trimester [31,32]. This suggests that adding fasting glucose levels to the diagnostic criteria may also have a high negative predictive value, potentially reducing the number of subjects requiring GDM screening. There is no consensus on a particular fasting glucose threshold for early GDM diagnosis. Therefore, well-designed trials are needed to establish a fasting glucose cut-off value and determine the risk of adverse maternal and fetal outcomes [33].

Basal fasting glucose decreases from week 5 to week 19 of gestation in all pre-gestational BMI groups [34]. However, being overweight or obese before pregnancy is associated with increased fasting glucose between 19 and 24 gestational weeks. Therefore, combining glycemic criteria with clinical criteria, such as maternal pre-gestational BMI, may also help to develop a reliable diagnostic algorithm.

During the COVID-19 pandemic, Spanish healthcare sites approved a consensus document allowing GDM diagnosis using only glucose and HbA1c [13]. Before the pandemic, a Spanish study evaluating diagnostic criteria and neonatal outcomes in 3200 subjects using only glucose (1t-FPG $\geq$ 100 mg/dL) for GDM or gestational hyperglycemia diagnosis in the first trimester found that 79.5% of pregnant individuals with early GDM, would not have been diagnosed using the OGTT and two-step screening method [35]. On the other hand, a Spanish cohort showed that GDM prevalence during the COVID-19 pandemic, when using the alternative diagnostic criteria (glucose and HbA1c measurements), was similar to GDM prevalence in 2019, when the standard diagnostic criteria was used. Although individuals referred for GDM screening during the pandemic had more GDM risk factors, obstetrical and perinatal outcomes were comparable to those before the pandemic [17].

Thus, our study provides more evidence suggesting that including glucose and HbA1c in the same blood test improves the predictive value for GDM.

Subsequent research is essential to determine optimal cut-off values, formulate algorithms that include glucose and HbA1c, and consider additional factors, such as ethnicity or BMI. This comprehensive approach aims to further refine the predictive algorithm.

4.3. Strengths

The study on the impact of GDM diagnostic criteria during the COVID-19 pandemic (DIABECOVID STUDY) has several strong points which are of relevance during a pandemic: The study addresses a critical issue during a pandemic; compares two methods (standard method and alternative method adopted during the pandemic) for GDM screening through a comprehensive analysis; provides comprehensive outcome measures and adjusts for confounders, such as maternal age, BMI, parity, ethnicity, smoking status, and family history. This enhances the reliability of the findings. In addition to assessing clinical outcomes, the study also examines the financial impact of the alternative screening method through a cost analysis. The data shows that the alternative screening method is cheaper, providing valuable information for healthcare providers and policymakers.

4.4. Limitations

Despite its strengths, the study on the impact of an alternative GDM screening method during the COVID-19 pandemic (DIABECOVID STUDY) has some limitations. On one hand, the study involved retrospective data collection during the initial phase of the COVID-19 pandemic. This may have introduced recall bias and limit the availability of complete data, potentially affecting the accuracy of results. There was also a selection bias; the decision of whether to use the standard or alternative diagnostic criteria was made by each healthcare site. This may have introduced selection bias, as sites that adopted the alternative method might have had different patient populations or policies that may have influenced outcomes. On the other hand, the study had incomplete and

missing data: exclusion of some participants due to missing perinatal outcomes may have affected sample's representativeness, potentially introducing bias if certain individuals were more likely to be excluded. Regarding missing data, it must be noted that some relevant data, such as family history of diabetes, had a high number of missing data, which may have impacted the accuracy of the results. Lastly, the study has limited applicability, as it was conducted in only one Spanish region (Catalonia) and results may not be applicable to other regions with different patient populations, healthcare practices, and pandemic situation.

Another limitation lies in the financial impact analysis, where only screening costs were considered. Additional costs associated with overdiagnosis in the standard screening method, including the number of medical visits, indirect or quality-of-life costs for pregnant individuals, and pregnancy outcomes, were not taken into account.

4.5. Further Research

The COVID-19 pandemic revealed the challenges pregnant individuals face in accessing GDM screening. Alternative screening methods, such as those implemented during the pandemic, provide an opportunity to rethink and optimize the GDM screening process.

One recent randomized controlled trial [36], comparing lower (92 mg/dl) versus higher (99 mg/dl) glycemic criteria for GDM diagnosis showed that using the lower glycemic criteria led to a higher diagnostic rate than using the higher glycemic criteria; therefore, use of healthcare services was greater in the group with lower glycemic criteria. Overall, the risks of giving birth to a large-for-gestational-age infant and other infant or maternal complications were not lower for the group with lower glycemic criteria. Although using the lower glycemic criteria led to a greater proportion of individuals receiving a GDM diagnosis and treatment, it did not lead to clear health benefits. However, using the lower glycemic criteria led to a greater use of healthcare services. The authors found more infants with hypoglycemia warranting treatment in the group with lower glycemic criteria. Neonatal hypoglycemia is associated with adverse neurodevelopmental outcomes later in life, so follow-up will be needed to determine whether GDM diagnosis and treatment leads to benefits or adverse outcomes later in life. Also, cost analyses will be necessary to aid decision-making.

New prospective clinical trials should aim to compare different GDM screening methods, including universal screening, risk factor-based screening, and alternative screening, while considering their impact on maternal and neonatal outcomes, healthcare costs, and healthcare system adaptability during crises. There is also a need to collect patient-reported outcomes and quality-of-life questionnaires to determine whether the new screening method has any positive impact on patients.

5. Conclusions

The DIABECOVID study found that the alternative screening method used during the pandemic resulted in a lower GDM prevalence, although obstetric and perinatal outcomes were comparable between the two groups. Notably, glycemic control and insulin treatment did not significantly differ between individuals receiving a GDM diagnosis and those who did not. However, the alternative screening method had a lower GDM diagnostic rate. Despite this, our study suggests that the alternative screening method for GDM screening can be effective in a population with a 12% incidence of GDM, as well as cheaper. These results need to be confirmed in prospective randomized controlled trials.

References

1. Tsakiridis I, Giouleka S, Mamopoulos A, Kourtis A, Athanasiadis A, Filopoulou D, et al. Diagnosis and management of gestational diabetes mellitus: An overview of national and international guidelines. *Obstet Gynecol Surv* [Internet]. 2021;76(6):367–81.
2. Kuo C-H, Li H-Y. Diagnostic strategies for gestational diabetes mellitus: Review of current evidence. *Curr Diab Rep* [Internet]. 2019;19(12).

3. Diagnostic testing for gestational diabetes mellitus (GDM) during the COVID-19 pandemic: Antenatal and postnatal testing advice [Internet]. Com.au. 2020.
4. Yamamoto JM, Donovan LE, Feig DS, Berger H. Temporary alternative screening strategy for gestational diabetes screening during the COVID-19 pandemic—the need for a middle ground. *Can J Diabetes* [Internet]. 2022;46(2):204–6.
5. Screening for GDM during COVID restrictions—Recommendations from New Zealand Society for the Study of Diabetes [Internet]. New Zealand Society for the Study of Diabetes. 2020.
6. Thangaratinam S, Khan R, Blott M, et al. Guidance for maternal medicine services in the evolving coronavirus (COVID-19) pandemic. April 2020;
7. Codina M, Corcoy R, Goya MM, Acosta Delgado D, Ballesteros Pérez M, Bandres Nivelá MO, et al. Actualización urgente: alternativa temporal para el diagnóstico de hiper-glucemia gestacional y el seguimiento de estas mujeres y aquellas con diabetes pregestacional durante la pandemia COVID-19. Consenso del Grupo Español de Diabetes y Embarazo (GEDE) de la Sociedad Española de Diabetes (SED) y la Sociedad Española de Ginecología y Obstetricia (SEGO). *Endocrinol Diabetes Nutr* [Internet]. 2020 ;67(8):545–52.
8. Royal College of Obstetricians and Gynaecologists. Guidance for Maternal Medicine Services in the Evolving Coronavirus (COVID-19) Pandemic. 2020.
9. Australasian Diabetes in Pregnancy Society. Diagnostic testing for Gestational Diabetes Mellitus (GDM) during the COVID 19 pandemic. Antenatal and Postnatal Testing Advice. 2020.
10. Thangaratinam S, Cooray SD, Sukumar N, Huda MSB, Devlieger R, Benhalima K, et al. ENDOCRINOLOGY IN THE TIME OF COVID-19: Diagnosis and management of gestational diabetes mellitus. *Eur J Endocrinol* [Internet]. 2020;183(2):G49–56.
11. Kasuga Y, Saisho Y, Ikenoue S, Ochiai D, Tanaka M. A new diagnostic strategy for gestational diabetes during the COVID-19 pandemic for the Japanese population. *Diabetes Metab Res Rev* [Internet]. 2020;36(8).
12. Vambergue A, Jacqueminet S, Lamotte M-F, Lamiche-Lorenzini F, Brunet C, Deruelle P, et al. Three alternative ways to screen for hyperglycaemia in pregnancy during the COVID-19 pandemic. *Diabetes Metab* [Internet]. 2020;46(6):507–10.
13. Codina M, Corcoy R, Goya MM. An update on gestational hyperglycemia diagnosis during the COVID-19 pandemic. *Endocrinol Diabetes Nutr (Engl)* [Internet]. 2020;67(8):545–52.
14. d’Emden M, McLeod D, Ungerer J, Appleton C, Kanowski D. Development of a fasting blood glucose-based strategy to diagnose women with gestational diabetes mellitus at increased risk of adverse outcomes in a COVID-19 environment. *PLoS One* [Internet]. 2020;15(12):e0243192.
15. van Gemert TE, Moses RG, Pape AV, Morris GJ. Gestational diabetes mellitus testing in the COVID-19 pandemic: The problems with simplifying the diagnostic process. *Aust N Z J Obstet Gynaecol* [Internet]. 2020;60(5):671–4.
16. McIntyre HD, Gibbons KS, Ma RCW, Tam WH, Sacks DA, Lowe J, et al. Testing for gestational diabetes during the COVID-19 pandemic. An evaluation of proposed protocols for the United Kingdom, Canada and Australia. *Diabetes Res Clin Pract* [Internet]. 2020;167(108353):108353.
17. Molina-Vega M, Gutiérrez-Repiso C, Lima-Rubio F, Suárez-Arana M, Linares-Pineda TM, Cobos Díaz A, et al. Impact of the gestational diabetes diagnostic criteria during the pandemic: An observational study. *J Clin Med* [Internet]. 2021;10(21):4904.
18. National Diabetes Data Group. Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* [Internet]. 1979;28(12):1039–57.
19. Control prenatal del embarazo normal. Guía de asistencia Práctica [Internet].
20. Protocol de seguiment de l'embaràs a Catalunya [Internet]. Gencat.cat.
21. Goya M, Codina M. Diabetes mellitus and pregnancy. Updated clinical practice guideline 2021. Executive summary. *Endocrinol Diabetes Nutr (Engl)* [Internet]. 2023;70:1–6.
22. Newcombe RG. Interval estimation for the difference between independent proportions: comparison of eleven methods. *Stat Med* [Internet]. 1998;17(8):873–90.
23. Riskin-Mashiah S, Younes G, Damti A, Auslender R. First-trimester fasting hyperglycemia and adverse pregnancy outcomes. *Diabetes Care* [Internet]. 2009;32(9):1639–43.
24. Mills JL, Jovanovic L, Knopp R, Aarons J, Conley M, Park E, et al. Physiological reduction in fasting plasma glucose concentration in the first trimester of normal pregnancy: The diabetes in early pregnancy study. *Metabolism* [Internet]. 1998;47(9):1140–4.
25. Langer O, Levy J, Brustman L, Anyaegbunam A, Merkatz R, Divon M. Glycemic control in gestational diabetes mellitus-How tight is tight enough: Small for gestational age versus large for gestational age? *Am J Obstet Gynecol* [Internet]. 1989;161(3):646–53.
26. Catalano PM, Mele L, Landon MB, Ramin SM, Reddy UM, Casey B, et al. Inadequate weight gain in overweight and obese pregnant women: what is the effect on fetal growth? *Am J Obstet Gynecol* [Internet]. 2014;211(2):137.e1-137.e7.

27. Amylidi S, Mosimann B, Stettler C, Fiedler GM, Surbek D, Raio L. First-trimester glycosylated hemoglobin in women at high risk for gestational diabetes. *Acta Obstet Gynecol Scand* [Internet]. 2016;95(1):93–7.
28. Roeder HA, Moore TR, Wolfson MT, Gamst AC, Ramos GA. Treating hyperglycemia in early pregnancy: a randomized controlled trial. *Am J Obstet Gynecol MFM* [Internet]. 2019;1(1):33–41.
29. Valadan M, Bahramnezhad Z, Golshahi F, Feizabad E. The role of first-trimester HbA1c in the early detection of gestational diabetes. *BMC Pregnancy Childbirth* [Internet]. 2022;22(1).
30. Benaiges D, Flores-Le Roux JA, Marcelo I, Mañé L, Rodríguez M, Navarro X, et al. Is first-trimester HbA1c useful in the diagnosis of gestational diabetes? *Diabetes Res Clin Pract* [Internet]. 2017;133:85–91.
31. Peng X, Liu M, Gang J, Wang Y, Ma X. Use of oral glucose tolerance testing and HbA1c at 6–14 gestational weeks to predict gestational diabetes mellitus in high-risk women. *Arch Gynecol Obstet* [Internet]. 2022;307(5):1451–7.
32. Phaloprakarn C, Tangjitgamol S. Use of oral glucose tolerance test in early pregnancy to predict late-onset gestational diabetes mellitus in high-risk women. *J Obstet Gynaecol Res* [Internet]. 2008;34(3):331–6.
33. Bhattacharya S, Nagendra L, Krishnamurthy A, Lakhani OJ, Kapoor N, Kalra B, et al. Early gestational diabetes mellitus: Diagnostic strategies and clinical implications. *Med Sci (Basel)* [Internet]. 2021;9(4):59.
34. Wei Y-M, Liu X-Y, Shou C, Liu X-H, Meng W-Y, Wang Z-L, et al. Value of fasting plasma glucose to screen gestational diabetes mellitus before the 24th gestational week in women with different pre-pregnancy body mass index. *Chin Med J (Engl)* [Internet]. 2019;132(8):883–8.
35. González González NL, González Dávila E, Bugatto F, Vega-Guedes B, Pintado P, Tascón L, et al. Fasting glucose for the diagnosis of gestational diabetes mellitus (GDM) during the COVID-19 pandemic. *Nutrients* [Internet]. 2022;14(16):3432.
36. Crowther CA, Samuel D, McCowan LME, Edlin R, Tran T, McKinlay CJ. Lower versus higher glycemic criteria for diagnosis of gestational diabetes. *N Engl J Med* [Internet]. 2022;387(7):587–98.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.